Number and proportion of *P. falciparum* gametocytes vary from acute infection to chronic

parasite carriage despite unaltered sexual commitment rate

Hannah van Dijk[1], Nathalia F Lima[1], Martin Kampmann[2], Michael Gabel[3], Usama Dabbas[1,2],

Safiatou Doumbo[4], Hamidou Cisse[4], Shanping Li[5], Myriam Jeninga[6], Richard Thomson-Luque[1],

Didier Doumtabe[4], Michaela Petter[6], Kassoum Kayentao[4], Aissata Ongoiba[4], Teun Bousema[7],

Peter D Crompton[5], Boubacar Traore[4], Frederik Graw[3,8], Silvia Portugal[1,2*]

[1] Center for Infectious Diseases, Parasitology, Heidelberg University Hospital, Heidelberg,

Germany

[2] Max Planck Institute for Infection Biology, Berlin, Germany

[3] BioQuant-Center for Quantitative Biology, Heidelberg University, Heidelberg, Germany

[4] Mali International Center of Excellence in Research, University of Sciences, Techniques and

Technologies of Bamako, Mali

[5] Laboratory of Immunogenetics, National Institute of Allergy and Infectious Diseases, National

Institutes of Health, Rockville, Maryland, USA

[6] Mikrobiologisches Institut-Klinische Mikrobiologie, Universitätsklinikum Erlangen

[7] Radboud University Medical Center, Nijmegen, the Netherlands

[8] Interdisciplinary Center for Scientific Computing, Heidelberg University, Heidelberg, Germany

[*]Corresponding author email: portugal@mpiib-berlin.mpg.de (SP)

The authors declare no conflict of interest

Abstract

Introduction In areas of seasonal malaria, resuming transmission every wet season relies on persistent asexual parasites during dry season that maintain the ability to produce gametocytes until the mosquito population resurges following the start of annual rains. Although human asymptomatic *P. falciparum* reservoirs in the dry season are widely recognized, the longitudinal dynamics of parasite sexual commitment and gametocytogenesis remain unclear.

Methods We compared the expression of genes related with sexual commitment and gametocytogenesis of *P. falciparum*, the density and proportion of *P. falciparum* gametocytes, and the blood concentrations of phospholipids potentially involved in gametocytogenesis from asymptomatic subjects at the end of the dry season versus subjects with symptomatic malaria in the wet season. Furthermore, we verified whether number and proportion of gametocytes in asymptomatic vs clinical cases had similar trends in areas of seasonal and perennial transmission settings. Finally, we adapted a within-host mathematical model considering asexual parasites, sexually committed parasites, and gametocytes to infer the dynamics of gametocyte number and proportion as *P. falciparum* infections progress.

Results We found that transcripts of genes specific of late-stage gametocytes were predominantly upregulated in asymptomatic infections at the end of the dry season, which also showed increased proportions of circulating mature gametocytes compared to clinical malaria cases. On the other hand, blood densities of gametocytes and asexual parasites were lower in chronic asymptomatic individuals compared to clinical malaria. In addition, the levels of parasite transcripts involved in sexual commitment were unaltered throughout the year.

Discussion Our experimental data in combination with mathematical modelling support a scenario in which gametocyte density and proportion diverge as infections progress from recent transmission to chronic carriage, without significant alterations in the rate of sexual commitment over time.

Key words *Plasmodium*, *P. falciparum*, gametocytes, dry season, asymptomatic, parasite sexual commitment, malaria mathematical model

Introduction

It has long been known that the malaria-causing parasite *Plasmodium falciparum* requires a mosquito host to complete its lifecycle and to be transmitted between humans (1). However, it remains unclear if, in the human host, the parasite can dynamically allocate resources to the differentiation of sexual-stage gametocytes that are transmitted to mosquitos, particularly in areas of seasonal malaria where the size of the mosquito population fluctuates dramatically during the dry and wet seasons. During blood feedings, infected Anopheles mosquitoes deposit *P. falciparum* sporozoites in the human dermis that find their way into circulation, reaching the liver where they infect hepatocytes. After 7-10 days of parasite replication, each infected hepatocyte releases thousands of merozoites into the blood stream (2), where they start continuous asexual intra-erythrocytic replication cycles of ~48 h, from merozoite invasion of red blood cells (RBCs), through the ring- and trophozoite-stages, until the multinucleated schizont releases new merozoites (3). A small proportion of intra-erythrocytic asexual parasites differentiate into sexual-stage gametocytes (4). Instead of another ~48 h asexual schizogony leading to the 8-30 new merozoites ready to continue asexual cycles (5), the parasite subsequently develops into sexually committed schizonts leading to 8-30 sexually committed merozoites. Upon infection of new RBCs each sexually committed merozoite produces one stage I gametocyte (6, 7) that through a period of 8-12 days will go through five developmental stages (8), where nuclear reshaping is driven by intranuclear microtubules (9, 10). Stages II to IV of gametocytaemia take place in the bone marrow or spleen (11, 12), while stage V gametocytes are again present in peripheral blood, each lasting up to ~20 days (13, 14) waiting for another mosquito bite. Following a blood meal gametocytes encounter the right environment for sexual reproduction in the mosquito midgut (15). They transform into female and male gametes that fuse into a zygote originating oocysts which invade the midgut wall to generate sporozoites. These sporozoites later reach the mosquito salivary glands and can restart the parasite lifecycle after being taken up in the next blood meal (16). The continued presence of mosquitoes and humans in a geographical region allows for perennial *P. falciparum* transmission.

However, in areas where the dry season is several months long, the mosquito population can

plummet to near zero (17), interrupting transmission. *P. falciparum* faces the challenge of surviving

in its human host for months while waiting for the return of mosquitoes as the wet season ensues

(18-20). During mosquito-free dry periods, gametocyte production is largely wasted, as

gametocytes cannot survive to bridge two wet seasons (13, 14). Although curtailing gametocyte

production rates during the dry season would seem beneficial to the parasite, such a strategy

would require a mechanism to reactivate gametocytogenesis with the return of rain and mosquitos.

Host factors such as age (21), haematocrit (22), haemoglobin type (23) and duration of malaria

symptoms (24) have been linked to gametocytaemia in individuals with clinical malaria; and higher

frequencies of gametocyte positive individuals have been observed in chronic asymptomatic

infections compared to acute infections (25-28). Additionally, various in vitro conditions have been

shown to modulate sexual commitment and thus gametocytogenesis (6, 29-31). The specific effect

of lysophosphatidylcholine (Lyso-PC) in repressing the pro-sexual commitment transcription factor

AP2G has been recently identified, with low Lyso-PC concentrations increasing the proportion of

sexual forms in vitro (31). Furthermore, gametocyte development 1 (GDV1) has been identified as

the de-repressing effector which counters the epigenetic silencing and promotes AP2G expression

leading to the induction of gametocytogenesis (32), and homeodomain protein 1 (HDP1) was

recently identified as a positive transcriptional regulator essential for gametocyte development (33).

Still, it is not fully understood how sexual investment and gametocytaemia is regulated in vivo, and

if or how transmission to mosquitoes can be optimized by the parasite to conserve resources.

Recently, we have shown that *P. falciparum* parasites isolated at the end of the dry season are

transcriptionally distinct from those of subjects with febrile malaria in the transmission season (18,

34), but alteration of sexual commitment and gametocytogenesis were not investigated. The

observed transcriptional differences reflected longer parasite circulation within each ~48 h intra-

erythrocytic cycle in the dry season, likely increasing splenic clearance of infected cells and thus

maintaining the low parasitaemias observed during the dry season (18). Here, we aimed to

examine sexual commitment and gametocytogenesis of parasites persisting through the dry

season and explored possible parasite adaptations during extended periods in the human host. We

quantified sexual commitment and female and male gametocyte densities in blood samples

collected from children at different times of the year with different clinical presentations and

seasonality patterns, and related them to total parasite density and infection length. Furthermore,

we determined longitudinally the Lyso-PC concentration in plasma of children, and assessed its

ability to induce *P. falciparum* sexual commitment and gametocytogenesis in vitro. Finally, we

adapted a previously described within-host mathematical model that considers asexual parasites,

sexually committed parasites, and gametocytes (35) to infer the dynamics of gametocyte number

and proportion as *P. falciparum* infections progress from recently transmitted to chronic

asymptomatic carriage over several months.

Materials and Methods

Gametocyte gene expression analysis. We used RNAseq data of leucocyte-depleted blood

pellets from 12 children with asymptomatic malaria at the end of the dry season and an age- and

gender-matched cohort presenting with their first malaria case in the ensuing transmission season

reported in (18). Originally 808 gametocyte specific genes were reported (36), and we counted 786

after excluding repetitions and converting nomenclature to its most recent version (Supplementary

Table 1). Out of these 786 genes, we detected normalized reads of 333 gametocyte-specific genes

in the RNAseq data set (18) to perform a principal component analysis. Using Deseq2, 205 genes

were determined to be differentially expressed between the dry season and malaria cases.

Analysis of early and late gametocyte markers in RNA-seq data. We compiled a list of early

and late gametocyte specific genes based on three RNA-seq studies comparing distinct parasite

stages, including data from Poran et al. 2017(37) obtained after request from the corresponding

author (compared day 2 versus day 9 of gametocytogenesis by single cell RNAseq); data from

Lopez-Barragan et al. 2011 (38) (compared day 4 versus day 15 of gametocytogenesis by bulk

RNAseq) and data from Jeninga et al., *under review* (compared day 4 of gametocytogenesis

versus day 10 of FACS sorted male and female gametocytes by bulk RNAseq). Poran et al.

selected early and late gametocyte genes as genes showing a 20-fold higher expression in early or

late gametocytes over the respective other group. Late gametocyte genes were also required to

show a 20-fold higher expression in late gametocytes compared to ring stage parasites, this was

not the case for early gametocyte genes. In Lopez-Barrgagan's data, early gametocyte genes were

defined as genes having a 10-fold expression increase in early gametocytes compared to late

gametocytes. Genes showing a 20-fold higher expression in late stages were designated as late

gametocyte genes. Expression differences to asexual stages were not considered. In Jeninga et al.

under review genes showing a 3-fold or higher expression difference to both asexual stages and

late gametocytes were selected as early gametocyte genes, regardless of gametocyte sex.

Intersecting the full list of early and late gametocyte genes with genes transcribed in circulating

parasites stages, defined by RNAseq data of Andrade et al., generated a final list of 109 early

gametocyte genes and 54 late gametocyte genes to evaluate expression differences of these

genes between dry and rainy seasons. Intersecting the early and late gametocyte list with all

gametocyte genes reported by Painter et al. we receive 456 gametocyte genes that are detected in

the wet vs dry season RNAseq dataset.

Study subjects and ethical approval. Approval by the Ethics Committee of Heidelberg University

Hospital, the Faculty of Medicine, Pharmacy and Odontostomatology (FMPOS) at the University of

Bamako, and the National Institute of Allergy and Infectious Diseases of the National Institutes of

Health Institutional Review Board was obtained to conduct the study registered at ClinicalTrials.gov

with the identifier NCT01322581. Clinical data and samples were obtained from 2017 to 2019 in a

cohort study previously described (39), conducted in Kalifabougou, Mali, where malaria

transmission occurs between June and December, and antimalarial drugs are provided by a single

clinic and pharmacy. All participants and the parents/guardians of included children provided

written informed consent. Exclusion criteria at enrolment included haemoglobin <7 g/dL, axillary

temperature ≥37.5°C, acute systemic illness, or use of antimalarial or immunosuppressive

medications in the 30 days preceding enrolment. Clinical malaria cases were prospectively

determined by passive surveillance and defined by axillary temperature ≥37.5°C, ≥2500 asexual

parasites/µL of blood, and no other evident cause of fever. Malaria episodes were treated with a 3-

day standard course of artemether/lumefantrine according to Malian national guidelines. Cross-

sectional clinical visits and blood draws were performed late in the transmission season (October),

and at the beginning (January), mid (March) and end (May) of the dry season.

Sample collection. Venous blood (4 mL if donor age was below 4 years, and 8 mL if above) was

drawn by venepuncture on wet and dry season cross-sectional visits, and at first malaria episode of

the transmission season. Samples are referred as MAL for malaria cases or by time point of

collection. Blood was collected in sodium citrate-containing cell preparation tubes (Vacutainer CPT

Tubes, BD) and transported to the laboratory where RBC pellet and plasma were separated and

stored at −80°C within three hours of blood draw. Plasma used for lipidomic analysis and induction

of gametocytogenesis was separated by centrifugation and immediately frozen in liquid N_2.

RNA extraction and quantification of parasite transcripts. RNA was isolated from 200 µl RBC

pellet by phenol-chloroform extraction using TRIzol (Ambion), followed by DNAse treatment using

the DNAse I Amplification Grade Enzyme (Invitrogen). RNA was quantified by absorption

measurements at 230, 260 and 280 nm on a Citation3 Imaging plate reader, followed by cDNA

synthesis using the Superscript IV Enzyme (Invitrogen) according to manufacturer instructions.

Taqman RT-qPCR multiplex assay (Applied Biosystems) was performed on a qTower (Analytic Jena)

with primers to quantify the female gametocyte marker ookinete surface protein P25

(PF3D7_1031000), the male gametocyte transcript PFMGET (Pf3D7_1469900), the master

regulator of sexual commitment Ap2G (PF3D7_1222600) and a parasite housekeeping gene

(Glycine-tRNA ligase, PF3D7_1420400). Amplification efficiency (E) was estimated on the basis of

the equation E=(10^(-1/slope)-1)x100 (Supplementary Figure 1). NF54 parasite line expressing

DyneinGFP and carrying an episomal P47mCherry plasmid (DynGFP/P47mCherry) was used for

sorting of male and female gametocytes through their respective specific mCherry and/or GFP expression (40). Standard curves of $10-10^6$/ml of sorter-purified female or male gametocytes were generated by 10-fold serial dilution in 50% haematocrit (Supplementary Figure 1). qPCR expression of female-specific P25 in female (mCherry+) sorted parasites was high and decreased with gametocyte densities, while male gametocytes (GFP+) show ~12-fold lower expression of P25, which may be explained by ~6% impurity originating from the opposite sex during sorting (Supplementary Figure 1). Likewise, PfMGET expression was approximately 10-fold higher expressed in male gametocytes (GFP+) compared to females (mCherry+), consistent with 5-7% impurity. Linear regression fit for the independent populations revealed a good fit with R^2-values of 0.93-0.97 by P25 expression and 0.96-0.98 by PfMGET expression (Supplementary Figure 1). The asexual parasites standard curve, was generated with Pf2004 asexual culture was counted by FACS and Glycine-tRNA ligase expression was quantified in dilution series of $10-10^7$ asexual late rings per ml. The combination of asexual and sexual standard curves was used for qPCR quantification of female or male proportion of total parasitaemias throughout the season. To account for fluorescence leakage between channels a compensation matrix was calculated based on single colour fluorescence measurements by an application specialist at Analytic Jena. Gexp5 (PF3D7_0936600) expression levels were quantified in a Taqman RT-qPCR duplex assay together with Glycine-tRNA ligase as a housekeeping gene. Expression levels of GDV1 (PF3D7_0935400), and MRSP1 (PF3D7_1335100) were determined in further Taqman RT-qPCR assays. All primers and probes sequences are listed in Supplementary Table 2. Relative quantification was performed by ΔΔCt method using Glycine-tRNA ligase as housekeeping gene and MAL samples a calibrator group.

Parasite culture and preparation of a standard curve for gametocyte quantification. Pf2004 parasites were cultured at 37° C in 5% O_2, 5% CO_2 and 90% N_2, at 5% haematocrit in RPMI-1640 medium (with L-glutamine and HEPES (Gibco)). Parasite culture medium was supplemented with 0.24% Sodium Bicarbonate (Gibco), 100µM Hypoxanthine (cc pro), 25 µg/mL gentamycin (Gibco)), and 0.25% Albumax II (Gibco). Medium was changed daily, parasitaemia assessed by Giemsa-

stained blood smear and the culture split according to experimental needs.

*PfDyn*GFP/*PfP47*mCherry parasite lines (41) were cultured in presence of 15 µg/ml of Blasticidin-

S-HCL (Invitrogen) and 2 µM Pyrimethamine (Sigma Aldrich) as described for non-integrating

plasmids and P52 integrating plasmid. Parasites were grown to >4% parasitaemia when 50 mM N-

Acetylglucosamine (Sigma Aldrich) was added to the culture to eliminate asexual stages. After 4

consecutive days of N-Acetylglucosamine treatment, culture in complete RPMI was resumed.

Gametocyte development was monitored daily by smear, and on day 5 after start of gametocyte

induction magnetic enrichment was performed with MACS columns (Miltenyi Biotec). Gametocytes

were sorted on a BD FACSAria III on day 8, keeping cells at 4° C in suspended activation buffer,

(gating strategy in Supplementary Figure 1). RNA of a defined numbers of sorted male and female

gametocytes was extracted and reverse transcribed, a standard curve prepared by serial dilution

and used for RT-qPCR as described above.

Phospholipid analysis. May and MAL plasma samples (shown in Figure 1) were analysed by

shotgun lipidomics at Lipotype as described in (42). Briefly, automated samples extraction was

followed by direct sample infusion and high-resolution Orbitrap mass spectrometry including lipid

class-specific internal standards to assure absolute quantification of LysoPC and LysoPE, using

LipotypeXplorer for identification of lipids in the mass spectra. Oct, Mar, May and MAL samples

(shown in Supplementary Figure 4) were analysed by Liquid chromatography-mass spectrometry

(LC-MS). Samples were extracted, processed, and analysed at the Lipidomics Unit at the Institute

of Physiological Chemistry in Mainz, using the liquid-liquid extraction protocol as described (43). All

values were normalized to the analysed plasma volume.

In vitro induction of *P. falciparum* sexual commitment and gametocytogenesis. In vitro

induction of *P. falciparum* sexual commitment was performed as described by (Brancucci et al.,

2015). Pf2004 parasites were synchronized to a 4 h window by repeated treatment with 5% sorbitol

and MACS purification. Parasites at 26 ± 2 h post invasion (hpi) were plated on 48-well plates in

500 µl culture volume at 0.8-1% parasitaemia and 2.5% haematocrit and cultured in minimum fatty

acid medium (mFA, RPMI1640 with 100 µM hypoxanthine, 25 µg/ml gentamycin, 0.39% fatty-acid-

free BSA, 30 µM palmitic acid and 30 µM oleic acid) supplemented with 25% plasma from

asymptomatic children at the end of the dry season (PIMay), or 25% plasma from malaria cases

during the rainy season (PIMAL), in addition to the conditions used by Brancucci et al. (mFA alone,

and mFA supplemented with 25% German plasma (PIDE)). After 24 h of treatment, the medium

was changed to culture medium and subsequently changed daily. At 48 h after beginning of

treatment 30 IU/ml heparin were added to block re-invasion of parasites and eliminate asexual

forms. At 0, 48 and 120 h post treatment RBC pellets were collected for RNA quantification of

gametocyte markers by RT-qPCR; parasitaemia and gametocytaemia determination by flow

cytometry (50 µl of culture stained with 5x SYBR Green II (Invitrogen) and 7,5 µM MitoTracker

(Applied Biosystems) for 30 min at 37 °C and acquired with CountBright Absolute Counting Beads

(Thermo Ficher) on a FACS Canto II or an LSR II and analysed using FlowJo software 10.2 or

higher versions (Tree Star)). Additionally, Giemsa-stained blood smears were prepared at 48 h and

120 h post-treatment and blindly analysed by microscopy, counting RBCs, parasites, and

gametocytes in 18 fields of view. Multiplication rate of parasites was calculated as $\frac{parasitemia\ 48\ h}{parasitemia\ 0\ h}$,

and conversion rate as $\frac{gametocytemia\ 120\ h}{parasitemia\ 48\ h}$.

Quantification of parasites in samples of Ugandan asymptomatic carriers by RT-qPCR. We

compared the Malian data from dry and wet season with data from 150 Ugandan individuals

enrolled in a 2-year longitudinal study including monthly standardized clinical evaluation and blood

collection (44) retrieved from ClinEpiDB

(https://clinepidb.org/ce/app/record/dataset/DS_51b40fe2e2122). Asymptomatic *P. falciparum*

positive individuals (20.4 years ± 17.2, median: 13.3 years) were included based on the following

criteria: i) ≥ two PCR+ time points, ii) ≤ 3 negative time points in between positive, iii) same clone

(determined by deep amplicon sequencing of atypical membrane antigen 1 (AMA1) as described in

(45) present at all time points of one asymptomatic infection. Additionally, we used samples from

an age-matched group of 38 clinical malaria cases (tympanic temperature > 38°C or history of

fever in the previous 24 hours and *P. falciparum* positivity by microscopy). *P. falciparum* parasite

and gametocyte density in blood samples were measured by RT-qPCR of multi copy conserved

var gene acidic terminal sequence (46) and P25 (P25, PF3D7_1031000), respectively. Average

levels of parasitaemia and gametocytaemia during each asymptomatic infection, were estimated as

the area under curve over the course of infection divided by length of infection. For symptomatic

infections, only the quantification before treatment on day of diagnosis was considered. For

comparison between two data sets, a multiple non-parametric comparison between groups of

asymptomatic and symptomatic in Mali and Uganda has been performed using Tukey-Correction

for multiple comparisons.

Mathematical model. The developed mathematical model extends a model used previously to

describe asexual parasite and gametocytes dynamics (35). The model considers 3 different main

parasite populations, i.e., distinguishing between asexual parasites (P), sexually committed

parasites (P_G) and gametocytes (G). Asexual parasites develop and replicate within a red blood

cell over a cycle of length a_L=48h after which mature schizonts burst and release a number of r_p =

18-32 merozoites that are able to invade new red blood cells. During this replication cycle,

parasites are assumed to be removed through splenic clearance and active immune responses. In

extension to the previous model, we assume that this removal of asexual and committed parasites

follows an age dependent death rate, σ_a that differs in distinct phases of acute and chronic *P.*

falciparum infection. The following iterative equation describes asexual parasite growth at discrete

time steps for parasite age, *a*, and time, *t*, increments of one hour.

$$P(a,t) = \begin{cases} P(a-1, t-1)e^{-\sigma_a} \,, a = 2,3,\dots,a_L \\ r_p \; P(a_L, t-1)e^{-\sigma_a} \,, a = 1 \end{cases} \tag{1}$$

$$P(a_s + 1, t) = (1-f)P(a_s, t-1)e^{-\sigma_a} \tag{2}$$

292   The rate of parasite removal, σ_a, is assumed to increase with parasite age within one replicative

293   cycle and is described by a sigmoidal curve given by:

$$\sigma_a = \sigma(a) = \frac{\sigma_{max}}{1 + e^{p_\sigma(T_\sigma - a)}} \tag{3}$$

The parameter σ_{max} defines the maximal rate of parasite removal, T_σ the age at which 50% of

parasites are removed, and p_σ a scaling parameter.

At parasite trophozoite stage of a_s= 20hpi parasites adhere to the endothelium when at the same

time a fraction, f, of parasites enters sexual commitment. Committed parasites continue the rest of

the replication cycle producing only committed parasites that enter the first gametocyte stage at the

sequestration age, a_s, of the following replication cycle.

$$P_G(a_s + 1, t) = f\, P(a_s, t - 1)e^{-\sigma_a} \tag{4}$$

$$P_G(a, t) = \begin{cases} P(a - 1, t - 1)e^{-\sigma_a}, a = 2, 3, \ldots, a_L \text{ except } a = a_s \text{ and } a = a_s + 1 \\ r_p\, P(a_L, t - 1)e^{-\sigma_a}, a = 1 \end{cases} \tag{5}$$

303   Gametocytes develop over 5 distinct stages of which only mature stage 5 is found in circulation.

304   The development of gametocytes is then modelled by:

$$G_1(t) = G_1(t - 1)e^{-(m - \delta_G)} + \frac{P_G(a_s - 1, t - 1)e^{-\sigma_a}(1 - e^{-(m + \delta_G)})}{m + \delta_G} \tag{6}$$

$$G_2(t) = G_2(t - 1)e^{-(m - \delta_G)} + \frac{m\,G_1\,(t - 1)\,(1 - e^{-(m + \delta_G)})}{m + \delta_G} \tag{7}$$

$$G_3(t) = G_3(t - 1)e^{-(m - \delta_G)} + \frac{m\,G_2\,(t - 1)\,(1 - e^{-(m + \delta_G)})}{m + \delta_G} \tag{8}$$

$$G_4(t) = G_4(t-1)e^{-(m-\delta_G)} + \frac{mG_3\,(t-1)\,(1-e^{-(m+\delta_G)})}{m+\delta_G} \qquad (9)$$

$$G_5(t) = G_5(t-1)e^{-(\delta_{G5})} + \frac{mG_4\,(t-1)\,(1-e^{-(\delta_{G5})})}{\delta_{G5}} \qquad (10)$$

Immature gametocytes, that are sequestered, mature with a maturation rate, m, and are lost by

death rate δ_G , while mature circulating gametocytes are removed by a splenic clearance rate δ_{G5}.

As pointed out above the whole model is simulated by discrete time-steps of one-hour increments.

Accounting for different phases of P. falciparum infection: In our model the start of the infection is

marked by the beginning of blood stage infection, where we assume an initial parasite load, IPL,

with parasite age normally distributed with mean, μ_{IPL} and standard deviation σ_{IPL}. During the acute

phase of infection parasitaemias increase rapidly leading to clinical symptoms within the first 2

weeks of blood stage infection (47). The acute phase of infection is accompanied by the generation

of an adaptive immune response, including the production of effective antibodies, that takes ~14

days to develop (48). To account for higher parasite removal due to the immune response, we

assume an increased maximal parasite removal rate σ_{max2} for age-dependent parasite loss when

an adaptive immune response is active. Increased parasite removal is considered to occur 14 days

post start of blood stage malaria infection and indicates the beginning of the transition phase from

the acute towards the chronic asymptomatic phase of infection. In addition, as selection for less

adhesive and less virulent parasite strains by an active immune response is considered to be

crucial for the maintenance of a malaria infection over the course of several dry season months

(18), we accounted for this effect in our model by an increased value of T_σ for Eq. (3) during the

chronic asymptomatic phase, i.e., replacing T_σ by $T_{\sigma,2}$. Altering this parameter shifts clearance

rates towards affecting older asexual parasite stages, leading to a constant, but oscillating number

of asexual parasite numbers in circulation. *Parameter estimation*: The model described in Eqs (1-

10) was applied to the data assuming a fixed parasite life-cycle period of a$_L$=48 h, a sequestration

age of a_S= 20 hpi, as well as a parameter of T_σ=32 h and p_σ=0.7 characterizing age-dependent

parasite removal during the acute infection phase. All other parameters were fitted to both

symptomatic and asymptomatic time points simultaneously using an approximative Bayesian

computing approach (49). Prior distributions for the individual parameters spanning biologically

reasonable ranges have been selected in accordance with previous studies (Supplementary Table

3). Due to the high model complexity and parameter dependencies, estimations of multiple pyABC-

runs diverge to different local minima for the individual parameters. To obtain robust results for the

parameters indicating phase-dependent changes, several independent pyABC fitting runs were

compared and the 6[th] generation was chosen where parameter posterior distributions would still

overlap. For the fitting, values of individual patients in the dry season are condensed to one time

point by calculating the average levels based on the area under the curve over the course of dry

season patient surveillance. Values were log10-transformed to ensure comparability of

parasitaemia and gametocytemia. In addition, wet and dry season measurements were assumed

to represent the parasite dynamics at 2- and 6-weeks post infection, respectively. Sensitivity of the

results with regard to these time points was investigated. In addition, different values for the

fraction of sexually committed parasites, f, were tested. All code to run and analyse the

mathematical model is available at https://github.com/GrawLab/MalariaReplication.

Supplementary Table 3. List of model parameters and respective prior distributions. All prior

distributions were uniformly distributed given the indicated ranges.

Parameter	Unit	Description	Prior range	Ref for prior range	89% CI of post range
ipl	Parasite /ml	Initial parasite load	1 - 10	(Hodgson et al. 2015) (Schats et al. 2015)	1.47 - 6.54
μ_{ipl}	h	Initial parasite age	4 - 10	(Eichner et al. 2001)	4.41 - 8.22
σ_{ipl}	h	Standard deviation of initial parasite age	4.5 - 6.5	(Cao et al. 2019)	4.89 - 5.81
r_P		Replication factor	18 - 32		$2.28 - 3.13 \times 10^1$
σ_{max}	h^{-1}	Maximal removal rates during acute phase of infection	0.01- 0.15	(Andrade et al. 2020)	$0.81 - 1.07 \times 10^{-1}$
σ_{max2}	h^{-1}	Maximal removal rates during transition phase of infection	0.01- 0.4		$2.13 - 2.39 \times 10^{-1}$
δ_G	h^{-1}	Death rate immature gametocytes	1e-05 - 1e-04	(Cao et al. 2019)	$3.53 - 8.60 \times 10^{-5}$
δ_{G5}	h^{-1}	Death rate mature gametocytes	$1.39 - 4.17 \times 10^{-3}$	(Cao et al. 2019)	$1.79 - 3.98 \times 10^{-3}$
m	h^{-1}	Gametocyte maturation rate	$1.6 - 4.6 \times 10^{-2}$	(Cao et al. 2019)	$2.76 - 4.29 \times 10^{-2}$

f	h^{-1}	Fraction of sexually committed parasites	$0.5 - 1.0 \times 10^{-2}$	(Cao et al. 2019), (Eichner et al. 2001)	$5.41 - 7.68 \times 10^{-3}$
a_L	h	Life cycle length	48		
a_s	h	Sequestration age	20		
T_σ	h	Age at which 50% of parasites are removed during acute phase of infection	32	(Andrade et al. 2020)	
$T_{\sigma 2}$	h	Age at which 50% of parasites are removed during chronic phase of infection			33.6 - 34.7
P_σ		Shape parameter	0.7	(Andrade et al. 2020)	
$\sigma_{max}/\sigma_{max2}$	h	Ratio of maximal removal rate	0.025 - 15		2.06 - 2.89

Results

P. falciparum **transcription and quantification of gametocyte density and proportions.** We

previously described differences in transcription between asexual *P. falciparum* of asymptomatic

Malian children at the end of the dry season versus age- and sex- matched Malians presenting

clinical malaria in the wet season (18), but we did not investigate transcripts specific to sexual

forms of the parasite. Now, we gathered a list of genes related to gametocytogenesis from multiple

RNAseq published and unpublished studies to query potential adaptations of *P. falciparum* sexual

stages to persistence periods. Painter et al. reported, with in vitro culture parasites, 786

gametocyte-related transcripts (36), and of those 333 showed detectable number of reads in our

dry vs wet season dataset (18) (Table 1 and Supplementary Table 1). Poran et al. described with

single cell RNAseq analysis early and late gametocyte transcripts based on a 20-fold differential

expression comparing day 2 versus day 9 of gametocytogenesis (late gametocyte genes were

defined by 20-fold higher expression compared to ring stage parasites, although this was not the

case for early gametocyte genes) (37). We found that 2 out of 3 transcripts and 15 out of 17 from

early- and late-gametocyte transcripts, respectively, were detected at the end of our dry season

asymptomatic versus wet season clinical case dataset (18) (Table 1 and Supplementary Table 1).

In addition, data from Lopez-Barragan et al. 2011 compared bulk RNAseq of day 4 versus day 15

of gametocytogenesis and identified early-gametocyte genes through a 10-fold increase of

expression in early compared to late gametocytes. Genes showing a 20-fold higher expression in

late stages were designated as late-gametocyte genes, but expression differences to asexual

stages were not considered (38). Forty-one and 58 transcripts were called early- and late-

gametocyte related genes (38) and of those 32 and 44 were detected in the end of our dry season

asymptomatic versus wet season clinical case dataset (18) (Table 1 and Supplementary table 1).

Finally, Jeninga et al. (personal communication, *under review*) obtained data comparing bulk

RNAseq data of day 4 versus day 10 of gametocytogenesis in FACS-sorted male and female

gametocytes defined early-female and early-male specific gametocyte transcripts with a 3-fold or

higher expression difference to both asexual stages and late gametocytes. Nineteen transcripts

were assigned to early gametocytes independent of sex, and 38 and 186 transcripts were identified

as early-female, and early-male gametocyte-related genes respectively. Of those, 15 and 64 were

detected at the end of our dry season asymptomatic versus wet season clinical cases dataset (18)

(Table 1 and Supplementary Table 1). The full list of 1048 sexual commitment- and

gametocytogenesis-related genes showed limited overlap between the different reports

(Supplementary Table 1) potentially due to the different timepoints and comparison conditions

defined in the different studies. Intersecting this list with transcripts of circulating parasites of the 12

asymptomatic Malians at the end of the dry season and 12 clinical malaria cases in the wet season

(18) resulted in 456 transcripts. Principal component analysis of expression values of these genes

showed a clear segregation of transcription profiles based on seasonality and clinical presentation

(Figure 1A), and produced clear patterns on a heatmap clustering malaria cases (MAL) or end of

the dry season asymptomatic samples (May) by similarity (Supplementary Figure 2). Of note,

transcripts characteristic of late stage gametocytes appeared especially enriched in upregulated

differentially expressed genes (DEGs) in samples of asymptomatic carriers in the dry season

versus malaria cases in the wet season. We identified 159 gametocyte-specific transcripts defining

early- or late-gametocyte development stages with quantifiable reads in samples of circulating

parasites persisting through the dry season and parasites causing malaria in the wet season (18).

Of those, 105 transcripts define early-stage gametocytes; and 54 transcripts are characteristic of

late-stage gametocytes (highlighted in Supplementary Table 1). Expression of these transcripts in

asymptomatic carriers at the end of the dry season (May) and malaria cases (MAL) in the wet

season showed an enrichment of upregulated transcripts of late-stage gametocytes in the dry

season. While 81,48% (44/54 genes, Chi^2 p-value < 0.00001) of late-stage gametocyte transcripts

were upregulated in the dry season, only 39,05% (41/105 genes, Chi^2 p-value = 0.231933) of

early-gametocyte transcripts and 33,4% (1131/3381 genes) of total *P. falciparum* transcripts were

upregulated in the dry season, respectively (Figure 1B). Transcripts of female (P25,

PF3D7_1031000) or male (PFMGET, Pf3d7_1469900) late stage gametocytes were upregulated

alongside other late sexual stage transcripts (highlighted in Supplementary Table 1), and P25

difference was also confirmed by qRT-PCR through the proportional quantification of female

gametocytes out of total parasites, while PFMGET qRT-PCR showed a non-significant trend of the

proportion of males out of total parasites between clinical malaria sample and asymptomatic at the

end of the dry season (Fig. 1C). On the other hand, transcripts known to regulate sexual

commitment (AP2G, PF3D7_1222600; and GDV1, PF3D7_0935400), to be present in sexually

committed schizonts (MSRP1, PF3D7_1335000), or to be markers of early gametocytes (GEXP5,

PF3D7_0936600) were not significantly different in the RNAseq analysis (highlighted in

Supplementary Table 1) nor qRT-PCR (Figure 1C). Decreased transcription of late-stage

gametocyte genes in symptomatic cases of malaria during the wet season coincided with lower

levels of phospholipids measured through shotgun lipidomics in the plasma of the same children

that is known to influence *P. falciparum* sexual commitment in vitro (31). Malaria cases in the wet

season (MAL) had lower levels of several species of the Lyso-PC (Figure 1D) and

lysophosphatidyle-thanolamine (Lyso-PE) (Figure 1D) families compared to asymptomatic carriers

at the end of the dry season (May). Decreasing Lyso-PC levels have been shown in vitro to

increase *P. falciparum* sexual commitment (31). Hence, we questioned if the reduction in Lyso-PC

and Lyso-PE species in plasma of malaria cases compared to persistent infections at the end of

the dry season could induce *P. falciparum* sexual conversion or gametocytaemia differently in vitro.

We exposed parasites of the *P. falciparum* 2004 line (Pf2004) to plasma of children collected

during clinical malaria cases (MAL) in the wet season, or to those of asymptomatically infected

children at the end of the dry season (May) alongside the conditions described by Brancucci et al.

(31) to induce or inhibit sexual convertion and gametocyteamia. As reported earlier (31), we

observed that 24 h after culturing Pf2004 without plasma or serum (noPI), and thus no Lyso-PC,

over minimal fatty acid medium (MFA), AP2G expression increased compared to parasites cultured

in the presence of German control plasma (PIDE) (Supplementary Figure 3). However, MFA

medium supplementation with 25% plasma from children presenting with clinical malaria in the wet

season (PIMAL), versus plasma from children with asymptomatic infections at the end of the dry

season (PIMay) resulted in similar APG2 induction 48 h after beginning of treatment

(Supplementary Figure 3), and similar conversion rate of parasites into gametocytes on day 5 after

treatment (defined by = d5gametocytes/ d2parasites) (Supplementary Figure 3). Absence of

plasma (noPI) resulted in higher gametocytaemia than observed within German pooled plasma

condition (PIDE) (Supplementary Figure 3), plasma from malaria cases (PIMAL) or asymptomatic

infections at the end of the dry season (PIMay), all revealing similar conversion rates. Although the

2-4 fold Lyso-PC differences between non-infected or asymptomatic children and malaria cases

were significant (Figure 1D), these were much lower than the several orders of magnitude included

in the standard curve used to establish that gametocyte conversion responds to decreasing Lyso-

PC in the culture medium (31).

***P. falciparum* gametocyte density is higher in clinical cases of malaria than in asymptomatic**

infection regardless of seasonality

We determined the density of total *P. falciparum* parasites (Glycine-tRNA ligase,

PF3D7_1420400), female (P25, PF3D7_1031000) and male (PFMGET, Pf3D7_1469900)

gametocytes, based on standard curves obtained from total and sorted gender-specific fluorescent

gametocytes (40), respectively (Supplementary Figure 1), and detected higher numbers of both

total parasites, and of female and male gametocytes (Figure 2A) per mL of blood in clinical cases

of malaria during the wet season (MAL) compared to asymptomatic carriers at the end of the dry

season (May). This highlighted that the increased proportion of gametocytes shown by the over-

expression of transcripts of late stage gametocytes (Figure 1) was linked to lower parasite and

gametocyte densities in asymptomatic infections at the end of the dry season.

Furthermore, we determined the dynamics of *P. falciparum* sexual commitment and

gametocytogenesis in persistently infected individuals continuously infected throughout the dry

season. We quantified the density of total parasites, and fully formed female and male

gametocytes by qRT-PCR at four different time points in asymptomatic children persistently

infected from the wet (Oct) until the end of the dry season (through Jan, Mar and May), and in a

parallel group of age-matched individuals presenting with clinical malaria during the ensuing wet

season (MAL). We observed that total parasite density was higher in malaria cases (MAL) than at

any time-point of the asymptomatic infections, whether in the wet (Oct) or the dry season (Jan, Mar

and May). In contrast, no differences were found between any of the time points of asymptomatic

infections (Oct, Jan, Mar and May) (Supplementary Figure 4). Similarly, we observed higher

densities of both female and male gametocytes in malaria cases (MAL) than at any time point of

the asymptomatic infections (Oct, Jan, Mar and May), and no differences within the persisting

period (Supplementary Figure 4). Conversely, the proportion of male gametocytes was higher at

later time points of persisting infections (Mar or May) than in malaria cases during the transmission

season (MAL) (Supplementary Figure 4), while AP2G was again not found to significantly vary

across season or clinical presentation (Supplementary Figure 4). We also examined associations

between plasma Lyso-PC and Lyso-PE, monitoring longitudinally ~20 children who were

continuously infected from the wet season until the end of the dry season. Quantification of Lyso-

PC 18:0 in malaria cases compared to asymptomatic infections progressing from the wet into the

dry season again resulted in decreased Lyso-PC values in clinical malaria cases (MAL) compared

to asymptomatic infections (Oct, Mar and May) (Supplementary Figure 4). We further detected a

decrease of Lyso-PC from the wet season (Oct) to the end of the dry season (May), but no

difference was observed between the two available time points of the dry season (Mar and May)

(Supplementary Figure 4). Furthermore, Lyso-PC 18:0 levels in persistent infections (Oct, Mar and

May) were comparable to non-infected individuals at the end of the dry season (May⁻), indicating

that asymptomatic infections have no impact on Lyso-PC 18:0 (Supplementary Figure 4).

Accordingly, a relation between Lyso-PC levels and parasitaemia or gametocytaemia was only

significant when considering clinical malaria plus all asymptomatic samples (Oct, Mar, May and

MAL) (r=−0.68 p< 0.0001; r=−0.56 p< 0.008 and r=−0.67 p< 0.0001 for parasitaemia, female and

male gametocytaemia respectively), but disappeared when each group was analysed

independently, or in all persisting asymptomatic infections analysed together (Supplementary

Figure 4).

Additionally, we investigated if the observed patterns of gametocyte densities and proportions were

generalizable to non-seasonal settings where infections also persist. While Mali shows a very

contrasting seasonal pattern with 5-6 months interruption of transmission every year (17), Uganda

has uninterrupted year-round transmission with peaks in times of stronger rain (44). RT-qPCR data

quantifying transcripts of total parasite (varATS) and gametocyte (P25, PF3D7_1031000) densities

from samples of 150 Ugandan individuals with asymptomatic parasitaemias persisting for over one

month (n=122) and age-matched individuals presenting with clinical malaria (n=38) had been

previously published (44) and were retrieved from ClinEpiDB (50). Determining average parasite

levels during the asymptomatic phase calculated by the area under the curve over the course of

infection normalized by the time of infection, indicated that parasite and gametocyte densities were

increased in clinical malaria cases (MAL) compared to asymptomatic infections (Asymp) (Figure

2B). In contrast, the proportion of gametocyte transcripts relative to total *P. falciparum* transcripts

showed again an inverse trend (Figure 2C), as seen within the Malian samples, where the

proportion of gametocytes was higher in persisting parasitaemias (Asymp) compared to malaria

cases (MAL) (Figure 1).

501
Progressing infection leads to apparent differences in gametocytogenesis. We observed

decreased densities but higher proportions of female or male gametocytes in asymptomatic versus

clinical malaria, without concomitant differences in the expression of regulators of sexual

commitment, or transcriptional markers of sexually committed schizonts or early gametocytes

(Figure 1, 2, Supplementary Figure 4 and Supplementary Table 1). Attempting to explain this, we

devised a mathematical model considering three different parasite populations - asexual parasites

(P), sexually committed parasites (P$_G$) and gametocytes (G), all of which can circulate or be

adherent - as previously developed (35) (Figure 3A). As the original model by Cao et al. (35) was

not able to sufficiently explain the observed dynamics when fitted to the data, we extended the

model in different ways. First, we incorporated our recent observation of increased clearance of

longer-circulating infected erythrocytes in persisting parasitaemias compared to recently

transmitted parasites (18), which promoted different removal rates of asexual and sexually-

committed parasites during distinct phases of acute and chronic *P. falciparum* infection (see

Methods and Source code 1). In addition, we distinguish between three phases *during P.*

falciparum infection in which the parasite multiplication factor varies due to immune response or

adhesive properties of the parasite, in agreement with a recent report (44) and Malian data

(Supplementary Figure 4). Using an iterative Bayesian fitting-approach to fit the model to the

symptomatic and asymptomatic parasitaemia and gametocytaemia levels of the Malian cohort, we

quantitatively assessed several biological parameters including sexual commitment rate,

gametocyte circulation time, parasite multiplication rate and gametocyte sequestration time,

thereby confirming previously determined ranges (35, 51) (Supplementary Figure 5). Our adapted

model was able to provide a dynamic depiction of parasitaemia levels, and gametocyte densities

and proportions, as we observed within the Malian data (Figure 3B). We found that the death rate

of asexual and sexually committed parasites increased at later stages of the 48 h cycle, shifting

toward the removal of older parasite stages and leading to a constant but oscillating number of

asexual parasite in circulation (highlighted in the box-inlay of Figure 3B). This oscillation, may be

explained by the variation seen in parasitaemias, especially during the chronic phase where we

pooled individual measurements collected across different time-points and calculated averages

from each individual by the area under the curve (see Methods). Keeping a constant fraction of

sexually committed parasites, f, throughout acute and chronic infection phases, our model predicts

a higher number and lower proportions of gametocytes in the acute phase - a trend that is inverted

during the transition phase of infection due to effective removal of asexual parasites by an active

immune response (Figure 3C and D). During the chronic phase of infection, when parasitaemias

remain constant or are slowly decreasing, gametocyte proportions are also decreasing until an

equilibrium is reached roughly 4-5 months after the start of blood stage infection.

Additionally, we tested the effect of increased sexual commitment during the chronic phase of

infection in our model. We observed that increasing sexual commitment We observed that

increasing sexual commitment from 0.54-0.76% (89% credibility interval of pyABC fitting approach)

to 2 or 20% during the chronic stage of infection led to steady and constant losses in parasitaemia,

which would, especially at very high sexual commitment rates, lead to parasite clearance and

substantially increased gametocyte proportions (Supplementary Figure 6). A high fraction of

sexually committed parasites during the chronic phase of infection would prevent a stable level of

parasitaemia over several months without reinfection. Overall, our model provides a dynamic

depiction of gametocyte densities and proportions as observed in the data from Mali and Uganda,

without the necessity to alter the sexual commitment rate (Fig 3C and D).

547
Discussion

The relative absence of mosquitoes during the months-long dry season and the short lifespan of *P.*

falciparum sexual-stage gametocytes (13, 14) renders the production of gametocytes ineffective

during most of the dry season. However, to ensure parasite reproduction, gametocytes must be

available to resume malaria transmission when mosquitos return with the wet season. Whether *P.*

falciparum maintains a constant rate of production of sexual forms or adjusts gametocytogenesis to

maximize transmission has been unclear. In this study, with samples from Malians exposed to

alternating 6-month dry and wet seasons, and samples from clinical malaria and asymptomatic *P.*

falciparum infection from year-round transmission region of Uganda, we showed that symptomatic

cases of malaria consistently have higher parasite and stage V gametocyte densities than

asymptomatic infections. However, the proportion of gametocytes is increased in the asymptomatic

persisting infections. Additionally, our data from Mali indicated that transcription of genes

characterizing sexually committed parasites and stage I gametocytes (52, 53) quantified by RT-

qPCR did not appear affected by the season or clinical symptoms (Figures 1 and 2), nor were

reads of over 100 genes characteristic of early gametocytes and detected by RNAseq of circulating

parasites particularly enriched for DEGs. This, clearly contrast with representative expressed

genes of late gametocytes that were very much enriched of transcripts upregulated in persisting

asymptomatic infections at the end of the dry season (Fig 1C and Supplementary Table 1).

Our work suggests that gametocytogenesis remains constant as infections progress from recent

transmission to chronic carriage, and it is the difference in asexual survival/replication rates that

promotes different gametocyte densities, independently of environmental factors.

Although it has been suggested that *P. falciparum* investment in gametocyte production adjusts to

host nutrient availability (54), drug pressure (24), transmission intensity (55), and even seasonality

(56, 57), results are often inconsistent and potentially affected by the dynamics of asexual parasite

growth, without necessarily indicating altered sexual commitment. In Sudan, following the long dry

season, gametocyte density estimates have been reported to increase prior to the highest

transmission peak (58), suggesting a higher gametocyte investment. However, in Burkina Faso,

gametocyte densities were higher at the peak and end of the transmission season rather than

during the dry season (59), while the efficiency of membrane feeding to infect mosquitoes was

increased early in the transmission season compared to peak of the malaria season or the dry

season (60). Nevertheless, these studies did not monitor longitudinally the same parasite

population during the dry and early wet season. In a recent study also in Burkina Faso,

transmission was shown to be more effective in persistent than in incident infections, but no data

was collected after day 35, and collections were all done during the wet season (25) when a mix of

more and less recently transmitted parasite clones is likely in individuals without symptoms.

In vitro, depletion of Lyso-PC or choline have been shown to induce AP2G expression, leading to

increased gametocyte production (31, 61), but associations in humans are inconsistent. In

Rwanda, low Lyso-PC associated with higher parasite densities (62); and in Ghana, although

correlations with parasitaemia were not reported, Lyso-PC levels in human blood samples were

inconsistently related with gametocyte-committed ring stages converting into gametocytes in vitro

(53). Our data show, like Orikiiriza et al.(62), an association of lower Lyso-PC levels with increased

parasitaemia and severity of malaria (Figure 1), but the association was lost at low parasitaemias,

and persistent infection at the end of the dry season did not affect Lyso-PC compared to non-

infected individuals (Supplementary Figure 4).

The effect of drugs on sexual conversion in vitro has been previously reported. Specifically, sub-

curative doses of dihydroartemisinin (DHA) stimulate AP2G expression and sexual conversion

when drug exposure occurs at the trophozoite stage, but not at the early ring stage. The observed

increase could not synergize nor add to increases promoted by depletion of choline, suggesting, as

DHA affects lipids, that the two effects might be the same (61). In vivo, it was recently shown that

soon after ART treatment AP2G transcripts increased in many but not all patients, and data

contrasted significantly between cohorts of different geographical regions, highlighting the complex

regulation of sexual conversion which is likely influenced by patient, parasite characteristics and

resistance status (63). Our RNAseq (S1 Table) and RT-qPCR (Fig 1H and 2D) data show constant

low-level expression of AP2G, the master regulator of gametocytogenesis, and equally unaltered

expression of transcripts characteristic of committed schizont and early gametocytes (S1 Table and

Fig 1H), suggesting that reductions in circulating Lyso-PC induced by uncomplicated clinical

malaria are not capable of affecting *P. falciparum* sexual commitment. Accordingly, we observed in

vitro that supplementation with plasma from clinical malaria cases (low Lyso-PC) or end of the dry

season persistently infected individuals (normal Lyso-PC) did not differentially induce AP2G

expression or the rate of sexual conversion (Fig 3), suggesting that Lyso-PC differences introduced

by *P. falciparum* infection are not sufficient to trigger the mechanisms demonstrated by Brancucci

et al. (31) and Portugaliza et al. (61). Nevertheless, it is possible that Lyso-PC levels in particular

bone marrow niches are sufficiently low to trigger parasites' sexual commitment, although this

would require that infected erythrocytes be at that niche to respond to the stimulus.

Previous reports highlighting differences in gametocytogenesis as a product of altered sexual

commitment did not account for longevity of infection, nor for dynamic changes within the asexual

compartment during a persistent infection. Greischar et al. demonstrated the difficulty of quantifying

true transmission investment by comparing existing quantification methods on simulated data

where true transmission investment is known (64), and Prajapati and colleagues relied on in vitro

culture of sexually committed circulating parasites to define samples with higher and lower

commitment rates (65). Also, our study has limitations as it is possible that our quantifications of

parasite and gametocyte densities are affected by imperfectly comparable primer efficiencies,

which would also affect comparisons of female, male and total gametocyte densities across

different transcripts measurements. Although we also do not have true longitudinal data, our model

can explain the observed patterns by assuming transitions from a faster to a slower growth rate, as

e.g. due to the development of immunity which has been recorded from malaria-therapy used to

treat neurosyphilis patients (66). Despite these limitations, our analyses show that the proportion of

sexual stages is inversely related in recent transmissions versus chronic infections, without

alteration in the expression of regulators of sexual commitment and while maintaining a constant

rate of sexual commitment. Our recent study in which we found that infections that persist beyond

the acute immune response exhibit prolonged circulation and increased splenic clearance (18)

allowed us to complement the model by Cao et al. (35) with an increased asexual parasite death

rate following the immuno-responsive period, which promoted the progressively increased

representation of sexual stages in later phases of the infection. Low gametocyte proportions early

on in infection can be explained by the fact that asexual replication efficiently increases while

adaptive immunity is absent and the lag of ~8 days of gametocyte development in the bone

marrow has its impact. In our model we propose that later immunity prevents fast asexual growth

and parasites persisting further are more efficiently removed in the spleen sustaining low

parasitaemias. By introducing these concepts into our model, we can observe the opposite lag

effect later in infection when the proportion of gametocytes finally reflects, despite keeping the

same sexual commitment rate, that of parasitaemias ~8 days earlier. In contrast, assuming

increased fractions of sexually committed parasites during the chronic phase of infection, our

model would predict a constant loss of parasitemia and eventual clearance of infection.

In conclusion, our data is consistent with a model in which *P. falciparum* maintains a constant rate

of sexual commitment such that the proportion of gametocytes increases over time as infection

persists and asexual parasite numbers decline more rapidly, a strategy that may help overcome

the vector-free period and increase transmission efficiency.

Acknowledgments

We thank the residents of Kalifabougou, Mali and Nangonera, Uganda for participating in the

studies. We acknowledge the support of the Flow Cytometry Core Facility of DKFZ, the Heidelberg

University Metabolomics Core Technology Platform, and the Clinical Lipidomics Unit of Mainz

University Medical Center. We thank Matthias Marti and Lauriane Sollelis at University of Glasgow

for help and expertise with in vitro induction of parasite sexual commitment and

gametocytogenesis. This work was supported by the European Research Council (ERC) under the

European Union's Horizon 2020 research and innovation programme (grant agreement No

759534), and the German Center for Infection research (DZIF). The Malian cohort study was

supported by the Division of Intramural Research, National Institute of Allergy and Infectious

Diseases of the National Institutes of Health; and the Ugandan study was supported by the

National Institutes of Health as part of the International Centers of Excellence in Malaria Research

program. Chica and Heinz-Schaller Foundation supported Frederik Graw.

Competing Interest Statement

The authors declare that they have no conflict of interest.

References

1.   Grassi B, Bignami A, Bastianelli G. Medical Zoology: Further Researches upon the Cycle of
Human Malaria in the Body of the Mosquito. Ind Med Gaz. 1899;34(3):104-7.
2.   Prudencio M, Rodriguez A, Mota MM. The silent path to thousands of merozoites: the
Plasmodium liver stage. Nat Rev Microbiol. 2006;4(11):849-56.
3.   Haldar K, Murphy SC, Milner DA, Taylor TE. Malaria: mechanisms of erythrocytic infection and
pathological correlates of severe disease. Annu Rev Pathol. 2007;2:217-49.
4.   Kafsack BF, Rovira-Graells N, Clark TG, Bancells C, Crowley VM, Campino SG, et al. A
transcriptional switch underlies commitment to sexual development in malaria parasites. Nature.
2014;507(7491):248-52.
5.   Arnot DE, Gull K. The Plasmodium cell-cycle: facts and questions. Ann Trop Med Parasitol.
1998;92(4):361-5.
6.   Bruce MC, Alano P, Duthie S, Carter R. Commitment of the malaria parasite Plasmodium
falciparum to sexual and asexual development. Parasitology. 1990;100 Pt 2:191-200.

7.   Inselburg J. Gametocyte formation by the progeny of single Plasmodium falciparum schizonts. J
Parasitol. 1983;69(3):584-91.
8.   Hawking F, Wilson ME, Gammage K. Evidence for cyclic development and short-lived maturity
in the gametocytes of Plasmodium falciparum. Trans R Soc Trop Med Hyg. 1971;65(5):549-59.
9.   Li J, Shami GJ, Cho E, Liu B, Hanssen E, Dixon MWA, et al. Repurposing the mitotic machinery
to drive cellular elongation and chromatin reorganisation in Plasmodium falciparum gametocytes.
Nature Communications. 2022;13(1).
10.   Parkyn Schneider M, Liu B, Glock P, Suttie A, McHugh E, Andrew D, et al. Disrupting assembly
of the inner membrane complex blocks Plasmodium falciparum sexual stage development. PLoS
Pathog. 2017;13(10):e1006659.
11.   Farfour E, Charlotte F, Settegrana C, Miyara M, Buffet P. The extravascular compartment of the
bone marrow: a niche for Plasmodium falciparum gametocyte maturation? Malar J. 2012;11:285.
12.   Tiburcio M, Silvestrini F, Bertuccini L, Sander AF, Turner L, Lavstsen T, et al. Early
gametocytes of the malaria parasite Plasmodium falciparum specifically remodel the adhesive
properties of infected erythrocyte surface. Cell Microbiol. 2013;15(4):647-59.
13.   Smalley ME, Sinden RE. Plasmodium falciparum gametocytes: their longevity and infectivity.
Parasitology. 1977;74(1):1-8.
14.   Gebru T, Lalremruata A, Kremsner PG, Mordmuller B, Held J. Life-span of in vitro differentiated
Plasmodium falciparum gametocytes. Malar J. 2017;16(1):330.
15.   Billker O, Lindo V, Panico M, Etienne AE, Paxton T, Dell A, et al. Identification of xanthurenic
acid as the putative inducer of malaria development in the mosquito. Nature. 1998;392(6673):289-92.
16.   Ross R. The role of the mosquito in the evolution of the malarial parasite: the recent researches
of Surgeon-Major Ronald Ross, I.M.S. 1898. Yale J Biol Med. 2002;75(2):103-5.
17.   Dicko A, Klion AD, Thera MA, Sagara I, Yalcouye D, Niambele MB, et al. The etiology of severe
anemia in a village and a periurban area in Mali. Blood. 2004;104(4):1198-200.
18.   Andrade CM, Fleckenstein H, Thomson-Luque R, Doumbo S, Lima NF, Anderson C, et al.
Increased circulation time of Plasmodium falciparum underlies persistent asymptomatic infection in the
dry season. Nat Med. 2020.
19.   Babiker HA, Abdel-Muhsin AM, Ranford-Cartwright LC, Satti G, Walliker D. Characteristics of
Plasmodium falciparum parasites that survive the lengthy dry season in eastern Sudan where malaria
transmission is markedly seasonal. Am J Trop Med Hyg. 1998;59(4):582-90.
20.   Portugal S, Tran TM, Ongoiba A, Bathily A, Li S, Doumbo S, et al. Treatment of Chronic
Asymptomatic Plasmodium falciparum Infection Does Not Increase the Risk of Clinical Malaria Upon
Reinfection. Clin Infect Dis. 2017;64(5):645-53.
21.   Adomako-Ankomah Y, Chenoweth MS, Tocker AM, Doumbia S, Konate D, Doumbouya M, et
al. Host age and Plasmodium falciparum multiclonality are associated with gametocyte prevalence: a 1-
year prospective cohort study. Malar J. 2017;16(1):473.
22.   Drakeley CJ, Secka I, Correa S, Greenwood BM, Targett GA. Host haematological factors
influencing the transmission of Plasmodium falciparum gametocytes to Anopheles gambiae s.s.
mosquitoes. Trop Med Int Health. 1999;4(2):131-8.
23.   Gouagna LC, Bancone G, Yao F, Yameogo B, Dabire KR, Costantini C, et al. Genetic variation
in human HBB is associated with Plasmodium falciparum transmission. Nat Genet. 2010;42(4):328-31.
24.   Price R, Nosten F, Simpson JA, Luxemburger C, Phaipun L, ter Kuile F, et al. Risk factors for
gametocyte carriage in uncomplicated falciparum malaria. Am J Trop Med Hyg. 1999;60(6):1019-23.
25.   Barry A, Bradley J, Stone W, Guelbeogo MW, Lanke K, Ouedraogo A, et al. Higher gametocyte
production and mosquito infectivity in chronic compared to incident Plasmodium falciparum infections.
Nat Commun. 2021;12(1):2443.
26.   Group WGS. Gametocyte carriage in uncomplicated Plasmodium falciparum malaria following
treatment with artemisinin combination therapy: a systematic review and meta-analysis of individual
patient data. BMC Med. 2016;14:79.
27.   Rek J, Blanken SL, Okoth J, Ayo D, Onyige I, Musasizi E, et al. Asymptomatic school-aged
children are important drivers of malaria transmission in a high endemicity setting in Uganda. J Infect
Dis. 2022.
28.   Tadesse FG, Slater HC, Chali W, Teelen K, Lanke K, Belachew M, et al. The Relative
Contribution of Symptomatic and Asymptomatic Plasmodium vivax and Plasmodium falciparum
Infections to the Infectious Reservoir in a Low-Endemic Setting in Ethiopia. Clin Infect Dis.
2018;66(12):1883-91.

29.   Williams JL. Stimulation of Plasmodium falciparum gametocytogenesis by conditioned medium
from parasite cultures. Am J Trop Med Hyg. 1999;60(1):7-13.

30.   Dyer M, Day KP. Regulation of the rate of asexual growth and commitment to sexual
development by diffusible factors from in vitro cultures of Plasmodium falciparum. Am J Trop Med Hyg.
2003;68(4):403-9.

31.   Brancucci NMB, Gerdt JP, Wang C, De Niz M, Philip N, Adapa SR, et al.
Lysophosphatidylcholine Regulates Sexual Stage Differentiation in the Human Malaria Parasite
Plasmodium falciparum. Cell. 2017;171(7):1532-44 e15.

32.   Filarsky M, Fraschka SA, Niederwieser I, Brancucci NMB, Carrington E, Carrio E, et al. GDV1
induces sexual commitment of malaria parasites by antagonizing HP1-dependent gene silencing.
Science. 2018;359(6381):1259-63.

33.   Campelo Morillo RA, Tong X, Xie W, Abel S, Orchard LM, Daher W, et al. The transcriptional
regulator HDP1 controls expansion of the inner membrane complex during early sexual differentiation of
malaria parasites. Nat Microbiol. 2022;7(2):289-99.

34.   Thomson-Luque R, Votborg-Novel L, Ndovie W, Andrade CM, Niangaly M, Attipa C, et al.
Plasmodium falciparum transcription in different clinical presentations of malaria associates with
circulation time of infected erythrocytes. Nat Commun. 2021;12(1):4711.

35.   Cao P, Collins KA, Zaloumis S, Wattanakul T, Tarning J, Simpson JA, et al. Modeling the
dynamics of Plasmodium falciparum gametocytes in humans during malaria infection. Elife. 2019;8.

36.   Painter HJ, Carrasquilla M, Llinas M. Capturing in vivo RNA transcriptional dynamics from the
malaria parasite Plasmodium falciparum. Genome Res. 2017;27(6):1074-86.

37.   Poran A, Notzel C, Aly O, Mencia-Trinchant N, Harris CT, Guzman ML, et al. Single-cell RNA
sequencing reveals a signature of sexual commitment in malaria parasites. Nature. 2017;551(7678):95-
9.

38.   Lopez-Barragan MJ, Lemieux J, Quinones M, Williamson KC, Molina-Cruz A, Cui K, et al.
Directional gene expression and antisense transcripts in sexual and asexual stages of Plasmodium
falciparum. BMC Genomics. 2011;12:587.

39.   Tran TM, Li S, Doumbo S, Doumtabe D, Huang CY, Dia S, et al. An intensive longitudinal
cohort study of Malian children and adults reveals no evidence of acquired immunity to Plasmodium
falciparum infection. Clin Infect Dis. 2013;57(1):40-7.

40.   Meerstein-Kessel L, Andolina C, Carrio E, Mahamar A, Sawa P, Diawara H, et al. A multiplex
assay for the sensitive detection and quantification of male and female Plasmodium falciparum
gametocytes. Malar J. 2018;17(1):441.

41.   Lasonder E, Rijpma SR, van Schaijk BC, Hoeijmakers WA, Kensche PR, Gresnigt MS, et al.
Integrated transcriptomic and proteomic analyses of P. falciparum gametocytes: molecular insight into
sex-specific processes and translational repression. Nucleic Acids Res. 2016;44(13):6087-101.

42.   Surma MA, Herzog R, Vasilj A, Klose C, Christinat N, Morin-Rivron D, et al. An automated
shotgun lipidomics platform for high throughput, comprehensive, and quantitative analysis of blood
plasma intact lipids. Eur J Lipid Sci Technol. 2015;117(10):1540-9.

43.   Lerner R, Post J, Loch S, Lutz B, Bindila L. Targeting brain and peripheral plasticity of the
lipidome in acute kainic acid-induced epileptic seizures in mice via quantitative mass spectrometry.
Biochim Biophys Acta Mol Cell Biol Lipids. 2017;1862(2):255-67.

44.   Andolina C, Rek JC, Briggs J, Okoth J, Musiime A, Ramjith J, et al. Sources of persistent
malaria transmission in a setting with effective malaria control in eastern Uganda: a longitudinal,
observational cohort study. Lancet Infect Dis. 2021.

45.   Briggs J, Teyssier N, Nankabirwa JI, Rek J, Jagannathan P, Arinaitwe E, et al. Sex-based
differences in clearance of chronic Plasmodium falciparum infection. Elife. 2020;9.

46.   Hofmann N, Mwingira F, Shekalaghe S, Robinson LJ, Mueller I, Felger I. Ultra-sensitive
detection of Plasmodium falciparum by amplification of multi-copy subtelomeric targets. PLoS Med.
2015;12(3):e1001788.

47.   Tran TM, Guha R, Portugal S, Skinner J, Ongoiba A, Bhardwaj J, et al. A Molecular Signature in
Blood Reveals a Role for p53 in Regulating Malaria-Induced Inflammation. Immunity. 2019;51(4):750-65
e10.

48.   Miller LH, Good MF, Milon G. Malaria pathogenesis. Science. 1994;264(5167):1878-83.

49.   Klinger E, Rickert D, Hasenauer J. pyABC: distributed, likelihood-free inference. Bioinformatics.
2018;34(20):3591-3.

50.   Ruhamyankaka E, Brunk BP, Dorsey G, Harb OS, Helb DA, Judkins J, et al. ClinEpiDB: an
open-access clinical epidemiology database resource encouraging online exploration of complex
studies. Gates Open Res. 2019;3:1661.
51.   Eichner M, Diebner HH, Molineaux L, Collins WE, Jeffery GM, Dietz K. Genesis, sequestration
and survival of Plasmodium falciparum gametocytes: parameter estimates from fitting a model to
malariatherapy data. Trans R Soc Trop Med Hyg. 2001;95(5):497-501.
52.   Tiburcio M, Dixon MW, Looker O, Younis SY, Tilley L, Alano P. Specific expression and export
of the Plasmodium falciparum Gametocyte EXported Protein-5 marks the gametocyte ring stage. Malar
J. 2015;14:334.
53.   Usui M, Prajapati SK, Ayanful-Torgby R, Acquah FK, Cudjoe E, Kakaney C, et al. Plasmodium
falciparum sexual differentiation in malaria patients is associated with host factors and GDV1-dependent
genes (vol 10, 2140, 2019). Nature Communications. 2019;10.
54.   Shanks GD. Hypothesis: dynamics of classical malaria epidemics show Plasmodium
falciparum's survival strategy. Am J Trop Med Hyg. 2015;92(3):561-4.
55.   Rono MK, Nyonda MA, Simam JJ, Ngoi JM, Mok S, Kortok MM, et al. Adaptation of
Plasmodium falciparum to its transmission environment. Nat Ecol Evol. 2018;2(2):377-87.
56.   Oduma CO, Ogolla S, Atieli H, Ondigo BN, Lee MC, Githeko AK, et al. Increased investment in
gametocytes in asymptomatic Plasmodium falciparum infections in the wet season. BMC Infect Dis.
2021;21(1):44.
57.   Rosenberg R, Andre RG, Ketrangsee S. Seasonal fluctuation of Plasmodium falciparum
gametocytaemia. Trans R Soc Trop Med Hyg. 1990;84(1):29-33.
58.   Gadalla AAH, Schneider P, Churcher TS, Nassir E, Abdel-Muhsin AMA, Ranford-Cartwright LC,
et al. Associations between Season and Gametocyte Dynamics in Chronic Plasmodium falciparum
Infections. Plos One. 2016;11(11).
59.   Ouedraogo AL, de Vlas SJ, Nebie I, Ilboudo-Sanogo E, Bousema JT, Ouattara AS, et al.
Seasonal patterns of Plasmodium falciparum gametocyte prevalence and density in a rural population of
Burkina Faso. Acta Trop. 2008;105(1):28-34.
60.   Ouedraogo AL, Goncalves BP, Gneme A, Wenger EA, Guelbeogo MW, Ouedraogo A, et al.
Dynamics of the Human Infectious Reservoir for Malaria Determined by Mosquito Feeding Assays and
Ultrasensitive Malaria Diagnosis in Burkina Faso. J Infect Dis. 2016;213(1):90-9.
61.   Portugaliza HP, Miyazaki S, Geurten FJ, Pell C, Rosanas-Urgell A, Janse CJ, et al. Artemisinin
exposure at the ring or trophozoite stage impacts Plasmodium falciparum sexual conversion differently.
Elife. 2020;9.
62.   Orikiiriza J, Surowiec I, Lindquist E, Bonde M, Magambo J, Muhinda C, et al. Lipid response
patterns in acute phase paediatric Plasmodium falciparum malaria. Metabolomics. 2017;13(4):41.
63.   Portugaliza HP, Natama HM, Guetens P, Rovira-Vallbona E, Some AM, Millogo A, et al.
Plasmodium falciparum sexual conversion rates can be affected by artemisinin-based treatment in
naturally infected malaria patients. EBioMedicine. 2022;83:104198.
64.   Greischar MA, Mideo N, Read AF, Bjornstad ON. Quantifying Transmission Investment in
Malaria Parasites. PLoS Comput Biol. 2016;12(2):e1004718.
65.   Prajapati SK, Ayanful-Torgby R, Pava Z, Barbeau MC, Acquah FK, Cudjoe E, et al. The
transcriptome of circulating sexually committed Plasmodium falciparum ring stage parasites forecasts
malaria transmission potential. Nat Commun. 2020;11(1):6159.
66.   Collins WE, Jeffery GM. A retrospective examination of sporozoite- and trophozoite-induced
infections with Plasmodium falciparum: development of parasitologic and clinical immunity during
primary infection. Am J Trop Med Hyg. 1999;61(1 Suppl):4-19.

A

B

C

D

Supplementary Figure 2

A

B

	Acute Phase	Transition phase	Chronic phase
Sexual commitment rate	0.54 - 0.76 %		
Gametocyte circulation time	12.04 - 22.28 days		
Parasite multiplication factor / 48 h	3.87 - 8.18	0.33 - 1.38	0.50 - 2.29
Gametocyte sequestration time	3.89 - 6.04 days		

A

B

Figure legends

Figure 1. *P. falciparum* transcription and quantification of gametocyte density and

proportions. A, Principal components analysis of RNA-Seq data of 333 gametocyte specific *P.*

falciparum genes from parasites collected at the end of the dry season and from clinical malaria

cases (n= 12 May, 12 MAL). **B**, Fold change between parasites collected at the end of the dry

season and from clinical malaria cases (n= 12 May, 12MAL) of early and late gametocyte

transcripts according to López-Barragán et al., 2011; Poran et al., 2017 and Petter unpublished

data. Each dot represents a gene and color identifies range in fold change, red: upregulated in

May, yellow: not DEG, and blue: downregulated in May). **C,** Proportion of female ($\female$) and male ($\male$)

gametocytes, relative expression of AP2G, GDV1, MSRP1, and GEXP5, detected by qRT-PCR of

RDT+ subclinical children at the end of the dry season (May n > 15) and of children with their first

clinical malaria episode (MAL n > 23) in the wet season. **D,** Plasma concentration of Lyso-PC (left)

and Lyso-PE (right) species of RDT$^+$ subclinical children at the end of the dry season (May) and of

children with their first clinical malaria episode (MAL) in the wet season. Lipid data represented as

median ± IQR; two-way ANOVA with Tukey multiple comparisons correction. Parasitaemia and

gametocytemia and AP2G, GDV1, MSRP1, and GEXP5 expression data represented as median ±

IQR; Mann-Whitney test.

Figure 2. *P. falciparum* gametocyte density is higher clinical cases of malaria than in

asymptomatic infection regardless of seasonality. A: Asexual parasite (left), Female (center)

and male (right) density detected by qRT-PCR of RDT$^+$ subclinical children at the end of the dry

season (May = 35) and of children with their first clinical malaria episode (MAL = 27) in the wet

season. D: gametocyte **B**, Parasite (left) and gametocyte (right) densities detected by qRT-PCR of

symptomatic (MAL n = 19 and 38 in Mali and Uganda respectively) and asymptomatic infections

(May n = 23 and Asymp n = 122) in Mali and Uganda. **C**, Gametocyte proportion detected by qRT-

PCR of symptomatic (MAL n = 19 and 38 in Mali and Uganda respectively) and asymptomatic

infections (May n = 23 and Asymp n = 122) in Mali and Uganda. Data represented as median ±

IQR; multiple non-parametric test with Tukey-correction for multiple comparisons.

Fig 3 | Progressing infection leads to apparent differences in gametocytogenesis. A,

Diagram of the mathematical model to describe parasite dynamics including individual

compartments, transitions and death rates; Asexual parasites (P) replicate in a 48 h cycle, being

cleared at an age dependent death rate (σ_a). A fraction of asexual parasites (f) transitions to the

sexually committed parasite compartment (P_G), goes through a round of replication and develops

into gametocytes (G). Gametocytes mature over 5 distinct stages in independent model

compartments. Asexual parasites (blue), sexually committed parasites (orange) and gametocytes

(grey) are indicated shaded if not circulating due to sequestration. **B**, Model prediction and clinical

data of asexual parasites. The small inlay indicates a zoom-in showing the oscillating parasite

dynamics within 48 h-intervals. **C**, Gametocytes density and **D**, proportion of gametocytes per mL

blood. Individual data points and median (black square) of wet (aqua circles) and dry (orange

circles) season data throughout acute (aqua shaded area) and chronic (salmon shaded area)

phase of infection are shown. Median fit (grey line) and 95% prediction interval of 100 model

predictions of the 6[th] PyABC generation are shown as grey shaded area.

Figure supplement legends

Supplementary Figure 1

A, Flow cytometry gating strategy to identify and sort PfDynGFP/ PfP47 gametocytes by mCherry

and GFP expression into three populations (female: mCherry-positive, male I: mCherry- GFP-

positive, male II: GFP-positive). **B,** Pre-sort gametocyte distribution, and post-sorting purity

verification of female and male gametocytes. **C,** Serial dilutions of P25, PfMGET, AP2g, and

Glycine-tRNA ligase synthetic DNA to determine curves' slopes and primer efficiency by qPCR. **D,**

Standard curves determined with serial dilutions of sorted-purified female and male gametocytes,

showing expression of female marker P25 (left) and male marker PfMGET (right).

Supplementary Figure 2

Heatmap showing normalized reads of 456 gametocyte specific genes of *P. falciparum* (rows) for

each subject (columns) collected at the end of the dry season (May) and at the first clinical malaria

case (MAL) in the ensuing transmission season.

Supplementary Figure 3

A, Relative expression of AP2G detected by qRT-PCR of Pf2004 24 h after 24 h culture with 25%

plasma collected during clinical malaria episodes (MAL n = 21), or at the end of the dry season

(May n = 19). **B,** Pf2004 conversion rate (gametocytes d5/parasites d2) (left) and gametocyte

density (right) detected by microscopy, after 24 h culture with 25% plasma collected during clinical

malaria episodes in the wet season (MAL n = 19) or at the end of the dry season (May n = 18),

alongside no plasma (noPl) and German (PIDE) conditions. Data represented as median ± IQR;

Mann-Whitney test.

Supplementary Figure 4

A, Density of total parasites (left), Female (centre), and male (right) gametocytes detected by qRT-

PCR of RDT[+] subclinical children at different times during the wet (Oct n = 10) and the dry season

(Jan, Mar and May n > 15), and during clinical malaria episodes (MAL n = 19). **B.** Proportion of

female (left) and male (centre) gametocytes, and relative expression of AP2G (right) determined by

qRT-PCR of RDT[+] subclinical children at different times during the wet (Oct n = 6) and the dry

season (Jan, Mar and May n > 11), and during clinical malaria episodes (MAL n = 19). **D.** Plasma

concentration of Lyso-PC 18:0 of RDT[+] subclinical children at different times during the wet (Oct n =

17) and the dry season (Mar and May n = 19), and during clinical malaria episodes (MAL n = 15).

Parasite gene expression and lipid data were compared through Kruskal-Wallis test with multiple

comparisons. **D**, Correlation between Lyso-PC 18:0 and parasite (left), female gametocyte (centre)

or male gametocyte (right) densities detected by qRT-PCR detected by qRT-PCR of RDT[+]

subclinical children at different times during the wet (Oct n = 10) and the dry season (Mar and May

n > 15), and during clinical malaria episodes (MAL n = 19). *P* and r determined by Pearson

correlation

Supplementary Figure 5

A, Posterior distribution of 12 model parameters generated of 100 fits of 6th pyABC generation of

one representative pyABC fitting run. **B,** Estimates of biological parameters in distinct phases of P.

falciparum infection shown as 89% credibility interval of the posterior distributions. Sexual

commitment rate was calculated by fx100%. The parasite multiplication factor is calculated by $\exp(-$

$a_L)(1-f)$ with the term $\exp(-a_L)(1-f)$ describing the fraction of surviving asexual parasite after death

and sexual conversion per replication cycle. Gametocyte sequestration time was calculated by 4/m

in relation to the 4 sequestered states (stage I to IV) (Cao et al. 2019). While calculations of the

model where done in 1 h increments, all estimates shown in S3 Figure. B are converted to per day.

Supplementary Figure 6

Model prediction and clinical data of asexual parasites and gametocytes with variable sexual

commitment rate during asymptomatic phase of infection. Asexual parasites, oscillating parasite

dynamics within 48h interval, gametocytes and the proportion of gametocytes per mL blood are

shown for sexual commitment (f) 2% in **A** and 20% in **B** during chronic phase of infection.

Individual data points and median (black square) of wet (aqua circles) and dry (orange circles)

season data throughout acute (aqua shaded area) and chronic (salmon shaded area) phase of

infection are shown. Median fit (grey line) and 95% prediction interval of 100 model predictions of

the 6[th] PyABC generation are shown as grey shaded area.

Table supplement legends

S1 Table. Gene Ids and descriptions of transcripts defined as gametocyte-specific by Painter et al.,

2017; transcripts detected in RNA sequencing dataset (Andrade et al., 2020) comparing persistent

P. falciparum in at the end of the dry season (May) to *P. falciparum* of clinical malaria in the wet

season (MAL); and transcripts defining early or late gametocyte development gathered from Poran

et al., 2017 after request from the corresponding author, López-Barragán et al., 2011, and Jeninga

et al., unpublished. (pink and blue shaded rows highlight transcripts up or down regulated in May

vs. MAL, respectively.)

S2 Table. Key Resources Table