Constant rate of *Plasmodium falciparum* sexual commitment promotes variable gametocyte proportions as infection progresses over time

Hannah van Dijk[1*], Nathalia F Lima[1*], Martin Kampmann[2], Michael Gabel[3], Usama Dabbas[1,2], Safiatou Doumbo[4], Hamidou Cisse[4], Shanping Li[5], Myriam Jeninga[6], Richard Thomson-Luque[1], Didier Doumtabe[4], Michaela Petter[6], Kassoum Kayentao[4], Aissata Ongoiba[4], Teun Bousema[7], Peter D Crompton[5], Boubacar Traore[4], Frederik Graw[3], Silvia Portugal[1,2†]

[1] Center for Infectious Diseases, Parasitology, Heidelberg University Hospital, Heidelberg, Germany
[2] Max Planck Institute for Infection Biology, Berlin, Germany
[3] BioQuant-Center for Quantitative Biology, Heidelberg University, Heidelberg, Germany
[4] Mali International Center of Excellence in Research, University of Sciences, Techniques and Technologies of Bamako, Mali
[5] Laboratory of Immunogenetics, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Rockville, Maryland, USA
[6] Mikrobiologisches Institut-Klinische Mikrobiologie, Universitätsklinikum Erlangen
[7] Radboud University Medical Center, Nijmegen, the Netherlands
*These authors contributed equally to this work
†Correspondence: Silvia Portugal portugal@mpiib-berlin.mpg.de

Abstract

In areas of seasonal malaria, resuming transmission every wet season relies on persistent asexual parasites during dry season that maintain the ability to produce gametocytes until the mosquito population resurges, following the start of annual rains. Although human asymptomatic *P. falciparum* reservoirs in the dry season are widely recognized, the longitudinal dynamics of parasite sexual commitment and gametocytogenesis are unclear. Here, we compared the density and proportion of *P. falciparum* gametocytes in blood of asymptomatic subjects during the dry season versus subjects with symptomatic malaria in the wet season. Blood concentrations of phospholipids potentially involved in gametocytogenesis and parasite transcriptomes were also compared. While blood densities of gametocytes and asexual parasites were lower during the dry season, we found that gametocytes were proportionally higher during the dry versus wet season. Levels of parasite transcripts involved in sexual commitment were similar throughout the year, and phospholipid content in the plasma throughout the year did not affect sexual commitment in vitro. We demonstrate experimentally and through mathematical modelling that gametocyte density and proportion diverge as infections progress from recently transmitted to chronic carriage, without significant alterations in the rate of sexual commitment over time.

Introduction

It has long been known that the malaria-causing parasite *Plasmodium falciparum* requires a mosquito host to complete its lifecycle and to be transmitted between humans (1). However, it remains unclear if in the human host the parasite can dynamically allocate resources to the differentiation of sexual-stage gametocytes that are transmitted to mosquitos, particularly in areas of seasonal malaria where the size of the mosquito population fluctuates dramatically during the dry and wet seasons. During blood feedings, infected Anopheles mosquitoes deposit *P. falciparum* sporozoites in the human dermis that find their way into circulation, reaching the liver where they infect hepatocytes. After 7-10 days of parasite replication, each infected hepatocyte releases thousands of merozoites into the blood stream (2), where they start continuous asexual intra-erythrocytic replication cycles of ~48 h, from merozoite invasion of erythrocytes, through the ring- and trophozoite-stages, until the multinucleated schizont releases new merozoites (3). A small proportion of intra-erythrocytic asexual parasites differentiate into sexual-stage gametocytes (4). Instead of another ~48 h asexual schizogony leading to the 8-30 new merozoites ready to continue asexual cycles (5), the parasite develops into sexually committed ring-stages (6). Through another round of replication, female or male sexually committed parasites will produce 8-30 merozoites that upon infection of new red blood cells (RBCs) each produce one stage I gametocyte that through a period of 8-12 days, will go through five developmental stages (7). Stages II to IV of gametocytaemia take place in the bone marrow or spleen (8, 9), while stage V gametocytes are again present in peripheral blood, each lasting up to ~20 days (10, 11) waiting for another mosquito bite. Following a blood meal gametocytes encounter the right environment for sexual reproduction in the mosquito midgut (12). They transform into female and male gametes that fuse into a zygote originating oocysts that invade the midgut wall to generate sporozoites that can, after reaching the mosquito salivary glands, restart the parasite lifecycle in the next blood meal (13). The continued presence of mosquitoes and humans in a geographical region allows for perennial *P. falciparum* transmission; however, in areas where the dry season is several months long, the mosquito population can plummet to near zero (14), thereby interrupting transmission.

P. falciparum faces the added challenge of surviving in its human host for months while

waiting for the return of mosquitoes with the ensuing wet season (15-17). During

mosquito-free dry periods gametocyte production is largely wasted, as gametocytes

cannot survive to bridge two wet seasons (10, 11). Although curtailing gametocyte

production rates during the dry season would seem beneficial to the parasite, such a

strategy would require a mechanism to reactivate gametocytogenesis with the return of

rain and mosquitos. Host factors such as age (18), haematocrit (19), haemoglobin type

(20) and duration of malaria symptoms (21) have been linked to gametocytaemia in

individuals with clinical malaria; and higher gametocyte densities have been observed in

chronic asymptomatic infections compared to acute infections (22-24). Additionally,

various in vitro conditions have been shown to modulate sexual commitment and thus

gametocytogenesis (6, 25-27). The specific effect of lysophosphatidylcholine (Lyso-PC) in

repressing the pro-sexual commitment transcription factor AP2G has been recently

identified, with low Lyso-PC concentrations increasing the proportion of sexual forms in

vitro (27). Furthermore, gametocyte development 1 (GDV1) has been identified as the

de-repressing effector which counters the epigenetic silencing and promotes AP2G

expression leading to the induction of gametocytogenesis (28). Still, it is not fully

understood how sexual investment and gametocytaemia is regulated in vivo, and if or

how transmission to mosquitoes can be optimized by the parasite to conserve

resources. Recently, we have shown that *P. falciparum* parasites isolated at the end of

the dry season are transcriptionally distinct from those of subjects with febrile malaria in

the transmission season (15, 29), but alteration of sexual commitment and

gametocytogenesis were not investigated. The observed transcriptional differences

reflected longer parasite circulation within each ~48 h intra-erythrocytic cycle in the dry

season, likely increasing splenic clearance of infected cells and thus maintaining the low

parasitaemia observed during the dry season (15). Here we aimed to examine sexual

commitment and gametocytogenesis of parasites persisting through the dry season and

explored possible parasite adaptations during mosquito-free periods. We quantified

sexual commitment and female and male gametocyte densities in blood samples

collected from children at different times of the year and with different clinical

presentations, and we related them to total parasite density and infection length.

Furthermore, we determined longitudinally the Lyso-PC concentration in plasma of

children, and assessed its ability to induce *P. falciparum* sexual commitment and

gametocytogenesis in vitro. Finally, we adapted a previously described within-host

mathematical model that considers asexual parasites, sexually committed parasites, and

gametocytes (30) to infer the dynamics of gametocyte number and proportion as *P.*

falciparum infections progress from recently transmitted to chronic asymptomatic

carriage over several months.

109
Results
Sexual stages differ between subclinical *P. falciparum* at the end of the dry season and

***P. falciparum* causing clinical malaria in the wet season.** We first investigated the

gametocyte-related transcriptome (31) within a whole genome RNA sequencing dataset

(15) comparing persistent *P. falciparum* detected by rapid diagnostic test (RDT)[+] in 12

asymptomatic Malian children at the end of the dry season (May) to *P. falciparum* of 12

age- and gender- matched Malians presenting clinical malaria with higher parasiatemias

in the wet season (MAL). Principal component analysis of 333 gametocyte-specific genes

with detectable number of reads (out of the 808 defined as gametocyte-related by ((31),

Table S1) showed segregation of transcription profiles based on seasonality (Fig. 1A).

Differentially expressed genes (DEGs) determined at a false discovery rate threshold of

5% resulted in 205 DEGs, with 146 transcripts up-regulated, and 59 transcripts down-

regulated in the dry season samples compared to clinical malaria samples (Fig. 1B and

Table S1). These transcriptional differences prompted us to quantify female and male

gametocytes in the blood of the two groups of children. First, we determined total

parasite (PFGLY, PF3D7_1420400) and total gametocyte densities (PFg17,

PF3D7_1319800) by qRT-PCR, and observed that these were higher in clinical malaria

cases in the transmission season (MAL) than in asymptomatic cases at the end of the dry

season (May) (Fig. 1C). Then, we determined the density of female (P25,

PF3D7_1031000) and male (PFMGET, Pf3D7_1469900) gametocytes, based on a

standard curve obtained from sorted gender-specific fluorescent gametocytes (32). This

analysis revealed higher numbers of both female and male gametocytes per mL of blood

in clinical cases of malaria during the wet season (MAL) compared to asymptomatic

carriers at the end of the dry season (May) (Fig. 1D). However, when we quantified

either gametocyte transcript relative to the total *P. falciparum* level present in each

subject's sample, normalized to a housekeeping gene transcribed by all parasite stages,

we observed an inverse trend. There were proportionally fewer female and male

gametocytes in malaria cases in the wet season (MAL) than at the end of the dry season

(May), although the difference was significant for the female comparison only (Fig. 1E).

We also quantified phospholipids known to influence *P. falciparum* sexual commitment

in vitro (27) in the plasma of the same children through shotgun lipidomic analysis.

Malaria cases in the wet season (MAL) showed lower levels of several species of the

Lyso-PC (Fig. 1F) and lysophosphatidylethanolamine (Lyso-PE) (Fig. 1G) families

compared to asymptomatic carriers at the end of the dry season (May), which could

potentially increase *P. falciparum* sexual commitment in malaria cases (MAL).

Nevertheless, transcription of the sexual commitment regulators (AP2G,

PF3D7_1222600; and GDV1, PF3D7_0935400), sexually committed schizonts (MSRP1,

PF3D7_1335000), or early gametocyte marker (GEXP5, PF3D7_0936600) were not

significantly different by qRT-PCR (Fig. 1H) nor were they differentially expressed in the

abovementioned RNAseq analysis (Table S1).

150
Fig. 1 | Sexual stages differ between subclinical P. falciparum at the end of the dry season and *P.*
***falciparum* causing clinical malaria in the transmission season. A**. Principal components analysis of RNA-
Seq data of 333 gametocyte specific *P. falciparum* genes from parasites collected at the end of the dry
season and from clinical malaria cases (n= 12 May, 12 MAL). **B**. Heatmap showing normalized reads of 205
gametocyte specific *P. falciparum* of differentially expressed genes (DEGs) (rows) for each subject
(columns) from *P. falciparum* collected at the end of the dry season (May) and at the first clinical malaria
case (MAL) in the ensuing transmission season. **C.** Asexual parasite (left) and gametocyte (right) densities
detected by qRT-PCR of RDT[+] subclinical children at the end of the dry season (May = 35) and of children
with their first clinical malaria episode (MAL = 27) in the wet season. **D.** Female (left) and male (right)
gametocyte densities detected by qRT-PCR of RDT[+] subclinical children at the end of the dry season (May =
35) and of children with their first clinical malaria episode (MAL= 27) in the wet season. **E.** Proportion of
female (left) and male (right) gametocytes detected by qRT-PCR of RDT[+] subclinical children at the end of
the dry season (May) and of children with their first clinical malaria episode (MAL) in the wet season. **F.**
Plasma concentration of Lyso-PC species of RDT[+] subclinical children at the end of the dry season (May)
and of children with their first clinical malaria episode (MAL) in the wet season. **G.** Plasma concentration
of Lyso-PE species of RDT[+] subclinical children at the end of the dry season (May) and of children with their
first clinical malaria episode (MAL) in the wet season. **H.** Relative expression of AP2G, GDV1, MSRP1, and
GEXP5, detected by qRT-PCR of RDT[+] subclinical children at the end of the dry season (May n > 15) and of
children with their first clinical malaria episode (MAL n > 23) in the wet season. Lipid data represented as
median ± IQR; two-way ANOVA with Tukey multiple comparisons correction. Parasitaemia and

gametocytemia and AP2G, GDV1, MSRP1, and GEXP5 expression data represented as median ± IQR;
Mann-Whitney test.

173
174
Sexual commitment and gametocytogenesis dynamic in persisting infections. We then

determined the dynamics of *P. falciparum* sexual commitment and gametocytogenesis

in persistently infected individuals, and examined associations between plasma Lyso-PC

and Lyso-PE, monitoring longitudinally ~20 children who were continuously infected

from the wet season until the end of the dry season. We quantified the number of total

parasites, and fully formed female and male gametocytes by qRT-PCR at four different

time points in asymptomatic children persistently infected since the wet season (Oct)

through the dry season until its end (Jan, Mar and May); and in a parallel group of age-

matched individuals presenting with clinical malaria during the ensuing wet season

(MAL). We observed that total parasite density was higher in malaria cases (MAL), than

at any time-point of the asymptomatic infections, whether in the wet (Oct) or the dry

season (Jan, Mar and May), while no differences were found between any of the time

points of asymptomatic infections (Oct, Jan, Mar and May) (Fig. 2A). Similarly, we

observed higher densities of both female and male gametocytes in malaria cases (MAL)

than at any time point of the asymptomatic infections (Oct, Jan, Mar and May), and

again no differences within the persisting period (Fig. 2B). Conversely, the proportion of

male gametocytes increased significantly at later time points of persisting infections

(Mar and May) than in malaria cases during the transmission season (MAL) (Fig. 2C).

Moreover, we found no significant difference in the sexual commitment regulator AP2G

between incident (MAL) or persistent infections at any point analyzed (Oct, Mar and

May) (Fig. 2D). Quantification of Lyso-PC 18:0 in malaria cases compared to

asymptomatic infections progressing from the wet into the dry season again resulted in

decreased Lyso-PC values in clinical malaria cases (MAL) compared to asymptomatic

infections (Oct, Mar and May) (Fig. 2E). We further detected a decrease of Lyso-PC from

the wet season (Oct) to the end of the dry season (May), but no difference was seen

between the two available time points of the dry season (Mar and May) (Fig. 2E).

Furthermore, Lyso-PC 18:0 levels in persistent infections (Oct, Mar and May) were

comparable to non-infected individuals at the end of the dry season, indicating that

asymptomatic infections have no impact on Lyso-PC 18:0 (Fig. 2E and Fig. S1).

Accordingly, a relation between Lyso-PC levels and parasitaemia or gametocytaemia was

only significant when considering clinical malaria plus all asymptomatic samples (Oct,

Mar, May and MAL) (r=$-$0.68 p< 0.0001; r=$-$0.56 p< 0.008 and r=$-$0.67 p< 0.0001 for

parasitaemia, female and male gametocytaemia respectively), but disappeared when

each group was analyzed independently, or in all persisting asymptomatic infections

analyzed together (Fig. 2F).

Fig. 2 | Sexual commitment and gametocytogenesis dynamic in persisting infections. A, Parasite, **B.** Female (left) and male (right) gametocyte, and **C.** Proportion of female (left) and male (right) gametocytes densities detected by qRT-PCR of RDT[+] subclinical children at different times during the wet (Oct n = 10) and the dry season (Jan, Mar and May n > 15), and during clinical malaria episodes (MAL n = 19). **D.** Relative expression of AP2G detected by qRT-PCR of RDT[+] subclinical children at different times during the wet (Oct n = 6) and the dry season (Jan, Mar and May n > 11), and during clinical malaria episodes (MAL n = 19). **E.** Plasma concentration of Lyso-PC 18:0 of RDT[+] subclinical children at different times during the wet (Oct n = 17) and the dry season (Mar and May n = 19), and during clinical malaria episodes (MAL n = 15). Parasite gene expression and lipid data were compared through Kruskal-Wallis test with multiple comparisons. **F**, Correlation between Lyso-PC 18:0 and parasite (left), female gametocyte (centre) or male gametocyte (right) densities detected by qRT-PCR detected by qRT-PCR of RDT[+] subclinical children at different times during the wet (Oct n = 10) and the dry season (Mar and May n > 15), and during clinical malaria episodes (MAL n = 19). *P* and r determined by Pearson correlation.

Decreased Lyso-PC during clinical malaria does not induce higher sexual commitment or gametocytaemia. We then questioned if the low levels of Lyso-PC and Lyso-PE species in plasma of malaria cases compared to persistent infections at the end of the dry season (Fig. 1F and 2E) could induce *P. falciparum* sexual conversion or gametocytaemia differently in vitro. We exposed parasites of the *P. falciparum* 2004 line (Pf2004) to plasma of children collected during clinical malaria cases (MAL) versus those

of children with asymptomatic infections at the end of the dry season (May) alongside the conditions described by Brancucci et al. to induce or inhibit sexual convertion and gametocyteamia. As reported earlier (27), we observed that 24 h after culturing Pf2004 without plasma or serum (noPl), and thus no Lyso-PC, over minimal fatty acid medium (MFA), AP2G expression increased compared to the same parasites cultured in the presence of German control plasma (PlDE) (Fig. 3A). However, MFA medium supplementation with 25% plasma from children presenting with clinical malaria during the wet season (PlMAL), vs. plasma from children with asymptomatic infections at the end of the dry season (PlMay) resulted in similar APG2 induction 48 h after beginning of treatment (Fig. 3A), and similar conversion rate of parasites into gametocytes on day 5 after treatment (defined by = d5gametocytes/ d2parasites) (Fig. 3B). Absence of plasma (noPl) resulted in higher gametocytaemia than observed on the German pooled plasma condition (PlDE) (Fig. 3B), plasmas from malaria cases (PlMAL) or asymptomatic infections at the end of the dry season (PlMay) revealed similar conversion rates (Fig. 3B). Although Lyso-PC differences between non-infected or asymptomatic children and malaria cases were significant (Fig. 1F-G and Fig. 2E) the 2-4 fold differences observed were much lower than the several orders of magnitude included in the standard curve used to establish that gametocyte conversion responds to low Lyso-PC in the culture medium (27), and were thus likely insufficient to impose alterations in sexual commitment or gametocytaemia.

Fig. 3 | Decreased Lyso-PC of clinical malaria cases does not induce higher sexual commitment or gametocytaemia in vitro. A, Relative expression of AP2G detected by qRT-PCR of Pf2004 24 h after 24 h culture with 25% plasma collected during clinical malaria episodes (MAL n = 21), or at the end of the dry season (May n = 19). **B**, Pf2004 conversion rate (gametocytes d5/parasites d2) (left) and gametocyte density (right) detected by microscopy, after 24 h culture with 25% plasma collected during clinical malaria episodes in the wet season (MAL n = 19) or at the end of the dry season (May n = 18), alongside no plasma (noPl) and German (PlDE) conditions.

Progressing infection leads to apparent differences in gametocytogenesis. We observed decreased densities (Fig. 1D and Fig. 2B) but higher proportions (Fig. 1E and Fig. 2C) of female and male gametocytes in the dry versus wet season, or in asymptomatic versus clinical malaria in the wet season (Fig. 2C), without concomitant differences in the expression of regulators of sexual commitment, or transcriptional markers of sexually committed schizonts or early gametocytes (Fig. 1H and Fig. 2E and Table S1). To further investigate this apparent conflict, we first ascertained if there were indeed indications of altered gametocytogenesis between parasites persisting through the dry season and parasites causing malaria in the wet season. Specifically, we investigated in more detail the transcriptional profiles of early and late-stage gametocytes. Using markers of early or late gametocyte development gathered from different sources ((33, 34) and Jeninga et al., unpublished Oct 2021) and intersecting them with the RNAseq data of clinical malaria cases and dry season samples shown in Fig. 1, we identified 163 gametocyte-specific transcripts with quantifiable reads in samples from individuals with circulating parasites persisting through the dry season and parasites causing malaria in the wet season. Of those, 109 transcripts define early gametocytes; and 54 transcripts are characteristic of late-stage gametocytes (highlighted in Table S1). Expression of these transcripts in asymptomatic carriers at the end of the dry season (May) and malaria cases (MAL) in the wet season showed an enrichment of upregulated transcripts of late-stage gametocytes in the dry season (79,6%) (43/54 genes), which clearly contrasted with the proportion of early-gametocyte transcripts or total *P. falciparum* transcripts upregulated at that time, which showed 38,5% (42/109 genes) and 33,4% (1131/3381 genes) upregulated in the dry season respectively (Fig. 4A).

The lack of difference in sexual commitment or early-stage gametocyte transcripts, along with the strong difference in late-stage gametocyte transcript between the end of the dry season (May) malaria cases (MAL) parasitaemias hints not at a difference in gametocyte production, but rather an effect during the 8-12 days of gametocyte development, in both/either the sexual and/or the asexual parasite compartments, suggesting that length of infection may play a role in the results observed. Hence, we

investigated if the observed patterns of female and male gametocyte densities and

proportions were generalizable to non-seasonal settings where infections can also

persist. While Mali shows a very contrasting seasonal pattern with 5-6 months

interruption of transmission every year (14), Uganda has uninterrupted year-round

transmission with peaks in times of stronger rain (35). RT-qPCR data quantifying

transcripts of total parasite (varATS) and gametocyte (P25, PF3D7_1031000) densities

from samples of 150 Ugandan individuals with asymptomatic parasitaemias persisting

for over one month (n=122) and age-matched individuals presenting with clinical

malaria (n=38) had been previously published (35) and were retrieved from ClinEpiDB

(36). The data indicated that parasite and gametocyte densities were increased in

clinical malaria cases (MAL) compared to asymptomatic infections (Asymp) (Fig. 4B and

C); while the proportion of gametocyte transcripts relative to total *P. falciparum*

transcripts showed again an inverse trend (Fig. 4D). As seen with the Malian samples,

the proportion of gametocytes was higher in persisting parasitaemias (Asymp)

compared to malaria cases (MAL) (Fig. 4D). Malian and Ugandan samples vary in

seasonality of transmission, but share the fact that clinical malaria cases are likely

recently transmitted (37), while asymptomatic parasitaemias have persisted longer than

one month in the Uganda cohort, and at least 5 months in the Malian setting. The

parallel in infection length between samples of incident clinical malaria and

asymptomatic infections in the two contexts led us to hypothesize that gametocyte

numbers and proportions change dynamically in the course of a *P. falciparum* infection,

leading to the observed patterns in Mali and Uganda. To address this, we devised a

mathematical model including three different parasite populations_ asexual parasites

(P), sexually committed parasites (P_G) and gametocytes (G), all of which can circulate or

be adherent as previously developed (30) (Fig. 4E). As the original model by Cao et al.

was not able to sufficiently explain the observed dynamics when fitted to the data, we

extended the model in different ways. First, we incorporated our recent observation of

increased clearance of longer-circulating infected erythrocytes in persisting

parasitaemias compared to recently transmitted parasites (15), which promoted

different removal rates of asexual and sexually-committed parasites during distinct

phases of acute and chronic *P. falciparum* infection in our model (see Methods). In

addition, we distinguish between three phases *during P. falciparum* infection in which

the parasite multiplication factor varies due to immune response or adhesive properties

of the parasite. Using an iterative Bayesian fitting-approach to fit the model to the

symptomatic and asymptomatic parasitaemia and gametocytaemia levels of the Malian

cohort, we quantitatively assessed several biological parameters including sexual

commitment rate, gametocyte circulation time, parasite multiplication rate and

gametocyte sequestration time, thereby confirming previously determined ranges (30,

38) (Fig. S2). Our adapted model was able to provide a dynamic depiction of

parasitaemia levels, and gametocyte densities and proportions, as we observed within

the Malian data (Fig. 4F-H). We defined that the death rate of asexual and sexually

committed parasites increased in later stages of the 48 h cycle, shifting toward the

removal of older parasite stages and leading to a constant but oscillating number of

asexual parasite in circulation (highlighted in the box-inlay of Fig. 4F). This can to some

extent explain the large variation seen in parasitaemias, especially during the chronic

phase, which additionally may arise from different collection time points after start of

infection for the individual samples (i.e., pooling of the individual measurements from

Oct-May and calculation of average for each individual by the area under the curve over

the course of blood collections during the asymptomatic phase and fixing it to one time

point for fitting (see Methods)). Keeping a constant sexual commitment rate, f,

throughout acute and chronic infection phases, our model predicts a higher number and

a lower proportions of gametocytes in the acute phase_ a trend that is inverted during

the transition phase of infection due to effective removal of asexual parasites by an

active immune response (Fig. 4G and H). During the chronic phase of infection, when

parasitaemias remain constant or are slowly decreasing, gametocyte proportions are

decreasing as well until reaching an equilibrium roughly 4-5 months after the start of

blood stage infection. Overall, our model provides a dynamic depiction of gametocyte

densities and proportions similar to the data we observed in Mali and in Uganda,

without altering the sexual commitment rate (Fig. 4C and D).

Fig. 4 | Progressing infection leads to apparent differences in gametocytogenesis. A, Fold change between parasites collected at the end of the dry season and from clinical malaria cases (May n= 12, MAL n= 12) of early and late gametocyte transcripts according to López-Barragán et al., 2011; Poran et al., 2017 and Petter unpublished data. Each dot represents a gene and color identifies range in fold change, red: upregulated in May, yellow: not DEG, and blue: downregulated in May). **B**, Parasitaemia, **C**, gametocytemia and **D**, proportion of gametocyte in symptomatic (MAL n = 19 and 38 in Mali and Uganda respectively) and asymptomatic infections (May n = 23 and Asymp n = 122) in Mali and Uganda quantified by RT-qPCR. Multiple Non-Parametric test with Tukey-Correction for multiple comparisons. **E**, Diagram of the mathematical model to describe parasite dynamics including individual compartments, transitions and death rates; Asexual parasites (P) replicate in a 48 h cycle, being cleared at an age dependent death rate (σ_a). A fraction of asexual parasites (f) transitions to the sexually committed parasite compartment (P_G), goes through a round of replication and develops into gametocytes (G). Gametocytes mature over 5 distinct stages in independent model compartments. Asexual parasites (blue), sexually committed parasites (orange) and gametocytes (grey) are indicated shaded if not circulating due to sequestration. **F**, Model prediction and clinical data of asexual parasites. The small inlay indicates a zoom-in showing the oscillating parasite dynamics within 48 h-intervals. **G**, gametocytes and **h**, the proportion of gametocytes per mL blood. Individual data points and median (black square) of wet (aqua circles) and dry (orange circles) season data throughout acute (aqua shaded area) and chronic (salmon shaded area) phase of infection are shown. Median fit (grey line) and 95% prediction interval of 100 model predictions of the 6[th] PyABC generation are shown as grey shaded area.

Discussion

The relative absence of mosquitoes during the months-long dry season and the short

lifespan of *P. falciparum* sexual-stage gametocytes (10, 11) renders the production of

gametocytes ineffective during most of the dry season; however, gametocytes must be

available to resume malaria transmission when mosquitos return with the wet season.

Whether *P. falciparum* maintains a constant rate of production of sexual forms or

adjusts gametocytogenesis to maximize transmission has been unclear. In this study,

with samples from Malians exposed to alternating 6-month dry and wet seasons, and

samples from incident malaria and persisting infection from year-round transmission

region of Uganda, we showed that incident cases of malaria consistently have higher

parasite and stage V gametocyte densities than asymptomatic infections; however, the

proportion of gametocytes is higher in the asymptomatic persisting infections (Fig. 1, 2

and 3). Additionally, our data from Mali indicated that transcription of genes

characterizing sexually committed parasites and stage I gametocytes (39, 40) quantified

by RT-qPCR did not appear affected by the season or clinical symptoms (Fig 1 and 2), nor

were reads of over 100 genes characteristic of early gametocytes and detected by

RNAseq of circulating parasites particularly enriched for DEGs. This, clearly contrast with

representative expressed genes of late gametocytes that were very much enriched of

transcripts upregulated in persisting asymptomatic infections at the end of the dry

season (Fig. 4 and Table S1).

Although it has been suggested that *P. falciparum* investment in gametocyte production

adjusts to host nutrient availability (41), drug pressure (21), transmission intensity (42),

and even seasonality (43, 44), results are often inconsistent and potentially affected by

the dynamics of asexual parasite growth, without necessarily indicating altered sexual

commitment. In Sudan, following the long dry season, gametocyte density estimates

have been reported to increase prior to the highest transmission peak (45), suggesting a

higher gametocyte investment. However, in Burkina Faso, gametocyte densities were

higher at the peak and end of the transmission season rather than during the dry season

(46), while the efficiency of membrane feeding to infect mosquitoes was increased early

in the transmission season compared to peak of the malaria season or the dry season

(47). Nevertheless, these studies did not monitor longitudinally the same parasite

population during the dry and early wet season. In a recent study also in Burkina Faso,

transmission was shown to be more effective in persistent than in incident infections,

but no data was collected after day 35, and collections were all done during the wet

season (24) when a mix of more and less recently transmitted parasite clones is likely in

individuals without symptoms.

In vitro, depletion of Lyso-PC or choline have been shown to induce AP2G expression,

leading to increased gametocyte production (27, 48), but associations in humans are

inconsistent. In Rwanda, low Lyso-PC associated with higher parasite densities (49); and

in Ghana, although correlations with parasitaemia were not reported, Lyso-PC levels in

human blood samples were inconsistently related with gametocyte-committed ring

stages converting into gametocytes in vitro (40). Our data show, like Orikiiriza et al., an

association of lower Lyso-PC levels with increased parasitaemia and severity of malaria

(Fig. 1F-G and 2E), but the association was lost at low parasitaemias, and persistent

infection at the end of the dry season did not affect Lyso-PC compared to non-infected

individuals (Fig. 2F and Fig. S1).

The effect of drugs on sexual conversion in vitro has been previously reported.

Specifically, sub-curative doses of dihydroartemisinin (DHA) stimulate AP2G expression

and sexual conversion when drug exposure occurs at the trophozoite stage, but not at

the early ring stage. The observed increase could not synergize nor add to increases

promoted by depletion of choline, suggesting, as DHA affects lipids, that the two effects

might be the same (48). Our RNAseq (Table S1) and RT-qPCR (Fig. 1H and 2D) data show

constant low-level expression of AP2G, the master regulator of gametocytogenesis, and

equally unaltered expression of transcripts characteristic of committed schizont and

early gametocytes (Table S1 and Fig. 1H), suggesting that reductions in circulating Lyso-

PC induced by uncomplicated clinical malaria are not capable of affecting *P. falciparum*

sexual commitment. Accordingly, we observed in vitro that supplementation with

plasma from clinical malaria cases (low Lyso-PC) or end of the dry season persistently

infected individuals (normal Lyso-PC) did not differentially induce AP2G expression or

the rate of sexual conversion (Fig. 3), suggesting that Lyso-PC differences introduced by

P. falciparum infection are not sufficient to trigger the mechanisms demonstrated by

Brancucci et al. and Portugaliza et al.. Nevertheless, it is possible that Lyso-PC levels in

particular bone marrow niches are sufficiently low to trigger parasites' sexual

commitment, although this would require that infected erythrocytes be at that niche to

respond to the stimulus.

Previous reports highlighting differences in gametocytogenesis as a product of altered

sexual commitment did not account for longevity of infection, nor for dynamic changes

within the asexual compartment during a persistent infection. Greischar et al.

demonstrated the difficulty of quantifying true transmission investment by comparing

existing quantification methods on simulated data where true transmission investment

is known (50), and Prajapati and colleagues relied on in vitro culture of sexually

committed circulating parasites to define samples with higher and lower commitment

rates (51). Here we show empirically, and reinforce with a mathematical model, that the

proportion of sexual stages is inversely related in recently transmitted versus chronic

infections, without alteration in the expression of regulators of sexual commitment and

while maintaining a constant rate of sexual commitment. Our recent study in which we

found that infections that persist beyond the acute immune response exhibit prolonged

circulation and increased splenic clearance (15) allowed us to complement the model by

Cao et al. with an increased asexual parasite death rate following the immuno-

responsive period, which promoted the progressively increased representation of sexual

stages in later phases of the infection.

Gametocyte proportions are low early on in infection because asexual replication

efficiently increases asexual parasite numbers while adaptive immunity is absent, and

the lag of ~8 days in gametocyte development in the bone marrow has its impact. In our

model we propose that later, as immunity prevents fast asexual growth and the

parasites persisting further are more efficiently removed in the spleen, the opposite lag

effect is then shown, and the proportion of gametocytes finally reflects, despite keeping

the same sexual commitment rate, that of parasitaemias ~8 days behind.

In conclusion, our data is consistent with a model in which *P. falciparum* maintains a

constant rate of sexual commitment such that the proportion of gametocytes increases

over time as infection persists and asexual parasite numbers decline more rapidly, a

strategy that may help overcome the vector-free period and increase transmission

efficiency.

465
Materials and Methods
Gametocyte-specific gene expression. We used RNAseq data of leucocyte-depleted

blood pellets from 12 children with asymptomatic malaria at the end of the dry season

and an age- and gender-matched cohort presenting with their first malaria case in the

ensuing transmission season reported in (15). Out of the 808 gametocyte specific genes

(31) we isolated normalized reads for 333 gametocyte specific genes that were

detectable in the RNAseq data set (15) to perform a principal component analysis. Using

Deseq2, 205 genes were determined to be differentially expressed between the dry

season and malaria cases and were plotted in the heatmap shown in Fig. 1B.

475
Study subjects and ethical approval. Approval by the Ethics Committee of Heidelberg

University Hospital, the Faculty of Medicine, Pharmacy and Odontostomatology

(FMPOS) at the University of Bamako, and the National Institute of Allergy and

Infectious Diseases of the National Institutes of Health Institutional Review Board was

obtained to conduct the study registered at ClinicalTrials.gov with the identifier

NCT01322581. Clinical data and samples were obtained from 2017 to 2019 in a cohort

study previously described (52), conducted in Kalifabougou, Mali, where malaria

transmission occurs between June and December, and antimalarial drugs are provided

by a single clinic and pharmacy. All participants and the parents/guardians of included

children provided written informed consent. Exclusion criteria at enrolment included

haemoglobin <7 g/dL, axillary temperature ≥37.5°C, acute systemic illness, or use of

antimalarial or immunosuppressive medications in the 30 days preceding enrolment.

Clinical malaria cases were prospectively determined by passive surveillance and

defined by axillary temperature ≥37.5°C, ≥2500 asexual parasites/μL of blood, and no

other evident cause of fever. Malaria episodes were treated with a 3-day standard

course of artemether/lumefantrine according to Malian national guidelines. Cross-

sectional clinical visits and blood draws were performed late in the transmission season

(October), and at the beginning (January), mid (March) and end (May) of the dry season.

Sample collection. Venous blood (4 mL if donor age was below 4 years, and 8 mL if above) was drawn by venepuncture on wet and dry season cross-sectional visits, and at first malaria episode of the transmission season. Samples are referred as MAL for malaria cases or by time point of collection. Blood was collected in sodium citrate-containing cell preparation tubes (Vacutainer CPT Tubes, BD) and transported to the laboratory where RBC pellet and plasma were separated and stored at −80°C within three hours of blood draw. Plasma used for lipidomic analysis and induction of gametocytogenesis was separated by centrifugation and immediately frozen in liquid N_2.

RNA extraction and quantification of parasite transcripts. RNA was isolated from 200 µl RBC pellet by phenol-chloroform extraction using TRIzol (Ambion), followed by DNAse treatment using the DNAse I Amplification Grade Enzyme (Invitrogen). RNA was quantified by absorption measurements at 230, 260 and 280 nm on a Citation3 Imaging plate reader, followed by cDNA synthesis using the Superscript IV Enzyme (Invitrogen) according to manufacturer instructions. Taqman RT-qPCR multiplex assay (Applied Biosystems) was performed on a qTower (Analytic Jena) to quantify the female gametocyte marker ookinete surface protein P25 (PF3D7_1031000), the male gametocyte transcript PFMGET (Pf3D7_1469900), the master regulator of sexual commitment Ap2G (PF3D7_1222600) and a parasite housekeeping gene (Glycine-tRNA ligase, PF3D7_1420400). All primers and probes sequences are listed in the key resources table. To account for fluorescence leakage between channels a compensation matrix was calculated based on single color fluorescence measurements by an application specialist at Analytic Jena. Gexp5 (PF3D7_0936600) expression levels were quantified in a Taqman RT-qPCR duplex assay together with Glycine-tRNA ligase as a housekeeping gene. Expression levels of GDV1 (PF3D7_0935400), PFg17 (PF3D7_1319800) and MRSP1 (PF3D7_1335100) were determined in further Taqman RT-qPCR assays. Relative quantification was performed by ΔΔCt method using Glycine-tRNA ligase as housekeeping gene and MAL samples a calibrator group.

Parasite culture and preparation of a standard curve for gametocyte quantification.

Pf2004 parasites were cultured at 37° C in 5% O_2, 5% CO_2 and 90% N_2, at 5% haematocrit

in RPMI-1640 medium (with L-glutamine and HEPES (Gibco)). Parasite culture medium

was supplemented with 0.24% Sodium Bicarbonate (Gibco), 100µM Hypoxanthine (cc

pro), 25 µg/mL gentamycin (Gibco)), and 0.25% Albumax II (Gibco). Medium was

changed daily, parasitaemia assessed by Giemsa-stained blood smear and the culture

split according to experimental needs. *PfDyn*GFP/*PfP47*mCherry parasite lines (32) were

cultured in presence of 15 µg/ml of Blasticidin-S-HCL (Invitrogen) and 2 µM

Pyrimethamine (Sigma Aldrich) as described for non-integrating plasmids and P52

integrating plasmid. Parasites were grown to >4% parasitaemia when 50 mM N-

Acetylglucosamine (Sigma Aldrich) was added to the culture to eliminate asexual stages.

After 4 consecutive days of N-Acetylglucosamine treatment, culture in complete RPMI

was resumed. Gametocyte development was monitored daily by smear and on day 5

after start of gametocyte induction magnetic enrichment was performed with MACS

columns (Miltenyi Biotec). Gametocytes were sorted on a BD FACSAria III on day 8,

keeping cells at 4° C in suspended activation buffer, (gating strategy in Fig S3). RNA of a

defined numbers of sorted male and female gametocytes was extracted and reverse

transcribed, a standard curve prepared by serial dilution and used for RT-qPCR as

described above.

Phospholipid analysis. May and MAL plasma samples (shown in Fig 1) were analyzed by

shotgun lipidomics at Lipotype as described in (53). Briefly, automated samples

extraction was followed by direct sample infusion and high-resolution Orbitrap mass

spectrometry including lipid class-specific internal standards to assure absolute

quantification of LysoPC and LysoPE, using LipotypeXplorer for identification of lipids in

the mass spectra. Oct, Mar, May and MAL samples (shown in Fig 2) were analyzed by

Liquid chromatography-mass spectrometry (LC-MS). Samples were extracted, processed,

and analyzed at the Lipidomics Unit at the Institute of Physiological Chemistry in Mainz,

using the liquid-liquid extraction protocol as described (54). All values have been

normalized to analyzed plasma volume.

In vitro induction of *P. falciparum* sexual commitment and gametocytogenesis. In vitro

induction of *P. falciparum* sexual commitment was performed as described by

(Brancucci et al., 2015). Pf2004 parasites were synchronized to a 4 h window by

repeated treatment with 5% sorbitol and MACS purification. Parasites at 26 ± 2 h post

invasion (hpi) were plated on 48-well plates in 500 µl culture volume at 0.8-1%

parasitaemia and 2.5% haematocrit and cultured in minimum fatty acid medium (mFA,

RPMI1640 with 100 µM hypoxanthine, 25 µg/ml gentamycin, 0.39% fatty-acid-free BSA,

30 µM palmitic acid and 30 µM oleic acid) supplemented with 25% plasma from

asymptomatic children at the end of the dry season (PlMay), or 25% plasma from

malaria cases during the rainy season (PlMAL), in addition to the conditions used by

Brancucci et al. (mFA alone, and mFA supplemented with 25% German plasma (PlDE)).

After 24 h of treatment, the medium was changed to culture medium and subsequently

changed daily. At 48 h after beginning of treatment 30 IU/ml heparin were added to

block re-invasion of parasites and eliminate asexual forms. At 0, 48 and 120 h post

treatment RBC pellets were collected for RNA quantification of gametocyte markers by

RT-qPCR; parasitaemia and gametocytemia determination by flow cytometry (50 µl of

culture stained with 5x SYBR Green II (Invitrogen) and 7,5 µM MitoTracker (Applied

Biosystems) for 30 min at 37 °C and acquired with CountBright Absolute Counting Beads

(Thermo Ficher) on a FACS Canto II or an LSR II and analyzed using FlowJo software 10.2

or higher versions (Tree Star)). Additionally, Giemsa-stained blood smears were

prepared at 48 h and 120 h post-treatment and blindly analyzed by microscopy,

counting RBCs, parasites, and gametocytes in 18 fields of view. Multiplication rate of

parasites was calculated as $\frac{parasitemia\ 48\ h}{parasitemia\ 0\ h}$, and conversion rate as $\frac{gametocytemia\ 120\ h}{parasitemia\ 48\ h}$.

Analysis of early and late gametocyte markers in RNA-seq data. We compiled a list of

early and late gametocyte specific genes based on three RNA-seq studies comparing

distinct parasite stages, data from Poran et al. 2017 (compared day 2 versus day 9 of

gametocytogenesis by single cell RNAseq); well as data from Lopez-Barragan et al. 2011

(compared day 4 versus day 15 of gametocytogenesis by bulk RNAseq) and Jeninga et

al., unpublished (compared day 4 of gametocytogenesis versus day 10 of FACS sorted

male and female gametocytes by bulk RNAseq). Poran et al. selected early and late

gametocyte genes as genes showing a 20-fold higher expression in early or late

gametocytes over the respective other group. Late gametocyte genes were also

required to show a 20-fold higher expression in late gametocytes compared to ring

stage parasites, this was not the case for early gametocyte genes. In Lopez-Barrgagan's

data, early gametocyte genes were defined as genes having a 10-fold expression

increase in early gametocytes compared to late gametocytes. Genes showing a 20-fold

higher expression in late stages were designated as late gametocyte genes. Expression

differences to asexual stages were not considered. In Jeninga et al. unpublished genes

showing a 3-fold or higher expression difference to both asexual stages and late

gametocytes were selected as early gametocyte genes, regardless of gametocyte sex.

Intersecting the full list of early and late gametocyte genes with genes transcribed in

circulating parasites stages, defined by RNAseq data of Andrade et al., generated a final

list of 109 early gametocyte genes and 54 late gametocyte genes to evaluate expression

differences of these genes between dry and rainy seasons.

Quantification of parasites in samples of Ugandan asymptomatic carriers by RT-qPCR.

We compared the Malian data from dry and wet season with data from 150 Ugandan

individuals enrolled in a 2-year longitudinal study including monthly standardized clinical

evaluation and blood collection (35) retrieved from ClinEpiDB

(https://clinepidb.org/ce/app/record/dataset/DS_51b40fe2e2122). Asymptomatic *P.*

falciparum positive individuals (20.4 years ± 17.2, median: 13.3 years) were included

based on the following criteria: i) ≥ two PCR+ time points, ii) ≤ 3 negative time points in

between positive, iii) same clone (determined by deep amplicon sequencing of atypical

membrane antigen 1 (AMA1) as described in (55) present at all time points of one

asymptomatic infection. Additionally, we used samples from an age-matched group of

38 clinical malaria cases (tympanic temperature > 38°C or history of fever in the

previous 24 hours and *P. falciparum* positivity by microscopy). *P. falciparum* parasite

and gametocyte density in blood samples were measured by RT-qPCR of multi copy

conserved *var* gene acidic terminal sequence (56) and Pf25 (P25, PF3D7_1031000),

respectively. Average levels of parasitaemia and gametocytemia during each

asymptomatic infection, were estimated as the area under curve over the course of

infection divided by length of infection. For symptomatic infection only the

quantification before treatment on day of diagnosis was considered. For comparison

between two data sets, a multiple non-parametric comparison between groups of

asymptomatic and symptomatic in Mali and Uganda has been performed using Tukey-

Correction for multiple comparisons.

Mathematical model. The developed mathematical model extends a model used

previously to describe asexual parasite and gametocytes dynamics (30). The model

considers 3 different main parasite populations, i.e., distinguishing between asexual

parasites (P), sexually committed parasites (P$_G$) and gametocytes (G) (Fig. 4E). Asexual

parasites develop and replicate within a red blood cell over a cycle of length a$_L$=48h

after which mature schizonts burst and release a number of r_P = 18-32 merozoites that

are able to invade new red blood cells. During this replication cycle, parasites are

assumed to be removed through splenic clearance and active immune response. In

extension to the previous model, we assume that this removal of asexual and

committed parasites follows an age dependent death rate, σ_a that differs in distinct

phases of acute and chronic *P. falciparum* infection. The following iterative equation

describes asexual parasite growth at discrete time steps for parasite age, *a*, and time, *t*,

increments of one hour.

$$P(a,t) = \begin{cases} P(a-1,t-1)e^{-\sigma_a} \ , a = 2,3,\dots,a_L \\ r_p \ P(a_L,t-1)e^{-\sigma_a} \ , a = 1 \end{cases} \tag{1}$$

$$P(a_s+1,t) = (1-f)P(a_s,t-1)e^{-\sigma_a} \tag{2}$$

638   The rate of parasite removal, σ_a, is assumed to increase with parasite age within one

639   replicative cycle and described by a sigmoidal curve given by:

$$\sigma_a = \sigma(a) = \frac{\sigma_{max}}{1+e^{p\sigma(T\sigma-a)}} \tag{3}$$

The parameter σ_{max} defines the maximal rate of parasite removal, T_σ, the age at which

50% of parasites are removed, and p_σ a scaling parameter.

At parasite trophozoite stage of a_s= 20hpi parasites adhere to the endothelium when at

the same time a fraction, f, of parasites enters sexual commitment. Committed

parasites continue the rest of the replication cycle producing only committed parasites

that enter the first gametocyte stage at the sequestration age, a_s, of the following

replication cycle.

$$P_G(a_s + 1, t) = f\, P(a_s, t-1)e^{-\sigma a} \tag{4}$$

$$P_G(a, t) = \begin{cases} P(a-1, t-1)e^{-\sigma a}, a = 2,3, \dots, a_L \text{ except } a = a_s \text{ and } a = a_s + 1 \\ r_p\, P(a_L, t-1)e^{-\sigma a}, a = 1 \end{cases} \tag{5}$$

650 Gametocytes develop over 5 distinct stages of which only mature stage 5 is found in

651 circulation. The development of gametocytes is modelled by:

$$G_1(t) = G_1(t-1)e^{-(m-\delta_G)} + \frac{P_G(a_s-1, t-1)e^{-\sigma a}(1-e^{-(m+\delta_G)})}{m+\delta_G} \tag{6}$$

$$G_2(t) = G_2(t-1)e^{-(m-\delta_G)} + \frac{m G_1\ (t-1)\ (1-e^{-(m+\delta_G)})}{m+\delta_G} \tag{7}$$

$$G_3(t) = G_3(t-1)e^{-(m-\delta_G)} + \frac{m G_2\ (t-1)\ (1-e^{-(m+\delta_G)})}{m+\delta_G} \tag{8}$$

$$G_4(t) = G_4(t-1)e^{-(m-\delta_G)} + \frac{m G_3\ (t-1)\ (1-e^{-(m+\delta_G)})}{m+\delta_G} \tag{9}$$

$$G_5(t) = G_5(t-1)e^{-(\delta_{G5})} + \frac{m G_4\ (t-1)\ (1-e^{-(\delta_{G5})})}{\delta_{G5}} \tag{10}$$

Immature gametocytes, that are sequestered, mature with a maturation rate, m, and

are lost by death rate δ_G, while mature circulating gametocytes are removed by a

splenic clearance rate δ_{G5}. As pointed out above the whole model is simulated by

discrete time-steps of one-hour increments.

Accounting for different phases of P. falciparum infection _ In our model the start of the

infection is marked by the beginning of blood stage infection, where we assume an

initial parasite load, IPL, with parasite age normally distributed with mean, μ_{IPL} and

standard deviation σ_{IPL}. During the acute phase of infection parasitaemias increase

rapidly leading to clinical symptoms within the first 2 weeks of blood stage infection

(57). The acute phase of infection is accompanied by the generation of an adaptive

immune response, including the production of effective antibodies, that takes ~14 days

to develop (58). To account for higher parasite removal due to the immune response,

we assume an increased maximal parasite removal rate σ_{max2} for age-dependent

parasite loss when an adaptive immune response is active. Increased parasite removal is

considered to occur 14 days post start of blood stage malaria infection and indicates the

beginning of the transition phase from the acute towards the chronic asymptomatic

phase of infection. In addition, as selection for less adhesive and less virulent parasite

strains by an active immune response is considered to be crucial for the maintenance of

a malaria infection over the course of several dry season months (15), we accounted for

this effect in our model by an increased value of T_σ for Eq. (3) during the chronic

asymptomatic phase, i.e., replacing T_σ by $T_{\sigma,2}$. This parameter change shifts clearance

rates towards affecting older asexual parasite stages, leading to a constant, but

oscillating number of asexual parasite numbers in circulation.

Parameter estimation _ The model described in Eqs (1-10) was applied to the data

assuming a fixed parasite life-cycle period of a_L=48 h, a sequestration age of a_S= 20 hpi,

as well as a parameter of T_σ=32 h and P_σ=0.7 characterizing age-dependent parasite

removal during the acute infection phase. All other parameters were fitted to the both

symptomatic and asymptomatic time points simultaneously using an approximative

Bayesian computing approach (59). Prior distributions for the individual parameters

spanning biologically reasonable ranges have been selected in accordance with previous

studies (Table 1). Due to the high model complexity and parameter dependencies,

estimations of multiple pyABC-runs diverge to different local minima for the individual

parameters. To obtain robust results for the parameters indicating phase-dependent

changes, several independent pyABC fitting runs were compared and the 6[th] generation

was chosen where parameter posterior distributions would still overlap. For the fitting,

values of individual patients in the dry season are condensed to one time point by

calculating the average levels based on the area under the curve over the course of dry

season patient surveillance. Values were log10-transformed to ensure comparability of

parasitaemia and gametocytemia. In addition, wet and dry season measurements were

assumed to represent the parasite dynamics at 2 and 6 weeks post infection,

respectively. Sensitivity of the results with regard to these time points was investigated.

Table 1. List of model parameters and respective prior distributions. All prior
distributions were uniformly distributed given the indicated ranges.

Parameter	Unit	Description	Prior range	Reference for prior range	89% CI of posterior range
ipl	Parasites /ml	Initial parasite load	1 - 10	(Hodgson et al 2015) (Schats et al 2015)	1.47 - 6.54
μ_{ipl}	h	Initial parasite age	4 - 10	(Eichner et al. 2001)	4.41 - 8.22
σ_{ipl}	h	Standard deviation of initial parasite age	4.5 - 6.5	(Cao et al. 2019)	4.89 - 5.81
r_p		Replication factor	18 - 32		$2.28 - 3.13 \times 10^1$
σ_{max}	h^{-1}	Maximal removal rates during acute phase of infection	0.01- 0.15	(Andrade et al. 2020)	$0.81 - 1.07 \times 10^{-1}$
σ_{max2}	h^{-1}	Maximal removal rates during transition phase of infection	0.01- 0.4		$2.13 - 2.39 \times 10^{-1}$
δ_G	h^{-1}	Death rate immature gametocytes	1e-05 − 1e-04	(Cao et al. 2019)	$3.53 - 8.60 \times 10^{-5}$
δ_{G5}	h^{-1}	Death rate mature gametocytes	$1.39 - 4.17 \times 10^{-3}$	(Cao et al. 2019)	$1.79 - 3.98 \times 10^{-3}$
m	h^{-1}	Gametocyte maturation rate	$1.6 - 4.6 \times 10^{-2}$	(Cao et al. 2019)	$2.76 - 4.29 \times 10^{-2}$
f	h^{-1}	Commitment rate	$0.5 - 1.0 \times 10^{-2}$	(Cao et al. 2019), (Eichner et al. 2001)	$5.41 - 7.68 \times 10^{-3}$
a_L	h	Life cycle length	48		
a_s	h	Sequestration age	20		
T_σ	h	Age at which 50% of parasites are removed during acute phase of infection	32	(Andrade et al. 2020)	
$T_{\sigma 2}$	h	Age at which 50% of parasites are removed during chronic phase of infection			33.6 - 34.7
P_σ		Shape parameter	0.7	(Andrade et al. 2020)	
$\sigma_{max}/\sigma_{max2}$	h	Ratio of maximal removal rates	0.025 - 15		2.06 - 2.89

701
702

Acknowledgments
We thank the residents of Kalifabougou, Mali and Nangonera, Uganda for participating

in the studies. We acknowledge the support of the Flow Cytometry Core Facility of DKFZ,

the Heidelberg University Metabolomics Core Technology Platform, and the Clinical

Lipidomics Unit of Mainz University Medical Center. We thank Matthias Marti and

Lauriane Sollelis at University of Glasgow for help and expertise with in vitro induction

of parasite sexual commitment and gametocytogenesis. This work was supported by the

European Research Council (ERC) under the European Union's Horizon 2020 research

and innovation programme (grant agreement No 759534), and the German Center for

Infection research (DZIF). The Malian cohort study was supported by the Division of

Intramural Research, National Institute of Allergy and Infectious Diseases of the National

Institutes of Health; and the Ugandan study was supported by the National Institutes of

Health as part of the International Centers of Excellence in Malaria Research program.

Chica and Heinz-Schaller Foundation supported Frederik Graw.

Competing Interest Statement
The authors declare that they have no conflict of interest.

References
1.   Grassi B, Bignami A, Bastianelli G. Medical Zoology: Further Researches upon the Cycle of Human Malaria in the Body of the Mosquito. Ind Med Gaz. 1899;34(3):104-7.
2.   Prudencio M, Rodriguez A, Mota MM. The silent path to thousands of merozoites: the Plasmodium liver stage. Nat Rev Microbiol. 2006;4(11):849-56.
3.   Haldar K, Murphy SC, Milner DA, Taylor TE. Malaria: mechanisms of erythrocytic infection and pathological correlates of severe disease. Annu Rev Pathol. 2007;2:217-49.
4.   Kafsack BF, Rovira-Graells N, Clark TG, Bancells C, Crowley VM, Campino SG, et al. A transcriptional switch underlies commitment to sexual development in malaria parasites. Nature. 2014;507(7491):248-52.
5.   Arnot DE, Gull K. The Plasmodium cell-cycle: facts and questions. Ann Trop Med Parasitol. 1998;92(4):361-5.
6.   Bruce MC, Alano P, Duthie S, Carter R. Commitment of the malaria parasite Plasmodium falciparum to sexual and asexual development. Parasitology. 1990;100 Pt 2:191-200.
7.   Hawking F, Wilson ME, Gammage K. Evidence for cyclic development and short-lived maturity in the gametocytes of Plasmodium falciparum. Trans R Soc Trop Med Hyg. 1971;65(5):549-59.
8.   Farfour E, Charlotte F, Settegrana C, Miyara M, Buffet P. The extravascular compartment of the bone marrow: a niche for Plasmodium falciparum gametocyte maturation? Malar J. 2012;11:285.
9.   Tiburcio M, Silvestrini F, Bertuccini L, Sander AF, Turner L, Lavstsen T, et al. Early gametocytes of the malaria parasite Plasmodium falciparum specifically remodel the adhesive properties of infected erythrocyte surface. Cell Microbiol. 2013;15(4):647-59.
10.   Smalley ME, Sinden RE. Plasmodium falciparum gametocytes: their longevity and infectivity. Parasitology. 1977;74(1):1-8.

11.   Gebru T, Lalremruata A, Kremsner PG, Mordmuller B, Held J. Life-span of in vitro
differentiated Plasmodium falciparum gametocytes. Malar J. 2017;16(1):330.
12.   Billker O, Lindo V, Panico M, Etienne AE, Paxton T, Dell A, et al. Identification of
xanthurenic acid as the putative inducer of malaria development in the mosquito. Nature.
1998;392(6673):289-92.
13.   Ross R. The role of the mosquito in the evolution of the malarial parasite: the recent
researches of Surgeon-Major Ronald Ross, I.M.S. 1898. Yale J Biol Med. 2002;75(2):103-5.
14.   Dicko A, Klion AD, Thera MA, Sagara I, Yalcouye D, Niambele MB, et al. The etiology of
severe anemia in a village and a periurban area in Mali. Blood. 2004;104(4):1198-200.
15.   Andrade CM, Fleckenstein H, Thomson-Luque R, Doumbo S, Lima NF, Anderson C, et
al. Increased circulation time of Plasmodium falciparum underlies persistent asymptomatic
infection in the dry season. Nat Med. 2020.
16.   Babiker HA, Abdel-Muhsin AM, Ranford-Cartwright LC, Satti G, Walliker D.
Characteristics of Plasmodium falciparum parasites that survive the lengthy dry season in eastern
Sudan where malaria transmission is markedly seasonal. Am J Trop Med Hyg. 1998;59(4):582-
90.
17.   Portugal S, Tran TM, Ongoiba A, Bathily A, Li S, Doumbo S, et al. Treatment of Chronic
Asymptomatic Plasmodium falciparum Infection Does Not Increase the Risk of Clinical Malaria
Upon Reinfection. Clin Infect Dis. 2017;64(5):645-53.
18.   Adomako-Ankomah Y, Chenoweth MS, Tocker AM, Doumbia S, Konate D, Doumbouya
M, et al. Host age and Plasmodium falciparum multiclonality are associated with gametocyte
prevalence: a 1-year prospective cohort study. Malar J. 2017;16(1):473.
19.   Drakeley CJ, Secka I, Correa S, Greenwood BM, Targett GA. Host haematological
factors influencing the transmission of Plasmodium falciparum gametocytes to Anopheles
gambiae s.s. mosquitoes. Trop Med Int Health. 1999;4(2):131-8.
20.   Gouagna LC, Bancone G, Yao F, Yameogo B, Dabire KR, Costantini C, et al. Genetic
variation in human HBB is associated with Plasmodium falciparum transmission. Nat Genet.
2010;42(4):328-31.
21.   Price R, Nosten F, Simpson JA, Luxemburger C, Phaipun L, ter Kuile F, et al. Risk factors
for gametocyte carriage in uncomplicated falciparum malaria. Am J Trop Med Hyg.
1999;60(6):1019-23.
22.   Tadesse FG, Slater HC, Chali W, Teelen K, Lanke K, Belachew M, et al. The Relative
Contribution of Symptomatic and Asymptomatic Plasmodium vivax and Plasmodium falciparum
Infections to the Infectious Reservoir in a Low-Endemic Setting in Ethiopia. Clin Infect Dis.
2018;66(12):1883-91.
23.   Group WGS. Gametocyte carriage in uncomplicated Plasmodium falciparum malaria
following treatment with artemisinin combination therapy: a systematic review and meta-analysis
of individual patient data. BMC Med. 2016;14:79.
24.   Barry A, Bradley J, Stone W, Guelbeogo MW, Lanke K, Ouedraogo A, et al. Higher
gametocyte production and mosquito infectivity in chronic compared to incident Plasmodium
falciparum infections. Nat Commun. 2021;12(1):2443.
25.   Williams JL. Stimulation of Plasmodium falciparum gametocytogenesis by conditioned
medium from parasite cultures. Am J Trop Med Hyg. 1999;60(1):7-13.
26.   Dyer M, Day KP. Regulation of the rate of asexual growth and commitment to sexual
development by diffusible factors from in vitro cultures of Plasmodium falciparum. Am J Trop Med
Hyg. 2003;68(4):403-9.
27.   Brancucci NMB, Gerdt JP, Wang C, De Niz M, Philip N, Adapa SR, et al.
Lysophosphatidylcholine Regulates Sexual Stage Differentiation in the Human Malaria Parasite
Plasmodium falciparum. Cell. 2017;171(7):1532-44 e15.
28.   Filarsky M, Fraschka SA, Niederwieser I, Brancucci NMB, Carrington E, Carrio E, et al.
GDV1 induces sexual commitment of malaria parasites by antagonizing HP1-dependent gene
silencing. Science. 2018;359(6381):1259-63.
29.   Thomson-Luque R, Votborg-Novel L, Ndovie W, Andrade CM, Niangaly M, Attipa C, et al.
Plasmodium falciparum transcription in different clinical presentations of malaria associates with
circulation time of infected erythrocytes. Nat Commun. 2021;12(1):4711.

30.   Cao P, Collins KA, Zaloumis S, Wattanakul T, Tarning J, Simpson JA, et al. Modeling the
dynamics of Plasmodium falciparum gametocytes in humans during malaria infection. Elife.
2019;8.
31.   Painter HJ, Carrasquilla M, Llinas M. Capturing in vivo RNA transcriptional dynamics from
the malaria parasite Plasmodium falciparum. Genome Res. 2017;27(6):1074-86.
32.   Lasonder E, Rijpma SR, van Schaijk BC, Hoeijmakers WA, Kensche PR, Gresnigt MS, et
al. Integrated transcriptomic and proteomic analyses of P. falciparum gametocytes: molecular
insight into sex-specific processes and translational repression. Nucleic Acids Res.
2016;44(13):6087-101.
33.   Lopez-Barragan MJ, Lemieux J, Quinones M, Williamson KC, Molina-Cruz A, Cui K, et al.
Directional gene expression and antisense transcripts in sexual and asexual stages of
Plasmodium falciparum. BMC Genomics. 2011;12:587.
34.   Poran A, Notzel C, Aly O, Mencia-Trinchant N, Harris CT, Guzman ML, et al. Single-cell
RNA sequencing reveals a signature of sexual commitment in malaria parasites. Nature.
2017;551(7678):95-9.
35.   Andolina C, Rek JC, Briggs J, Okoth J, Musiime A, Ramjith J, et al. Sources of persistent
malaria transmission in a setting with effective malaria control in eastern Uganda: a longitudinal,
observational cohort study. Lancet Infect Dis. 2021.
36.   Ruhamyankaka E, Brunk BP, Dorsey G, Harb OS, Helb DA, Judkins J, et al. ClinEpiDB:
an open-access clinical epidemiology database resource encouraging online exploration of
complex studies. Gates Open Res. 2019;3:1661.
37.   Hofmann NE, Karl S, Wampfler R, Kiniboro B, Teliki A, Iga J, et al. The complex
relationship of exposure to new Plasmodium infections and incidence of clinical malaria in Papua
New Guinea. Elife. 2017;6.
38.   Eichner M, Diebner HH, Molineaux L, Collins WE, Jeffery GM, Dietz K. Genesis,
sequestration and survival of Plasmodium falciparum gametocytes: parameter estimates from
fitting a model to malariatherapy data. Trans R Soc Trop Med Hyg. 2001;95(5):497-501.
39.   Tiburcio M, Dixon MW, Looker O, Younis SY, Tilley L, Alano P. Specific expression and
export of the Plasmodium falciparum Gametocyte EXported Protein-5 marks the gametocyte ring
stage. Malar J. 2015;14:334.
40.   Usui M, Prajapati SK, Ayanful-Torgby R, Acquah FK, Cudjoe E, Kakaney C, et al.
Plasmodium falciparum sexual differentiation in malaria patients is associated with host factors
and GDV1-dependent genes (vol 10, 2140, 2019). Nature Communications. 2019;10.
41.   Shanks GD. Hypothesis: dynamics of classical malaria epidemics show Plasmodium
falciparum's survival strategy. Am J Trop Med Hyg. 2015;92(3):561-4.
42.   Rono MK, Nyonda MA, Simam JJ, Ngoi JM, Mok S, Kortok MM, et al. Adaptation of
Plasmodium falciparum to its transmission environment. Nat Ecol Evol. 2018;2(2):377-87.
43.   Oduma CO, Ogolla S, Atieli H, Ondigo BN, Lee MC, Githeko AK, et al. Increased
investment in gametocytes in asymptomatic Plasmodium falciparum infections in the wet season.
BMC Infect Dis. 2021;21(1):44.
44.   Rosenberg R, Andre RG, Ketrangsee S. Seasonal fluctuation of Plasmodium falciparum
gametocytaemia. Trans R Soc Trop Med Hyg. 1990;84(1):29-33.
45.   Gadalla AAH, Schneider P, Churcher TS, Nassir E, Abdel-Muhsin AMA, Ranford-
Cartwright LC, et al. Associations between Season and Gametocyte Dynamics in Chronic
Plasmodium falciparum Infections. Plos One. 2016;11(11).
46.   Ouedraogo AL, de Vlas SJ, Nebie I, Ilboudo-Sanogo E, Bousema JT, Ouattara AS, et al.
Seasonal patterns of Plasmodium falciparum gametocyte prevalence and density in a rural
population of Burkina Faso. Acta Trop. 2008;105(1):28-34.
47.   Ouedraogo AL, Goncalves BP, Gneme A, Wenger EA, Guelbeogo MW, Ouedraogo A, et
al. Dynamics of the Human Infectious Reservoir for Malaria Determined by Mosquito Feeding
Assays and Ultrasensitive Malaria Diagnosis in Burkina Faso. J Infect Dis. 2016;213(1):90-9.
48.   Portugaliza HP, Miyazaki S, Geurten FJ, Pell C, Rosanas-Urgell A, Janse CJ, et al.
Artemisinin exposure at the ring or trophozoite stage impacts Plasmodium falciparum sexual
conversion differently. Elife. 2020;9.
49.   Orikiiriza J, Surowiec I, Lindquist E, Bonde M, Magambo J, Muhinda C, et al. Lipid
response patterns in acute phase paediatric Plasmodium falciparum malaria. Metabolomics.
2017;13(4):41.

50.   Greischar MA, Mideo N, Read AF, Bjornstad ON. Quantifying Transmission Investment in
Malaria Parasites. PLoS Comput Biol. 2016;12(2):e1004718.
51.   Prajapati SK, Ayanful-Torgby R, Pava Z, Barbeau MC, Acquah FK, Cudjoe E, et al. The
transcriptome of circulating sexually committed Plasmodium falciparum ring stage parasites
forecasts malaria transmission potential. Nat Commun. 2020;11(1):6159.
52.   Tran TM, Li S, Doumbo S, Doumtabe D, Huang CY, Dia S, et al. An intensive longitudinal
cohort study of Malian children and adults reveals no evidence of acquired immunity to
Plasmodium falciparum infection. Clin Infect Dis. 2013;57(1):40-7.
53.   Surma MA, Herzog R, Vasilj A, Klose C, Christinat N, Morin-Rivron D, et al. An
automated shotgun lipidomics platform for high throughput, comprehensive, and quantitative
analysis of blood plasma intact lipids. Eur J Lipid Sci Technol. 2015;117(10):1540-9.
54.   Lerner R, Post J, Loch S, Lutz B, Bindila L. Targeting brain and peripheral plasticity of the
lipidome in acute kainic acid-induced epileptic seizures in mice via quantitative mass
spectrometry. Biochim Biophys Acta Mol Cell Biol Lipids. 2017;1862(2):255-67.
55.   Briggs J, Teyssier N, Nankabirwa JI, Rek J, Jagannathan P, Arinaitwe E, et al. Sex-based
differences in clearance of chronic Plasmodium falciparum infection. Elife. 2020;9.
56.   Hofmann N, Mwingira F, Shekalaghe S, Robinson LJ, Mueller I, Felger I. Ultra-sensitive
detection of Plasmodium falciparum by amplification of multi-copy subtelomeric targets. PLoS
Med. 2015;12(3):e1001788.
57.   Tran TM, Guha R, Portugal S, Skinner J, Ongoiba A, Bhardwaj J, et al. A Molecular
Signature in Blood Reveals a Role for p53 in Regulating Malaria-Induced Inflammation. Immunity.
2019;51(4):750-65 e10.
58.   Miller LH, Good MF, Milon G. Malaria pathogenesis. Science. 1994;264(5167):1878-83.
59.   Klinger E, Rickert D, Hasenauer J. pyABC: distributed, likelihood-free inference.
Bioinformatics. 2018;34(20):3591-3.

Figure S1. Plasma concentration of Lyso-PC 16:0 and 18:0 of RDT[+] asymptomatic children at different times during the wet (Oct) and the dry season (Mar and May), and during clinical malaria episodes (MAL).

A

B

	Acute Phase	Transition phase	Chronic phase
Sexual commitment rate	0.54 - 0.76 %		
Gametocyte circulation time	12.04 - 22.28 days		
Parasite multiplication factor / 48 h	3.87 - 8.18	0.33 - 1.38	0.50 - 2.29
Gametocyte sequestration time	3.89 - 6.04 days		

Figure S2. A. Posterior distribution of 12 model parameters generated of 100 fits of 6th pyABC generation of one representative pyABC fitting run. **B.** Estimates of biological parameters in distinct phases of *P. falciparum* infection shown as 89% credibility interval of the posterior distributions. Sexual commitment rate was calculated by f×100% and gametocyte circulation time by 1//24 converting hours into days by the factor of 24. Parasite multiplication factor is calculated by exp(– aL) (1– f) where the term exp(– aL)(1–f) describes the fraction of surviving asexual parasites after death and sexual conversion per replication cycle. Gametocyte sequestration time was calculated by 4/m/24 where 4 indicates four sequestered state (stage I to IV) and 24 converts hours into days (Cao et al. 2019).

Figure S3. Flow cytometry gating strategy to identify and sort PfDynGFP/ PfP47 gametocytes by mCherry and GFP expression into three populations (female: mCherry-positive, male I: mCherry- GFP-positive, male II: GFP-positive).