Bacterial diet modulates tamoxifen-induced death via host fatty acid metabolism

Cédric Diot[1], Aurian P. García-González[1], Andre F. Vieira[2], Melissa Walker[1], Megan Honeywell[1], Hailey Doyle[1], Olga Ponomarova[1], Yomari Rivera[1], Huimin Na[1], Hefei Zhang[1], Michael Lee[1], Carissa P. Olsen[2], Albertha J.M. Walhout[1*]

[1]Department of Systems Biology, University of Massachusetts Chan Medical School, Worcester, MA, 01605, USA

[2]Department of Chemistry and Biochemistry, Worcester Polytechnic Institute, 100 Institute Road, Worcester, MA, 01609, USA

Corresponding author: marian.walhout@umassmed.edu

Abstract

Tamoxifen is a selective estrogen receptor (ER) modulator that is used to treat ER-positive breast cancer, but that at high doses kills both ER-positive and ER-negative breast cancer cells. We recapitulate this off-target effect in *Caenorhabditis elegans*, which does not have an ER ortholog. We find that different bacteria dramatically modulate tamoxifen toxicity in *C. elegans*, with a three-order of magnitude difference between animals fed *Escherichia coli*, *Comamonas aquatica*, and *Bacillus subtilis*. Remarkably, host fatty acid (FA) biosynthesis mitigates tamoxifen toxicity, and different bacteria provide the animal with different FAs, resulting in distinct FA profiles. Surprisingly these bacteria modulate tamoxifen toxicity by different death mechanisms, some of which are modulated by FA supplementation and others by antioxidants. Together, this work reveals a complex interplay between microbiota, FA metabolism and tamoxifen toxicity that may provide a blueprint for similar studies in more complex mammals.

Introduction

The bacteria that inhabit our body, known as our microbiota, influence many biological processes in both health and disease, and greatly contribute to our metabolic capacity [1,2]. The gut microbiota are among the first cells to encounter orally ingested nutrients and xenobiotics. Therefore, it is perhaps not surprising that the microbiota can also impact our response to medications[1]. Several mechanisms have been identified by which microbes influence the host's drug response. First, the microbial composition can change in response to drugs. Known as dysbiosis, such altered composition can affect the host's response to therapeutic drugs[3,4]. Second, bacteria can alter host drug availability, for instance by drug sequestration or modification[1,5]. Third, the metabolic crosstalk between the microbiota and its host can modify host physiology, and consequently drug efficacy[6]. All these processes can change drug action and it is therefore important to systematically identify which bacteria modulate the efficacies of which drugs, and to dissect the mechanisms involved.

Systematic identification of microbial effects on the host's drug response in humans or mammalian model systems is challenging because of their complex diet and microbiota, as well as relatively low scalability. Recently, we and others, have developed the nematode *Caenorhabditis elegans* and its bacterial diet as an interspecies model system to rapidly assess the animal's response to different drugs and how this response can be modulated by different bacteria[7-9]. The application of high-throughput genetic screens not only in the host, but also in the bacteria it consumes, make this a powerful model[4,10-12]. Recent studies illustrate the insights on host-bacteria-drug interactions that can be obtained using this model system. In a

screen with 11 chemotherapeutic drugs and two bacterial diets, we found that three

drugs, 5-fluoro-2'-deoxyuridine (FUDR), 5-fluoro-uracil (5-FU) and camptothecin (CPT)

elicited a reproductive phenotype, two of which were modulated by two bacterial

species, in opposite directions [7]. Specifically, a diet of *Escherichia coli* rendered the

animals more sensitive to FUDR, but less sensitive to CPT, relative to a diet of

Comamonas aquatica[7]. We found that bacterial pyrimidine metabolism, specifically the

generation of 5-fluorouridine monophosphate (FUMP), was critical to FUDR toxicity in

the animal[7,8].

Tamoxifen is a triphenylethylene belonging to the class of drugs known as

selective estrogen receptor modulators (SERMs). SERMs bind to and either activate or

repress the estrogen receptor (ER), a nuclear hormone receptor (NHR) transcription

factor[13]. Tamoxifen is commonly used to treat ER-positive breast cancers, as well as

other estrogen-dependent ailments such as ovarian or endometrial carcinoma [14,15]. A

significant proportion of ER-negative breast cancers also appear responsive to

tamoxifen, but the underlying biology remains to be uncovered[16]. There has also been

increasing clinical interest in the potential of tamoxifen for treating various other cancer

types[17]. The potential repurposing of tamoxifen for the treatment of non-ER-positive

breast malignancies requires an understanding of the potential alternative mechanisms

of action of this drug, especially in ER-negative models.

The *C. elegans* genome encodes more than 250 NHRs[18]. Most of these are

paralogs of HNF4α and there is no ER ortholog. We find that tamoxifen is toxic in

C. elegans at higher doses and that three bacterial species, *E. coli, C. aquatica,* and

Bacillus subtilis, confer dramatically different levels of sensitivity to tamoxifen toxicity,

with an EC_{50} ranging three orders of magnitude. We identify a role for host fatty acid

(FA) metabolism in modulating tamoxifen toxicity. We then find that the FA composition

of *C. elegans* is greatly modulated by bacterial diet, and that FA supplementation

modulates drug toxicity in a diet-specific manner. Antioxidant supplementation and

expression profiling further point to a role of lipid oxidative stress. Perhaps most

strikingly, we find that different bacteria modulate *C. elegans* tamoxifen toxicity through

different death mechanisms: on *E. coli* and *C. aquatica*, but not on *B. subtilis,* apoptosis

deficient *ced-3* and *ced-4* mutants potentiate drug toxicity. Taken together, we show

that a complex interplay between FA metabolism, oxidative stress, and differential

potentials to engage cell-death pathways, modulate tamoxifen toxicity.

Results

Tamoxifen kills both ER-positive and ER-negative cancer cells

We first confirmed that tamoxifen can kill ER-negative breast cancer cells[19]. Specifically,

we compared the effects of tamoxifen on ER-positive T-47D cells, which rely on ER for

proliferation, to the effects of tamoxifen on ER-negative MDA-MB-231 cells[20]. Drug

toxicity can be assessed through two parameters: growth rate (GR) and fractional

viability (FV). The GR index is calculated by the relative number of live cells over time in

presence or absence of drug and integrates both cytostatic and lethal effects of a

drug[21]. In contrast, FV only considers the lethal effect of a drug[22]. We first calculated the

GR index and, as expected, observed a decrease in the growth rate of T-47D cells

incubated with tamoxifen in a dose-dependent manner (**Fig. 1a**). While lower doses of

tamoxifen did not affect the growth rate of MDA-MB-231 cells, we found that

concentrations greater than 10 µM resulted in a reduction in the GR, indicating either a

reduced proliferation rate or an increased death rate (**Fig. 1a**). When we computed FV,

we found that at tamoxifen doses greater than 10 µM, killed both T-47D and MDA-MB-

231 cells effectively (**Fig. 1b**). This indicates that tamoxifen toxicity relies on two distinct

mechanisms depending on the dose: at low doses, tamoxifen slows the growth of ER-

positive, but not ER-negative cells, whereas at high doses, tamoxifen kills independently

of whether the cells express ER. Since this cell death does not depend on ER, it can be

considered an off-target effect.

Bacteria differentially modulate tamoxifen toxicity in *C. elegans*

C. elegans does not have an obvious ER ortholog[18,23]. Therefore, we wondered whether

we could use *C. elegans* to study off-target tamoxifen toxicity in an intact animal. Since

we previously found that the bacterial diet consumed by *C. elegans* can greatly affect

the response to chemotherapeutic drugs[7], we supplemented increasing doses of

tamoxifen to larval stage 1 (L1)-arrested animals fed either of three different bacterial

diets: *E. coli, C. aquatica* or *B. subtilis*. Visual inspection showed dramatic differences in

tamoxifen toxicity, depending on bacterial diet: there was little effect on animal

development on *E. coli*, while animals fed *C. aquatica* displayed developmental arrest or

delay at high drug concentrations, and animals fed *B. subtilis* were exquisitely sensitive

to tamoxifen, even at low micromolar concentrations (**Fig. 1c** and **Supplementary Fig.**

1). We next used L1 arrest as a proxy for tamoxifen toxicity and plotted the proportion of

animals that failed to develop after incubation on tamoxifen-containing plates for 48

hours as a function of the drug concentration. The resulting dose response curves

(DRCs) confirmed that tamoxifen is more than three orders of magnitude more toxic to

animals fed *B. subtilis* than to animals fed *E. coli* (**Fig. 1d**). Notably, bacterial lawns on

NGM plates were not affected by the presence of tamoxifen (**Supplementary Fig. 2**).

Together, these results show that tamoxifen is toxic to *C. elegans* at high doses and

that bacterial diet greatly modulates this toxicity.

Bacteria modulate drug bioavailability

To gain insight into the mechanism by which bacteria modulate tamoxifen toxicity in

C. elegans, we performed genetic screens in both *E. coli and C. aquatica*. Specifically,

we used the *E. coli* Keio mutant collection, which contains 3,985 single-gene deletion

mutant strains[24], and a *C. aquatica* mutant collection of 5,760 strains that we previously

generated by transposon-based mutagenesis[11] (**Fig. 2a**). To enable the identification of

bacterial mutants that either increase or decrease tamoxifen toxicity in *C. elegans*, we

screened each bacterial mutant collection with two doses of tamoxifen: a toxic dose

(200 μM and 300 μM on animals fed *C. aquatica* and *E. coli,* respectively), and a non-

toxic dose (100 μM) on which animals develop on either diet. We did not identify any *E.*

coli mutants that reproducibly altered drug toxicity in *C. elegans*. This indicates that

active *E. coli* metabolism is unlikely to play a dominant role in modifying tamoxifen

toxicity in the animal. However, we did find four *C. aquatica* mutants that changed the

animal's response to tamoxifen, two of which decreased and two that increased toxicity

(**Fig. 2b**). The two mutants that increased the severity of tamoxifen toxicity harbor the

transposon insertion in the *tadC* gene*,* which encodes a component of the type II

secretion system[25], and in the *exbD* gene, which encodes a protein involved in

TonB-dependent transport[26]. The two mutants that decreased tamoxifen toxicity harbor

the transposon in a gene encoding a hypothetical protein with no annotated functional

domains and in the *acrR* gene, which encodes a transcriptional repressor of the *acrA/B*

multidrug efflux pump[27](**Fig. 2c**). This pump is associated with efflux of hydrophobic

xenobiotics[28]. Since we identified a transcriptional repressor of this pump, this may

suggest that the pump removes tamoxifen from the bacteria. These results suggest that

tamoxifen transport, rather than active bacterial metabolism, affects the amount of drug

taken up by *C. aquatica,* and, therefore, drug bioavailability in *C. elegans* fed this

bacterial diet. To test this, we measured tamoxifen accumulation in wild-type and

mutant strains of *C. aquatica* exposed to the drug by gas chromatography - mass

spectrometry (GC-MS). We found that the four *C. aquatica* mutants harbor different

levels of tamoxifen, and that these levels correlate with tamoxifen toxicity in *C. elegans*

(**Fig. 2d**).

Next, we compared tamoxifen levels in the three bacterial species, and in

animals fed these different diets. We found that bacteria that confer increased toxicity

accumulate 5-10-fold more tamoxifen, which translates to an 5-10-fold increased drug

accumulation in *C. elegans* (**Fig. 2e**, **f**)*.* However, these differences do not explain the

dramatic tamoxifen toxicity in animals fed *B. subtilis.*

Taken together, different bacterial species deliver different amounts of tamoxifen

to *C. elegans*, indicating that bioavailability affects, in part, toxicity in the animal.

However, these effects are not sufficient to explain the large differences in toxicity

depending on bacterial diet, nor do they provide insight into the mechanisms of

tamoxifen toxicity in *C. elegans.*

C. elegans fatty acid synthesis protects against tamoxifen toxicity

Because differences in drug bioavailability are not sufficient to explain the differences in tamoxifen toxicity in animals fed the three different bacteria, and since tamoxifen is only toxic to *C. elegans* fed *E. coli* at high micromolar doses (**Fig. 1d**), we hypothesized that *C. elegans* metabolism may affect drug toxicity. To test this, we used RNAi of 1,495 *C. elegans* metabolic genes to identify genes that increased or decreased drug toxicity when knocked down. These include genes with known metabolic functions that make up the genome-scale iCEL1314 metabolic network model, as well as additional genes predicted to encode metabolic enzymes[29]. We did not find any genes that, upon RNAi knockdown, rendered the animals less sensitive to tamoxifen. We did, however, identify three genes for which RNAi knockdown led to increased tamoxifen toxicity (**Fig. 3a**). These genes include *dhs-19,* which encodes a short-chain dehydrogenase of unknown function that localizes to lipid droplets [30] and is predicted to function in retinol metabolism [31]. The other two genes, *elo-3* and *elo-6,* both encode elongases predicted to be involved in long chain FA biosynthesis [32,33]. We generated a *dhs-19* deletion mutant by CRISPR/Cas9 genome editing and confirmed that this strain is indeed more sensitive to tamoxifen (**Fig. 3b** and **Supplementary Fig. 3**).

While mutant strains are available for *elo-3* and *elo-6*, we could not use these animals because their development is strongly impaired even in absence of drug exposure (data not shown). Both *elo-3* and *elo-6* function in FA biosynthesis (**Fig. 3c**). We therefore wondered if these two genes specifically affect tamoxifen toxicity, or if FA metabolism is more broadly involved even though no other genes were captured in the

RNAi screen. To test this, we performed full tamoxifen DRC experiments for the

knockdown of 33 genes encoding enzymes predicted to be involved in FA elongation

and desaturation[33] (**Fig. 3c**). In addition, we included eight genes encoding acyl-CoA

synthetases that catalyze the addition of co-enzyme A (CoA) to FAs and eight genes

encoding enzymes that remove CoA. We calculated the difference between the area

under the curve (ΔAUC) for each RNAi condition + tamoxifen relative to *E. coli* (**Fig. 3d**).

Statistical significance was evaluated on the original DRCs (**Supplementary Fig. 4**, and

Supplementary Tables 1, 2). Remarkably, knockdown of most FA biosynthesis genes

tested significantly increased tamoxifen toxicity (**Fig. 3e, Supplementary Fig. 4,** and

Supplementary Table 1). These results show that *C. elegans* FA synthesis is broadly

involved in mitigating tamoxifen toxicity. This points to a systems-level involvement of

FA biosynthesis rather than a single FA species.

Next, we asked whether genes involved in mitochondrial and peroxisomal FA

degradation also modulate tamoxifen toxicity and found that, of 33 genes tested, only

ech-8 appeared to increase tamoxifen toxicity, but this effect was not statistically

significant (**Fig. 3f, Supplementary Fig. 5**). Knockdown of 13 of the 33 FA degradation

genes tested led to a decrease in tamoxifen toxicity, with only *daf-22* RNAi being

statistically significant (**Supplementary Table 2**). This result shows that the

biosynthesis, but not degradation, of long chain FAs is critical to mitigate tamoxifen

toxicity in *C. elegans.*

Bacterial diets modulate *C. elegans* fatty acid profiles

Previously, it has been shown that different strains of *E. coli* diets affect the FA

composition of *C. elegans*[34]. Since *C. elegans* FA synthesis modulates tamoxifen

toxicity, we hypothesized that the three bacterial species used herein may elicit different

FA profiles in the animal. To test this, we measured both bacterial and *C. elegans* FA

profiles by GC-MS (**Supplementary Tables 3**, **4**). The bacterial FA profiles showed

dramatic differences: *E. coli* mainly contained the saturated FAs (SFAs) palmitic acid

and stearic acid, cyclopropane FAs (CPFAs), and the mono-unsaturated FAs (MUFAs)

oleic acid, palmitoleic acid and cis-vaccenic acid; *C. aquatica* mainly contained the

MUFAs oleic acid and palmitoleic acid; and *B. subtilis* almost exclusively contains

monomethyl branched chain FAs (mmBCFAs), which agrees with previous findings[35]

(**Fig. 4a**).

The FA profiles of *C. elegans* fed the three bacteria reflected the three bacterial

diets in some but not all respects. One striking observation is that the proportion of each

of the different poly-unsaturated FAs (PUFAs) is diet-independent (**Fig. 4b**). This

indicates that the animal can effectively convert different dietary FAs into PUFAs by

elongation and/or desaturation. Oleic acid is the primary substrate for PUFA synthesis

and is present in *E. coli* and *C. aquatica* but absent in *B. subtilis* (**Fig. 4a**). This could

indicate that PUFAs are generated by a combination of FA metabolic steps that yield

this even-chained MUFA from odd-chained mmBCFAs in animals fed *B. subtilis* or

synthesizes them de novo.

Another striking observation is that most other types of FAs are quite different in

proportion in *C. elegans* fed different bacterial diets. Most notable are the near absence

of MUFAs, and presence of C15 and C17-ante-iso mmBCFAs in animals fed *B. subtilis*

(**Fig. 4b**). The differences between animals fed the other two bacteria are more subtle,

with a greater proportion of the MUFA cis-vaccenic acid in *C. aquatica*-fed animals,

relative to those fed *E. coli* (**Fig. 4b**). Altogether, the different bacterial diets result in a

different proportion of the different types of FAs: high overall levels of MUFAs in animals

fed *E. coli* or *C. aquatica* and high overall levels of mmBCFAs in animals fed *B. subtilis*

(**Fig. 4b**, bottom). Taken together, these results show that *C. elegans* maintains its

PUFA content in a narrow and well-defined regime, while it can tolerate different

proportions of the other types of FAs.

Dietary fatty acids modulate tamoxifen toxicity

Our findings so far lead us to consider two models: either tamoxifen treatment could

alter the animal's FA profile, leading to diet-dependent toxicity, or differences in *C.*

elegans FA composition elicited by bacterial diet may modulate tamoxifen toxicity. To

discriminate

between these two models, we first measured bacterial and *C. elegans* FAs in the

presence or absence of tamoxifen and found that the drug did not modulate the animal's

FA profiles (**Supplementary Table 5**). Thus, we did not find evidence to support the

first model.

To evaluate the second model, we first asked whether supplementation of a FA

cocktail, containing equimolar amounts of the PUFAs arachidonic and linoleic acid, the

SFAs lauric and myristic acid, and the MUFA oleic acid[36], would modulate tamoxifen

toxicity in animals fed either of the three bacterial diets. We found that FA cocktail

supplementation suppressed tamoxifen toxicity but only in animals fed *C. aquatica*; it

had no effect on tamoxifen toxicity in animals fed *E. coli* or *B. subtilis* (**Fig. 5a-c**). Next,

we tested the effects of the supplementation of individual FAs and found that MUFAs

such as cis-vaccenic acid increased tamoxifen toxicity on animals fed *E. coli*, but not *C.*

aquatica (**Fig. 5d**, **e**, and **Supplementary Fig. 6**). Since *C. aquatica*-fed animals

contain more cis-vaccenic acid than *E. coli*-fed animals (33% vs 20%, **Fig. 4b**), this

suggests that this difference may contribute to increased tamoxifen toxicity on the *C.*

aquatica diet.

268   Supplementation of individual PUFAs such as α-linoleic acid decreased

tamoxifen toxicity on animals fed *C. aquatica* (**Fig. 5f**, **g,** and **Supplementary Fig. 6**).

On an *E. coli* diet PUFAs had no effect. However, animals are generally tolerant to

tamoxifen, which limits the ability to observe decreased tamoxifen toxicity. Since the

proportion of PUFAs is similar in animals fed each of the three bacterial diets (**Fig. 4b**),

this indicates that the ratio of PUFAs to other FAs is more relevant than absolute levels.

In contrast to *E. coli* and *C. aquatica*, supplementation of either MUFAs or PUFAs

abolished the growth of *B. subtilis*, and therefore these FA could not be tested for

tamoxifen toxicity modulation in *C. elegans* fed these bacteria (**Fig. 5h**, **i** and

Supplementary Fig. 7). Together with the observation that knockdown of genes

involved in FA synthesis increases tamoxifen toxicity, these results point to a complex

interplay between dietary FA, FA synthesis and tamoxifen toxicity.

280   We next asked whether bacterial metabolism was required for the effects of

supplemented FAs on tamoxifen toxicity. We generated metabolically inactive bacterial

powders by mechanical lysis followed by lyophilization as previously described[7], and

seeded these powders onto tamoxifen-containing plates. We observed that the

dietary-dependent effects were maintained: *E. coli* powder-fed animals developed in

presence of 600 μM tamoxifen, while animals fed *C. aquatica* powder did not. Similar to

feeding live *C. aquatica*, supplementation of FA cocktail allowed animals to develop in

presence of 600 μM tamoxifen when fed metabolically inactive *C. aquatica*

(**Supplementary Fig. 8**). *B. subtilis* powder did not support *C. elegans* development

and therefore could not be tested for tamoxifen toxicity (**Supplementary Fig. 8**). These

results indicate that active bacterial metabolism does not affect tamoxifen toxicity in *C.*

elegans, which is also supported by the lack of metabolic gene mutants identified in the

bacterial mutant screens. Instead, these observations further support the idea that

different bacteria provide the animal with different FAs and that these FAs differently

modulate tamoxifen toxicity.

E. coli-fed animals harbor high levels of CPFAs, while *B. subtilis*-fed animals

contain high levels of mmBCFAs (**Fig. 4b**). Therefore, we asked whether these FAs

could modulate tamoxifen toxicity in *C. elegans*. We used bacterial genetics to assess

the importance of CPFAs because they are exclusively synthesized in bacteria by CFA

enzymes[37]. Because CPFAs specifically accumulate in *E. coli* (**Fig. 4b**)*,* and because

this diet renders the animals less sensitive to tamoxifen toxicity, we reasoned that

CPFAs may be protective. We fed *C. elegans* two independent *E. coli* Δ*cfa* strains from

the Keio library [24] and did not observe and effect on tamoxifen toxicity (**Supplementary**

Fig. 9). Next, we tested whether mmBCFAs, which specifically accumulate in *B. subtilis*,

would enhance tamoxifen toxicity. mmBCFAs occur in two types, iso-mmBCFAs and

ante-iso-mmBCFAs[38]. We were not able to test mmBCFAs by supplementation due to

cost and supply issues. However, because ante-iso mmBCFAs are synthesized from

isoleucine supplied in growing media, we could assess their contribution to tamoxifen

toxicity by growing *B. subtilis* in the absence of isoleucine[38]. There was no difference in

tamoxifen toxicity in animals fed *B. subtilis* grown in LB (rich in isoleucine) or minimal

M9 media supplemented with glucose as carbon source (**Supplementary Fig. 10**).

While not unequivocally ruling out the contribution of mmBCFAs, these data show that a

switch from ante-iso to iso-mmBCFAs in the food uptake does not modulate tamoxifen

toxicity.

Previously, it has been shown that tamoxifen inhibits glucosylceramide synthesis

in human cells[39]. In *C. elegans*, mmBCFAs are converted into the sphingolipid d17-

isoglucosylceramide by ELO-5, and this conversion is important to support development

and growth [40]. Since we found that *elo-5* RNAi enhances tamoxifen toxicity (**Fig. 3e**), we

next asked whether tamoxifen elicits a d17-isoglucosylceramide deficiency that explains

drug toxicity. In *C. elegans*, knockdown of *elo-5* can be rescued by combining it with a

mutation in components of the NPRL2/3 complex[40] (**Supplementary Figure 11**). If

tamoxifen toxicity is due to the lack of d17-isoglucosylceramide, we would expect that

mutants in the NPRL-2/3 complex would mitigate toxicity. However, we found that *nprl-3*

deletion mutants showed equal sensitivity to tamoxifen as wild type animals

(**Supplementary Fig. 11**). This result shows that tamoxifen does not affect *C. elegans*

by depletion of d17-isoglucosylceramide.

How do FAs and FA biosynthesis modulate tamoxifen toxicity in *C. elegans*?

Since

FAs are easily oxidized and rapidly turnover in cellular and organellar membranes[41], we

hypothesized that tamoxifen treatment causes FA oxidation, and that this oxidation is

mitigated by replenishing oxidized FAs with newly synthesized FAs. Vitamin E is a

potent antioxidant that inhibits FA oxidation and that stabilizes the FA cocktail[42]. We

found that vitamin E had no effect on tamoxifen toxicity in animals fed either *E. coli* or *C.*

aquatica, indicating that it is the FAs in the FA cocktail that mitigate toxicity in animals

fed *C. aquatica*. However, vitamin E supplementation did suppress tamoxifen toxicity in

animals fed *B. subtilis* (**Fig. 5j-l**), when N-acetylcysteine (NAC), another antioxidant that

can rescue the growth of *C. elegans* on iron-deficient *E. coli* mutants[43] did not affect

tamoxifen toxicity (**Fig. 5m**). These results indicate that FA oxidation, but not the

general accumulation of reactive oxygen species (ROS), affects tamoxifen toxicity in

C. elegans when fed *B. subtilis*. Taken together, these results indicate that FA or

vitamin E supplementation can mitigate tamoxifen toxicity differently depending on

which bacteria the animals are fed, which suggests that different, diet-dependent toxicity

mechanisms are involved.

Tamoxifen induces the expression of type II detoxification genes

We next used expression profiling by RNA-seq to ask whether tamoxifen

supplementation modulates the expression of genes involved in FA metabolism or

oxidative stress in *C. elegans*. To avoid capturing mRNA changes due to differences

developmental rate, we used a tamoxifen dose at which no developmental arrest was

observed, but in which animals are delayed by ~12 hours. The development of animals

exposed to tamoxifen was monitored hourly, and samples were collected 12h after

control animals. A total of 558 and 738 genes were up- and down regulated,

respectively (>2-fold change, P< 0.01, **Supplementary Table 6**). Tamoxifen

supplementation did not change the expression of genes involved in FA biosynthesis or

degradation genes with the exceptions of *fat-5,* which increased by three-fold, and *ech-*

6 which was reduced by ~two-fold (**Fig. 6a**). Together with the observation that

tamoxifen supplementation did not alter the animal's FA profiles, these results show that

tamoxifen does not modulate FA metabolism. Instead, it suggests that the animal's FA

state modulates tamoxifen toxicity.

Several genes that change in expression in response to tamoxifen

supplementation pointed to potential mechanisms of toxicity. First, tamoxifen induced

the expression of R05D8.9 (38-fold, adjusted *p*-value: 1.2×10^{-49}), which encodes a

predicted 17beta-hydroxysteroid dehydrogenase 14 (17βHSD14) homolog (**Fig. 6a** and

Supplementary Table 6). In humans, 17βHSD14 enzymes are associated with the

response to tamoxifen in breast cancer[44]. These enzymes are further known to

interconvert different steroids. We tested whether the induction of this gene is functional

and found that R05D8.9 knockdown increased tamoxifen toxicity (**Fig. 6b**). This

observation suggests that R05D8.9 is involved in the detoxification of tamoxifen. The

expression of several CYP genes (*cyp-35B3, -35B2, -35B1,* and *-34A7*), which encode

cytochrome P450 enzymes involved in type II detoxification, is induced by tamoxifen

supplementation (179-, 55-, 29-, and 10-fold, adjusted *p*-values: 1.6×10^{-15}, 9.2×10^{-16},

9.2×10^{-16} and 4.1×10^{-4}) (**Fig. 6a** and **Supplementary Table 6**). Cytochrome P450

enzymes have been reported to modify tamoxifen in humans[45], and are a source of

oxidative stress as their enzymatic cycle involves significant leakage of electrons [46].

Again, we tested whether these expression changes are functional and found that

knockdown of *cyp-35B3* increased drug tamoxifen toxicity, while knockdown of *cyp-*

35B1 decreased toxicity at high doses (**Fig. 6c**, **d**). These results demonstrate the

functional involvement of CYP-450 enzymes in detoxifying tamoxifen toxicity, and

further suggest, that different CYP-450 proteins can exhibit differential tamoxifen-

detoxifying and ROS-generating potentials, which is in agreement with previous

observations[45,47]. Taken together, tamoxifen does not transcriptionally rewire FA

metabolism. Instead, it induces the expression of ROS-generating detoxifying enzymes,

which may contribute to FA oxidation.

Bacteria elicit different tamoxifen-induced death mechanisms

Lipid metabolism has been linked to different mechanisms of cell death, especially

apoptosis and ferroptosis[48,49]. In fact, it has been shown that cell survival is dependent

on lipid homeostasis, and both excessive MUFAs or SFAs have been shown to be pro-

apoptotic, depending on the cell lines[49]. Further, several MUFAs have been reported to

protect against ferroptosis[50], which is characterized by lipid peroxidation, while others,

specifically cis-vaccenic acid have been reported to induce cell death[51]. PUFAs, most

notably di-homo-gamma-linoleic acid, have been shown to induce ferroptosis in both

human cells and in *C. elegans*[52]. Given these connections, we asked whether apoptosis

contributes to tamoxifen toxicity in *C. elegans*. To test this, we examined tamoxifen

toxicity in *ced-3* and *ced-4* mutants that are deficient in apoptosis[53]. CED-3 is a caspase

that is essential for apoptosis, and CED-4 is an activator of CED-3. Remarkably, loss of

ced-3, and to a lesser extent *ced-4,* greatly increased sensitivity to tamoxifen in animals

fed either *E. coli* or *C. aquatica* but had no effect on animals fed *B. subtilis* (**Fig. 6e-g**

and **Supplementary Fig. 12**). This result suggests that, when fed *E. coli* or *C. aquatica,*

tamoxifen toxicity employs apoptosis, and that *ced-3* and *ced-4* mutant animals switch

to another, more potent mode of death that is suppressed when apoptosis is functional.

We found that a high dose of vitamin E could mitigate tamoxifen toxicity in *ced-3* mutant

animals fed *C. aquatica* but not *E. coli* (**Supplementary Fig. 12**). This indicates that, in

the absence of *ced-3*, tamoxifen toxicity in animals fed *C. aquatica* involves FA

oxidation, similar to what we observed with wild type animals fed *B. subtilis*.

Finally, we tested the effects of Vitamin E and FA cocktail on a dose of tamoxifen

that is toxic independently of ER in human cells (31 µM, **Fig. 1a, b**). Tamoxifen toxicity

was severely suppressed by either supplement in both ER-negative MDA-MB-231 and

ER-positive cells (**Fig. 6h**, **Supplementary Fig. 13**). Furthermore, ZVAD, a pan-

caspase inhibitor that prevents apoptosis, reproducibly lowered the drug toxicity in

MDA-MB-231 but not in T-47D cells. This suggests that, like in *C. elegans*, the ER-

independent toxicity of tamoxifen can be elicited through different cell death pathways in

human cells.

Discussion

In this study, we present *C. elegans* as a model to study off-target, ER-independent,

mechanisms of tamoxifen toxicity using concentrations in the range of drug-levels

accumulating in patients[54,55]. We found that FA biosynthesis and composition of the

animals are major drivers of tamoxifen toxicity, and that this toxicity relies on the

potential of the drug to induce oxidative stress and lipid peroxidation. Moreover, we

found that both the mechanism and the degree of tamoxifen toxicity are greatly

modulated by bacteria, where *B. subtilis* induces a non-apoptotic mechanism that is

three orders of magnitude more severe than apoptotic mechanism elicited by tamoxifen

in the presence of either *E. coli* or *C. aquatica* (**Fig. 6**).

In the *C. elegans* model, bacteria form a source of nutrition, can be pathogenic,

and can inhabit the gut as the animal ages[56]. Both in the laboratory and in the wild, the

animals are constantly exposed to bacterial metabolism which provides nutrients and

contributes to animal physiology[57]. Our findings illuminate the possible effects the

microbiota may have on tamoxifen off-target effect toxicity and the treatment of ER-

independent malignancies[17]. It is tempting to speculate that tamoxifen activity could be

modulated by perturbations of the microbiota using antibiotics or probiotics. Notably,

B. subtilis, which is present in a large proportion of commercially available probiotics[58]

may affect patients treated with tamoxifen. Our findings of the role of FA metabolism

and oxidative stress further suggest that studies of nutritional intake of fats and vitamin

E in humans may shed light on whether the off-target mechanisms discovered herein

are conserved.

We uncovered complex relationships between bacteria, host FA metabolism,

apoptosis, and tamoxifen toxicity. The literature about the relationship between FAs and

cell survival integrates two major components: distinct FA-species display different

oxidation potentials[59], and different potentials to induce alternative cell-death

mechanisms[49,50]. Our data using lipophilic ROS-scavenging vitamin E and apoptosis-

deficient animals suggest that tamoxifen-toxicity relies on these two entangled

components. MUFAs are generally described as 'healthy' FAs, as they buffer the

oxidative stress of ROS-sensitive PUFAs[50]. To our knowledge, there is only one report

on the effects of the MUFA cis-vaccenic acid, which found that its accumulation induces

cell death, when other MUFAs tested in parallel do not[51]. Overall, our observation of

extreme tamoxifen toxicity in *B. subtilis*-fed animals that do not provide the animal with

MUFAs, and in *C. aquatica*-fed animals that accumulate cis-vaccenic acid are

consistent with the literature. Further studies are required to systematically determine

the effects of different FAs and lipids on cell survival and tamoxifen toxicity in human

cells.

Knockdown of or mutations in *dhs-19* enhance tamoxifen toxicity in animals fed

E. coli. The precise molecular or metabolic function of DHS-19 is unknown. It encodes a

dehydrogenase and is a marker of lipid droplets[60], which are essential for FA

homeostasis in all eukaryotes[61]. Based on enzyme homology, DHS-19 is predicted to

act in retinoid metabolism, through the interconversion of retinol and retinal[33]. This

reaction precedes the synthesis of retinyl palmitate which consumes palmitoyl-CoA,

which is a precursor for the synthesis of palmitoleic and vaccenic acid, both of which

potentiate tamoxifen toxicity in *C. elegans* when animals are fed *E. coli*. Retinoids are

also potent ROS scavengers and have been shown to act synergically with vitamin E to

reduce lipid oxidative stress[62]. Together, this suggests that the increase in tamoxifen

toxicity by *dhs-19* perturbation could result from a complex integration of metabolic cues

involving oxidative stress and retinoid metabolism.

C. elegans is a remarkable animal with a fixed lineage in which wild type animals

develop deterministically to adults with precisely 959 somatic nuclei[63,64]. Therefore, one

would assume that all adult animals are essentially the same. Here we show that wild

type animals can exhibit marked differences in their FA composition, depending on

which bacterial diet they are fed. To our knowledge, *C. elegans* FA composition has

only been studied in animals fed *E. coli*, and the profiles observed in our experiments

are consistent with the only available previous study[34]. Here, we observe much more

drastic changes in FA composition in animals fed *C. aquatica* or *B. subtilis*, relative to

those fed *E. coli*. First, the MUFA, mmBCFA or CPFAs content of the animals strongly

reflects the dietary input; a diet of *C. aquatica* is rich in MUFAs, and a diet of *B. subtilis*

is rich in mmBCFAs, and these FAs accumulate in the animals. This result

demonstrates that the animal's membranes are permissive to varying levels of these

types of FAs. In contrast, the PUFA composition of *C. elegans* is largely independent of

its bacterial diet. While PUFAs are essential dietary FAs in humans, *C. elegans*

synthesize PUFAs de novo[65]. This result indicates that *C. elegans* PUFA levels are

sensed, and that metabolic fluxes in FA biosynthesis pathways are adjusted to maintain

relative levels in a tight regime. It is therefore perhaps not surprising that PUFA

deficiencies have been associated with developmental and neuronal defects in *C.*

elegans[66]. Future studies are required to reveal to what the extent wild type *C. elegans*

exhibit other fitness changes in response to differences in FA composition.

How does cell-death connect to organism death? Fifty years after Horvitz and

Sulston first described apoptosis in *C. elegans*[63], it remains unclear how cell death

mechanisms communicate and how they relate to organismal phenotypes including

death. Other open questions that remain to be answered are: What is the exact nature

of the triggers directing decisions in the induction of alternative cell-death mechanisms?

And how do bacteria affect cell-death potentials? This study provides a foundation for

future studies to address fundamental questions related to cell-death and organismal

fate.

Methods

C. elegans

C. elegans N2 (Bristol) was used as the wild-type strain. Prior to all experiments, animals were maintained on Nematode Growth Media (NGM) and fed a diet of *E. coli* OP50. To prepare L1-synchronized animals, eggs purified by dissolving gravid animals in NaOH-buffered bleach were collected and incubated for 20h in M9 buffer for hatching. The *C. elegans ced-3(n717)* mutant strain was kindly provided by the Francis lab (UMass Medical School) and was first described in (8598288). The *C. elegans dhs-19(VL1313)* mutant strain was generated using CRISPR-Cas9 genome editing[67].

Bacterial strains

E. coli BW25113, *E. coli* HT115, *C. aquatica* DA1877, and *B. subtilis* sp168 were grown in LB, overnight at 37°C to stationary phase under agitation (200 rpm). Alternatively, *B. subtilis* sp168 was grown in M9 at 37°C to stationary phase (~24h). The dietary effects on tamoxifen toxicity described herein were independent of the *E. coli* strain tested. Experiments were conducted using *E. coli* BW25113, unless stated otherwise. Bacterial powders were generated by mechanical lysis followed by desiccation as previously described[7]. Powders were seeded onto tamoxifen containing NGM plates supplemented antibiotics and prepared without peptone.

Bacterial genetic screens

Bacterial genetic screens were performed in 96-well plates containing NGM

supplemented with 200 and 300 µM tamoxifen for *E. coli* mutants, 100 and 300 µM

tamoxifen for *C. aquatica* mutants. Wells were seeded with individual mutant strains,

and approximately twenty L1-arrested animals were added to each well on the following

day. After 2 and 3 days at 20°C, hits were scored visually using a dissecting microscope

as bacterial mutants leading to a difference in animal drug response relative to

wildtype-fed animals. Primary hits were then retested in 48-well plates, in two

independent experiments, on NGM supplemented with 0.5% DMSO, 50, 100, 150, 300,

or 600 µM tamoxifen for *E. coli* and *C. aquatica* mutants. Bacterial mutants that showed

robust differences on host drug response were genotyped. *E. coli* and *C. aquatica* and

mutants were genotyped using gene-specific primers flanking -200bp and +200bp

relative to the start codon and semi-random two step PCR, respectively.

Tamoxifen-containing NGM plate preparation

Tamoxifen citrate (Selleckchem, S1972) was dissolved to 150 mM in sterile dimethyl

sulfoxide (DMSO), aliquoted and stored at -20°C. Stock solutions were diluted in series

to 200X the final concentration and added at a final volume of 0.5% to NGM agar

heated to 55°C. Drug containing NGM plates were dried overnight at room temperature.

NGM was supplemented with antibiotics for experiments in 96-well plate format, and 2

mM IPTG for RNAi experiments. 96-well and 48-well plates were seeded with 15 µL and

30 µL of bacterial cultures, respectively. Plates were then dried for 6h at room

temperature and kept overnight at room temperature before being used. *E. coli* cultures

were concentrated 1-2X, and *C. aquatica* DA1877 and *B. subtilis* sp168 strains were

concentrated 2-4X prior to seeding onto NGM agar plates, to normalize for post-seeding

bacterial growth.

Importantly, batch-to-batch effects were observed as the study was conducted

with differences in drug toxicity depending on the batch. However, dietary differences on

tamoxifen drug toxicity were systematically observed.

Tamoxifen measurements

Tamoxifen levels were measured on an Agilent 7890B/5977B single quadrupole GC-MS

equipped with an HP-5ms Ultra Inert capillary column. Briefly, 5,000 animals were

mechanically lysed in 80% methanol using a FastPrep-24 bead beater (MP

Biomedicals). Lysates were centrifuged for 10 min at 20,000 × g, and 250 µL of the

supernatants were dried using a SpeedVac concentrator SPD111V (Thermo Fisher

Scientific). Derivatization was performed by incubating the extracts with methoxyamine

hydrochloride   (Sigma-Aldrich)   [37°C - 1.5h],   and   then   with   N-methyl-N-

(trimethylsilyl)trifluoroacetamide (Sigma-Aldrich) [37°C, 3h and then 25°C, 5h]. Peak

areas were quantified using samples within a linear response range, after mean

normalization to total metabolites and blank subtraction. Sample size biases were then

corrected by normalizing tamoxifen signals to the metabolomic weight of the sample

they originate from (i.e., the sum of all 23-metabolites measured in the run of interest).

For bacteria, lawns were recovered 24h post-seeding, which corresponds to the

time at which animals are exposed to tamoxifen in the experiments presented in this

manuscript. For *C. elegans*, animals were first grown to the young-adult stage in

absence of tamoxifen, on *E. coli*, *C. aquatica* or *B. subtilis*. They were then transferred

onto tamoxifen containing plates, seeded with the same bacteria 24h before. Animals

were recovered after 12h of exposure.

RNAi screen

We tested a collection of 1,645 individual *E. coli* HT115 strains expressing dsRNA

against a unique *C. elegans* metabolic gene[29]. The screens were performed using 96-

well plates containing NGM supplemented with tamoxifen, 50 µg/mL ampicillin, and 2

mM IPTG. We used a low dose of 300 µM tamoxifen to identify knockdowns that

aggravate tamoxifen toxicity, and a high dose of 600 µM tamoxifen identify knockdowns

that mitigate tamoxifen toxicity. Approximatively 20 L1-arrested animals were added to

each well. Two days later, hits were scored visually using a dissecting microscope as

knockdowns leading to a difference in animal drug response relative to animals fed an

E. coli strain containing the empty dsRNA vector backbone (pL4440).

C. elegans **dose-response curves**

Dose-response curves (DRCs) were performed in 48-well plates with six to twelve

technical replicates. Fifty to one hundred L1-arrested animals were plated onto each

well and incubated at 20°C for 48 hours. Animals that developed beyond the L1 stage

were counted under a dissecting microscope. The percentage of animals that did not

developed was calculated relative to the total number of animals developed in the

control condition. For ΔAUCs in **Fig. 3d-f**, we used the average AUCs from the DRCs of

the *E. coli* BW25113 and *E. coli* HT115 pL4440 control conditions.

Bacterial growth curves

Bacteria were first grown overnight in LB at 37°C and shaking at 200 rpm. Next, bacterial cultures were diluted 1:200 and grown in LB supplemented with 1% of FA supplements (or ethanol vehicle as a control). Cultures were then grown for 20h at 37°C and shaking at 200 rpm, in an Eon™ Microplate Spectrophotometer (BioTek) and absorbance at 600 nm was measured every 900 s.

Fatty acid and vitamin E supplementations

The FA cocktail contains equimolar amounts of the PUFAs arachidonic and linoleic acid, the SFAs lauric and myristic acid, and the MUFA oleic acid[36] (Sigma-Aldrich, F7050). Undiluted fatty acid cocktail and PUFAs solutions were supplemented at 1% v.v. in LB for overnight growth. MUFAs were first diluted 1:10 in ethanol and supplemented at 1% v.v. in LB for overnight growth. Undiluted vitamin E was supplemented from 0.1% to 10% in LB for overnight growth. MUFAs references: cis-vaccenic acid (VA; Cayman chemicals, 20023), palmitoleic acid (PA; Fisher Scientific, AC376910010), oleic acid (Sigma-Aldrich, 364525). PUFAs references: α-Linolenic Acid (ALA; Cayman chemicals, 90210), γ-Linolenic Acid (GLA; Cayman chemicals, 90220), dihomo-γ-Linolenic Acid (DGLA; Cayman chemicals, 90230). Antioxidants: Vitamin E (Sigma-Aldrich, T3251), N-acetyl-L-cystein (NAC; Sigma-Aldrich, A7250).

C. elegans **FA measurements**

Fatty acid measurements in bacteria and animals were performed as described before[68]. Briefly, lipids were extracted from whole animals using a 2:1

chloroform:methanol mixture and dried in liquid nitrogen. A calibrated phospholipid or

triacylglycerol standard was added to the extracted lipids before separation by

solid-phase exchange chromatography. Purified lipids were then converted into fatty

acid methyl esters by incubation with methanol/2.5% H_2SO_4 before GC/MS analysis. All

data are presented as percentage of the total fraction of fatty acids measured in the

sample.

Mammalian cell culture

MDA-MB-231 cells were obtained from the American Type Culture Collection (ATCC,

Manassas, VA) and cultured in Dulbecco's modified Eagle's medium

(Corning, 10-017-CV). T-47D cells were grown in RPMI-1640 media (Gibco, 11875119).

All cell culture media were supplemented with 10% FBS (Hyclone, SH30910.03, lot

AYG161519), 2 mM glutamine (Corning, 25-005-CI), and penicillin/streptomycin

(Corning, 30-002-CI). Cells were cultured in a humidified incubator at 37°C with 5%

CO_2. In addition, cell lines were maintained at low passage numbers and regularly

tested for mycoplasma.

Mammalian drug sensitivity assays

Measurement of fractional viability (FV) and normalized growth rate inhibition (GR) were

performed using the FLICK assay as previously described [69]. Briefly, cells were seeded

in 96-well optical-bottom black-walled plates (Greiner Bio-One, 655090) in 90 μL of

media and allowed to adhere overnight. Density was optimized for each cell line

depending on the growth rate of the cells and the duration of the assay. Dilutions of

tamoxifen (Selleckchem, S1972) were prepared at 10x final concentration in media

containing 50 µM SYTOX green (Thermo Fisher Scientific, S7020). 10 µL of drug

combinations were supplemented to the cell culture media and fluorescence signals

(excitation = 504 nm, emission = 523 nm) were recorded at the indicated time points on

a Tecan Spark multimode plate reader. Total cell fluorescence was evaluated by lysing

untreated cells at the start and experimental conditions at the end of the assay, using a

supplement of 0.1% Triton X-100 (Fisher Scientific, BP151-500) diluted in PBS.

Supplements: Ferrostatin-1 (Sigma-Aldrich, SML0583), ZVAD[70].

Gene expression profiling by RNA-seq

Total RNA was extracted using TRIzol (ThermoFisher), followed by DNase I (NEB)

treatment and purified using the Direct-zol RNA mini-prep kit (Zymo research). RNA

quality was verified by agarose gel electrophoresis. Briefly, multiplexed libraries were

prepared using Cel-seq2[71]. Two to three biological replicates were sequenced with a

NextSeq 500/550 High Output Kit v2.5 (75 Cycles) on a Nextseq500 sequencer. Paired

end sequencing was performed; 13 cycles for read 1, six cycles for the illumina index

and 73 cycles for read 2. Analysis was performed as previously detailed[72], using a

homemade DolphinNext pipeline[73].

Data availability

RNA-seq data have been deposited at GEO and is publicly available as of the date of

publication under the following accession number: GSE186785. This study does not

report original code.

References

Koppel, N., Maini Rekdal, V. & Balskus, E. P. Chemical transformation of xenobiotics by the human gut microbiota. *Science* **356**, doi:10.1126/science.aag2770 (2017).

Spanogiannopoulos, P., Bess, E. N., Carmody, R. N. & Turnbaugh, P. J. The microbial pharmacists within us: a metagenomic view of xenobiotic metabolism. *Nat Rev Microbiol* **14**, 273-287, doi:10.1038/nrmicro.2016.17 (2016).

Maier, L. *et al.* Extensive impact of non-antibiotic drugs on human gut bacteria. *Nature* **555**, 623-628, doi:10.1038/nature25979 (2018).

Diot, C., Garcia-Gonzalez, A. P. & Walhout, A. J. M. *C. elegans* and its bacterial diet: an interspecies model to explore the effects of microbiota on drug response. *Drug Discovery Today: Disease Models* **28**, 21-26 (2018).

Zhang, X., Han, Y., Huang, W., Jin, M. & Gao, Z. The influence of the gut microbiota on the bioavailability of oral drugs. *Acta Pharm Sin B* **11**, 1789-1812, doi:10.1016/j.apsb.2020.09.013 (2021).

Selwyn, F. P., Cui, J. Y. & Klaassen, C. D. RNA-Seq Quantification of Hepatic Drug Processing Genes in Germ-Free Mice. *Drug Metab Dispos* **43**, 1572-1580, doi:10.1124/dmd.115.063545 (2015).

Garcia-Gonzalez, A. P. *et al.* Bacterial metabolism affects the *C. elegans* response to cancer chemotherapeutics. *Cell* **169**, 431-441 e438, doi:10.1016/j.cell.2017.03.046 (2017).

Scott, T. A. *et al.* Host-microbe co-metabolism cictates cancer drug efficacy in *C. elegans*. *Cell* **169**, 442-456 e418, doi:10.1016/j.cell.2017.03.040 (2017).

Rosener, B. *et al.* Evolved bacterial resistance against fluoropyrimidines can lower chemotherapy impact in the *Caenorhabditis elegans* host. *Elife* **9**, doi:10.7554/eLife.59831 (2020).

Watson, E., MacNeil, L. T., Arda, H. E., Zhu, L. J. & Walhout, A. J. M. Integration of metabolic and gene regulatory networks modulates the *C. elegans* dietary response. *Cell* **153**, 253-266 (2013).

Watson, E. *et al.* Interspecies systems biology uncovers metabolites affecting *C. elegans* gene expression and life history traits. *Cell* **156**, 759-770 (2014).

Garcia-Gonzalez, A. P. & Walhout, A. J. M. Worms, bugs and drugs: *Caenorhabditis elegans* as a model for host-microbe-drug interactions. *Curr Opin Systems Biol* **6**, 46-50 (2017).

Shelly, W., Draper, M. W., Krishnan, V., Wong, M. & Jaffe, R. B. Selective estrogen receptor modulators: an update on recent clinical findings. *Obstet Gynecol Surv* **63**, 163-181, doi:10.1097/OGX.0b013e31816400d7 (2008).

Jordan, V. C. Tamoxifen: a most unlikely pioneering medicine. *Nat Rev Drug Discov* **2**, 205-213, doi:10.1038/nrd1031 (2003).

Shagufta & Ahmad, I. Tamoxifen a pioneering drug: An update on the therapeutic potential of tamoxifen derivatives. *Eur J Med Chem* **143**, 515-531, doi:10.1016/j.ejmech.2017.11.056 (2018).

Manna, S. & Holz, M. K. Tamoxifen Action in ER-Negative Breast Cancer. *Sign Transduct Insights* **5**, 1-7, doi:10.4137/STI.S29901 (2016).

Clifford, R. E., Bowden, D., Blower, E., Kirwan, C. C. & Vimalachandran, D. Does tamoxifen have a therapeutic role outside of breast cancer? A systematic review of the evidence. *Surg Oncol* **33**, 100-107, doi:10.1016/j.suronc.2020.02.006 (2020).

Reece-Hoyes, J. S. *et al.* A compendium of *C. elegans* regulatory transcription factors: a resource for mapping transcription regulatory networks. *Genome Biol* **6**, R110 (2005).

Bekele, R. T. *et al.* Oxidative stress contributes to the tamoxifen-induced killing of breast cancer cells: implications for tamoxifen therapy and resistance. *Sci Rep* **6**, 21164, doi:10.1038/srep21164 (2016).

Lacroix, M. & Leclercq, G. Relevance of breast cancer cell lines as models for breast tumours: an update. *Breast Cancer Res Treat* **83**, 249-289, doi:10.1023/B:BREA.0000014042.54925.cc (2004).

Hafner, M., Niepel, M., Chung, M. & Sorger, P. K. Growth rate inhibition metrics correct for confounders in measuring sensitivity to cancer drugs. *Nat Methods* **13**, 521-527, doi:10.1038/nmeth.3853 (2016).

Schwartz, H. R. *et al.* Drug GRADE: An Integrated Analysis of Population Growth and Cell Death Reveals Drug-Specific and Cancer Subtype-Specific Response Profiles. *Cell Rep* **31**, 107800, doi:10.1016/j.celrep.2020.107800 (2020).

Taubert, S., Ward, J. D. & Yamamoto, K. R. Nuclear hormone receptors in nematodes: evolution and function. *Mol Cell Endocrinol* **334**, 49-55 (2011).

Baba, T. *et al.* Construction of *Escherichia coli* K-12 in-frame, single-gene knockout mutants: the Keio collection. *Molecular Systems Biology* **2**, 2006 0008, doi:10.1038/msb4100050 (2006).

Kachlany, S. C. *et al.* Nonspecific adherence by *Actinobacillus actinomycetemcomitans* requires genes widespread in bacteria and archaea. *J Bacteriol* **182**, 6169-6176, doi:10.1128/jb.182.21.6169-6176.2000 (2000).

Ahmer, B. M., Thomas, M. G., Larsen, R. A. & Postle, K. Characterization of the exbBD operon of *Escherichia coli* and the role of ExbB and ExbD in TonB function and stability. *J Bacteriol* **177**, 4742-4747, doi:10.1128/jb.177.16.4742-4747.1995 (1995).

Ma, D., Alberti, M., Lynch, C., Nikaido, H. & Hearst, J. E. The local repressor AcrR plays a modulating role in the regulation of acrAB genes of *Escherichia coli* by global stress signals. *Mol Microbiol* **19**, 101-112 (1996).

Kobylka, J., Kuth, M. S., Muller, R. T., Geertsma, E. R. & Pos, K. M. AcrB: a mean, keen, drug efflux machine. *Ann N Y Acad Sci* **1459**, 38-68, doi:10.1111/nyas.14239 (2020).

Bhattacharya, S. *et al.* A metabolic regulatory network for the *Caenorhabditis elegans* intestine. *iScience* **25**, 104688, doi:10.1016/j.isci.2022.104688 (2022).

Liu, Y. *et al.* Hydroxysteroid dehydrogenase family proteins on lipid droplets through bacteria, *C. elegans*, and mammals. *Biochim Biophys Acta Mol Cell Biol Lipids* **1863**, 881-894, doi:10.1016/j.bbalip.2018.04.018 (2018).

Yilmaz, L. S. *et al.* Modeling tissue-relevant *Caenorhabditis elegans* metabolism at network, pathway, reaction, and metabolite levels. *Mol Syst Biol* **16**, e9649, doi:10.15252/msb.20209649 (2020).

Watts, J. L. Fat synthesis and adiposity regulation in *Caenorhabditis elegans*. *Trends Endocrinol Metab* **20**, 58-65, doi:10.1016/j.tem.2008.11.002 (2009).

Walker, M. D. *et al.* WormPaths: *Caenorhabditis elegans* metabolic pathway annotation and visualization. *Genetics* **219**, doi:10.1093/genetics/iyab089 (2021).

Brooks, K. K., Liang, B. & Watts, J. L. The influence of bacterial diet on fat storage in *C. elegans. PLoS One* **4**, e7545, doi:10.1371/journal.pone.0007545 (2009).

Kaneda, T. Fatty acids of the genus Bacillus: an example of branched-chain preference. *Bacteriol Rev* **41**, 391-418 (1977).

Sharma, S. *et al.* Free fatty acids induce Lhb mRNA but suppress Fshb mRNA in pituitary LbetaT2 gonadotropes and diet-induced obesity reduces FSH levels in male mice and disrupts the proestrous LH/FSH surge in female mice. *Endocrinology* **154**, 2188-2199, doi:10.1210/en.2012-2218 (2013).

Chen, Y. Y. & Ganzle, M. G. Influence of cyclopropane fatty acids on heat, high pressure, acid and oxidative resistance in Escherichia coli. *Int J Food Microbiol* **222**, 16-22, doi:10.1016/j.ijfoodmicro.2016.01.017 (2016).

Klein, W., Weber, M. H. & Marahiel, M. A. Cold shock response of *Bacillus subtilis:* isoleucine-dependent switch in the fatty acid branching pattern for membrane adaptation to low temperatures. *J Bacteriol* **181**, 5341-5349, doi:10.1128/JB.181.17.5341-5349.1999 (1999).

Cabot, M. C., Giuliano, A. E., Volner, A. & Han, T. Y. Tamoxifen retards glycosphingolipid metabolism in human cancer cells. *FEBS Lett* **394**, 129-131, doi:10.1016/0014-5793(96)00942-8 (1996).

Zhu, H., Shen, H., Sewell, A. K., Kniazeva, M. & Han, M. A novel sphingolipid-TORC1 pathway critically promotes postembryonic development in *Caenorhabditis elegans*. *Elife* **2**, e00429, doi:10.7554/eLife.00429 (2013).

Dancy, B. C., Chen, S. W., Drechsler, R., Gafken, P. R. & Olsen, C. P. 13C- and 15N-Labeling Strategies Combined with Mass Spectrometry Comprehensively Quantify Phospholipid Dynamics in *C. elegans. PLoS One* **10**, e0141850, doi:10.1371/journal.pone.0141850 (2015).

42 Traber, M. G. & Atkinson, J. Vitamin E, antioxidant and nothing more. *Free Radic Biol Med* **43**, 4-
15, doi:10.1016/j.freeradbiomed.2007.03.024 (2007).
43 Zhang, J. *et al.* A delicate balance between bacterial iron and reactive oxygen species supports
optimal *C. elegans* development. *Cell Host Microbe* **26**, 400-411 e403,
doi:10.1016/j.chom.2019.07.010 (2019).
44 Sivik, T. *et al.* 17beta-Hydroxysteroid dehydrogenase type 14 is a predictive marker for tamoxifen
response in oestrogen receptor positive breast cancer. *PLoS One* **7**, e40568,
doi:10.1371/journal.pone.0040568 (2012).
45 Singh, M. S., Francis, P. A. & Michael, M. Tamoxifen, cytochrome P450 genes and breast cancer
clinical outcomes. *Breast* **20**, 111-118, doi:10.1016/j.breast.2010.11.003 (2011).
46 Hrycay, E. G. & Bandiera, S. M. Involvement of Cytochrome P450 in Reactive Oxygen Species
Formation and Cancer. *Adv Pharmacol* **74**, 35-84, doi:10.1016/bs.apha.2015.03.003 (2015).
47 Zangar, R. C., Davydov, D. R. & Verma, S. Mechanisms that regulate production of reactive
oxygen species by cytochrome P450. *Toxicol Appl Pharmacol* **199**, 316-331,
doi:10.1016/j.taap.2004.01.018 (2004).
48 Galluzzi, L. *et al.* Molecular mechanisms of cell death: recommendations of the Nomenclature
Committee on Cell Death 2018. *Cell Death Differ* **25**, 486-541, doi:10.1038/s41418-017-0012-4
(2018).
49 Garbarino, J. & Sturley, S. L. Saturated with fat: new perspectives on lipotoxicity. *Curr Opin Clin*
Nutr Metab Care **12**, 110-116, doi:10.1097/MCO.0b013e32832182ee (2009).
50 Magtanong, L. *et al.* Exogenous Monounsaturated Fatty Acids Promote a Ferroptosis-Resistant
Cell State. *Cell Chem Biol* **26**, 420-432 e429, doi:10.1016/j.chembiol.2018.11.016 (2019).
51 Nakamura, Y. *et al.* Elovl6 regulates mechanical damage-induced keratinocyte death and skin
inflammation. *Cell Death Dis* **9**, 1181, doi:10.1038/s41419-018-1226-1 (2018).
52 Perez, M. A., Magtanong, L., Dixon, S. J. & Watts, J. L. Dietary Lipids Induce Ferroptosis in
Caenorhabditis elegans and Human Cancer Cells. *Dev Cell* **54**, 447-454 e444,
doi:10.1016/j.devcel.2020.06.019 (2020).
53 Yuan, J. Y. & Horvitz, H. R. The *Caenorhabditis elegans* genes *ced-3* and *ced-4* act cell
autonomously to cause programmed cell death. *Dev Biol* **138**, 33-41, doi:10.1016/0012-
1606(90)90174-h (1990).
54 Kisanga, E. R. *et al.* Tamoxifen and metabolite concentrations in serum and breast cancer tissue
during three dose regimens in a randomized preoperative trial. *Clin Cancer Res* **10**, 2336-2343,
doi:10.1158/1078-0432.ccr-03-0538 (2004).
55 Robinson, S. P., Langan-Fahey, S. M., Johnson, D. A. & Jordan, V. C. Metabolites,
pharmacodynamics, and pharmacokinetics of tamoxifen in rats and mice compared to the breast
cancer patient. *Drug Metab Dispos* **19**, 36-43 (1991).
56 Felix, M. A. & Braendle, C. The natural history of *Caenorhabditis elegans*. *Curr Biol* **20**, R965-
969, doi:10.1016/j.cub.2010.09.050 (2010).
57 Yilmaz, L. S. & Walhout, A. J. M. Worms, bacteria and micronutrients: an elegant model of our
diet. *Trends Genet* **30**, 496-503 (2014).
58 Joerger, R. D. & Ganguly, A. Current Status of the Preharvest Application of Pro- and Prebiotics
to Farm Animals to Enhance the Microbial Safety of Animal Products. *Microbiol Spectr* **5**,
doi:10.1128/microbiolspec.PFS-0012-2016 (2017).
59 Gardner, H. W. Oxygen radical chemistry of polyunsaturated fatty acids. *Free Radic Biol Med* **7**,
65-86, doi:10.1016/0891-5849(89)90102-0 (1989).
60 Zhang, P. *et al.* Proteomic study and marker protein identification of *Caenorhabditis elegans* lipid
droplets. *Mol Cell Proteomics* **11**, 317-328, doi:10.1074/mcp.M111.016345 (2012).
61 Olzmann, J. A. & Carvalho, P. Dynamics and functions of lipid droplets. *Nat Rev Mol Cell Biol* **20**,
137-155, doi:10.1038/s41580-018-0085-z (2019).
62 Tesoriere, L. *et al.* Antioxidant activity of all-trans-retinol in homogeneous solution and in
phosphatidylcholine liposomes. *Arch Biochem Biophys* **307**, 217-223,
doi:10.1006/abbi.1993.1581 (1993).
63 Sulston, J. E. & Horvitz, H. R. Post-embryonic cell lineages of the nematode, *Caenorhabditis*
elegans. *Dev Biol* **56**, 110-156, doi:10.1016/0012-1606(77)90158-0 (1977).
64 Sulston, J. E., Schierenberg, E., White, J. G. & Thomson, J. N. The embryonic cell lineage of the
nematode *Caenorhabditis elegans*. *Dev Biol* **100**, 64-119 (1983).

65 Watts, J. L. & Browse, J. Genetic dissection of polyunsaturated fatty acid synthesis in
Caenorhabditis elegans. *Proc Natl Acad Sci U S A* **99**, 5854-5859, doi:10.1073/pnas.092064799
(2002).
66 Connor, A. J. & Watts, J. L. in *Omega fatty acids in brain and neurological health (second edition)*
Ch. 7, 107-116 (Academic Press, 2019).
67 Kim, H. *et al.* A co-CRISPR strategy for efficient genome editing in *Caenorhabditis elegans.*
Genetics **197**, 1069-1080, doi:10.1534/genetics.114.166389 (2014).
68 Perez, C. L. & Van Gilst, M. R. A 13C isotope labeling strategy reveals the influence of insulin
signaling on lipogenesis in *C. elegans. Cell Metab* **8**, 266-274, doi:10.1016/j.cmet.2008.08.007
(2008).
69 Richards, R., Honeywell, M. E. & Lee, M. J. FLICK: an optimized plate reader-based assay to
infer cell death kinetics. *STAR Protoc* **2**, 100327, doi:10.1016/j.xpro.2021.100327 (2021).
70 Slee, E. A. *et al.* Benzyloxycarbonyl-Val-Ala-Asp (OMe) fluoromethylketone (Z-VAD.FMK) inhibits
apoptosis by blocking the processing of CPP32. *The Biochemical journal* **315 (Pt 1)**, 21-24,
doi:10.1042/bj3150021 (1996).
71 Hashimshony, T. *et al.* CEL-Seq2: sensitive highly-multiplexed single-cell RNA-Seq. *Genome Biol*
17, 77, doi:10.1186/s13059-016-0938-8 (2016).
72 Giese, G. E. *et al. Caenorhabditis elegans* methionine/S-adenosylmethionine cycle activity is
sensed and adjusted by a nuclear hormone receptor. *Elife* **9**, doi:10.7554/eLife.60259 (2020).
73 Yukselen, O., Turkyilmaz, O., Ozturk, A. R., Garber, M. & Kucukural, A. DolphinNext: a
distributed data processing platform for high throughput genomics. *BMC Genomics* **21**, 310,
doi:10.1186/s12864-020-6714-x (2020).
836

Acknowledgements

We thank members of the Walhout lab and Job Dekker for discussion and critical

reading of the manuscript. We thank Leslie Shaw for providing the T-47D cell line. This

work was supported by grants from the National Institutes of Health DK068429 and

GM122502 to A.J.M.W., GM127559 to M.J.L., AG058950 to C.P.O. and GM122393 to

A.G.G. Several bacterial and nematode strains used in this work were provided by the

CGC, which is funded by the NIH Office of Research Infrastructure Programs (P40

OD010440).

Author contributions

C.D., A.G.G. and A.J.M.W. conceived the study. C.D. and A.G.G. performed most

experiments with technical help from M.W., H.D., O.P., Y.R. and A.F.V. performed the

fatty acid measurements under supervision of C.P.O. H.N. engineered the Δ*dhs-19*

mutant strain. M.H. and C.D. performed the mammalian cell experiments under the

supervision of M.L. C.D. and H.Z. performed and analyzed the RNA-seq experiments.

The manuscript was written by C.D. and A.J.M.W. with editing by all other authors.

Ethics declarations

The authors have no competing interests.

Supplementary information

Supplementary information includes thirteen Supplementary Figures and six

Supplementary Tables.

Fig. 1: Tamoxifen Kills ER-Negative Breast Cancer Cells and Animals

a, **b** The growth rate index (**a**) and fractional viability (**b**) of ER-positive (T-47D) and ER-negative (MDA-MB-231) breast cancer cells plotted as a function of increasing concentrations of tamoxifen.

c Bright-field images showing *C. elegans* supplemented with increasing doses of tamoxifen, fed *E. coli*, *C. aquatica*, and *B. subtilis*. Images were taken at 2x magnification after 48h exposure to tamoxifen.

d Dose response curves (DRCs) of *C. elegans* supplemented with increasing doses of tamoxifen, fed *E. coli*, *C. aquatica*, and *B. subtilis.* Data are represented as mean ±SEM of three biological replicates.

Fig. 2: *C. aquatica* transport mutants modulate tamoxifen toxicity in *C. elegans*

a Schematic of the bacterial genetics screens performed.

b DRCs of tamoxifen toxicity in animals fed *C. aquatica* mutants identified in the primary screen

c Schematic of the function of the genes mutated in the strains identified in **Fig. 3a-b**. Genes encode for factors involved in the type II secretion system (*tadC*), the TonB-dependent transport (*exbD*), or the *acrA/B* multidrug efflux pump (transcriptional repressor *acrR*).

d Quantification of tamoxifen levels accumulating in *C. aquatica* strains grown on tamoxifen-containing NGM plates.

e Quantification of tamoxifen levels accumulating in wild-type *E. coli*, *C. aquatica*, and *B. subtilis* strains grown on tamoxifen-containing NGM plates.

f Quantification of tamoxifen levels in animals fed wild-type *E. coli*, *C. aquatica*, and *B. subtilis* strains in tamoxifen-containing NGM plates.

Diot *et al.* Fig. 3

Fig. 3: Host long chain fatty acid synthesis modulates tamoxifen toxicity

a Screening a metabolic gene RNAi library identified three genes (*dhs-19*, *elo-3* and *elo-5*) that, when knocked down, increase TAM toxicity. Animals were fed *E. coli* HT115 expressing double stranded RNA as indicated. Control indicates *E. coli* containing vector control plasmid (pL4440). Bright-field images were taken at 2x magnification after a 48h exposure to tamoxifen.

b Tamoxifen dose response curves comparing Δ*dhs-19* mutant animals to wild type animals. Statistical significance was assessed by performing 2way anovas using Graphpad Prism (v9) on original DRCs and is reported in **Supplementary Table 1.** See also **Supplementary Fig. 4**.

c *C. elegans* fatty acid biosynthesis pathway. Gene names are colored by enzymatic function: FA elongation in blue, FA desaturation in pink, and acyl-CoA synthesis in purple.

d Cartoon illustrating calculation of differential area under curve (ΔAUC) used in panels E and F. AUCs were calculated using average DRCs. Statistical significance was assessed by performing 2way anovas using Graphpad Prism (v9) on original DRCs and is reported in **Supplementary Table 2**. See also **Supplementary Fig. 5**.

e, **f** Bar graph showing ΔAUC values obtained from average full DRCs for animals exposed to RNAi of indicated genes.

Diot, *et al.* Figure 4

Fig. 4: Long chain fatty acid profiles of bacteria and *C. elegans* fed different bacteria

a, b Bubble plots showing long chain fatty acid proportions measured by GC-MS in the indicated bacteria (a), and animals fed the indicated bacteria (b). Bubble sizes are indicative of FA ratios within the sample of interest and are not indicative of absolute quantities. Ratios are available in **Supplementary Table 3**, **4**. Data are represented as mean of three to six biological replicates. Statistical pairwise comparisons are available in **Supplementary Table 5**.

Fig. 5: Long chain fatty acid supplementation modulates tamoxifen toxicity

a-m Unless otherwise indicated, fatty acids and vitamin E were supplemented to bacterial culture medium, as described previously (Devkota et al., 2021).

a-c DRCs of tamoxifen toxicity with and without FA cocktail supplementation on animals fed *E. coli* (**a**), *C. aquatica* (**b**) or *B. subtilis* (**c**).

d, **e** DRCs of of tamoxifen toxicity with and without PUFA (ALA) supplementation on animals fed *E. coli* (**d**) or *C. aquatica* (**e**).

f Growth curves of *B. subtilis* supplemented with PUFAs ALA or GLA. For clarity, no error bars are reported in the graph, n=3.

g, **h** DRCs of tamoxifen toxicity with and without MUFA (VA) supplementation on animals fed *E. coli* (**g**) or *C. aquatica* (**h**).

i Growth curves of *B. subtilis* supplemented with MUFAs VA, PA, or OA. No error bars are reported in the graph, n=3.

j-l DRCs of tamoxifen toxicity with and without vitamin E supplementation on animals fed *E. coli* (**j**), *C. aquatica* (**k**) or *B. subtilis* (**l**). **m** DRCs of tamoxifen toxicity with and without the antioxidants vitamin E and N-acetyl-cysteine (NAC). Culture: NAC was supplemented in the bacterial culture medium as for vitamin E. Medium: NAC was supplemented to the NGM.

DRC: dose response curve, FA: fatty acid, MUFA: mono-unsaturated FA, PUFA: poly-unsaturated FA, ALA: alpha-linoleic acid, GLA: gamma-linoleic acid, DGLA: di-homo-gamma-linoleic acid, VA: cis-vaccenic acid, PA: palmitoleic acid, OA: oleic acid. Statistical analysis of DRCs was conducted by performing 2-way Anova using GraphPad Prism (v9) and are reported in **Supplementary Table 6**. Larger panel of fatty acid supplements was tested in **Supplementary Fig. 7**.

Fig. 6: Bacteria affect the modulation of tamoxifen toxicity by apoptosis

a Volcano plot distribution of genes which expression changed significantly in animals fed *E. coli* and exposed to 400 µM tamoxifen (FDR<0.1, >2-fold change). Full list is provided in **Supplementary Table 6**. Pink indicates genes encoding enzymes participating in FA metabolism, and red indicates genes encoding for enzymes that may be involved in the detoxification of tamoxifen. See also **Supplementary Fig. 12**.

b-d DRCs of tamoxifen toxicity in animals fed *E. coli* expressing double stranded RNA as indicated. Control indicates *E. coli* containing vector control plasmid (pL4440). Error bars represent SEM, n=3. Statistical analysis of DRCs was conducted by performing 2-way Anova using GraphPad Prism (v9). Notably, the significance of *CYP35-B1* RNAi (**c**) rescue was not captured on the entire curve where no substantial toxicity can be overserved on the *E. coli* control, but rather on the highest drug-concentration where reliable toxicity can observed.

e-g DRCs of tamoxifen toxicity on apoptosis deficient mutant animals fed *E. coli* (**e**), *C. aquatica* (**f**) or *B. subtilis* (**g**).

h Kinetics of tamoxifen toxicity in ER-negative (MDA-MB-231) breast cancer cells in presence of absence of FA cocktail, Ferrostatin-1 (ferroptosis inhibitor), Vitamin E, or ZVAD (apoptosis inhibitor). See also **Supplementary Fig. 13**.

Fig. 7: Model of Interplay Between Bacteria, Fatty Acid Metabolism, and Death Pathways Affecting Tamoxifen Toxicity