1

4   **Fossils Do Not Substantially Improve, and May Even Harm, Estimates of Diversification**

5   **Rate Heterogeneity**

7   Jeremy M. Beaulieu[1] and Brian C. O'Meara[2]

9   *[1]Department of Biological Sciences, University of Arkansas, Fayetteville, Arkansas, 72701 USA*

10   *[2]Department of Ecology and Evolutionary Biology, University of Tennessee, Knoxville,*

11   *Tennessee, 37996-1610 USA*

*Email for correspondence: jmbeauli@uark.edu

Abstract — The fossilized birth-death (FBD) model is a naturally appealing way of directly

incorporating fossil information when estimating diversification rates. However, an important yet

often overlooked property of the original FBD derivation is that it distinguishes between two

types of sampled lineages. Here we first discuss and demonstrate the impact of severely

undersampling, and even not including fossils that represent samples of lineages that also had

sampled descendants. We then explore the benefits of including fossils, generally, by

implementing and then testing two-types of FBD models, including one that converts a fossil set

into stratigraphic ranges, in more complex likelihood-based models that assume multiple rate

classes across the tree. Under various simulation scenarios, including a scenario that exists far

outside the set of models we evaluated, including fossils rarely outperforms analyses that exclude

them altogether. At best, the inclusion of fossils improves precision but does not influence bias.

Similarly, we found that converting the fossil set to stratigraphic ranges, which is one way to

remedy the effects of undercounting the number of k-type fossils, results in turnover rates and

extinction fraction estimates that are generally underestimated. While fossils remain essential for

understanding diversification through time, in the specific case of understanding diversification

given an existing, largely modern tree, they are not especially beneficial.

Keywords — fossilized birth-death, state speciation extinction, MiSSE, fossils, stratigraphic

ranges, turnover rate

Diversification models, while fascinating to biologists, frequently lurch close to

extinction themselves. For instance, Nee et al. (1994) demonstrated that estimating speciation

and extinction rates from a molecular chronogram is theoretically possible. This was quickly

followed by work from Kubo and Iwasa (1995), which showed that if rates varied through time

there is an infinite number of alternative sets of time-varying speciation and/or extinction rates

that produce the same number of lineages at any given point in time (an observation largely

ignored at the time). State speciation and extinction models (-SSE; Maddison et al. 2007) were

derived as a seemingly robust framework for estimating the direct effects of discrete traits on

diversification rates. However, Rabosky and Goldberg (2015) found that if a tree evolved under a

heterogeneous branching process, completely independent from the evolution of the focal

character, SSE models will almost always return erroneously strong support for a model of state-

dependent diversification. While this issue was partially rescued by the hidden state models of

Beaulieu and O'Meara (2016), there remains some confusion as to whether SSE models remain a

viable means of assessing state-dependent diversification (Rabosky and Goldberg, 2017; but see

Caetano et al. 2018).

With Louca and Pennell (2020) comes a new salvo of criticism regarding diversification

methods, which echo and expand on the earlier points by Kubo and Iwasa (1995), as well as past

concerns about the possibility of estimating extinction rates generally (i.e., Rabosky, 2010). One

postulated source of salvation has been the inclusion of fossils (Mitchell et al. 2019; Louca and

Pennell, 2020; but see Černý et al. 2021). At a minimum, the inclusion of fossils should

drastically reduce the number of congruent models by excluding any congruent model that

assumes no extinction. We also know that fossils, even without formal models, have contributed

greatly to the understanding of diversification processes — take the discovery of mass

extinctions, for example. For these reasons, the fossilized birth-death (FBD) model (Stadler

2010) is seen as an appealing way of directly incorporating fossil information when estimating

diversification rates, because it naturally assumes that the fossil information represents samples

of extinct lineages in the past, in addition to the species sampled at the present. The purpose of

this point of view is to investigate the improvement of estimation of diversification processes at

just the tips of a tree.

Importance of Sampling Ancestors

There are two main ways to incorporate fossils. The first is to use individual fossils

placed as points on the tree (Fig. 1): they can be tips and/or along branches. The second is to

represent stratigraphic ranges, which define the continuous duration of a species using only the

oldest and youngest fossil appearance, ignoring how many fossils are sampled in between

(Stadler et al. 2018). We first focus on the individual fossil process, as this has the most

information, before examining the interval approach. An important, but frequently overlooked,

property of FBD for individual fossils is that it distinguishes between two types of sampled

lineages (Fig. 1). The first are referred to as m fossils, which represent sampled branches that

went completely extinct before the present and did not give rise to any additional sampled

lineages. The second type are referred to as k fossils, which represent samples of lineages that

had sampled descendants (other fossils, or extant tips). In other words, these are fossils that

represent ancestors sampled on internal branches that eventually lead to sampled species. The

model assumes a sampling rate of fossils that does not vary across taxa or time — a questionable

assumption, but probably no worse than similar simplifications that speciation rate or extinction

rate is similarly invariant. However, while the model makes this distinction among these fossil

types, the rate by which each is sampled is governed by the same global sampling rate, ψ, such

that $\psi_m = \psi_k$.

94
Figure 1. a) An example of a fossil set sampled from a complete tree generated by the birth
death process, which includes both surviving and extinct members. b) The sampled phylogeny
pruned down from (a) to only includes sampled tips and fossils. The fossilized birth-death model
distinguishes between two types of sampled lineages. An *m* fossil (denoted by a blue dot)
represents sampled branches that went completely extinct before the present and did not give rise
to any additional sampled lineages (i.e., "childless" fossils). A *k* fossil (denoted by an orange
dot) represents samples of lineages that had sampled descendants (other fossils, or extant tips;
"parent" fossils).
103

The linking of these parameters makes sense. When a dinosaur gets washed into a river,

the probability of it being excavated later by a paleontologist does not depend on whether a

descendant seventy million years later is sampled (making our unlucky dinosaur a *k* fossil) or not

(making this an *m* fossil). However, in the model, an assumption is not only that the probability

of sampling the fossil is the same regardless of whether it has descendants or not, but also that

the probability of including it in the phylogeny is the same whether or not it has descendants. For

example, Paleobiology Database (PBDB; accessed October 2021) has 67 collections for

Tyrannosaurus rex (Osborn 1905) and only eight for *Stegosaurus stenops* (Marsh 1887). For the

FBD models to fully apply, there should be 67/8 times as many *T. rex* fossils as *S. stenops* fossils

included in the tree. The FBD model of Stadler et al (2018), with its coarsening of the required

data to only include stratigraphic ranges, reduces the need for not missing individual fossils but

still makes assumptions of sampling of intervals.

Of course, there are other biases and variation in fossilization rate (the whole field of

taphonomy studies this rich and varied process), but we are concerned with a potential bias of

experimentalists undersampling *k* fossils. Even with approaches that infer whether a fossil is an

m- or *k*-type from the data (Zhang et al. 2016), typically only one fossil per species is included

(i.e., Ford and Benson 2020), and often infer zero *k* fossils on empirical data. This would have

predictable effects on the estimates of ψ as well as the extinction rate itself, even if the methods

place fossils perfectly. To better understand the dynamics of the different fossil types, here is Eq.

(5) from Stadler (2010), which is probability density of a tree, *T*, with *n* extant taxa, $m \geq 0$

fossils samples, and $n \geq 0$ samples conditioned on the time since the most recent common

ancestor of all taxa in the tree, x_1:

$$f[T|t_{mrca} = x_1] = \frac{\lambda^{n+m+2}\psi^{k+m}}{(1-\hat{p}_0(x_1))^2} p_1(x_1) \prod_{i=1}^{n+m+1} p_1(x_i) \prod_{i=1}^{m} \frac{p_0(y_i)}{p_1(y_i)}.$$

For specifics on what the probabilities $p_0(t)$, $p_1(t)$, and $\widehat{p_0}(t)$ represent, we refer readers to the

original derivation in Stadler (2010); for the purpose of this discussion, just note that they do not

include *k*. We highlight two important observations here. The first is that the information

provided by the *m*-type samples are used throughout the equation, including the time at which

the extinct tip was sampled, y_i, as well as the time at which they split from their common

ancestor, x_i. The second observation is that the only information provided by the *k*-type fossils is

simply how numerous they are in the fossil set. In fact, it is irrelevant where exactly they occur

on a branch or which branches have them — that is, the likelihood is the same if they are

dispersed evenly across the tree or sampled in one 10,000-year interval on one edge. This point is

best illustrated by rewriting the numerator in the first term in the equation above to separate out k

and m,

$f[T|t_{mrca} = x_1] = \psi^k \frac{\lambda^{n+m+2} \psi^m}{(1-\widehat{p_0}(x_1))^2} p_1(x_1) \prod_{i=1}^{n+m+1} p_1(x_i) \prod_{i=1}^{m} \frac{p_0(y_i)}{p_1(y_i)}.$

Moving the ψ^k to the front of the equation shows that the effect that k-type samples have on the

probability is simply based on a factor of ψ^k. What this also says is that when both m- and k-type

fossils are included, the overall log-likelihood can be calculated as, $logL = logL_{m\ only} + [k *$

$log(\psi)]$. As an example, take the tree and set of sampled extinct lineages presented in Figure 2d.

The overall log-likelihood for the maximum likelihood estimate (MLE) for this tree is -1224.937.

The log-likelihood assuming only m-type fossils, but using the same parameter estimates, is

-983.6938, and we can add $107*log(\widehat{\psi}=0.1049145)$ to obtain the overall log-likelihood of

-1224.937 (for more details on these calculations see Supplemental Materials).

On the one hand, if a tree was generated by a Yule process (i.e., no extinction), the

inclusion of only k-type samples has no effect on the parameter estimates. Their inclusion will

simply slide the overall log-likelihood down, again, by a factor $k*log(\psi)$. On the other hand, in

the case of non-zero extinction rates, if m-type fossils are included and k-type not added, this will

have a profound impact on the parameter estimates, particularly with regards to ψ and μ. Under

the FBD formulation, the linking of $\psi_m = \psi_k$ forces the model to interpret the lack of k-type

fossils as evidence of a low sampling rate. This will be in tension with the presence of only m-

type samples, such that to explain a low sampling rate, but with many m-type fossils in the set,

the extinction rate must have been substantial. As shown in Figure 2e-f, this is exactly what

happens when the *k*-type fossils are removed from the fossil sample set. The likelihood surface

erroneously shifts away from the generating parameters towards regions of parameter space of

very high extinction rates (based on estimates of turnover rate, $\lambda + \mu$, and extinction fraction,

μ/λ).

Figure 2. Contour plots of the likelihood surface under various scenarios for the same tree
(topology shown in d) simulated under a birth-death process. Each surface is constrained such
that turnover rate ($\tau = \lambda + \mu$) and extinction fraction ($\varepsilon = \mu/\lambda$) are set given a pair of fixed
values, but ψ, the rate of fossil preservation, is free to find their MLE. We sampled 5000 pairs of
points for τ and ε from a latin hypercube sampling design. When provided with a sample of
extinct lineages that is perfectly consistent with the generating model, the FDB performs and
behaves well and generally reduces the variance in turnover rate and extinction fraction (b, c),
relative to ignoring fossils completely (a). However, if *k* fossils are removed completely (e,f), the
likelihood surface begins to erroneously shift away from the generating parameters towards
regions of parameter space of very high extinction rates as values of ψ increase. The dashed
vertical line represents the generating value for turnover rate (τ=0.70) and the dashed horizontal
line represents the generating value for extinction fraction (ε=0.75).

In any case, when provided with a sample of extinct lineages that is perfectly consistent

with the generating model, the FDB performs and behaves exactly as it should (Fig. 2b-c), but so

does a model that ignores fossils completely (Fig. 2a). Our concern, however, is what happens

when theory meets practice. An important element of the FBD model is the possibility of having

sampled ancestors, long a question in paleontology (see Foote 1996). There is a rich literature

investigating the theoretical plausibility (Foote 1996; Gavryushkina et al. 2014) and statistical

identifiability of ancestors in the fossil record (Gingerich 1979; Fisher 2008; Parins-Fukuchi et al

2019; Wagner 2019; Parins-Fukuchi 2021). Many of these predict that sampled ancestors may be

common. For example, Zhang et al (2016) ran simulations that resulted in approximately half to

three-quarters of the fossils being ancestral to descendant samples (other fossils and/or modern

taxa). We simulated over a wide range of conditions (Fig. 3) and found that under most

conditions, sampled ancestors should outnumber other kinds of fossils. To address a reviewer's

question about empirical findings, we examined a few papers that had used the FBD model for

empirical data. Sampled ancestors outnumbering other kinds of fossils is not the pattern typically

found with FBD studies. For example, Gavryushkina et al. (2017) found 18 of 35 fossils were

ancestral, but Slater et al. (2017) and Pyron (2017) appeared to find fewer than 2% of fossils as

being ancestral, based on nonzero terminal branch lengths of their main trees. However, we note

that with FBD, if people are presenting consensus branch lengths as an average, then it is

possible that most of the posterior probability is for zero length branches (thus an ancestral

fossil), but with a handful of samples with the fossil not being ancestral to lead to a nonzero

average branch length.

There are many other potential reasons for mismatch between empirical and simulated

data. We focus here on just one possible explanation, undersampling of k-type fossils. The

likelihood equations of Stadler et al. (2010) all assume that a dataset is not censored to only

include m-type fossils, though some implementations of the FBD allow for using priors

prohibiting sampled ancestors (see comparison conducted by Matzke and Wright, 2016). But, as

far as we can ascertain, there is no conditioning for this censoring in the likelihood equations,

and it may lull empiricists into believing that sampling ancestors is not a necessary requirement.

Of course, as noted above, fossils need not only be treated as individual fossils placed as points

on the tree. A recent extension of the FBD explicitly treats the same fossil sets above as multiple

samples of the same species that specify distinct stratigraphic ranges (Stadler et al. 2018). All

that is required, then, is the oldest and youngest fossil within a range, as opposed to the total

number of fossils, which is then marginalized out, potentially rendering issues related to

counting k fossils somewhat moot in most cases.

These are the topics we seek to investigate here. Specifically, we explore the benefits of

including fossils in more complex models that assume multiple rate classes across the tree in

comparison to excluding them altogether. In short, will fossils help diversification analyses

largely based on modern phylogenies? Our hypotheses at the outset were that 1) well-sampled

fossils would have a very beneficial effect on estimation of diversification parameters versus

having no fossils and that models that exclude k-type fossils would perform poorly, likely being

outcompeted by models with no fossils; 2) we expected approaches with stratigraphic samples

would have less benefit (reduced data, especially about fossilization rate) but also would be

relatively insensitive to sampling issues; and 3) we expected that fossils would be especially

valuable in a situation where the true model was more complex than any model used in analysis,

which approximates the situation in reality. To keep analyses feasible, we assumed that we knew

the topology and branch lengths of the tree, and timing and placement of any fossils used,

perfectly. Improvement of empirical trees from inclusion of fossil taxa, or harms coming from

incorrect fossil placement or uncertain timings, are beyond the scope of this study.

Figure 3. Histogram of the fraction of *k*-type fossils sampled across a large set of simulations
over a wide range of conditions in relation to *k*-type fossils used in empirical FBD studies.
Specifically, we generated a grid that make up all possible combinations of elements from
vectors of turnover rate (incrementing $\tau = 0.1$ to 1.0, sampling every 0.1), extinction fraction
(incrementing $\varepsilon = 0.05$ to 0.95, sampling every 0.1), and sampling rate ($\psi = 0.1$, 0.05, 0.10,
0.20). We repeated each generating parameter 20 times, resulting in 8000 simulation replicates in
total, with the histogram representing the distribution of the fraction of *k*-type fossils sampled
across all replicates. Despite the wide range of simulation conditions, empirical studies (blue
dots) appear to have fewer sampled ancestors. The red line along the x-axis represents the
approximate range of simulated values from Zhang et al. (2016), and the blue dot corresponding
to the study of Zhang et al. (2016) represents the median fraction across the range used in their
analysis of Hymenoptera.

Fossilized *SSE

Incorporating the sampling rate, ψ, into a birth-death model that allows multiple discrete

shifts has been implemented before (e.g., fossil BAMM, Mitchell et al. 2019). However, here we

incorporate fossils within our hidden state speciation and extinction framework (HiSSE;

Beaulieu and O'Meara, 2016), that includes any number of observed and/or hidden states. We

focus our tests here exclusively using MiSSE (see Vasconcelos et al. 2021), a likelihood-based,

hidden state only model, like BAMM, but without priors (which could help, hurt, or have little

effect on inference), but also, importantly, without the discontinuous inheritance of extinction

probability that made the likelihoods used in BAMM mathematically incorrect (see Moore et al.

2016). It is relatively straightforward to extend the equations of the canonical FBD model of

Stadler (2010) to the state speciation and extinction models of Maddison et al. (2007) so that

they also include both state transitions and the tree-wide rate of fossil sample rate parameter, ψ:

$\frac{dE_i}{dt} = \mu_i - \left(\lambda_i - \mu_i - \psi + \sum_{i \neq j} q_{ij}\right)E_i(t) + \mu_i E_i(t)^2 + \sum_{i \neq j} q_{ij}E_j(t),$

$\frac{dD_i}{dt} = -\left(\lambda_i - \mu_i - \psi + \sum_{i \neq j} q_{ij}\right)D_i(t) + 2\lambda_i D_i(t)E_i(t) + \sum_{i \neq j} q_{ij}D_j(t).$ (1)

The probability $E_i(t)$ is the probability that a lineage starting at time t in state i leaves no

descendants at the present day (t=0), and $D_i(t)$ is the probability of a lineage in state i at time t

before the present (t>0) evolved the exact branching structure as observed. These ordinary

differential equations are generalized so that any number of observed or hidden states can be

included in the model. For character-based models, like HiSSE and MuHiSSE, i and j represent

the different observed and hidden states combinations specified in the model, whereas with

MiSSE i and j represent hidden states only. For an extant tip, the initial condition for $D_i(0)$ is ρ_i,

which defines the probability that an extant individual observed in state i is sampled in the tree

and $1 - \rho_i$ for $E_i(0)$. For an m-type fossil, $D_i(t)=\psi E_i(t_e)$, and $E_i(t)=E_i(t_e)$, where t_e represents

the time at which the sampled extinct lineage was sampled; for k-type fossil, $D_i(t)=\psi D(t)$, and

$E_i(t)=E_i(t)$. At nodes, N, the initial condition is the combined probability of its two descendant

branches, L and R, such that, $D_{N,i} = D_{L,i}D_{R,i}\lambda_i$. The overall likelihood is the sum of $D_{N,i}$

calculated at the root.

When the fossil set represents stratigraphic ranges, the likelihood calculation is much

more involved and requires designating three specific types of edge segments. First, we note that

our implementation reflects the "symmetric speciation only" portion of Stadler et al. (2018; see

Corollary 13), where only bifurcating speciation events are allowed — that is, a speciation event

reflects the extinction of an ancestral species, and two new descendant species arise. This greatly

reduces the complexity of the original model formulation, which at its most complex, also

includes both anagenetic (ancestral species is replaced by a new descendant) and asymmetric (a

single descendant "buds" off a persistent ancestral lineage) speciation processes with their own

rates. Second, we note that a stratigraphic range is defined only by the oldest (o) and youngest

(y) fossils that are unequivocally assigned to a species across a distinct time interval. Thus, a

stratigraphic interval records the known duration of species on branches. These ranges are

represented by a single fossil and a tip (i.e., fossil is $y=0$), by a branch fossil and an extinct tip

($y=t_e$), by two fossils representing a sampled ancestral stratigraphic range along an edge ($o>y$), or

even by a single fossil (i.e., $o=y$). Fossils that occur within a given range are ignored.

When an edge segment is not associated with a stratigraphic range, the branch calculation

and starting conditions proceed exactly as described in Eq. 1 above. However, when an edge

segment represents a stratigraphic range, we must modify $D_i(t)$ to remove the possibility of an

unobserved speciation and subsequent extinction event within the interval given that we know

that the entire segment [o,y] belongs to the same species:

$\frac{d\widehat{D}_i}{dt} = -\left(\lambda_i - \mu_i - \psi + \sum_{i \neq j} q_{ij}\right)\widehat{D}_i(t) + \sum_{i \neq j} q_{ij}\widehat{D}_i(t).$ (2)

When y=0, such that it is an extant species, the initial condition for $\widehat{D}_i(0)$ is ρ_i; if y is an m-type

fossil, $\widehat{D}_i(t)$=$\psi E_i(t_e)$; if y denotes the beginning of a sampled ancestor stratigraphic range then

$\widehat{D}_i(t)$=$D(t)$. Finally, for edge segments that represent intervening time intervals between two

stratigraphic ranges without an observed speciation event, we modify the branch calculation to

account for zero or more unobserved speciation events. Again, given the assumption that a

stratigraphic range represents the sampling of a single species across a distinct interval of time,

the presence of two stratigraphic ranges along the same edge would imply that at least one

unobserved speciation event had occurred somewhere within this interval. To account for this, at

the rootward end, y_a, which represents the youngest fossil of the older of the older of the two

stratigraphic ranges, we correct $D_i(t)$ following the Stadler et al. (2018):

$D_i(t) = D_i(y_a) * \left(1 - \frac{\widehat{D}_i(y_a)}{D_i(y_a)}\right).$ (3)

This probability then becomes the starting condition for the next stratigraphic range. Note that

none of three specific branch types requires altering $E_i(t)$ from what is presented in Eq. 1,

because this probability is based solely on time and is therefore unaffected by the tree topology.

As before, at nodes, N, the initial condition is the combined probability of its two descendant

branches, L and R, such that, $D_{N,i} = D_{L,i}D_{R,i}\lambda_i$. The overall likelihood is, again, the sum of $D_{N,i}$

obtained at the root, but we must also marginalize over the number of fossils within a

stratigraphic range. This is done by multiplying the sum of $D_{N,i}$ at the root by $\psi^{K'}$ and $e^{\psi L_s}$,

where K' represents the total number of sampled fossils that represent the start and end times of a

stratigraphic range (if $o=y$ for a given range then this counts as only one fossil), and where L_s

represents the sum of all stratigraphic range lengths ($\sum_{i=1}^{n} o_i - y_i$).

For both the original FBD, and FBD with stratigraphic ranges, we condition the overall

likelihood by $\lambda_i [1 - E_i(t \mid \psi = 0)]^2$, where and $E_i(t \mid \psi = 0)$ is the probability that a

descendant lineage of the root survived to the present and was sampled assuming no sampling in

the past. We note that fossil BAMM does not condition on survival in this way, or even at all,

when the tree includes extant taxa. For character-based models, such as HiSSE and MuHiSSE,

we weight the overall likelihood by the probability that each possible state gave rise to the

observed data (see FitzJohn et al. 2009). However, it was pointed out by Herrera-Alsina et al.

(2018) that at the root, the individual likelihoods for each possible state should be conditioned

prior to averaging the individual likelihoods across states. It is unclear to us which procedure is

correct, but it does seem that both weighting schemes behave exactly as they should in the case

of character-independent diversification — that is, the overall likelihood reduces to the

likelihood of the tree multiplied by the likelihood of a trait model. We have also tested the

behavior of both (not shown), and the likelihood differences are very small, and the parameter

estimates in simulation are nearly indistinguishable from one another.

In the absence of character information, assuming a model with a single birth rate and

death rate, and a $\psi>0$ the likelihoods are identical to that of Stadler (2010) and Stadler et al.

(2018). To ensure that our math and the implementations are correct with two or more birth and

death rates, we relied on the property of SSE models described by Beaulieu and O'Meara (2016)

and Caetano et al. (2018): when a trait has no differential effect on the diversification process

(i.e., the character-independent model), the overall likelihood is the product of the tree likelihood

and likelihood of a Markov model applied to the character data (it is the sum of the two

likelihoods in log space). Thus, for a binary character that is independent of the diversification

process on a tree with two rate classes and a non-zero fossil preservation rate, the BiSSE

likelihood should reduce to the product of the likelihood of a MiSSE model with two rate classes

and the likelihood of the transition rate model applied in the character data. As we show in the

Supplemental Materials, we can confirm that our implementation of the FBD for MiSSE, HiSSE,

and MuHiSSE (two binary characters; Nakov et al. 2019) are indeed correct. We also show that

the different branch calculations for FBD with stratigraphic ranges conforms to the analytical

calculations provided by Stadler et al. (2018). All models are available in the R package *hisse*

(Beaulieu and O'Meara 2016).

How Do Fossils Help in a World Where All Models are Simply Wrong?

Mitchell et al (2019) showed that with their fossil implementation within BAMM, which

allows for shifts in discrete regimes of speciation and extinction, there were substantial

improvements to the rate estimates, most notably with the extinction fraction. These results seem

at odds with our results presented thus far, as well as results from a set of additional single rate

regime simulations where trees were generated using a range of speciation and extinction rates

(see Table S1; Supplemental Materials). While their simulations were focused on understanding

the behavior of a much more complex model, it should be noted that the most striking

improvement in extinction fraction occurred when $\psi=1$ fossils/Myr, which is an extremely high

rate of preservation. In our case, the maximum value $\psi=0.1$ fossils/Myr resulted in a fossil set

that was at least as numerous as the number of extant taxa ($N>200$), which seemed reasonable to

us. Still, even when the addition of fossils resulted in doubling the amount of data available to

the birth-death model, there was very little impact on the rate estimates, especially when the

generating extinction fraction was high ($\varepsilon = 0.75$; Figure 2). We also wondered how other

functions of speciation and extinction, such as turnover rates and net diversification rates, are

impacted not only by the inclusion of fossils, but also by the inclusion of biased samples. The

simulations of Mitchell et al. (2019) demonstrated that estimates of speciation are insensitive to

the inclusion of fossils, with extinction being strongly impacted, which suggests that this would

also impact additional functions of speciation and extinction in different ways.

Similar to the single regime simulations above, when we first simulated under our MiSSE

model under scenarios of moderate difficulty, the main effect of including fossils when

estimating of functions of speciation and extinction (namely, the turnover rate, $\tau_i = \lambda_i + \mu_i$ and

the extinction fraction, $\varepsilon_i = \mu_i/\lambda_i$) was, again, to simply reduce the variance of the estimates

(i.e., calculated as the mean of the squared errors from the $x_{\text{model-averaged}}$) across the different rate

regimes, not the bias (i.e., calculated as the mean of $x_{\text{model-averaged}} - x_{\text{true}}$; see Supplemental

Materials for more details). As expected, removing k fossils completely from the set results in

very high biases related to turnover rates specifically, as ψ increases, and rather severe

downward biases generally in estimates of net diversification rates ($r_i = \lambda_i - \mu_i$). With regards

to converting the fossil set to stratigraphic ranges, there was a curious and general tendency for a

downward bias in both turnover rate and extinction fraction.

An easy and valid criticism of these types of simulations is that they are too simplistic.

Trees were generated under an SSE model that shifted between, at most, two different rate

classes, with the rate of these shifts set by a single transition rate, q. While we were genuinely

surprised that extant-only trees performed as well as trees that also included fossil sample data

under the same conditions, the simulation scenarios are hardly realistic. The processes that

generate most empirical trees are likely very complex, likely carrying signatures of non-random

extinction, even mass extinctions, with diversification rates varying substantially among lineages

and across time. The simplicity of our MiSSE models is out of mathematical convenience and

tractability. Even still, we might expect that even when the true model is not included in the set

of models evaluated, the inclusion of fossil information attached to branches distributed across

the tree would provide more weight and better parameter estimation for the more complex

models that are included in the set. Most models will work fine with data that meet their

assumptions, but with complex and messy data, maybe fossils are needed to rescue their

performance.

To explore this further we devised a simulation scenario that was meant to closely imitate

processes and conditions that are more realistic in empirical settings, which would also prove

challenging to our MiSSE model, but also be biased in a way that would favor including fossils.

Specifically, we simulated 100 trees where we assumed that there were four discrete "regimes"

of turnover rates ($\tau_A= 0.3$, $\tau_B= 0.6$, $\tau_C= 1.5$, $\tau_D= 1.0$ events/Myr) and extinction fractions ($\varepsilon_A=$

0.7, $\varepsilon_B= 0.9$, $\varepsilon_C= 0.95$, $\varepsilon_D= 0.8$) that controls the diversification dynamics. We also assumed a

rather extreme heterogeneous transition matrix, $\mathbf{Q}$, that governed the dynamics of transitions, q_{ij},

among these four regimes,

$$\mathbf{Q} = \begin{array}{c} A \\ B \\ C \\ D \end{array} \begin{bmatrix} - & 0.01 & 0.01 & 0.01 \\ 0.05 & - & 0.04 & 0.02 \\ 0.02 & 0.03 & - & 0.03 \\ 0.06 & 0.02 & 0.08 & - \end{bmatrix}$$

The magnitude of these rates was chosen somewhat at random, but cumulatively ensured that

approximately 10% of the branches across the entire history of the tree (including extinct

lineages) contained at least one change in rate regime. We also encoded two mass extinction

events. The first occurred after the first 40-time units and removed, at random, 70% of lineages

alive at that time point. The second occurred 30-time units later, 70-time units in total from the

start of the simulation and removed 90% of the lineages alive at the time point (see Fig. 4a).

Finally, simulations were terminated once the tree reached 267 taxa. However, we non-randomly

chose 67 taxa and removed them from the final tree. The purpose of this procedure was to

assume that our final tree had a biased sampling fraction of 75%. The bias was generated by

simulating a trait under Brownian motion, then normalizing the values so that they were between

0 and 1 and had a phylogenetic signal. These values were then used as probabilities for removing

taxa.

We know fossilization rates vary, sometimes dramatically, by time and taxa (Wagner and

Marcott 2013). Incorporating this would make the simulation far more realistic as well, but it

would provide additional complications given the difficulties in separating variable extinction

rates from variable preservation rates (e.g., Foote et al. 2019; Louca et al. 2021). The fact that the

fossilization process matches that of the model gives the fossils a chance to perform better than

not including them. That is, can well-modeled fossils help an analysis where the true

diversification model is more complex and different than any analyzed? If fossils cannot help in

this case, they probably will not help. Had we used a more complex fossilization process, and

fossils failed to help in that case, it could just be that we tried too extreme a difference and set up

a bias against the utility of fossils. In all cases, we assumed that all the models started with the

true trees and that any fossils that were sampled were placed perfectly, both in time and on the

tree. These assumptions are quite optimistic; our goal was to find the impact of fossils on

diversification rate estimates, not assess the impact of fossils on all aspects of tree inference

(which would require simulation of morphology and other traits, state-based fossilization rates,

tree inference, homology assessment, and more), nor make it difficult for fossils by adding

realistic issues of difficulty with placement, taphonomic biases, and more.

For each simulation replicate, we fit four MiSSE models where we set $f_i = 0.75$ to

account for incomplete sampling (though assuming, incorrectly, that all taxa have the same

sampling rate; see Beaulieu 2020). Specifically, we fit a single regime model, a model that only

allows turnover rate to vary ($\tau_A \neq \tau_B$), a model that only allowed extinction fraction to vary, and

a model that allowed both to vary ($\tau_A \neq \tau_B$ and $\varepsilon_A \neq \varepsilon_B$). Each model was evaluated using a two-

step optimization routine. The first step consists of a bounded stochastic simulated annealing run

for 5000 iterations, followed by a bounded subplex routine that searches parameter space until

the maximum likelihood is found. This model set ensured that, as with real data, the set of

models were far simpler than the true generating model. We also fit the same set of models for

increasing values of ψ (i.e., 0.01, 0.05, 0.10 events/Myr), and for different sampling of k-type

fossils (i.e., full, half k, only m-type fossils used), including converting the set to stratigraphic

intervals. When converting a fossil set into a set of stratigraphic ranges, we simply take the

oldest and youngest fossil on each edge, removing all others, prior to pruning unsampled extinct

lineages from the full tree. We chose to summarize results based on the diversification rates

model-averaged across only the tips that survived to the present, because the rate regimes in the

generating model do not map to the rate regimes in the model set, namely, regime D in a four-

rate model does not easily map to regime A or B in a two-rate model. For a given model, the

marginal probability of each rate regime is obtained for every tip, and the rates for each regime

are averaged together using the marginal probability as a weight: a weighted average of these

rates is then obtained across all models using Akaike weights. The use of model-averaging is

particularly advantageous here because the tip rates are not conditional on a single best model

that would be far less complex than the model that generated the data. Different models reflect

different aspects of the data. By fitting multiple models, each good model might tell us

something about how the data evolved. Model-averaging is one way to summarize these different

elements in a comprehensible way.

449
Figure 4. The uncertainty surrounding estimates of turnover rate when the models being
evaluated does not include the true model. Specifically, we assumed that there were four discrete
"regimes" of turnover rates (τ_A= 0.3, τ_B= 0.6, τ_C= 1.5, τ_D= 1.0) and extinction fractions (ε_A= 0.7,
ε_B= 0.9, ε_C= 0.95, ε_D= 0.8) that controls the diversification dynamics; we also assumed a rather
extreme transition dynamics among these four regimes (see text). We also encoded two mass
extinction events as shown by the lineage through time plot (a), and we also assumed biased
sampling among the tips that survived to the present. The boxplots represent the distribution of
model-averaged rates based on the true rate classes at the tips. (b-c) Despite the true model being
far outside the models being evaluated, the turnover rates at the tips hints at there being more
complexity within the data than even the most complex two-rate models in the set might suggest.
However, the inclusion of fossils does not seem to perform remarkably better than not including
them.
462
Despite the true generating model being far outside the set of models evaluated, the use

of model-averaging at least hints at there being more complexity within the data than even the

most complex two-rate models allow. When the model-averaged tip rates are aligned with the

four true tip rate regimes for each simulation iteration there are consistent significant positive

slopes across all treatments (Fig. 4). For fossil-based analyses, particularly for those where the

number of fossils was greater than the number of extant tips (i.e., $\psi = 0.10$ fossils/Myr), there are

clearer differences between the four rate regimes compared to the extant-only estimates.

However, as with all other analyses presented here, when the fossil set is perfectly consistent

with the FBD process by which the samples were taken, the main effect of the fossils is to reduce

the overall uncertainty in the tip rate estimates compared to extant-only inferences.

There were some improvements with respect to the parameter estimates with the

inclusion of the full fossil sets overall, most notably at two ends of the rate distribution under the

generating model (i.e., $\tau_A = 0.3$ and $\tau_C = 1.5$ events/Myr). Incidentally, both rate classes were by

far the most frequent states across the entire history of almost every simulated tree. The highest

turnover rate regime also had the highest extinction fraction, and this coupled with several mass

extinctions, resulted in very few survivors at the tips. Even with the addition of a substantial

number of fossils placed throughout the tree, the improvement in inferring this particular rate

category from model-average rates was minimal. From a qualitative standpoint, by ignoring

biases in rate and focusing on comparisons among tips, it seems that all treatments, regardless of

the fossil makeup, can consistently assign the correct sign differences among pairs of tip rates.

That is, even when the trees were analyzed with only half of the k-type fossils used, if they were

removed entirely, or converted to stratigraphic ranges, the expected proportion of sign

differences are consistent with the extant-only and full fossil set estimates (Table S2). Not

surprisingly, as the number of k-type fossils are removed, the model-averaged estimates of

turnover rate become greatly inflated, from erroneously high estimates of the extinction, and are

associated with much greater uncertainty. When the fossil sets are converted to stratigraphic

ranges, the turnover rates are, again, generally underestimated, with the improvement of the

estimates with the lower turnover rate regimes being simply incidental. Taken together, this is

consistent with the MiSSE scenarios above (Fig. S1), as well as the constant birth-death models

(Fig. 2e-f), which suggests that: 1) there is only a modest improvement in the rate estimates

when perfectly placed fossils are included; 2) gross omissions of k-type samples in a FBD

erroneously shifts estimates towards areas of very high turnover rates and extinction fraction;

and 3) converting to stratigraphic intervals results in erroneous shifts towards lower turnover

rates and extinction fraction.

A Caution About Stratigraphic Ranges

Our simulation results highlight a curious pattern in that when a fossil set is converted to

stratigraphic ranges, estimates of turnover rate and extinction fraction are increasingly

underestimated as the generating ψ increases. A clear illustration of the pattern is shown in the

likelihood surfaces depicted in Figure 5a-b. When the generating ψ is low (ψ=0.01 fossils/Myr;

Fig. 5a), the likelihood surface includes the generating parameters. However, when ψ is

increased by an order of magnitude (ψ=0.1 fossils/Myr; Fig. 5a-b), the likelihood surface begins

to shift away from the generating parameters towards regions of parameter space represented by

lower turnover rates and extinction fraction. While this downward bias in this particular example

is somewhat subtle, this behavior is consistent across all models and all simulation scenarios

examined here. It should be noted that it is possible that we have incorrectly interpreted the math

described in Stadler et al. (2018). Admittedly, the implementation of this model was quite

challenging, and is further complicated by the fact that FBD with stratigraphic ranges is not

naturally nested within the canonical FBD model. That is, if all fossils are treated as single

representatives of a stratigraphic range, the likelihood does not simply revert to the likelihood of

the canonical FBD model. This is because of the assumption in the FBD with stratigraphic

ranges that corrects for unobserved speciation occurring between two distinct stratigraphic

ranges on the same edge (see Eq. 3). That being said, in the constant birth-death case, we

implemented the model using the analytic equations presented in Stadler et al. (2018) as well as

using the ordinary differential equations above and the likelihoods were identical across a range

of parameter values.

519
Figure 5. (a-b) Contour plots of the likelihood surface for the fossilized birth-death with
stratigraphic ranges using the same tree and data set as shown in Figure 2. When the generating
ψ is low, the likelihood surface includes the generating parameters, but when ψ is increased by
an order of magnitude (ψ=0.1), the likelihood surface also begins to drift away from the
generating parameters towards regions of parameter space represented by lower turnover rates
and extinction fraction. (c) A stratigraphic range records the known duration of species on
branches, and so we assume that no unobserved speciation or extinction takes place along these
intervals. Since the same rates apply to all branches, and since turnover rate is a measure of the
total number of speciation and extinction events, as the number of stratigraphic ranges and
duration of each increase, turnover rates will increasingly be underestimated as the proportion of
the stratigraphic ranges relative to the total tree length increases. Interestingly, this does not seem
to impact estimates of extinction fraction (d).

In the absence of an implementation error, we suspect that the underestimation of

turnover rate is likely due to the assumptions made when computing the probability of the

portion of an edge representing a stratigraphic range. Under this model a stratigraphic range

records the known duration of species on branches, and so we assume that no unobserved

speciation or extinction takes place along these intervals. Since the same rates apply to all

branches, and since turnover rate is a measure of the total number of speciation and extinction

events, we might expect that as the number of stratigraphic ranges and duration of each increase,

the rate estimates would likely reflect a balance between a large proportion of time in a tree

where no unobserved speciation and extinction events have occurred. Thus, we would predict

that the downward bias in turnover rates would be a function of the proportion of the tree

consisting of stratigraphic ranges. We conducted a simulation where we evolved a set of 100

trees for 50 Myr, assuming the same generating rates as the ones that produced the trees in

Figure 2. We also used a range of fossil sampling rates starting from $\psi=0.05$ fossils/Myr and

increasing at 0.01 increments until reaching $\psi=0.15$ fossils/Myr. Consistent with our hypothesis,

we found that as the proportion of the stratigraphic ranges relative to the total tree length

increased (i.e., stratigraphic ranges accounted for increasingly more of the time represented in

the tree), we found that this was significantly and negatively associated with an increased

downward bias in turnover rates ($R^2=0.15$, slope=-0.272, $P<0.001$). Interestingly, we did not find

a similar trend with extinction fraction ($R^2<0.001$, slope=0.09, $P=0.246$). While these results

only speak to our implementation of the "symmetric speciation only" portion of Stadler et al.

(2018; see Corollary 13), where only bifurcating speciation events are allowed, we suspect this

behavior is general for all stratigraphic range models. Even if allowing for anagenetic and/or

asymmetric speciation modes, we suspect turnover rate, which is a measure of the frequency of

both speciation and extinction events, will be generally underestimated given all three speciation

modes remove the possibility of an unobserved speciation and subsequent extinction event along

a stratigraphic range edge.

Concluding Remarks

Our results show that, if the trees fit the generating model well, then adding fossil taxa

correctly might help with precision in rate estimations, but not with bias. However, for most

cases if there is undersampling of fossils along branches (k-type fossils) the results are worse

than ignoring fossils altogether, even when the placement of the sampled fossils was correct

temporally and topologically. When placing the fossil set not as individual points on the tree, but

rather as stratigraphic ranges, there is tendency for some measures of diversification to be

systematically underestimated (i.e., turnover rate). All these results were obtained from a

simulation framework where the diversification models used were far simpler than the simulated

data. We also expected that at least the full fossil set would be able to somewhat rescue the

results given that they were nearly three-times more numerous than extant species in certain

scenarios (i.e., scenario 4). As we have emphasized elsewhere (see O'Meara and Beaulieu 2021),

moving forward, focusing our comparisons of diversification rates at the tips of the tree remains

a fruitful avenue of inquiry, even without fossils, given that most of the data from a phylogenetic

tree of modern taxa exists nearer the present.

We hasten to emphasize that fossils remain key for understanding macroevolution — that

is, important extinct groups like trilobites, sauropods, and extinct lycophytes are essential to our

understanding of evolution in deep time. However, the idea that neontological studies of

diversification are dramatically enhanced by sprinkling carefully chosen fossils on a tree and

applying sophisticated fossilized birth-death models to understand rate variation does not seem

supported by our simulations. At best, fossils have minor effects. At worst, they lead to less

accurate inferences than removing them altogether. We even carefully biased our study to

purposely give fossils the best possible chance, namely, by perfectly identifying each fossil,

placing them with full certainty in a phylogeny, and assuming that their true sampling rate was

constant through time. All this is impossible with empirical data, which is likely the product of

variable fossilization rates (Wagner and Marcott 2013) that further complicates teasing apart

variable extinction from variable preservation rates (see Foote et al. 2019; Louca et al. 2021).

Taken together, our simulations provide us with little hope for the utility of fossils in real-world

applications, at least with regards to estimating diversification rate heterogeneity across a tree.

One area where we suspect fossils will continue to be of great importance in

neontological applications is in molecular divergence time analyses (e.g., Heath et al. 2014).

Much of the information used in fitting most diversification models is the timing between events.

So, if the relaxed clock does not properly account for clades with a faster rate of molecular

substitution, most methods to smooth branches will yield ages that are much older than they

should be (since the branches are still too long relative to the variation implicit in the clock; see

Beaulieu et al. 2015), which will tend to also decrease the net diversification rate across the tree.

However, having more fossils throughout the tree, we suspect, might alleviate this issue. It thus

remains an open question (though with reasonable advocacy for various sides) whether methods

that jointly estimate the topology, branching times, and diversification parameters using an FBD-

type model give a better estimate of diversification rate parameters than alternate approaches

(e.g., *r8s*; Sanderson 2002), which infer a chronogram agnostic to the diversification rate process

and is subsequently used as input in diversification analysis programs. Our study simply

demonstrates that in the case of a perfectly accurate (albeit perhaps undersampled) tree, fossils

do not help substantially at estimating diversification process rates at the tips.

One motivation for this research, in the light of recent and well-earned concern about

diversification methods using phylogenies of extant taxa (Kubo and Iwasa, 1995; Maddison and

FitzJohn, 2015; Rabosky and Goldberg, 2015; Louca and Pennell, 2020; Louca et al. 2021) is to

help guide empiricists seeking to create improvements in diversification rate estimation. There is

understandable interest in fossils to improve diversification models. After all, we owe the

discovery of extinction in the nineteenth century to fossils, and major events like mass

extinctions or the Cambrian explosion were first and by far best known from purely the fossil

record. While we did not investigate, and remain pessimistic, about the feasibility of estimating

diversification processes through time, we were optimistic about the potential impact of fossils

on rates at the tips as a first step. In our own reading of the foundational work of Stadler (2010),

which was motivated by our own curiosity about a path for improvement, we were struck by the

m versus k fossil distinction and what this might mean for its usability in practice, as well as how

stratigraphic intervals might mitigate this issue. We were surprised that, at least for our particular

use case, fossils provided at best little benefit to, and at worst harmed, inferences of

diversification at the tips from trees of mostly modern taxa. This does not imply they may not

help substantially in other areas of phylogenetics of extant taxa, but as a solution to this

particular issue, they are less helpful than we hoped, and researchers might be more fruitful

attempting other solutions first.

Funding

This work was funded by the National Science Foundation grants DEB–1916558 and DEB-1916539.

Supplementary Materials

Data available from the following github repository:

https://github.com/thej022214/Fossils_impact_BO

Acknowledgements

We thank members of the Beaulieu and O'Meara labs for their comments and discussions of the ideas presented here.

References

Beaulieu J.M. 2020. The problem with clade-specific sampling fractions. Available at:https://rdrr.io/cran/hisse/f/inst/doc/Clade-specific-sampling.pdf.

Beaulieu J.M., O'Meara B.C. 2016. Detecting hidden diversification shifts in models of trait-dependent speciation and extinction. Systematic Biology 65:583-601.

Beaulieu J.M., O'Meara B.C., Crane P., Donoghue M.J. 2015. Heterogeneous rates of molecular evolution and diversification could explain the Triassic age estimate for angiosperms. Systematic Biology 64:869-878.

Burnham K.P., and Anderson D.R. 2003. Model selection and multimodel inference: a practical information-theoretic approach. New York:Springer.

Caetano D.S., O'Meara B.C., Beaulieu J.M. 2018. Hidden state models improve state-dependent

diversification approaches, including biogeographic models. Evolution 72:2308-2324.

Černý D., Madzia D., Slater G.J. 2021. Empirical and methodological challenges to the model-

based inference of diversification rates in extinct clades. Systematic Biology

https://doi.org/10.1093/sysbio/syab045.

Fisher D.C. 2008. Stratocladistics: Integrating temporal data and character data in phylogenetic

inference. Annual Reviews of Ecology, Evolution, and Systematics 39:365-385.

FitzJohn R.G., Maddison W.P., Otto S.P. 2009. Estimating trait-dependent speciation and

extinction rates from incompletely resolved phylogenies. Systematic Biology 58:595–

611.

Foote M. 1996. On the probability of ancestors in the fossil record. Paleobiology 2:141-151.

Foote M., Sadler P.M., Cooper R.A., Crampton J.S. 2019. Completeness of the known

graptoloid palaeontological record. Journal of the Geological Society 176:1038-1055.

Ford D.P., Benson R.B.J. 2020. The phylogeny of early amniotes and the affinities of

Parareptilia and Varanopidae. Nature Ecology and Evolution 4:57-65.

Gavryushkina A., Heath T.A., Ksepka D.T., Stadler T., Welch D., Drummond A.J. 2017.

Bayesian total-evidence dating reveals the recent crown radiation of penguins. Systematic

Biology 66:57-73.

Gavryushkina A., Welch D., Stadler T., Drummond A.J. 2014. Bayesian inference of sampled

ancestor tree for epidemiology and fossil calibration. PLoS Computational Biology

10:e1003919

Gingerich P.D. 1979. Stratophenetic approach to phylogeny reconstruction in vertebrate

paleontology. In Cracraft J., Eldredge N., editors. Phylogenetic analysis and

paleontology. New York: Columbia University Press. p. 41–77.

Heath T.A., Huelsenbeck J.P., Stadler T. 2014. The fossilized birth-death process for coherent

calibration of divergence-time estimates. Proceedings of the National Academy of

Sciences, USA 111:E2957-E2966.

Herrera-Alsina L., van Els P., Etienne R.S. 2018. Detecting the dependence of diversification on

multiple traits from phylogenetic trees and trait data. Systematic Biology, 68:317-328.

Kubo T., Iwasa Y. 1995. Inferring the rates of branching and extinction from molecular

phylogenies. Evolution 49:694-704.

Louca S., McLaughlin A., MacPherson A., Joy J.B., Pennell M.W. 2021. Fundamental

identifiability limits in molecular epidemiology. Molecular Biology and Evolution

38:4010-4024.

Louca S., Pennell M.W. 2020. Extant timetrees are consistent with a myriad of diversification

histories. Nature 580:502-506.

Maddison W.P., R.G. FitzJohn. 2015. The unsolved challenge to phylogenetic correlation tests

for categorical characters. Systematic Biology 64: 127-136.

Maddison W.P., Midford P.E., Otto S.P. 2007. Estimating a binary character's effect on

speciation and extinction. Systematic Biology 56:701–710.

Marsh O.C. Principal characters of American Jurassic dinosaurs. Part IX: The skull and dermal

armor of Stegosaurus. American Journal of Science 34: 413–417.

Matzke N.J, Wright A. 2016. Inferring node dates from tip dates in fossil Canidae: the

importance of tree priors. Biology Letters 12:20160328.

Mitchell J.S., Etienne R.S., Rabosky D.L. 2019. Inferring diversification rate variation from

phylogenies with fossils. Systematic Biology 68:1-18.

Moore B.R., Hohna S., May M.R., Rannala B., Huelsenbeck J.P. 2016. Critically evaluating the

theory and performance of Bayesian analysis of macroevolutionary mixtures.

Proceedings of the National Academy of Sciences, USA 113:9569-9574.

Nakov T., Beaulieu J.M., Alverson A.J. 2019. Diatoms diversify and turn over faster in

freshwater than marine environments. Evolution 73:2497-2511.

Nee S., Holmes E.C., May R.M., Harvey P.H. 1994. Extinction rates can be estimated from

molecular phylogenies. Philosophical Transactions of the Royal Society B 344:77–82.

Osborn H.F. 1905. Tyrannosaurus and other Cretaceous carnivorous dinosaurs. Bulletin

American Museum of Natural History 21:259-265.

Parins-Fukuchi C. 2021. Morphological and phylogeographic evidence for budding speciation:

an example in hominins. Biology Letters 17, doi: https://doi.org/10.1098/rsbl.2020.0754

Parins-Fukushi C., Greiner E., MacLatchy L.M., Fisher D.C. 2019. Phylogeny, ancestors, and

anagenesis in the hominin fossil record. Paleobiology 45:378-393.

Pyron R.A. 2017. Novel approaches for phylogenetic inference from morphological data and

total-evidence dating in squamate reptiles (lizards, snakes, and amphisbaenians).

Systematic Biology 66:38-56.

Rabosky D.L. 2010. Extinction rates should not be estimated from molecular phylogenies.

Evolution 64:1816-1824.

Rabosky D.L., Goldberg E.E. 2015. Model inadequacy and mistaken inferences of trait-

dependent speciation. Systematic Biology 64:340–355.

Rabosky D.L., Goldberg E.E. 2017. FiSSE: A simple nonparametric test for the effects of a

binary character on lineage diversification rates. Evolution, 71:1432–1442.

Sanderson M.J. 2002. Estimating absolute rates of molecular evolution and divergence times: A

penalized likelihood approach. Molecular Biology Evolution 14:1218–1231.

Slater G.J., Goldbogen J.A., Pyenson N.D. 2017. Independent evolution of baleen whale

gigantism linked to Plio-Pleistocene ocean dynamics. Proceedings of the Royal Society B

284:20170546

Stadler T. 2010. Sampling-through-time in birth-death trees. Journal of Theoretical Biology

267:396-404.

Stadler T., Gavryushkina A., Warnock R.C.M., Drummond A.J., Heath T.A. 2018. The fossilized

birth-death model for the analysis of stratigraphic range data under different speciation

modes. Journal of Theoretical Biology 447:41-55.

Vasconcelos T., O'Meara B.C., Beaulieu J.M. 2021. Hidden state-only speciation and extinction

models provide accurate tip-estimates of diversification rates. Submitted.

Wagner P.J. 2019. Phylogenetic analyses and the fossil record: Tests and inferences, hypotheses

and models. Paleobiology 26:341-371.

Wagner P.J., Marcot J.D. 2013. Modelling distributions of fossil sampling rates over time, space

and taxa: assessment and implications for macroevolutionary students. Methods in

Ecology and Evolution 4:703-713.

Zhang C., Stadler T., Klopstein S., Heath T.A., Ronquist F. 2016. Total-evidence dating under

the fossilized birth-death process. Systematic Biology 65:228-249.

 Figure Captions

 Figure 1. a) An example of a fossil set sampled from a complete tree generated by the birth

 death process, which includes both surviving and extinct members. b) The sampled phylogeny

 pruned down from (a) to only includes sampled tips and fossils. The fossilized birth-death model

 distinguishes between two types of sampled lineages. An *m* fossil (denoted by a blue dot)

 represents sampled branches that went completely extinct before the present and did not give rise

 to any additional sampled lineages (i.e., "childless" fossils). A *k* fossil (denoted by an orange

 dot) represents samples of lineages that had sampled descendants (other fossils, or extant tips;

 "parent" fossils).

 Figure 2. Contour plots of the likelihood surface under various scenarios for the same tree

 (topology shown in d) simulated under a birth-death process. Each surface is constrained such

 that turnover rate ($\tau = \lambda + \mu$) and extinction fraction ($\varepsilon = \mu/\lambda$) are set given a pair of fixed

 values, but ψ, the rate of fossil preservation, is free to find their MLE. We sampled 5000 pairs of

 points for τ and ε from a latin hypercube sampling design. When provided with a sample of

 extinct lineages that is perfectly consistent with the generating model, the FDB performs and

 behaves well and generally reduces the variance in turnover rate and extinction fraction (b, c),

 relative to ignoring fossils completely (a). However, if *k* fossils are removed completely (e,f), the

 likelihood surface begins to erroneously shift away from the generating parameters towards

 regions of parameter space of very high extinction rates as values of ψ increase. The dashed

 vertical line represents the generating value for turnover rate (τ=0.70) and the dashed horizontal

 line represents the generating value for extinction fraction (ε=0.75).

Figure 3. Histogram of the fraction of *k*-type fossils sampled across a large set of simulations

over a wide range of conditions in relation to *k*-type fossils used in empirical FBD studies.

Specifically, we generated a grid that make up all possible combinations of elements from

vectors of turnover rate (incrementing τ = 0.1 to 1.0, sampling every 0.1), extinction fraction

(incrementing ε = 0.05 to 0.95, sampling every 0.1), and sampling rate (ψ = 0.1, 0.05, 0.10,

0.20). We repeated each generating parameter 20 times, resulting in 8000 simulation replicates in

total, with the histogram representing the distribution of the fraction of *k*-type fossils sampled

across all replicates. Despite the wide range of simulation conditions, empirical studies (blue

dots) appear to have fewer sampled ancestors. The red line along the x-axis represents the

approximate range of simulated values from Zhang et al. (2016), and the blue dot corresponding

to the study of Zhang et al. (2016) represents the median fraction across the range used in their

analysis of Hymenoptera.

Figure 4. The uncertainty surrounding estimates of turnover rate when the models being

evaluated does not include the true model. Specifically, we assumed that there were four discrete

"regimes" of turnover rates (τ_A= 0.3, τ_B= 0.6, τ_C= 1.5, τ_D= 1.0) and extinction fractions (ε_A= 0.7,

ε_B= 0.9, ε_C= 0.95, ε_D= 0.8) that controls the diversification dynamics; we also assumed a rather

extreme transition dynamics among these four regimes (see text). We also encoded two mass

extinction events as shown by the lineage through time plot (a), and we also assumed biased

sampling among the tips that survived to the present. The boxplots represent the distribution of

model-averaged rates based on the true rate classes at the tips. (b-c) Despite the true model being

far outside the models being evaluated, the turnover rates at the tips hints at there being more

complexity within the data than even the most complex two-rate models in the set might suggest.

However, the inclusion of fossils does not seem to perform remarkably better than not including

them.

Figure 5. (a-b) Contour plots of the likelihood surface for the fossilized birth-death with

stratigraphic ranges using the same tree and data set as shown in Figure 2. When the generating

ψ is low, the likelihood surface includes the generating parameters, but when ψ is increased by

an order of magnitude ($\psi=0.1$), the likelihood surface also begins to drift away from the

generating parameters towards regions of parameter space represented by lower turnover rates

and extinction fraction. (c) A stratigraphic range records the known duration of species on

branches, and so we assume that no unobserved speciation or extinction takes place along these

intervals. Since the same rates apply to all branches, and since turnover rate is a measure of the

total number of speciation and extinction events, as the number of stratigraphic ranges and

duration of each increase, turnover rates will increasingly be underestimated as the proportion of

the stratigraphic ranges relative to the total tree length increases. Interestingly, this does not seem

to impact estimates of extinction fraction (d).