1

**Fossils Do Not Substantially Improve, and May Even Harm, Estimates of Diversification**

**Rate Heterogeneity**

Jeremy M. Beaulieu[1] and Brian C. O'Meara[2]

[1]*Department of Biological Sciences, University of Arkansas, Fayetteville, Arkansas, 72701 USA*

[2]*Department of Ecology and Evolutionary Biology, University of Tennessee, Knoxville,*

*Tennessee, 37996-1610 USA*

[*]Email for correspondence: jmbeauli@uark.edu

*Abstract* — There is a prevailing view that the inclusion of fossil data could remedy

identifiability issues related to models of diversification, by drastically reducing the number of

congruent models. The fossilized birth-death (FBD) model is an appealing way of directly

incorporating fossil information when estimating diversification rates. Here we explore the

benefits of including fossils by implementing and then testing two-types of FBD models in more

complex likelihood-based models that assume multiple rate classes across the tree. We also

assess the impact of severely undersampling, and even not including fossils that represent

samples of lineages that also had sampled descendants (i.e., $k$-type fossils), as well as converting

a fossil set to represent stratigraphic ranges. Under various simulation scenarios, including a

scenario that exists far outside the set of models we evaluated, including fossils rarely

outperforms analyses that exclude them altogether. At best, the inclusion of fossils improves

precision but does not influence bias. We also found that severely undercounting the number of

$k$-type fossils produces highly inflated rates of turnover and extinction fraction. Similarly, we

found that converting the fossil set to stratigraphic ranges results in turnover rates and extinction

fraction estimates that are generally underestimated. While fossils remain essential for

understanding diversification through time, in the specific case of understanding diversification

given an existing, largely modern tree, they are not especially beneficial.

Keywords — fossilized birth-death, state speciation extinction, MiSSE, fossils, stratigraphic

ranges, turnover

Diversification models, while fascinating to biologists, frequently lurch close to

extinction themselves. For instance, Nee et al. (1994) demonstrated that estimating speciation

and extinction rates from a molecular timetree is theoretically possible. This was quickly

followed by work from Kubo and Iwasa (1995), which showed that if rates varied through time

there is an infinite number of alternative sets of time-varying speciation and/or extinction rates

that produce the same number of lineages at any given point in time (an observation largely

ignored at the time). State speciation and extinction models (-SSE; Maddison et al. 2007) were

derived as a seemingly robust framework for estimating the direct effects of discrete traits on

diversification rates. However, Rabosky and Goldberg (2015) found that if a tree evolved under a

heterogeneous branching process, completely independent from the evolution of the focal

character, SSE models will almost always return erroneously strong support for a model of state-

dependent diversification. While this issue was partially rescued by the hidden state models of

Beaulieu and O'Meara (2016), there remains some confusion as to whether SSE models remain a

viable means of assessing state-dependent diversification (Rabosky and Goldberg, 2017; but see

Caetano et al. 2018).

With Louca and Pennell (2020) comes a new salvo of criticism regarding diversification

methods, which echo and expand on the earlier points by Kubo and Iwasa (1995), as well as past

concerns about the possibility of estimating extinction rates generally (i.e., Rabosky, 2010). One

postulated source of salvation has been the inclusion of fossils (Mitchell et al. 2019; Louca and

Pennell, 2020; but see Černý et al. 2021). At a minimum, the inclusion of fossils should

drastically reduce the number of congruent models by excluding any congruent model that

assumes no extinction. We also know that fossils, even without formal models, have contributed

greatly to the understanding of diversification processes — take the discovery of mass

extinctions, for example. For these reasons, the fossilized birth-death (FBD) model (Stadler

2010) is seen as an appealing way of directly incorporating fossil information when estimating

diversification rates, because it naturally assumes that the fossil information represents samples

of extinct lineages in the past, in addition to the species sampled at the present. An important, but

frequently overlooked, property of FBD is that it distinguishes between two types of sampled

lineages (Fig. 1). The first are referred to as *m* fossils, which represent sampled branches that

went completely extinct before the present and did not give rise to any additional sampled

lineages. The second type are referred to as *k* fossils, which represent samples of lineages that

had sampled descendants (other fossils, or extant tips). In other words, these are fossils that

represent ancestors sampled on internal branches that eventually lead to sampled species. The

model assumes a sampling rate of fossils that does not vary across taxa or time — a questionable

assumption, but probably no worse than similar simplifications that speciation rate or extinction

rate is similarly invariant. However, while the model makes this distinction among these fossil

types, the rate by which each is sampled is governed by the same global sampling rate, $\psi$, such

that $\psi_m = \psi_k$.

The linking of these parameters makes sense. When a dinosaur gets washed into a river,

the probability of it being excavated later by a paleontologist does not depend on whether a

descendant seventy million years later is sampled (making our unlucky dinosaur a *k* fossil) or not

(making this an *m* fossil). However, in the model, an assumption is not only that the probability

of sampling the fossil is the same regardless of whether it has descendants or not, but also that

the probability of including it in the phylogeny is the same whether or not it has descendants. For

example, Paleobiology Database (PBDB; accessed October 2021) has 67 collections for

*Tyrannosaurus rex* (Osborn 1905) and only eight for *Stegosaurus stenops* (Marsh 1887). For the

model to fully apply, there should be 67/8 times as many *T. rex* fossils as *S. stenops* fossils

included in the tree.

**Figure 1.** a) An example of a fossil set sampled from a complete tree generated by the birth-
death process, which includes both surviving and extinct members. b) The sampled phylogeny
pruned down from (a) to only includes sampled tips and fossils. The fossilized birth-death model
distinguishes between two types of sampled lineages. An *m* fossil (denoted by a blue dot)
represents sampled branches that went completely extinct before the present and did not give rise
to any additional sampled lineages (i.e., "childless" fossils). A *k* fossil (denoted by an orange
dot) represents samples of lineages that had sampled descendants (other fossils, or extant tips;
"parent" fossils).
103

Of course, there are other biases and variation in fossilization rate (the whole field of

taphonomy studies this rich and varied process), but we are concerned with a potential bias of

experimentalists undersampling *k* fossils. This would have predictable effects on the estimates of

$\psi$ as well as the extinction rate itself. To better understand the dynamics of the different fossil

types, here is Eq. (5) from Stadler (2010), which is probability density of a tree, *T*, with *n* extant

taxa, $m \geq 0$ fossils samples, and $n \geq 0$ samples conditioned on the time since the most recent

common ancestor of all taxa in the tree, $x_1$:

$$f[T|t_{mrca} = x_1] = \frac{\lambda^{n+m+2}\psi^{k+m}}{(1-\hat{p}_0(x_1))^2} p_1(x_1) \prod_{i=1}^{n+m+1} p_1(x_i) \prod_{i=1}^{m} \frac{p_0(y_i)}{p_1(y_i)}$$

For specifics on what the probabilities $p_0(t)$, $p_1(t)$, and $\hat{p_0}(t)$ represent, we refer readers to the

original derivation in Stadler (2010); for the purpose of this discussion, just note that they do not

include $k$. We highlight two important observations here. The first is that the information

provided by the $m$-type samples are used throughout the equation, including the time at which

the extinct tip was sampled, $y_i$, as well as the time at which they split from their common

ancestor, $x_i$,. The second observation is that the only information provided by the $k$-type fossils

is simply how numerous they are in the fossil set. In fact, it is irrelevant where exactly they occur

on a branch or which branches have them — that is, the likelihood is the same if they are

dispersed evenly across the tree or sampled in one 10,000-year interval on one edge. This point is

best illustrated by modifying the numerator in the first term in the equation above to separate out

$k$ and $m$,

$$f[T|t_{mrca} = x_1] = \psi^k \frac{\lambda^{n+m+2}\psi^m}{(1-\hat{p_0}(x_1))^2} p_1(x_1) \prod_{i=1}^{n+m+1} p_1(x_i) \prod_{i=1}^{m} \frac{p_0(y_i)}{p_1(y_i)}.$$

Moving the $\psi^k$ to the front of the equation shows that the effect that $k$-type samples have on the

probability is simply based on a factor of $\psi^k$. What this also says is that when both $m$- and $k$-type

fossils are included, the overall log-likelihood can be calculated as, $logL = logL_{m\ only} + [k *$

$log(\psi)]$. As an example, take the tree and set of sampled extinct lineages presented in Figure 2d.

The overall log-likelihood for the maximum likelihood estimate (MLE) for this tree is -1224.937.

The log-likelihood assuming only $m$-type fossils, but using the same parameter estimates, is

**Figure 2.** Contour plots of the likelihood surface under various scenarios for the same tree
(topology shown in d) simulated under a birth-death process. Each surface is constrained such
that turnover ($\tau = \lambda + \mu$) and extinction fraction ($\varepsilon = \mu/\lambda$) are fixed, but $\psi$, the rate of fossil
preservation, is free to find their MLE given a pair of fixed parameter values. When provided
with a sample of extinct lineages that is perfectly consistent with the generating model, the FDB
performs and behaves well and generally reduces the variance in turnover and extinction fraction
(b, c), relative to ignoring fossils completely (a). However, if $k$ fossils are removed completely
(e,f), the likelihood surface begins to erroneously shift away from the generating parameters
towards regions of parameter space of very high extinction rates as values of $\psi$ increase. The
dashed vertical line represents the generating value for turnover ($\tau$=0.70) and the dashed
horizontal line represents the generating value for extinction fraction ($\varepsilon$=0.75).

-983.6938, and we can add $107*\log(\hat{\psi}=0.1049145)$ to obtain the overall log-likelihood of

-1224.937 (for more details on these calculations see Supplemental Materials).

On the one hand, if a tree was generated by a Yule process (i.e., no extinction), the

inclusion of only $k$-type samples has no effect on the parameter estimates. Their inclusion will

simply slide the overall log-likelihood down, again, by a factor $k*\log(\psi)$. On the other hand, in

the case of non-zero extinction rates, if $m$-type fossils are included and $k$-type not added, this will

have a profound impact on the parameter estimates, particularly with regards to $\psi$ and $\mu$. Under

the FBD formulation, the linking of $\psi_m = \psi_k$ forces the model to interpret the lack of $k$-type

fossils as evidence of a low sampling rate. This will be in tension with the presence of only $m$-

type samples, such that to explain a low sampling rate, but many $m$-type fossils in the set, the

extinction rate must have been substantial. As shown in Figure 2e-f, this is exactly what happens

when the $k$-type fossils are removed from the fossil sample set. The likelihood surface

erroneously shifts away from the generating parameters towards regions of parameter space of

very high extinction rates (based on estimates of turnover, $\lambda + \mu$, and extinction fraction, $\mu/\lambda$).

In any case, when provided with a sample of extinct lineages that is perfectly consistent

with the generating model, the FDB performs and behaves exactly as it should (Fig. 2b-c), but so

does a model that ignores fossils completely (Fig. 2a). Including fossils should help, especially

with extinction rates (as shown by Mitchell et al. 2019). Our concern, however, is what happens

when theory meets practice. It seems to us that aside from a few clades with remarkably rich

fossil records (e.g., terrestrial arthropods, marine invertebrates, cetaceans), a vast majority, at

best, may only have a small handful of fossils that can be placed with any confidence. This

would mean that the decrease in the uncertainty (but likely not the bias) surrounding

diversification rate estimates that include fossils would be small, though the inclusion of $m$-type

fossils would help with our estimation of extinction rates. However, these benefits would quickly

fade when estimating rates using a few fossils in a more parameter-rich model that contains

many discrete changes in rates throughout the tree. We also suspect that, in practice, it is

generally hard to know whether a fossil is best placed exactly on a branch or as an extinct sister

to it given insufficient morphological evidence generally (see Parins-Fukuchi 2021). Many

studies that combine molecular and morphological information to place fossils in the context of

modern taxa, particularly within plant groups, often place fossils as extinct sisters, suggesting

that most data sets probably mistakenly contain far more m-type fossils than k-type ones. While

some studies have begun placing fossils on edges or tips (e.g., Heath et al. 2014; Gavryushkina et

al. 2017), much of the recent work is done in a Bayesian framework where it is hoped this

uncertainty is integrated out with no ill effects. Similarly, a recent extension of the FBD

explicitly treats the same fossil sets as multiple samples of the same species that specify distinct

stratigraphic ranges (Stadler et al. 2018). All that is required, then, is the oldest and youngest

fossil within a range, as opposed to the total number of fossils, which is then marginalized out,

potentially rendering issues related to counting $k$ fossils moot in most cases.

These are the topics we seek to investigate here. Specifically, we explore the benefits of

including fossils in more complex models that assume multiple rate classes across the tree in

comparison to excluding them altogether. We also assess the impact of severely undersampling

and even not including $k$-type fossils into these analyses. In short, will fossils help diversification

analyses largely based on modern phylogenies? Our hypotheses at the outset were that well-

sampled fossils would have a very beneficial effect on estimation of diversification parameters

versus having no fossils and that models that exclude $k$-type fossils would perform poorly, likely

being outcompeted by models with no fossils. We expected approaches with stratigraphic

samples would have less benefit (reduced data, especially about fossilization rate) but also would

be relatively insensitive to sampling issues. We expected that fossils would be especially

valuable in a situation where the true model was more complex than any model used in analysis,

which approximates the situation in reality.

**Fossilized -SSE**

Incorporating the sampling rate, $\psi$, into a birth-death model that allows multiple discrete

shifts has been implemented before (e.g., fossil BAMM, Mitchell et al. 2019). However, here we

incorporate fossils within our hidden state speciation and extinction framework (HiSSE;

Beaulieu and O'Meara, 2016), that includes any number of observed and/or hidden states. We

focus our tests here exclusively using MiSSE (see Vasconcelos et al. 2021), a hidden state only

model, like BAMM, but without priors (which could help, hurt, or have little effect on

inference), but also, importantly, without the discontinuous inheritance of extinction probability

that makes the likelihoods used in BAMM mathematically incorrect (see Moore et al. 2016). It is

relatively straightforward to extend the equations of the canonical FBD model of Stadler (2010)

to the state speciation and extinction models of Maddison et al. (2007) so that they also include

both state transitions and the tree-wide rate of fossil sample rate parameter, $\psi$:

$\frac{dE_i}{dt} = \mu_i - \left(\lambda_i - \mu_i - \psi + \sum_{i \neq j} q_{ij}\right)E_i(t) + \mu_i E_i(t)^2 + \sum_{i \neq j} q_{ij} E_j(t),$

$\frac{dD_i}{dt} = -\left(\lambda_i - \mu_i - \psi + \sum_{i \neq j} q_{ij}\right)D_i(t) + 2\lambda_i D_i(t)E_i(t) + \sum_{i \neq j} q_{ij} D_j(t).$     (1)

The probability $E_i(t)$ is the probability that a lineage starting at time $t$ in state $i$ leaves no

descendants at the present day ($t$=0), and $D_i(t)$ is the probability of a lineage in state $i$ at time $t$

before the present ($t$>0) evolved the exact branching structure as observed. These ordinary

differential equations are generalized so that any number of observed or hidden states can be

included in the model. For character-based models, like HiSSE and MuHiSSE, $i$ and $j$ represent

the different observed and hidden states combinations specified in the model, whereas with

MiSSE $i$ and $j$ represent hidden states only. For an extant tip, the initial condition for $D_i(0)$ is $\rho_i$,

which defines the probability that an extant individual observed in state $i$ is sampled in the tree

and $1 - \rho_i$ for $E_i(0)$. For an $m$-type fossil, $D_i(t)=\psi E_i(t_e)$, and $E_i(t)=E_i(t_e)$, where $t_e$ represents

the time at which the sampled extinct lineage was sampled; for $k$-type fossil, $D_i(t)=\psi D(t)$, and

$E_i(t)=E_i(t)$. At nodes, $N$, the initial condition is the combined probability of its two descendant

branches, $L$ and $R$, such that, $D_{N,i} = D_{L,i}D_{R,i}\lambda_i$. The overall likelihood is the sum of $D_{N,i}$

calculated at the root.

When the fossil set represents stratigraphic ranges, the likelihood calculation is much

more involved and requires designating three specific types of edge segments. First, we note that

our implementation reflects the "symmetric speciation only" portion of Stadler et al. (2018; see

Corollary 13), where only bifurcating speciation events are allowed — that is, a speciation event

reflects the extinction of an ancestral species, and two new descendant species arise. This greatly

reduces the complexity of the original model formulation, which at its most complex, also

includes both anagenetic (ancestral species is replaced by a new descendant) and asymmetric (a

single descendant "buds" off a persistent ancestral lineage) speciation processes with their own

rates. Second, we note that a stratigraphic range is defined only by the oldest ($o$) and youngest

($y$) fossils that are unequivocally assigned to a species across a distinct time interval. Thus, a

stratigraphic interval records the known duration of species on branches. These ranges are

represented by a single fossil and a tip (i.e., fossil is $y=0$), by a branch fossil and an extinct tip

($y=t_e$), by two fossils representing a sampled ancestral stratigraphic range along an edge ($o>y$), or

even by a single fossil (i.e., $o=y$). Fossils that occur within a given range are ignored.

When an edge segment is not associated with a stratigraphic range, the branch calculation

and starting conditions proceed exactly as described in Eq. 1 above. However, when an edge

segment represents a stratigraphic range, we must modify $D_i(t)$ to remove the possibility of an

unobserved speciation and subsequent extinction event within the interval given that we know

that the entire segment [$o,y$] belongs to the same species:

$\frac{d\widehat{D}_i}{dt} = -\left(\lambda_i - \mu_i - \psi + \sum_{i \neq j} q_{ij}\right)\widehat{D}_i(t) + \sum_{i \neq j} q_{ij}\widehat{D}_i(t).$    (2)

When $y=0$, such that it is an extant species, the initial condition for $\widehat{D}_i(0)$ is $\rho_i$; if $y$ is an $m$-type

fossil, $\widehat{D}_i(t)=\psi E_i(t_e)$; if $y$ denotes the beginning of a sampled ancestor stratigraphic range then

$\widehat{D}_i(t)=D(t)$. Finally, for edge segments that represent intervening time intervals between two

stratigraphic ranges without an observed speciation event, we modify the branch calculation to

account for zero or more unobserved speciation events. Again, given the assumption that a

stratigraphic range represents the sampling of a single species across a distinct interval of time,

the presence of two stratigraphic ranges along the same edge would imply that at least one

unobserved speciation event had occurred somewhere within this interval. To account for this, at

the rootward end, $y_a$, which represents the youngest fossil of the older of the older of the two

stratigraphic ranges, we correct $D_i(t)$ following the Stadler et al. (2018):

$D_i(t) = D_i(y_a) * \left(1 - \frac{\widehat{D}_i(y_a)}{D_i(y_a)}\right)$ .    (3)

This probability then becomes the starting condition for the next stratigraphic range. Note that

none of three specific branch types requires altering $E_i(t)$ from what is presented in Eq. 1,

because this probability is based solely on time and is therefore unaffected by the tree topology.

As before, at nodes, $N$, the initial condition is the combined probability of its two descendant

branches, $L$ and $R$, such that, $D_{N,i} = D_{L,i}D_{R,i}\lambda_i$. The overall likelihood is again the sum of $D_{N,i}$

obtained at the root, but we must also marginalize over the number of fossils within a

stratigraphic range. This is done by multiplying the sum of $D_{N,i}$ at the root by $\psi^{K'}$ and $e^{\psi L_s}$,

where $K'$ represents the total number of sampled fossils that represent the start and end times of a

stratigraphic range (if $o=y$ for a given range then this counts as only one fossil), and where $L_s$

represents the sum of all stratigraphic range lengths ($\sum_{i=1}^{n} o_i - y_i$ ).

For both the canonical FBD, and FBD with stratigraphic ranges, we condition the overall

likelihood by $\lambda_i[1 - E_i(t \mid \psi = 0)]^2$, where and $E_i(t \mid \psi = 0)$ is the probability that a

descendant lineage of the root survived to the present and was sampled assuming no sampling in

the past. We note that fossil BAMM does not condition on survival in this way, or even at all,

when the tree includes extant taxa. For character-based models, such as HiSSE and MuHiSSE,

we weight the overall likelihood by the probability that each possible state gave rise to the

observed data (see FitzJohn et al. 2009). However, it was pointed out by Herrera-Alsina et al.

(2018) that at the root, the individual likelihoods for each possible state should be conditioned

prior to averaging the individual likelihoods across states. It is unclear to us which procedure is

correct, but it does seem that both weighting schemes behave exactly as they should in the case

of character-independent diversification — that is, the overall likelihood reduces to the

likelihood of the tree multiplied by the likelihood of a trait model. We have also tested the

behavior of both (not shown), and the likelihood differences are very small, and the parameter

estimates in simulation are nearly indistinguishable from one another.

In the absence of character information, assuming a model with a single birth rate and

death rate, and a $\psi > 0$ the likelihoods are identical to that of Stadler (2010) and Stadler et al.

(2018). To ensure that our math and the implementations are correct with two or more birth and

death rates, we relied on the property of SSE models described by Beaulieu and O'Meara (2016)

and Caetano et al. (2018): when a trait has no differential effect on the diversification process

(i.e., the character-independent model), the overall likelihood is the product of the tree likelihood

and likelihood of a Markov model applied to the character data (it is the sum of the two

likelihoods in log space). Thus, for a binary character that is independent of the diversification

process on a tree with two rate classes and a non-zero fossil preservation rate, the BiSSE

likelihood should reduce to the product of the likelihood of a MiSSE model with two rate classes

and the likelihood of the transition rate model applied in the character data. As we show in the

Supplemental Materials, we can confirm that our implementation of the FBD for MiSSE, HiSSE,

and MuHiSSE (two binary characters; Nakov et al. 2019) are indeed correct. We also show that

the different branch calculations for FBD with stratigraphic ranges conforms to the analytical

calculations provided by Stadler et al. (2018). All models are available in the R package *hisse*

(Beaulieu and O'Meara 2016).

**Impact of Fossils Under Various MiSSE Models**

298           Under a single rate model of birth and death, the inclusion of fossil data reduces

uncertainty, but not necessarily the bias, in the estimates of functions of speciation and

extinction, namely, the turnover rate ($\tau = \lambda + \mu$; Beaulieu and O'Meara 2016) and the extinction

fraction ($\varepsilon = \mu/\lambda$). We explored these results further by conducting additional single rate regime

simulations where trees were generated using a speciation rate of $\lambda$=0.4 and setting the extinction

rate to take on three values, $\mu$= 0.1, 0.2, 0.3, to ensure three different values of extinction

fraction, $\varepsilon$ = 0.25, 0.50, and 0.75, respectively. In each scenario, trees were grown until they

reached 200 taxa. We then used a range of fossil sampling rates ($\psi$=0.01, 0.05, and 0.10

fossils/Myr), as well as varying the number of $k$-type fossils, where either only half were

randomly chosen to be included in the analysis (i.e., half $k$ fossils missing), none were included

at all (i.e., only *m*-type fossils included), or converted to stratigraphic ranges. When converting a

fossil set into a set of stratigraphic ranges, we simply take the oldest and youngest fossil on each

edge, removing all others, prior to pruning unsampled extinct lineages from the full tree.

In all cases, and across three metrics of diversification (turnover, extinction fraction, and

net diversification), when comparing extant-only versus a full fossil sample under a given

sampling scenario, there was minimal improvement in the rate estimates. Perhaps the most

dramatic differences occurred when the extinction fraction was the smallest (i.e., $\varepsilon = 0.25$) and

when the fossil sampling rates were highest ( $\psi=0.10$ fossils/Myr), which resulted in the full

fossil set having both the bias and the variance being an order of magnitude smaller than

estimates from extant-only. However, when extinction fraction was the highest, at $\varepsilon = 0.75$, the

inclusion of fossils had very minor improvements in terms of bias and variance. When a portion

of the *k*-type fossils were removed, there was a notable upward bias, but not the variance, in the

rate estimates (see Table S1), which is consistent with the contour plot shown in Figure 2. We

also note that when the fossil set is converted to stratigraphic ranges, there is rather severe

downward bias in estimates of both turnover rate and extinction fraction as both extinction

fraction and $\psi$ increases. Interestingly, bias in the estimates of net diversification remains

nominal generally across every combination of generating parameters (see Table S1).

By contrast, Mitchell et al (2019) showed that with their fossil implementation within

BAMM, which allows for shifts in discrete regimes of speciation and extinction, there were

substantial improvements to the rate estimates, most notably with the extinction fraction. These

results seem at odds with our results presented thus far. While their simulations were focused on

understanding the behavior of a much more complex model, it should be noted that the most

striking improvement in extinction fraction occurred when $\psi=1$ fossils/Myr, which is an

extremely high rate of preservation. In our case, the maximum value $\psi$=0.1 fossils/Myr resulted

in a fossil set that was at least as numerous as the number of extant taxa ($N$>200), which seemed

reasonable to us. Still, even when the addition of fossils resulted in doubling the amount of data

available to the birth-death model, there was very little impact on the rate estimates, especially

when the generating extinction fraction was high ($\varepsilon$ = 0.75). We also wondered how other

functions of speciation and extinction, such as turnover rates and net diversification rates, are

impacted not only by the inclusion of fossils, but also by the inclusion of biased samples. The

simulations of Mitchell et al. (2019) demonstrated that estimates of speciation are insensitive to

the inclusion of fossils, with extinction being strongly impacted, which suggests that this would

also impact additional functions of speciation and extinction in different ways.

We explored this by performing simulations to test the behavior of our MiSSE model

when fossils are included. As mentioned above, the underlying model is like BAMM in that it

ignores character information altogether. Our "hidden state only" model assumes that a "shift" in

diversification denotes a lineage tracking some unobserved, hidden state. We evaluated the

performance of our MiSSE model by simulating trees under three scenarios and then estimating

the fit and bias of the inferred rates from these trees. The first simulation scenario (scenario 1),

we introduced two turnover rate regimes, where regime $B$ ($\tau_B$=0.50 events/Myr) had a turnover

rate that was 2x the rate of regime $A$ ($\tau_A$=0.25 events/Myr). The extinction fraction was set at

0.75 for both regimes. In the second scenario (scenario 2), the rates were set such that there were

two unique turnover rate and extinction fraction regimes. Specifically, we assumed regime $B$

($\tau_B$=0.50 events/Myr) had a turnover rate that was 2x the rate of regime $A$ ($\tau_A$=0.25 events/Myr),

but where regime $A$ had a lower extinction fraction ($\varepsilon_A$ =0.55) than regime $B$ ($\varepsilon_B$=0.75). With the

turnover rates and extinction fractions structured in this way the net diversification rates between

**Figure 3.** The uncertainty surrounding estimates of turnover ($\tau_A$ and $\tau_B$), extinction fraction ($\varepsilon_A$ and $\varepsilon_B$), and net diversification ($r_A$ and $r_B$) when the generating model assumes: (a-c) regime $B$ had a turnover rate that was 2x the rate of regime $A$ ($\tau_A$=0.25, $\tau_B$=0.50) and extinction fraction was set at 0.75 for both regimes; (d-f) there were two unique turnover rate and extinction fraction regimes, specifically, regime $B$ had a turnover rate that was 2x the rate of regime $A$ ($\tau_A$=0.25, $\tau_B$=0.50), but where regime $A$ had a lower extinction fraction than regime $B$ ($\varepsilon_A$ =0.55, $\varepsilon_B$=0.75); or (g-h) both regime $A$ and $B$ had identical turnover rates ($\tau_A$=$\tau_B$=0.375) and extinction fractions ($\varepsilon_A = \varepsilon_B$=0.75). In all cases, different assumptions about the fossils were fit to the same data set, and parameters were model averaged across a set of four models that variously allowed turnover and extinction fraction to vary (see text). The solid red lines represent the generating value for each parameter in each regime; the right y-axis represent the net diversification scale.

these regimes are nearly identical. Finally, we simulated a scenario (scenario 3) that assumed

both regime $A$ and $B$ had identical turnover rates ($A=B$=0.375 events/Myr) and extinction

fractions ($\varepsilon_A = \varepsilon_B$=0.75). This scenario was meant to examine the impact that including fossils

has on the support for more complex models when the true model contains homogeneous

diversification rates. In all three scenarios we assumed that the transition between regimes $A$ and

$B$ was $q$=0.005 transitions/Myr.

For each simulation replicate, within each scenario, we fit four MiSSE models that

broadly capture the complexity of the scenarios. Specifically, we fit a single regime model, a

model that only allows turnover to vary ($\tau_A \neq \tau_B$), a model that only allowed extinction fraction

to vary, and a model that allowed both to vary ($\tau_A \neq \tau_B$ and $\varepsilon_A \neq \varepsilon_B$). Each model was evaluated

using a two-step optimization routine. The first step consists of a bounded stochastic simulated

annealing run for 5000 iterations, followed by a bounded subplex routine that searches parameter

space until the maximum likelihood is found. Since the state space allowed each model to be

nested within the most complex model where turnover and extinction fraction were allowed to

vary, we model-averaged the rate estimates using the AIC weights. However, before doing so,

we culled the model set to remove any resulting redundant model fits. If the maximum likelihood

estimates for, say, $\tau_A$ and $\tau_B$, in the turnover varying only model, take on the same value, it is

essentially the same as including a single turnover rate model twice, and this would lower the

weight of other models as a consequence. It is recommended in these situations to remove the

more complex of the two from the set (see Burnham and Anderson 2003).

As with the single regime simulations above, the main effect of including fossils is to

reduce the variance of the estimates, not the bias. In fact, scenario 2, which assumed differences

in both turnover and extinction fraction, but not net diversification, proved difficult regardless of

whether fossils were included into the analyses. There was a consistent downward bias in the

rates associated with shifts to regime $B$ even as $\psi$ increased, meaning even when many fossils

were included. With regards to converting the fossil set to stratigraphic ranges, there was, again,

a general tendency for a downward bias in both turnover and extinction fraction. However, when

compared to extant-only and a full fossil set assuming the canonical FBD, the estimates of net

diversification for stratigraphic ranges generally performed better, albeit with higher variance

associated with the estimates compared to the other two analysis types. We do note that with

scenario 3, which assumed no rate differences among regimes $A$ and $B$, the use of stratigraphic

ranges consistently and erroneously inferred higher net diversification rates for regime $A$.

Finally, removing $k$ fossils completely from the set results in very high biases related to turnover

rates specifically, as $\psi$ increases, and rather severe downward biases generally in estimates of net

diversification rates.

**How Do Fossils Help in a World Where All Models are Simply Wrong?**

An easy and valid criticism of the simulations described above is that they are too

simplistic. In other words, trees were generated under an SSE model that shifted between, at

most, two different rate classes, with the rate of these shifts set by a single transition rate, $q$.

While we were genuinely surprised that extant-only trees performed as well as trees that also

included fossil sample data under the same conditions, the simulation scenarios are hardly

realistic. The processes that generate most empirical trees are likely very complex, likely

carrying signatures of non-random extinction, even mass extinctions, with diversification rates

varying substantially among lineages and across time. The simplicity of our MiSSE models is

out of mathematical convenience and tractability. Even still, we might expect that even when the

true model is not included in the set of models evaluated, the inclusion of fossil information

attached to branches distributed across the tree would provide more weight and better parameter

estimation for the more complex models that are included in the set. Most models will work fine

with data that meets their assumptions, but with complex and messy data, maybe fossils are

needed to rescue their performance.

To explore this further we devised a fourth simulation scenario that was meant to closely

imitate processes and conditions that are more realistic in empirical settings, which would also

prove challenging to our MiSSE model, but also be biased in a way that would favor including

fossils. Specifically, we assumed that there were four discrete "regimes" of turnover rates ($\tau_A=$

0.3, $\tau_B=$ 0.6, $\tau_C=$ 1.5, $\tau_D=$ 1.0 events/Myr) and extinction fractions ($\varepsilon_A=$ 0.7, $\varepsilon_B=$ 0.9, $\varepsilon_C=$ 0.95,

$\varepsilon_D=$ 0.8) that controls the diversification dynamics. We also assumed a rather extreme

heterogeneous transition matrix, $\mathbf{Q}$, that governed the dynamics of transitions, $q_{ij}$, among these

four regimes,

425
426 $$\mathbf{Q} = \begin{array}{c} A \\ B \\ C \\ D \end{array} \begin{bmatrix} - & 0.01 & 0.01 & 0.01 \\ 0.05 & - & 0.04 & 0.02 \\ 0.02 & 0.03 & - & 0.03 \\ 0.06 & 0.02 & 0.08 & - \end{bmatrix}$$
427

The magnitude of these rates was chosen somewhat at random, but cumulatively ensured that

approximately 10% of the branches across the entire history of the tree (including extinct

lineages) contained at least one change in rate regime. We also encoded two mass extinction

events. The first occurred after the first 40-time units and removed, at random, 70% of lineages

alive at that time point. The second occurred 30-time units later, 70-time units in total from the

start of the simulation and removed 90% of the lineages alive at the time point (see Fig. 4a).

Finally, simulations were terminated once the tree reached 267 taxa. However, we non-randomly

chose 67 taxa and removed them from the final tree. The purpose of this procedure was to

assume that our final tree had a biased sampling fraction of 75%. The bias was generated by

simulating a trait under Brownian motion, then normalizing the values so that they were between

0 and 1 and had a phylogenetic signal. These values were then used as probabilities for removing

taxa.

We know fossilization rates vary, sometimes dramatically, by time and taxa (Wagner and

Marcott 2013). Incorporating this would make the simulation far more realistic as well, but it

would provide additional complications given the difficulties in separating variable extinction

rates from variable preservation rates (e.g., Foote et al. 2019). The fact that the fossilization

process matches that of the model gives the fossils a chance to perform better than not including

them. That is, can well-modeled fossils help an analysis where the true diversification model is

more complex and different than any analyzed? If fossils cannot help in this case, they probably

will not help. Had we used a more complex fossilization process, and fossils failed to help in that

case, it could just be that we tried too extreme a difference and set up a bias against the utility of

fossils.

Our model set contained the same models as before (though now with $f_i = 0.75$ to account

for incomplete sampling, though assuming, incorrectly, that all taxa have the same sampling rate;

see Beaulieu 2020), thus ensuring that the generating model was not included in the model set.

We also fit the same set of models for increasing values of $\psi$ (i.e., 0.01, 0.05, 0.10

events/Myr), and for different sampling of $k$-type fossils (i.e., full, half $k$, only $m$-type fossils

used), including converting the set to stratigraphic intervals. We chose to summarize results

based on the diversification rates model-averaged across only the tips that survived to the

**Figure 4.** The uncertainty surrounding estimates of turnover when the models being evaluated does not include the true model. Specifically, we assumed that there were four discrete "regimes" of turnover rates ($\tau_A = 0.3$, $\tau_B = 0.6$, $\tau_C = 1.5$, $\tau_D = 1.0$) and extinction fractions ($\varepsilon_A = 0.7$, $\varepsilon_B = 0.9$, $\varepsilon_C = 0.95$, $\varepsilon_D = 0.8$) that controls the diversification dynamics; we also assumed a rather extreme transition dynamics among these four regimes (see text). We also encoded two mass extinction events (a), and we also assumed biased sampling among the tips that survived to the present. The boxplots represent the distribution of model-averaged rates based on the true rate classes at the tips. (b-c) Despite the true model being far outside the models being evaluated, the turnover rates at the tips at hints at there being more complexity within the data than even the most complex two-rate models in the set might suggest. However, the inclusion of fossils does not seem to perform remarkably better than not including them.

present, because the rate regimes in the generating model do not map to the rate regimes in the

model set, namely, regime *D* in a four-rate model does not easily map to regime *A* or *B* in a two-

rate model. For a given model, the marginal probability of each rate regime is obtained for every

tip, and the rates for each regime are averaged together using the marginal probability as a

weight: a weighted average of these rates is then obtained across all models using Akaike

weights. The use of model-averaging is particularly advantageous here because the tip rates are

not conditional on a single best model that would be far less complex than the model that

generated the data. Different models reflect different aspects of the data. By fitting multiple

models, each good model might tell us something about how the data evolved. Model-averaging

is one way to summarize these different elements in a comprehensible way.

Despite the true generating model being far outside the set of models evaluated, the use

of model-averaging at least hints at there being more complexity within the data than even the

most complex two-rate models allow. When the model-averaged tip rates are aligned with the

four true tip rate regimes for each simulation iteration there are consistent significant positive

slopes across all treatments. For fossil-based analyses, particularly for those where the number of

fossils outpaced the number of extant tips (i.e., $\psi = 0.10$ fossils/Myr), there are clearer

differences between the four rate regimes compared to the extant-only estimates. However, as

with all other analyses presented here, when the fossil set is perfectly consistent with the FBD

process by which the samples were taken, the main effect of the fossils is to reduce the overall

uncertainty in the tip rate estimates compared to extant-only inferences.

There were some improvements with respect to the bias with the inclusion of the full

fossil sets overall, most notably at two ends of the rate distribution under the generating model

(i.e., $\tau_A = 0.3$ and $\tau_C = 1.5$ events/Myr). Incidentally, both rate classes were by far the most

frequent states across the entire history of almost every simulated tree. The highest turnover rate

regime also had the highest extinction fraction, and this coupled with several mass extinctions,

resulted in very few survivors at the tips. Even with the addition of a substantial number of

fossils placed throughout the tree, the improvement in inferring this particular rate category from

model-average rates was minimal. From a qualitative standpoint, by ignoring biases in rate and

focusing on comparisons among tips, it seems that all treatments, regardless of the fossil

makeup, can consistently assign the correct sign differences among pairs of tip rates. That is,

even when the trees were analyzed with only half of the $k$-type fossils used, if they were removed

entirely, or converted to stratigraphic ranges, the expected proportion of sign differences are

consistent with the extant-only and full fossil set estimates (Table S2). However, as the number

of *k*-type fossils are removed, the model-averaged estimates of turnover become greatly inflated,

from erroneously high estimates of the extinction, and are associated with much greater

uncertainty. Also, when the fossil sets are converted to stratigraphic ranges, the turnover rates

are, again, generally underestimated, with the improvement of the estimates with the lower

turnover regimes being simply incidental. Taken together, this is consistent with the MiSSE

scenarios above (Fig. 3), as well as the constant birth-death models (Fig. 2e-f), which suggests

that gross omissions of *k*-type samples in a FBD erroneously shifts estimates towards areas of

very high turnover rates and extinction fraction. Moreover, converting to stratigraphic intervals

results in erroneous shifts towards lower turnover rates and extinction fraction.

**A Caution About Stratigraphic Ranges**

Our simulation results highlight a curious pattern in that when a fossil set is converted to

stratigraphic ranges, estimates of turnover and extinction fraction are increasingly

underestimated as the generating $\psi$ increases. A clear illustration of the pattern is shown in the

likelihood surfaces depicted in Figure 5a-b. When the generating $\psi$ is low ($\psi$=0.01 fossils/Myr;

Fig. 5a), the likelihood surface includes the generating parameters. However, when $\psi$ is

increased by an order of magnitude ($\psi$=0.1 fossils/Myr; Fig. 5a-b), the likelihood surface begins

to shift away from the generating parameters towards regions of parameter space represented by

lower turnover rates and extinction fraction. While this downward bias in this particular example

is somewhat subtle, this behavior is consistent across all models and all simulation scenarios

examined here. It should be noted that it is possible that we have incorrectly interpreted the math

described in Stadler et al. (2018). Admittedly, the implementation of this model was quite

challenging, and is further complicated by the fact that FBD with stratigraphic ranges is not

**Figure 5.** (a-b) Contour plots of the likelihood surface for the fossilized birth-death with stratigraphic ranges using the same tree and data set as shown in Figure 2. When the generating $\psi$ is low, the likelihood surface includes the generating parameters, but when $\psi$ is increased by an order of magnitude ($\psi=0.1$), the likelihood surface also begins to drift away from the generating parameters towards regions of parameter space represented by lower turnover rates and extinction fraction. (c) A stratigraphic range records the known duration of species on branches, and so we assume that no unobserved speciation or extinction takes place along these intervals. Since the same rates apply to all branches, and since turnover is a measure of the total number of speciation and extinction events, as the number of stratigraphic ranges and duration of each increase, turnover rates will increasingly be underestimated as the proportion of the stratigraphic ranges relative to the total tree length increases. Interestingly, this does not seem to impact estimates of extinction fraction (d).

naturally nested within the canonical FBD model. That is, if all fossils are treated as single

representatives of a stratigraphic range, the likelihood does not simply revert to the likelihood of

the canonical FBD model. This is because of the assumption about how to correct for unobserved

speciation occurring along an intervening edge segment of two distinct stratigraphic ranges on

the same edge. This makes unit testing of this model challenging. That being said, in

the constant birth-death case, we implemented the model using the analytic equations presented

in Stadler et al. (2018) as well as using the ordinary differential equations above and the

likelihoods were identical across a range of parameter values.

In the absence of an implementation error, we suspect that the underestimation of

turnover is likely due to the assumptions made when computing the probability of the portion of

an edge representing a stratigraphic range. Under this model a stratigraphic range records the

known duration of species on branches, and so we assume that no unobserved speciation or

extinction takes place along these intervals. Since the same rates apply to all branches, and since

turnover is a measure of the total number of speciation and extinction events, we might expect

that as the number of stratigraphic ranges and duration of each increase, the rate estimates would

likely reflect a balance between a large proportion of time in a tree where no unobserved

speciation and extinction events have occurred. Thus, we would predict that the downward bias

in turnover rates would be a function of the proportion of the tree consisting of stratigraphic

ranges. We conducted a simulation where we evolved a set of 100 trees for 50 Myr, assuming the

same generating rates as the ones that produced the trees in Figure 2. We also used a range of

fossil sampling rates starting from $\psi$=0.05 fossils/Myr and increasing at 0.01 increments until

reaching $\psi$=0.15 fossils/Myr. Consistent with our hypothesis, we found that as the proportion of

the stratigraphic ranges relative to the total tree length increased (i.e., stratigraphic ranges

accounted for increasingly more of the time represented in the tree), we found that this was

significantly and negatively associated with an increased downward bias in turnover rates

($R^2$=0.15, slope=-0.272, $P$<0.001). Interestingly, we did not find a similar trend with extinction

fraction ($R^2$<0.001, slope=0.09, $P$=0.246).

**Concluding Remarks**

Our results show that, if the trees fit the generating model well, then adding fossil taxa correctly might help with precision in rate estimations, but not with bias. However, for most cases if there is undersampling of fossils along branches ($k$-type fossils) the results are worse than ignoring fossils altogether, even when the placement of the sampled fossils was correct temporally and topologically. These results occurred in a simulation framework where the diversification models used were far simpler than the simulated data and where we expected fossils to be able to somewhat rescue the results given that they were nearly 3-times more numerous than extant species in certain scenarios (i.e., scenario 4).

We hasten to emphasize that fossils remain key for understanding macroevolution — that is, important extinct groups like trilobites, sauropods, and extinct lycophytes are essential to our understanding of evolution in deep time. However, the idea that neontological studies of diversification is dramatically enhanced by sprinkling carefully chosen fossils on a tree and applying sophisticated fossilized birth-death models to understand rate variation does not seem supported by our simulations. At best, fossils have minor effects. At worst, they lead to less accurate inferences than removing them altogether. We even carefully biased our study to purposely give fossils the best possible chance, namely, by perfectly identifying each fossil, placing them with full certainty in a phylogeny, and assuming that their true sampling rate was constant through time. All this is impossible with empirical data, which is likely the product of variable fossilization rates (Wagner and Marcott 2013) that further complicates teasing apart variable extinction from variable preservation rates (see Foote et al. 2019). Taken together, our simulations provide us with little hope for the utility of fossils in real-world applications, at least with regards to estimating diversification rate heterogeneity across a tree.

One area where we suspect fossils will continue to be of great importance in

neontological applications is in molecular divergence time analyses. Much of the information

used in fitting most diversification models is the timing between events. So, if the relaxed clock

does not properly account for clades with a faster rate of molecular substitution, most methods to

smooth branches will yield ages that are much older than they should be (since the branches are

still too long relative to variation implicit in the clock; see Beaulieu et al. 2015), which will tend

to also decrease the net diversification rate across the tree. However, having more fossils

throughout the tree, we suspect, might alleviate this issue. It thus remains an open question

whether methods that jointly estimate the topology, branching times, and diversification

parameters using an FBD-type model give a better estimate of diversification rate parameters

than alternate approaches (e.g., *r8s*; Sanderson 2002) that infer a chronogram agnostic to the

diversification rate process, which is then used as input for subsequent diversification analyses.

Our study simply demonstrates that in the case of a perfectly accurate (albeit perhaps

undersampled) tree, fossils do not help substantially and, in some circumstances, can hurt.

One motivation for this research, in the light of recent and well-earned concern about

diversification methods using phylogenies (Kubo and Iwasa, 1995; Maddison and FitzJohn,

2015; Rabosky and Goldberg, 2015; Louca and Pennell, 2020) is to help guide empiricists

seeking to create improvements in diversification rate estimation. There is understandable

interest in fossils to improve diversification models. In our own reading of the foundational work

of Stadler (2010), which was motivated by our own curiosity about a path for improvement, we

were struck by the $m$ versus $k$ fossil distinction and what this might mean for its usability in

practice, as well as how stratigraphic intervals might mitigate this issue. We were surprised that,

at least for our particular use case, fossils provided at best little benefit to, and at worst harmed,

inferences of diversification from trees of mostly modern taxa. This does not imply they may not

help substantially in other areas of phylogenetics of extant taxa, but as a solution to the field's

present diversification difficulties, they are less helpful than we hoped, and researchers might be

more fruitfully rewarded attempting other solutions first.

**Funding**

This work was funded by the National Science Foundation grants DEB–1916558 and DEB-

1916539.

**Supplementary Materials**

Data available from the following github repository:

https://github.com/thej022214/Fossils_impact_BO

**Acknowledgements**

We thank members of the Beaulieu and O'Meara labs for their comments and discussions of the

ideas presented here. We would also like to thank Caroline Parins-Fukuchi and James Pease for

edits at various stages of this work.

## References

**References**

Beaulieu J.M. 2020. The problem with clade-specific sampling fractions. Available

at:https://rdrr.io/cran/hisse/f/inst/doc/Clade-specific-sampling.pdf.

Beaulieu J.M., O'Meara B.C. 2016. Detecting hidden diversification shifts in models of trait-

dependent speciation and extinction. Systematic Biology 65:583-601.

Beaulieu J.M., O'Meara B.C., Crane P., Donoghue M.J. 2015. Heterogeneous rates of molecular

evolution and diversification could explain the Triassic age estimate for angiosperms.

Systematic Biology 64:869-878.

Burnham K.P., and Anderson D.R. 2003. Model selection and multimodel inference: a practical

information-theoretic approach. New York:Springer.

Caetano D.S., O'Meara B.C., Beaulieu J.M. 2018. Hidden state models improve state-dependent

diversification approaches, including biogeographic models. Evolution 72:2308-2324.

Černý D., Madzia D., Slater G.J. 2021. Empirical and methodological challenges to the model-

based inference of diversification rates in extinct clades. Systematic Biology

https://doi.org/10.1093/sysbio/syab045.

FitzJohn R.G., Maddison W.P., Otto S.P. 2009. Estimating trait-dependent speciation and

extinction rates from incompletely resolved phylogenies. Systematic Biology 58:595–

611.

Foote M., Sadler P.M., Cooper R.A., Crampton J.S. 2019. Completeness of the known

graptoloid palaeontological record. Journal of the Geological Societ 176:1038-1055.

Gavryushkina A., Heath T.A., Ksepka D.T., Stadler T., Welch D., Drummond A.J. 2017.

Bayesian total-evidence dating reveals the recent crown radiation of penguins. Systematic

Biology 66:57-73.

Heath T.A., Huelsenbeck J.P., Stadler T. 2014. The fossilized birth-death process for coherent

calibration of divergence-time estimates. Proceedings of the National Academy of

Sciences, USA 111:E2957-E2966.

Herrera-Alsina L., van Els P., Etienne R.S. 2018. Detecting the dependence of diversification on

multiple traits from phylogenetic trees and trait data. Systematic Biology, 68:317-328.

Kubo T., Iwasa Y. 1995. Inferring the rates of branching and extinction from molecular

phylogenies. Evolution 49:694-704.

Louca S., Pennell M.W. 2020. Extant timetrees are consistent with a myriad of diversification

histories. Nature 580:502-506.

Maddison W.P., R.G. FitzJohn. 2015. The unsolved challenge to phylogenetic correlation tests

for categorical characters. Systematic Biology 64: 127-136.

Maddison W.P., Midford P.E., Otto S.P. 2007. Estimating a binary character's effect on

speciation and extinction. Systematic Biology 56:701–710.

Marsh O.C. Principal characters of American Jurassic dinosaurs. Part IX: The skull and dermal

armor of Stegosaurus. American Journal of Science 34: 413–417.

Mitchell J.S., Etienne R.S., Rabosky D.L. 2019. Inferring diversification rate variation from

phylogenies with fossils. Systematic Biology 68:1-18.

Moore B.R., Hohna S., May M.R., Rannala B., Huelsenbeck J.P. 2016. Critically evaluating the

theory and performance of Bayesian analysis of macroevolutionary mixtures.

Proceedings of the National Academy of Sciences, USA 113:9569-9574.

Nakov T., Beaulieu J.M., Alverson A.J. 2019. Diatoms diversify and turn over faster in

freshwater than marine environments. Evolution 73:2497-2511.

Nee S., Holmes E.C., May R.M., Harvey P.H. 1994. Extinction rates can be estimated from

molecular phylogenies. Philosophical Transactions of the Royal Society B 344:77–82.

Osborn H.F. 1905. Tyrannosaurus and other Cretaceous carnivorous dinosaurs. Bulletin

American Museum of Natural History 21:259-265.

Parins-Fukuchi C. 2021. Morphological and phylogeographic evidence for budding speciation:

an example in hominins. Biology Letters 17, doi: https://doi.org/10.1098/rsbl.2020.0754

Rabosky D.L. 2010. Extinction rates should not be estimated from molecular phylogenies.

Evolution 64:1816-1824.

Rabosky D.L., Goldberg E.E. 2015. Model inadequacy and mistaken inferences of trait-

dependent speciation. Systematic Biology 64:340–355.

Rabosky D.L., Goldberg E.E. 2017. FiSSE: A simple nonparametric test for the effects of a

binary character on lineage diversification rates. Evolution, 71:1432–1442.

Sanderson M.J. 2002. Estimating absolute rates of molecular evolution and divergence times: A

penalized likelihood approach. Molecular Biology Evolution 14:1218–1231.

Stadler T. 2010. Sampling-through-time in birth-death trees. Journal of Theoretical Biology

267:396-404.

Stadler T., Gavryushkina A., Warnock R.C.M., Drummond A.J., Heath T.A. 2018. The fossilized

birth-death model for the analysis of stratigraphic range data under different speciation

modes. Journal of Theoretical Biology 447:41-55.

Vasconcelos T., O'Meara B.C., Beaulieu J.M. 2021. Hidden state-only speciation and extinction

models provide accurate tip-estimates of diversification rates. Submitted.

Wagner P.J., Marcot J.D. 2013. Modelling distributions of fossil sampling rates over time, space

and taxa: assessment and implications for macroevolutionary students. Methods in

Ecology and Evolution 4:703-713.

**Figure Captions**

**Figure 1.** a) An example of a fossil set sampled from a complete tree generated by the birth-

death process, which includes both surviving and extinct members. b) The sampled phylogeny

pruned down from (a) to only includes sampled tips and fossils. The fossilized birth-death model

distinguishes between two types of sampled lineages. An *m* fossil (denoted by a blue dot)

represents sampled branches that went completely extinct before the present and did not give rise

to any additional sampled lineages (i.e., "childless" fossils). A *k* fossil (denoted by an orange

dot) represents samples of lineages that had sampled descendants (other fossils, or extant tips;

"parent" fossils).

**Figure 2.** Contour plots of the likelihood surface under various scenarios for the same tree

(topology shown in d) simulated under a birth-death process. Each surface is constrained such

that turnover ($\tau = \lambda + \mu$) and extinction fraction ($\varepsilon = \mu/\lambda$) are fixed, but $\psi$, the rate of fossil

preservation, is free to find their MLE given a pair of fixed parameter values. When provided

with a sample of extinct lineages that is perfectly consistent with the generating model, the FDB

performs and behaves well and generally reduces the variance in turnover and extinction fraction

(b, c), relative to ignoring fossils completely (a). However, if *k* fossils are removed completely

(e,f), the likelihood surface begins to erroneously shift away from the generating parameters

towards regions of parameter space of very high extinction rates as values of $\psi$ increase. The

dashed vertical line represents the generating value for turnover ($\tau$=0.70) and the dashed

horizontal line represents the generating value for extinction fraction ($\varepsilon$=0.75).

**Figure 3.** The uncertainty surrounding estimates of turnover ($\tau_A$ and $\tau_B$), extinction fraction ($\varepsilon_A$

and $\varepsilon_B$), and net diversification ($r_A$ and $r_B$) when the generating model assumes: (a-c) regime $B$

had a turnover rate that was 2x the rate of regime $A$ ($\tau_A$=0.25, $\tau_B$=0.50) and extinction fraction

was set at 0.75 for both regimes; (d-f) there were two unique turnover rate and extinction fraction

regimes, specifically, regime $B$ had a turnover rate that was 2x the rate of regime $A$ ($\tau_A$=0.25,

$\tau_B$=0.50), but where regime $A$ had a lower extinction fraction than regime $B$ ($\varepsilon_A$ =0.55, $\varepsilon_B$=0.75);

or (g-h) both regime $A$ and $B$ had identical turnover rates ($\tau_A$=$\tau_B$=0.375) and extinction fractions

($\varepsilon_A$ = $\varepsilon_B$=0.75). In all cases, different assumptions about the fossils were fit to the same data set,

and parameters were model averaged across a set of four models that variously allowed turnover

and extinction fraction to vary (see text). The solid red lines represent the generating value for

each parameter in each regime; the right y-axis represent the net diversification scale.

**Figure 4.** The uncertainty surrounding estimates of turnover when the models being evaluated

does not include the true model. Specifically, we assumed that there were four discrete "regimes"

of turnover rates ($\tau_A$= 0.3, $\tau_B$= 0.6, $\tau_C$= 1.5, $\tau_D$= 1.0) and extinction fractions ($\varepsilon_A$= 0.7, $\varepsilon_B$= 0.9,

$\varepsilon_C$= 0.95, $\varepsilon_D$= 0.8) that controls the diversification dynamics; we also assumed a rather extreme

transition dynamics among these four regimes (see text). We also encoded two mass extinction

events (a), and we also assumed biased sampling among the tips that survived to the present. The

boxplots represent the distribution of model-averaged rates based on the true rate classes at the

tips. (b-c) Despite the true model being far outside the models being evaluated, the turnover rates

at the tips at hints at there being more complexity within the data than even the most complex

two-rate models in the set might suggest. However, the inclusion of fossils does not seem to

perform remarkably better than not including them.

**Figure 5.** (a-b) Contour plots of the likelihood surface for the fossilized birth-death with

stratigraphic ranges using the same tree and data set as shown in Figure 2. When the generating

$\psi$ is low, the likelihood surface includes the generating parameters, but when $\psi$ is increased by

an order of magnitude ($\psi$=0.1), the likelihood surface also begins to drift away from the

generating parameters towards regions of parameter space represented by lower turnover rates

and extinction fraction. (c) A stratigraphic range records the known duration of species on

branches, and so we assume that no unobserved speciation or extinction takes place along these

intervals. Since the same rates apply to all branches, and since turnover is a measure of the total

number of speciation and extinction events, as the number of stratigraphic ranges and duration of

each increase, turnover rates will increasingly be underestimated as the proportion of the

stratigraphic ranges relative to the total tree length increases. Interestingly, this does not seem to

impact estimates of extinction fraction (d).