Article type: Research article

Title: Morphology of immatures of the thelytokous ant, *Monomorium triviale* Wheeler

(Formicidae: Myrmicinae: Solenopsidini) with descriptions of the extraordinary last-

instar queen larvae.

Authors: Naoto Idogawa[1*], Ayako Gotoh[2], and Shigeto Dobata[3]

Affiliation:

[1] Laboratory of Insect Ecology, Graduate School of Agriculture, Kyoto University, Kyoto

606-8502, Japan.

[2] Department of Biology, Faculty of Science and Engineering and Institute for Integrative

Neurobiology, Konan University, 8-9-1 Okamoto, Higashinada-ku, Kobe, 658-8501, Japan.

[3] Department of General Systems Studies, Graduate School of Arts and Sciences, The

University of Tokyo, Komaba, Meguro, Tokyo 153-8902, Japan.

*Corresponding author: E-mail: idogawa.naoto.na@alumni.tsukuba.ac.jp

ORCID digit of the author:

Naoto Idogawa: 0000-0003-4055-8204

Ayako Gotoh: 0000-0002-2885-9971

Shigeto Dobata: 0000-0003-1586-6758

Running title:

SPECTACULAR MORPHOLOGY OF QUEEN LARVAE IN A MONOMORIUM ANT (58

/ 60 words)

Abstract (179 words)

The ant genus *Monomorium* is one of the most species-rich but taxonomically problematic groups in the hyperdiverse subfamily Myrmicinae. An East Asian species, *M. triviale* Wheeler, W. M., produces both reproductive queens and sterile workers via obligate thelytokous parthenogenesis. Here, we describe the immature forms of *M. triviale* based on light and scanning electron microscopy observations, with a note on the striking caste dimorphism in the last larval instar. The last-instar queen larvae were easily recognized by their large size, "aphaenogastoroid" body shape, and rows of doorknob-like tubercles on the lateral and dorsal body surface. This type of queen-specific structure has not been found in ants in general, let alone congeneric species found in Japan. In stark contrast to the queen larvae, worker larvae showed a "pheidoloid" body shape and a body surface similar to other ants. The worker larvae were estimated to have three instars, consistent with previously described congeners. The pupae of both castes also had no cocoon, a characteristic commonly described in other Myrmicinae species. In total, the developmental period from egg to adult worker averaged 59 days under 25°C. We discuss possible functions of the tubercles of queen larvae based on previous studies.

Keywords: larval morphology, Pharaoh ant, parthenogenesis, phenotypic plasticity, protuberance, chaetotaxy

Introduction

Ants (Hymenoptera: Formicidae) are characterized by the division of labor between two phenotypes of females: reproductive queens and sterile workers (Hölldobler & Wilson, 1990). These two castes share the same genetic background and diverge epigenetically in response to socio-environmental factors during immature stages (Rajakumar *et al.* 2018). Furthermore, ant larvae play an important role in colony dynamics and physiological integration by digesting food and distributing nutrients to their nestmates (Dussutour & Simpson 2009; Schultner *et al.* 2017; Masuko 2019). Such behavior emphasizes the importance of the morphological study on the immature stages of ants. However, our knowledge of the immature stages of ants, including number of larval instars, growth rate, and developmental periods is surprisingly limited. Even for the worker caste, complete larval descriptions are available in only 0.5% of >13,000 extant species (Bharti *et al.* 2019; Bolton 2021). The reproductive castes, i.e., queens and males, are usually produced in small numbers during limited seasons, meaning there is even less information on their immature stages.

Here, we provide a detailed description of the immature stages of a thelytokous ant, *Monomorium triviale* Wheeler, with a note on their preimaginal caste dimorphism. *Monomorium* Mayr is a globally distributed and speciose genus in the hyperdiverse subfamily Myrmicinae. Although many species have been transferred to other genera by recent phylogenetic studies (Ward *et al.* 2014; Sparks *et al.* 2019), approximately 300 species have been described and a large number of species have yet to be described (Pontieri & Linksvayer 2019). In this genus, the larval morphology has been described from a dozen of species (Wheeler & Wheeler 1955, 1973, 1980; Bernard 1955; Ettershank 1965), but there is no information on the number of larval instars except for well-studied tramp species, the pharaoh ant, *M. pharaonis* L. (Alvares *et al.* 1993; Pontieri *et al.* bioRxiv) and the flower ant, *M. floricola* Jerdon (Solis *et al.* 2010a). The East Asian species, *M. triviale,* is one of the recently

found parthenogenetic *Monomorium* species (Gotoh *et al*. 2012; Idogawa *et al*. 2021; Ito *et al*.

2021). Males of *M. triviale* have never been reported before, and virgin queens produce both

workers and next-generation queens via thelytoky.

In the present study, the number of larval instars of *M. triviale* were estimated based

on the distributional pattern of body size. The morphological features of each larval instar

were also investigated with light and scanning electron microscopy. Additionally, histological

observation was performed on the newly discovered queen-specific structure. Finally, the

developmental period required for each stage was examined by rearing experiments.

Materials and Methods

Collection of Samples   The nests of *M. triviale* were collected in a thicket consisting

mostly of deciduous broad-leaved trees located in the northern suburbs of Kyoto, Japan

(35°03'33" N, 135°47'01" E, alt. 90 m) from 2017 to 2021. To obtain all stages of immatures,

the nests were transferred into artificial nests in the laboratory and were kept at 25°C.

Determination of Larval Instars   A number of previous studies previous studies on ant

larvae have estimated the number of larval instars from the size distribution pattern of head

capsule width (e.g., Parra & Haddad 1989). However, Masuko (2017) points out that size-

based estimations can overlook the instars with low abundance (especially the first instar). In

the genus *Monomorium*, the shape and distribution of hairs have been reported to be useful for

the identification of larval instars (Solis *et al*. 2010a). Therefore, we estimated the number of

larval instars in *M. triviale* using a combination of morphometrical and chaetotaxical

characteristics.

To bracket the lower and upper growth limit of *M. triviale* larvae, we explicitly

identified the individuals belonging to the first and last instar. Sixty-two larvae with egg

chorion (i.e., just after hatching = definitely first instar) were collected from the egg piles in

the nests. Thirty-five prepupae (i.e., just before pupation = definitely last instar) were also

collected. We photographed these larvae under a digital microscope (VHX-900; Keyence) and

measured the maximum head capsule width using ImageJ software (Schneider *et al*. 2012).

Then, we measured 582 randomly chosen larvae and plotted the distributional pattern of the

head width in a histogram. The distinct peaks of the histogram were detected to find the

intermediate instars between the first and last instar. At least five individuals representative of

each peak were extracted and morphological differences such as body hair number and type

were examined (see next section). Finally, the number of larval instars was determined on the

consensus of qualitative and quantitative traits. The data calculation and visualization was

performed with R ver. 4.0.0 software (R Core Team 2021).

During this experiment, we found worker larvae that were clearly larger than their last

instar larvae. These larvae were reared in the laboratory until their eclosion and confirmed to

be the last instar queen larvae (approx. 50 individuals). We were unable to estimate the

number of larval instars for queens in the above manner because *M. triviale* colonies produce

only a very small number of queen larvae (1–10 individuals) during the early summer.

Observation of the immature stages   To characterize the immature stages, the external

morphology was investigated with a binocular microscope (SZ40; OLYMPUS Optical, Tokyo,

Japan), digital microscope (VHX-900; Keyence, Osaka, Japan), and scanning electron

microscope (SEM: VE-8800; Keyence, Osaka, Japan). For the binocular and digital

microscope observations, living immatures were carefully fixed on the slide glasses using

sticky earthquake-resistant gel pads (Seiwa-pro, Osaka, Japan). For the SEM observation,

fresh samples were frozen in a deep freezer at -20°C for 15 min then quickly mounted on the

aluminum stage with conductive carbon double-sided tape. All observations and photography

were performed as soon as possible after sample preparation. Voucher specimens of eggs,

larvae, pupae, and nestmate imagos were deposited in the Laboratory of Insect Ecology,

Kyoto University, Kyoto, Japan. All terminology used in our larval descriptions follows

Wheeler and Wheeler (1976) and Solis *et al.* (2010a) and measures are given as mean ± SD

followed by the number (*n*) of observations. For comparison of caste dimorphism, three

Monomorium species available in Japan, *M. intrudens* Smith, *M. chinense* Santschi, and *M.*

hiten Terayama, were additionally observed in the same manner.

Since we found caste dimorphism in the external morphology of the last instar larvae,

we examined the histological features of the queen larvae. The larvae were fixed in FAA

(pure ethanol:formaldehyde:acetic acid = 16:6:1) for 24 hours and were dehydrated in a

graded ethanol series before embedding in paraffin. They were cut at 4 μm and stained with

hematoxylin and eosin as described in (Gotoh *et al.* 2016). Histological observations were

performed with Leica DM IL LED inverted contrasting microscope and Leica MC 170 HD

camera (Leica Microsystems, Wetzlar, Germany).

Developmental Periods   Rearing experiments were performed to determine the

developmental periods of each immature stage of *M. triviale*. Due to the difficulty in

distinguishing between first and second instar larvae, these stages are grouped as "young

larvae". From the 18 field-collected source nests, a single queen and 10 nestmate workers

were isolated in an artificial nest for determination of the egg period. The newly oviposited

eggs and hatched larvae were counted daily for approximately 50 days.

For determination of the larval and pupal periods, 10 immatures just before the target

stage (i.e., eggs for young larvae period, young larvae for last instar larvae period, last instar

larvae for pupae period) were isolated with 10 workers. The numbers of individuals belonging

to each stage were counted every day. From the 11 field-collected source nests, 7 artificial

nests were prepared for young larvae, 8 nests for last instar larvae, and 7 nests for pupae.

Observations were continued for a maximum of one month until all individuals molted to the

next stage or died. All experimental nests were isolated in the plastic container (36 × 36 × 14

mm) with gypsum on the bottom and were kept in the laboratory at 25°C. The water and food

(mealworms, *Tenebrio molitor* L. cut into approximately 5 mm length) was replenished every

3 days.

Results

Determination of Larval Instars

The head capsule width of larvae just after hatching (i.e., definitely first instar; mean

$\pm$ SD = 0.123 $\pm$ 0.007 mm, n = 62) and prepupae (i.e., definitely last instar; 0.173 $\pm$ 0.006 mm,

n = 35) bracket the lower and upper limit of the larval head width. Between these limits, an

intermediate peak was detected around 0.135–0.140 mm in the histogram of larval head width

(n = 590, bin width = 0.005 mm), suggesting the presence of an additional larval instar.

Furthermore, the number and shape of larval hairs in these three head width ranges were

clearly different; first instar: 100–150 simple smooth hairs, intermediate: 500–600 simple

smooth hairs, last instar: 400–500 anchor-tipped hairs. According to the agreement between

morphometrical and chaetotaxical features, we estimated that there are three instars of *M.*

triviale worker larvae (Table 1). The larval head width of first (0.122 $\pm$ 0.006 mm, range:

0.107–0.137 mm, n = 84), second (0.137 $\pm$ 0.006 mm, range: 0.124–0.152 mm, n = 226) and

third (0.175 $\pm$ 0.008 mm, range: 0.146–0.191 mm, n = 116) instar larvae overlapped (Fig. 1).

The average growth rate between the larval instars was 1.20 (1.12 between the first and

second instars; 1.27 between the second and third instars).

Description of the immatures of *M. triviale*

Egg

The egg is oval in profile and presents a whitish translucent chorion (Fig. 2A). Length $0.31 \pm$

0.02 mm, range 0.24–0.35 mm, width 0.19 $\pm$ 0.01 mm, range 0.16–.022 mm (all n = 183).

The length:width ratio for the species is 1.62 on average.

First instar

Body: Whitish, profile 'pheidoloid' *sensu* Wheeler & Wheeler (1976) which is characterized by short and straight abdomen, short prothorax, and ends rounded to ventral direction (Fig. 2B); distinct anterior somites; head subcircular; anus subterminal (Fig. 2C). Body 0.328 ± 0.036 mm long (0.237–0.416 mm), 0.161 ± 0.014 mm wide (0.123–0.20 mm, $n = 84$). Length through spiracles 0.44 ± 0.34 mm ($n = 6$). Body hair (varying 100–150 in number; $n = 5$) unbranched, 8.4 ± 2.0 µm long ($n = 25$ of 5 individuals), distance between two adjacent hairs 25.6 ± 9.1 µm ($n = 25$ from 5 individuals, Fig. 2D). Body with 10 spiracles; distance between two spiracles 29.2 ± 9.0 µm ($n = 5$). Spiracle opening unornamented; diameter of the first spiracle 0.8 ± 0.3, other spiracles 0.5 ± 0.2 µm ($n = 5$).

Head capsule: Cranium subcircular shaped, 0.125 ± 0.011 mm in height (0.091–0.149 mm), 0.122 ± 0.006 mm in width (0.107–0.137 mm, $n = 84$); cranium index = 102.3 ± 9.6. Head surface smooth with 26 unbranched hairs: 4 along the ventral border of the clypeus; 12 hairs on each gena; 4 on frons; 2 on vertex; and 4 along the occipital border. Length of hairs; 8.7 ± 1.9 µm ($n = 25$ of 5 individuals).

Mouthparts: Clypeus clearly defined, with no sensilla (Fig. 2E); labrum bilobed, 62.1 ± 1.7 µm wide ($n = 5$), with six sensilla on the anterior surface. Mandibles subtriangular with broad base in shape, blade ending in an apical tooth, lacking medial teeth (Fig. 2E), 0.034 ± 0.002 mm long (0.029–0.038 mm, $n = 9$). Maxilla conoidal in shape, 36.1 ± 2.1 µm in width ($n = 10$). Galea with two sensilla. Maxillary palpus skewed peg with two sensilla. Labium elliptical, 52.3 ± 2.8 µm wide ($n = 5$). Labial palpus with a sensillum on the top.

Second instar

Body: Whitish, profile pheidoloid in shape (Fig. 3A); distinct anterior somites; head

subcircular; anus subterminal (Fig. 3B). Body about 0.459 ± 0.074 mm long (0.281–0.593

mm), 0.189 ± 0.021 mm wide (0.149–0.250 mm, $n = 226$). Length through spiracles $0.77 \pm$

0.18 mm ($n = 5$). Body hair (varying 500–600 in number; $n = 5$) unbranched, 6.8 ± 2.0 μm

long ($n = 25$ of 5 individuals), distance between two adjacent hairs 13.8 ± 2.9 μm ($n = 25$

from 5 individuals, Fig. 3C). Body with 10 spiracles; distance between two spiracles $46.8 \pm$

4.8 μm ($n = 5$). Spiracle opening unornamented; diameter of the first spiracle 2.1 ± 0.3, other

spiracles 0.6 ± 0.1 μm ($n = 5$).

Head capsule: Cranium subcircular shaped, 0.146 ± 0.016 mm in height (0.108– 0.189 mm),

0.137 ± 0.006 mm in width (0.124–0.152 mm, $n = 226$); cranium index $= 106.4 \pm 10.5$. Head

surface smooth with 26 unbranched hairs: 4 along the ventral border of the clypeus; 12 hairs

on each gena; 4 on frons; 2 on vertex; and 4 along the occipital border. Length of hairs; $12.9 \pm$

3.3 μm ($n = 25$ of 5 individuals).

Mouthparts: Clypeus clearly defined, with no sensilla; Labrum bilobed, 67.3 ± 1.3 μm wide

($n = 5$, Fig. 3D), with six sensilla on the anterior surface. Mandibles 'ectatommoid' as defined

by Wheeler & Wheeler (1976): "Subtriangular; with a medial blade arising from the anterior

surface and bearing one or two medial teeth; apex curved medially to form a tooth" in shape,

sclerotized and bearing at least two teeth, 0.042 ± 0.002 mm long (0.039–0.046 mm, $n = 13$).

Maxilla conoidal in shape, 40.1 ± 1.6 μm in width ($n = 8$). Galea with two sensilla. Maxillary

palpus skewed peg with two sensilla. Labium elliptical, 51.5 ± 2.5 μm wide ($n = 5$). Labial

palpus with a sensillum on the top.

Third instar

Body: Whitish, profile pheidoloid in shape (Fig. 4A); distinct anterior somites (Fig. 4B); head

subcircular (Fig. 4C); anus subterminal. Body about 0.847 ± 0.192 mm long (0.414–1.20 mm),

0.360 ± 0.087 mm wide (0.192–0.608 mm, $n = 164$). Length through spiracles 0.95 ± 0.05

mm ($n = 5$). Body hair (varying 400–500 in number; $n = 5$) deeply branched, 32.5 ± 4.4 μm

long ($n = 25$ of 5 individuals), distance between two adjacent hairs 26.4 ± 5.3 μm ($n = 25$

from 5 individuals, Fig. 4D). Body with 10 spiracles; distance between two spiracles $57.8 \pm$

9.2 μm ($n = 5$). Spiracle opening unornamented; diameter of the first spiracle 2.2 ± 0.4, other

spiracles 0.7 ± 0.2 μm ($n = 5$, Fig. 4E).

Head capsule: Cranium subcircular shaped, 0.189 ± 0.017 mm in height (0.134–0.227 mm),

0.175 ± 0.008 mm in width (0.146–0.191 mm, $n = 164$); cranium index $= 108.3 \pm 8.0$. Head

surface smooth with Twenty-six branched hairs: 4 along the ventral border of the clypeus; 12

hairs on each gena; 4 on frons; 2 on vertex; and 4 along the occipital border. Length of hairs;

29.1 ± 4.7 μm ($n = 25$ of 5 individuals).

Mouthparts: Clypeus clearly defined, with no sensilla (Fig. 4D); Labrum bilobed, 65.5 ± 1.3

μm wide ($n = 5$), with six sensilla on the anterior surface. Mandibles ectatommoid in shape,

with three distinct teeth, more sclerotized than those of the second instar, 0.067 ± 0.003 mm

long (0.063–0.071 mm, $n = 5$). Maxilla conoidal in shape, 46.9 ± 1.2 μm in width ($n = 8$).

Galea with two sensilla. Maxillary palpus skewed peg with two sensilla. Labium elliptical,

55.5 ± 4.2 μm wide ($n = 5$). Labial palpus with a sensillum on the top.

Last instar queen

Body: Whitish, profile 'aphaenogastroid' *sensu* Wheeler & Wheeler (1976) which is

characterized by constriction between thorax and abdomen (Fig. 5A); distinct anterior

somites; head subcircular; anus subterminal (Fig. 5B). Body about 1.756 ± 0.329 mm long

(0.742–2.291 mm), 0.873 ± 0.194 mm wide (0.408–1.214 mm, $n = 116$). Length through

spiracles 1.8–2.1 mm ($n = 2$). Abdominal somites hairless (Fig 5A-C), prothoracic body hair

(varying 50–70 in number; $n = 2$) unbranched, 52.8 ± 14.9 µm long ($n = 20$ of 2 individuals,

Fig. 5D, E), distance between two adjacent hairs 25.3 ± 5.9 µm ($n = 20$ from 2 individuals).

Body with 10 spiracles; distance between two spiracles 118.5–213.4 µm ($n = 1$). Spiracle

opening unornamented; diameter of the first spiracle 2.7–3.1, other spiracles 1.3–1.8 µm ($n =$

2).

Head capsule: Cranium subcircular shaped (Fig. 5D), 0.194 ± 0.027 mm in height (0.096–

0.256 mm), 0.187 ± 0.013 mm in width (0.16–0.220 mm, $n = 116$); cranium index = 103.9 ±

12.7. Head surface smooth with Twenty-six unbranched hairs: 4 along the ventral border of

the clypeus; 12 hairs on each gena; 4 on frons; 2 on vertex; and 4 along the occipital border.

Length of hairs; 28.3 ± 9.3 µm ($n = 20$ of 2 individuals).

Mouthparts: Clypeus clearly defined, with no sensilla; Labrum bilobed, 77.5 ± 2.2 µm wide

($n = 5$), with six sensilla on the anterior surface. Mandibles ectatommoid in shape, with 3

teeth, 0.069–0.070 mm long ($n = 2$). Maxilla conoidal in shape, 51.8 ± 3.4 µm in width ($n = 8$).

Galea with two sensilla. Maxillary palpus skewed peg with two sensilla. Labium elliptical,

59.1 ± 2.2 µm wide ($n = 5$). Labial palpus with a sensillum on the top.

Protuberances: Total 37 protuberances; 9 pairs on dorsal, 6 pairs on lateral and unpaired 7

on mid-ventral (Table 2). Paired doorknob-like dorsal tubercles on 2–3rd thoracic and 1–4th

abdominal somites, 72.3 ± 9.4 µm diameter, 134.0 ± 52.7 µm long ($n = 12$). Lateral doorknob-

like tubercles on 2–3rd thoracic and 1–7th abdominal somites, 73.4 ± 9.9 µm diameter, 103.7

± 33.9 µm long ($n = 36$). Ventral protuberances "bosses" as defined by Wheeler & Wheeler

(1976); "an elevated structure with a rounded terminus" in shape. Anterior four bosses on the

3rd thoracic and 1–3rd abdominal somites clearly defined by outlines, 260.7 ± 95.0 µm wide,

105.4 ± 40.5 µm long, 116.5 ± 35.9 µm high ($n = 8$). Posterior three bosses on the 4–6th

abdominal somites slightly raised from body without distinct border. All protuberances

surface uneven with minute papillae (Fig. 5F, G). We did not find any opening or duct-like

structures on the surface of protuberances.

Pupae

Exarate, with no cocoon, 0.99–1.12 mm long in workers ($n = 10$, Fig. 6A) and 2.20–2.49 mm

in queens ($n = 2$, Fig. 6B), milky-white when young; eyes becoming black and body gradually

darkened to a yellowish-brown as they develop into imagos. Queen wings vestigial (i.e.,

brachypterous, Fig. 6C).

Caste dimorphism in congeneric species

The last instar larvae of the three observed *Monomorium* species, *M. hiten, M.*

intrudens and *M. chinense* shared distinct anterior somites, subcircular head shape and

subterminal anus. The worker larvae of all three species indicated the pheidoloid shape in

profile and were covered with anchor-shaped hairs on the thoracic and abdominal somites

(Fig. 7A–C). In *M. hiten*, the larvae of reproductive males and females (= queens) also had

pheidoloid shape (Fig. 7D). However, in *M. intrudens* (Fig. 7E) and *M. chinense* (Fig. 7F), the

reproductive larvae indicated "attoid" shape as defined by Wheeler & Wheeler (1976);

"diameter approximately equal to distance from labium to anus" in profile. The anchor-

shaped hairs were not found in the reproductive larvae of all three species.

Internal morphology of last instar queen larvae

Histological observations of the queen larva showed that the dorsal and lateral

doorknob-like tubercles (Fig. 8A, B) and ventral bosses (Fig. 8C, D) were formed by

extended epidermis and cuticle. There are no muscles or innervation inside the dorsal and

lateral tubercles and ventral bosses. The thickness of the cuticle and epidermis was 8.6 ± 1.8

μm (n = 7) and 11.6 ± 3.3 μm (n = 7) in the doorknob-like tubercles, two times thicker than

the other part of the larval body, which was 4.2 ± 0.2 μm (n = 9) and 4.9 ± 1.0 μm (n = 9).

The ventral bosses were formed by slightly thicker cuticle 5.78 ± 1.11 μm (n = 5) and

markedly thicker epidermis 16.42 ± 4.09 μm (n = 5). In the ventral bosses, the small cells

(possibly fat bodies) surrounded by epidermis were observed (Fig. 8D). We did not find any

specialized cells and duct-like structures in the doorknob-like tubercles and in the ventral

bosses.

Developmental periods

During the 53-day experimental period, 211 eggs and 166 first instar larvae appeared

in the artificial nests (Table 3). The average duration between oviposition and hatching was

13.39 ± 2.28 days. Out of 70 induced eggs, 32 young (= first and second instar) larvae

hatched and all successfully turned into the last instar larvae in an average of 7.31 ± 1.00 days.

Seventy-two of 80 induced young larvae molted to the last instar larvae and 51 of them turned

into prepupae in 18.31 ± 3.29 days, and all molted to pupae in 4.43 ± 1.10 days. Thirty-three

of 70 induced last instar larvae turned to pupae and 22 individuals successfully emerged as

worker imago in 16.87 ± 1.55 days. In total, the overall developmental period of *M. triviale*

worker was estimated to be 59 days.

Discussion

Based on the combination of the body size and body hair features, three larval instars

were estimated for *M. triviale*. This is the general lowest number of larval instars in

Formicidae reported from four subfamilies; Dorylinae, Ectatomminae, Formicinae and

Myrmicinae (Solis *et al.* 2010a). In Myrmicinae, three larval instars were known from 16

species including the congeners, *M. pharaonis* and *M. floricola* (Solis *et al.* 2010a; Bharti &

Gill 2011; Bharti *et al.* 2019). In *M. triviale*, calculated growth rates between the instars are

consistent with the Dyar principle of a constant ratio of growth by molting between 1.1 and

1.9 (Parra & Haddad, 1989). The growth rate during the larval period of *M. triviale* was 1.19,

which is smaller than that of *M. floricola* (1.23; Solis *et al.* 2010a) and *M. pharaonis* (1.36;

Alvares *et al.* 1993). This difference may reflect the difference in body size of adult workers,

ca. 1.5 mm for *M. triviale*, 1.7–2.0 mm for *M. floricola* (Wetterer 2010a), and 2.2–2.4 mm for

M. pharaonis (Wetterer 2010b).

The total developmental period of *M. triviale* was estimated to be about 59 days (egg:

13 days + larva: 29 days + pupa: 17 days). These results are slightly longer than those of the

tropical congeners *M. pharaonis* (egg: 11 days + larva: 22 days + pupa: 12 days, 27°C;

Pontieri *et al.* bioRxiv) and *M. hiten* (egg: 12 days + larva: 17 days + pupa: 20 days, 24°C; Ito

et al. 2021). It should be noted that the larval period of *M. triviale* may be extended under

natural conditions because the larvae can stop their development and overwinter in the

temperate zone of Japan.

Adams *et al.* (2021) suggested that the combination of quantitative and binary traits is

effective in distinguishing the sex and caste of ant larvae. Our results showed that this

approach is also viable in *M. triviale*. Although the body size of each larval instar overlapped

(Fig.1), three larval instars could be separated by the number and shape of their body hairs

(Table 1). The larvae of some *Monomorium* species have deeply branched anchor-shaped

hairs (Solis *et al.* 2010a; Penick *et al.* 2012). In *M. triviale*, the third instar worker larvae

possess these anchor-shaped hairs all over their body. On the other hand, the last instar larvae

of the queen have very few simple hairs only on the prothorax. Moreover, the three

congeneric species *M. intrudens*, *M. hiten* and *M. chinense* also showed the same caste-

specific pattern: worker larvae had branched hairs, but reproductive larvae were almost

hairless. Such caste dimorphism of body hairs was previously known in *M. minimum*

(Wheeler & Wheeler 1955) and *M. pharaonis* (Edwards 1991). The absence of anchor-tipped

hairs on the last instar could be a useful trait to distinguish the reproductive larvae of

Monomorium species.

The caste dimorphism in the last larval instar of *M. triviale* is more striking than in

other observed congeneric species. The queen larvae of *M. triviale* have "aphaenogastroid"

shape in profile, the lateral and dorsal rows of doorknob-like tubercles on thoracic and

abdominal segments and semi-elliptical protrusion on ventral abdominal segments. The

distinct segmental patterns of these structures among lateral, dorsal and ventral sides,

especially their absence on the first thoracic segment and the terminal abdominal segments

(Table 2), might provide a good opportunity to investigate the caste-specific developmental

processes.

Unusual forms of ant larvae have been described from diverse subfamilies (Table 4).

In the Myrmicinae, the caste dimorphism similar to that in *M. triviale* has been reported in

Crematogaster species. Some late-instar larvae of *C. rivai* var. *luctuosa* and *C. scutellaris*

have rows of lateral protuberances on the abdominal segments (Menozzi 1930; Eidmann

1926), and it is speculated that these larvae differentiate into queens (Casevitz-Weulersse

1984). Contrastingly, in some genera of Dolichoderinae, protuberances have been found only

on worker larvae (e. g. Solis *et al.* 2010b). Although their developmental homology remains

unknown, this phylogenetic distribution would be suggestive of their evolutionary

convergence.

Wheeler & Wheeler (1976) proposed five possible functions of protuberances in the

ant larvae: (1) support of the body position and direction (2) defense against cannibalism

among larvae (3) attachments to ceilings and walls of the nest (4) organs for trophallaxis with

adult workers (5) structures for holding food on the body surface. These hypotheses have not

been sufficiently tested, with only a few exceptions such as Masuko (2019) and Fox *et al.*

(2017). Our morphological and histological observations could not find any specialized

structures such as duct-like openings and secretory cells in the protuberances of *M. triviale*

queen larvae. As in many other cases, the function of queen-specific tubercles of the *M.*

triviale larvae is still unclear at this time. Behavioral observations of the interaction between

the workers and the queen larvae are required in the future. Further examination of the

function of these unusual structures of larvae will help us understand the hidden but essential

roles larvae play in complex ant societies.

Acknowledgments

We are grateful to Kenji Matsuura who allowed us to use his laboratory. We thank

Fuminori Ito for providing his *M. hiten* nest, Tomoya Kamada for his help during histological

observation, and Matthew Kamiyama for his English corrections of this manuscript. Finally,

we also thank Christian Peeters and Keiichi Masuko for their valuable comments on larval

protuberances. This work was supported by a Japan Society for the Promotion of Science

(JSPS) Research Fellowship for Young Scientists to NI (19J22242), JSPS KAKENHI Grant

Number JP20K06080 to AG and a grant from the Secom Science and Technology Foundation

to SD.

References

Adams, R.M.M., Larsen, R.S., Stylianidi, N., Cheung, D., Qiu, B., Murray, S.K., Zhang, G. &

Boomsma, J.J. (2021) Hairs distinguish castes and sexes: identifying the early

ontogenetic building blocks of a fungus-farming superorganism (Hymenoptera:

Formicidae). *Myrmecological News*, 31, 201–216.

https://doi.org/10.25849/MYRMECOL.NEWS_031:201

Alvares, L.E., Bueno, O.C. & Fowler, H.G. (1993) Larval instars and immature development

of a Brazilian population of pharaoh's ant, *Monomorium pharaonis* (L.) (Hym.,

Formicidae). *Journal of Applied Entomology*, 116, 90–93.

https://doi.org/10.1111/j.1439-0418.1993.tb01171.x

Bernard, F. (1955) Morphologie et comportement des fourmis lestobiotiques du genre

Epixenus Emery. *Insectes Sociaux*, 2, 273–283.

https://doi.org/10.1007/BF02331506

Bharti, H. & Gill, A. (2011) SEM studies on immature stages of *Pheidole indica* Mayr, 1879

(Hymenoptera: Formicidae) from India. *Halteres*, 3, 38–44.

Bharti, H., Kaur, P. & Bharti, M. (2019) Description of the ant larvae of two species of the

genus *Myrmica* Latreille, 1804 (Hymenoptera: Formicidae) from Indian Himalayas.

Asian Myrmecology, 11, e011004.

https://doi.org/10.20362/am.011004

Bolton, B. (2021) An online catalog of the ants of the world. Available from: http://antcat.org

(accessed 31 October 2021)

Casevitz-Weulersse, J. (1984) Les larves à expansions latérales de *Crematogaster*

(Acrocoelia) *scutellaris* (Olivier) (Hym. Formicidae). *Actes des Colloques Insectes*

Sociaux, 1, 131–138.

Dussutour, A. & Simpson, S.J. (2009) Communal nutrition in ants. *Current Biology*, 19, 740–

744.

https://doi.org/10.1016/j.cub.2009.03.015

Edwards, J.P. (1991) Caste regulation in the pharaoh's ant *Monomorium pharaonis*:

recognition and cannibalism of sexual brood by workers. *Physiological Entomology*, 16,

263–271.

https://doi.org/10.1111/j.1365-3032.1991.tb00565.x

Eidmann, H. (1926) Die ameisenfauna der balearen. *Zoomorphology*, 6, 694–742.

https://doi.org/10.1007/BF00409039

Ettershank, G. (1965) A New Species of *Megalomyrmex* from the Chilean Andes (Formicidae,

Hymenoptera). *Psyche*, 72, 55–58.

https://doi.org/10.1155/1965/32130

Fox, E.G.P., Smith, A.A., Gibson, J.C. & Solis, D.R. (2017) Larvae of trap-jaw ants,

Odontomachus LATREILLE, 1804 (Hymenoptera: Formicidae): morphology and

biological notes. *Myrmecological News*, 25, 17–28.

https://doi.org/10.25849/myrmecol.news_025:017

Gotoh, A., Billen, J., Hashim, R. & Ito, F. (2016) Degeneration patterns of the worker

spermatheca during morphogenesis in ants (Hymenoptera: Formicidae). *Evolution &*

Development, 18, 96–104.

https://doi.org/10.1111/ede.12182

Gotoh, A., Billen, J., Tsuji, K., Sasaki, T. & Ito, F. (2012) Histological study of the

spermatheca in three thelytokous parthenogenetic ant species, *Pristomyrmex punctatus*,

Pyramica membranifera and *Monomorium triviale* (Hymenoptera: Formicidae). *Acta*

Zoologica, 93, 200–207.

https://doi.org/10.1111/j.1463-6395.2010.00498.x

Hölldobler, B. & Wilson, E.O. (1990) *The Ants*, Harvard University Press, Cambridge, MA,

732 pp.

Idogawa, N., Sasaki, T., Tsuji, K. & Dobata, S. (2021) Comprehensive analysis of male-free

reproduction in *Monomorium triviale* (Formicidae: Myrmicinae). *PLOS ONE*, 16(4),

e0246710.

https://doi.org/10.1371/journal.pone.0246710

Ito, F., Makita, S., Nakao, H., Hosokawa, R., Kikuchi, T. & Yamane, S. (2021) Thelytokous

parthenogenesis by dealate queens in the myrmicine ant *Monomorium hiten* distributed

in Nansei Islands, western Japan, with description of the male. *Asian Myrmecology*, 14,

e014001.

Masuko, K. (1989) Larval hemolymph feeding in the ant *Leptanilla japonica* by use of a

specialized duct organ, the "larval hemolymph tap" (Hymenoptera: Formicidae).

Behavioral Ecology and Sociobiology, 24, 127–132.

https://doi.org/10.1007/BF00299644

Masuko, K. (2017) Larval instars of the ant *Strumigenys solifontis* Brown (Hymenoptera:

Formicidae): the fallacy of size distribution. *Journal of Natural History*, 51, 115–126.

https://dx.doi.org/10.1080/00222933.2016.1254299

Masuko, K. (2019) Larval hemolymph feeding and hemolymph taps in the ant *Proceratium*

itoi (Hymenoptera: Formicidae). *Myrmecological News*, 29, 21–34.

https://doi.org/10.25849/myrmecol.news_029:021

Menozzi, C. (1930) Formiche della Somalia italiana meridionale. *Memorie della Società*

Entomologica Italiana, 9, 76–130.

Parra, J.R.P. & Haddad, G.M.L. (1989) Determinação do número de instares de insetos.

Piracicaba, Fundação de Estudos. Agrários Luiz de Queiroz, 49 p.

Peeters, C. & Hölldobler, B. (1992) Notes on the morphology of the sticky "doorknobs" of

larvae in an Australian *Hypoponera* sp. (Formicidae; Ponerinae). *Psyche*, 99, 23–30.

https://doi.org/10.1155/1992/96238

Penick, C.A., Copple, R.N., Mendez, R.A. & Smith, A.A. (2012) The role of anchor-tipped

larval hairs in the organization of ant colonies. *PLOS ONE*, 7, e41595.

https://doi.org/10.1371/journal.pone.0041595

Petralia, R.S. & Vinson, S.B. (1979) Comparative anatomy of the ventral region of ant larvae,

and its relation to feeding behavior. *Psyche*, 86, 375–394.

https://doi.org/10.1155/1979/70316

Pontieri, L. & Linksvayer, T.A. (2019) *Monomorium*. In: *Encyclopedia of Social Insects*.

Springer International Publishing, Cham, pp. 1–6.

https://doi.org/10.1007/978-3-319-90306-4_171-1

Pontieri, L., Rajakumar, A., Rafiqi, A.M. & Larsen, R.S. (2020) From egg to adult: a

developmental table of the ant *Monomorium pharaonis*. bioRxiv.

https://doi.org/10.1101/2020.12.22.423970

Rajakumar, R., Koch, S., Couture, M., Favé, M.-J., Lillico-Ouachour, A., Chen, T., De Blasis,

G., Rajakumar, A., Ouellette, D. & Abouheif, E. (2018) Social regulation of a

rudimentary organ generates complex worker-caste systems in ants. *Nature*, 562, 574–

577.

https://doi.org/10.1038/s41586-018-0613-1

R Core Team (2021) R: A Language and Environment for Statistical Computing. R

Foundation for Statistical Computing, Vienna. Available from: https://www.R-

project.org (accessed 31 October 2021).

Solis, D. R., Gonçalves Paterson Fox, E., Mayumi Kato, L., Massuretti de jesus, C.,

Teruyoshi Yabuki, A., Eugênia de Carvalho Campos, A. & Correa Bueno, O. (2010a)

Morphological description of the immatures of the ant, *Monomorium floricola*. *Journal of Insect Science*, 10, 1–17.

https://doi.org/10.1673/031.010.1501

Solis, D.R., Fox, E.G.P., Rossi, M.L. & Bueno, O.C. (2010b) Description of the immatures of

Linepithema humile Mayr (Hymenoptera: Formicidae). *Biological Research*, 43, 19–30.

https://doi.org/10.4067/S0716-97602010000100004

Schneider, C.A., Rasband, W.S. & Eliceiri, K.W. (2012) NIH Image to ImageJ: 25 years of

image analysis. *Nature Methods*, 9, 671–675.

https://doi.org/10.1038/nmeth.2089

Schultner, E., Oettler, J. & Helanterä, H. (2017) The role of brood in eusocial Hymenoptera.

The Quarterly Review of Biology, 92, 39–78.

https://doi.org/10.1086/690840

Shattuck, S.O. (1992) Generic revision of the ant subfamily Dolichoderinae. *Sociobiology*, 21,

1–181.

Sparks, K.S., Andersen, A.N. & Austin, A.D. (2019) A multi-gene phylogeny of Australian

Monomorium Mayr (Hymenoptera: Formicidae) results in reinterpretation of the genus

and resurrection of *Chelaner* Emery. *Invertebrate Systematics*, 33, 225–236.

https://doi.org/10.1071/IS16080

Taylor, R. W. (1965) A monographic revision of the rare tropicopolitan ant genus

Probolomyrmex Mayr (Hymenoptera: Formicidae). *Transactions of the Royal

Entomological Society of London*, 117, 345–365.

https://doi.org/10.1111/j.1365-2311.1965.tb00044.x

Taylor, R. W. (1967) A monographic revision of the ant genus *Ponera* Latreille

(Hymenoptera: Formicidae). *Pacific Insects Monograph*, 13, 1–112.

Villet, M.H., Hanrahan, S.A. & Walther, C. (1990) Larval structures associated with larva-to-

adult trophallaxis in *Platythyrea* (Hymenoptera : Formicidae). *International Journal of*

Insect Morphology and Embryology, 19, 243–256.

https://doi.org/10.1016/0020-7322(90)90010-M

Ward, P.S., Brady, S.G., Fisher, B.L. & Schultz, T.R. (2015) The evolution of myrmicine

ants: Phylogeny and biogeography of a hyperdiverse ant clade (Hymenoptera:

Formicidae). *Systematic Entomology*, 40, 61–81.

https://doi.org/10.1111/syen.12090

Wetterer, J.K. (2010a) Worldwide spread of the pharaoh ant, *Monomorium floricola*

(Hymenoptera: Formicidae). *Myrmecological News*, 13, 19–27.

Wetterer, J.K. (2010b) Worldwide spread of the pharaoh ant, *Monomorium pharaonis*

(Hymenoptera: Formicidae). *Myrmecological News*, 13, 115–129.

Wheeler, G.C. & Wheeler, J. (1955) The Ant Larvae of the Myrmicine Tribe Solenopsidini.

American Midland Naturalist, 54, 119–141.

Wheeler, G.C. & Wheeler, J. (1966) The Ant Larvae of the Subfamily Dolichoderinae:

Supplement. *Annals of the Entomological Society of America*, 59, 726–732.

Wheeler, G.C. & Wheeler, J. (1971) Ant larvae of the subfamily Ponerinae: second

supplement. *Annals of the Entomological Society of America*, 64(6), 1197–1217.

Wheeler, G.C. & Wheeler, J. (1973) The ant larvae of six tribes: second supplement

(Hymenoptera: Formicidae: Myrmicinae). *Journal of the Georgia Entomological Society*,

8:27–39.

Wheeler, G.C. & Wheeler, J. (1976) Ant larvae: review and synthesis. *Memoirs of the*

Entomological Society of Washington, 7, 1–108.

Wheeler, G.C. & Wheeler, J. (1980) Supplementary Studies on ant Larvae: Ponerinae,

Myrmicinae and Formicinae. *Transactions of the American Entomological Society*, 106,

527–545.

Wheeler, W.M. (1918) A study of some ant larvae, with a consideration of the origin and

meaning of the social habit among insects. *Proceedings of the American Philosophical*

Society, 57, 293–343.

FIGURE 1. Relationship between the maximum head capsule width (mm) and the maximum body length (mm) of *Monomorium triviale* larvae. The 1st instar (orange circle, $n = 84$), 2nd instar (blue triangle, $n = 226$), 3rd instar worker (green square, $n = 164$) and last instar queen (brown cross, $n = 116$) larvae were determined by morphology and chaetotaxy. Histogram bin width of head capsule width and body length is 0.005 mm and 0.1 mm respectively.

FIGURE 2. Egg and first instar larva of *Monomorium triviale*. **A.** side view of egg; **B.** habitus of first instar larva in lateral view; **C.** habitus of first instar larva in ventral view; **D.** lateral body surface of first instar larva, showing the spiracles (white arrowheads) and unbranched hairs (black arrowheads) on the thoracic (T2–T3) and abdominal (A1) segments; **E.** mouthparts of first instar larvae: labrum (lr), maxilla (mx), mandible (M), labium (lb) and labial palp (pl). Scale bars: 33.3 μm (A, B, C); 14.2 μm (D, E).

FIGURE 3. Second instar larva of *Monomorium triviale*. **A.** habitus in lateral view; **B.** habitus in ventral view; **C.** lateral body surface, showing the spiracles (white arrowheads) and unbranched hairs (black arrowheads) on the thoracic (T2–T3) and abdominal (A1) segments; **D.** mouthparts: labrum (lr), maxilla (mx), labium (lb), maxillary palp (pm) and labial palp (pl). Scale bars: 50 μm (A, B); 14.2 μm (C, D).

FIGURE 4. Third instar larva of *Monomorium triviale*. **A.** habitus in lateral view; **B.** habitus in ventral view; **C.** head and mouthparts: antenna (an), clypeus (cl), labrum (lr), maxilla (mx), labium (lb), maxillary palp (pm) and labial palp (pl); **D.** anchor-shaped body hairs on dorsal thoracic somite; **E.** lateral body surface, showing the spiracles (black arrowheads) on the thoracic (T2–T3) and abdominal (A1) segments. Scale bars: 100 µm (A, B); 25 µm (C, D, E).

FIGURE 5. Last instar larva of *Monomorium triviale* queen. **A.** habitus in lateral view; **B.** habitus in ventral view; **C.** habitus in dorsal view; **D.** head and mouthparts: antenna (an), clypeus (cl), labrum (lr), maxilla (mx), labium (lb), maxillary palp (pm) and labial palp (pl); **E.** body hairs; **F.** doorknob-like tubercle; **G.** mid-ventral boss. Scale bars: 142 μm (A, B, C); 20 μm (D, E, F, G).

FIGURE 6. Pupae of *Monomorium triviale*. **A.** habitus of worker pupa in lateral view; **B.** habitus of queen pupa in lateral view; **C.** thorax of queen pupa in dorsal view. All scale bars = 200 μm.

FIGURE 7. Habitus of last instar *Monomorium* larvae in ventral view. **A.** *M. hiten* worker; **B.** *M. intrudens* worker; **C.** *M. chinense* worker; **D.** *M. hiten* reproductive; **E.** *M. intrudens* reproductive; **F.** *M. chinense* reproductive. All scale bars = 200 μm.

FIGURE 8. Histological section through the protuberances of last instar queen larvae, showing epidermis (ep) and cuticle (cu). **A.** cross section of dorsal doorknob-like tubercles; **B.** cross section of lateral doorknob-like tubercles; **C.** longitudinal section of mid-ventral bosses; **D.** cross section of mid-ventral bosses. All scale bars = 50 µm.

Table 1. Morphological and chaetotaxical characteristics in different castes and instars of *M. triviale*.

Larval instar	Head width (mm)	Body length (mm)	Hair number	Hair type	Protuberances
1st instar	0.122 ± 0.006 (0.107–0.137)	0.328 ± 0.036 (0.237–0.416)	100–150	simple smooth	absent
2nd instar	0.137 ± 0.006 (0.124–0.152)	0.459 ± 0.074 (0.281–0.593)	500–600	simple smooth	absent
3rd instar worker	0.175 ± 0.008 (0.146–0.191)	0.847 ± 0.192 (0.414–1.20)	400–500	anchor-tipped	absent
Last instar queen	0.187 ± 0.013 (0.16–0.220)	1.756 ± 0.329 (0.742–2.291)	almost hairless	simple smooth	present

TABLE 2. Protuberance position and size in the last instar queen larvae of *Monomorium triviale*.

Somites	Lateral tubercle size (µm)		Dorsal tubercle size (µm)		Ventral boss size (µm)		
	diameter	length	diameter	length	width	length	height
Thoracic 1	-		-		-		
Thoracic 2	61-76	116-141	80-95	184-210	-		
Thoracic 3	77-87	102-120	67-82	160-210	101-217	52-75	43-79
Abdominal 1	60-86	130-161	70-73	135-173	181-277	66-109	116-122
Abdominal 2	66-93	119-168	69-73	113-135	238-349	89-138	123-152
Abdominal 3	68-92	107-127	62-69	82-95	293-430	134-180	137-159
Abdominal 4	73-82	92-111	59-67	46-65	indistinct		
Abdominal 5	66-81	79-90	-		indistinct		
Abdominal 6	55-75	54-70	-		indistinct		
Abdominal 7	55-66	38-58	-		-		
Abdominal 8	-		-		-		
Abdominal 9	-		-		-		
Abdominal 10	-		-		-		

TABLE 3. The developmental periods of the immature stages of *M. triviale* and the number of observed individuals for each stage.

Developmental stage	Source nests	The numbers of individuals			Developmental period (day)	
		Total observed	Turned to target stage	Molted to next stage	mean ± SD	range
Egg	18	NA	211	166	13.39 ± 2.28	8-22
Young larva	7	70	32	32	7.31 ± 1.00	6-9
Old larva	8	80	72	51	18.31 ± 3.29	13-26
Prepupa	8	51	51	51	4.43 ± 1.10	1-7
Pupa	7	70	33	23	16.87 ± 1.55	13-20

TABLE 4. The list of previously reported unusual larval structures. references cited: 1, Masuko 1989; 2, Masuko 2019; 3, Taylor 1965; 4, Wheeler 1918; 5, Villet *et al*. 1990; 6, Petralia & Vinson; 1979; 7, Taylor 1967; 8, Wheeler & Wheeler 1971; 9, Peeters & Hölldobler 1992; 10, Fox *et al.* 2017; 11, Wheeler & Wheeler 1966; 12, Shattuck 1992; 13, Solis *et al.* 2010b; 14, Eidmann 1926; 15, Menozzi 1930; 16, Casevitz-Weulersse 1984.

Subfamily	Segment	Position	Structure	Caste	Instar	Function	Genera	References
Leptanillinae	abdominal	dorsal	duct-like opening	-	last	trophallaxis	*Leptanilla*	1
Proceratiinae	abdominal	dorsal	slit-like openings	-	last	trophallaxis	*Proceratium*	2
		posterodorsal	sucker-like appendage	-	-	attachment	*Probolomyrmex*	3
Pseudomyrmecinae	thoracic, abdominal	ventral	large protuberance	-	all	trophallaxis	*Tetraponera*	4
Ponerinae	abdominal	ventral	single conical tubercle	worker	late	trophallaxis	*Platythyrea*	5
			protuberances	-	-	food holding	*Odontomachus, Pachycondyla, Leptogenys*	6
		dorsal	paired doorknob tubercles	-	varies	support, attachment	*Brachyponera, Cryptopone, Hypoponera, Odontomachus,*	7, 8, 9, 10

							Ponera, Myopias, Simopelta, Belonopelta	
Dolichoderinae	abdominal	dorsal	no-paired tubercles	worker	all	-	*Froggattella*	11, 12
			single tubercle	worker	all	-	*Linepithema*	12, 13
		posterodorsal	single tubercle	worker	-	-	*Tapinoma*	11, 12
	abdominal	lateral	well-developed dilatations	queen?	late	-	*Crematogaster*	14, 15, 16
Myrmecinae	thoracic, abdominal	lateral, dorsal	doorknob-like tubercles	queen	last	-	*Monomorium*	This study
	thoracic, abdominal	ventral	boss-shaped protuberances	queen	last	-	*Monomorium*	This study