Whole-genomes from the extinct Xerces Blue butterfly can help identify declining insect species

Toni de-Dios [1,2+], Claudia Fontsere [1,3+], Pere Renom [1+], Josefin Stiller [3], Laia Llovera[1], Marcela Uliano-Silva[4], Alejandro Sánchez-Gracia[5], Charlotte Wright[4], Esther Lizano [1,6], Berta Caballero[7], Arcadi Navarro [1,8], Sergi Civit [9], Robert K. Robbins[10], Mark Blaxter[4], Tomàs Marquès-Bonet [1,6,8,11*], Roger Vila [1*], Carles Lalueza-Fox [1,7*]

1Institute of Evolutionary Biology (CSIC-Universitat Pompeu Fabra), 08003 Barcelona, Spain

2Institute of Genomics, University of Tartu, Riia 23B, 51010 Tartu, Estonia

3Section for Evolutionary Genomics, The Globe Institute, Faculty of Health and Medical Sciences, University of Copenhagen, Copenhagen K, Denmark

3Centre for Biodiversity Genomics, University of Copenhagen, DK-2100, Denmark

4Wellcome Sanger Institute, Hinxton, Saffron Walden CB10 1RQ, UK

5Departament de Genetics, Microbiology and Statistics-Institut de Recerca de la Biodiversitat (IRBio), Universitat de Barcelona, Barcelona 08028, Spain

6Institut Català de Paleontologia Miquel Crusafont, Universitat Autònoma de Barcelona, 08193 Cerdanyola del Vallès, Barcelona, Spain

7Museu de Ciències Naturals de Barcelona, 08019, Barcelona, Spain

8 Catalan Institution of Research and Advanced Studies (ICREA), 08010 Barcelona, Spain

9Departament de Genetics, Microbiology and Statistics-Institut de Recerca de la Biodiversitat (IRBio), Universitat de Barcelona, Barcelona 08028, Spain

10 Department of Entomology, National Museum of Natural History, Smithsonian Institution, Washington, DC 20013-7012, USA

11 CNAG-CRG, Centre for Genomic Regulation, Barcelona Institute of Science and Technology (BIST), 08036 Barcelona, Spain

+ equally contributed

*Corresponding Authors

Carles Lalueza-Fox
Tomas Marquès-Bonet
Roger Vila

Emails: (T.d.D) tonidedios94@gmail.com; (C.F.) claudia.fontsere@sund.ku.dk; (P.R.) cogitoergomitto@gmail.com; (L.Ll.) laia.llovera@crg.eu ; (J.S.) josefinstiller@gmail.com; (M.U.-S.) mu2@sanger.ca.uk; (A.S.G.) elsanchez@ub.edu; (C.W.) cw22@sanger.ac.uk; (E.L.) esther.lizano@upf.edu; (B.C.) bcaballerolo@bcn.cat; (A.N). arcadi.navarro@upf.edu (S.C.) svives@ub.edu; (R.K.R.) ROBBINSR@si.edu; (M.B.) mb35@sanger.ac.uk; (T.M.-B) tomas.marques@upf.edu; (R.V.) roger.vila@csic.es; (C.L.-F) carles.lalueza.fox@gmail.com;

Running Head: Genomes of the extinct Xerces Blue butterfly

41   **Keywords:** Extinction; Museomics; Butterflies; Population Genomics; Xerces Blue;

42   Genomic Erosion

44   Abstract

The Xerces Blue (*Glaucopsyche xerces*) is considered to be the first butterfly to

become extinct at global scale in historical times. It was notable for its chalky lavender

wings with conspicuous white spots on the ventral wings. The last individuals were

collected in their restricted habitat, in the dunes near the Presidio military base in San

Francisco, in 1941. We sequenced the genomes of four 80 to 100-year-old Xerces

Blue, and seven historical and one modern specimens of its closest relative, the Silvery

Blue (*G. lygdamus*). We compared these to a novel annotated genome of the Green-

Underside Blue (*G. alexis*). Phylogenetic relationships inferred from complete

mitochondrial genomes indicate that Xerces Blue was a distinct species that diverged

from the Silvery Blue lineage at least 850,000 years ago. Using nuclear genomes, both

species experienced population growth during the Eemian interglacial period, but the

Xerces Blue decreased to a very low effective population size subsequently, a trend

opposite to that observed in the Silvery Blue. Runs of homozygosity and deleterious

load in the Xerces Blue were significantly greater than in the Silvery Blue, suggesting a

higher incidence of inbreeding. These signals of population decline observed in Xerces

Blue could be used to identify and monitor other insects threatened by human

activities, whose extinction patterns are still not well known.

63   Introduction

The Xerces Blue butterfly (*Glaucopsyche xerces)*[1] was native to the coastal sand

dunes of San Francisco in association with the common Deerwood (*Acmispon glaber),*

which was the preferred food source for larval stage[2]. It was notable for its iridescent

blue colouration on the dorsal (upper) wing surface, and conspicuous, variable white

spots on the ventral surface[3]. With the growth of San Francisco and the destruction of

sand dune habitats, the Xerces Blue became restricted to a few sites in what is now

Golden Gate National Recreation Area. The last specimens were reportedly collected

by entomologist W. Harry Lange on March 23, 1941[3]. It is considered the first butterfly

to have been driven to global extinction by human activities[3].

The Xerces Blue and the closely related Silvery Blue (*Glaucopsyche lygdamus*) were

recently proposed to be distinct species based on mtDNA data from a single Xerces

Blue specimen[4]. However, two nuclear genes analysed (ribosomal 28S and histone

H3) were invariable and genome-wide data were unavailable for the Xerces Blue,

hampered by the inherent difficulties of retrieving genome-wide data from historical

insect specimens[5,6] and the absence of a suitable reference genome. The genus

Glaucopsyche consists of 18 extant species distributed across the temperate regions

of the northern hemisphere. To provide a relevant reference, we generated an

annotated genome from the Palearctic Green-Underside Blue butterfly *Glaucopsyche*

alexis[7]. Using DNA extracted from five Xerces Blue and seven Silvery Blue

(*Glaucopsyche lygdamus*) historical specimens from the vicinity of San Francisco, and

also from a modern Silvery Blue male from Canada, we generated whole genome

resequencing data for both species and investigated their relationships and historical

population genetics.

Results

Historic and modern butterfly genomes

We extracted DNA from 12 historical specimens (5 *G. xerces*, 7 *G. lygdamus*) (Table

S1). One Xerces Blue sample did not yield detectable DNA in two independent

extractions. For each of the successful extracts we prepared a single library which was

shotgun sequenced on the HiseqX Illumina platform. We mapped 124,101,622 and

184,084,237 unique DNA reads of Xerces Blue and Silvery Blue, respectively, against

the *G. alexis* reference genome (Table 1 and Table S2). The DNA reads exhibited

typical ancient DNA features, such as short mean read length (ranging from 47.55 to

67.41 bases on average, depending on the specimen (Fig. S1)) and post-mortem

deamination patterns at the 5' and 3' ends (Fig. S2 and S3). As listed in the original

museum records, we found one Silvery Blue and two Xerces Blue females (Table S2).

Inter-individual comparisons suggested no close kinship link among the studied

individuals.

The historical genomes covered 49.3% (Xerces Blue) and 55.2% (Silvery Blue) of the

G. alexis reference genome, largely because repetitive chromosomal regions cannot

be confidently assessed with short, ancient DNA sequence reads (Fig. S4 – S7). To

estimate the mappable fraction of the reference *G. alexis* genome, we randomly

fragmented it to 50 to 70 nucleotides and mapped the generated fragments back to the

complete genome. An average of 57.8% of the *G. alexis* genome was covered with

these read lengths (Table S2). We suggest that reduced coverage from the historical

specimens may be due to genomic divergence of *G. xerces* and *G. lygdamus* from the

G. alexis reference (Fig. S8). The annotation of genes located in those unrecoverable

regions provided a putative list of 14 nuclear genes with diverse functions obtained

from BLAST, that should be further explored to understand the uniqueness of the

extinct species (Table S4).

Phylogenetic relationships

Maximum likelihood phylogenetic inference using whole mitochondrial genomes

showed that the Xerces Blue specimens form a monophyletic clade, as do the Silvery

Blue specimens (Fig. 1A). We inferred a time-calibrated Bayesian phylogenetic tree

from protein-coding genes Analysis and 12 related butterflies in Polyommatinae

subfamily, revealing high support for the sister group relationship (posterior

probability=1). We found the specie *Shijimiaeoides (Sinia) divina* inside the

Glaucopsyche clade, in agreement with previous phylogenetic studies, and we

subsequently labelled it as *G. divina*[8]. Because there are no known fossils to calibrate

the time since divergence, we first used a molecular clock that spanned the range of

rates frequently used for arthropod mitochondrial genes (1.5-2.3% divergence/Ma).

Our dated analysis yielded an origin of this subgroup of Polyommatinae at 12.4 Ma

(8.82-16.27 Ma 95% HPD interval) and divergence of the Xerces Blue from the Silvery

Blue at 900,000 years ago (0.61-1.19 Ma 95% HPD interval, Fig. 1B). A second

estimate based on larger-scale fossil-based calibrations[9] fixed the origin of the

subgroup to ca. 33 Ma[10], inferred the subsequent divergence of the Xerces Blue and

Silvery Blue to 2.40 Ma (1.95-2.73 Ma 95% HPD interval, Fig 1B). The recent

speciation of Xerces and Silvery blue is not obviously due to infection with the

Wolbachia, as no evidence of infection of the sampled specimens with this alpha-

proteobacterium is detected in the raw read data.

Principal Component Analysis (PCA) using PCAngsd[11] and nuclear genome

polymorphisms for the three Glaucopsyche species supports the relationships among

them; the historical specimens are equally distant to G. alexis in the first PC, explaining

52.81% of the variance (Fig. S9). The second PC separates the Xerces Blue from the

Silvery Blue specimens.

Demographic history and diversity

We used the Pairwise Sequentially Markovian coalescent (PSMC) algorithm[12] to

evaluate the demographic histories of both butterfly species, first exploring the two

specimens with highest coverage (L05 and L13) (Fig. 2). We found an increase in

effective population size in both species that is roughly coincident with the interglacial

Marine Isotopic Stage 7 (approximately from 240,000 to 190,000 years ago[13]). After

this timepoint the trends differ. We estimated a continuous decrease in Xerces Blue

population size in parallel to the Wisconsin Glacial Episode, which started about

75,000 years ago. However, both the modern and the historical Silvery Blue do not

appear to have been negatively affected by this event (Fig. S10), suggesting different

adaptive strategies to cope with cooling temperatures and/or food plant availability.

Second, we generated PSMC curves from the remaining lower-coverage individuals

and down-sampled data from specimen L05 to 50% and 75% of the total coverage to

explore the effects of coverage on estimation of heterozygous sites. Although there

was a reduction in the effective population size estimates, as expected, the temporal

trajectories in lower-coverage individuals were similar to their respective, higher-

coverage Xerces Blue and Silvery Blue references (Fig. S10).

We subsequently explored the heterozygosity of each individual and found that Xerces

Blue had 22% less heterozygosity on average than the Silvery Blue historical samples,

a difference that is statistically significant (T-test; p=0.0072) (Fig. S8, Table S2). We

searched for runs of homozygosity (RoH) that can indicate the existence of inbreeding

in a dwindling population. The total fraction of the genome presenting RoH, although

limited, is much higher in Xerces Blue (up to 6% of the genome) than in Silvery Blue,

especially in short RoH of size between 100 and 500 kb (Fig. 3 and Fig. S11),

consistent with background inbreeding. The limited presence of long RoH discards

consanguinity as a common scenario in Xerces Blue.

We identified amino acid-changing alleles that may be suggestive of a deleterious

genetic load associated with long-term low population numbers in the Xerces Blue. The

average Ka/Ks ratio is higher in Xerces Blue than in Silvery Blue; the former also

carries a higher fraction of nonsense and functionally high-to-moderate effect variants

in homozygosity and RoHs with an increased concentration of high-to-moderate effect

variants (Fig. 4), as predicted with a functional prediction toolbox, SnpEff[14].

Discussion

We have used a modern reference genome and ancient DNA genome sequence data

from museum specimens to explore the relationships and historical population genetic

history of an extinct butterfly, the Xerces Blue; to our knowledge, this is the first ancient

genome ever generated from an extinct insect. Based upon a near-complete mtDNA

genome from a Xerces Blue specimen, Grewe et al. (2021)[4] proposed that the Xerces

Blue and the Silvery Blue were distinct species. We confirm this finding using full

mitochondrial genomes and extensive nuclear genomic data from multiple specimens.

Given the lack of evidence for *Wolbachia* infection, the recent speciation of Xerces

Blue and Silvery Blue seems unrelated to cytoplasmic incompatibility cause by this

endosymbiont[15,16]; a detailed analysis of genomic architectures could help identify

barriers to introgression between these species.

Our analyses indicate that the Xerces Blue had experienced a severe demographic

decline for tens of thousands of years, likely associated with changing climatic factors.

Thus, the destruction of the Xerces Blue habitat by humans was likely the final blow in

the extinction process. We provide evidence for low population size in Xerces Blue,

correlated with low genetic variation, a higher proportion of runs of homozygosity and

increased frequency of deleterious, amino acid-changing alleles [17–19]. However, there

was no genetic evidence of recent inbreeding.

Inbreeding genetic signals in the form of long chromosomal sections with no variation

sometimes occur in critically endangered species [20,21] and in extinct species such as

the last Mammoths from Wrangel Island[22] or the Altai Neanderthal[23]. The PSMC shows

a continuous low effective population size for Xerces Blue; demographic declines are

also seen in some extinct species, including Wrangel Mammoths[19] but not in others

such as the Woolly Rhino that showed a pre-extinction demographic stability and

relatively low inbreeding signals[24]. In many endangered species there is little

concordance between genome diversity, population sizes and conservation status[21];

this decoupling was also observed in the genomes of the extinct passenger pigeon that

despite being one of the world's most numerous vertebrates, showed a surprisingly low

genetic diversity[25]. Despite being notoriously abundant, insects, and in particular

butterflies, are very sensitive to climate fluctuations; therefore, we suggest that insects

with observations of demographic traits indicative of long-term low effective population

size such as those found in Xerces Blue should be considered to be especially

vulnerable to extinction events.

Our study further demonstrates the value of museum insect specimens for estimating

temporal changes in genetic diversity at a population scale[26]. Despite being notoriously

abundant, insects, and in particular butterflies, are very sensitive to climate

fluctuations. We suggest that insects with genetic observations of long-term low

effective population size such as those found in Xerces Blue should be considered to

be especially vulnerable to extinction events. However, being the insect numbers

usually very high, it is likely that their genomic signals of extinction could be different to

those described in vertebrates in many cases. Therefore, this is a subject that should

be further explored with genomic data from other declining insects.

Methods

Historical butterfly specimens

The Xerces Blue specimens analysed belong to the Barnes collection deposited at the

Smithsonian National Museum of Natural History. Two of them were collected on April

26th, 1923. The Silvery Blue specimens were mostly collected between 1927 and

1948, in Haywood City, Santa Cruz, Oakland, San José, Fairfax and Marin County

(these locations surround San Francisco Bay) (Table S1).

DNA extraction and sequencing of Xerces Blue and Silvery Blue specimens

All DNA extraction and initial library preparation steps (prior to amplification) were

performed in a dedicated clean lab, physically-isolated from the laboratory used for

post-PCR analyses. Strict protocols were followed to minimize the amount of human

DNA in the ancient DNA laboratory, including the wearing a full body suit, sleeves,

shoe covers, clean shoes, facemask, hair net and double gloving, as well as frequent

bleach cleaning of benches and instruments. DNA extraction was performed from 12

abdominal samples of historical Xerces Blue and Silvery Blue, as well as a modern

Silvery Blue specimen from Canada. Experimental procedures are described in detail

in the Supplementary Material.

***Glaucopsyche alexis* genome sequencing and annotation**

Glaucopsyche alexis was chosen as a congeneric reference to compare the

demographic histories of both the Xerces Blue and the Silvery Blue. We generated a

G. alexis reference genome from a male specimen collected in Alcalá de la Selva in

Teruel (Spain). Its genome has a sequence length of 619,543,730 bp on 24

chromosomes – including the Z sex chromosome – and the mitochondrial genome.

The genome sequence is biologically complete (BUSCO Lepidoptera completeness

97.1%)[27]. The *G. alexis* genome was sequenced at the Sanger Institute as part of the

Darwin Tree of Life Project following the extraction, sequencing and assembly

protocols developed for Lepidoptera[7].

Xerces Blue and Silvery Blue mapping and variant calling

The ancient DNA reads were clipped using AdapterRemoval2[28], and only reads longer

than 25bp were kept. Filtered reads were mapped against the *G. alexis* assembly with

Burrows-Wheeler Aligner (BWA)[29], with parameters optimised for the analysis of aDNA

(Supplementary Materials). Basic mapping statistics were generated using

Qualimap2[30] (Table S2). We used bedtools[31] to assess genome coverage across the

reference using windows of 1mbp for the nuclear fraction of the genome (Fig. S4 – S6),

as well as depth of coverage (Fig. S7), read length (Fig. S1) and edit distance

distribution (Fig. S8). Authenticity of the sequences was assessed by characterising

aDNA damage patterns with pmdtools[32] and MapDamage2[33] (Fig S2 and S3).

We used snpAD[34], a program for genotype calling in ancient specimens. The mapped

sequences were transformed from bam-format into snpAD-format files, priors for base

composition estimated, and genotypes were called using standard settings. The VCFs

were combined and concatenated with CombineVariants and GatherVcfs from GATK[35]

and filtered with vcftools[36] to keep only sites within the mappable fraction of the

genome previously obtained with minimum read depth of 2, max read depth of 30,

genotype quality > 30, maximum missingness of 0.6, minor allele frequency of 5% and

excluding indels and multiallelic sites.

Genotype likelihoods were obtained with ANGSD[37] using the GATK model with the

following parameters for all the samples: -uniqueOnly 1 -remove_bads 1 -

only_proper_pairs 1 -trim 10 -C 50 -baq 1 -minInd 5 -skipTriallelic 1 -GL 2 -minMapQ

30 -doGlf 2 -doMajorMinor 1 -doMaf 2 -minMaf 0.05 -SNP_pval 1e-6.

Sex determination

The sex of the specimens was determined by differential coverage of the Lepidopteran

Z chromosome (females are the heterogametic sex in the Lepidoptera and show

reduced coverage on the Z chromosome) (Table S2).

Mitochondrial phylogenetic tree and divergence dating

Haploid variants were called using bcftools[38] with a ploidy of 1, filtering low quality

indels and variants, after which a consensus sequence was exported. We downloaded

14 complete mitochondrial genomes for Polyommatinae from NCBI (Table S3).

All mitochondrial genomes were annotated with MitoFinder[39] using *Shijimiaeoides*

divina as the reference. The 11 protein-coding genes were aligned with the codon-

aware aligner MACSE[40] and the ribosomal rRNAs were aligned with MAFFT I-ins-i[41].

We first investigated phylogenetic relationships among five *G. xerces* and eight *G.*

lygdamus individuals, with *G. alexis* as the outgroup. We used IQ-TREE2[42] to select

the best fitting nucleotide substitution model for each partition and merge similar

partitions[43], built a maximum likelihood tree and assessed support with 1000 ultrafast

bootstrap replicates[44].

To infer a time-calibrated phylogenetic hypothesis, we selected one individual of

Xerces Blue (L003) and Silvery Blue (RVcoll10-B005) and analysed with 13 other

Polyommatinae species. We used BEAST2[45] with the bModelTest[46] package to

perform phylogenetic site model averaging for each of the merged partitions. Because

there is no accepted molecular clock rate for butterflies and no fossils to apply in this

part of the phylogeny, we used two strategies to apply time constraints to the analysis.

First, we used two published molecular clock rates for the mitochondrial COX1

gene (1.5% divergence/Ma estimated for various invertebrates[47], and the 'standard'

insect mitochondrial clock 2.3% divergence/Ma[48]). We applied a strict clock with a

normal prior set up to span 1.5-2.3% with the 95% HPD interval (mean=1.9%,

sigma=0.00119). Second, we borrowed the age of the most recent common ancestor

of our sampled taxa from fossil-calibrated analyses across butterflies[10,49]. We fixed the

root age to 33 Ma and allowed the remaining node ages to be estimated using a strict

clock. Analyses were run twice from different starting seeds for 10 million MCMC

generations and trees were sampled every 1000 generations. Runs were checked for

convergence with Tracer and all effective sample size (ESS) values were >200. Runs

were combined with the BEAST2 package LogCombiner[50], after removing the first 10%

of topologies as burn-in, and a maximum credibility tree was generated with

TreeAnnotator[50]. Phylogenetic analyses were performed on the National Life Science

Supercomputing Center - Computerome 2.0 (www.computerome.dk).

322   Xerces Blue and Silvery Blue population histories

We used the Pairwise Sequentially Markovian Coalescent (PSMC) model[12] to explore

the demographic history of both butterfly species. We obtained a consensus fastq

sequence of the mappable fraction of the genome for each autosomal chromosome

(total of 22 chromosomes of *G. alexis* assembly). Only positions with a depth of

coverage above 4X and below 15X were kept. Posteriorly a PSMC was built using the

following parameters: -N25 -t15 -r5 -p "28*2+3+5". We used 1 year for the generation

time and a mutation rate of 1.9×10^{-9}, estimated in *Heliconius melpomene*[51]. Considering

that calling consensus sequences from low coverage samples (<10x) can

underestimate heterozygous sites[52], and given the different coverage between

samples, we corrected by False Negative Rate the samples with coverage lower than

the coverage of L005 (for Xerces Blue) and L013 (for Silvery Blue), as recommended

by the developers of the software, so that all samples are comparable with each other.

However, since in our dataset we do not reach a coverage >20x, we acknowledge that

we are not capturing the whole diversity and thus our PSMC might infer lower historical

effective population sizes.

339   Population stratification and average genome heterozygosity

Principal Component Analysis (PCA) was performed using PCAngsd[11] after obtaining

genotype likelihoods with ANGSD including all individuals. To assess global levels of

heterozygosity, the unfolded SFS was calculated for each sample separately using

ANGSD[37] and realSFS with the following quality filter parameters: -uniqueOnly 1 -

remove_bads 1 -only_proper_pairs 1 -trim 10 -C 50 -baq 1 -minMapQ 30 -minQ 30 -

setMaxDepth 200 - doCounts 1 -GL 2 -doSaf 1.

Runs of Homozygosity (RoH)

RoH were called based on the density of heterozygous sites in the genome using the

implemented Hidden Markov Model (HMM) in bcftools[38] roh with the following

parameters: -G30 --skip-indels --AF-dflt 0.4 --rec-rate 1e^{-9} from the mappable fraction of

the genome with the filtered VCF file. We kept the RoH with a phred score > 85. We

divided the RoH into different size bins: very short RoH (<100 kb), short RoH (100-500

kb), intermediate RoH (500 kb-1Mb) and long (1-5 Mb or > 5Mb). Short RoH reflect LD

patterns, intermediate size RoH describe background inbreeding due to genetic drift

and long RoH appear in the case of recent inbreeding[53].

Deleterious load

358 We used the *G. alexis* annotations to create a SNPeff database that we used to

359 annotate our callings. Using SNPeff[14] again and the set of variants discovered by

360 angsd, we predicted the putative effect of those variant in the analysed individuals

361 (Table S2). In addition to wide genome mutations, we specifically focused on mutations

362 present in homozygosis, heterozygosis and the previously annotated RoH.

Unrecoverable regions

To further explore how the genomic divergence can influence our genome

reconstruction success, we undertook a similar approach as the genome of the

Christmas Island rat[54], and explored the chromosomal regions in the *G. alexis*

reference that were significantly depleted of Xerces DNA reads. We used bedtools[31]

and some in-home bash scripting to calculate the mean coverage per gene of the *G.*

alexis genome for Xerces Blue sequencing DNA reads. We first used bedtools'

algorithms *bamtobed* and *genomecov* to estimate the genome-wide per-site coverage

of the reference genome in these two species. Then, we extracted the coordinates of

all protein coding genes from the annotation file (gff file) and used the intersect to

estimate the average coverage of each protein coding gene. We performed a

functional analysis of all genes uncovered in *G. alexis*, excluding those that are present

in *G. lygdamus* with more than 5x coverage (as we were looking for evolutionary

novelties in the Xerces Blue lineage alone) using profile- IntersProScan[55] and

sequence similarity-based (blasp) searches[56].

379 *Wolbachia* screening

Wolbachia are endosymbiotic alpha-proteobacteria that are present in about 70% of

butterfly species and induce diverse reproductive alterations, including genetic barriers

when two different strains infect the same population or when two populations – one

infected and one uninfected – meet[15]. As potential evidence for a reproductive barrier

promoting the separation of Xerces Blue and Silvery Blue, we searched for *Wolbachia*

DNA reads in our specimens, taking advantage of the high coverage and the shotgun

approach. First, we collapsed unique reads from the butterfly-free sequences with

BBmap[57] and removed from the dataset low complexity sequences using Prinseq[58].

Afterwards, we used Kraken2[59] to assign reads against the standard plus human

Kraken2 database (bacteria, archaea, fungi, protozoa and viral). The historical

specimens did not display enough reads assigned to Wolbachia for us to suspect of the

presence of the bacteria in those samples (Table S6).

393 Acknowledgments

C.L.-F. is supported by a PID2021-124590NB-100 grant (MCIU/AEI/FEDER, UE) of

Spain; T.M.-B. is supported by funding from the European Research Council (ERC)

(grant agreement No. 864203), BFU2017-86471-P (MINECO/FEDER, UE), "Unidad de

Excelencia María de Maeztu", funded by the AEI (CEX2018-000792-M), Howard

Hughes International Early Career, and Generalitat de Catalunya, GRC 2017-SGR-

880; R.V. is supported by grant PID2019-107078GB-I00, funded by

MCIN/AEI/ 10.13039/501100011033, and by GRC 2017-SGR-991 (Generalitat de

Catalunya). We are grateful to the SCIENCE Faculty at University of Copenhagen for

free access to Computerome 2.0. This research was funded in whole, or in part, by the

Wellcome Trust Grants 206194 and 218328 (MU, MB). For the purpose of Open

Access,

the author has applied a CC BY public copyright license to any Author Accepted

Manuscript version arising from this submission.

Competing Interest Statement: The authors declare no competing interest.

Author Contributions: C.L.-F., R.V., R.R. and P.R. conceived the project. R.R.

studied and sampled the specimens. L.L. and E.L. performed experimental work. T.d-

D., C.F., J.S., A.S.G., M.U.-S., C.W., S.C. and P.R. undertook different computational

analyses. C.L.-F., T.M.-B., A.N. and M.B. coordinated different computational teams.

J.S. worked in visualization. C.L.-F., J.S. and R.R. wrote the manuscript with input from

all coauthors.

Data Accessibility

The genetic data generated is publicly available; the accession numbers for the Xerces

Blue and Silvery Blue genomes reported in this study are in the European Nucleotide

Archive (ENA): PRJEB47122. Data on *G. alexis* are available in INSDC under

BioProject PRJEB43798 and genome assembly accessions GCA_905404095.1

(primary haplotype) and GCA_905404225.1 (secondary, alternate haplotype).

References

1.   Boisduval, J. A. Lépidoptères de la Californie. *Ann. Soc. Ent. Fr.* **21**, 275–324

(1852).

2.   Tilden, J. W. San Francisco's vanishing butterflies. *Lepid. News* **10**, 133–1445

(1956).

3.   Downey, J. C. Analysis of Variation in a recently extinct polymorphic lycaenid

butterfly, Glaucopsyche Xerces. *Bull. So. Calif. Acad. Sci.* **55**, 153–170 (1956).

4.   Grewe, F., Kronforst, M. R., Pierce, N. E. & Moreau, C. S. Museum genomics

reveals the Xerces blue butterfly (Glaucopsyche xerces) was a distinct species

driven to extinction . *Biol. Lett.* **17**, 20210123 (2021).

5.   Thomsen, P. F. *et al.* Non-destructive sampling of ancient insect DNA. *PLoS*

One **4**, e5048 (2009).

6.   Staats, M. *et al.* Genomic Treasure Troves: Complete Genome Sequencing of

Herbarium and Insect Museum Specimens. *PLoS One* **8**, e69189–e69189

(2013).

7.   Hinojosa Galisteo, J. C. & Vila, R. The genome sequence of the green-

underside blue, *Glaucopsyche alexis* (Poda, 1761). *Wellcome Open Res.* **6**, 274

(2021).

8.   Lukhtanov, V. A. & Gagarina, A. V. Molecular Phylogeny and Taxonomy of the

Butterfly Subtribe Scolitantidina with Special Focus on the Genera

Pseudophilotes, Glaucopsyche and Iolana (Lepidoptera, Lycaenidae). *Insects* **13**

(12), 1110 (2022).

9.   Espeland, M. *et al.* A Comprehensive and Dated Phylogenomic Analysis of

Butterflies. *Curr. Biol.* **28**, 770-778.e5 (2018).

10.   Chazot, N. *et al.* Priors and Posteriors in Bayesian Timing of Divergence

Analyses: The Age of Butterflies Revisited. *Syst. Biol.* **68**, 797–813 (2019).

11.   Meisner, J. & Albrechtsen, A. Inferring population structure and admixture

proportions in low-depth NGS data. *Genetics* **210**, 719–731 (2018).

12.   Li, H. & Durbin, R. Inference of human population history from individual whole-

genome sequences. *Nature* **475**, 493–496 (2011).

13.   Batchelor, C. L. *et al.* The configuration of Northern Hemisphere ice sheets

through the Quaternary. *Nat. Commun.* **10**, 3713 (2019).

14.   Cingolani, P. *et al.* A program for annotating and predicting the effects of single

nucleotide polymorphisms, SnpEff: SNPs in the genome of Drosophila

melanogaster strain w1118; iso-2; iso-3. *Fly* **6**, 80–92 (2012).

15.   Telschow, A., Hammerstein, P. & Werren, J. H. The effect of Wolbachia versus

genetic incompatibilities on reinforcement and speciation. *Evolution* **59**, 1607–

1619 (2005).

16. Sucháčková Bartoňová, A. *et al.* Wolbachia affects mitochondrial population

structure in two systems of closely related Palaearctic blue butterflies. *Sci. Rep.*

11, 3019 (2021).

17. Szpiech, Z. A. *et al.* Long Runs of Homozygosity Are Enriched for Deleterious

Variation. *Am. J. Hum. Genet.* **93**, 90–102 (2013).

18. Spielman, D., Brook, B. W. & Frankham, R. Most species are not driven to

extinction before genetic factors impact them. *Proc. Natl. Acad. Sci. U. S. A.*

101, 15261–15264 (2004).

19. Palkopoulou, E. *et al.* Complete Genomes Reveal Signatures of Demographic

and Genetic Declines in the Woolly Mammoth. *Curr. Biol.* **25**, 1395–1400 (2015).

20. van der Valk, T., Díez-Del-Molino, D., Marques-Bonet, T., Guschanski, K. &

Dalén, L. Historical Genomes Reveal the Genomic Consequences of Recent

Population Decline in Eastern Gorillas. *Curr. Biol.* **29**, 165-170.e6 (2019).

21. Díez-del-Molino, D., Sánchez-Barreiro, F., Barnes, I., Gilbert, M. T. P. & Dalén,

L. Quantifying Temporal Genomic Erosion in Endangered Species. *Trends Ecol.*

Evol. **33**, 176–185 (2018).

22. Rogers, R. L. & Slatkin, M. Excess of genomic defects in a woolly mammoth on

Wrangel island. *PLoS Genet.* **13**, e1006601 (2017).

23. Prüfer, K. *et al.* The complete genome sequence of a Neanderthal from the Altai

Mountains. *Nature* **505**, 43–49 (2014).

24. Lord, E. *et al.* Pre-extinction Demographic Stability and Genomic Signatures of

Adaptation in the Woolly Rhinoceros. *Curr. Biol.* **30**, 3871-3879.e7 (2020).

25. Murray, G. G. R. *et al.* Natural selection shaped the rise and fall of passenger

pigeon genomic diversity. *Science.* **358**, 951–954 (2017).

26. Lalueza-Fox, C. Museomics. *Curr. Biol.* **32**, R1214–R1215 (2022).

27. Manni, M., Berkeley, M. R., Seppey, M. & Zdobnov, E. M. BUSCO: Assessing

Genomic Data Quality and Beyond. *Curr. Protoc.* **1**, e323 (2021).

28.   Schubert, M., Lindgreen, S. & Orlando, L. AdapterRemoval v2: rapid adapter

trimming, identification, and read merging. *BMC Res. Notes* **9**, 88 (2016).

29.   Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows-

Wheeler transform. *Bioinformatics* **25**, 1754–1760 (2009).

30.   Okonechnikov, K., Conesa, A. & García-Alcalde, F. Qualimap 2: Advanced

multi-sample quality control for high-throughput sequencing data. *Bioinformatics*

32, 292–294 (2016).

31.   Quinlan, A. R. & Hall, I. M. BEDTools: A flexible suite of utilities for comparing

genomic features. *Bioinformatics* **26**, 841–842 (2010).

32.   Skoglund, P. *et al.* Separating endogenous ancient DNA from modern day

contamination in a Siberian Neandertal. *Proc. Natl. Acad. Sci. U. S. A.* **111**,

2229–2234 (2014).

33.   Jónsson, H., Ginolhac, A., Schubert, M., Johnson, P. L. F. & Orlando, L.

MapDamage2.0: Fast approximate Bayesian estimates of ancient DNA damage

parameters. *Bioinformatics* **29**, 1682–1684 (2013).

34.   Prüfer, K. SNPAD: An ancient DNA genotype caller. *Bioinformatics* **34**, 4165–

4171 (2018).

35.   McKenna, A. *et al.* The genome analysis toolkit: A MapReduce framework for

analyzing next-generation DNA sequencing data. *Genome Res.* **20**, 1297–1303

(2010).

36.   Danecek, P. *et al.* The variant call format and VCFtools. *Bioinformatics* **27**,

2156–2158 (2011).

37.   Korneliussen, T. S., Albrechtsen, A. & Nielsen, R. ANGSD: Analysis of Next

Generation Sequencing Data. *BMC Bioinformatics* **15**, 356 (2014).

38.   Danecek, P. *et al.* Twelve years of SAMtools and BCFtools. *Gigascience* **10**,

giab008 (2021).

39.   Allio, R. *et al.* MitoFinder: Efficient automated large-scale extraction of

mitogenomic data in target enrichment phylogenomics. *Mol. Ecol. Resour.* **20**,

892–905 (2020).

40. Ranwez, V., Douzery, E. J. P., Cambon, C., Chantret, N. & Delsuc, F. MACSE

v2: Toolkit for the alignment of coding sequences accounting for frameshifts and

stop codons. *Mol. Biol. Evol.* **35**, 2582–2584 (2018).

41. Katoh, K. & Standley, D. M. MAFFT multiple sequence alignment software

version 7: Improvements in performance and usability. *Mol. Biol. Evol.* **30**, 772–

780 (2013).

42. Minh, B. Q. *et al.* IQ-TREE 2: New Models and Efficient Methods for

Phylogenetic Inference in the Genomic Era. *Mol. Biol. Evol.* **37**, 1530–1534

(2020).

43. Kalyaanamoorthy, S., Minh, B. Q., Wong, T. K. F., Von Haeseler, A. & Jermiin,

L. S. ModelFinder: Fast model selection for accurate phylogenetic estimates.

Nat. Methods **14**, 587–589 (2017).

44. Hoang, D. T., Chernomor, O., Von Haeseler, A., Minh, B. Q. & Vinh, L. S.

UFBoot2: Improving the ultrafast bootstrap approximation. *Mol. Biol. Evol.* **35**,

518–522 (2018).

45. Bouckaert, R. *et al.* BEAST 2.5: An advanced software platform for Bayesian

evolutionary analysis. *PLoS Comput. Biol.* **15**, e1006650 (2019).

46. Bouckaert, R. R. & Drummond, A. J. bModelTest: Bayesian phylogenetic site

model averaging and model comparison. *BMC Evol. Biol.* **17**, 42 (2017).

47. Quek, S. P., Davies, S. J., Itino, T. & Pierce, N. E. Codiversification in an ant-

plant mutualism: Stem texture and the evolution of host use in Crematogaster

(Formicidae: Myrmicinae) inhabitants of Macaranga (Euphorbiaceae). *Evolution*

58, 554–570 (2004).

48. Van Zandt Brower, A. Phylogeny of Heliconius Butterflies Inferred from

Mitochondrial DNA Sequences (Lepidoptera: Nymphalidae). *Mol. Phylogenet.*

Evol. **3**, 159–174 (1994).

49.   Wiemers, M., Chazot, N., Wheat, C. W., Schweiger, O. & Wahlberg, N. A

544   complete time-calibrated multi-gene phylogeny of the european butterflies.

545   *Zookeys* **2020**, 97–124 (2020).

546   50.   Drummond, A. J. & Rambaut, A. BEAST: Bayesian evolutionary analysis by

547   sampling trees. *BMC Evol. Biol.* **7**, 1–8 (2007).

548   51.   Martin, S. H. *et al.* Natural selection and genetic diversity in the butterfly

549   heliconius melpomene. *Genetics* **203**, 525–541 (2016).

550   52.   Keightley, P. D. *et al.* Estimation of the Spontaneous Mutation Rate in Heliconius

551   melpomene. *Mol. Biol. Evol.* **32**, 239–243 (2015).

552   53.   Ceballos, F. C., Joshi, P. K., Clark, D. W., Ramsay, M. & Wilson, J. F. Runs of

553   homozygosity: windows into population history and trait architecture. *Nat. Rev.*

554   *Genet.* **19**, 220–234 (2018).

555   54.   Lin, J. *et al.* Probing the genomic limits of de-extinction in the Christmas Island

556   rat. *Curr. Biol.* **32**, 1650-1656.e3 (2022).

557   55.   Jones, P. *et al.* InterProScan 5: genome-scale protein function classification.

558   *Bioinformatics* **30**, 1236–1240 (2014).

559   56.   Gish, W. & States, D. J. Identification of protein coding regions by database

560   similarity search. *Nat. Genet.* **3**, 266–272 (1993).

561   57.   Bushnell, B. BBMap. (2014).

562   58.   Schmieder, R. & Edwards, R. Quality control and preprocessing of metagenomic

563   datasets. *Bioinformatics* **27**, 863–864 (2011).

564   59.   Wood, D. E., Lu, J. & Langmead, B. Improved metagenomic analysis with

565   Kraken 2. *Genome Biol.* **20**, 257 (2019).

568 **Tables**

Sample Identifier	Generated Reads	Q25 Unique Mapped Reads	Breadth of Coverage	Average Depth Covered Regions
L002	300,294,248	23,337,751	37.27%	5.105
L003	405,198,060	32,547,820	36.86%	6.78
L004	357,165,438	28,722,185	38.77%	6.55
L005	776,312,378	56,459,037	45.7%	12.42
L006	359,520,168	28,498,720	40.07%	6.18
L007	348,916,870	26,758,356	34.79%	6.21
L008	508,120,156	32,107,192	42,08%	7.422
L009	322,955,384	39,312,617	40.6%	8.02
L011	236,886,534	24,165,282	38.6%	5.40
L012	328,359,669	18,683,738	33.37%	4.29
L013	385,635,644	52,612,937	47.2%	12.3

Table 1. Mapping Statistics of the analysed historical specimens. Mapping

statistics of the 4 historical *G. xerces* (L003, L005, L007 and L009) and the 7 historical

G. lydagmus (L002, L004, L006, L008, L011, L012 and L013) specimens mapped

against the *G. alexis* reference genome. Average depth is displayed for the covered

regions of each individual.

Figure captions

Fig. 1. Phylogenetic placement of the Xerces Blue. A: Maximum likelihood tree from

whole mitochondrial genomes of Xerces Blue, Silvery Blue and Green-Underside Blue.

Node labels are bootstrap support values. Asterisk indicates the position of a

previously published mitochondrial DNA genome (MW677564.1)[4]. B: Time-calibrated

phylogeny from Bayesian inference using mitochondrial protein-coding genes of

Xerces Blue and related butterflies. Node values show median age estimates from

dating analysis with a molecular clock (above nodes) or from fixing the age of the root

(below nodes). Bars are 95% HPD intervals for node ages. All posterior probabilities

were 1, except for one node annotated in black. Images show upper sides (left) and

undersides (right) of male specimens for a selection of species. We used MN974526

as a proxy for *P. argus*; however, after detecting a high sequence identity (>99%) with

a previously published *P. melissa* mitogenome, we have decided to label it as *Plebejus*

sp. in the tree.

Fig. 2. PSMC plot of one Xerces Blue (*Glaucopsyche xerces*) (L05) specimen and

one Silvery Blue specimen (*Glaucopsyche lygdamus*). The two historical samples

are those with higher average coverage. Individual PSMC plots were bootstrapped 100

times each (lighter lines). One year of generation time and a mutation rate of $\mu=1.9 \times 10^{-9}$

were used. The peak of the Marine Isotopic Stage 7 interglacial is marked in yellow.

Fig. 3. Runs of Homozygosity (RoH) in the genomes of Xerces Blue and Silvery

Blue (modern and historical). a: Percentage of the autosomal genome in RoH by size

bins: very short RoH (<100 Kb), short RoH (100-500 Kb), intermediate RoH (500Kb-

1Mb) and long (1-5Mb). Short RoH reflect LD patterns, intermediate size RoH describe

background inbreeding due to genetic drift and long RoH appear in the case of very

recent inbreeding due to consanguinity. Error bars show the standard deviation. b:

Distribution of RoH in the autosomal genome of a Xerces specimen, L05 c) Distribution

of RoH in the autosomal genome of a Silvery specimen L13.

Fig. 4. Functional effect prediction on the fixed amino acid-changing alleles

observed in Xerces Blue and Silvery Blue. a: Wide genome Ka/Ks ratio comparison.

b: High-to-moderate effect variant comparison in homozygous sites. c: High-to-

moderate effect variant comparison in heterozygous sites. d: Presence of high-to-

moderate variants in regions of the genome in RoH. Error bars show the standard

deviation.

Specie   *G. lygdamus*   *G. xerces*