Sexual Dimorphic Metabolic and Cognitive Responses of C57BL/6 Mice to Fisetin or

Dasatinib and Quercetin Cocktail Oral Treatment

Yimin Fang[1], David Medina[2], Robert Stockwell[2], Sam McFadden[1], Kathleen Quinn[1],

Mackenzie R. Peck[1], Andrzej Bartke[2,3], Kevin N. Hascup[1,3,4], Erin R. Hascup[1,4,*]

[1] Department of Neurology, Dale and Deborah Smith Center for Alzheimer's Research

and Treatment, Neuroscience Institute, Southern Illinois University School of Medicine,

Springfield, IL 62702, USA.

[2] Department of Internal Medicine, Southern Illinois University School of Medicine,

Springfield, IL 62702, USA.

[3] Department of Medical Microbiology, Immunology and Cell Biology, Southern Illinois

University School of Medicine, Springfield, IL 62702, USA.

[4] Department of Pharmacology, Southern Illinois University School of Medicine,

Springfield, IL 62702, USA.

*Corresponding Author: Erin R. Hascup, Department of Neurology, Dale and Deborah

Smith Center for Alzheimer's Research and Treatment, Southern Illinois University

School of Medicine, P.O. Box 19628, Springfield, IL 62794-6928, USA Tel: 217-545-

6988, Email: ehascup@siumed.edu

Abstract

Senolytic treatment in aged mice clears senescent cell burden leading to functional improvements. However, less is known regarding the effects of these compounds when administered prior to significant senescent cell accumulation. From 4-13 months of age, C57BL/6 male and female mice received monthly oral dosing of either 100 mg/kg Fisetin or a 5 mg/kg Dasatinib (D) plus 50 mg/kg Quercetin (Q) cocktail. During treatment, several aspects of healthy aging were assayed including glucose metabolism using an insulin and glucose tolerance test, cognitive performance using Morris water maze and novel object recognition, and energy metabolism using indirect calorimetry. Afterwards, mice were euthanized for plasma and tissue specific markers of senescence-associated secretory phenotype (SASP) and white adipose tissue accumulation (WAT). Sexually dimorphic treatment effects were observed. Fisetin treated male mice had reduced SASP, enhanced glucose and energy metabolism, improved cognitive performance, and increased mRNA expression of adiponectin receptor 1 and glucose transporter 4. D+Q treatment had minimal effects in male C57BL/6 mice, but was detrimental to females causing increased SASP expression along with accumulation of WAT depots. Reduced energy metabolism and cognitive performance were also noted. Fisetin treatment had no effect in female C57BL/6 mice potentially due to a slower rate of biological aging. In summary, the senolytic treatment in young adulthood, has beneficial, negligible, or detrimental effects in C57BL/6 mice dependent upon sex and treatment. These observations should serve as a note of caution in this rapidly evolving and expanding field of investigation.

Graphical Abstract

Male and female C57BL/6 mice were treated with once monthly oral doses of either Dasatinib (D) + Quercetin (Q) or Fisetin from 4-13 months of age. Females treated with D+Q had increased adiposity and SASP markers (red spheres) along with decreased metabolism (blue flame) and cognitive performance. Males treated with Fisetin had reduced SASP markers (blue spheres) as well as improved metabolism (red flame) and cognition. No effects were observed in females treated with Fisetin or males treated with D+Q.

Introduction

Fisetin and quercetin (Q) are plant-derived flavonoids that offer cytoprotection against cellular stress and act as anti-inflammatory, chemopreventive, chemotherapeutic, and senotherapeutic agents (1, 2). Additionally, Q incites immune response against allergic diseases (2, 3). Dasatinib (D) is a tyrosine kinase inhibitor used to treat leukemia (4) and is routinely used in combination with Q to improve the senotherapeutic potency. Although both prooxidan flavonoids are structurally analogous, Fisetin has higher senolytic activity than Q whose potency is also reliant on endogenous trace elements such as iron and copper (5).

As the first senolytic drugs, they were discovered to selectively clear senescent cells generated during cellular senescence (6). Cell senescence is a phenomenon defined as a stable cell cycle arrest, and is associated with fundamental aging processes and age-related diseases (7). Although this process is essential for preventing replication of damaged DNA thereby suppressing tumor formation for cancer, accumulation of senescent cells with aging produces senescence-associated secretory phenotype (SASP) (7, 8). SASP includes proinflammatory cytokines, chemokines, growth factors, and proteases (9, 10) caused by chronic inflammation (10-12), DNA damage (13, 14), mitochondrial dysfunction (15), immune cell dysfunction (16), ROS generation (13, 17), and brain protein aggregation (18, 19). All of which may be factors that predispose individuals to a multitude of age-related disorders (7).

Fisetin and D+Q selectively clear senescent cells (6), thereby delaying aging-associated disorders and improving healthspan and lifespan. This has been observed after reducing senescent cell burden in progeroid or aged (twenty-four-month-old)

C57BL/6 mice (20-22). Moreover, deletion of senescent cells from the brain genetically

or pharmacologically with senolytic drugs led to functional improvements in mouse

models of neurodegenerative diseases such as Parkinson's and Alzheimer's disease

(18, 23-25). These studies have shown senotherapeutics can reduce senescent cell

burden and have positive impacts on animals with accelerated aging, advanced age, or

neurodegenerative disorders. Accordingly, senotherapeutics are currently marketed as

anti-aging therapies where young, healthy adults can take these products as dietary

supplements.

However, less is known about the effects of these compounds when

administered prior to significant senescent cell accumulation. Thus, the experiments

were designed to examine the long-term effects of monthly oral Fisetin or D+Q

treatment when initiated in young (four-month-old) C57BL/6 mice. We examined

morphological, metabolic, physical, and cognitive components that are known to be

affected by senescent cell accumulation. The results presented here indicate that

monthly administration of Fisetin or D+Q had sexually dimorphic effects which also

depended on treatment type in C57BL/6 mice. We highlight a potential new mechanism

of action involving the beneficial roles of glucose and adiponectin signaling both

peripherally and centrally by early and long-term administration of Fisetin on male

C57BL/6 mice.

Results

Administration of Fisetin or D+Q altered the SASP profile of C57BL/6 mice in a sex dependent manner.

It has been reported that senolytic drug treatment decreased senescent burden in twenty-four-month-old C57BL/6 mice leading to functional improvements (20). However, it was unknown whether the same treatment administered monthly initiated during young adulthood (four months of age) would improve the overall health of C57BL/6 mice as they aged. To examine this, Fisetin or a cocktail of D+Q were orally administered in C57BL/6 mice starting at four months of age and continued once each month for nine months. All mice underwent glucose tolerance test (GTT) and insulin tolerance test (ITT) to evaluate glucose homeostasis, indirect calorimetry to determine energy metabolism, grip force to monitor muscle strength, and Morris water maze (MWM) and novel object recognition (NOR) to ascertain cognitive abilities. At approximately thirteen months of age, the mice were sacrificed to determine body composition and collect plasma and tissues for analyses of SASP markers, lipid metabolism, glucose and adiponectin signaling, and synaptic plasticity. The experimental paradigm is shown in Fig 1a.

To investigate whether the compounds used in this study decreased SASP profiles and other senescent markers, the gene expressions were measured in gonadal white adipose tissue (WAT) by RT-PCR. To evaluate aging effects, additional groups of treatment naïve young (four months of age) and old (twenty-one months of age) mice were studied. We assayed for SASP markers including $p16^{Ink4\alpha}$, $p21^{Cip1}$, TNFα, MCP 1, IL-6, and IL-10. Although IL-10 is a potent anti-inflammatory cytokine, studies have

shown IL-10 activates cellular senescence mechanisms (26) and is secreted from

senescent cells (27). The results showed that in both the male and female mice, the

transcriptional levels of these SASP markers were lower in the treatment naïve young

compared to old C57BL/6 mice (Fig. 1b-g). The vehicle treated littermate control

C57BL/6 mice sacrificed at thirteen months of age had similar mRNA levels of the

above mentioned SASP markers as the untreated twenty-one-month-old mice (Fig. 1b-

g). This is consistent with recent tissue and age-specific RNA sequencing in both sexes

of C57BL/6 mice where a significant increase in differentially expressed genes was first

evident in gonadal WAT that occurred by mid-age and was maintained at ages older

than twenty months (28). Others have shown p16$^{Ink4\alpha}$ and p21^{Cip1} have similar mRNA

expression levels at 12 and 24 months of age in the hypothalamus, heart, liver, and

kidney of female CB6F1 mice (29).

We also observed that C57BL/6 males had higher levels of mRNA expression of

TNFα and IL-6 (Fig. 1d, f) compared to littermate females. Interestingly, the effects of

the treatments were sex-dependent. The treatment with Fisetin or D+Q reduced mRNA

levels of p16$^{Ink4\alpha}$ in the male mice, but had no effects in the female mice (Fig. 1b). The

treatment with Fisetin reduced the mRNA levels of p21^{Cip1}, TNFα, and IL-6 in the male

mice, but had no effect in the female mice (Figs. 1c, 1d, & 1f), while the treatment with

D+Q raised the mRNA levels of p21^{Cip1}, MCP1, IL-6, and IL-10 (Figs. 1c & 1e-g) in the

female mice.

The same SASP mRNA levels were examined in the hippocampus (SI Fig. 1a-f).

mRNA levels were higher in both sexes of treatment naïve twenty-one-month-old

C57BL/6 mice compared to four-month-old littermates. p21^{Cip1} was the only SASP

marker increased in the hippocampus of female C57BL/6 mice compared to male

littermates. In general, mRNA hippocampal expression patterns after senotherapeutic

treatment was similar to observations in gonadal WAT. In male C57BL/6 mice, Fisetin

treatment reduced mRNA levels of p16$^{Ink4\alpha}$ and p21^{Cip1} (SI Fig 1a-b), but did not affect

the other assayed cytokines (Fig 1c-f). Reduced hippocampal mRNA levels of p21^{Cip1}

(SI Fig. 1b) were also observed in female C57BL/6 mice. D+Q treatment increased

mRNA levels of TNFα (SI Fig. 1c) and IL-10 (SI Fig. 1e) in both sexes of C57BL/6 mice.

Plasma concentrations of TNFα, MCP1, IL-6, and IL-10 (SI Fig. 2a-d) were

elevated in both sexes of treatment naïve twenty-one-month-old C57BL/6 mice

compared to four-month-old littermates. The plasma concentration of TNFα was

increased in males, while IL-10 was increased in females compared to sex-matched

C57BL/6 littermates. Fisetin treatment decreased the concentration of MCP1 (SI Fig.

2b) in males while increasing IL-10 (SI Fig. 2d) concentration in female C57BL/6 mice.

D+Q treatment had no effect in males, but increased the concentration of MCP1, IL-6,

and IL-10 (SI Fig. b-d) in female C57BL/6 mice.

Altogether, the data showed that Fisetin treatment reduced the levels of SASP

markers in the plasma as well as the peripheral and central tissue of male C57BL/6

mice, while little to no changes were observed in female littermates. The D+Q treatment

increased, rather than decreased, SASP markers in female C57BL/6 mice. In male

C57BL/6 mice, D+Q treatment had minimal to no effects in plasma and WAT, but

increased hippocampal TNFα mRNA expression.

D+Q treatment led to increased body weight and adiposity in female C57BL/6 mice while Fisetin treatment improved glucose metabolism and increased plasma adiponectin in male C57BL/6 mice.

Given the observation that C57BL/6 mice responded to the treatment in a sexually dimorphic manner, it was important to know how the physiological parameters including body composition, glucose utilization, and energy metabolism were affected by senotherapeutic treatment. The results showed that body weight in the males was not changed by either treatment, while it was increased in the female C57BL/6 mice receiving monthly D+Q administration (Fig. 2a-b). Body composition analysis indicated that the increased body weight of D+Q treated female C57BL/6 mice was attributed to an elevation of adiposity including subcutaneous (SC) and gonadal WAT depots (Figs. 2c-e). No changes in body weight or composition were observed after nine months of Fisetin treatment in female C57BL/6 mice, and no changes in these parameters were detected in male C57BL/6 mice receiving either of the treatments (Figs 2a-g). Body composition also showed sex differences whereby female C57BL/6 mice had a higher percentage of SC and gonadal WAT while males had a larger perirenal depot (Fig. 2d-f).

The accumulation of WAT in D+Q treated female mice could result from upregulated lipid deposition or downregulated thermogenic activity in brown adipose tissue (BAT). Thus, the expressions of genes related to lipid metabolism in gonadal WAT, and to uncoupling of the mitochondrial electron transport chain in BAT were examined. The results showed that Acetyl-CoA carboxylase (ACC), the rate-limiting enzyme for lipid synthesis, was up-regulated in gonadal WAT of D+Q treated female

C57BL/6 mice (Fig. 2h). In contrast to WAT, uncoupling protein 1 (UCP1), a key

regulator of non-shivering thermogenesis, and its upstream activator PPARγ (30, 31)

were downregulated in BAT of D+Q treated female mice (Fig. 2i-j). Sex effects were

observed with higher ACC in female C57BL/6 gonadal WAT and UCP1 in male

interscapular BAT. The data from WAT and BAT support that D+Q treated female mice

had higher lipid synthesis activity, mediated by increased ACC, and lower thermogenic

activity, via reduced UCP1 and PPARγ. These changes could cause WAT

accumulation subsequently leading to the observed SASP increases after D+Q

treatment in female C57BL/6 mice.

Considering adipocytes are involved in glucose metabolism, we wanted to

determine treatment effects on peripheral glucose utilization. The results showed that

Fisetin treatment improved glucose clearance as measured by GTT (Figs. 2k, m) in

male mice. Although D+Q treated female mice accumulated more adipose tissues they

did not have impaired glucose metabolism (Figs. 2l-m). No differences in insulin

sensitivity between treated and littermate control mice were observed as determined

with ITT (Figs. 2n-p). Additionally, plasma insulin concentrations (Fig. 2q) were similar

across treatments in both sexes of C57BL/6 mice. Plasma adiponectin (an adipokine

that is involved in regulating glucose levels) concentration was significantly increased in

the D+Q and Fisetin treated males, but not after either treatment in female C57BL/6

mice (Fig. 2r). The concentration of plasma insulin was elevated while adiponectin was

decreased in male C57BL/6 mice compared to female littermates. This is consistent

with the literature where adiponectin levels in male mice (32) and humans (33) are

lower compared to females. The data indicated that monthly Fisetin treatment improved

peripheral glucose metabolism in the male mice with increased adiponectin levels and

enhanced glucose clearance. In female C57BL/6 mice, D+Q treatment increased WAT

accumulation without affecting glucose metabolism. This may be reflective of the GTT

assay being conducted in the middle rather than at the end of the experimental design.

Fisetin treatment enhanced energy metabolism in male C57BL/6 mice, while D+Q

treatment reduced energy metabolism in female C57BL/6 mice.

214   Energy metabolism is strongly associated with aging and is a contributing factor

to adipose accumulation (34, 35). In the present study, the treatment effects on energy

metabolism were assessed by indirect calorimetry. Oxygen consumption (VO_2) (Figs.

3a, c) and energy expenditure (EE) (Fig. 3d, f) were increased after Fisetin treatment in

male C57BL/6 mice, while both parameters were decreased by D+Q treatment in

female littermates (Figs. 3b-c, e-f). Thus, Fisetin enhanced energy metabolism in male

C57BL/6 mice, while D+Q reduced energy metabolism in the female mice, potentially

contributing to their WAT accumulation. Respiratory quotient (RQ) is the ratio of carbon

dioxide produced and oxygen consumed by the body, reflecting substrate utilization for

energy generation (36). In the current study, RQ for Fisetin treated male C57BL/6 mice

was reduced indicating a shift in metabolic substrate utilization from mixed lipids and

carbohydrates (RQ ~ 0.8) to predominantly lipids (RQ ~ 0.7) (36) (Fig. 3g, i). The RQ of

female C57BL/6 mice treated with D+Q did not change (Fig. 3f). Female C57BL/6 mice

had higher levels of all three measures compared male littermates. In summary,

monthly Fisetin treatment in male C57BL/6 mice increased energy metabolism as

indicated by enhanced VO_2 and EE and shifted substrate utilization to predominantly

lipids. In contrast, D+Q treatment decreased energy metabolism in female C57BL/6

mice as indicated by reduced VO_2 and EE without a concomitant change in substrate

utilization.

Fisetin treatment improved spatial cognitive performance in male C57BL/6 mice,

while D+Q treatment impaired object recognition in female C57BL/6 mice.

Although deletion of senescent cells by the senolytic compounds led to functional

improvements in models of neurodegenerative diseases (23, 24), it remains to be

elucidated whether the monthly senolytic treatment could affect cognitive performance

in normal aging mice. To investigate the effects of the treatment on cognitive functions,

the MWM spatial learning and memory paradigm and a NOR task were performed. The

results from MWM indicated that over 5 training sessions, the path efficiency was

increased (Fig. 4a, c), and the corrected integrated path length (CIPL) (Fig. 4d, f) was

reduced significantly in the Fisetin-treated male C57BL/6 mice indicative of improved

learning. Neither Fisetin nor D+Q treatment affected MWM learning in the female

C57BL/6 mice (Figs. 4b-c, e-f). Platform entries during the delayed probe challenge was

not affected by either treatment in both sexes (Fig. 4g) suggesting spatial memory recall

was unaffected. The results of NOR showed that the exploration preference for the

novel object measured by retention index was decreased in the D+Q treated female

mice (Fig. 4h), indicating reduced memory recall in this group. Since senotherapeutic

treatment had differential effects on energy expenditure between the sexes, we wanted

to verify changes to cognitive performance were not due to motor function. Mean

swimming speed was similar for all treatment groups and across sexes for the training

sessions and probe challenge (SI Figs. 3a-d). Additionally, no treatment differences in

grip strength were observed (SI Fig. 3e). Female C57BL/6 mice had better grip strength

compared to littermate males, which was consistent with other reports in C57BL/6 mice

(37).

Considering the effects of the treatments on cognitive performance, several

genes related to memory formation and synaptic plasticity (38) were examined by RT-

PCR in the hippocampus. Genes associated with synaptic vesicle release and

glutamatergic signaling were not affected by either treatment in male or female C57BL/6

mice (SI Fig. 4). Next, we examined glucose transporter 4 (Glut4) hippocampal mRNA

expression because central glucose metabolism is directly related to cognitive

performance. Glut4 was increased in the hippocampus of Fisetin treated male C57BL/6

mice while D+Q had no effect in either sex (Fig. 4i). Peripherally, D+Q and Fisetin

treatment increased Glut4 liver expression in male C57BL/6 mice only, but neither

treatment altered expression in gonadal WAT (Figs. 4j-k). Since plasma adiponectin

concentration was increased after senotherapeutic treatment in male C57BL/6 mice, we

also examined adiponectin receptor 1 (AdipoR1) mRNA expression in both the

hippocampus and peripheral tissues. Adiponectin signaling modulates glucose

metabolism while providing anti-inflammatory effects both of which improve cognitive

performance. Fisetin treated male C57BL/6 mice had increased hippocampal AdipoR1

expression, but D+Q treatment did not alter expression levels of either sex (Fig. 4l).

Peripherally, neither senotherapeutic treatment affected AdipoR1 expression in the liver

nor gonadal WAT of male C57BL/6 mice (Figs 4m-n). D+Q treatment reduced AdipoR1

expression in the liver while increasing expression in gonadal WAT of female C57BL/6

mice (Figs. 4m-n). This dichotomy is a result of tissue specific AdipoR1 regulation on

lipid storage and utilization (39). Decreased AdipoR1 signaling in the liver increases

hepatic lipogenesis causing triglyceride release and accumulation. Increased AdipoR1

expression in WAT reduces lipolysis. Together, this helps explain the increased body

and WAT weight seen in C57BL/6 females mice receiving D+Q treatment. Overall, the

effects of serotherapeutic treatment on Glut4 and AdipoR1 expression is consistent with

changes to peripheral glucose metabolism (Figs. 2) and energy expenditure (Figs. 3).

Comparing the rate of biological aging between sexes of C57BL/6 mice.

Finally, to better understand why senotherapeutic treatment, particularly Fisetin,

was only effective in males, we compared physical (SI Fig. 5a), metabolic (SI Fig. 5b-f),

and cognitive (SI Fig. 5g-i) in four and thirteen-month-old C57BL/6 mice to determine

potential differences in the rates of biological aging. Compared to female littermates,

four-month-old mice male C57BL/6 mice have worse metabolic parameters as observed

by GTT, oxygen consumption, and energy expenditure. Physical and cognitive

parameters are similar between sexes at this age. By 13 months of age, an aging effect

is observed whereby physical, metabolic, and cognitive parameters were reduced in

both sexes of C57BL/6 mice. Since swimming speed declines with age this can

influence the duration to the platform and the number of platform crosses on the MWM

(SI Figs. 5g, i, j). Although the CIPL calculation avoids variability in results due to

swimming speed differences, both sexes of 13-month-old C57BL/6 mice had worse

spatial navigation learning and memory recall compared to their younger sex-matched

littermates (SI Fig. 5h, k). These baseline differences between sexes of C57BL/6 mice

may partially explain why Fisetin was effective only in males.

Discussions and conclusions

Administration of senolytic compounds is a novel approach to removing

senescent cell accumulation and thereby reducing inflammation. These compounds are

being investigated as potential lifespan and healthspan extending therapies and may

also have applications in the treatment of numerous metabolic and neurodegenerative

disorders (18, 23-25). Since Fisetin and Q are naturally occurring plant flavonoids, they

are readily available as dietary supplements for use without a prescription. There is little

information on the efficacy of these compounds when taken at young ages and for

prolonged periods of time. The data presented here indicates that these compounds

have sexually dimorphic effects, each with health benefits and risks. The monthly

dosing strategy used in this study was based on previous publications showing that

markers of cell senescence are still reduced after a four week off-treatment period (7).

Prior research suggests that senolytic compounds work either by clearing senescent

cells or reversing cellular senescence rather than preventing cells from entering a

senescent state. Since senescent cell accumulation occurs over weeks, maintaining an

effective concentration dosage is not required. Rather, intermittent dosing may be

preferable thereby avoiding adverse effects.

Monthly Fisetin treatment was efficacious in male C57BL/6 mice when started at

four months of age. This dosing strategy had little to no effect in female littermates

when compared to vehicle treatment. Nine months of Fisetin treatment efficiently

reduced SASP in WAT (Fig. 1), hippocampus (SI Fig. 1), and plasma (SI Fig. 2), of

thirteen-month-old male C57BL/6 mice as lower mRNA expression of $p16^{Ink4\alpha}$, $p21^{Cip1}$,

TNFα, and IL-6 were observed. Male C57BL/6 mice receiving Fisetin treatment also had

improved glucose metabolism as supported by better glucose clearance (Figs. 2k, m)

after five treatments and elevated plasma adiponectin (Fig. 2r) collected upon

euthanization at thirteen months of age. Improved energy metabolism was also

supported by increased VO_2 and EE (Figs. 3a, c-d, f) along with reduced RQ (Fig. 3g, i)

when measured at 12 months of age. Monthly Fisetin treatment also reduced SASP

markers to similar levels as four-month-old treatment naïve male C57BL/6 mice. The

metabolic improvements also mirrored levels observed in younger littermates. But

physical parameters (swimming speed and grip strength) as well as learning and

memory recall was still worse after Fisetin treatment when compared with younger male

C57BL/6 mice. This suggests administering Fisetin treatment at younger ages has a

stronger anti-inflammatory (potentially mediated through increased plasma adiponectin

signaling) rather than an overall anti-aging effect. This cannot be definitively concluded

without frailty or lifespan measurements which were outside the scope of the study.

Also, inflammation and aging are not mutually exclusive whereby chronic inflammation

contributes to worse health and the onset numerous age-related ailments and

neurodegenerative disorders. Approaches to reduce chronic inflammation have the

potential to extend quality of life during aging.

In the present study, D+Q treatment showed minimal efficacy in male C57BL/6

mice and was detrimental to female mice. In females, SASP markers (Fig. 1) and body

weight were increased attributed to larger WAT depots (Fig. 2a-e). Several factors

contributed to this increased abdominal adiposity. These include increased ACC (the

rate-limiting enzyme in lipid synthesis) and AdipoR1 in WAT (Fig. 2h and Fig. 4n),

reduced UCP1 and PPARγ activity in BAT (Figs. 2i-j), reduced energy metabolism and

liver AdipoR1 expression (Fig. 3 and Fig. 4m). Because obesity can lead to adipose

tissue entering a senescent-like state at a young age (40), the accumulation of

abdominal WAT in D+Q treated female mice could result in increased SASP. This

possibility is supported by expression of SASP-related genes in D+Q treated females

resembling or even exceeding the values measured in much older (twenty-one-month-

old) untreated females (Fig. 1, SI Fig. 1, and SI Fig. 2). The reasons for the detrimental

outcome of D+Q treatment in the female mice remain to be fully elucidated, but are not

unique to this study. A recent publication using a similar D+Q dosing regimen reported

similar observations in a C57BL/6 mouse model of hepatocellular carcinoma (41).

Others have proposed that continuous senescent cell removal beginning at young ages

does not always activate cell-replacement mechanisms, but rather induces regenerative

responses causing fibrosis (42). A better understanding of cell-type specific effects of

senescent cell removal is warranted.

Sexually dimorphic biological aging may partially explain the senotherapeutic

differences observed in C57BL/6 mice. Four-month-old male and female C57BL/6

treatment naïve mice have similar SASP mRNA and plasma concentrations (Fig. 1 and

SI Fig. 2), but males have worse metabolism (SI Figs. 5b, d-e). SASP became more

prominent in male C57BL/6 mice compared with age-matched thirteen-month females,

particularly proinflammatory cytokines TNFα and IL-6 (Fig. 1d, f and SI Fig. 2a, c).

Since Fisetin shows therapeutic efficacy by reducing inflammation leading to metabolic

improvement, these SASP factors were not at harmful levels in female C57BL/6 mice.

This suggests male C57BL/6 mice age biologically faster and may explain why Fisetin

treatment was ineffective at the ages tested in female littermates. Unfortunately, data is

only beginning to emerge regarding sexually dimorphic aging characteristics due in large part from male centric sampling in both preclinical and clinical research. Sparse evidence suggests females maintain better cellular health throughout the majority of their life, but become frailer when approaching death compared with males (43).

In C57BL/6 mice, cell senescence usually becomes prominent by approximately 14 months of age (7), and treatment with senolytic drugs at this or a later ages cleared senescent cells and led to various beneficial effects (20, 21). However, it was unclear whether chronic treatment with senolytic drugs started at an earlier age (prior to reported senescent cell accumulation) in C57BL/6 mice would have similar effects as the reported observations during later age (20, 21). This gap in knowledge is particularly relevant given that Fisetin and Q are available without a prescription and may be taken by younger adults. Our data revealed that such treatment indeed had impacts on the mice as they reached median lifespan. However, the senotherapeutic treatment effects were sexually dimorphic. C57BL/6 male, not female, mice receiving Fisetin treatment had beneficial responses mediated through reduced SASP and improved glucose and energy metabolism. Female, not male, mice receiving a D+Q cocktail treatment had detrimental responses with reduced energy metabolism, increased WAT accumulation, and elevated SASP.

D+Q was the first well-characterized senotherapeutic agent proven to reduce senescent cell burden and extend lifespan. Despite Q and Fisetin being structural analogs, Q had minimal to no effects when initiated at a young age in male C57BL/6 mice. Although both are potent redox-active flavonoids, Fisetin has higher prooxidant activity. Furthermore, Q's senolytic potency is augmented by trace elements such as

copper and iron that are known to accumulate during aging and in senescent cells (5).

Since senolytic drugs were administered at a young age prior to reported senescent cell

accumulation, iron and copper accumulation was not a threshold to enhance Q's

potency. Fisetin's prooxidant activities are not amplified in the presence of copper or

iron, which may explain why health benefits were only observed with this

senotherapeutic in male C57BL/6 mice.

Moreover, Fisetin-treated male C57BL/6 mice not only had improved markers of

peripheral health, but also their spatial learning during the MWM training sessions (Figs.

4a, c, d, f) compared to vehicle treated C57BL/6 mice. These cognitive improvements,

however, did not reach levels observed in 4-month-old treatment naïve mice (Fig. 4g

and SI Fig. 5i). Although we did not observe changes in markers of synaptic plasticity

associated with hippocampal memory formation (SI Fig. 4), we did observe elevated

expression of Glut4 (Fig. 4i-j), hippocampal AdipoR1 (Fig. 4l), and plasma adiponectin

concentrations (Fig. 2r) suggesting enhanced glucose transport and adiponectin

signaling in the brain. This novel finding that Fisetin treatment concurrently improved

peripheral as well as central glucose and adiponectin signaling has important

implications for metabolic and cognitive function. In the periphery, adiponectin increases

insulin sensitivity, stimulates fatty acid oxidation and glucose uptake, suppresses

hepatic glucose production (44), and inhibits inflammation (45). In the brain, adiponectin

signaling via AdipoR1 has anti-inflammatory effects and is involved with cognitive and

neuroprotective mechanisms (46). Monthly Fisetin treatment in male mice led to

increased adiponectin levels in circulation, which likely contributed to improvements of

peripheral glucose and lipid metabolism. Circulating adiponectin passes through the

blood brain barrier where it can enhance cognitive function by upregulating glucose

transport and AdipoR1 signaling (46). However, neither hippocampal Glut4 (Fig. 4i),

AdipoR1 (Fig. 4l), nor the examined synaptic plasticity markers (SI Fig. 4) were affected

by D+Q treatment in the female mice even though object recognition memory retention

was reduced (Fig. 4h). The reduced cognitive performance of these animals might be

attributed to the increased adiposity after D+Q treatment, but future studies are needed

to identify the mechanisms involved.

Cell senescence is strongly correlated with aging, but currently no diagnostic

testing is available to quantify senescent cells accumulation. It seems possible that the

earlier in life senotherapeutics are taken, the better likelihood of delaying age-related

decline and disease risk might be expected. Senolytics such as Fisetin and D+Q are

known to reduce senescent cell burden (7), but the optimal age to begin treatment is

currently unknown. Our study indicates that both biological aging and sex may also

determine the therapeutic outcomes of senolytic treatment. When the treatment was

started at 4 months of age, before the reported senescent cell accumulation, Fisetin had

beneficial effects in male C57BL/6 mice while a D+Q cocktail had adverse

consequences in female C57BL/6 mice.

These observations provide novel information with translational relevance. First,

senolytic drugs taken at an age before significant senescent cell burden to reduce or

prevent their prevalence later in life may be detrimental to overall health. Second, males

and females can have differential responses to the same senolytic treatment when

initiated at younger ages. Third, a particular senolytic treatment may have beneficial,

negligible or detrimental effects depending on the age, sex, or disease state. Fourth,

oral senolytic treatment not only impacted peripheral physiological responses, but also

affected cognitive performance, possibly via increased central adiponectin signaling and

glucose transport. Overall, these observations should serve as a note of caution in this

rapidly evolving and expanding field of investigation.

Designing senotherapeutics with the role of sex and age of onset could

conceivably provide a useful preventative treatment strategy for dealing with

development of neurodegenerative disorders. Supporting this possibility, development

of metabolic syndrome in midlife was reported to increases the risk for Alzheimer's

disease (AD) later on (47). Besides clearance of senescent cells, senotherapeutics may

ameliorate or delay cognitive decline through mechanisms associated with improved

peripheral and central glucose metabolism. Considering metabolic dysregulation has

been observed in the APP/PS1 model of AD (48), Additional studies are warranted to

determine if prodromal senolytic treatment could provide cognitive benefits in

neurodegenerative diseases.

Material and methods

Chemicals

Unless otherwise noted, all chemicals were obtained from Sigma-Aldrich (St. Louis, MO) including Quercetin (Cat# RHR1488). Fisetin was purchased from Selleckchem (Houston, TX; Cat #S2298), and Dasatinib from LC laboratories (Woburn, MA; Cat# D-3307).

Animals

Male and female C57BL/6 mice were maintained in our established breeding colonies. Protocols for animal use were approved by the Institutional Animal Care and Use Committee at Southern Illinois University School of Medicine. Mice were group housed on a 12:12 h light-dark cycle, and food (Chow 5001 with 23.4% protein, 5% fat, and 5.8% crude fiber; LabDiet PMI Feeds) and water were available ad libitum. Sentinel mice located in the same rooms as our breeder and experimental mice were tested triannually for pyrogens and pathogens and none were detected. All assays, including tissue collection, were conducted one week after monthly senotherapeutic dosing. As shown in Fig 1a, glucose metabolism was assessed via ITT and GTT after six treatments at the animal age of nine months. Grip Strength was measured after seven treatments while NOR and MWM was performed after eight treatments. Energy metabolism was measured by indirect calorimetry at the animal age of twelve. A week following the final senotherapeutic treatment at 13 months, body weight was recorded then euthanized with an overdose of isoflurane. A cardiac puncture was used to collect blood for assessment of circulating SASP markers and proteins. Mice were rapidly

decapitated and the peripheral tissues were weighed to calculate their contribution to

total body weight. Tissue was immediately flash frozen and stored at -80˚C until

processing as previously described (49).

Senolytic drug treatment

Senotherapeutic concentrations and dosing strategy were based on previous

publications (20, 50-52). C57BL/6 mice were dosed with 100 mg/kg of Fisetin while a

cocktail of 5 mg/kg of D + 50 mg/kg of Q, or vehicle (2% DMSO in canola oil) by oral

administration (53). The treatments were given once per month from 4-13 months of

age.

Glucose tolerance test (GTT) and insulin tolerance test (ITT)

GTT or ITT was carried out as described previously (49). Sixteen-hour-fasted mice

underwent GTT by intraperitoneal (i.p.) injection with 1 g glucose per kg of body weight

(BW). Blood glucose levels were measured at 0, 15, 30, 45, 60, and 120 min with a

PRESTO glucometer (AgaMatrix). For ITT, nonfasted mice were injected i.p. with 1 IU

porcine insulin from sigma (St. Louis, MO) (Cat# I5523) per kg of BW. Blood glucose

levels were measured at 0, 15, 30, 45, 60, and 120 min. The data for GTT are presented

as absolute value, and for ITT are presented as a percentage of baseline glucose.

Indirect calorimetry

Indirect calorimetry was performed as previously described (49) using AccuScan

Metabolic System (AccuScan Instruments). In this system, mice are housed individually

in metabolic chambers with ad libitum access to food and water. After a twenty-four-hour

acclimation period, VO_2, VCO_2, EE and RQ measurements were collected every ten min

per animal and averaged for each hour.

Morris water maze (MWM) training and probe challenge

The MWM was used to assess spatial learning and memory recall, and performed as

previously described (54). Mice were trained to utilize visual cues placed around the room

to repeatedly swim to a static, hidden escape platform (submerged one cm below the

opaque water surface), regardless of starting quadrant. The MWM paradigm consisted of

5 consecutive training days with three, 90 s trials/day and a minimum inter-trial-interval of

20 min. Starting quadrant was randomized for each trial. After two days without testing,

the escape platform was removed and all mice entered the pool of water from the same

starting position for a single 60 s probe challenge to test long-term memory recall. The

ANY-maze video tracking system (Stoelting Co., Wood Dale, IL) was used to record

mouse navigation during the training and probe challenge. The three trials for each

training day were averaged for each mouse.

Novel Object Recognition (NOR)

The NOR was used to evaluate memory retention based on the premise that mice

spend more time exploring a novel object rather than a familiar object if memory

capabilities remain intact. Mice were habituated in the open field chamber for 30 min on

the first day. Twenty-four hours after, the mouse was returned to the chamber and

presented with two similar objects for 5 min. A 24-hour inter-session-interval was used

between introduction and retention phases to assess long-term memory retrieval.

During the retention phase one of the familiar objects is replaced with a novel object

and the mouse is given 5 minutes of exploration. The ANY-maze video tracking system

(Stoelting Co., Wood Dale, IL) was used to record mouse navigation during the

familiarization and retention phases and time spent at each object. The time spent

exploring the novel object was divided by the total time spent exploring both objects to

calculate the retention index.

Grip Strength

Using their forepaws, mice grasp a wire grid connected to an isometric force transducer

(Harvard Apparatus; Holliston, MA) and are gently pulled horizontally away by the tail.

Force is measured in grams and normalized according to body weight. Three trials with

a minimum inter-trial-interval of twenty minutes are averaged per mouse.

Assessment of blood chemistry

Plasma was collected from animals anesthetized with isoflurane by cardiac puncture at

sacrifice. The blood was mixed with EDTA, followed by centrifugation at 10,000 g for 15

min at 4°C for plasma collection. Per the manufacturer's protocol, insulin or adiponectin

was measured with respective ELISA kits (Crystal Chem, Elk Grove Village, IL; Cat#

90080 and 80569). Plasma cytokine levels were assayed using a multiplex immunoassay

on a MESO QuickPlex SQ 120 with accompanying software (Meso Scale Diagnostics,

LLC).

RT–PCR

mRNA expression was analyzed by quantitative RT–PCR as previously described (49)

using cDNA synthesis kits and SYBR green mix from Bio-Rad (Cat# 1708897 and

1725121) performed with the StepOne Real-Time PCR System (Thermo Fisher

Scientific). RNA was extracted using an RNeasy mini kit or RNeasy Lipid Tissue Mini Kit

(Qiagen) following the manufacturer's instructions. Relative expression was calculated as

previously described and primers were purchased from Integrated DNA Technologies

(Supplemental Information Table 1).

Statistical analysis

Statistical analyses were conducted using a two-way ANOVA to test for significance of

sex (males × females), treatment (Vehicle × Fisetin × D+Q), and an interaction between

sex and treatment. Difference between two groups were assessed with unpaired two-

tailed Student's t tests with significance defined as $p \leq 0.05$. The area under the curve

was calculated using the two-way ANOVA. Data are presented as means ± SEM. All

statistical analyses and graphs were completed using Prism 9 (GraphPad Inc, La Jolla,

CA, USA).

Acknowledgements

This work was supported by the National Institutes of Health NIA R01-AG057767 and NIA R01-AG061937, Dale and Deborah Smith Center for Alzheimer's Research and Treatment, Kenneth Stark Endowment (YF, SF, KQ, MRP, KNH, ERH), NIA R21-AG062985, and American Diabetes Association 1-19-IBS-126 (DM, RS, AB). We would like to thank Melissa Roberts for performing and analyzing the plasma cytokine multiplex assay and Lisa Hensley for editorial support.

Author Contributions:

YF, DM, RS, SM, KQ, and MRP conducted the experiments and analyzed the data. YF, AB, KNH, and ERH conceived the study, designed the experiments, interpreted the data, and wrote the manuscript. All authors approved the final version of the manuscript.

Data Availability

All data is available upon reasonable request.

Data Citation

All data was generated by the authors. No external sources of data were used to create any figures presented in this manuscript.

Conflict of Interest

The authors declare no conflicts of interest.

Reference

1.   Grynkiewicz G & Demchuk OM (2019) New Perspectives for Fisetin. *Front Chem* 7:697.

2.   Jafarinia M*, et al.* (2020) Quercetin with the potential effect on allergic diseases. *Allergy Asthma*
Clin Immunol 16:36.

3.   Mlcek J, Jurikova T, Skrovankova S, & Sochor J (2016) Quercetin and Its Anti-Allergic Immune
Response. *Molecules* 21(5).

4.   Aguilera DG & Tsimberidou AM (2009) Dasatinib in chronic myeloid leukemia: a review. *Ther Clin*
Risk Manag 5(2):281-289.

5.   Wang Y, He Y, Rayman MP, & Zhang J (2021) Prospective Selective Mechanism of Emerging
Senolytic Agents Derived from Flavonoids. *J Agric Food Chem* 69(42):12418-12423.

6.   Zhu Y*, et al.* (2015) The Achilles' heel of senescent cells: from transcriptome to senolytic drugs.
Aging Cell 14(4):644-658.

7.   Kirkland JL & Tchkonia T (2020) Senolytic drugs: from discovery to translation. *J Intern Med*
288(5):518-536.

8.   Campisi J (2005) Senescent cells, tumor suppression, and organismal aging: good citizens, bad
neighbors. *Cell* 120(4):513-522.

9.   Kuilman T & Peeper DS (2009) Senescence-messaging secretome: SMS-ing cellular stress. *Nat Rev*
Cancer 9(2):81-94.

10.   Coppe JP, Desprez PY, Krtolica A, & Campisi J (2010) The senescence-associated secretory
phenotype: the dark side of tumor suppression. *Annu Rev Pathol* 5:99-118.

11.   Lasry A & Ben-Neriah Y (2015) Senescence-associated inflammatory responses: aging and cancer
perspectives. *Trends Immunol* 36(4):217-228.

12.   Freund A, Orjalo AV, Desprez PY, & Campisi J (2010) Inflammatory networks during cellular
senescence: causes and consequences. *Trends Mol Med* 16(5):238-246.

13.   McHugh D & Gil J (2018) Senescence and aging: Causes, consequences, and therapeutic avenues.
J Cell Biol 217(1):65-77.

14.   Chen JH, Hales CN, & Ozanne SE (2007) DNA damage, cellular senescence and organismal ageing:
causal or correlative? *Nucleic Acids Res* 35(22):7417-7428.

15.   Chapman J, Fielder E, & Passos JF (2019) Mitochondrial dysfunction and cell senescence:
deciphering a complex relationship. *FEBS Lett* 593(13):1566-1579.

16.   Prata L, Ovsyannikova IG, Tchkonia T, & Kirkland JL (2018) Senescent cell clearance by the immune
system: Emerging therapeutic opportunities. *Semin Immunol* 40:101275.

17.   Passos JF, Miwa S, & von Zglinicki T (2013) Measuring reactive oxygen species in senescent cells.
Methods Mol Biol 965:253-263.

18.   Musi N*, et al.* (2018) Tau protein aggregation is associated with cellular senescence in the brain.
Aging Cell 17(6):e12840.

19.   Ogrodnik M*, et al.* (2019) Obesity-Induced Cellular Senescence Drives Anxiety and Impairs
Neurogenesis. *Cell Metab* 29(5):1233.

20.   Yousefzadeh MJ*, et al.* (2018) Fisetin is a senotherapeutic that extends health and lifespan.
EBioMedicine 36:18-28.

21.   Xu M*, et al.* (2018) Senolytics improve physical function and increase lifespan in old age. *Nat Med*
24(8):1246-1256.

22.   Baker DJ*, et al.* (2011) Clearance of p16Ink4a-positive senescent cells delays ageing-associated
disorders. *Nature* 479(7372):232-236.

23.   Bussian TJ*, et al.* (2018) Clearance of senescent glial cells prevents tau-dependent pathology and
cognitive decline. *Nature* 562(7728):578-582.

24. Zhang P*, et al.* (2019) Senolytic therapy alleviates Abeta-associated oligodendrocyte progenitor
cell senescence and cognitive deficits in an Alzheimer's disease model. *Nat Neurosci* 22(5):719-
728.
25. Khatoon S, Agarwal NB, Samim M, & Alam O (2021) Neuroprotective Effect of Fisetin Through
Suppression of IL-1R/TLR Axis and Apoptosis in Pentylenetetrazole-Induced Kindling in Mice. *Front*
Neurol 12:689069.
26. Huang YH*, et al.* (2020) Interleukin-10 induces senescence of activated hepatic stellate cells via
STAT3-p53 pathway to attenuate liver fibrosis. *Cell Signal* 66:109445.
27. Maciel-Baron LA*, et al.* (2016) Senescence associated secretory phenotype profile from primary
lung mice fibroblasts depends on the senescence induction stimuli. *Age (Dordr)* 38(1):26.
28. Ou MY, Zhang H, Tan PC, Zhou SB, & Li QF (2022) Adipose tissue aging: mechanisms and
therapeutic implications. *Cell Death Dis* 13(4):300.
29. Hudgins AD*, et al.* (2018) Age- and Tissue-Specific Expression of Senescence Biomarkers in Mice.
Front Genet 9:59.
30. Cannon B & Nedergaard J (2004) Brown adipose tissue: function and physiological significance.
Physiol Rev 84(1):277-359.
31. Kalinovich AV, de Jong JM, Cannon B, & Nedergaard J (2017) UCP1 in adipose tissues: two steps
to full browning. *Biochimie* 134:127-137.
32. Gui Y, Silha JV, & Murphy LJ (2004) Sexual dimorphism and regulation of resistin, adiponectin, and
leptin expression in the mouse. *Obes Res* 12(9):1481-1491.
33. Isobe T*, et al.* (2005) Influence of gender, age and renal function on plasma adiponectin level: the
Tanno and Sobetsu study. *Eur J Endocrinol* 153(1):91-98.
34. Azzu V & Valencak TG (2017) Energy Metabolism and Ageing in the Mouse: A Mini-Review.
Gerontology 63(4):327-336.
35. Westbrook R, Bonkowski MS, Strader AD, & Bartke A (2009) Alterations in oxygen consumption,
respiratory quotient, and heat production in long-lived GHRKO and Ames dwarf mice, and short-
lived bGH transgenic mice. *J Gerontol A Biol Sci Med Sci* 64(4):443-451.
36. Prentice RL*, et al.* (2013) An exploratory study of respiratory quotient calibration and association
with postmenopausal breast cancer. *Cancer Epidemiol Biomarkers Prev* 22(12):2374-2383.
37. Baumann CW, Kwak D, & Thompson LV (2019) Sex-specific components of frailty in C57BL/6 mice.
Aging (Albany NY) 11(14):5206-5214.
38. Reshetnikov VV*, et al.* (2020) Genes associated with cognitive performance in the Morris water
maze: an RNA-seq study. *Sci Rep* 10(1):22078.
39. Lee B & Shao J (2014) Adiponectin and energy homeostasis. *Rev Endocr Metab Disord* 15(2):149-
156.
40. Tchkonia T*, et al.* (2010) Fat tissue, aging, and cellular senescence. *Aging Cell* 9(5):667-684.
41. Raffaele M*, et al.* (2021) Mild exacerbation of obesity- and age-dependent liver disease
progression by senolytic cocktail dasatinib + quercetin. *Cell Commun Signal* 19(1):44.
42. Grosse L*, et al.* (2020) Defined p16(High) Senescent Cell Types Are Indispensable for Mouse
Healthspan. *Cell Metab* 32(1):87-99 e86.
43. Hagg S & Jylhava J (2021) Sex differences in biological aging with a focus on human studies. *Elife*
10.
44. Roy B & Palaniyandi SS (2021) Tissue-specific role and associated downstream signaling pathways
of adiponectin. *Cell Biosci* 11(1):77.
45. Bloemer J*, et al.* (2018) Role of Adiponectin in Central Nervous System Disorders. *Neural Plast*
2018:4593530.
46. Rastegar S*, et al.* (2019) Expression of adiponectin receptors in the brain of adult zebrafish and
mouse: Links with neurogenic niches and brain repair. *J Comp Neurol* 527(14):2317-2333.

47. Baumgart M*, et al.* (2015) Summary of the evidence on modifiable risk factors for cognitive decline
and dementia: A population-based perspective. *Alzheimers Dement* 11(6):718-726.
48. Hascup ER*, et al.* (2019) Diet-induced insulin resistance elevates hippocampal glutamate as well
as VGLUT1 and GFAP expression in AbetaPP/PS1 mice. *J Neurochem* 148(2):219-237.
49. Fang Y*, et al.* (2020) Lifespan of long-lived growth hormone receptor knockout mice was not
normalized by housing at 30 degrees C since weaning. *Aging Cell* 19(5):e13123.
50. Palmer AK*, et al.* (2015) Cellular Senescence in Type 2 Diabetes: A Therapeutic Opportunity.
Diabetes 64(7):2289-2298.
51. Saccon TD*, et al.* (2021) Senolytic Combination of Dasatinib and Quercetin Alleviates Intestinal
Senescence and Inflammation and Modulates the Gut Microbiome in Aged Mice. *J Gerontol A Biol*
Sci Med Sci 76(11):1895-1905.
52. Roos CM*, et al.* (2016) Chronic senolytic treatment alleviates established vasomotor dysfunction
in aged or atherosclerotic mice. *Aging Cell* 15(5):973-977.
53. Dalterio S & Bartke A (1979) Perinatal exposure to cannabinoids alters male reproductive function
in mice. *Science* 205(4413):1420-1422.
54. Hascup KN, Findley CA, Sime LN, & Hascup ER (2020) Hippocampal alterations in glutamatergic
signaling during amyloid progression in AbetaPP/PS1 mice. *Sci Rep* 10(1):14503.

Figure legends

Figure 1: Senotherapeutic treatment altered the SASP profile in C57BL/6 mice. (a) Experimental paradigm. (b-g) Gene expressions in gonadal fat depot. Data are represented as means ± *SEM* (*n* = 8–12). A two-way ANOVA was used to determine P-values for the categorial variables (S = Sex and T = Treatment) and their interaction (S x T), which are shown for each graph.   *$p \leq 0.05$, **$p \leq 0.01$, ***$p \leq 0.001$ based on a two-tailed Student's *t* test.

Figure 2: D+Q treatment increased abdominal WAT in female mice while Fisetin treatment improved glucose metabolism and increased plasma adiponectin in male mice. Mouse body weight (a-b, BW) along with percentage of total (c) and different adipose tissue depots present in relation to BW (d-g). Gene expression of ACC in gonadal WAT (h) as well as UCP1 and PPARγ in interscapular BAT (i-j) depots. Glucose tolerance test (GTT), insulin tolerance test (ITT), and their respective area under the curve (AUC) after five treatments (k-p). Plasma insulin and adiponectin concentrations (q-r) taken at time of euthanization after 10 treatments. Data are means ± SEM (n = 8–20). A two-way ANOVA was used to determine P-values for the categorial variables (S = Sex and T = Treatment) and their interaction (S x T), which are shown for each bar graph. *$p \leq 0.05$, **$p \leq 0.01$ based on a two-tailed Student's t test.

Figure 3: Fisetin treatment enhanced energy metabolism in male mice, while D+Q treatment decreased energy metabolism in female mice. Twenty-four-hour male and female oxygen consumption (a-c; VO_2), energy expenditure (d-f; EE), and respiratory quotient (g-i; RQ) along with their corresponding AUC after 9 treatments. Data are presented as means ± SEM (n = 16-20). A two-way ANOVA was used to determine P-

values for the categorial variables (S = Sex and T = Treatment) and their interaction (S

x T), which are shown for each bar graph. **$p \leq 0.01$ based on a two-tailed Student's t

test.

Figure 4: Fisetin treatment improved spatial learning in male mice, while D+Q

treatment impaired object recognition memory recall in female mice. Assessment

of cognition and memory recall was determined by MWM and NOR. MWM path

efficiency (a-c) and corrected integrated path length (d-f; CIPL) in male and female

C57BL/6 mice during 5 training sessions. Number of platform crosses during the

delayed probe challenge (g). Retention index of the familiar object after a delayed

novelty introduction (h). Hippocampal, liver, and gonadal WAT mRNA expression levels

of Glut4 (i-k) and AdipoR1 (l-n) were measured at time of euthanization after 10

treatments. Data are presented as means ± SEM (n = 16-20). A two-way ANOVA was

used to determine P-values for the categorial variables (S = Sex and T = Treatment)

and their interaction (S x T), which are shown for each bar graph. *$p \leq 0.05$, **$p \leq 0.01$

based on a two-tailed Student's t test.

Supplemental Information: Figure 1: Effects of senotherapeutic treatment on the

hippocampal SASP profile in C57BL/6 mice. mRNA expression levels of

hippocampal SASP markers (a-f) were measured at time of euthanization after 10

treatments. Data are presented as means ± SEM (n = 8-12). A two-way ANOVA was

used to determine P-values for the categorial variables (S = Sex and T = Treatment)

and their interaction (S x T), which are shown for each bar graph.   *$p \leq 0.05$, **$p \leq$

0.01, ***$p<0.001$ based on a two-tailed Student's t test.

Supplemental Information Figure 2: Senotherapeutic treatment altered plasma

cytokine concentrations in C57BL/6 mice. Plasma concentrations of TNFα, MCP1,

IL-6, and IL-10 (a-d) from time of euthanization after 10 senotherapeutic treatments.

Data are represented as means ± SEM (n = 8). A two-way ANOVA was used to

determine P-values for the categorial variables (S = Sex and T = Treatment) and their

interaction (S x T), which are shown for each bar graph. $*p \leq 0.05$, $**p \leq 0.01$,

$***p < 0.001$ based on a two-tailed Student's t test.

Supplemental Information Figure 3: Effects of senotherapeutic treatment on

physical performance in C57BL/6 mice. MWM swimming speed during the training

days and probe challenge (a-d) and forepaw grip strength (e). Data are presented as

means ± SEM (n = 16-20). A two-way ANOVA was used to determine P-values for the

categorial variables (S = Sex and T = Treatment) and their interaction (S x T), which are

shown for each bar graph.

Supplemental Information Figure 4: Hippocampal mRNA expression of genes

involved with synaptic plasticity were unaltered after senotherapeutic treatment.

Hippocampal mRNA expression from time of euthanization after 10 senotherapeutic

treatments in C57BL/6 mice. Data are represented as means ± SEM (n = 16-20). A

two-way ANOVA was used to determine P-values for the categorial variables (S = Sex

and T = Treatment) and their interaction (S x T), which are shown for each bar graph.

$**p \leq 0.01$ based on a two-tailed Student's t test.

Supplemental Information Figure 5: Effects of aging on physical, metabolic, and

cognitive parameters. Nontreated four-month-old C57BL/6 mice were compared to

sex-matched control treated mice. Data are represented as means ± SEM (n = 16-20).

A two-way ANOVA was used to determine P-values for the categorial variables (S =

Sex and A = Age) and their interaction (S x A), which are shown for each bar graph.

$*p \leq 0.05$, $**p \leq 0.01$, $***p < 0.001$ based on a two-tailed Student's t test.

Supplemental Information: Table 1: A list of forward and reverse primers used in this

study.

Figure 1

Figure 2

Figure 3

 Figure 4

Supplemental Information: Figure 1

 Supplemental Information: Figure 2

Supplemental Information: Figure 3

Supplemental Information: Figure 4

 Supplemental Information: Figure 5

Supplemental Information: Table 1

Gene	Protein	Forward	Reverse
CDKN2A	p16$^{Ink4\alpha}$	TACCCCGATTCAGGTGAT	TTGAGCAGAAGAGCTGCTACGT
CDKN1A	p21^{Cip1}	CCTGGTGATGTCCGACCTG	CCATGAGCGCATCGCAATC
TNFα	TNFα	GCCTCTTCTCATTCCTGCTTG	CTGATGAGAGGGAGGCCATT
CCL2	MCP1	CCACTCACCTGCTGCTACTCAT	TGGTGATCCTCTTGTAGCTCTCC
IL6	IL-6	TAGTCCTTCCTACCCCAATTTCC	TTGGTCCTTAGCCACTCCTTC
IL10	IL-10	GCTCTTACTGACTGGCATGAG	CGCAGCTCTAGGAGCATGTG
ACACA	ACC	ATGGGCGGAATGGTCTCTTTC	TGGGGACCTTGTCTTCATCAT
UCP1	UCP1	AGGCTTCCAGTACCATTAGGT	CTGAGTGAGGCAAAGCTGATTT
PPARG	PPARγ	ACCCCCTGCTCCAGGAGAT	TGCAATCAATAGAAGGAACACGT
NPTX	Nptx	ACACCATGAAAGTGGGAGGTAACT	ACTGGTCTTGTCCTTACTTCCGGA
SNAP25	Snap25b	CTCATCGAGTGGGGCAACA	GTGATGCCATTTGCCATCGTT
Syt7	Syt7	ACTCCATCATCGTGAACATCATC	TATGTCGAAGGCGAAAGAC
Slc8a2	Na$^+$ / Ca^{++} Exchanger, Member 2	AACAGCACCTTCTACGTGG	GTCCTTCTGCGTCTCCC
GRIN2A	GluN2A	ACGTGACAGAACGCGAACTT	TCAGTGCGGTTCATCAATAACG
GRIN2B	GluN2B	CAGCAAAGCTCGTTCCCAAAA	GTCAGTCTCGTTCATGGCTAC
CREB1	Creb1b	AGCAGCTCATGCAACATCATC	AGTCCTTACAGGAAGACTGAACT
SLC16A1	MCT1	TGTTAGTCGGAGCCTTCATTTC	CACTGGTCGTTGCACTGAATA
GRIA1	Gria1 Flip	ACACCATGAAAGTGGGAGGTAACT	ACTGGTCTTGTCCTTACTTCCGGA
GRIA1	Gria1 Flop	GTCCGCCCTGAGAAATCCA	GCACTCGCCCTTGTCGTA
SLC2A4	Glut4	ACACTGGTCCTAGCTGTATTCT	CCAGCCACGTTGCATTGTA
ADIPOR1	AdipoR1	ACGTTGGAGAGTCATCCCGTAT	CTCTGTGTGGATGCGGAAGAT
B2M	B2M	TTCTGGTGCTTGTCTCACTGA	CAGTATGTTCGGCTTCCCATTC