Title

- **Full title:** Developmental dynamics of the neural crest-mesenchymal axis in creating the thymic microenvironment.
- **Short title:** Developmental mesenchymal dynamics in the thymus.

Authors

Adam E. Handel[1,2]†, Stanley Cheuk[1,3]†, Fatima Dhalla[1], Stefano Maio[1], Tania Hübscher[4], Ioanna Rota[1], Mary E. Deadman[1], Olov Ekwall[3,5], Matthias Lütolf[4], Kenneth Weinberg[6], Georg Holländer[1,7,8]*

Affiliations

1. Department of Paediatrics and the Weatherall Institute of Molecular Medicine, University of Oxford, Oxford, United Kingdom.

2. Nuffield Department of Clinical Neurosciences, University of Oxford, Oxford, United Kingdom.

3. Department of Rheumatology and Inflammation Research, University of Gothenburg, Gothenburg, Sweden.

4. Laboratory of Stem Cell Bioengineering, Swiss Federal Institute of Technology in Lausanne, Lausanne, Switzerland.

5. Department of Pediatrics, University of Gothenburg, Gothenburg, Sweden.

6. Division of Stem Cell Transplantation and Regenerative Medicine Department of Pediatrics, Stanford University, Stanford, USA.

7. Paediatric Immunology, Department of Biomedicine, University of Basel and University Children's Hospital Basel, Basel, Switzerland.
8. Department of Biosystems Science and Engineering, ETH Zurich, Basel, Switzerland.

* Corresponding author: georg.hollander@paediatrics.ox.ac.uk
† These authors contributed equally.

Abstract

The thymic stroma is composed of epithelial and non-epithelial cells that collectively provide separate microenvironments controlling the homing of blood-born precursors to the tissue, and their subsequent differentiation to functionally mature and correctly selected T cells. While thymic epithelial cells are well characterized for their role in thymopoiesis, a comparably comprehensive analysis of the non-epithelial thymic stroma is lacking. Here we explore at single cell resolution the complex composition and dynamic changes that occur over time in the non-epithelial stromal compartment. We detail across different developmental stages in human and mouse thymus, and in an experimental model of Di George syndrome, the most common form of human thymic hypoplasia, the separate transcriptomes of mouse mesothelium, fibroblasts, neural crest cells, endothelial and vascular mural cells. The detected gene expression signatures identify novel stromal subtypes and relate their individual molecular profiles to separate differentiation trajectories and functions. Specifically, we demonstrate an abundance and unprecedented heterogeneity of diverse fibroblast subtypes that emerge at discrete developmental stages and vary in their expression of key regulatory signalling circuits and components of the extracellular matrix. Taken together, these findings highlight the dynamic complexity of the non-epithelial thymus stroma and link the cells' specific gene expression profiles to separate instructive roles essential for normal thymus organogenesis and tissue maintenance.

Teaser

Single cell profiling of thymic stroma identifies a dynamic contribution from neural crest cells to the thymic mesenchyme.

Introduction

Thymic T cell lineage commitment, development, maturation, and repertoire selection are instructed by a stromal scaffold that includes thymic epithelial cells (TEC) (*1*), endothelial cells (*2*) and mesenchymal cells (*3, 4*). The TEC compartment is both phenotypically and transcriptionally well characterized providing at single cell resolution a detailed account of the cells' developmental dynamics and functions (*5, 6*). In addition, the thymus microenvironment is also composed of stromal cells of mesenchymal origin, including fibroblasts, endothelial cells and vascular mural cells. Derived primarily from either mesoderm or ectodermal neural crest cells, these thymic mesenchymal cells interact with TEC and thus create unique cellular niches that control thymopoiesis. This critical function of mesenchymal cells is accomplished via the production of extracellular matrix components, morphogens and key growth factors (*3, 4, 7*). Hence, the thymic mesenchyme is indispensable for the organ's correct formation and function (*1, 7, 8*).

Fibroblasts constitute the largest component of the non-epithelial thymus stroma. (NETS) Though first described as distinct cell type over 150 years ago, the specific contributions of fibroblasts to organ formation, maintenance and function have only recently begun to be unraveled (*8*). Utilizing single-cell genomic technologies for the comparison of diverse tissues, fibroblasts were noted to display a significant heterogeneity with both cross-organ communalities and tissue-specific differences (*9*). Likewise, endothelial cells and vascular mural cells display organotypic features that have only recently been appreciated when resolving the cells' distinct transcriptomes at single-cell resolution (*10*). In addition to their essential role in providing oxygen, nutrients, cells and other cargo to tissues, blood vessels also express in a context-specific fashion diverse transcriptomic profiles that include sets of growth factors inducing, specifying, patterning, and guiding organ formation and homeostasis (*11*). A third stromal component of non-epithelial origin are neural crest cells which enter the anlage as a migratory population as early as embryonic day 12 where they differentiate into distinct cell types, including vasculature-associated pericytes juxtaposed between endothelia and the other components of the stromal scaffold (*12*).

A detailed phenotypic, transcriptomic, and functional genomic description of the diverse population of non-epithelial thymus stromal cells is to date still wanting. We have therefore employed flow cytometry and single cell multiomics technologies to detail the complexity and developmental dynamics of thymic mesenchymal cells in both mouse and human tissue. Our results highlight a previously unappreciated heterogeneity among cells belonging to the NETS under physiological conditions and identify distinct yet selective defects of these cells in a genetic mouse model of the 22q11 deletion syndrome, the most common human condition associated with congenital thymus hypoplasia.

Results

Single cell sequencing reveals high levels of complexity within the thymic mesenchyme

We first sought to delineate both the frequency and diversity of NETS cells (phenotypically defined as Ter119⁻CD45⁻EpCAM⁻) in the thymus of 4-week-old mice. These cells accounted for approximately half of the total thymus stroma cellularity and distinct subpopulations were identified using the differential expression of glutamyl aminopeptidase Ly51, glycoprotein podoplanin (gp38) and dipeptidyl peptidase-4 (DPP4, CD26) (Fig. 1a and Fig. S1a) (*3, 13*). The Ly51^{hi}gp38⁻ phenotype identified neural crest-derived pericytes which surround blood vessels adjacent to endothelial cells (*4, 12*). The gp38 positive stromal cells expressed a reduced level of Ly51 and could be further differentiated into separate subpopulations based on their CD26 expression: gp38⁺CD26⁺ cells localized to the thymus capsule whereas gp38⁺CD26⁻ were enriched in the medulla (Fig. 1b) (*13*).

To delineate the heterogeneity of the thymic mesenchyme in an unbiased fashion and independent of a limited number of phenotypic markers, we next generated transcriptomic libraries from 5,878 single Ter119⁻CD45⁻EpCAM⁻ cells isolated from 4-week old thymi (Extended Data Fig. 1a). We identified 12 distinct cell subtypes based on their separate gene expression profiles (Fig. 1c, d, Fig. S1b). (For clarity, we refer to transcriptionally defined stromal cell clusters as subtypes whereas the terms populations and subpopulations specify cells that have been defined by cytometry). A Uniform Manifold Approximation and Projection (UMAP) analysis of non-TEC stroma identified four separate cell clusters of different sizes. The two largest comprised several individual cell subtypes that were transcriptomically defined as either capsular or medullary fibroblasts (Fig. 1c and Fig. S1c i and ii) (*13*). The capsular cluster closely resembling the gene expression profile of capsular fibroblasts consisted of four subsets that separated from mesothelial cells defined – *inter alia* - by their expression of *Msln* and *Upk3b,* encoding the glycosylphosphatidylinositol-anchored cell-surface adhesion protein mesothelin and the membrane integral protein Uroplakin, respectively (*14*). Within the capsular fibroblast clusters, subtype 1 (designated CapFb1) was characterized by the high expression of *Svep1*, *Sfrp1* and *Dpep1*, which encode a multidomain cellular adhesion molecule (*15*), the secreted frizzled-related protein 1 modulating stromal to epithelial signaling via Wnt inhibition (*16*), and a membrane-bound dipeptidase involved in the metabolism of glutathione and other similar proteins (*17*). The capsular subtypes 2 (CapFb2) and 4 (CapFb4) were characterized by the expression of *Pi16*, *Mfap5* and *Fstl1*, which encode a peptidase inhibitor of largely unknown function, the microfibrillar-associated protein 5 related to extracellular matrix remodeling and inflammation (*18*), and the secreted extracellular glycoprotein follistatin-like 1 . CapFb4 also highly expressed *Timp2* encoding the tissue inhibitor of metalloproteinase 2 relevant for tissue remodeling (*19*), *Anxa3* translating into the membrane-associated Annexin 3 protein activating the epithelial-to-mesenchymal transition (EMT) program and Wnt signaling pathway (*20*), and *Fbn1* encoding Fibrillin 1, a major component of extracellular microfibrils. The CapFb3 subtype typically expressed *Igf1* encoding Insulin-growth factor 1 regulating tissue homeostasis via cell proliferation, differentiation, maturation, and survival (*21*), *Gdf10* translating into the transforming growth factor-β (TGF-β) superfamily member growth differentiation factor-10 (GDF10) (*22*), and *Olfml3* encoding the secreted glycoprotein olfactomedin-like 3 which has matrix-related functions central to embryonic development (*23*).

The second UMAP cluster incorporated two distinct medullary fibroblast subtypes, pericytes and vascular smooth muscle cells. The medullary fibroblast subtypes 1 (MedFb1) and 2 (MedFb2) displayed similar gene expression profiles although transcripts for the out-at-first protein

(encoded by *Oaf*) and the alpha 1 chain of collagen XV (*Col15a1*) were detected at higher levels
in MedFb1 while transcripts for extracellular superoxide dismutase 3 (*Sod3*) were particularly
evident in MedFb2. Both subtypes also comprised transcripts for IL-33 and Cxcl16, which are
important for dendritic cell activation and NKT cell migration, respectively (*24, 25*). Transcripts
related to antigen processing and presentation were enriched in MedFb2 (Fig. S1b). However,
contrary to a recent observation (*13*), tissue restricted antigens were not generally more frequent
in medullary fibroblasts when compared to other thymic NETS (Fig. S1e).
156
Pericytes were identified by their characteristic expression of *Kcnj8* encoding member 8 of the J
subfamily of the potassium inwardly rectifying channels, which form part of the ATP/ADP-
binding potassium channel of these cells (*26*). Vascular smooth muscle cells (VSM) were
characterized by the expression of contractile elements, including *Acta2*, *Tpm1* and *Myh11*,
whereas endothelial cells displayed a high number of transcripts for *Gpihbp1* and *Pecam1* which
encode the glycosylphosphatidylinositol anchored high density lipoprotein binding protein 1 and
the intercellular junction protein platelet and endothelial cell adhesion molecule (PCAM, aka
CD31), respectively. Neural crest-derived cells (NCC) were characterized by their expression of
Foxd3 and *Sox10* which are critical for the cells' specification and development (*27, 28*). Finally,
actively proliferating cells were identified by the expression of different cell cycle-related genes
including *Mki67* encoding the nuclear protein Ki67.
168
We determined the gene expression profiles of the 4 FACS-defined non-TEC thymic stromal
subpopulations (Fig. S1a) and deconvoluted the individual transcriptomes by projection onto the
single cell UMAP data (Fig. 1e, Table 1). Stroma cells expressing CD31 identified the cluster
defined as endothelial cells and the Ly51$^+$Gp38$^-$ subpopulation represented the pericyte and
vascular smooth muscle cell clusters (Fig. 1e, f). The gp38$^+$CD26$^+$ subpopulation included all of
the 4 capsular fibroblasts subtypes, whereas the gp38$^+$CD26$^-$ subpopulation was mainly enriched
for the medullary fibroblast subtypes but also included CapFb1 cells (Fig. 1e, f). We showed that
this fibroblast heterogeneity patterned the thymic extracellular matrix by staining for two key
extracellular matrix molecules (type I collagens and fibronectin), which were most highly
expressed in capsular fibroblasts (Fig. 1g).
Hence, the single cell RNA-seq based identification of thymic stromal cells unmasked a
previously unrecognized heterogeneity of individual subsets among gp38$^+$ NETS which could not
be identified by conventional flowcytometry-based phenotyping.
182
183 **Thymic organogenesis is characterised by dynamic mesenchymal changes**
184
The thymus undergoes significant micro-architectural changes during organogenesis, including
the compartmentalisation into distinct cortical and medullary domains and the formation of a
complex vascular network (*29*). We therefore investigated how these morphological changes
paralleled compositional alterations of the mesenchymal stroma (Fig. 2a-b). At embryonic day
(E)12.5, NETS accounted for more than 90% of all CD45$^-$ thymic cells with Ly51$^+$gp38$^-$ cells
being by far the most dominant subpopulation. The frequency of TEC gradually increased parallel
to thymus growth and reached a relative maximum at E16.5 when epithelia represented 60% of
the thymic stroma. Earlier during thymus organogenesis, the non-TEC stroma lacked the
heterogeneity observed at E16.5 and thereafter. For example, gp38$^+$CD26$^-$ and gp38$^+$CD26$^-$
fibroblasts dominated the stromal compartment at both E12.5 and E13.5, endothelial cells were
only discovered at E13.5 and Ly51$^+$gp38$^-$ pericytes were not detected prior to E16.5 (Fig. 2a, Fig.
S2a). The subpopulation of gp38$^+$CD26$^+$ capsular fibroblasts was identified as early as E13.5 and
increased in frequency thereafter (Fig. 2a).
198

We next used single cell RNA-seq to detail changes in the heterogeneity of individual NETS
subtypes and to determine the cells' developmental trajectories. We generated libraries on a total
of 36,208 single stromal cells isolated from embryonic (E12.5, 13.5, 16.5), new-born and young
adult thymus tissue, which collectively reiterated the clusters observed in the thymus of 4-week-
old mice and provided sufficient resolution to identify additional heterogeneity (Fig. 2b-d, Table
1). Complex dynamic changes in the frequency of individual subtypes occurred over time
between the early developmental stages and the completion of a mature thymus
microenvironment. For example, CapFb1a and CapFb2b appeared early but their frequencies
gradually decreased during organogenesis whereas all of the medullary fibroblasts (with the
notable exception of MedFb1a) increased parallel to the emergence of mTEC (*30*). NCC were
largely absent after E16.5 but other NETS subtypes remained either mainly unchanged or
displayed a bi-model variation in frequency between E12.5 and 4 weeks of age (Fig. 2c i and ii).
This finding is in agreement with lineage tracing studies demonstrating the cells' developmental
potential to differentiate into VSM and pericytes.(*12, 31*) Thus, single cell RNA-seq revealed
complex and dynamic changes in the relative number of individual NETS subtypes that would be
captured incompletely by classical cell surface phenotyping, such as CD26 (Fig. S2b).
215
We leveraged the splicing information obtained from single cell transcriptomes to determine the
developmental trajectories of individual NETS subtypes. This analysis identified the CapFb1a and
CapFb2b subtypes as the principal precursors for other capsular fibroblasts (Fig. 2d) and
suggested MedFb1a to serve as a precursor for other fibroblast subtypes in the emerging medulla
(*30*). This analysis also recognized CapFb3 fibroblasts as intermediates between mesothelial cells
and other fibroblast subtypes, a finding consistent with the concept that fibroblasts can be derived
from mesothelial cells (*32*). However, CapFb3 were distinct from mesothelia as they lacked the
expression of *Msln* and *Upk3b* (Fig. 1d) (*14*).
224
The single cell transcriptome data was also used to infer gene regulatory network activities of
individual NETS subtypes (Fig. 2e). A transcription factor motif analysis of these gene regulatory
networks was executed to identify potential cell type-specific transcription factors (*33*). In
keeping with their proposed differentiation from mesothelial cells, CapFb3 fibroblasts expressed
gene regulatory networks controlled by the transcription factors *Hoxa5* and *Wt1*, which typically
are active in mesothelial cells (Fig. 2e, Fig. S2c) (*14*). CapFb4 were highly enriched for a *Creb5*-
controlled gene regulatory network that has previously been identify to modulates the
differentiation of fibroblasts to myofibroblasts (*34*) and to control age-related thymic fibrosis (*35*)
(Fig. S2d). MedFb2b expressed *Irf7* encoding the Interferon-regulatory factor 7 (IRF7), a master
regulator of type I IFN secretion that interacts with Smad3 to regulate TGF-β signalling for
collagen production (Fig. 2e, Fig. S2e) (*36*).
236
We assessed the expression of canonical Wnt signalling transcripts and growth factors known to
be important in thymic stromal interactions with thymocytes (Fig. S3) (*37*). Several Wnt ligands
displayed distinct expression patterns among cells of the NETS. For example, *Wnt4* transcripts
were detected in mesothelium, *Wnt5a* in CapFb2b, CapFb3 and CapFb4, *Wnt6* in NCC, and
Wnt10b in CapFb4. Wnt modulators were also highly expressed in particular non-epithelial
stroma cells, including *Rspo1* in mesothelium and *Sfrp5* in NCC. Several cell subtypes of the
NETS acted as prominent sources of key growth factors, including *Bmp4* transcribed in CapFb1c,
Bmp7 in CapFb4 and mesothelium, *Fgf10* in CapFb1a, CapFb1b and CapFb1c, and *Tgfb1* in
endothelial cells. In keeping with their role in regulating the extracellular matrix, fibroblast
subtypes showed high expression of key extracellular matrix transcripts, with collagens (*e.g.*
Col1a2, *Col3a1* and *Col14a1*) primarily expressed in capsular fibroblasts, and laminins (*Lama2*
and *Lama4*) in a mixture of capsular (CapFb1a, CapFb1b and CapFb1c) and medullary fibroblasts

(MedFb1b and MedFb2b). This distinctive expression of growth and differentiation factors, and
components of the extracellular matrix demonstrated that heterogeneity within the NETS
compartment determined modularity in the expression of key molecules, thus implicating
different developmental and functional niches.
253
Ligand-receptor pairing analysis identifies interactions between neural crest-derived
mesenchyme and endothelial cells
256
Given that NCCs are known to differentiate into perivascular cell types, we aimed to uncover the
ligand-receptor signaling and the subsequent transcriptomic networks that control the
differentiation of NCCs into pericytes and VSM and pericytes (*12, 31*). To this end, NicheNet
identified intercellular ligand-receptor interactions associated with cell type-specific transitions
across early (E12.5 and E13.5) to later stages (E16.5) in embryonic thymus formation (*38*). This
analysis demonstrated that ligands expressed by endothelial cells, including the adhesive and
multimeric glycoprotein von Willebrand factor (vWF) and Transforming Growth Factor Beta 1
(TGFB1), influenced gene expression in thymic NCC (Fig. 3a, b). Conversely, heterotypic
interactions between junctional adhesion molecule-3 (*Jam3)* produced by NCC and its receptor
Jam2 on endothelial cells identified a candidate ligand-receptor pair that orchestrated the changes
in the gene expression profile of embryonic endothelial cells (Fig. 3c). Together, these inferred
ligand-receptor interactions suggested that reciprocal cellular relationships between vascular
structures and NCC shape the perivascular thymic stroma during embryogenesis.
270
Reduced cellularity and complexity of mesenchymal stroma are features of an experimental
22q.11.2 Deletion Syndrome model
273
A spontaneously occurring heterozygous deletions of 1.5Mb to 3Mb size following recombination
between four blocks of low copy repeats within 22q11.2 account for the loss of up to 106 genes
(*39*). This mutation constitutes the most common molecular etiology of the 22q11.2 deletion
syndrome (DS; previously referred to as DiGeorge Syndrome (DGS)) which manifests clinically
as a range of features that include either athymia resulting in T cell deficiency or thymus
hypoplasia compromising immunological fitness (*39*). Regions of mouse chromosome 16 are
syntenic to the human 22q11.2 (*40*) and include *Tbx1,* encoding a T-box transcription factor and
Crkl, encoding an adapter protein implicated in fibroblast growth factor and focal cell adhesion
signaling (*39*). Compound haploinsufficiency of *Tbx1* and *Crkl* in mice results in typical
hallmarks of 22q11.2DS, including thymic hypoplasia (*41*).
284
The abnormal migration of cephalic NCC has been identified as a possible cause for the
pharyngeal patterning defects observed in 22q11.2DS which is recapitulated in mice compound
heterozygous for *Tbx1* and *Crkl* (designated *Tbx1*[+/-]*Crkl*[+/-]). Gene products of these two loci have
been alleged to interact in a dosage-sensitive fashion (*41*). *Tbx1* expression in NETS was
exclusively confined to E12.5 and detected in a subset of cortical fibroblast subtypes, especially
CapFb4, and proliferating cells. Yet, *Crkl* transcripts were mainly detected in CapFb1 and
CapFb4 subtypes early during thymus development but could also be identified in a small fraction
of these and other NETS later in development (Fig. S4a-c).
293
The thymi of mice compound heterozygous for a loss of *Tbx1* and *Crkl* were hypoplastic and
revealed already at E13.5 significantly fewer haematopoietic, epithelial and mesenchymal cells
than their wild type controls, which was not the case for either single *Tbx1* or *Crkl* mutants (Fig.
4a-c and Fig. S4d). At birth, hematopoietic cells and all phenotypically identified major NETS
subpopulations were reduced in mutant mice and remained diminished in 4-week-old mice with

the notable exception of gp38$^+$CD26$^+$ capsular fibroblasts. In contrast, the cellularity of TEC and
endothelial cells varied over time and were not uniformly reduced in mutant mice at these times
(Fig. 4b, c; Fig. S4e). Thus, several NETS subpopulations were consistently reduced in mice
heterozygous for *Tbx1* and *Crkl*.
303
We next compared the transcriptome of individual epithelial and NETS cells isolated from new-
born *Tbx1$^{+/-}$Crkl$^{+/-}$* mice (Fig. 5a and Fig. S5a; epithelial and non-epithelial stromal cells were
annotated as previously published (*5*) and shown in Figure 2, respectively). *Tbx1* and *Crkl*
compound heterozygosity substantially changed the composition of the thymus stroma resulting
in a reduction of 8 of the 18 individual subtypes in *Tbx1$^{+/-}$Crkl$^{+/-}$* mice. Specifically and in
contrast to the results obtained by flow cytometry, the relative cellularity of several capsular and
medullary fibroblast subtypes together with that of pericytes and VSM was lessened (Fig. 5b). In
addition, the frequencies of mature cortical (mcTEC) and intertypical TEC (itTEC) were reduced
in mutant mice whereas that of perinatal cortical TEC (pcTEC), post-Aire mTEC (pamTEC) and
neural TEC (nTEC) were enriched (Fig. 5c).
314
315   To assess the functional consequences of a compound heterozygous loss of *Crkl* and *Tbx1*, we
316   performed an enrichment analysis for differentially expressed gene sets (Fig. 5d, Fig S5b). This
317   analysis revealed that transcripts for gene products relevant for cell migration were reduced in
318   several capsular and medullary fibroblast as well as in pericytes. In VSM, fewer transcripts for
319   contractile elements (e.g. *Myl6* and *Tpm2*; Fig 5d) were observed.
320
321   **Compound *Tbx1* and *Crkl* heterozygosity is associated with accelerated aging of thymic**
322   **mesenchyme**
323
22q11DS patients display an accelerated thymic senescence (*42*), a process implicated to be
caused by an age-dependent chronic systemic inflammatory condition known as inflammaging
(*43*). To appraise the effects of inflammaging on NETS, we applied to our data an ageing score
computed from age-driven transcriptomic changes common across many tissues (*44*). In contrast
to the heterogeneous effect of ageing on TEC subsets (*5*), the ageing module score of non-TEC
stroma progressively increased from early embryonic stages to young adulthood (Fig. 6a). These
changes demonstrated a switch from an abundant expression of transcripts belonging to
biosynthetic pathways to gene products associated with angiogenesis and immunological
crosstalk (Fig. S6a).
333
Altered transcription factor network activity has been linked to the process of senescence (*44*).
Using the transcriptomes from NETS isolated from mice at different ages, a decrease in
transcripts related to gene networks controlled by Sox 4 and its close relative Sox11 were
observed. In parallel, gene networks controlled by IRF7 were gradually activated over time and
beyond what would be expected from their enrichment in MedFb2b subsets (Fig. 6b and Fig.
S2e). In contrast, *Fos* and *Fosb* controlled gene networks peaked at birth but were subsequently
weakened (Fig. S6b), a pattern previously noted in other tissues (*45*).
341
The ageing module score analysis was extended to include thymic stromal cells isolated from
Tbx1$^{+/-}$Crkl$^{+/-}$ mice at postnatal day 0. Accelerated aging (as discernible by an increased score)
was noted in mutant mice for the population of mesenchymal but not endothelial and epithelial
cells (Figure 6c). Within the mesenchymal compartment, accelerated aging was not uniform as an
increased score was observed in only 7 of the 19 distinct stromal subtypes including pericytes,
VSM and 2 cortical and medullary fibroblast subtypes (Figure 6d). Hence, these studies showed
age-related transcriptomic changes across distinct NETS subtypes of *Tbx1$^{+/-}$ Crkl$^{+/-}$* mice.

Although there was only a very small number of NCCs present at postnatal day 0 in either wild type or $Tbx1^{+/-}Crkl^{+/-}$ mice, NCCs migration and differentiation are known to be impaired in 22q.11.2 Deletion Syndrome (*46*). It is therefore possible that the alterations within the NETS compartment observed in $Tbx1^{+/-} Crkl^{+/-}$ mice may be driven by aberrant differentiation of NCCs into mesenchymal cells, particularly perivascular cell types (*31*).

Neural crest cells differentiate into perivascular cells in the human prenatal thymus

To extend the analysis of the thymus stroma to human tissue, we used single NETS nuclei to investigate their gene expression profiles and correlated these to chromatin accessibility. For this purpose, we employed a multiomics analysis that investigated 528 individual non-TEC thymic stroma nuclei isolated from two donors at 14- and 17-weeks post-conception (Fig. S7 and S8). This analysis identified seven distinct cell clusters, corresponding to NCCs (NCC-I and NCC-II), capsular fibroblasts, VSM, endothelial cells, medullary fibroblasts and pericytes. The frequency of cells in cluster 6 increased from 14- to 17-weeks post-conception, whereas the frequency of those in clusters 3 and 5 decreased within that time span (Fig. S9). We found the expected gradients in *PDGFRA* and *PDGFRB* expression across clusters composed of fibroblasts, pericytes and VSM (Fig. 7a) and *PECAM1* expression in endothelial cells (Fig. 7b).

 Given the dynamic changes in NCC frequency in the thymus over development and the changes observed in NCC-derived perivascular structures in $Tbx1^{+/-}Crkl^{+/-}$ thymi, we focused further on the chromatin and transcriptomic landscape of the NCC subclusters. NCC development is understood as a stepwise series of bifurcating cell fate decisions that lead to multiple cell identities and traits (*47*). Once specified in their fate, NCCs undergo an epithelial-to-mesenchymal transition and migrate throughout the embryo. NCC-II showed genome-wide high accessibility for sequences with SOX10 transcription factor binding motifs whereas the NCC-I displayed only an intermediate degree of accessibility for this motif (Fig. S10).

To further investigate the dynamics of SOX10 activity, we integrated gene expression profiles with chromatin accessibility in single cells to identify key enhancer-TSS interactions driving *SOX10* expression in human NCCs (Fig. 7c). We identified two sequences, which corresponded to two upstream orthologous enhancer elements (called U2 and U3) previously implicated in the control of *SOX10* expression in mouse NCCs and their progeny (*48*). Having identified that *SOX10* regulation was shared throughout the two NCC subclusters, we further examined heterogeneity amongst these NCCs to establish whether these could represent NCCs in different states of differentiation.

The comparative analysis of the gene expression profiles among the two NCC subclusters revealed an enrichment of gene pathways associated with cellular motility and vascular development for NCC-I (*e.g.* *BCL2*, *FGF13* and *RHOJ*; Fig. S11a). In contrast, NCC-II was enriched for gene pathways related to neuronal development and thus marked cells with gene expression profiles characteristic of bona fide NCC (*e.g.* *NES*, *NCAM2* and *GRID2*; Fig. S11a). This difference in gene expression profiles suggested that NCC-I may constitute a population of NCCs differentiating into mesenchymal and perivascular cell types. In support of this notion, NCC-II showed in comparison to NCC-I a significantly higher expression of *TFAP2A*, a key transcription factor in early NCC development, whereas *NR2F2*, a transcription factor involved in NCC migration, was most highly expressed in NCC-I (Fig. S11b) (*49*). Hence, the multiomics analysis of NES cells in human thymi captured the process of NCC differentiation into other cell types, a finding that could not be observed in mouse thymic stromal samples analyzed in this study.

Discussion

Stromal cells with separate functions emerge from all germ-layers during development to populate organs where they instruct the tissue's essential activities, for example via the differential production of extracellular matrix components, the release of growth and differentiation factors, and the creation of signalling niches that provide critical molecular cues (*50*). In addition to cross-organ communalities, stromal cells with seemingly identical phenotypes also display a heterogeneity both within and across tissues as revealed by dissimilarities in transcripts encoding pathway elements, transporters and cell-surface markers (*51*). Previous studies of the thymic stroma in both mice and humans could identify only a limited number of phenotypically distinct thymic mesenchyme subtypes (*13, 37, 52*) despite the cells' acknowledged roles as critical components in maintaining tissue structure and TEC function (*7*).

Employing gene expression profiles at single cell resolution, we now show an unprecedented heterogeneity among thymic mesenchymal cells and identify dynamic changes in the frequency of these cells across a large range of developmental stages. Notably, the observed diversity is not replicated using flow cytometry as separate since transcriptionally defined stromal subtypes display identical phenotypic features due to a limited number of suitable cell surface markers. This limitation has hindered a comprehensive understanding of how non-epithelial thymic stromal cells contribute to local tissue microenvironments which control discrete stages of intrathymic T cell differentiation.

Early in development, the thymus stroma is mostly composed of cells belonging to the NETS and, with the notable exception of NCC, continue to structure the scaffold also in adult mice where they contribute together with TEC to the non-haematopoietic stroma. We identify within the NETS separate cell types as decoded by unique transcriptional fingerprints, including endothelial cells, vascular mural cells, NCCs, mesothelial cells and fibroblasts. Among the fibroblast population, at least 11 distinct capsular and medullary subtypes are recognized, thus largely extending the previously identified heterogeneity defined mostly by phenotypic markers and bulk RNA sequencing (*13, 37, 52*). These subtypes display dynamic changes in their relative representation over time and demonstrate RNA splicing patterns that identify CapFb1a, CapFb2b and MedFb1a as fibroblast subtypes with precursor potential and CapFb3 to originate from mesothelial cells as this fibroblast subtype continues to expresses several mesothelium-specific biomarkers, including *Wt1*, *Cxcl13* and *Rspo1* (Fig. S12; ref. (*14*)).

Cells with gene expression profiles typical of arterial, capillary and venous vasculature are detected already at embryonic day E12.5 when the colonization of the thymus by haematopoietic precursor cells has been initiated independent of an established vasculature (*53*). After E15.5, the frequency of pericytes and vascular smooth muscle cells increases which coincides with the histological evidence of vessel formation. The gate keeper molecules P-selectin, ICAM-1, VCAM-1 and CCL25 enable the entry of T cell precursor into the thymic microenvironment and we find these molecules expressed by all thymic endothelia in the postnatal thymus (*54*). The expression level of the adhesion molecules increases parallel to the age of the mouse but differs between distinct anatomical sites along the vasculature (Fig. S13). This expression pattern specifies that T cell precursors enter the thymic microenvironment via postcapillary venules in a gated and temporally-controlled way (*55*) and that the efficiency of this process may differ between developmental stages. The further development of these haematopoietic cells is regulated by membrane bound Kit ligand which we find expressed by all endothelia and thus also at those anatomical locations where haematopoietic precursors enter the thymus microenvironment (Fig. S12).

450
Endothelial cells also regulate in a non-redundant fashion the egress of mature thymocytes via the
expression of sphingosine-1-phosphate (S1P) lyase (encoded by *Sgpl1)*, lipid phosphate
phosphatase 3 (*Pllp3*) and spinster homologue 2 (*Spns2*) that modify SIP availability and enable
the molecular transport, respectively (*56-58*). Transcripts for *Ppl3* and *Spns1* are detected in all
endothelial cell types even paradoxically at a time of development when T cell export has not yet
commenced (Fig. S13). This expression pattern designates the anatomical site from where
thymocytes can exit and indorses the molecular mechanism by which this process is controlled
(*59*).
459
The NETS collectively promotes the proliferation and differentiation of TEC either indirectly via
ligands that engage for example the platelet-derived growth factor receptor-alpha (PDGFRα) (*60*)
or directly via different signalling ligands, including Wnts, BMP4, Fgf7 and Fgf10 (*3, 61, 62*).
For example, BMP is expressed by CapFb1c together with Fgf10 and upregulates FOXN1, a
transcription factor indispensable for TEC differentiation and function (*61-63*). Wnt4 which also
stimulates the up-regulation of FOXN1 both in an auto- and paracrine way is not expressed by
any of the identified thymic fibroblast subtypes but detected in mesothelial cells, thymocytes and
epithelia within the thymus (*62, 64*). Moreover, three of the four capsular fibroblast subtypes
express a range of Wnt ligands albeit none that had previously been implicated in stimulating
FOXN1 expression. The transcriptome of individual thymic fibroblast subtypes also infers that
Wnt-mediated signals are furthermore either positively modulated by R-Spondin 1 secreted by
mesothelia or negatively delimited by Kremen and Dickkopf-1 which are expressed by other
cellular components within the stroma (*65, 66*). Moreover, CapFb4 fibroblasts which express
endosialin (*CD248*) have previously been implicated in the maintenance and regeneration of TEC
(*67*).
475
A major feature of fibroblasts is their capacity to express extracellular matrix components which
form scaffolds that differ regionally in their composition, shape, biophysical characteristics and
functions (*50*). The heterogeneity and distinct spatial distribution of individual stromal cells
therefore accounts for the diverse properties of the extracellular matrix with collagen expression
restricted to capsular fibroblasts and transcripts for laminins detected more widely across capsular
and medullary fibroblast subtypes (Fig. S3). Patterning of the extracellular matrix is critical in
supporting thymic organogenesis, suggesting that thymic mesenchymal diversity will likely have
a broad impact on thymic *in vivo* function (*8*).
484
Single cell sequencing of human thymus tissue identifies a heterogeneity among NETS that is
similar to the variance observed in mice and thus constitutes a trans-species phenomenon (*37, 52*).
Surprisingly, a relatively large population of NCCs is still detected at 14- and 17-weeks post-
conception, i.e. at a time when thymus morphogenesis has ended and full function has been
attained (*68*). This finding thus contrasts the results observed in mouse tissue at a corresponding
developmental stage since the thymus of mice largely lacks NCCs as early as E16.5. This
incongruity suggests that contrary to mice human NCCs have a more enduring role in shaping the
NETS compartment. The analysis of human thymus tissue at late foetal stages of thymus
organogenesis reveals two distinct NCC subtypes each with a seemingly different developmental
potential, bespoke chromatin accessibility and proliferation dynamics. One of the two NCC
subtypes, i.e. NCC-I, represents a subtype that is poised to adopt a mesenchymal and perivascular
cell fate. Notably, the corresponding cell type is not identified in the mouse thymus. This
suggests that the transition between neural crest-derived cells and perivascular cells in mice must
either be rapid and profound, or, alternatively, occurs at an embryonic time-point that is not

captured in our dataset as the lineage mapping of NCCs has previously identified their
differentiation into VSM and pericytes(*12, 31*).
501
The murine model of 22q11DS reveals major quantitative and qualitative changes in the thymus
stroma which for the first time are highlighted by a decreased proportional representation of
several TEC subtypes and mesenchymal cells, the latter including a selection of capsular and
medullary fibroblasts and the NCC-derived pericytes and VSM. However, *Tbx1* and *Crkl*
transcripts are only solidly detected in a selection of mouse thymus fibroblast subtypes at the
earliest stages of organogenesis and are notably absent in thymic NCCs. Indeed, NCCs retain
their relative frequency in the presence of compound *Crkl* and *Tbx1* haploinsufficiency. Hence,
the observed modifications in vascular mural cells are not the result of a reduced thymic NCC
frequency or changes in *Crkl* and *Tbx*-controlled gene expression in these cells having migrated to
the thymus. Rather, our data suggests that they are the consequence of altered TGFβ receptor
mediated signalling in pericytes, VSM and likely their immediate precursors as this requires the
involvement of Crkl, with *Tgfb1* expressed by endothelial cells and *Tgfb3* by capsular fibroblasts
(Fig. S14) (*69*). Yet, the absence of normal Crkl-dependent signalling in different
haploinsufficient fibroblast subtypes may in addition and indirectly impair pericytes and VSM
development, thus arguing for a hitherto unexplored aspect of intra-thymic cellular crosstalk.
517
Overall, our findings have identified previously underappreciated levels of cellular heterogeneity
and developmental dynamics within the non-TEC thymic stromal compartment. Cellular diversity
within this compartment is present both in murine and human thymic development, but shows
clear trans-species differences worthy of further investigation. Many of these cell populations are
disrupted in 22q11.2 DS, a syndrome known to cause defective thymic organogenesis and
function. Further work should focus on identifying the precise function of each fibroblast cell
subpopulation, along with their contribution to overall thymic development and function.
525

Materials and Methods

Mice

All mice were maintained under specific pathogen free conditions and according to United Kingdom Home Office regulations and federal regulations and permissions, depending on where the mice were housed. Wild-type C57BL/6 mice originated were bred in-house. A mouse line carrying a germ-line Crkl null allele (Crkl[tm1d(EUCOMM)Hmgu]/ImoJ) was generated with Cre-mediated recombination in the epiblast by crossing the Crkl-flox mice (Crkl[tm1c(EUCOMM)Hmgu]/ImoJ) (*70*) with Meox2 Cre knock-in strain (*71*), followed by backcrosses with wild-type C57BL/6 mice to segregate out Meox2Cre. Tbx1[lacz/+] (*72*) mice were obtained from Prof. Antonio Baldini via Prof. Peter Scambler at University College London. Tbx1 and Crkl compound heterozygous mice (Tbx1[+/-]Crkl[+/-]) were generated by crossing between Tbx1[+/-] males with Crkl[+/-] females. Embryos of specific embryonic ages were obtained through timed mating where the presence of vaginal plug was defined as embryonic day (E)0.5.

Isolation of mouse thymic stromal cells and preparation for flow cytometry

Thymic cell suspensions were obtained via enzymatic digestion of thymic lobes using Liberase (Roche) and DNaseI (Roche). To enrich for non-haematopoietic stromal cells in thymic digests from adult mice, cell suspensions were counted and stained with anti-CD45 microbeads (Miltenyi Biotec) for 15 min on ice, before negative selection using the AutoMACS (Miltenyi Biotec) system. Enriched samples or non-enriched samples were then stained for cell surface markers for 30 min at 4°C. For intracellular staining, the Foxp3 Transcription Factor Staining Buffer Kit (eBioscience) was used according to the manufacturer's instructions. Combinations of UEA-1 lectin (Vector Laboratories) labelled with BV605 and the following antibodies were used to stain the cells: TER-119::BV421 (BioLegend), CD45::AF700 (30-F11, BioLegend), EpCAM::PerCPCy5.5 (G8.8, BioLegend), Ly51::PE (6C3, BioLegend), CD80::PECy5 (16-10A1, BioLegend), CD26::PECy7 (H194-112, BioLegend), MHCII::APCCy7 (M5/114.15.2, BioLegend), MHCII::BV421 (M5/114.15.2, BioLegend), CD31::AF488 (MEC13.3, BioLegend), podoplanin (gp38)::AF647(PMab-1, BioLegend). DAPI or the LIVE/DEAD Fixable Aqua Dead Cell Stain Kit was used (Thermo Fisher Scientific) for the assessment of cell viability. After staining, cells were acquired and sorted using a FACS Aria III (BD Biosciences) and analysed using FlowJo v10 and GraphPad Prism 8. Statistical analyses were performed using t-tests, with correction for multiple comparisons where appropriate. A p-value or the adjusted P-value of $\leq$ 0.05 was considered statistically significant.

Immunofluorescent microscopy for extracellular matrix proteins

Thymus from a 5-weeks-old female WT C57BL/6J mouse was used. The standard procedure for immunofluorescence on tissue sections was described here (https://www.biorxiv.org/content/10.1101/2021.03.21.436320v1). Briefly, organs are collected in PBS, fixed in 4% paraformaldehyde overnight at 4°C on a rotating shaker. Organs were then washed in PBS, and lobes separated for the next steps. Paraffin infiltration was done using a Tissue-Tek VIP 6 AI Vacuum Infiltration Processor (Sakura). Lobes were then embedded in paraffin and 4μm sections cut with a Hyrax M25 microtome (Zeiss).
Before immunostaining, de-waxing and antigen retrieval in citrate buffer at pH6.0 (using a heat-induced epitope retrieval PT module, ThermoFischer Scientific) were performed. Sections were then blocked and permeabilized for 30min in 1% BSA, 0.2% Triton X-100 in PBS and blocked for 30min in 10% donkey serum (Gibco) in PBS at RT. Sections were incubated with primary

antibodies overnight at 4°C in 1.5% donkey serum in PBS. Sections were washed twice in 1%
BSA, 0.2% Triton X-100 in PBS and incubated with secondary antibodies at RT for 45min.
Finally, sections were washed twice in 0.2% Triton X-100 in PBS and mounted with
Fluoromount-G (SouthernBiotech). Pictures were acquired with a CCD DFC 3000 black and
white camera on an upright Leica DM5500 scanning microscope.
Antibodies: goat anti-Fibronectin (Santa Cruz, sc-6953, 1/250); rabbit anti-Collagen 1 (Abcam,
ab21286, 1/250); Donkey anti-goat Alexa 488 (ThermoFischer Scientific, A-11055, 1/500);
Donkey anti-rabbit Alexa 647 (ThermoFischer Scientific, A-31573, 1/500). Nuclei staining: DAPI
(Sigma-Aldrich, 1µg per ml).
585
586 **Immunofluorescent microscopy for CD26 and podoplanin**
587
Freshly isolated thymic lobes were frozen in OCT compound (Tissue-Tek) and cryosectioned at a
thickness of 10 µm. Tissue sections were fixed with ice cold acetone for 5 min and blocked with
Avidin/Biotin Blocking Kit (Vector laboratories) and Protein block (Protein block (Dako)
according to manufacturer's protocol. Tissue sections were then incubated with primary
antibodies at 4 ºC overnight: rabbit anti-mouse CD26 (DPP4) (EPR5883(2), Abcam) and biotin
anti-mouse podoplanin (8.1.1, Biolegend). Secondary antibody staining was performed at room
temperature for 30 min with anti-rabbit:AF488 (Invitrogen) and streptavidin-AF555 (Invitrogen)).
Nuclei were stained with Hoechst 34580 in PBS (according to manufacturer's protocol). Sections
were mounted with ProLong Gold Antifade Mountant (Thermo Fisher Scientific) and acquired
using an LSM700 confocal microscope (Carl Zeiss AG). Image analysis was performed with
ImageJ software (Rasband WS, ImageJ, US National Institutes of Health, Bethesda, Md).
599
600 **Single cell RNA sequencing**
601
Total thymic non-epithelial stromal cells (Live Ter119-CD45-EpCAM-) thymic cells from E12.5,
E13.5, E16.5, P0 and 4-week-old wild type mice were sorted and kept on ice before they were
counted. 18,000 cells per sample were loaded onto a Chromium Single Cell B Chip (10x
Genomics) followed by library preparation using Chromium Single Cell 3′ solution (10x
Genomics) and sequencing by NovaSeq6000. (28+98) (Illumina). For the Tbx1$^{LacZ/+}$Crkl$^{+/-}$
dataset, total non-haematopoietic stromal cells (Live Ter119-CD45-) from P0 Tbx1$^{LacZ/+}$Crkl$^{+/-}$
(n=3) and their wildtype littermates (n=3) were sorted and fixed using RNAprotect Cell Reagent
(Qiagen) for storage before sample submission to the Oxford Genomics Centre, where all
downstream steps were performed including 10x Genomics Chip loading, library preparation and
sequencing.
612
613 **Single cell RNA sequencing analysis**
614
Sequencing reads were processed using Cell Ranger (version 3.1.0). Cells were retained for
downstream analysis if there was expression of >1,000 genes, <5% of UMIs mapped to
mitochondrial genes, cells were called as singlets by DoubletFinder, and cells did not cluster into
Ptprc (CD45)-expressing clusters or other contaminant clusters (such as thymic epithelial cells or
clusters present only in one replicate) (*73*). Seurat was used to remove batch effect between
samples using canonical correlation analysis-based integration (*74*). Cells were projected into
two-dimensional space using Uniform Manifold Approximation and Projection (UMAP). Clusters
were called using a resolution of 0.8 and cell label transfer between datasets was undertaken using
Seurat. Differential analysis between clusters used Wilcoxon-rank sum testing and over different
ages used the Kruskal-Wallis analysis of variance. P-values were corrected for multiple
hypothesis testing using the Benjamini-Hochberg method. GENIE3 and RcisTarget were used to

identify gene-regulatory networks on highly variable genes expressed in at least 5% of cells with
subsequent module expression calculated using Seurat (*75*). RNA velocity analysis was
undertaken using Velocyto and scVelo (*76, 77*). Ligand-receptor-target networks were inferred
using Nichenetr, with differential expression assessed between early (E12.5/13.5) and late (E16.5)
embryogenesis (*38*). Gene ontology analysis was undertaken using clusterProfiler (version 4.0.0)
(*78*).
632
633 **Single nuclei multiomics of human fetal thymic stroma**
634
Human fetal thymi, obtained from terminations of pregnancy at 14 and 17 post conception weeks
were enzymatically dissociated using Liberase (Roche) and DNaseI (Roche). The resultant cell
suspension was stained with the following antibodies directed against cell surface antigens for 30
minutes at 4°C: CD45::BV421 (H130, BioLegend) and HLA-DR::PE-Cy7 (L243, BioLegend);
7AAD (BioLegend) was used as a viability marker. Live CD45- MHCII intermediate-high cells
were sorted in 250, 000 cell aliquots using a FACS Aria III (BD Biosciences). Samples were then
processed using the 10x Genomics Multiomics ATAC (Assay for Transposase-Accessible
Chromatin using sequencing) + Gene Expression kit according to the manufacturer's protocol
with some adaptations. Specifically, nuclei were isolated using a 0.1x diluted nuclei extraction
buffer for 6 minutes before being captured into droplets on the 10x Genomics Chromium platform
and sequenced on an Illumina NovaSeq machine.
This study of human thymic tissue has been granted ethical approval and is publicly listed (IRAS
ID 156910, CPMS ID 19587).
648
649 **Multiomics analysis**
650
Sequencing data were processed using Cell Ranger ARC (version 1.0.1). Counts and ATAC data
were analysed using Seurat (version 4.0.3) and Signac (version 1.2.1) (*74, 79*). Barcodes were
filtered to high quality cells (ATAC library size 1,000-100,000, RNA library size 1,000-31,622,
ATAC peaks 1,000-31,622, RNA features 1,000-10,000 and proportion of mitochondrial RNA
reads ≤0.15). ATAC peaks were recalled across each sample for all cells. Clusters were called on
integrated RNA data used a clustering threshold of 0.8 and projected onto a joint UMAP plot of
RNA and ATAC components generated using Seurat and Signac. Differential gene expression
between clusters was estimated using the default method in Seurat. Differentially accessible peaks
were identified using the likelihood ratio method with correction for ATAC library size. Motif
activity was estimated using chromVAR with the JASPAR2020 motif dataset (*80*). RNA-ATAC
links were analysed using Signac and Seurat in 50kb windows around genes of interest.
662
663

References

1. G. Anderson, E. J. Jenkinson, N. C. Moore, J. J. Owen, MHC class II-positive epithelium and mesenchyme cells are both required for T-cell development in the thymus. *Nature* **362**, 70-73 (1993).
2. K. D. James, E. J. Cosway, B. Lucas, A. J. White, S. M. Parnell, M. Carvalho-Gaspar, A. V. Tumanov, G. Anderson, W. E. Jenkinson, Endothelial cells act as gatekeepers for LTβR-dependent thymocyte emigration. *Journal of Experimental Medicine* **215**, 2984-2993 (2018).
3. K. M. Sitnik, K. Kotarsky, A. J. White, W. E. Jenkinson, G. Anderson, W. W. Agace, Mesenchymal cells regulate retinoic acid receptor-dependent cortical thymic epithelial cell homeostasis. *J Immunol* **188**, 4801-4809 (2012).
4. K. M. Sitnik, K. Wendland, H. Weishaupt, H. Uronen-Hansson, A. J. White, G. Anderson, K. Kotarsky, W. W. Agace, Context-Dependent Development of Lymphoid Stroma from Adult CD34(+) Adventitial Progenitors. *Cell Rep* **14**, 2375-2388 (2016).
5. J. Baran-Gale, M. D. Morgan, S. Maio, F. Dhalla, I. Calvo-Asensio, M. E. Deadman, A. E. Handel, A. Maynard, S. Chen, F. Green, R. V. Sit, N. F. Neff, S. Darmanis, W. Tan, A. P. May, J. C. Marioni, C. P. Ponting, G. A. Holländer, Ageing compromises mouse thymus function and remodels epithelial cell differentiation. *Elife* **9**, (2020).
6. C. Bornstein, S. Nevo, A. Giladi, N. Kadouri, M. Pouzolles, F. Gerbe, E. David, A. Machado, A. Chuprin, B. Tóth, O. Goldberg, S. Itzkovitz, N. Taylor, P. Jay, V. S. Zimmermann, J. Abramson, I. Amit, Single-cell mapping of the thymic stroma identifies IL-25-producing tuft epithelial cells. *Nature* **559**, 622-626 (2018).
7. W. E. Jenkinson, E. J. Jenkinson, G. Anderson, Differential requirement for mesenchyme in the proliferation and maturation of thymic epithelial progenitors. *J Exp Med* **198**, 325-332 (2003).
8. T. Nitta, H. Takayanagi, Non-Epithelial Thymic Stromal Cells: Unsung Heroes in Thymus Organogenesis and T Cell Development. *Front Immunol* **11**, 620894 (2020).
9. M. B. Buechler, R. N. Pradhan, A. T. Krishnamurty, C. Cox, A. K. Calviello, A. W. Wang, Y. A. Yang, L. Tam, R. Caothien, M. Roose-Girma, Z. Modrusan, J. R. Arron, R. Bourgon, S. Müller, S. J. Turley, Cross-tissue organization of the fibroblast lineage. *Nature* **593**, 575-579 (2021).
10. J. Kalucka, L. P. M. H. de Rooij, J. Goveia, K. Rohlenova, S. J. Dumas, E. Meta, N. V. Conchinha, F. Taverna, L.-A. Teuwen, K. Veys, M. García-Caballero, S. Khan, V. Geldhof, L. Sokol, R. Chen, L. Treps, M. Borri, P. de Zeeuw, C. Dubois, T. K. Karakach, K. D. Falkenberg, M. Parys, X. Yin, S. Vinckier, Y. Du, R. A. Fenton, L. Schoonjans, M. Dewerchin, G. Eelen, B. Thienpont, L. Lin, L. Bolund, X. Li, Y. Luo, P. Carmeliet, Single-Cell Transcriptome Atlas of Murine Endothelial Cells. *Cell* **180**, 764-779.e720 (2020).
11. S. Rafii, J. M. Butler, B. S. Ding, Angiocrine functions of organ-specific endothelial cells. *Nature* **529**, 316-325 (2016).
12. S. M. Müller, C. C. Stolt, G. Terszowski, C. Blum, T. Amagai, N. Kessaris, P. Iannarelli, W. D. Richardson, M. Wegner, H. R. Rodewald, Neural crest origin of perivascular mesenchyme in the adult thymus. *J Immunol* **180**, 5344-5351 (2008).
13. T. Nitta, M. Tsutsumi, S. Nitta, R. Muro, E. C. Suzuki, K. Nakano, Y. Tomofuji, S. Sawa, T. Okamura, J. M. Penninger, H. Takayanagi, Fibroblasts as a source of self-antigens for central immune tolerance. *Nat Immunol* **21**, 1172-1180 (2020).
14. T. Xie, Y. Wang, N. Deng, G. Huang, F. Taghavifar, Y. Geng, N. Liu, V. Kulur, C. Yao, P. Chen, Z. Liu, B. Stripp, J. Tang, J. Liang, P. W. Noble, D. Jiang, Single-Cell

Deconvolution of Fibroblast Heterogeneity in Mouse Pulmonary Fibrosis. *Cell Rep* **22**,
3625-3640 (2018).

15.   N. Morooka, S. Futaki, R. Sato-Nishiuchi, M. Nishino, Y. Totani, C. Shimono, I. Nakano,
H. Nakajima, N. Mochizuki, K. Sekiguchi, Polydom Is an Extracellular Matrix Protein
Involved in Lymphatic Vessel Remodeling. *Circ. Res.* **120**, 1276-1288 (2017).

16.   A. Clemenceau, C. Diorio, F. Durocher, Role of Secreted Frizzled-Related Protein 1 in
Early Mammary Gland Tumorigenesis and Its Regulation in Breast Microenvironment.
Cells **9**, 208 (2020).

17.   P. A. Eisenach, E. Soeth, C. Röder, G. Klöppel, J. Tepel, H. Kalthoff, B. Sipos,
Dipeptidase 1 (DPEP1) is a marker for the transition from low-grade to high-grade
intraepithelial neoplasia and an adverse prognostic factor in colorectal cancer. *Br J Cancer*
109, 694-703 (2013).

18.   M. Vaittinen, M. Kolehmainen, U. Schwab, M. Uusitupa, L. Pulkkinen, Microfibrillar-
associated protein 5 is linked with markers of obesity-related extracellular matrix
remodeling and inflammation. *Nutrition & Diabetes* **1**, e15-e15 (2011).

19.   J. M. Ngu, G. Teng, H. C. Meijndert, H. E. Mewhort, J. D. Turnbull, W. G. Stetler-
Stevenson, P. W. Fedak, Human cardiac fibroblast extracellular matrix remodeling: dual
effects of tissue inhibitor of metalloproteinase-2. *Cardiovasc Pathol* **23**, 335-343 (2014).

20.   R. Du, B. Liu, L. Zhou, D. Wang, X. He, X. Xu, L. Zhang, C. Niu, S. Liu,
Downregulation of annexin A3 inhibits tumor metastasis and decreases drug resistance in
breast cancer. *Cell Death Dis* **9**, 126 (2018).

21.   S. Yakar, M. L. Adamo, Insulin-like growth factor 1 physiology: lessons from mouse
models. *Endocrinol Metab Clin North Am* **41**, 231-247, v (2012).

22.   T. Zhou, L. Yu, J. Huang, X. Zhao, Y. Li, Y. Hu, Y. Lei, GDF10 inhibits proliferation and
epithelial-mesenchymal transition in triple-negative breast cancer via upregulation of
Smad7. *Aging (Albany NY)* **11**, 3298-3314 (2019).

23.   S. Zhao, J. Zhang, X. Hou, L. Zan, N. Wang, Z. Tang, K. Li, OLFML3 expression is
decreased during prenatal muscle development and regulated by microRNA-155 in pigs.
Int J Biol Sci **8**, 459-469 (2012).

24.   E. Germanov, L. Veinotte, R. Cullen, E. Chamberlain, E. C. Butcher, B. Johnston, Critical
role for the chemokine receptor CXCR6 in homeostasis and activation of CD1d-restricted
NKT cells. *J Immunol* **181**, 81-91 (2008).

25.   M. Kurokawa, S. Matsukura, M. Kawaguchi, K. Ieki, S. Suzuki, S. Watanabe, T. Homma,
M. Yamaguchi, H. Takeuchi, M. Adachi, Interleukin-33-activated dendritic cells induce
the production of thymus and activation-regulated chemokine and macrophage-derived
chemokine. *Int Arch Allergy Immunol* **161 Suppl 2**, 52-57 (2013).

26.   C. Bondjers, L. He, M. Takemoto, J. Norlin, N. Asker, M. Hellström, P. Lindahl, C.
Betsholtz, Microarray analysis of blood microvessels from PDGF-B and PDGF-Rbeta
mutant mice identifies novel markers for brain pericytes. *Faseb j* **20**, 1703-1705 (2006).

27.   E. M. Southard-Smith, L. Kos, W. J. Pavan, Sox10 mutation disrupts neural crest
development in Dom Hirschsprung mouse model. *Nat Genet* **18**, 60-64 (1998).

28.   L. Teng, N. A. Mundell, A. Y. Frist, Q. Wang, P. A. Labosky, Requirement for Foxd3 in
the maintenance of neural crest progenitors. *Development* **135**, 1615-1624 (2008).

29.   H. R. Rodewald, Thymus organogenesis. *Annu Rev Immunol* **26**, 355-388 (2008).

30.   D. B. Klug, C. Carter, I. B. Gimenez-Conti, E. R. Richie, Cutting edge: thymocyte-
independent and thymocyte-dependent phases of epithelial patterning in the fetal thymus.
J Immunol **169**, 2842-2845 (2002).

31.   K. Foster, J. Sheridan, H. Veiga-Fernandes, K. Roderick, V. Pachnis, R. Adams, C.
Blackburn, D. Kioussis, M. Coles, Contribution of neural crest-derived cells in the
embryonic and adult thymus. *J Immunol* **180**, 3183-3189 (2008).

32. R. Dixit, X. Ai, A. Fine, Derivation of lung mesenchymal lineages from the fetal
mesothelium requires hedgehog signaling for mesothelial cell entry. *Development* **140**,
4398-4406 (2013).
33. S. Aibar, C. B. González-Blas, T. Moerman, V. A. Huynh-Thu, H. Imrichova, G.
Hulselmans, F. Rambow, J.-C. Marine, P. Geurts, J. Aerts, J. van den Oord, Z. K. Atak, J.
Wouters, S. Aerts, SCENIC: single-cell regulatory network inference and clustering.
Nature Methods, (2017).
34. M. Noizet, E. Lagoutte, M. Gratigny, M. Bouschbacher, I. Lazareth, H. Roest Crollius, X.
Darzacq, C. Dugast-Darzacq, Master regulators in primary skin fibroblast fate
reprogramming in a human ex vivo model of chronic wounds. *Wound Repair Regen* **24**,
247-262 (2016).
35. N. M. Thalji, M. A. Hagler, H. Zhang, G. Casaclang-Verzosa, A. A. Nair, R. M. Suri, J. D.
Miller, Nonbiased Molecular Screening Identifies Novel Molecular Regulators of
Fibrogenic and Proliferative Signaling in Myxomatous Mitral Valve Disease. *Circ
Cardiovasc Genet* **8**, 516-528 (2015).
36. J. Qing, C. Liu, L. Choy, R. Y. Wu, J. S. Pagano, R. Derynck, Transforming growth factor
beta/Smad3 signaling regulates IRF-7 function and transcriptional activation of the beta
interferon promoter. *Mol Cell Biol* **24**, 1411-1425 (2004).
37. J. L. Bautista, N. T. Cramer, C. N. Miller, J. Chavez, D. I. Berrios, L. E. Byrnes, J.
Germino, V. Ntranos, J. B. Sneddon, T. D. Burt, J. M. Gardner, C. J. Ye, M. S. Anderson,
A. V. Parent, Single-cell transcriptional profiling of human thymic stroma uncovers novel
cellular heterogeneity in the thymic medulla. *Nature Communications* **12**, 1096 (2021).
38. R. Browaeys, W. Saelens, Y. Saeys, NicheNet: modeling intercellular communication by
linking ligands to target genes. *Nat Methods* **17**, 159-162 (2020).
39. D. M. McDonald-McGinn, K. E. Sullivan, B. Marino, N. Philip, A. Swillen, J. A.
Vorstman, E. H. Zackai, B. S. Emanuel, J. R. Vermeesch, B. E. Morrow, P. J. Scambler,
A. S. Bassett, 22q11.2 deletion syndrome. *Nat Rev Dis Primers* **1**, 15071 (2015).
40. A. Puech, B. Saint-Jore, B. Funke, D. J. Gilbert, H. Sirotkin, N. G. Copeland, N. A.
Jenkins, R. Kucherlapati, B. Morrow, A. I. Skoultchi, Comparative mapping of the human
22q11 chromosomal region and the orthologous region in mice reveals complex changes
in gene organization. *Proc Natl Acad Sci U S A* **94**, 14608-14613 (1997).
41. D. L. Guris, G. Duester, V. E. Papaioannou, A. Imamoto, Dose-dependent interaction of
Tbx1 and Crkl and locally aberrant RA signaling in a model of del22q11 syndrome. *Dev
Cell* **10**, 81-92 (2006).
42. K. Lima, T. G. Abrahamsen, I. Foelling, S. Natvig, L. P. Ryder, R. W. Olaussen, Low
thymic output in the 22q11.2 deletion syndrome measured by CCR9+CD45RA+ T cell
counts and T cell receptor rearrangement excision circles. *Clin Exp Immunol* **161**, 98-107
(2010).
43. R. Thomas, W. Wang, D. M. Su, Contributions of Age-Related Thymic Involution to
Immunosenescence and Inflammaging. *Immun Ageing* **17**, 2 (2020).
44. B. A. Benayoun, E. A. Pollina, P. P. Singh, S. Mahmoudi, I. Harel, K. M. Casey, B. W.
Dulken, A. Kundaje, A. Brunet, Remodeling of epigenome and transcriptome landscapes
with aging in mice reveals widespread induction of inflammatory responses. *Genome Res*
29, 697-709 (2019).
45. J. W. Kasik, Y. J. Wan, K. Ozato, A burst of c-fos gene expression in the mouse occurs at
birth. *Mol Cell Biol* **7**, 3349-3352 (1987).
46. L. Kochilas, S. Merscher-Gomez, M. M. Lu, V. Potluri, J. Liao, R. Kucherlapati, B.
Morrow, J. A. Epstein, The role of neural crest during cardiac development in a mouse
model of DiGeorge syndrome. *Dev Biol* **251**, 157-166 (2002).

47. E. Theveneau, R. Mayor, Neural crest delamination and migration: From epithelium-to-
mesenchyme transition to collective cell migration. *Dev. Biol.* **366**, 34-54 (2012).
48. T. Werner, A. Hammer, M. Wahlbuhl, M. R. Bösl, M. Wegner, Multiple conserved
regulatory elements with overlapping functions determine Sox10 expression in mouse
embryogenesis. *Nucleic Acids Res* **35**, 6526-6538 (2007).
49. R. M. Williams, I. Candido-Ferreira, E. Repapi, D. Gavriouchkina, U. Senanayake, I. T.
C. Ling, J. Telenius, S. Taylor, J. Hughes, T. Sauka-Spengler, Reconstruction of the
Global Neural Crest Gene Regulatory Network In Vivo. *Developmental Cell* **51**, 255-
276.e257 (2019).
50. M. V. Plikus, X. Wang, S. Sinha, E. Forte, S. M. Thompson, E. L. Herzog, R. R. Driskell,
N. Rosenthal, J. Biernaskie, V. Horsley, Fibroblasts: Origins, definitions, and functions in
health and disease. *Cell* **184**, 3852-3872 (2021).
51. M. D. Lynch, F. M. Watt, Fibroblast heterogeneity: implications for human disease. *J Clin*
Invest **128**, 26-35 (2018).
52. J. E. Park, R. A. Botting, C. Dominguez Conde, D. M. Popescu, M. Lavaert, D. J. Kunz, I.
Goh, E. Stephenson, R. Ragazzini, E. Tuck, A. Wilbrey-Clark, K. Roberts, V. R. Kedlian,
J. R. Ferdinand, X. He, S. Webb, D. Maunder, N. Vandamme, K. T. Mahbubani, K.
Polanski, L. Mamanova, L. Bolt, D. Crossland, F. de Rita, A. Fuller, A. Filby, G.
Reynolds, D. Dixon, K. Saeb-Parsy, S. Lisgo, D. Henderson, R. Vento-Tormo, O. A.
Bayraktar, R. A. Barker, K. B. Meyer, Y. Saeys, P. Bonfanti, S. Behjati, M. R.
Clatworthy, T. Taghon, M. Haniffa, S. A. Teichmann, A cell atlas of human thymic
development defines T cell repertoire formation. *Science* **367**, (2020).
53. C. Liu, F. Saito, Z. Liu, Y. Lei, S. Uehara, P. Love, M. Lipp, S. Kondo, N. Manley, Y.
Takahama, Coordination between CCR7- and CCR9-mediated chemokine signals in
prevascular fetal thymus colonization. *Blood* **108**, 2531-2539 (2006).
54. M. L. Scimone, I. Aifantis, I. Apostolou, H. von Boehmer, U. H. von Andrian, A multistep
adhesion cascade for lymphoid progenitor cell homing to the thymus. *PNAS* **103**, 7006-
7011 (2006).
55. D. L. Foss, E. Donskoy, I. Goldschneider, The Importation of Hematogenous Precursors
by the Thymus Is a Gated Phenomenon in Normal Adult Mice. *Journal of Experimental*
Medicine **193**, 365-374 (2001).
56. Y. Maeda, H. Yagi, K. Takemoto, H. Utsumi, A. Fukunari, K. Sugahara, T. Masuko, K.
Chiba, S1P lyase in thymic perivascular spaces promotes egress of mature thymocytes via
up-regulation of S1P receptor 1. *Int. Immunol.* **26**, 245-255 (2013).
57. B. Bréart, W. D. Ramos-Perez, A. Mendoza, A. K. Salous, M. Gobert, Y. Huang, R. H.
Adams, J. J. Lafaille, D. Escalante-Alcalde, A. J. Morris, S. R. Schwab, Lipid phosphate
phosphatase 3 enables efficient thymic egress. *Journal of Experimental Medicine* **208**,
1267-1278 (2011).
58. S. Fukuhara, S. Simmons, S. Kawamura, A. Inoue, Y. Orba, T. Tokudome, Y. Sunden, Y.
Arai, K. Moriwaki, J. Ishida, A. Uemura, H. Kiyonari, T. Abe, A. Fukamizu, M.
Hirashima, H. Sawa, J. Aoki, M. Ishii, N. Mochizuki, The sphingosine-1-phosphate
transporter Spns2 expressed on endothelial cells regulates lymphocyte trafficking in mice.
J Clin Invest **122**, 1416-1426 (2012).
59. M. Matloubian, C. G. Lo, G. Cinamon, M. J. Lesneski, Y. Xu, V. Brinkmann, M. L.
Allende, R. L. Proia, J. G. Cyster, Lymphocyte egress from thymus and peripheral
lymphoid organs is dependent on S1P receptor 1. *Nature* **427**, 355-360 (2004).
60. W. E. Jenkinson, S. W. Rossi, S. M. Parnell, E. J. Jenkinson, G. Anderson, PDGFRalpha-
expressing mesenchyme regulates thymus growth and the availability of intrathymic
niches. *Blood* **109**, 954-960 (2007).

61.   P. T. Tsai, R. A. Lee, H. Wu, BMP4 acts upstream of FGF in modulating thymic stroma
862   and regulating thymopoiesis. *Blood* **102**, 3947-3953 (2003).

863   62.   G. Balciunaite, M. P. Keller, E. Balciunaite, L. Piali, S. Zuklys, Y. D. Mathieu, J. Gill, R.
864   Boyd, D. J. Sussman, G. A. Holländer, Wnt glycoproteins regulate the expression of
865   FoxN1, the gene defective in nude mice. *Nat Immunol* **3**, 1102-1108 (2002).

866   63.   S. Žuklys, A. Handel, S. Zhanybekova, F. Govani, M. Keller, S. Maio, C. E. Mayer, H. Y.
867   Teh, K. Hafen, G. Gallone, T. Barthlott, C. P. Ponting, G. A. Holländer, Foxn1 regulates
868   key target genes essential for T cell development in postnatal thymic epithelial cells.
869   *Nature Immunology* **17**, 1206-1215 (2016).

870   64.   F. Brunk, I. Augustin, M. Meister, M. Boutros, B. Kyewski, Thymic Epithelial Cells Are a
871   Nonredundant Source of Wnt Ligands for Thymus Development. *J. Immunol.* **195**, 5261-
872   5271 (2015).

873   65.   M. Osada, E. Ito, H. A. Fermin, E. Vazquez-Cintron, T. Venkatesh, R. H. Friedel, M.
874   Pezzano, The Wnt Signaling Antagonist Kremen1 is Required for Development of
875   Thymic Architecture. *Clinical and Developmental Immunology* **13**, 602150 (2006).

876   66.   M. Osada, L. Jardine, R. Misir, T. Andl, S. E. Millar, M. Pezzano, DKK1 Mediated
877   Inhibition of Wnt Signaling in Postnatal Mice Leads to Loss of TEC Progenitors and
878   Thymic Degeneration. *PLOS ONE* **5**, e9062 (2010).

879   67.   L. Sun, C. Sun, Z. Liang, H. Li, L. Chen, H. Luo, H. Zhang, P. Ding, X. Sun, Z. Qin, Y.
880   Zhao, FSP1(+) fibroblast subpopulation is essential for the maintenance and regeneration
881   of medullary thymic epithelial cells. *Sci Rep* **5**, 14871 (2015).

882   68.   J. F. George, Jr., H. W. Schroeder, Jr., Developmental regulation of D beta reading frame
883   and junctional diversity in T cell receptor-beta transcripts from human thymus. *J Immunol*
884   **148**, 1230-1239 (1992).

885   69.   H. Wurdak, L. M. Ittner, K. S. Lang, P. Leveen, U. Suter, J. A. Fischer, S. Karlsson, W.
886   Born, L. Sommer, Inactivation of TGFbeta signaling in neural crest stem cells leads to
887   multiple defects reminiscent of DiGeorge syndrome. *Genes Dev* **19**, 530-535 (2005).

888   70.   E. Lopez-Rivera, Y. P. Liu, M. Verbitsky, B. R. Anderson, V. P. Capone, E. A. Otto, Z.
889   Yan, A. Mitrotti, J. Martino, N. J. Steers, D. A. Fasel, K. Vukojevic, R. Deng, S. E.
890   Racedo, Q. Liu, M. Werth, R. Westland, A. Vivante, G. S. Makar, M. Bodria, M. G.
891   Sampson, C. E. Gillies, V. Vega-Warner, M. Maiorana, D. S. Petrey, B. Honig, V. J.
892   Lozanovski, R. Salomon, L. Heidet, W. Carpentier, D. Gaillard, A. Carrea, L. Gesualdo,
893   D. Cusi, C. Izzi, F. Scolari, J. A. van Wijk, A. Arapovic, M. Saraga-Babic, M. Saraga, N.
894   Kunac, A. Samii, D. M. McDonald-McGinn, T. B. Crowley, E. H. Zackai, D. Drozdz, M.
895   Miklaszewska, M. Tkaczyk, P. Sikora, M. Szczepanska, M. Mizerska-Wasiak, G.
896   Krzemien, A. Szmigielska, M. Zaniew, J. M. Darlow, P. Puri, D. Barton, E. Casolari, S. L.
897   Furth, B. A. Warady, Z. Gucev, H. Hakonarson, H. Flogelova, V. Tasic, A. Latos-
898   Bielenska, A. Materna-Kiryluk, L. Allegri, C. S. Wong, I. A. Drummond, V. D'Agati, A.
899   Imamoto, J. M. Barasch, F. Hildebrandt, K. Kiryluk, R. P. Lifton, B. E. Morrow, C.
900   Jeanpierre, V. E. Papaioannou, G. M. Ghiggeri, A. G. Gharavi, N. Katsanis, S. Sanna-
901   Cherchi, Genetic Drivers of Kidney Defects in the DiGeorge Syndrome. *N Engl J Med*
902   **376**, 742-754 (2017).

903   71.   M. D. Tallquist, P. Soriano, Epiblast-restricted Cre expression in MORE mice: a tool to
904   distinguish embryonic vs. extra-embryonic gene function. *Genesis* **26**, 113-115 (2000).

905   72.   E. A. Lindsay, F. Vitelli, H. Su, M. Morishima, T. Huynh, T. Pramparo, V. Jurecic, G.
906   Ogunrinu, H. F. Sutherland, P. J. Scambler, A. Bradley, A. Baldini, Tbx1
907   haploinsufficieny in the DiGeorge syndrome region causes aortic arch defects in mice.
908   *Nature* **410**, 97-101 (2001).

909   73.   C. S. McGinnis, L. M. Murrow, Z. J. Gartner, DoubletFinder: Doublet Detection in
910   Single-Cell RNA Sequencing Data Using Artificial Nearest Neighbors. *Cell Syst* **8**, 329-
911   337.e324 (2019).
912   74.   T. Stuart, A. Butler, P. Hoffman, C. Hafemeister, E. Papalexi, W. M. Mauck, 3rd, Y. Hao,
913   M. Stoeckius, P. Smibert, R. Satija, Comprehensive Integration of Single-Cell Data. *Cell*
914   **177**, 1888-1902.e1821 (2019).
915   75.   S. Aibar, C. B. González-Blas, T. Moerman, V. A. Huynh-Thu, H. Imrichova, G.
916   Hulselmans, F. Rambow, J.-C. Marine, P. Geurts, J. Aerts, J. van den Oord, Z. K. Atak, J.
917   Wouters, S. Aerts, SCENIC: single-cell regulatory network inference and clustering.
918   *Nature Methods* **14**, 1083-1086 (2017).
919   76.   V. Bergen, M. Lange, S. Peidli, F. A. Wolf, F. J. Theis, Generalizing RNA velocity to
920   transient cell states through dynamical modeling. *Nature Biotechnology*, (2020).
921   77.   G. La Manno, R. Soldatov, A. Zeisel, E. Braun, H. Hochgerner, V. Petukhov, K.
922   Lidschreiber, M. E. Kastriti, P. Lönnerberg, A. Furlan, J. Fan, L. E. Borm, Z. Liu, D. van
923   Bruggen, J. Guo, X. He, R. Barker, E. Sundström, G. Castelo-Branco, P. Cramer, I.
924   Adameyko, S. Linnarsson, P. V. Kharchenko, RNA velocity of single cells. *Nature* **560**,
925   494-498 (2018).
926   78.   T. Wu, E. Hu, S. Xu, M. Chen, P. Guo, Z. Dai, T. Feng, L. Zhou, W. Tang, L. Zhan, X.
927   Fu, S. Liu, X. Bo, G. Yu, clusterProfiler 4.0: A universal enrichment tool for interpreting
928   omics data. *The Innovation* **2**, 100141 (2021).
929   79.   T. Stuart, A. Srivastava, C. Lareau, R. Satija, Multimodal single-cell chromatin analysis
930   with Signac. *bioRxiv*, 2020.2011.2009.373613 (2020).
931   80.   A. N. Schep, B. Wu, J. D. Buenrostro, W. J. Greenleaf, chromVAR: inferring
932   transcription-factor-associated accessibility from single-cell epigenomic data. *Nature*
933   *Methods* **14**, 975-978 (2017).
934
935

Acknowledgments

Funding:
Swiss National Science Foundation grant IZLJZ3_171050; 310030_184672 (GAH)
Medical Research Council grant MR/S036407/1 (GAH, AEH)
Wellcome Trust grant 105045/Z/14/Z (GAH)
The work was generously supported by a donor of Stanford's Center for Definitive and
Curative Medicine 22q11 Deletion Syndrome Consortium (GAH)
National Institute for Health Research Oxford Biomedical Research Centre grant (GAH)
NIHR Clinical Lectureship (AEH, FD)
Health Research Bridging Salary Scheme (AEH)
Wellcome Trust grant (FD)
International postdoc fellowship from the Swedish research council (Vetenskapsrådet)
(SC)
The human embryonic and fetal material was provided by the Joint MRC/Wellcome Trust
(grant # MR/006237/1) Human Developmental Biology Resource (www.hdbr.org).

Author contributions:
Conceptualization: AEH, SC, ML, KW, GAH
Investigation: SC, FD, SM, TH, MED
Analysis: AEH, SC
Visualization: AEH, SC, FD, IR, TH, GAH
Supervision: ML, GAH
Writing—original draft: AEH, SC, GAH
Writing—review & editing: AEH, SC, FD, SM, TH, IR, MED, OE, ML, KW, GAH

Competing interests: Authors declare that they have no competing interests.

965 **Figures and Tables**
966
967

968
Fig. 1 Heterogeneity of the thymic stroma at 4 weeks of age. (**a**) Representative FACS plot of
live Ter119⁻ CD45⁻ EpCAM⁻ thymic stromal cells at 4 weeks old (left), the relative frequency
(middle) and cellularity (right) of CD31⁺, Ly51+gp38⁻, CD26-gp38⁺ and CD26+gp38⁺ cells
among total stroma or non-epithelial stroma. (b) Immunofluorescence staining of thymic
mesenchymal cells (Red: gp38, Green: CD26, Blue: DAPI). (**c**) A UMAP plot of Ter119-CD45-
EpCAM- cells from 4-week-old mice. (**d**) A heatmap of top 5 differentially expressed genes
between each cluster. (**e**) An overlay on UMAP plot and (**f**) violin plots of the expression of genes
specific to each FACS-isolated subpopulation from bulk RNA-seq and pie charts showing the

proportion of cell types which express gene signatures characteristic of specific FACS-isolated
populations as inferred from bulk RNASeq. CapFb = capsular fibroblast; En = endothelium;
MedFb = medullary fibroblast; Mes = mesothelium; NCC = neural crest; Pc = pericyte; Pr =
proliferating cell; VSM = vascular smooth muscle. (**g**) Immunofluorescence microscopy showing
the distribution of type-I and fibronectin in a 5-week-old mouse thymus. FACS data shown in
panel a were representative of one experiment (n=5) out of two independent experiments (total
n=7), and mean value and SD are shown in the corresponding bar graphs (n=5).
984
985

Fig. 2 Characteristics of the thymic stroma over developmental time. (**a**) (upper) representative FACS plot of the non-epithelial thymic stroma from E12.5 to week 4. (lower) and the relative frequency of stromal subpopulations among the total stroma (left) or the non-TEC stroma (right). (**b**) A combined UMAP plot of Ter119-CD45-EpCAM- cells from mice at E12.5, E13.5, E16.5, P0 and 4 weeks of age. (**c**) (i) Individual UMAP plots for each developmental timepoint, with cell number down-sampled to the smallest sample size (n = 1,997). (**ii**) Scaled proportional representation of each age in each cluster. (**d**) A UMAP plot with each cell coloured by the maximum transcription factor gene-regulatory network expression. Gene-regulatory network centroids are labelled. (**e**) An RNA velocity plot with velocity streamlines projected onto UMAP plot. CapFb = capsular fibroblast; EnA = arterial endothelium; ENCC = capillary endothelium; EnVL = venous/lymphatic endothelium; MedFb = medullary fibroblast; Mes = mesothelium; NCC = neural crest; Pc = pericyte; Pr = proliferating cell; VSM = vascular smooth

999 muscle. E12.5 and E13.5 were sorted and analysed from thymi pooled from two litters consist of
1000 at least 3 embryos per litters. For E16.5, P0 and week 4, data shown consist of cells sorted from
1001 two thymi per timepoint. Mean value and SD were shown in the bar charts (a).
1002
1003

1004
Fig. 3 Ligand-receptor interaction of neural crest derived cells and other thymic cell types.
(a) (Upper) Heatmap of top ligand-receptor interactions of cell types signalling to NCC showing
the Pearson's r for ligand activity in promoting ageing transition between E12.5/13.5 and E16.5.
(Lower) Violin plots of the cell type-specific expression of the top two ligands, Tgfb1 and Vwf.
(**b**) A UMAP plot showing weighted connections for the top 25% ligand-receptor-target networks
for connections from neural crest cells. The width of the line is proportional to the strength of
ligand-receptor interactions. (**c**) UMAP and violin plots of *Jam2* and *Jam3* expression.
1012

1013
Fig. 4. Reduction of non-epithelial thymic stroma in Tbx1+/-Crkl+/- thymi. (a) Total
cellularity, Number of CD45[+], EpCAM[+] and total mesenchymal cells at E13.5 wildtype (n=9) and
Tbx1[+/-]Crkl[+/-] (n=5) thymi. **(b)** Total thymic cellularity, absolute number of CD45+, EpCAM+,
CD31+, Ly51[+]gp38[-], CD26[-]gp38[+], CD26[+]gp38[+] cells and the frequency of stromal
subpopulations from wildtype (P0: n=9, 4-week: n=6) and Tbx1[+/-]Crkl[+/-] (P0: n=5, 4-week: n=6)
neonates **(b)** and 4-week old mice **(c)**. Data was normalised to the mean of wildtype litter mates
from two independent experiments (n=3 from each experiment). Mean values were shown in the
bar charts. Unpaired t-test, $p<0.05(*)$, $0.01(**)$, $0.001(***)$.

1022
Fig. 5 Thymic stromal cells from Tbx1$^{+/-}$Crkl$^{+/-}$ and wild-type mice differ transcriptomically
within cellular populations. A total of 10,300 cells from *Tbx1$^{+/-}$Crkl$^{+/-}$* and 11,151 cells from
their wildtype littermates were analysed. (**a**) A UMAP plot showing thymic stromal cell types.
Scatter plot of genotype-specific enrichment of cell type frequency in Tbx1$^{+/-}$Crkl$^{+/-}$ and wild-type
within NETS (**b**) and TEC (**c**). Error bars show 95% confidence intervals. (**d**) Gene ontology
analysis of genes more highly expressed within each cell type in (**i**) Tbx1$^{+/-}$Crkl$^{+/-}$ and (**ii**) wild-
type thymi. (e) Scatter plot of the difference in ageing score of stromal cell subsets between
Tbx1$^{+/-}$Crkl$^{+/-}$ and wild-type. CapFb = capsular fibroblast; EnA = arterial endothelium; ENCC =
capillary endothelium; EnVL = venous/lymphatic endothelium; itTEC = intertypical TEC;
McTEC = mature cortical TEC; MedFb = medullary fibroblast; Mes = mesothelium; mmTEC =
mature medullary TEC; NCC = neural crest; pAmTEC = post-AIRE medullary TEC; Pc =
pericyte; pcTEC = perinatal cortical TEC; pr = proliferating cell; prTEC = proliferating TEC;
sTEC = structural TEC; Thy = thymocyte; TlTEC = tuft-like TEC; VSM = vascular smooth
muscle. Symbol in red showed cell subtypes significantly enriched in Tbx1$^{+/-}$Crkl$^{+/-}$ and symbol in
blue showed population showed cell subtypes enriched in wildtype (**b,c**). Enrichments were

calculated using Fisher's exact test with 95% confidence intervals and significance was adjusted
for multiple hypothesis testing using Benjamini-Hochberg correction (b,c)
1040

1041
1042
1043

Fig. 6 Age-specific transcriptomic programmes differ over development in thymic stroma.
(**a**) A violin plot showing the overall expression of genes associated with ageing in a tissue-independent manner within the NES across different ages (*44*). (**b**) Violin plots showing the expression of Sox4, Sox11 and Irf7 gene-regulatory networks. (**c**) Violin plot showing the ageing score of fibroblasts, endothelial cells and TECs from Tbx1$^{+/-}$Crkl$^{+/-}$ and wild-type thymi at P0. Symbols in red showed cell subtypes significantly increased ageing score in Tbx1$^{+/-}$Crkl$^{+/-}$ as compared to wildtype and symbols in blue showed population showed significantly reduced ageing score (**d**). Differences in gene module scores were estimated using Wilcoxon tests and were adjusted for multiple hypothesis testing using Benjamini-Hochberg correction (c,d).

1053
1054

1055
Fig. 7 Single nuclei multiomics analysis of human thymic NES demonstrates diverse cell
populations. (a) UMAP plot of 528 nuclei showing joint projection of transcriptomic and
chromatin accessibility data. **(b)** UMAP plot showing marker gene expression for *PDGFRA*,
PDGFRB and *PECAM1*. **(c)** Links plot of the *SOX10 locus* showing chromatin accessibility data,
SOX10 gene expression (violin plot on right), accessible chromatin peaks (blue peaks were
significant for at least one cell type at FDR<0.05) and the correlations between chromatin
accessibility and *SOX10* gene expression (shaded by the strength of each link). Significance of
motif activity and chromatin accessibility were calculated using likelihood ratios, correcting for
the size of chromatin accessibility libraries. P-values were adjusted for multiple hypothesis testing
using Benjamini-Hochberg correction.
1066

Cell type	4-week cluster	All ages cluster	Top genes	Peak age
Capsular fibroblasts	CapFb1	CapFb1a	*Itm2a, Clec3b, Capn6*	E12.5
		CapFb1b	*Cdo1, Ptn, Itm2a*	P0
		CapFb1c	*Lpl, Thbs1, Mt2*	P0
	CapFb2	CapFb2a	*Adamts2, Mfap4, Bgn*	E12.5
		CapFb2b	*Col1a1, Col14a1, Mfap5*	E12.5
	CapFb3	CapFb3	*Igf1, Dcn, Cpxm1*	W4
	CapFb4	CapFb4	*Fbn1, Mfap5, Dpt*	W4
Medullary fibroblasts	MedFb1	MedFb1a	*Sele, Col15a1, Tenm4*	E13.5
		MedFb1b	*Col15a1, Col26a1, Serpine2*	W4
	MedFb2	MedFb2a	*Tmem176a, Des, Tmem176b*	W4
		MedFb2b	*Oasl2, Isg15, Iigp1*	W4
Proliferating fibroblasts	Pr	Pr	*Stmn1, H2afz, Hmgb2*	E13.5
Pericytes	Pc	Pc	*Colec11, Gucy1a1, Ebf1*	E16.5
Vascular smooth muscle	VSM	VSM	*Myh11, Tpm1, Nrip2*	W4
Neural crest cells	NCC	NCC	*Mal, Plp1, Dbi*	E12.5
Endothelium	En	EnA	*Fbln5, Icam2, Egfl8*	W4
		EnC	*Gpihbp1, Rgcc, Fabp4*	W4
		EnVL	*Selp, Pecam1, Aqp1*	W4
Mesothelium	Mes	Mes	*Upk3b, 2010300C02Rik, Krt19*	W4

1067 **Table 1. Thymic stromal cell types identified by single cell RNA-seq.** The three top genes (by
1068 area under the curve) and peak age are shown for each cluster. All clusters show significant
1069 differences in proportional makeup of the non-TEC thymic stromal across different ages (Fisher's
1070 test with 10,000 permutations: all p < 0.0001).