Title Page

WATLAS: high throughput and real-time tracking of many small birds in the Dutch

Wadden Sea

Allert I. Bijleveld[1*], Frank van Maarseveen[1], Bas Denissen[1], Anne Dekinga[1], Emma

Penning[1,2], Selin Ersoy[1,2], Pratik Gupte[1,2], Luc de Monte[1], Job ten Horn[1], Roeland A. Bom[1],

Sivan Toledo[3], Ran Nathan[4], and Christine E. Beardsworth[1]

[1] Department of Coastal Systems, NIOZ Royal Netherlands Institute for Sea Research,

NL-1790 AB, Den Burg, Texel, the Netherlands

[2] Groningen Institute for Evolutionary Life Sciences (GELIFES), University of

Groningen, P.O. Box 11103, 9700 CC Groningen, the Netherlands

[3] Blavatnik School of Computer Science, Tel-Aviv University, Tel Aviv 67798, Israel

[4] Movement Ecology Laboratory, The Alexander Silberman Institute of Life Sciences,

The Hebrew University of Jerusalem, Jerusalem 91904, Israel

*Correspondence to allert.bijleveld@nioz.nl

Abstract

Background. Tracking animal movement is important for understanding how animals

interact with their (changing) environment, and crucial for predicting and explaining how

animals are affected by anthropogenic effects. The Wadden Sea is a UNESCO World

Heritage Site and a region of global importance for millions of small shorebirds. Due to

climate change and anthropogenic activity, understanding and predicting movement and

space-use in areas like the Wadden Sea is increasingly important. Monitoring and predicting

animal movement, however, requires high-resolution tracking of many individuals. While

high-resolution tracking has been made possible through GPS, trade-offs between tag weight

and battery life limit its use to larger species.

Methods. Here, we introduce WATLAS (the Wadden Sea deployment of the ATLAS

tracking system) capable of monitoring the movements of hundreds of (small) birds

simultaneously in the Dutch Wadden Sea. WATLAS employs an array of receiver stations

that can detect and localise small, low-cost tags at fine spatial (meters) and temporal

resolution (seconds). From 2017-2021, we tracked red knots, sanderlings, bar-tailed godwits,

and common terns. We use parts of these data to give four examples on its performance and

how WATLAS can be used to study numerous aspects of animal behaviour, such as, space-

use (both intra- and interspecific), among-individual variation, and social networks.

Results. After describing the WATLAS system, we first illustrate space-use of red knots

across the study area and how the tidal environment affects their movement. Secondly, we

show large among-individual differences in distances travelled per day, and thirdly illustrate

how high-throughput WATLAS data allows calculating a proximity-based social network.

Finally, we demonstrate that using WATLAS to monitor multiple species can reveal

differential space use. For example, despite sanderlings and red knots roosting together, they

foraged in different areas of the mudflats.

Conclusions. The high-resolution tracking data collected by WATLAS offers many

possibilities for research into the drivers of bird movement in the Wadden Sea. WATLAS

could provide a tool for impact assessment, and thus aid nature conservation and management

of the globally important Wadden Sea ecosystem.

Keywords: animal tracking, ATLAS, biologging, biotelemetry, conservation, high-throughput movement ecology, reverse-GPS, shorebirds, space use, Wadden Sea UNESCO World Heritage Site, waders.

Background

Movement is a fundamental aspect of life and tracking wild animals under natural conditions has become central to animal behaviour, ecology, and conservation science [1-5]. Animal tracking has revealed extreme and large-scale migratory journeys [6, 7] and detailed patterns of habitat use [8, 9], as well as elucidated mechanisms of navigation [10-12], predator-prey dynamics [13], and social interactions [14]. Insights from animal tracking studies are regularly incorporated in policy and conservation management [5, 15]. For example, identifying important areas for the protection of migration routes [16, 17], detecting wildlife crime [18, 19], and quantifying the human-wildlife conflict [20].

The introduction of the 'movement ecology' framework [2], coupled with the rapid development of new tracking technologies and data-processing tools [21, 22] has led to an exponential increase in animal movement ecology research [23]. These developments, particularly the miniaturization of tags capable of generating high-throughput localization data for many individuals simultaneously, allow for novel opportunities to address contemporary questions on individual, group, population, and community level behaviours in the wild [24, 25]. For instance, studies on intra-specific variability [26-28], collective behaviour [29], and interactions among individuals and species with their physical, biotic, and anthropogenic environments [30]. Furthermore, the ongoing miniaturization of tags allows the tracking of ever smaller species, and thus may give a more complete pictures how different species use their habitat [1, 25].

The most common tracking methods, which allow monitoring the movement of

animals at high temporal and spatial resolution in the wild, are based on Global Navigation

Satellite Systems (GNSS), such as the Global Positioning System (GPS). GPS employs a

network of orbiting satellites with known locations, which transmit signals to a receiving tag

that uses this information to estimate its position. The principal drawback of GPS is that it

does not directly provide a means of reporting position information back to the researcher.

The position information is either stored and retrieved later or downloaded via an auxiliary

radio-frequency link. In many cases, loggers cannot be retrieved, and the energy costs for

transmitting localization data from the animal to the researcher are large [31]. Therefore,

trade-offs between data retrieval, sampling frequency, battery size and tag weight often limit

the use of these tags and the biological insights gained to larger species [1, 32].

When retrieving loggers is impractical, time-of-arrival (reverse-GPS) and direction-

of-arrival systems provide an alternative tracking method in which the locations of

transmitters need to be estimated and the location of receivers are known. Examples of

reverse-GPS systems are presented by MacCurdy et al. [33], Weller-Weiser et al. [34] and

Krüger et al. [35]. MOTUS [36] and ARTS [37] are examples of direction-of-arrival

transmitter localization. ARGOS [38] is a Doppler-based transmitter-localization satellite

system. Because energy costs of transmissions are low and data processing is handled outside

of the tag at the receiver end, tags can be small and lightweight while maintaining high-

throughput signal transmissions. The smallest VHF-tag compatible with MOTUS, for

instance, weighs 0.13 g (www.lotek.com). The spatial resolution of Doppler and direction-of-

arrival systems, however, is often coarse. ARGOS, for example, provides a spatial resolution

of 250 - 1500 m [38]. MOTUS and ARTS systems can, by triangulating signal-strength of

VHF-tag detections, provide localizations at an accuracy of several hundred meters.

However, the detection ranges are limited to hundreds of meters up to a few km, and

localization error increases with the distance between the tag and receivers [36, 37, 39, 40].

Successful localizations thus require a dense receiver network that limits the size of study

area. Time-of-arrival transmitter localization tends to be more accurate, but similarly requires

a dense receiver network.

ATLAS (Advanced Tracking and Localization of Animals in real-life Systems) is a

recently developed reverse-GPS system [34, 41] that builds on the time-of-arrival wildlife

tracking system of MacCurdy et al. [33], and allows real-time tracking and data collection of

many small individuals simultaneously. ATLAS comprises an array of stationary receivers

that continuously listen for transmissions from small tags. Locations are calculated based on

differences in tag-signal arrival times at minimally three receiver stations. Tags are light-

weight (0.8 g) and relatively inexpensive (25 €), which facilitates tracking small species and

hundreds of individuals simultaneously [42]. Location data is available without retrieval of

the tag and in real time, which avoids the need to recapture tagged animals for data retrieval,

and allows for locating tagged individuals for auxiliary behavioural observations [43] or for

confirming mortality [9]. Whereas GNSS systems and ARGOS allow (near) global tracking,

reverse-GPS systems like ATLAS produces location data at a more local scale and, for best

performance, requires 'line of sight' between receiver and tag [44]. In open landscapes, with

published detection ranges up to 40 km for ATLAS [11], its spatial scale is limited by the

extent and density of receiver stations. ATLAS has been used to track over 50 different

species in four countries, including sites of high scientific or conservation value, such as the

Hula Valley in Israel [11, 34] and the Dutch Wadden Sea.

The Wadden Sea is recognized as a UNESCO World Heritage Site for providing a

rich habitat for marine mammals [45], fish [46], invertebrates [47], birds [48], and especially

migratory shorebirds [49]. Shorebirds form an important component of the Wadden Sea

ecosystem, which they use for breeding [50], refuelling during migratory journeys [51], and

finding food and safety during their non-breeding periods [52-55]. Millions of shorebirds

depend heavily on the worms, snails and shellfish that are found on and in the sediments of

the mudflats [56]. Perhaps uniquely, over the past decade, the Wadden Sea has been subject

to a large scale benthic macrofauna monitoring survey (Synoptic Intertidal BEnthic Survey

(SIBES) [57, 58]), which maps food resources for shorebirds [52, 59]. Combining resource

mapping with the simultaneous tracking of many birds offers novel opportunities for studies

on space use, trophic interactions and collective behaviour in the wild [60]. Many of the

shorebird species are declining in numbers [48], and appear particularly susceptible to the

effects of habitat destruction, disturbance, overexploitation of resources, and global climate

change [61]. Detailed studies of shorebird space use, in conjunction with knowledge of

resource landscapes will offer novel ecological insights and, in combination with monitoring

anthropogenic activities, will allow quantifying if and how animals are impacted, which may

assist in evidence-based conservation efforts in this important region [62]. Moreover, the

Wadden Sea with its flat and open landscape, large numbers of birds, and conservation value,

is an ideal candidate for an ATLAS system.

Here, we introduce WATLAS, which is the Wadden Sea deployment of the ATLAS

tracking system. In 2017, WATLAS started with 5 receivers and has since grown to have 26

receivers in 2021 (Supplementary Fig. S1 in Additional File 1), making it the largest ATLAS

system in the world. The 26 receiver stations are located in the western Dutch Wadden Sea

and encompass 1,326 km^2 (Fig. 1) with a focus on the mudflats surrounding Griend, an

important shorebird high-tide roosting site and nature reserve. WATLAS allows simultaneous

tracking of several hundred animals at fine temporal and spatial resolution comparable to

GPS tracking within a specific study region [63]. So far, WATLAS has been used to track red

knots *Calidris canutus* (~120 g), sanderlings *Calidris alba* (~50 g), bar-tailed godwits *Limosa*

lapponica (~240 g), and common terns *Sterna hirundo* (~130 g), but there is scope to track an

even larger range of species. In this paper, we will introduce WATLAS, along with several

use-cases for the system demonstrating the strengths of the system for studies of movement

across levels of organization: from individuals, to species, to populations, and even

communities. First, to investigate space use and environmental drivers of movement, we

show how space use of red knots tracked in 2019 varies across the entire study area and on a

smaller spatial scale across tidal cycles. Second, we give an example of among-individual

variation in distance travelled for red knots tagged in 2020. Third, we show how fine-scale

movement data WATLAS provides, and how this allows estimating social interactions

(proximity-based networks) in red knots. Fourth, as an example of community tracking, we

show differences in home ranges between sanderlings and red knots, tracked simultaneously

near Richel and Griend. We end by discussing how WATLAS offers possibilities for both

fundamental and applied research into the natural and anthropogenic drivers of bird

movement in the Wadden Sea.

Methods

The deployment of WATLAS, which is the Wadden Sea ATLAS system, comprises an array

of receivers that continuously listen for tag transmissions. When a transmission is detected,

the receiver records the arrival time. These arrival time measurements are sent to a

centralized server where location estimates can be computed when at least three receivers

detect the signal. Receivers can detect a transmission from any tag in the system at any time,

so the tags can transmit as frequently as a localization is needed. Beacons (tags that transmit

in fixed known locations – see BEACONS) enable clock-synchronisation across receiver

stations.

RECEIVERS

The WATLAS system currently consists of 26 receiver stations located in the western

Wadden Sea (Fig. 1). Fourteen receivers were installed on buildings and other stable

structures where power was available, which allowed receivers to be operational year-round.

One year-round receiver was placed high on a dune and powered with twelve 100 W solar

panels (EnjoySolar) that connected to four 100 Ah AGM batteries (Beaut). Eleven receivers

were placed temporarily on the mudflats. Because of the increased likelihood of weather

damage in winter, the temporary receivers (Fig. 2) were only in place between July and

November each year. One of these temporary receivers was placed on an anchored pontoon

that housed a solar powered field station (Fig. 2C). The other ten temporary receivers were

attached to scaffolds (Fig. 2A and B) and powered with four 100 W monocrystalline solar

panels (EnjoySolar) and a 100 W wind turbine (Ampair), which were connected to three 100

Ah AGM batteries (Beaut). For visibility and safety, a solar powered LED-light was placed

on top of the scaffold.

Fig. 1 Map of the study area with the receiver array and beacons in 2019. The locations of

temporary receivers vary between years (Supplementary Fig. S1 in Additional File 1). Land

is shown in dark grey, water in blue and mudflat in light grey. The red square in the inset

shows the study area within Europe. Rectangle A shows the area of Richel and Griend as

shown in Figures 7 and 9. Rectangle B shows the area of Griend as shown in Figures 4 and

6. The coordinate system refers to UTM 31N. © map data from Rijkswaterstaat.

Fig. 2 Examples of the temporary solar- and wind powered receiver stations placed on

mudflats shown at A) low tide and B) high tide, and C) on the field station.

Each receiver had a 1.5 m Ultra High Frequency (UHF) antenna (Diamond X-50N) mounted

on a 6 m aluminium scaffold. To increase the range of tag detections, receiver antennas were

placed as high as possible. Antenna height for temporary receivers in 2019 was on average

9.5 m (range: 8.4 – 11.7 m) above sea level, and on average 18.7 m (range: 10.9 to 44.4 m)

for year-round receivers.

A coaxial cable connected the antenna to a water-proof cabinet (53 x 43 x 20 cm,

Supplementary Fig. S2 in in Additional File 1) via a custom built external Low Noise

Amplifier (LNA). The LNA includes a helical bandpass filter to protect against static

discharge from thunderclouds. These LNAs connect to a custom front-end unit that acts as a

bandpass filter, radio frequency limiter, and power supply to the LNAs [64]. Next comes the

Software Defined Radio (SDR) consisting of an USRP N200 with WBX40 daughter board

(Ettus Research). This SDR precisely timestamps incoming signal detections using a GPS

disciplined oscillator (GPSDO, Ettus Research). The GPSDO was connected to an external

amplified ceramic patch antenna (Ettus Research), which allowed the clock rates of all

receivers to be synchronized with the atomic clocks from GPS-satellites. Signal transmissions

are processed by an onboard computer (Intel NUC i7) that runs Linux. Some computers ran

Ubuntu 16 others Ubuntu 20. The number of unique an ATLAS system can detect depends on

the processing capability of this Intel NUC. With our onboard computers, we estimate that we

can reliably track 300 unique tags that send a transmission every second simultaneously. All

receiver stations were connected to the internet using a cellular modem (Huawei E3372

4G/LTE dongle) and an externally mounted antenna (GTT OS-UMTS-0103-C0) to send

detection reports to a central server at NIOZ Royal Netherlands Institute for Sea Research.

This server runs software that estimates tag locations from time-stamped tag and beacon

detections [34]. All data are stored in an online database running MySQL (v5.7,

https://www.mysql.com/). Localizations are visualised on www.nioz.nl/watlas in real-time.

Moreover, system health (e.g. battery voltage, power usage, and temperature) can also be

monitored remotely and in real time for each receiver.

TAGS

Tags consist of an assembled Printed Circuit Board (PCB), a battery, an antenna, and a

protective coating (Fig. 3). The PCBs are based on a CC1310 or CC1350 microcontroller

with a built-in Radio-Frequency (RF) transceiver that can transmit a code unique to the tag at

433 MHz. In this study, we used 0.6 g PCB-boards (v2.6.1 and 2.6.2f) [42]. The radio signal

is emitted through a 17 cm long antenna made of gold plated multistranded steel wire with a

plastic coating, which can handle mechanical stress in a marine environment. Tags were

coated with a mixture of two-component epoxy (3M Scotch Weld DP270). To reduce tag

weight, the epoxy was mixed with low density glass spheres at a ratio of 1:2. PCB's are fitted

with a (Hall) sensor allowing the tag to be switched on and off with a magnet placed next to

the tag. Tags operate at a voltage of 1.8 to 3.8 V and can be fitted with a range of batteries.

For example, a pair of silver oxide batteries (0.26 g), single lithium coin-cell batteries ranging

from CR1025 (0.7 g) to CR2477 (10.5 g), or a pair of AA batteries (24 g). At signal

transmission costs of approximately 0.4 mJ, the capacity of the battery determines the

number of transmissions that can be sent, and together with the frequency of transmissions,

sets the tag's operational lifetime (longevity). In 2017 and 2018 we used tag transmission

intervals of 1 or 3 s, and in later years 6 s. With WATLAS, we have used CR1620 (1.3 g) and

CR2032 batteries (3.0 g), which resulted in final tag weights of respectively 2.4 and 4.4 g

(Fig. 3). With a signal transmission interval of 6 s this corresponds to an estimated longevity

of 3 and 8 months for the CR1620- and CR2032-loaded tags, respectively (Supplementary

Fig. S3 in Additional File 1).

Fig. 3 *WATLAS tags and batteries. A) tag without battery and coating and a one-euro coin*

for scale. B) 2.4 g coated tag with CR1620 battery. C) a 4.4 g coated tag with CR2032

battery. The batteries of the tags are shown on the left.

BEACONS

Beacons were built as standard WATLAS tags, but fitted with one lithium C cell and a helical

bandpass filter to protect against static discharge from thunderclouds and connected to a

vertical colinear antenna providing 7dB gain in the horizontal plane (Diamond X-50N)

identical to the antennas on receivers. The transmission interval of beacons was set to 1 s.

Seven beacons were mounted on 6 m aluminium scaffold poles (Supplementary Fig. S4 in

Additional File 1). To ensure that each receiver detected at least one beacon consistently,

beacons were placed across the study area (Fig. 1 and Supplementary Fig. S1 in Additional

File 1). During deployment, the locations of receivers and beacons were recorded with dGPS

at 1.5 cm accuracy (Topcon HiPer SR).

WATLAS COSTS

The most substantial costs are setting up an initial array of receivers. The costs of ATLAS components fluctuate, and more economical configurations are being developed. However, at the time of writing the receiver cabinet with the radio frequency electronics costs about 4,500 €. For a temporary receiver station that requires an independent power supply, there is an additional cost of 5,000 € that includes equipment for generating wind and solar power, batteries, and scaffolding. Tag cost is dominated by costs of assembling the electronics, and this largely depends on the numbers of tags produced in a batch: 100 € each at 20 pcs and 22 € each at 200 pcs (www.circuithub.com). The labour costs of tag assembly can easily cost an equal amount. Operational costs can be quite substantial as well, such as those for mobile data transfer. For example, between August and November 2018 receivers transferred an average of 14 GB of data per month (7 to 18 GB per receiver per month). Per receiver, the monthly costs for an unlimited data plan were 35 €.

TRACKING SHOREBIRDS WITH WATLAS

We present examples from red knots tracked in 2018 (N = 79), 2019 (N = 221) and 2020 (N=44), and sanderlings tracked in 2018 (N = 35). Red knots were caught on Richel (53.28° N, 5.01° E) and Griend (53.25° N, 5.25° E) (Fig. 1) with mist nets during new moon periods each year between July and October. Most sanderling (N = 33) were caught on Griend by means of canon netting on 26 July 2018, but some by mist-netting on 12 August 2018 (N = 2). All birds were banded with unique combinations of colour-rings and released after gluing a WATLAS tag to their rump with cyanoacrylate glue (Fig. 4). Red knots were fitted with 4.4 g tags (Fig. 3C) that were on average 3.2 % (SD = 0.2) of body mass. Sanderling were fitted with 2.4 g tags (Fig. 3B) that were on average 4.4 % (SD = 0.4) of body mass. Because red knots and sanderlings can double in body mass while fuelling for migration, we believe the

additional weight of the tag (<5% of body mass) did not cause substantial detrimental effects

[65, 66]. All birds were released from Richel and Griend.

ACCURACY OF WATLAS AND TRACKING DURATION

In another study [63], the bias, accuracy and precision of WATLAS was quantified. In brief,

WATLAS location estimates were comparable to GPS with localization errors of 4 - 6 m.

Accuracy was higher if more receivers detected the tag and localization error was reduced to

a few meters. The tag's location in respect to the receiver array configuration had a large

effect, with less accurate estimates occurring when the tag was on the outskirts or outside the

array of receivers. The proportion of localizations of a stationary test tag 1.2 m above ground

was over 90% of the expected fixes, but for tagged red knots on the ground this proportion

reduced to 51%. Additionally, the proportion of accurate localizations of tagged birds was

higher when they were central to the array, and lower on the outskirts of the array [Fig. 6 in

63]. Nonetheless, with the small interval between tag transmissions, a fix rate of 51% still

resulted in near continuous tracking, and even near the edges of the array provided reliable

and useful localizations.

At a 6s interval, the theoretical longevity of the 4.4 g tags attached to red knots was 8

months. Indeed, this matched the maximum duration of 239 days that red knots were detected

in 2019 (Fig. 9 in [42]). Fig. 9 in Toledo et al. [42], however, also shows a steady decline in

the number of unique tags tracked, which was particularly pronounced directly after release.

Reasons for this decline are multiple and include tags failing, birds dying, and birds leaving

the tracking area. Another important aspect is that the period of tagging coincided with the

period that red knots moult into their winter plumage [67]. Because we glue the tags to the

birds, birds in moult will lose their tags and localizations will cease. In 2021, we recaptured a

red knot that was tagged the previous year and did not show visible signs where the tag had

been attached.

Fig. 4 A colour-ringed red knot in winter plumage, bearing a WATLAS tag glued to its rump;

the tag antenna can be seen extending beyond the tail to the right of the image as indicated

with the red arrow. WATLAS tags allow free movement of the wings and fall off as the

feathers underneath the tag regrow. The inset shows this bird's localizations around Griend,

collected between 15 September and 21 September 2017. See rectangle B in Fig. 1 for

placement of the inset within the study area. © map data from Rijkswaterstaat, and photo

taken on 16 September 2017 by Benjamin Gnep.

PRE-PROCESSING WATLAS DATA TO IMPROVE POSITION ESTIMATES

The accuracy of WATLAS localizations is comparable to conventional GNSS systems [63]

and to the Hula Valley ATLAS system [34]. However, in common with other positioning

systems, WATLAS data can contain some inaccurate localization estimates. Sources of the

localization errors can be due to temporary mismatches in clock synchronization between

receivers, RF-interference, or tag signal collisions. To reduce errors in positioning estimates,

filtering and smoothing localization data is common practice in movement ecology [68].

Here, we used a simple filter-smoothing process on the estimates that are reported by the

localization algorithm, namely variance in the Easting and Northing. We removed

localizations that had variances in Easting and Northing above 2,000 m^2. Additionally, we

smoothed the data with 'runmed' in R [69] by computing a 5-point median smooth across the

localizations [52, 68]. After data processing the precision and error of localizations can be

reduced to several metres [63].

Results

We will demonstrate examples for how WATLAS opens-up possibilities for studying space-

use and environmental drivers of movement, among-individual variation in distance travelled,

intra-specific (social) interactions, and community tracking with interspecific space use in the

wild. All analyses were done in R v4.0.2 [69].

EXAMPLE 1. ESTIMATING SPACE-USE

To show how WATLAS can be used to investigate space use and e.g. identify hotspots, we

created heatmaps of the localisations of 221 red knots tracked between 1 August 2019 and 1

November 2019 (92 days). We created heatmaps at two spatial scales: The large spatial scale

of the entire study area in grid cells of 500 x 500 m, as well as the smaller spatial scale

around Richel and Griend with grid cells of 250 x 250 m. To additionally illustrate how

WATLAS data can be used to investigate environmental drivers of space use, we created

heatmaps on a smaller spatial scale separately for the different phases of the tidal cycle. The

tidal-phases were selected based on the water level (NAP; Amsterdam Ordnance Datum) at

the tide gauge at West-Terschelling including a delay of 30 min due to the distance from

Griend (53.37° N, 5. 22° E): high tide (> 100 cm NAP), first ebb tide (outgoing tide between

50 and 100 cm NAP), second ebb tide (outgoing tide between -50 and 50 cm NAP), low tide

(< -50 cm NAP), first flood tide (incoming tide between -50 and 50 cm NAP), second flood

tide (incoming tide between 50 and 100 cm NAP). These tags were programmed to transmit

every 6 s, thus each location represents at least 6 s of space use.

The large-scale heatmap confirmed that Richel and Griend, where the knots were

caught, are hotspots. Nonetheless, red knots spread-out across the entire study area (Fig. 5). It

should be noted, however, that the number of localizations, being related to the density of

receiving stations, is higher in the central part of the area than on the border. As can be seen

from the localizations over relatively deep gullies, several birds moved between islands and

the mainland (Supplementary Fig. S5 in Additional File 1). Likewise, localizations across the

North Sea suggest that birds crossed in the direction of the United Kingdom. In some cases,

these birds were detected up to 34 km from the closest receiver.

On a smaller spatial scale, the heatmaps for different phases of the tidal cycle around

Richel and Griend showed how the tidal dynamic affects space use of red knots on a

population level (Fig. 6). With the outgoing tide, the birds moved out on the now-exposed

mudflats in search of invertebrate prey. Interestingly, space-use differed between ebb and

flood tides even though the water level was the same. This can, for instance, be seen by

comparing Fig. 6C with 6E, and Fig. 6B with 6F. The fewest localizations were observed

during the low tide period, probably because birds spread out and even moved outside the

tracking area. With the incoming tide the birds returned and aggregated on Richel and

Griend.

Fig. 5 *Large-scale space use of 221 red knots tracked between 1 August and 1 November*

2019 (92 days) within the entire study area. The colour scale represents the number of

localizations in 500 x 500 m grid cells. Note that the colour scale is logarithmic. Water is

coloured blue, land dark grey, and mudflats light grey with a solid line indicating their

boundery. Because the tags send a signal at 1/6 Hz, each localization represents a minimum

of 6 s of space use for red knots. The coordinate system refers to UTM 31N. © map data from

Rijkswaterstaat.

Fig. 6 Small-scale space use during different phases of the tidal cycle for 221 red knots near

Richel and Griend (locations labelled in panel B) between 1 August and 1 November 2019

(92 days). Panels show different phases of the tidal cycle from high tide (panel A), through

ebb tide (panels B and C), to low tide (panel D) and flood tide (panels E and F). The colour

scale represents the number of localizations in 250 x 250 m grid cells. The boundary of

 mudflats are indicated with a grey dashed line (i.e. the Lowest Astronomical Tide, LAT). The

blue line indicates the lowest water level within the different tidal phases. Land is indicated

with a solid black line. Because the tags send a signal at 1/6 Hz, each localization represents

a minimum of 6 s of space use for red knots. See rectangle A in Fig. 1 for placement of this

map within the study area. The coordinate system refers to UTM 31N. © map data from

Rijkswaterstaat.

EXAMPLE 2. AMONG-INDIVIDUAL VARIATION IN MOVEMENT

To illustrate the large-scale application of WATLAS tracking and to explore among-

individual variation in space use, we selected data from seven red knots (out of 44) on 17

October 2020. These tags were programmed to transmit every 6 s. Additionally, we

calculated cumulative distance between successive localizations to reveal variation in daily

distances travelled for all individuals tracked that day.

Red knots were successfully localized in large parts of the study area, though gaps in

the tracks also occurred (Fig. 7). These gaps happened especially where the density of

receivers was low, i.e. when the tag was not within the detection range of at least three

receivers. The tracking data revealed substantial differences among individuals in the

distance travelled over a 24-hour period, which ranged between 42 and 328 km day^{-1} for all

44 birds (mean±SD: 131±49 km d^{-1}, histogram in Fig. 7). There were also differences in the

number of localizations between birds (mean = 2,111 bird^{-1}; range = 144 - 3,680), which

significantly explained distance travelled per day (slope = 26.3 m per localization, p<0.01).

Nonetheless, when dividing the distance travelled by the number of localizations per bird,

there were still large among-individual differences (mean = 83.1 m per localization; range =

27.3 - 316.5), which shows that the among-individual variation in distance travelled is not

merely caused by differences in the number of successful localizations.

Fig. 7 *Tracks for a subset of seven individual red knots that differ in the spatial scale of*

space use across the entire study area. Data collected over 24 hours on 17 October 2020 are

shown. The inset shows the histogram of cumulative distance travelled for all 44 birds

tracked on this day. © map data from Bing, OpenStreetMap.

EXAMPLE 3. FINE SCALE MOVEMENT AND INTRA-SPECIFIC INTERACTIONS

To illustrate the application of high-resolution WATLAS data for investigating social

interactions, we selected seven red knots (out of 79) tracked near Griend (Fig. 1) between two

high tides at 00:50 and 13:16 CEST on 31 August 2018. These data, recorded at 1/3Hz, were

aggregated into 30 s timesteps, and the mean coordinates were calculated. Within these time

steps, social proximity was defined as being within 50 m of each other [70]. The social

network was created in R with the library 'spatsoc' [71].

The fine-scale movement patterns of the red knots confirmed (Fig. 8) that birds

roosted on Griend during high tide and, as the water recedes with the ebb tide, moved out

onto the exposed mudflat to forage. While foraging, the birds walked across the mudflat,

which resulted in areas with dense localizations. Birds flew between different areas to forage

as can be seen by the areas of dense localizations connected by lines with sparser

localizations (flight). An animation of the fine-scale movement with the incoming tide can be

found in the supplement (Additional file 2). Tracking the fine-scale movements of many

tagged animals allows for the investigation of inter-individual interactions. For instance, the

proximity-based social network of our subset of seven red knots revealed that some

individuals were often in close proximity (e.g. birds with tag IDs 409 and 412; Fig. 8),

whereas some individuals were rarely close to the other individuals. The individual with tag

ID 458 was mostly static, hence rarely close to any other tagged individual within this period.

The comparison between the tracks and proximity network further shows the merit of

collecting fine-scale high-resolution tracking data. Visually, for example, the individual with

tag ID 409 seems to have much higher overlap with the individual with tag ID 418 than with

the individual with tag ID 412. Nonetheless, the spatiotemporal proximity network shows that

individuals with tag IDs 409 and 412 have the highest overlap.

Fig. 8 *Detailed movements from a subset of seven tracked individual red knots around*

Griend. Data between two high tides at 00:50 and 13:16 CEST *on 31 August 2018 are*

shown. The inset shows the proximity network for these seven birds based on a spatial

proximity of 50 m (see methods). In total, 79 individuals were localized in this timeframe for

which an animation of their movement relative to the tide can be found in the Supplementary

Information. See rectangle B in Fig. 1 for placement of this map within the study area. © map

data from Bing, OpenStreetMap.

EXAMPLE 4. COMMUNITY TRACKING

Tracking individuals from different species within one region, allows investigations of inter-

specific space use. To show how space use, comprised of individual movements, scales up to

community-level space use, we analyse differences in the home ranges of sanderlings and red

knots. Kernel densities were calculated with the R-library 'amt' [72] for 35 sanderling and 35

red knots tracked simultaneously between 12 and 19 August 2018. More red knots had been

successfully tracked in that period, but to equalize sample sizes with the sanderlings, were

randomly selected. Note that the mean number of localizations in this period for sanderlings

was 1,294 fixes (SD 2,745) and that for red knots was 9,521 fixes (SD 9,868).

The home-range analyses show that although sanderlings and red knots both roost on

Richel and Griend, they differ in their low-tide distribution (Fig. 9). The home range of

sanderlings appeared larger than that of red knots and included intertidal flats near Richel and

extended more to the east of Griend. The differences in space use between sanderlings and

red knots might be related to differences in the behaviour and spatial distribution of their

prey. For instance, red knots forage on patchily distributed and relatively sessile shellfish,

whereas sanderlings forage on shrimp that are mobile and follow the tide.

Fig. 9 *Space use by sanderlings and red knots. The colour scale shows home range estimates*

with kernel densities for A) 35 sanderlings and B) 35 red knots during low tide between 12

and 19 August 2018. The islets of Richel and Griend are roosting sites for these sanderlings

and red knots. Water is coloured blue, land dark grey, and mudflats light grey. See rectangle

A in Fig. 1 for placement of this map within the study area. The coordinate system refers to

UTM 31N. © map data from Rijkswaterstaat.

Discussion

With an array of 26 receiver stations located in the western Dutch Wadden Sea, WATLAS

covers 1,326 km^2 and is currently the largest deployment of an ATLAS tracking system

worldwide. With examples from red knots and sanderlings, we illustrated various

applications of the high spatial and temporal resolution movement data obtained by

WATLAS. Moreover, we provided examples of how high-throughput movement data can be

utilized to study important aspects of animal movement ecology and space use, such as

among-individual variation in behaviour, intra-specific interactions (social networks) as well

as inter-specific interactions (community assembly). For regional-scale studies on small

animals, reverse-GPS systems like ATLAS, are promising.

TECHNICAL CONSIDERATIONS

For successful localization, reverse-GPS tracking like ATLAS, requires at least three

receivers to detect the tag's signal, and signal detection requires a 'line of sight'. A study on

the accuracy of WATLAS localizations, showed that tags (~1.2 m from ground) were mostly

detected by receivers within 5 km of the tag [63]. Near Richel and Griend, where the

distances between receivers were smallest, localizations were most numerous. Near the edges

of the array and on the large-scale of the entire study area, where the distance between

receivers was largest (Fig. 1), tags were localized less often, causing gaps in the tracks (Fig.

7). To avoid missing localizations, the density of receivers can be increased, and the array

should surround the main area of interest [63].

Tags attached to animals in flight generally have larger detection ranges than animals

on the ground, due to their usually greater height. For instance, in another ATLAS system

[11], Egyptian Fruit Bats *Rousettus aegyptiacus* were detected during flight up to 40 km

away from the receivers. In our study system, we recorded similar detection ranges of birds in

flight, and we were able to localise them across the entire study area up to 34 km from the

nearest receiver.

The most substantial costs for reverse-GPS are setting up the initial array of receivers.

Once a reverse-GPS tracking system is in place, the relatively cheap tags allows tracking

large and representative samples of animal populations at low costs [31]. The number of

unique tags ATLAS systems can detect simultaneously is limited, but so far WATLAS has

tracked 232 tags simultaneously without problems. This limit is set by the processing

capability of the computer within the receiver, as well as interference of overlapping

transmissions between tags. The percentage of missed tag transmissions increases

exponentially as a function of the number of transmitters within range (see Fig. 3.5 in [73]).

Note that both limitations are not an intrinsic limitation of ATLAS, but a limitation of the

current implementation. More powerful processors in the receivers will, for instance, allow

more and simultaneous tag detections.

Another advantage of ATLAS systems is the weight of the trackers. So far, the

lightest ATLAS tag (including battery and coating) weighs as little as 0.8 g [42], which

allows tracking small and light-weight individuals that were previously too small to track

remotely at high spatial accuracy. Other tracking systems that allow tracking of smaller free-

living individuals, include MOTUS [36], and light-level geolocation data loggers [74]. These

devices can provide high temporal resolution data or be used to track birds over large areas.

Compared to ATLAS, however, the spatial accuracy of localizations with MOTUS and

geolocation loggers is large (kms) and retrieving data from geolocation loggers requires

recapturing the tracked animals. Recapturing can be problematic and prevents real-time

observations and analyses of tracked animals. Another promising tracking system is ICARUS

[75, 76], but this is under development and tags are estimated to be larger than the lightest

tags of ATLAS.

Compared to GPS systems, reverse-GPS systems have dramatically reduced tag

energy consumption, which allows high temporal resolution localizations to be collected with

small batteries [31] and tags can be expected to last for longer for a given number of

localizations [73]. Because of the eight months maximum lifetime of WATLAS-tags used in

this study, the tags are glued to the backs of birds and will fall off during body moult. From

an ethical perspective this is preferred over e.g. full body harnesses [77], because animals

need to cope with the added weight [78] and potential aerodynamic discomfort only

temporarily [79].

Despite the promise of ATLAS, it is necessary to note some limitations of the system.

Compared to other more global tracking systems, ATLAS is limited to much smaller scales

and therefore is unsuitable for some study designs. Furthermore, the ATLAS system was

developed and is being applied by multidisciplinary teams of scientists for movement ecology

research [41]. Each system has been installed and is maintained by scientists, and often

requires collaborations with landowners and organisations for establishing receiver locations.

This is in stark contrast to plug-and-play tags like many GPS or ARGOS tags. Other

biologgers also collect auxiliary information, such as altitude and acceleration [21]. While

not in use (yet) in our WATLAS system, ATLAS tags have also been successfully fitted with

air-pressure, temperature, and humidity sensors [42] and an on-board accelerometer is under

development. However, remote transfer of data from such sensors remain challenging.

OPPORTUNITIES FOR ANIMAL MOVEMENT RESEARCH AND CONSERVATION

Simultaneously tracking many free-living small birds at high spatial and temporal resolution,

allows for novel studies on e.g., among-individual variation, collective behaviour, and inter-

specific interactions in the wild [60]. Moreover, because individuals from different species

and different trophic levels can be tracked simultaneously, exciting opportunities exist for

studying movement ecology at the community level [80]. Clearly, the list of examples given

here is not exhaustive for what is possible with an ATLAS system like WATLAS. Many

individual or environmental factors can be linked with movement data which can be used to

describe or predict behaviour. Specifically, within the Dutch Wadden Sea, of particular

interest is the ability to combine shorebird tracking with knowledge on their food resources.

In this study, we illustrated differences in home range between sanderlings and red knots.

Because sanderling prefer shrimp and red knot prefer shellfish [55], the differences in space-

use are likely related to differences in the distribution of their preferred prey. Linking space-

use of these birds with the large-scale mapping of benthic resources in collaboration with the

SIBES project [57, 58], will offer exciting opportunities for understanding consumer-resource

interactions and space-use in intertidal ecosystems. Having long-term tracking systems like

WATLAS established also allows researchers to sample behaviour opportunistically. In 2020,

more than 3,000 red knots with a highly pathogenic avian influenza virus were found dead in

the Wadden sea [81]. Avian influenza is a disease of global concern, which can affect both

domestic and wild populations. However, none of the found red knots were tagged with

WATLAS but collecting data on movement and social interactions could provide valuable

insights into disease transmission [82-84].

Despite strong legal protection and management measures being in place, many

activities occur in the Dutch Wadden Sea that can have detrimental effects on its inhabitants

[85], such as commercial fishing, kite surfing, tourism, military exercises, and mining. In

combination with large-scale phenomena, such as sea level rise and global warming [86],

these anthropogenic activities can cause disturbances and habitat destruction, and thus

contribute to population declines [87, 88]. The causes underlying the declines of shorebird

population numbers in the Wadden Sea are often debated, partly because of our limited

understanding of environmental processes and animal space use, which leads to tension and

possibly conflict between stakeholders and management [61, 85, 89, 90]. The development of

WATLAS has opened-up possibilities for quantifying space use of many small shorebirds

directly, automatically, and at high spatiotemporal accuracy. WATLAS could thus aid in

studies of impact assessment on shorebird space use, such as assessing the effects of

disturbances and habitat loss due to land subsidence and sea level rise. More generally,

WATLAS could facilitate evidence-based conservation, and aid the management of this

UNESCO world heritage site.

Conclusions

In this study, we introduced WATLAS as a high-utility tracking system in the Dutch Wadden

Sea, capable of tracking hundreds of small individuals simultaneously at high spatiotemporal

resolution. After the initial investment for an array of receivers (which can be substantial), the

costs per tag are low, which facilitates regional, long-term studies on movement ecology and

space use of many individuals and multiple species and facilitates collaboration between

researchers across research institutes. Additionally, maintenance and operating costs need to

be considered. Because tags are lightweight as well as cheap, WATLAS can facilitate studies

on, for instance, collective behaviour, social information use, and movement ecology of

entire communities of free-living animals. WATLAS can also support evidence-based nature

conservation and management, for example with assessing the impact of anthropogenic

activities on space use of shorebirds. More generally, with WATLAS, animals can function

as sentinels informing us about the state of the Wadden Sea ecosystem [62, 76], and thus aid

nature conservation and management of this globally important ecosystem.

DECLARATIONS

Acknowledgements

Many people and organisations are involved in hosting the WATLAS equipment, without

whom this study would be impossible. We therefore thank Hoogheemraadschap Hollands

Noorderkwartier, Koninklijke Nederlandse Redding Maatschappij, Staatsbosbeheer, Marine

Eco Analytics, Koninklijke Luchtmacht, Het Posthuys, Natuurmonumenten, Wetterskip

Fryslan, Afsluitdijk Wadden Center, Vermilion, Rijkswaterstaat, Carl Zuhorn, Lenze Hofstee

and Lydia de Loos. We thank Natuurmonumenten for access to Griend and using their

facilities. Also, we thank Hein de Vries, Klaas Daalder, Hendrik-Jan Lokhorst, Bram Fey,

Wim-Jan Boon from the RV Navicula and RV Stern, as well as the many other NIOZ staff

and volunteers that facilitated this work. We would particularly like to thank Anita Koolhaas,

Hinke and Cornelis Dekinga for their help with building the receiver stations. We thank Jeras

de Jonge, Martin Laan, Sander Asjes, and Aris van der Vis for their technical help, and

Benjamin Gnep for his beautiful photos and persistently replacing broken LNA's. Thanks to

Marten Tacoma for visualizing the tracking data in real-time on www.nioz.nl/watlas and

Ingrid de Raad for help posting WATLAS-related news. We also thank the Minerva

Foundation and the Minerva Center for Movement Ecology for supporting the development

and maintenance of all ATLAS systems, and for Yotam Orchan and Yoav Bartan for their

most valuable technical assistance. Finally, we thank Theunis Piersma, two anonymous

referees and the associate editor for providing valuable comments and suggestions for

improvements on an earlier version of this manuscript.

Ethics approval and consent to participate

Permits to catch, handle and tag sanderlings and red knots was granted to the NIOZ under

protocol number NIOAVD8020020171505.

Consent for publication

Not applicable

Availability of data and materials

Upon publication, the datasets supporting the conclusions of this article will be included

within the article and its supplementary files.

Competing interests

Not applicable

Funding

This work was partly funded by the Dutch Research Council grand VI.Veni.192.051 awarded

to AIB, and grants from the Israel Science Foundation to RN and ST (ISF grants 965/15 and

1919/19) for supporting the development and maintenance of all ATLAS systems.

Authors' contributions

AIB conceived the idea for the manuscript, analysed the data and led the writing of the

manuscript; RN and ST developed the ATLAS system and provided support throughout data

collection; FvM, BD and AD built and maintained the WATLAS system and tags, and

provided operational support; AIB, AD, LdM, JtH, CEB and RAB deployed the receiver

stations; AIB, AD, EP, SE, PG, LdM, JtH, RAB, and CEB caught and tagged birds; all

authors contributed critically to the drafts and gave final approval for publication.

Additional file information

Additional file 1

Microsoft Word file (.docx)

Supplementary figures S1-S5.

Additional file 2

MPEG animation (.mp4)

Animation of red knot movement near Griend between two consecutive high tides.

Different individuals are indicated with different colours. However, due to the large number

of individuals (N=79), colours can be used more than once between individuals.

References

1.   Kays R, Crofoot MC, Jetz W, Wikelski M. Terrestrial animal tracking as an eye on

life and planet. Science. 2015;348(6240):aaa2478.

2.   Nathan R, Getz WM, Revilla E, Holyoak M, Kadmon R, Saltz D, et al. A movement

ecology paradigm for unifying organismal movement research. Proc Natl Acad Sci USA.

2008;105(49):19052-9.

3.   Tucker MA, Böhning-Gaese K, Fagan WF, Fryxell JM, Van Moorter B, Alberts SC,

et al. Moving in the anthropocene: global reductions in terrestrial mammalian movements.

Science. 2018;359(6374):466-9.

4.   Allen AM, Singh NJ. Linking movement ecology with wildlife management and

conservation. Front Ecol Evol. 2016;3:155.

5.   Hays GC, Bailey H, Bograd SJ, Bowen WD, Campagna C, Carmichael RH, et al.

Translating marine animal tracking data into conservation policy and management. Trends

Ecol Evol. 2019;34(5):459-73.

6.   Gill RE, Tibbitts TL, Douglas DC, Handel CM, Mulcahy DM, Gottschalck JC, et al.

Extreme endurance flights by landbirds crossing the Pacific Ocean: ecological corridor rather

than barrier? Proc R Soc B. 2009;276(1656):447-57.

7.   Lindström Å, Alerstam T, Andersson A, Bäckman J, Bahlenberg P, Bom R, et al.

Extreme altitude changes between night and day during marathon flights of great snipes. Curr

Biol. 2021;31(15):3433-9.

8.   Dickie M, McNay SR, Sutherland GD, Cody M, Avgar T. Corridors or risk?

Movement along, and use of, linear features varies predictably among large mammal predator

and prey species. J Anim Ecol. 2020;89(2):623-34.

9.   Beardsworth CE, Whiteside MA, Laker PR, Nathan R, Orchan Y, Toledo S, et al. Is

habitat selection in the wild shaped by individual-level cognitive biases in orientation

strategy? Ecol Lett. 2021;24:751-60.

10.   Guilford T, Biro D. Route following and the pigeon's familiar area map. J Exp Biol.

2014;217(2):169-79.

11.   Toledo S, Shohami D, Schiffner I, Lourie E, Orchan Y, Bartan Y, et al. Cognitive

map–based navigation in wild bats revealed by a new high-throughput tracking system.

Science. 2020;369(6500):188-93.

12.   Harten L, Katz A, Goldshtein A, Handel M, Yovel Y. The ontogeny of a mammalian

cognitive map in the real world. Science. 2020;369(6500):194-7.

13.   Fortin D, Beyer HL, Boyce MS, Smith DW, Duchesne T, Mao JS. Wolves influence

elk movements: behavior shapes a trophic cascade in Yellowstone National Park. Ecology.

2005;86(5):1320-30.

14.   Strandburg-Peshkin A, Farine DR, Couzin ID, Crofoot MC. Shared decision-making

drives collective movement in wild baboons. Science. 2015;348(6241):1358-61.

15.   Choi C-Y, Peng H-B, He P, Ren X-T, Zhang S, Jackson MV, et al. Where to draw the

line? Using movement data to inform protected area design and conserve mobile species. Biol

Conserv. 2019;234:64-71.

16.   Middleton AD, Sawyer H, Merkle JA, Kauffman MJ, Cole EK, Dewey SR, et al.

Conserving transboundary wildlife migrations: recent insights from the Greater Yellowstone

Ecosystem. Frontiers in Ecology and the Environment. 2020;18(2):83-91.

17.   Pekarsky S, Schiffner I, Markin Y, Nathan R. Using movement ecology to evaluate

the effectiveness of multiple human-wildlife conflict management practices. Biol Conserv.

2021;262:109306.

18.   Murgatroyd M, Redpath SM, Murphy SG, Douglas DJT, Saunders R, Amar A.

Patterns of satellite tagged hen harrier disappearances suggest widespread illegal killing on

British grouse moors. Nat Commun. 2019;10(1):1094.

19.   de Knegt HJ, Eikelboom JAJ, van Langevelde F, Spruyt WF, Prins HHT. Timely

poacher detection and localization using sentinel animal movement. Sci Rep.

2021;11(1):4596.

20.   Preisler HK, Ager AA, Wisdom MJ. Analyzing animal movement patterns using

potential functions. Ecosphere. 2013;4(3):1-13.

21.   Williams HJ, Taylor LA, Benhamou S, Bijleveld AI, Clay TA, de Grissac S, et al.

Optimising the use of biologgers for movement ecology research. J Anim Ecol.

2020;89(1):186-206.

22.   Joo R, Boone ME, Clay TA, Patrick SC, Clusella-Trullas S, Basille M. Navigating

through the R packages for movement. J Anim Ecol. 2020;89:248-67.

23.   Joo R, Picardi S, Boone ME, Clay TA, Patrick SC, Romero-Romero VS, et al. A

decade of movement ecology. arXiv preprint arXiv:200600110. 2020.

24.   Börger L, Bijleveld AI, Fayet AL, Machovsky-Capuska GE, Patrick SC, Street GM,

et al. Biologging special feature. J Anim Ecol. 2020;89(1):6-15.

25.   Nathan R, Monk CT, Arlinghaus R, Adam T, Alós J, Assaf M, et al. Big-data

approaches lead to an increased understanding of the ecology of animal movement. Science.

2022;375(6582):eabg1780.

26.   Hertel AG, Royauté R, Zedrosser A, Mueller T. Biologging reveals individual

variation in behavioural predictability in the wild. J Anim Ecol. 2021;90(3):723-37.

27.   Shaw AK. Causes and consequences of individual variation in animal movement.

Mov Ecol. 2020;8(1):12.

28.   Verhoeven MA, Loonstra A, Senner NR, McBride AD, Both C, Piersma T. Variation

from an unknown source: large inter-individual differences in migrating black-tailed godwits.

Front Ecol Evol. 2019;7:31.

29.   Strandburg-Peshkin A, Farine DR, Crofoot MC, Couzin ID. Habitat and social factors

shape individual decisions and emergent group structure during baboon collective movement.

Elife. 2017;6:e19505.

30.   Bedriñana-Romano L, Hucke-Gaete R, Viddi FA, Johnson D, Zerbini AN, Morales J,

et al. Defining priority areas for blue whale conservation and investigating overlap with

vessel traffic in Chilean Patagonia, using a fast-fitting movement model. Sci Rep.

2021;11(1):2709.

31.   MacCurdy RB, Bijleveld AI, Gabrielson RM, Cortopassi KA. Automated wildlife

radio tracking. In: Zekavat SA, Buehrer RM, editors. Handbook of position location: theory,

practice, and advances. 2nd ed. Hoboken, New Jersey, USA: John Wiley & Sons, Inc.; 2019.

32.   MacCurdy RB, Gabrielson RM, Cortopassi KA. Automated wildlife radio tracking.

In: Zekavat SA, Buehrer RM, editors. Handbook of position location: theory, practice, and

advances. Hoboken, New Jersey, USA: John Wiley & Sons, Inc.; 2011. p. 1129-67.

33.   MacCurdy RB, Gabrielson RM, Spaulding E, Purgue A, Cortopassi KA, Fristrup KM.

Automatic animal tracking using matched filters and time difference of arrival. J Commun.

2009;4:487-95.

34.   Weller-Weiser A, Orchan Y, Nathan R, Charter M, Weiss AJ, Toledo S.

Characterizing the accuracy of a self-synchronized reverse-GPS wildlife localization system.

15th International Conference on Information Processing in Sensor Networks: IEEE; 2016. p.

1-12.

35.   Krüger SW. An inexpensive hyperbolic positioning system for tracking wildlife using

off-the-shelf hardware: North-West University (South Africa), Potchefstroom Campus; 2017.

36.   Taylor P, Crewe T, Mackenzie S, Lepage D, Aubry Y, Crysler Z, et al. The Motus

wildlife tracking system: a collaborative research network to enhance the understanding of

wildlife movement. Avian Cons Ecol. 2017;12(1):8.

37.   Kays R, Tilak S, Crofoot M, Fountain T, Obando D, Ortega A, et al. Tracking animal

location and activity with an automated radio telemetry system in a tropical rainforest. The

Computer Journal. 2011;54(12):1931-48.

38.   CLS. Argos User's Manual. 2015 last accessed: 28 march 2022. available at

http://www.argos-system.org/.

39.   Fisher KE, Dixon PM, Han G, Adelman JS, Bradbury SP. Locating large insects using

automated VHF radio telemetry with a multi-antennae array. Methods Ecol Evol.

2021;12(3):494-506.

40.   Rutz C, Morrissey MB, Burns ZT, Burt J, Otis B, St Clair JJ, et al. Calibrating

animal☐borne proximity loggers. Methods Ecol Evol. 2015;6(6):656-67.

41.   Toledo S, Kishon O, Orchan Y, Shohat A, Nathan R. Lessons and experiences from

the design, implementation, and deployment of a wildlife tracking system. 15th International

Conference on Information Processing in Sensor Networks2016. p. 51-60.

42.   Toledo S, Mendel S, Orchan Y, Bartan Y, Margalit S, Levi A, et al. Vildehaye: A

family of versatile, widely-applicable, and field-proven lightweight wildlife tracking and

sensing tags. 2022 Proceedings of the International Conference on Information Processing in

Sensor Networks (IPSN); Milan, Italy (unpublished).

43.   Ersoy S, Beardsworth CE, Dekinga A, van der Meer M, Piersma T, Groothuis TG, et

al. Exploration speed in captivity predicts foraging specialisation in free-living red knots

Calidris canutus. J Anim Ecol. 2022;91:356-66.

44.   Xia H, Bertoni HL, Maciel LR, Lindsay-Stewart A, Rowe R. Radio propagation

characteristics for line-of-sight microcellular and personal communications. IEEE

Transactions on antennas and propagation. 1993;41(10):1439-47.

45.   Aarts G, Brasseur S, Poos JJ, Schop J, Kirkwood R, van Kooten T, et al. Top-down

pressure on a coastal ecosystem by harbor seals. Ecosphere. 2019;10(1):e02538.

46.   van der Veer HW, Dapper R, Henderson PA, Jung AS, Philippart CJ, Witte JIJ, et al.

Changes over 50 years in fish fauna of a temperate coastal sea: Degradation of trophic

structure and nursery function. Estuar Coast Shelf S. 2015;155:156-66.

47.   Beukema J, Dekker R. Half a century of monitoring macrobenthic animals on tidal

flats in the Dutch Wadden Sea. Mar Ecol Prog Ser. 2020;656:1-18.

48.   van Roomen M, Laursen K, van Turnhout C, van Winden E, Blew J, Eskildsen K, et

al. Signals from the Wadden sea: population declines dominate among waterbirds depending

on intertidal mudflats. Ocean Coast Manag. 2012;68:79-88.

49.   van de Kam J, Ens B, Piersma T, Zwarts L. Shorebirds. An illustrated behavioral

ecology. Utrecht, The Netherlands: KNNV Publishers; 2004.

50.   Allen AM, Ens BJ, Van de Pol M, Van der Jeugd H, Frauendorf M, Oosterbeek K, et

al. Seasonal survival and migratory connectivity of the Eurasian Oystercatcher revealed by

citizen science. The Auk. 2019;136(1).

51.   Rakhimberdiev E, Duijns S, Karagicheva J, Camphuysen CJ, Dekinga A, Dekker R,

et al. Fuelling conditions at staging sites can mitigate Arctic warming effects in a migratory

bird. Nat Commun. 2018;9(1):1-10.

52.   Bijleveld AI, MacCurdy RB, Chan Y-C, Penning E, Gabrielson RM, Cluderay J, et al.

Understanding spatial distributions: negative density-dependence in prey causes predators to

trade-off prey quantity with quality. Proc R Soc B. 2016;283(1828):20151557.

53.   van Gils JA, Spaans B, Dekinga A, Piersma T. Foraging in a tidally structured

environment by red knots (*Calidris canutus*): ideal, but not free. Ecology. 2006;87(5):1189-

202.

54.   Bakker W, Ens BJ, Dokter A, van der Kolk H-J, Rappoldt K, van de Pol M, et al.

Connecting foraging and roosting areas reveals how food stocks explain shorebird numbers.

Estuar Coast Shelf S. 2021;259:107458.

55.   Piersma T, Hoekstra R, Dekinga A, Koolhaas A, Wolf P, Battley PF, et al. Scale and

intensity of intertidal habitat use by knots *Calidris canutus* in the Western Wadden Sea in

relation to food, friends and foes. Neth J Sea Res. 1993;31:331-57.

56.   Zwarts L, Blomert A-M, Wanink JH. Annual and seasonal variation in the food

supply harvestable by knot *Calidris canutus* staging in the Wadden Sea in late summer. Mar

Ecol Prog Ser. 1992:129-39.

57.   Bijleveld AI, van Gils JA, van der Meer J, Dekinga A, Kraan C, van der Veer HW, et

al. Designing a benthic monitoring programme with multiple conflicting objectives. Methods

Ecol Evol. 2012;3(3):526-36.

58.   Compton TJ, Holthuijsen S, Koolhaas A, Dekinga A, ten Horn J, Smith J, et al.

Distinctly variable mudscapes: distribution gradients of intertidal macrofauna across the

Dutch Wadden Sea. J Sea Res. 2013;82:103-16.

59.   Oudman T, Piersma T, Ahmedou Salem MV, Feis ME, Dekinga A, Holthuijsen S, et

al. Resource landscapes explain contrasting patterns of aggregation and site fidelity by red

knots at two wintering sites. Mov Ecol. 2018; 6(1):24.

60.   King AJ, Fehlmann G, Biro D, Ward AJ, Fürtbauer I. Re-wilding collective

behaviour: an ecological perspective. Trends Ecol Evol. 2018;33(5):347-57.

61.   Boere GC, Piersma T. Flyway protection and the predicament of our migrant birds: a

critical look at international conservation policies and the Dutch Wadden Sea. Ocean Coast

Manag. 2012;68:157-68.

62.   Piersma T, Lindström Å. Migrating shorebirds as integrative sentinels of global

environmental change. Ibis. 2004;146:61-9.

63.   Beardsworth CE, Gobbens E, Maarseveen Fv, Denissen B, Dekinga A, Nathan R, et

al. Validating a high-throughput tracking system: ATLAS as a regional-scale alternative to

GPS. bioRxiv. 2021.

64.   Melamed I, Toledo S. A robust, selective, and flexible RF front-end for wideband

sampling receivers. ICT Express. 2017;3(2):96-100.

65.   Naef-Daenzer B, Widmer F, Nuber M. A test for effects of radio-tagging on survival

and movements of small birds. Avian Science. 2001;1(1):15-23.

66.   Kenward RE. A manual for wildlife radio tagging: Academic press; 2001.

67.   Dietz MW, Rogers KG, Piersma T. When the seasons don't fit: speedy molt as a

routine carry-over cost of reproduction. PLoS One. 2013;8(1):e53890-e.

68.   Gupte PR, Beardsworth CE, Spiegel O, Lourie E, Toledo S, Nathan R, et al. A guide

to pre-processing high-throughput animal tracking data. J Anim Ecol. 2022;91:287-307.

69.   R Core Team. R: a language and environment for statistical computing. R foundation

for statistical computing, Vienna, Austria. URL https://www.R-project.org2020.

70.   Farine DR, Whitehead H. Constructing, conducting and interpreting animal social

network analysis. J Anim Ecol. 2015;84(5):1144-63.

71.   Robitaille AL, Webber QMR, Vander Wal E. Conducting social network analysis

with animal telemetry data: applications and methods using spatsoc. Methods Ecol Evol.

2019;10(8):1203-11.

72.   Signer J, Fieberg J, Avgar T. Animal movement tools (amt): R package for managing

tracking data and conducting habitat selection analyses. Ecol Evol. 2019;9(2):880-90.

73.   MacCurdy RB, Bijleveld AI, Gabrielson RM, Cluderay J, Spaulding EL, Oudman T,

et al. Automatic, intensive wildlife radiotracking. Untying the knot: Mechanistically

understanding the interactions between social foragers and their prey2015. p. 41-56.

74.   Bridge ES, Kelly JF, Contina A, Gabrielson RM, MacCurdy RB, Winkler DW.

Advances in tracking small migratory birds: a technical review of light-level geolocation. J

Field Ornithol. 2013;84(2):121-37.

75.   Wikelski M, Kays RW, Kasdin NJ, Thorup K, Smith JA, Swenson GW. Going wild:

what a global small-animal tracking system could do for experimental biologists. J Exp Biol.

2007;210(2):181-6.

76.   Jetz W, Tertitski G, Kays R, Mueller U, Wikelski M, Åkesson S, et al. Biological

earth observation with animal sensors. Trends Ecol Evol. 2022;37(4):293-8.

77.   Chan Y-C, Brugge M, Tibbitts TL, Dekinga A, Porter R, Klaassen RH, et al. Testing

an attachment method for solar-powered tracking devices on a long-distance migrating

shorebird. J Ornithol. 2016;157(1):277-87.

78.   Portugal SJ, White CR. Externally attached biologgers cause compensatory body

mass loss in birds. Methods Ecol Evol. 2021;00:1-9.

79.   Bowlin MS, Henningsson P, Muijres FT, Vleugels RHE, Liechti F, Hedenström A.

The effects of geolocator drag and weight on the flight ranges of small migrants. Methods

Ecol Evol. 2010;1(4):398-402.

80.   Schlägel UE, Grimm V, Blaum N, Colangeli P, Dammhahn M, Eccard JA, et al.

Movement☐mediated community assembly and coexistence. Biol Rev. 2020;95(4):1073-96.

81.   EFSA, ECDC, EURL, Adlhoch C, Fusaro A, Gonzales JL, et al. Avian influenza

overview December 2020 – February 2021. EFSA Journal. 2021;19(3):e06497.

82.   Fortuna MA, Popa-Lisseanu G, Ibanez C, Bascompte J. The roosting spatial network

of a bird-predator bat. Ecology. 2009;90(4):934-44.

83.   Altizer S, Bartel R, Han BA. Animal migration and infectious disease risk. Science.

2011;331(6015):296-302.

84.   van Dijk JGB, Hoye BJ, Verhagen JH, Nolet BA, Fouchier RAM, Klaassen M.

Juveniles and migrants as drivers for seasonal epizootics of avian influenza virus. J Anim

Ecol. 2014;83(1):266-75.

85.   Kabat P, Bazelmans J, van Dijk J, Herman PMJ, van Oijen T, Pejrup M, et al. The

Wadden Sea Region: Towards a science for sustainable development. Ocean Coast Manag.

2012;68:4-17.

86.   van de Pol M, Ens BJ, Heg D, Brouwer L, Krol J, Maier M, et al. Do changes in the

frequency, magnitude and timing of extreme climatic events threaten the population viability

of coastal birds? J Appl Ecol. 2010;47(4):720-30.

87.   Kraan C, van Gils JA, Spaans B, Dekinga A, Bijleveld AI, van Roomen M, et al.

Landscape-scale experiment demonstrates that Wadden Sea intertidal flats are used to

capacity by molluscivore migrant shorebirds. J Anim Ecol. 2009;78(6):1259-68.

88.   van Gils JA, Piersma T, Dekinga A, Spaans B, Kraan C. Shellfish dredging pushes a

flexible avian top predator out of a marine protected area. PLoS Biol. 2006;4(12):2399-404.

89.   Wolff WJ. The exploitation of living resources in the Dutch Wadden Sea: a historical

overview. Helgol Mar Res. 2005;59(1):31-8.

90.   Floor JR, van Koppen CSA, Lindeboom HJ. A review of science–policy interactions

in the Dutch Wadden Sea — The cockle fishery and gas exploitation controversies. J Sea

Res. 2013;82:165-75.