Nuclear progesterone receptor regulates blood coagulation facilitating ovulation in zebrafish

Jing Huang [a], Chao Sun [a], Nan Nan Zhao [a], Jordan A. Shavit [b], Yong Zhu [a, c, *], Shi Xi Chen [a, d, *]

[a] State Key Laboratory of Marine Environmental Science, College of Ocean and Earth Sciences, Xiamen University, Xiamen 361102, China
[b] Division of Pediatric Hematology/Oncology, Department of Pediatrics, University of Michigan, Ann Arbor, MI 48109, USA
[c] Department of Biology, East Carolina University, 1000 E 5th Street, Greenville, NC 27858, USA
[d] State-Province Joint Engineering Laboratory of Marine Bioproducts and Technology, Xiamen University, Xiamen 361102, China

* Corresponding authors:
Shi Xi Chen, College of Ocean and Earth Sciences, Xiamen University, Xiamen 361102, China. Phone: +86-13515969358, E-mail address: chenshixi@xmu.edu.cn.
Yong Zhu, Department of Biology, East Carolina University, 1000 E 5th Street, Greenville, NC 27858, USA. Phone: 252-328-6504, E-mail address: zhuy@ecu.edu.

Author Contributions:
JH performed most experiments, analyzed the data, and wrote the draft with the assistance of CS and NNZ. JAS generated the *f5* zebrafish knockout, provided protocols and experimental suggestions, discussed experimental results, and revised the manuscript. YZ supervised the project, conducted the fertility experiment, and revised the manuscript. SXC conceived and supervised the project and revised the manuscript.

Competing Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial conflicts.

Classification: Biological Sciences Physiology

Keywords: Pgr, coagulation, ovulation, metalloproteases,

This PDF file includes:

Main Text
Figures 1 to 10

Abstract

Ovulation is a remodeling process including blood capillary rupture and coagulation. Until now, there is no regulation and functional studies of coagulation factors in ovulation. Here, we report dramatic increases of coagulation factors (*f5, f3a*) in zebrafish preovulatory follicles. This upregulation was induced by progestin (DHP: 17α, 20β-dihydroxy-4-pregnen-3-one), a native ligand for nuclear progestin receptor (Pgr) that is essential for ovulation in zebrafish; but was abolished in $pgr^{-/-}$. In addition, promoter activities of *f5* and *f3a* were significantly enhanced by progestin via zebrafish Pgr. Similarly, we found promoter activities of human *F5* were significantly stimulated by progesterone (P4) via human PGRB. Moreover, a dramatic increase of erythrocyte numbers in capillaries on ovarian follicles was associated with ovulation. Importantly, heparin, an anticoagulant, inhibited ovulation. Furthermore, reduced fecundity and impaired ovulation were observed in $f5^{+/-}$ female zebrafish. Together, our results provide plausible evidence for an exceptional function of coagulation factors in ovulation.

Significance Statement

Precise activation of metalloprotease is essential for ovulation. This activation is controlled by serum-derived protease inhibitors. However, the association between coagulation factors and ovulation is unclear. We demonstrated that nuclear progestin receptor (Pgr), a well-established initiator for ovulation, was a major regulator for a dramatic increase of coagulation factors (*f5* & *f3a*) in preovulatory follicles of zebrafish. Blood coagulation in the capillaries of preovulatory follicles occurred prior to follicular rupture in wildtype zebrafish. Interestingly, this blood coagulation was blocked in Pgr knockout fish, which can't ovulate. Inhibition of blood coagulation significantly reduced ovulation. Decreased fertility was also found in F5 heterozygous zebrafish. Our findings reveal critical roles of steroid receptor-regulated coagulation factors in blood coagulation that controls ovulation.

1. Introduction

Ovulation is a tissue remodeling process involving the rupture of follicular layers, basal membrane, and many blood capillaries intertwined in the follicular cells. This dramatic remodeling process share features with inflammatory responses [1]. Blood coagulation is part of the innate defense in response to the inflammation during ovulation [2]. However, reports that imply links between blood coagulation and ovulation are still rare. Formatting of blood clots at apical vessels was found shortly before follicle rupture in animals [1, 3, 4, 5]. Stop of the blood flow likely due to blood clots were observed in apex vessels of ovarian follicles during the late ovulatory phase in women [6]. In addition, an increase of activity and expression of thrombin (a protease essential for fibrin formation) and its receptors were also reported in luteinized granulosa cells and follicular fluid in human [7, 8]. Increased fibrinogen secretion in bovine granulosa cells was thought to be related to the proteolytic activity needed for follicle rupture [9]. Circulating F3 (also known as tissue factor) was elevated in women with polycystic ovary syndrome that is characterized by chronic oligo-ovulation or anovulation [10]. However, the mechanism or contribution of the blood coagulation factor toward ovulation is still unknown [9].

Progesterone (P4) and its nuclear progesterone receptor (PGR) are well-established initiators for vertebrate ovulation [11, 12, 13, 14, 15, 16]. Recently, we identified coagulation factors *f3a* and *f5* as potential downstream targets of Pgr in a transcriptomic analysis of preovulatory follicles from zebrafish [17]. Interestingly, changes in these coagulation factors during ovulation are conserved among human, mice, and zebrafish [17, 18, 19]. Possible involvements of P4 or PGR in the regulation of coagulation have been hinted from several studies using progestin contraceptives in human. F3 was upregulated in endometriosis and

at the endometrial bleeding sites in women using long-term progestin-only contraception

[20]. Orally administered progesterone reduced spontaneous miscarriages and improved

fertility among women who had a history of recurrent miscarriages due to abnormal

bleeding in early pregnancy [21]. Increased risk of venous thromboembolic events was

associated with combined oral contraceptives, particularly those containing the progestin

desogestrel [22, 23, 24]. A recent genetic study indicated that the present-day carriers of the

Neandertal variant of PGR had less bleeding during early pregnancy because of high levels

of PGR expression [25]. However, evidence of P4 induced or PGR mediated regulation of

coagulation factors, especially in ovarian follicles during ovulation, is lacking.

In this study, we examined the expressions, hormonal regulations, and functions of

f5 and *f3a* genes in preovulatory follicles of zebrafish. The gene of *f3* was duplicated in

zebrafish (*f3a* and *f3b*) [26]. We first demonstrated the expression and localization of *f5, f3a,*

and *f3b* in preovulatory follicular cells. We found that progestin (DHP, 17α, 20β-

dihydroxy-4-pregnen-3-one, a native ovulation-inducing ligand in zebrafish) via Pgr

regulated *f5* and *f3a* expression. We also found that follicular cells with strong Pgr

expression were adjacent to blood capillaries, and inhibition of blood coagulation reduced

ovulation. Finally, fertility and ovulation were reduced in $f5^{+/-}$ zebrafish mutants *in vivo*.

Our study provides credible evidence that coagulation factors (*f5* & *f3a*) in preovulatory

follicular cells are regulated by progestin (DHP) via Pgr, which are well-known upstream

initiators for ovulation. Our results also suggest a previously unknown role of these

coagulation factors in ovulation and female fertility.

2.   Results

2.1 Dramatic increase of *f5* an*d f3a* transcripts in follicular cells of preovulatory follicles during a natural spawning cycle *in vivo*

Expression of *f5* and *f3a* mRNA was exclusively restricted to the follicular cells of mature follicles (IVb) and was not found in the germ cells (oocytes) (Fig. 1B). In contrast, *f3b* mRNA was expressed in both follicular cells and oocytes. The expression of *f5, f3a*, and *f3b* in mature follicles (IVb), just prior to ovulation, collected at 06:40 were ~120-fold, ~2-fold, and ~3,145-fold higher than those in immature but fully-grown follicles (IVa, collected at 12:00, Fig. 1C). Since the increase in *f3b* expression was modest, we focused the rest of our studies on *f5* and *f3a*. Results from WISH showed that both *f5* and *f3a* were nearly undetectable in immature follicles (IVa, Fig. 1D, E). In contrast, strong signals of both *f5* and *f3a* transcripts were detected in the follicular cells of the mature follicles (IVb) with the antisense probes (Fig. 1D, E), while no signal was detectable using sense probes (data not shown).

2.2 Progestin via Pgr upregulates *f5* and *f3a* transcripts

2.2a Transcripts of *f5* and *f3a* were upregulated mainly by progestin in preovulatory follicles

The progestin (DHP) is a well-established native ligand for Pgr and an upstream hormonal inducer for ovulation in zebrafish [14, 32, 37]. As predicted, the increase of *f5* and *f3a* transcripts were dose- and time-dependent in fully grown immature follicles (IVa)

exposed to external progestin (Fig. 2A-D). Both late vitellogenesis follicles (III, ~7 fold)

and fully-grown immature follicles (IVa, ~337 fold) responded to progestin signal to

increase of *f5* transcripts (Fig. 3A). In comparison, only fully-grown immature follicles

(IVa) were capable to respond to the progestin signal to increase *f3a* transcripts, but not the

early developing follicles (stage I-III, Fig. 3B). LH is the primary endocrine factor for

inducing ovulation in all animals including zebrafish [38]. As expected, increased *f5*

expression (~ 4 fold) was observed in IVa immature follicles exposed to hCG (50 IU/mL,

Fig. 3C), but not for *f3a* (Fig. 3D). In comparison, progestin (DHP) had much stronger

stimulatory effects on the expression of *f5* (~ 318 fold) and *f3a* (~ 10 fold) than hCG (Fig.

3C, D).

Significantly reduced *f5 and f3a* expression was found in mature preovulatory

follicles (IVb) collected from *pgr*[-/-] fish compared to those in wildtype siblings (Fig. 3E).

Importantly, DHP-induced *f5* and *f3a* expression were blocked in fully-grown immature

follicles (IVa) from *pgr*[-/-] fish (Fig. 3F).

2.2b Progestin via Pgr enhances promoter activity of *f5* and *f3a*

As expected [32], progestin-induced dose-dependent, Pgr-mediated activation of the

MMTV promoters (Fig. 4A). In the presence of Pgr, progestin significantly increased the

f5 and f3a promoter activities, and the increase was dose-dependent (Fig. 4B, C). No

change of promoter activity was observed when pcDNA3.1(+)-Pgr was replaced by a

control vector (pcDNA3.1(+), Fig. 4B, C). The fold increase for *f5* was much bigger than

that of the *f3a*. In the presence of human PGRB, P4 also significantly increased human *F5*

promoter activity, and the increase was dose-dependent. No change of promoter activity

was observed when pCMV-HA-hPGRB was replaced by a control vector (pCMV-HA, Fig.

4D)

2.3 Coagulation factors participate in hemostasis and ovulation

2.3a Follicular cells with strong Pgr expression localize adjacent to blood vessels

Intriguingly, we found almost all the follicular cells with strong Pgr expression

were located adjacent to capillary vessel networks on the surface of the IVa follicles in

transgenic zebrafish line *Tg(pgr:eGFP/ fli1:DsRed)* (Fig. 5).

2.3b Rupture of blood vessels during ovulation

The ovulation process started with the appearance of a small hole in follicular cells

and a ruptured blood vessel (indicated by a red arrow in Fig. 6A, supplemental video clip

A) with a concomitant appearance of the ovulating oocyte on the surface of the ovaries.

When the hole of the broken follicular cell layer reached a certain size, the oocyte was

quickly released from the follicular layer (Fig. 6B-E, supplemental video clip A).

2.3c Increased numbers of erythrocytes in the lumen of blood vessels of mature

preovulatory follicles as ovulation proceeds

The numbers of erythrocytes in capillaries of mature preovulatory follicles (IVb)

increased as they proceeded to follicular rupture, from ~06:40, (Fig. 7A, D, C) to ~07:00

(Fig. 7B, E, G). This effect was absent in the same mature follicles (IVb) collected from

anovulatory $pgr^{-/-}$ female siblings around 07:00 (Fig. 7C, F, G).

2.3d Heparin inhibits ovulation *in vitro*

The intact IVb follicles or ovulated follicles from *Tg(fli1:eGFP)* were easily

recognizable by the interconnecting blood vessels with strong GFP expression and

dramatic changes in size (Fig. 8A-D). As expected, the ovulation rate decreased to 41 %

when follicles were exposed to EDTA (2 mM), an inhibitor for metalloproteases that are

required for ovulation [36]. Intriguingly, heparin (100 µg/mL), an anticoagulant, significantly

reduced the rate of ovulation to 66 % (Fig. 8E). Another anticoagulant, warfarin (50

µg/mL), had a reduced but not significant adverse effect on ovulation rate (83 %) (Fig. 8E).

2.3e Reduced fertility in $f5^{+/-}$ zebrafish females *in vivo*

Although *f5* homozygous mutants live to early adulthood, most die by 4-6 months

of age and do not breed well [39]. Therefore, we examined female fertility in $f5^{+/-}$ mutants in

a consecutive spawning test. We observed significantly reduced numbers of embryos per

spawning and total embryos over two weeks in comparison to wildtype sibling females.

(Fig. 9A, B). The spawning frequency of $f5^{+/-}$ female fish also significantly decreased

compared to that of wildtype siblings (Fig. 9C). One $f5^{+/-}$ female only spawned once in two

weeks (Fig. 9C). The spawning intervals of $f5^{+/-}$ females were significantly longer in

comparison to that of wildtype siblings (Fig. 9D).

3.   Discussion

We have provided comprehensive evidence demonstrating that the expression of

coagulation factors *f5* and *f3a* in preovulatory follicles is controlled by progestin and its

cognate receptor Pgr. Increased expression of these coagulation factors is likely critical for

the regulation of ovulation and female fertility in addition to their well-established roles in

hemostasis. Our conclusions are based on the following evidence: 1) Coagulation factors,

f5 and *f3a*, are expressed in the follicular cells and increase dramatically prior to ovulation

during natural spawning cycles in zebrafish; 2) Expression of these cofactors in

preovulatory follicles is controlled by progestin via Pgr, which are essential and upstream

regulators for ovulation; 3) Follicular cells with strong Pgr expression are adjacent to blood

capillaries; 4). As follicular cells near rupture, a significant increase in the numbers of

erythrocytes in blood vessels suggests hemostasis which is correlated with a dramatic

increase of *f5* and *f3a* in preovulatory follicular cells; 5) Inhibition of coagulation leads to

reduced ovulation; 6) Low fertility is found in $f5^{+/-}$ zebrafish female mutants. Thus, we

provided an ovulation model that summarized findings of our lab as well as those from

others (Fig.10)

3.1 Progestin via Pgr regulates coagulation factors

A major finding of the present study is the regulation of *f5* and *f3a* by progestin via

Pgr in the preovulatory follicles during ovulation (Fig.10). F3 is expressed in a wide array

of tissues [40]. It has been demonstrated that physiological concentrations of P4 could

increase both protein and mRNA of F3 in breast cancer cell lines ZR-75 and T47D [41].

Increased F3 expression was correlated with an increase of procoagulant activity, which

was hypothesized to be responsible for the increased risk of thrombosis in breast cancer

patients [42]. Studies of the uterus have also shown that RU486, an antagonist of PGR, not

only blocks but also reverses progestin-enhanced expression of F3 mRNA and protein in

stromal cells [43]. In line with these studies, PGR binding sites were identified in the promoter

region of the human F3 gene (Supplemental Table 5) [44]. The increase of *f3a* expression in

the preovulatory follicles is primarily controlled by Pgr (current study). A difference in the

weak activity of *f3a* promoters in a human cell line in comparison to progestin-induced

strong *f3a* promoter activities found in zebrafish follicles *in vitro* may be due to missing

core *cis*-regulatory elements in the cloned promoter sequence of zebrafish *f3a*

(Supplemental Table 5). Nevertheless, our results suggest that *f3a* expression is controlled

by progestin via Pgr (Supplemental Table 5, Fig.10).

Until now, no information is available for progestin regulation of F5. Estrogen and

estrogen receptors regulate proteins and transcripts of *F5* in mice bone marrow/bone [45, 46].

The present study demonstrates that a new extrahepatic site of *f5* expression is regulated

by progestin through Pgr. Several putative Pgr binding sites were identified in promoter

regions of human, mouse, and zebrafish *F5* (Supplemental Table 6). Moreover, progestin

enhances promoter activities of *f5* in zebrafish and F5 in humans via PGR. Whether P4

and/or PGR regulate *F5* expression in the human ovary will be an interesting topic for

further investigation.

3.2 Coagulation factor regulate hemostasis and ovulation

Functions and mechanisms of coagulation factors in hemostasis are well established

[47, 48, 49]. However, the role of hemostasis that occurred during ovulation had been

overlooked. Formation of blood clots at apical blood vessels was found shortly before

follicular rupture in several mammalian species [1, 3, 4, 5, 6]. In addition, the permeability of

blood vessels increased 3-fold prior to ovulation in rats [50], which likely facilitates material

exchanges between the follicles and blood vessels thus anticipating hemostasis. We

observed ruptured blood vessels on the surface of ovulating follicles. In addition, we also

found a significant increase of erythrocytes in capillaries on the surface of ovulating

follicles, indicative of hemostasis, as follicles progress toward rupture/ovulation. Lack of

functional F5 or F8 caused various bleeding symptoms, including recurrent and serious

ovulation-related blood losses in women [51]. Occurrence of hemostasis in advance of follicle

rupture could preemptively stop excessive bleeding caused by ovulation-induced capillary

ruptures. The granulosa cells appear to be primary cells for synthesizing F3a and F5 as

these cells account for most cells that express Pgr (this study & [11, 37]). Intriguingly, we also

found that strong Pgr expressing theca cells were near capillaries on the follicles. Whether

these theca cells are key switchers of F3a or F5-initiated hemostasis in the capillaries

around ovarian follicles will be an interesting topic to examine in the future.

Activation of multiple metalloproteases within the ovulating follicles at the right

times is a critical step for sequential digestion of extracellular matrix proteins of the

follicular cells that are required for ovulation [1, 38]. Metalloproteases are synthesized in a

latent form (pro-enzyme) [52]. Their activation and substrate binding are precisely regulated

by the timing of prodomain cleavage of the enzyme and removal of endogenous tissue

inhibitors of metalloproteases (Timps). Protease inhibitors (Timps and other inhibitors) are

typically present at high concentrations in the serum to prevent metalloprotease from acting

accidentally [53]. Vasoconstriction of blood vessels within the thecal layer at the apex of the

follicles was suggested to be critical for the removal of the protease inhibition from the

circulation [54]. We hypothesize that coagulation that occurs shortly before ovulation in the

follicles may have a similar function. In addition to the removal of protease inhibitors from

serum, the cleavage of propeptide from laten form of metalloproteinase is required for

metalloprotease activation and thus ovulation (Fig.10). An F3-dependent pathway of

thrombin generation was also shown in bovine, equine, and human follicular fluid [7, 55, 56,

57]. Granulosa cells were shown to express and secrete fibrinogen, which is catalyzed by

thrombin to become fibrin [9]. Fibrin formation could stimulate activities of

plasmin/plasminogen activators (plat and plau) signal, which is well-established for

plasminogen (plg) activation [58]. Furthermore, the function of Plat, Plau, and Plg in

metalloproteinase activations during ovulation is also recognized [59, 60]. Therefore, we

hypothesize that a dramatic increase of F3a and F5 expression in the follicular cells may

trigger the formation of fibrin, followed by the activation of Plat, Plau, and Plg, which

further activate the metalloproteinases. Taken together, as blood coagulation occurs, blood

flow decreases, then stops completely, effectively terminating inhibitory effects of protease

inhibitors from the serum, triggering metalloprotease activations to start the degradation of

the follicular extracellular matrix for preparing the release of a mature oocyte (Fig.10). So

far, a few reports are available suggesting a link between coagulation factors and the

ovulation process. A patient diagnosed with severe F5 deficiency had to undergo both

ovariectomies to remove both ovaries because of the presence of hemorrhagic anovulatory

follicles and a life-threatening hemorrhage in the pouch of Douglas [61]. Circulating F3 was

elevated in women with polycystic ovary syndrome that is the most common form of

anovulatory infertility [10]. Inconsistent with findings from medaka [36], the rate of ovulation

was suppressed by EDTA treatment in the current study likely via inhibition of

metalloproteases. Heparin, a known anticoagulant, also suppressed ovulation. Another

finding in the present study is that reduced fecundity and impaired ovulation were observed

in $f5^{+/-}$ female zebrafish *in vivo*. Together, these results argue strongly for conserved

functions of coagulation factors in the regulation of ovulation in vertebrate ovaries.

In summary, our results demonstrate that *f5* and *f3a* expression are dramatically

induced by progestin via a Pgr-dependent mechanism in preovulatory follicles (Fig. 10).

Moreover, these coagulation factors regulate ovulation and female fertility in addition to

their roles in hemostasis. Future studies on the regulation and functions of coagulation

factors in other organisms including mammalian follicles or ovarian fluid during ovulation

likely lead to developments of diagnostic and therapeutic modalities for patients with

gynecological problems associated with mutations and/or dysregulation of coagulation.

4. Materials and Methods

4.1 Animals & Reagents

Zebrafish used are offspring of Tübingen strain unless stated otherwise. Detailed information for knockouts, transgenic lines, key chemicals, and reagents are listed in Supplemental Table 1 & Table 2. Fish were fed three times daily with a commercial food (Otohime B2; Marubeni Nisshin Feed, Tokyo, Japan), supplemented with newly hatched brine shrimp, and were maintained in recirculating system (ESEN, Beijing, China) at $28 \pm 0.5\,°C$ under a controlled photoperiod (lights on at 08:00, off at 22:00). The daily ovulation cycle in mature adult zebrafish is established according to an established protocol [27]. Briefly, each mature adult female and male zebrafish were paired in a spawning tank. These fish typically start spawning soon after lights on in the morning. Around noon, fish water was replaced, and fish were fed at least three times daily with the commercial food and supplemented with newly hatched brine shrimp. Around one hour prior to lights off, fish water was replaced again, a mesh inner tank was inserted into each spawning tank to prevent cannibalism typical for zebrafish adults. After one week, most fish form a daily ovulation cycle. When applicable, siblings from heterozygous in-cross were used for comparison. All experimental protocols were approved by the Institutional Animal Care and Use Committee (IACUC) at Xiamen University.

4.2 Collection of fully-grown follicles (stage IV) during a daily spawning cycle

Fully-grown immature follicles (stage IVa, >650 μm, with visible germinal vesicles, i.e., GV, appears opaque) or mature follicles (stage IVb, GV disappears, appears

transparent) were collected at four representative timepoints during a daily spawning cycle

(Fig. 1A): 12:00 (IVa follicles); 20:00 (IVa follicles); 05:00 (the onset of oocyte maturation,

IVa follicles); and 06:40 (completion of oocyte maturation but prior to ovulation, IVb

follicles). To be simple and accurate in our description, we will use "IVa" to indicate fully-

grown immature stage IVa follicles; "IVb" to indicate mature but yet ovulated follicles

(prior to ovulation) throughout this manuscript.

To sacrifice fish humanely, the spinal cord and blood supply behind the gill cover

were cut off swiftly using sharp scissors following a lethal dose of anesthetization (MS-

222, 300 mg/L buffered solutions). Ovaries were immediately placed in 60 % L-15 media

containing 15 mM HEPES at pH 7.2. Individual follicles were obtained according to an

established procedure [27, 28]. At each timepoint, a total of 25 IVa or IVb follicles (containing

both germ cell and surrounding follicular cells) from 4 females were pooled as one sample.

In addition, the follicular cells (~100 follicles) or their enclosed oocytes (~20 follicles)

were collected separately from IVb follicles according to an established method [29]. These

samples were immediately homogenized in RNAzol solution (Molecular Research Center,

Cincinnati, USA) and stored at a -80 ℃ freezer until further processing.

4.3 Hormone treatments

According to established follicle classification system [27], follicles at the different

stages were collected at 14:00 from several females as described above. Intact follicles with

no damage were transferred into a 24-well culture plate containing 60 % L-15 medium.

These follicles were incubated at 28 °C with various doses of hormones or various exposure

times. Several progestin concentrations (DHP, 1-1,000 nM) with a 2 hrs incubation or

several exposure times (0.5-6 hrs) with 100 nM DHP were examined. Based on the result

of dose- and time-dependent experiments, follicles at different stages were incubated with

DHP (100 nM) for 2 hrs. Besides, stage IVa follicles were incubated with hCG (50 IU/mL)

[28] or DHP (100 nM) alone or in combination for 2 hrs. Stage IVa follicles collected from

$pgr^{-/-}$ were incubated with DHP (100 nM) for 2 hrs. After incubation, minimal 25 follicles

with no damage were collected and homogenized in RNAzol, stored in a -80 °C freezer

until RNA purification. Minimal three replicates were collected at each timepoint.

4.4 Real-time quantitative PCR (qPCR)

RevertAid cDNA synthesis kit (Thermo Scientific, Waltham, USA) was used for

reverse transcription (500 ng total RNA/sample). All qPCRs were carried out in a 20 μL

reaction on the qTOWER 2.2 real-time PCR system (Analytik Jena AG, Jena, Germany)

using the PowerUp SYBR green master mix (Applied Biosystems, USA). Each PCR

primer of a pair was designed to target different exons of the gene (Supplemental Table 3).

Expression of the target genes was determined as relative values using the comparative C_t

method [30] with the *ef1α* gene serving as an internal control.

4.5 Whole-mount *in situ* hybridization (WISH)

A DNA fragment corresponding to partial *f5* or *f3a* cDNAs was amplified from

ovarian cDNAs by PCR using the gene-specific primers (see Supplemental Table 3 for

detail) and cloned into the pGEM-T Easy vector (Promega, USA). Correct sequence and

orientations of inserts were validated by Sanger sequencing. Antisense or sense

digoxigenin-labeled cRNA probes for *f3a* or *f5* were synthesized by PCR using the plasmid

cDNA templates, SP6 or T7 RNA polymerase depending on the insert orientation. IVa or

IVb follicles were collected at 20:00 and 06:40 (immediately prior to ovulation),

respectively, and fixed in 4 % PFA overnight at 4 ℃. Following washing with PBST, the

follicles were treated with proteinase K (5 µg/mL) for 2 min at 37 °C, then washed and

subsequently hybridized with digoxigenin-labeled *f5* or *f3a* sense or antisense probes at

65 °C overnight using a previously established protocol [31]. Following WISH, follicles were

photographed using an M165FC dissecting microscope and DFC 550 digital camera (Leica,

Wetzlar, Germany).

4.6 Dual-luciferase reporter assay

The Pgr expression and reporter assay plasmid constructs were listed in

Supplemental Table 4. Human embryonic kidney cells (HEK 293T) were used for the

transactivation assays [32, 33]. Briefly, cells were maintained at 37 °C with 5 % CO_2 in phenol

red-free Dulbecco's modified eagle's medium (Hyclone, USA) containing

penicillin/streptomycin (Meilunbio, Dalian, China) and supplemented with 10 % fetal

bovine serum (FBS, Gibco, Brazil). Lipofectamine 3000 (Life Technologies) was used for

transient transfection according to manufactures' protocols. Cells were transiently

transfected with a firefly luciferase reporter plasmid (3 µg/vector/60mm petri dish), the

pRL-TK vector (150 ng/60mm petri dish) containing the Renilla luciferase reporter gene (as an internal control for transfection efficiency), and a Pgr expression vector or a control vector with no insert (1.5 μg/vector/60mm petri dish). After an overnight incubation at 37 °C, the medium was replaced with luciferase assay medium (DMEM without phenol red, supplemented with 10 % charcoal-stripped FBS) containing various concentrations of progestins (100 pM-1 μM DHP for zebrafish Pgr, P4 or human PGRB, respectively) or vehicle. Following additional 24 hrs of incubation at 37 °C, the cells were washed with PBS and harvested to determine firefly and Renilla luciferase activities according to the Dual-Luciferase reporter assay system (Promega, USA) on a luminometer (GloMax 20/20, Promega, USA). Firefly luciferase activities were normalized using Renilla luciferase data. After normalization for transfection efficiency, induction factors were calculated as the ratios of the average luciferase value of the steroid-treated samples versus vehicle-treated samples.

4.7 Blood capillaries in stage IV follicles

IVa immature follicles were obtained from paired female fish (*Tg(pgr:eGFP/ fli1:DsRed)*). In this line, the blood capillaries are labeled with DsRed, while the follicular cells are labeled with eGFP. The nucleus of follicles was co-stained with Hoechst 33342 in a 29-mm glass-bottom dish (Cellvis, Hangzhou, China). Images were captured using an SP8 confocal laser scanning microscope (Leica, Germany). IVb mature follicles were obtained from ovaries undergoing ovulation from faired female fish (*Tg(fli1:eGFP)*) at

7:00. A series of images during ovulation was recorded using a Leica digital camera and

fluorescent dissecting microscope described previously.

4.8 Erythrocyte numbers inside the capillaries on the surface of preovulatory follicles.

Erythrocytes inside the capillaries on the surface of ovarian follicles were stained

with *o*-dianisidine according to an established protocol [34]. Ovaries with IVb follicles were

collected at 6:40 or 7:00 from siblings of *pgr*$^{+/-}$ in-cross, producing both wildtype (*pgr*$^{+/+}$)

and *pgr*$^{-/-}$. As sampling moves closely toward 7:00, IVb follicles will be further in the

ovulation process in *pgr*$^{+/+}$ females. In contrast, IVb follicles in *pgr*$^{-/-}$ females will never

initiate ovulation [14, 16], which serves as an excellent control. After collection, the whole

ovaries were immediately stained for 15 min at room temperature in the dark in 0.6 mg/ml

o-dianisidine, 0.01 M sodium acetate (pH 4.5), 0.65 % H_2O_2, and 40 % (vol/vol) ethanol.

Stained ovaries were washed 3 times for 2 min each with PBST. At least 3 ovary samples

were collected at each timepoint for each genotype. Minimal 10 IVb follicles were

randomly chosen from each sample for image capture using DM 2500 LED microscope

and DFC 7000T digital camera (Leica), then the number of erythrocytes under the field of

view were counted by a reviewer who was blinded from sampling conditions and genotypes.

4.9 *In vitro* ovulation assay

Ovaries with IVb mature follicles from *Tg(fli1:eGFP)* zebrafish were collected at

06:40, 20 min before ovulation, and were placed in 60 % L-15 medium. Each ovary was

immediately cut into several small pieces and then transferred into a 12-well plate with

medium containing 2 mM EDTA, 100 µg/mL heparin, or 50 µg/mL warfarin, to examine

their inhibitory effect on ovulation. The dose of each drug was chosen based on previous

reports [35, 36]. Follicles were immediately dispersed by pipetting using a glass pipette (~ 1

mm in diameter). At least 50 IVb preovulatory follicles per well were used in each

experiment. The follicles were cultured at 28 °C in a 2 mL culture medium for 2 hrs.

Ovulation was continuously monitored until 09:00 under a fluorescent dissecting

microscope (Leica M165FC), and the numbers of oocytes that had successfully ovulated

were recorded. Minimal replicates for each treatment were eight.

4.10 Consecutive spawning test

The spawning of zebrafish was optimized under enhanced feeding conditions as

described previously [27]. Multiple pairs and spawning tanks were set up. Ten to twenty

mature females (three months old) of $f5^{+/-}$ or wildtype ($f5^{+/+}$) female siblings were used.

Each female was individually paired and housed with a fertile wildtype male. Around 21:00

(1 hrs prior to lights off), water in spawning tanks was replaced, and a mesh inner tank was

inserted into each spawning tank to prevent the eating of eggs by the adults. The number

of fertilized eggs was recorded for each pair of fish around noon of the next day. Then,

water would be replaced, and fish would be fed at least three times a day. The $f5^{+/-}$ and

wildtype ($f5^{+/+}$) females were continuously monitored for at least 3 weeks. Data from the

last two weeks were used for analyses.

4.11 Statistical analyses

Data analyses were performed using GraphPad Prism version 8. Depending on the experimental setup, either independent (unpaired two-tailed) Student's t-test or one-way ANOVA (one-tailed) followed by the Dunnett's post hoc test was used to assess statistical differences in comparison to the control group. Each experiment (e.g. qPCR, reporter assay, WISH, drug exposure, reporter assay, erythrocytes count, *in vitro* ovulation, fertility test) was repeated at least once.

Acknowledgments

We want to thank Dr. Mingyu Li (Xiamen University, China) and Dr. Yu Xue (Minnan Normal University, China) for providing zebrafish transgenic lines (*Tg(fli1:eGFP)*, *Tg(fli1:DsRed)*), Dr. Shuang-Bo Kong (Xiamen University, Xiamen, China) for sharing human PGRB expression vector (pCMV-HA-hPGRB). This work was supported by Fundamental Research Funds for the Central Universities (No. 20720200114), NSFC 31672628, and NSFC 41976092 to SXC, NIH GM100461 to YZ, and NIH R35 HL150784 to JAS.

References

1. Duffy DM, Ko C, Jo M, Brannstrom M, Curry TE. Ovulation: parallels with inflammatory processes. *Endocr. Rev.* **40**, 369-416 (2019).

2. Esmon CT. The interactions between inflammation and coagulation. *Br. J. Haematol.* **131**, 417-430 (2005).

3. Kanzaki H, Okamura H, Okuda Y, Takenaka A, Morimoto K, Nishimura T. Scanning electron-microscopic study of rabbit ovarian follicle microvasculature using resin injection corrosion casts. *J. Anat.* **134**, 697-704 (1982).

4. Cavender JL, Murdoch WJ. Morphological-studies of the microcirculatory system of periovulatory ovine follicles. *Biol. Reprod.* **39**, 989-997 (1988).

5. Zackrisson U, Lofman CO, Janson PO, Wallin A, Mikuni M, Brannstrom M. Alterations of follicular microcirculation and apex structure during ovulation in the rat. *Eur. J. Obstet. Gynecol. Reprod. Biol.* **157**, 169-174 (2011).

6. Brannstrom M, *et al*. Preovulatory changes of blood flow in different regions of the human follicle. *Fertil. Steril.* **69**, 435-442 (1998).

7. Gentry PA, Plante L, Schroeder MOB, LaMarre J, Young JE, Dodds WG. Human ovarian follicular fluid has functional systems for the generation and modulation of thrombin. *Fertil. Steril.* **73**, 848-854 (2000).

8. Kim YS, *et al*. Proteomic analysis of recurrent spontaneous abortion: Identification of an inadequately expressed set of proteins in human follicular fluid. *Proteomics* **6**, 3445-3454 (2006).

9. Parrott JA, Whaley PD, Skinner MK. Extrahepatic expression of fibrinogen by granulosa cells: potential role in ovulation. *Endocrinology* **133**, 1645-1649 (1993).

10. Gonzalez F, Kirwan JP, Rote NS, Minium J. Elevated circulating levels of tissue factor in polycystic ovary syndrome. *Clin. Appl. Thromb./Hemostasis* **19**, 66-72 (2013).

11. Natraj U, Richards JS. Hormonal-regulation, localization, and functional-activity of the progesterone-receptor in granulosa-cells of rat preovulatory follicles.

492   *Endocrinology* **133**, 761-769 (1993).

493   12.   Lydon JP*, et al.* Mice lacking progesterone-receptor exhibit pleiotropic

494   reproductive abnormalities. *Genes Dev.* **9**, 2266-2278 (1995).

495   13.   Hagiwara A, Ogiwara K, Katsu Y, Takahashi T. Luteinizing hormone-induced

496   expression of ptger4b, a Prostaglandin E-2 receptor indispensable for ovulation of

497   the medaka Oryzias latipes, is regulated by a genomic mechanism involving nuclear

498   progestin receptor. *Biol. Reprod.* **90**, 1-14 (2014).

499   14.   Zhu Y, Liu DT, Shaner ZC, Chen SX, Hong WS, Stellwag EJ. Nuclear progestin

500   receptor (Pgr) knockouts in zebrafish demonstrate role for Pgr in ovulation but not

501   in rapid non-genomic steroid mediated meiosis resumption. *Front. Endocrinol.* **6**,

502   37 (2015).

503   15.   Kubota K*, et al.* Rethinking progesterone regulation of female reproductive

504   cyclicity. *Proc. Natl. Acad. Sci. U. S. A.* **113**, 4212-4217 (2016).

505   16.   Tang HP*, et al.* Gene knockout of nuclear progesterone receptor provides insights

506   into the regulation of ovulation by LH signaling in zebrafish. *Sci. Rep.* **6**, 28545

507   (2016).

508   17.   Liu DT, Brewer MS, Chen SX, Hong WS, Zhu Y. Transcriptomic signatures for

509   ovulation in vertebrates. *Gen. Comp. Endocrinol.* **247**, 74-86 (2017).

510   18.   Wissing ML*, et al.* Identification of new ovulation-related genes in humans by

511   comparing the transcriptome of granulosa cells before and after ovulation triggering

512   in the same controlled ovarian stimulation cycle. *Hum. Reprod.* **29**, 997-1010

513   (2014).

514   19.   Carletti MZ, Christenson LK. Rapid effects of LH on gene expression in the mural

granulosa cells of mouse periovulatory follicles. *Reproduction* **137**, 843-855 (2009).

20. Runic R, Schatz F, Wan L, Demopoulos R, Krikun G, Lockwood CJ. Effects of norplant on endometrial tissue factor expression and blood vessel structure. *J. Clin. Endocrinol. Metab.* **85**, 3853-3859 (2000).

21. Haas DM, Hathaway TJ, Ramsey PS. Progestogen for preventing miscarriage in women with recurrent miscarriage of unclear etiology. *Cochrane Database Syst. Rev.*, 1-50 (2019).

22. Poulter NR, Meirik O, Chang CL, Farley TMM, Marmot MG. Venous thromboembolic disease and combined oral contraceptives: results of international multicentre case-control study. . *Lancet* **346**, 1575-1582 (1995).

23. Farley TMM, Meirik O, Chang CL, Marmot MG, Poulter NR. Effect of different progestagens in low estrogen oral-contraceptives on venous thromboembolic disease. *Lancet* **346**, 1582-1588 (1995).

24. Jick H, Jick SS, Gurewich V, Myers MW, Vasilakis C. Risk of idiopathic cardiovascular death and nonfatal venous thromboembolism in women using oral-contraceptives with differing progestagen components. *Lancet* **346**, 1589-1593 (1995).

25. Zeberg H, Kelso J, Paabo S. The neandertal progesterone receptor. *Mol. Biol. Evol.* **37**, 2655-2660 (2020).

26. Stein C, Caccamo M, Laird G, Leptin M. Conservation and divergence of gene families encoding components of innate immune response systems in zebrafish. *Genome Biol.* **8**, 1-23 (2007).

27. Liu DT, Carter NJ, Wu XJ, Hong WS, Chen SX, Zhu Y. Progestin and nuclear

progestin receptor are essential for upregulation of metalloproteinase in zebrafish preovulatory follicles. *Front. Endocrinol.* **9**, 517 (2018).

28. Liu DT, Hong WS, Chen SX, Zhu Y. Upregulation of adamts9 by gonadotropin in preovulatory follicles of zebrafish. *Mol. Cell. Endocrinol.* **499**, 110608 (2020).

29. Hanna RN, Zhu Y. Controls of meiotic signaling by membrane or nuclear progestin receptor in zebrafish follicle-enclosed oocytes. *Mol. Cell. Endocrinol.* **337**, 80-88 (2011).

30. Schmittgen TD, Livak KJ. Analyzing real-time PCR data by the comparative C-T method. *Nat. Protoc.* **3**, 1101-1108 (2008).

31. Hauptmann G, Gerster T. Multicolor whole-mount in situ hybridization. *Methods Mol. Biol.* **137**, 139-148 (2000).

32. Chen SX*, et al.* Molecular cloning and functional characterization of a zebrafish nuclear progesterone receptor. *Biol. Reprod.* **82**, 171-181 (2010).

33. Zhang YT, Hong WS, Liu DT, Qiu HT, Zhu Y, Chen SX. Involvement of membrane progestin receptor Beta (mPRbeta/Paqr8) in sex pheromone progestin-induced expression of luteinizing hormone in the pituitary of male chinese black sleeper (Bostrychus Sinensis). *Front. Endocrinol.* **9**, 397 (2018).

34. Detrich HW*, et al.* Intraembryonic hematopoietic-cell migration during vertebrate development. *Proc. Natl. Acad. Sci. U. S. A.* **92**, 10713-10717 (1995).

35. Hanumanthaiah R, Thankavel B, Day K, Gregory M, Jagadeeswaran P. Developmental expression of vitamin K-dependent gamma-carboxylase activity in zebrafish embryos: effect of warfarin. *Blood Cells Mol. Dis.* **27**, 992-999 (2001).

36. Ogiwara K, Takano N, Shinohara M, Murakami M, Takahashi T. Gelatinase A and

membrane-type matrix metalloproteinases 1 and 2 are responsible for follicle rupture during ovulation in the medaka. *Proc. Natl. Acad. Sci. U. S. A.* **102**, 8442-8447 (2005).

37. Hanna RN*, et al*. Characterization and expression of the nuclear progestin receptor in zebrafish gonads and brain. *Biol. Reprod.* **82**, 112-122 (2010).

38. Takahashi T, Hagiwara A, Ogiwara K. Follicle rupture during ovulation with an emphasis on recent progress in fish models. *Reproduction* **157**, R1-R13 (2019).

39. Weyand AC*, et al*. Analysis of factor V in zebrafish demonstrates minimal levels needed for early hemostasis. *Blood Adv.* **3**, 1670-1680 (2019).

40. Osterud B, Bjorklid E. Sources of tissue factor. *Semin. Thromb. Hemost.* **32**, 11-23 (2006).

41. Kato S*, et al*. Progesterone increases tissue factor gene expression, procoagulant activity, and invasion in the breast cancer cell line ZR-75-1. *J. Clin. Endocrinol. Metab.* **90**, 1181-1188 (2005).

42. Richer JK, Jacobsen BM, Manning NG, Abel MG, Wolf DM, Horwitz KB. Differential gene regulation by the two progesterone receptor isoforms in human breast cancer cells. *J. Biol. Chem.* **277**, 5209-5218 (2002).

43. Lockwood CJ, Krikun G, Papp C, Aigner S, Nemerson Y, Schatz F. Biological mechanisms underlying ru-486 clinical effects - inhibition of endometrial stromal cell tissue factor content. *J. Clin. Endocrinol. Metab.* **79**, 786-790 (1994).

44. Bravo ML*, et al*. Progesterone regulation of tissue factor depends on MEK1/2 activation and requires the proline-rich site on progesterone receptor. *Endocrine* **48**, 309-320 (2015).

45.   Moverare S, Skrtic S, Lindberg MK, Dahlman-Wright K, Ohlsson C. Estrogen

increases coagulation factor V mRNA levels via both estrogen receptor-alpha and -

beta in murine bone marrow/bone. *Eur. J. Endocrinol.* **151**, 259-263 (2004).

46.   Cleuren AC, Van der Linden IK, De Visser YP, Wagenaar GT, Reitsma PH, Van

Vlijmen BJ. 17alpha-Ethinylestradiol rapidly alters transcript levels of murine

coagulation genes via estrogen receptor alpha. *J. Thromb. Haemost.* **8**, 1838-1846

(2010).

47.   Spronk HM, Govers-Riemslag JW, ten Cate H. The blood coagulation system as a

molecular machine. *Bioessays* **25**, 1220-1228 (2003).

48.   Esmon CT. The impact of the inflammatory response on coagulation. *Thromb. Res.*

114, 321-327 (2004).

49.   Hathcock JJ. Flow effects on coagulation and thrombosis. *Arterioscler. Thromb.*

Vasc. Biol. **26**, 1729-1737 (2006).

50.   Mitsube K, Brannstrom M, Haraldsson B. Modulation of microvascular

permeability in the preovulatory rat ovary by an ovulatory gonadotropin stimulus.

Fertil. Steril. **99**, 903-909 (2013).

51.   Spiliopoulos D, Kadir RA. Congenital factor V and VIII deficiency in women: a

systematic review of literature and report of two new cases. *Blood Coagul.*

Fibrinolysis **27**, 237-241 (2016).

52.   Zhu Y. Metalloproteases in gonad formation and ovulation. *Gen. Comp. Endocrinol.*

314, 113924 (2021).

53.   Travis J, Salvesen GS. Human plasma proteinase inhibitors. *Annu. Rev. Biochem.*

52, 655-709 (1983).

54.   Migone FF, Cowan RG, Williams RM, Gorse KJ, Zipfel WR, Quirk SM. In vivo

imaging reveals an essential role of vasoconstriction in rupture of the ovarian

follicle at ovulation. *Proc. Natl. Acad. Sci. U. S. A.* **113**, 2294-2299 (2016).

55.   Yamada M, Gentry PA. Hemostatic profile of bovine ovarian follicular fluid. *Can.*

J. Physiol. Pharmacol. **73**, 624-629 (1995).

56.   Yamada M, Gentry PA. The hemostatic profile of equine ovarian follicular fluid.

Thromb. Res. **77**, 45-54 (1995).

57.   Roach LE, Petrik JJ, Plante L, LaMarre J, Gentry PA. Thrombin generation and

presence of thrombin receptor in ovarian follicles. *Biol. Reprod.* **66**, 1350-1358

(2002).

58.   Medved L, Nieuwenhuizen W. Molecular mechanisms of initiation of fibrinolysis

by fibrin. *Thromb. Haemost.* **89**, 409-419 (2003).

59.   Beers WH, Strickland S, Reich E. Ovarian plasminogen activator: Relationship to

ovulation and hormonal regulation. *Cell* **6**, 387-394 (1975).

60.   Ogiwara K, Minagawa K, Takano N, Kageyama T, Takahashi T. Apparent

involvement of plasmin in early-stage follicle rupture during ovulation in medaka.

Biol. Reprod. **86**, 113 (2012).

61.   Montefusco MC*, et al.* Clinical and molecular characterization of 6 patients

affected by severe deficiency of coagulation factor V: broadening of the mutational

spectrum of factor V gene and in vitro analysis of the newly identified missense

mutations. *Blood* **102**, 3210-3216 (2003).

Figure 1. Dramatic increase of *f5* an*d f3a* transcripts in follicular cells of preovulatory follicles during a daily spawning cycle *in vivo*.

(**A**) Schematic illustration of daily spawning cycle in zebrafish synchronized with lights on. Oocyte maturation in zebrafish typically occurs around 06:40, while ovulation occurs around 07:00.

(**B**) Cell-specific expression of *f5, f3a, and f3b* in mature follicles examined by RT-PCR. Transcripts of *f5* and *f3a* were expressed in the follicular cells but not in the oocytes, while the transcript of *f3b* was expressed in both follicular cells and oocytes of mature preovulatory follicles. MK: molecular makers; I: intact follicles including both surrounding follicular cells and germ cell; O: denuded oocyte (without follicular cells); F: follicular cells (with no germ cell); N: no template control.

(**C**) Dramatic increase of *f5, f3a,* and *f3b* transcripts in preovulatory follicles during a daily

spawning cycle *in vivo*. Fully-grown ovarian follicles (stage IVa or IVb) were collected

from wildtype mature female zebrafish at four representative timepoints during a daily

spawning cycle (see Fig. 1A). Expression of *f3a, f3b,* and *f5* mRNA were determined by

qPCR and normalized to the internal housekeeping gene (*ef1α*). Data are expressed as the

mean ± SEM (N = 4) relative to the respective transcript levels of *f5* in IVa follicles

collected at 12:00. P-values were calculated by one-tailed one-way ANOVA followed by

Dunnett's test against respective controls (samples collected at 12:00).

(**D, E**) Vividly increase of *f5* and *f3a* transcripts in the follicular cells of mature follicles.

Fully-grown but immature (stage IVa) and mature follicles but prior to ovulation (stage

IVb) were collected at 20:00 and 06:40, respectively. Strong signals of *f5* and *f3a*

transcripts were only observed in mature follicles (IVb) by a whole-mount *in situ*

hybridization, while no signals were observed in the immature follicles (IVa, D, E). The

positive ISH signals were localized in the follicular cells (F, G). Scale bar, 250 μm (D, E)

and 50 μm (F, G).

Figure 2. Transient increase of *f5* and *f3a* expression in preovulatory follicles in

response to stimulation of progestin *in vitro*.

(**A, C**) Effects of various doses of progestin (DHP) (2 hrs exposure) on *f5* and *f3a*

expression in fully grown immature preovulatory follicles (stage IVa) *in vitro*. (**B, D**)

Effects of DHP (100 nM) on *f5* and *f3a* expression in preovulatory follicles (IVa) at various

timepoints *in vitro*. P-values were calculated by one-tailed one-way ANOVA followed by

Dunnett's test against respective controls (0 timepoint or vehicle control).

Figure 3. Increase of *f5* and *f3a* transcripts in late-stage follicles is mediated by Pgr.
Follicles were incubated with progestin (DHP, 100nM), or LH (hCG, 50 IU/ml) alone, or in combination for 2 hrs at 28 ℃ *in vitro* before qPCR analysis. (**A**) An increase of *f5* transcript was observed in late-stage (III and IVa) follicles from wildtype ($pgr^{+/+}$) zebrafish, but not in early-stage follicles (I or II) from wildtype zebrafish, in response to progestin stimulation. (**B**) An increase of *f3a* expression was observed only in fully-grown immature follicles (IVa) from wildtype ($pgr^{+/+}$) zebrafish, but not in developing follicles (I-III) from wildtype fish, in response to progestin stimulation. (**C**) Synergistic increase of *f5* mRNA by hCG and DHP in fully grown immature follicles (IVa) from wildtype zebrafish. (**D**) Progestin but not hCG stimulated the expression of *f3a* mRNA in full-grown immature follicles (IVa) from wildtype zebrafish. (**E**) Both *f3a* and *f5* transcripts in preovulatory follicles (stage IVb) of $pgr^{-/-}$ were significantly lower than that of $pgr^{+/+}$ (WT). (**F**) Stimulatory effect of progestin (DHP) for *f3a* and *f5* expression was abolished in the stage

680   IVa follicles of *pgr⁻ᐟ⁻*. The expression of *f5* and *f3a* transcripts were determined by qPCR

681   and normalized to an internal housekeeping gene (*ef1α*). Data are expressed as the mean ±

682   SEM (N = 3) relative to the respective transcript levels of target gene measured in wildtype,

683   or vehicle treatment group. P-values were calculated by two-tailed Students' t-test.

Figure 4. Progestin via Pgr enhances promoter activity of *f5* or *f3a*.

HEK293T cells were transiently co-transfected with, a firefly luciferase reporter vector

containing putative Pgr binding elements (3 μg/vector/60mm petri dish), a pRL-TK vector

(150 ng/60mm petri dish) containing the Renilla luciferase reporter gene (as a control for

transfection efficiency), and a Pgr expression vector (1.5 μg/vector/60mm petri dish) or an

empty control expression vector (1.5 μg/vector/60mm petri dish). In the presence of

zebrafish Pgr (zPgr), progestin significantly increased the MMTV (**A**), *zf5* (**B**), and *zf3a*

(**C**) promoter activities, and the increase was dose-dependent. (**D**) In the presence of human

PGRB (hPGRB), P4 also significantly increased human *hF5* promoter activity, and the

increase was dose-dependent. The cells were incubated for 24 hrs with increasing

696 concentrations of progestins (DHP or P4: 100 pM to 1 µM). Extracts of the treated cells

697 were assayed for luciferase activity. Values are shown relative to the luciferase activity of

698 the vehicle treatment group. Data are expressed as mean $\pm$ SEM (N = 3). P-values were

699 calculated by one-tailed one-way ANOVA followed by Dunnett's test against respective

700 controls (vehicle controls, zero dose).

Figure 5. Follicular cells with strong Pgr expression are adjacent to blood capillaries.
Images of a representative stage IVa follicle from a double transgenic zebrafish line
Tg(pgr:eGFP/ fli1:DsRed) are shown. (**A**) Nucleus staining of follicular cells by Hoechst
33342. (**B**) Pgr promoter drives GFP expression in follicular cells. (**C**) Red fluorescence-
labeled vascular endothelial cells on the follicles. (**D**) Merged images to show that
follicular cells with strong GFP expression driven by the Pgr promoter are adjacent to blood

709   capillaries. Red arrowheads indicate follicular cells with strong GFP expression driven by

710   the Pgr promoter. Scale bars: 100 μm.

Figure 6. Rupture of blood capillaries and follicular cells during ovulation.

Ovaries undergoing ovulation were collected from *Tg(fli1:eGFP)* female at 07:00. (**A-F**),

a series of images were taken when a mature (stage IVb) ovarian follicle started ovulation.

Interconnected blood vessels labeled with GFP were observed easily on the surface of

ovarian follicles. A red arrow indicates a ruptured blood vessel on the surface of an

717   ovulating follicle. Red arrowheads and red dotted lines indicate a hole left in the remaining

718   follicular cells after ovulation. The white dotted line represents the remaining follicular

719   cells. Scale bars: 250 μm. Oc, oocyte. (Please also see supplemental video clip A for

720   additional details).

Figure 7. Increased erythrocytes in capillaries on the surface of mature follicles during progression but prior to rupture.

(**A, D, B, E**) Erythrocytes within the surface capillaries of mature follicles (IVb) from wildtype ($pgr^{+/+}$) female zebrafish collected at 6:40 or 7:00 were visualized using *o*-dianisidine staining as follicles progressing towards rupture and ovulation. (**C, F**) Erythrocytes were also visualized in similar mature follicles that were collected at 7:00 from $pgr^{-/-}$ females, in which ovulation never occurs. (**G**) At least 30 mature follicles were randomly chosen from at least 3 different females of each condition, and the number of erythrocytes in capillaries were counted by a blinded observer. Data are expressed as mean ± SEM (N ≥ 30). P-values were calculated by two-tailed Student's t-test. Scale bars: 500 μm (A-C) or 100 μm (D-F).

Figure 8. Heparin inhibits ovulation *in vitro*.

Ovulation rate was determined using mature follicles (IVb) isolated from transgenic female

zebrafish (*Tg(fli1:eGFP)*). The intact IVb follicles were easily recognizable by the

interconnecting blood vessels with strong GFP expression (**A, C**). After ovulation, oocytes

were easily recognizable by the absence of interconnecting GFP-positive vessels and

expansion of the vitelline envelope (chorion) (**B, D**). Follicular layers separated from

ovulated oocytes could also be easily distinguished due to interconnecting blood vessels

and a dramatic size reduction (**B, D**). (**A, C**) Mature follicles (IVb) were collected from

Tg(fli1:eGFP) female zebrafish. (**B, D**) Representative images of ovulated mature follicles

were taken 2 hrs after incubation at 28 °C. Arrows indicate the remaining follicular layer

following ovulation. Arrowheads indicate expanded vitelline envelopes. Oc, oocyte. (**E**)

mature follicles were incubated in a medium containing EDTA (2 mM), heparin (100

µg/mL), warfarin (50 µg/mL), or vehicle control, and rates of ovulation were determined

748   2 hrs following incubation. Data are expressed as the mean $\pm$ SEM (N $\geqslant$ 7). P-values were

749   calculated by two-tailed Student's t-test. Scale bars: 1 mm (A, B) or 250 μm (C, D).

Figure 9. Reduced fecundity in *f5*^{+/-} female zebrafish.

(**A**) Reduced numbers of live embryos per spawning for each *f5*$^{+/-}$ female in comparison to that of wildtype (*f5*$^{+/+}$) females. (**B**) Reduced total numbers of embryos for each *f5*$^{+/-}$ female fish in comparison to wildtype (*f5*$^{+/+}$) females. (**C**) Reduced spawning times in *f5*$^{+/-}$ females in comparison to that of wildtype (*f5*$^{+/+}$) females. (**D**) Increased spawning intervals in *f5*$^{+/-}$ females in comparison to wildtype (*f5*$^{+/+}$) females. Data are expressed as the mean ± SEM (N ⩾ 14). P-values were calculated by two-tailed Student's t-test.

Figure 10. A proposed model for Pgr mediated upregulation of F5 and F3a in blood coagulation that facilitates ovulation in zebrafish.

Luteinizing hormone (LH) induces ovulation via luteinizing hormone receptor (Lhcgr) in preovulatory follicles, namely in stage IV follicles in zebrafish. LH signaling is mainly mediated by progestin (DHP, i.e., 17α,20β-dihydroxy-progesterone, an active ovulation inducing steroid in zebrafish) and its receptor (Pgr) in the production and activation of metalloproteases. The illustration on the left side is showing the upregulation of metalloproteases in preovulatory follicles, to be more specific, during the transition from stage IVa to IVb follicles, which is an important preparation step for ovulation. However, these proteases are produced in a latent form, i.e., inactive form of metalloproteases (iMPs, see Zhu, 2021 for detail), and inhibited by tissue inhibitors of metalloproteases (TIMP) presented at high levels in and delivered by blood capillaries to the follicles. The illustration on the right side is showing a dramatic increase of coagulation factors *f5* and *f3a* shortly before ovulation, specifically in the late stage of IVb follicles, which are also mainly

upregulated by progestin via Pgr. Once F5 and F3a are activated, they accelerate blood

coagulation, which in turn stops blood flow and therefore effectively terminates TIMP

inhibition of metalloproteases from the bloodstream. In addition, the production of fibrin

(not shown) due to the formation of blood clots stimulates activities of

plasmin/plasminogen activators, which are well-known inducers for metalloprotease

activation. The combined force of these two pathways triggers rapid activation of

metalloproteases, breakdown of extracellular matrix and basal membrane, rupture of

follicular cell layers, the final release of a mature oocyte.