Nuclear Progestin Receptor Mediated Linkage of Blood Coagulation and Ovulation

Jing Huang [a], Chao Sun [a], Dong Teng Liu[a], Nan Nan Zhao [a], Jordan A. Shavit [b], Yong Zhu [a, c, *], Shi Xi Chen [a, d, *]

[a] State Key Laboratory of Marine Environmental Science, College of Ocean and Earth Sciences, Xiamen University, Xiamen 361102, China

[b] Departments of Pediatrics and Human Genetics, University of Michigan, Ann Arbor, MI 48109, USA

[c] Department of Biology, East Carolina University, 1000 E 5th Street, Greenville, NC 27858, USA

[d] State-Province Joint Engineering Laboratory of Marine Bioproducts and Technology, Xiamen University, Xiamen 361102, China

Short title: Pgr regulate coagulation factor for ovulation

Keywords: Pgr, progestin, coagulation factor, ovulation

* **Corresponding authors and request for reprints:**

Dr. Shi Xi Chen

College of Ocean and Earth Sciences, Xiamen University, Xiamen 361102, China.

Phone: +86-13515969358, E-mail address: chenshixi@xmu.edu.cn.

Dr. Yong Zhu

Department of Biology, East Carolina University, 1000 E 5th Street, Greenville, NC

27858, USA. Phone: 252-328-6504, E-mail address: zhuy@ecu.edu.

Financial Support:

This work was supported by Fundamental Research Funds for the Central

Universities (No. 20720200114), NSFC 31672628, and NSFC 41976092 to SXC, NIH

GM100461 to YZ, and NIH R35 HL150784 to JAS.

Author Contributions:

JH performed most experiments, analyzed the data, and wrote the draft with the

assistance of CS, DTL and NNZ. JAS generated the *f5* zebrafish knockout, provided

protocols and experimental suggestions, discussed experimental results, and revised the

manuscript. YZ supervised the project, conducted the fertility experiment, and revised the

manuscript. SXC conceived and supervised the project and revised the manuscript.

Disclosure summary: The authors have nothing to disclose.

Abstract

Ovulation is a dramatic remodeling process including rupture of blood capillaries and clotting, but coagulation is not thought to directly regulate this process. Herein, we report remarkable increases of coagulation factors V (*f5*, ~3145-fold) and tissue factor (*f3a*, ~120-fold) in zebrafish ovarian follicle cells during ovulation. This increase was mediated through the nuclear progestin receptor (Pgr), which is essential for ovulation in zebrafish. This increase was totally abolished in ovarian follicular cells from $pgr^{-/-}$ mutants. In addition, promoter activities of *f5* and *f3a* were significantly enhanced by progestin (DHP) via Pgr. Similar regulation of human *F5* promoter activity was induced via human PGRB, suggesting a conserved mechanism. Site-directed mutagenesis of the zebrafish *f5* promoter further demonstrated a direct regulation of coagulation factors via progestin response elements. Moreover, a stark increase of erythrocytes occurred in capillaries meshed in wildtype preovulatory follicles, but was absent in $pgr^{-/-}$ mutants. Interestingly, the anticoagulant heparin inhibited ovulation *in vitro*. Furthermore, reduced fecundity was observed in $f5^{+/-}$ female zebrafish. Taken together, our study provides plausible evidence for steroid regulation of coagulation factors, and a new hypothesis for blood clotting triggered ovulation in vertebrates.

1.   Introduction

Ovulation is a tissue remodeling process involving the rupture of follicular layers, basal membrane, and many blood capillaries intertwined in the follicular cells. This dramatic remodeling process share several features with inflammatory responses (1). Blood coagulation was suggested to be part of the innate defense in response to inflammation during ovulation (2). Formation of blood clots at apical vessels was found shortly before follicle rupture in several animal species including sheep, rabbits, and rodents (1,3-5). Cessation of blood flow likely due to blood coagulation was also observed in apical vessels of ovarian follicles during the late ovulatory phase in women (6). In addition, an increase of activity and expression of thrombin (a protease essential for fibrin formation) and its receptors were also reported in luteinized granulosa cells and follicular fluid in human (7,8). Increased fibrinogen secretion in bovine granulosa cells was thought to be related to the proteolytic activity needed for follicle rupture. Circulating F3 (also known as tissue factor) was elevated in women with polycystic ovary syndrome that is characterized by chronic oligo-ovulation or anovulation (9). Although these reports imply possible association between blood coagulation and the ovulation process, the mechanisms for this role in ovulation are still unclear.

Progestin and its nuclear progesterone receptor (PGR) are well-established initiators for vertebrate ovulation (10-15). Recently, we identified coagulation factors *f3a* and *f5* as potential downstream targets of Pgr in a transcriptomic analysis of preovulatory follicles from zebrafish (16). Interestingly, changes in these coagulation factors during ovulation appear to be conserved among human, mice, and zebrafish (16-18). Possible involvements of progestin or PGR in the regulation of coagulation have been hinted from

several studies using progestin contraceptives. F3 was upregulated in endometriosis and at endometrial bleeding sites in women using long-term progestin-only contraception (19). Orally administered progesterone reduced spontaneous miscarriages and improved fertility among women who had a history of recurrent miscarriages due to abnormal bleeding in early pregnancy (20). Increased risk of venous thromboembolic events was associated with combined oral contraceptives, particularly those containing the progestin desogestrel (21,22). A recent genetic study indicated that the present-day carriers of the Neandertal variant of PGR had less bleeding during early pregnancy because of high levels of PGR expression (23). However, evidence of progestin induced or PGR mediated regulation of coagulation factors, especially in ovarian follicles during ovulation, is lacking.

In this study, we examined the expression, hormonal regulation, and potential function of coagulation factors (*f5* & *f3*) in preovulatory follicles of zebrafish. The *f3* gene is duplicated in zebrafish (*f3a* and *f3b*) (24). We first demonstrated the expression and localization of *f5, f3a,* and *f3b* in preovulatory follicular cells. We found that progestin (DHP, 17α, 20β-dihydroxy-4-pregnen-3-one, a native ovulation-inducing ligand in zebrafish) via Pgr regulated *f5* and *f3a* expression in preovulatory follicles. We also found that follicular cells with strong Pgr expression were adjacent to blood capillaries, and inhibition of blood coagulation reduced ovulation. Finally, fertility and ovulation were reduced in $f5^{+/-}$ zebrafish mutants *in vivo*. Our study provides credible evidence that coagulation factors (*f5* & *f3a*) in preovulatory follicular cells are regulated by progestin (DHP) via Pgr, which are well-known upstream initiators for ovulation. Our

results also suggest a previously unknown role of these coagulation factors in ovulation

and female fertility.

2.   Materials and Methods

2.1 Animals & Reagents

Zebrafish used are offspring of Tübingen strain unless stated otherwise. Detailed

information for knockouts, transgenic lines, key chemicals, and reagents are listed in

Supplemental Table 1 & Table 2 (25). Fish were fed three times daily with a commercial

food (Otohime B2; Marubeni Nisshin Feed, Tokyo, Japan), supplemented with newly

hatched brine shrimp, and were maintained in recirculating system (ESEN, Beijing,

China) at 28 ± 0.5 □ under a controlled photoperiod (lights on at 08:00, off at 22:00). The

daily ovulation cycle in mature adult zebrafish is established according to an established

protocol (26). Briefly, each mature adult female and male zebrafish were paired in a

spawning tank. These fish typically start spawning soon after lights on in the morning.

Around noon, fish water was replaced, and fish were fed at least three times daily with

the commercial food and supplemented with newly hatched brine shrimp. Around one

hour prior to lights off, fish water was replaced again, a mesh inner tank was inserted into

each spawning tank to prevent cannibalism typical for zebrafish adults. After one week,

most fish form a daily ovulation cycle. When applicable, siblings from heterozygous in-

cross were used for comparison. All experimental protocols were approved by the

Institutional Animal Care and Use Committee (IACUC) at Xiamen University.

2.2 Collection of fully-grown follicles (stage □) during a daily spawning cycle

Fully-grown immature follicles (stage IVa, >650 μm, with visible germinal vesicles, i.e., GV, appears opaque) or mature follicles (stage IVb, GV disappears, appears transparent) were collected at four representative timepoints during a daily spawning cycle (Fig. 1A): 12:00 (IVa follicles); 20:00 (IVa follicles); 05:00 (the onset of oocyte maturation, IVa follicles); and 06:40 (completion of oocyte maturation but prior to ovulation, IVb follicles). To be simple and accurate in our description, we will use "IVa" to indicate fully-grown immature stage IVa follicles; "IVb" to indicate mature but yet ovulated follicles (prior to ovulation) throughout this manuscript.

To sacrifice fish humanely, the spinal cord and blood supply behind the gill cover were cut off swiftly using sharp scissors following a lethal dose of anesthetization (MS-222, 300 mg/L buffered solutions). Ovaries were immediately placed in 60 % L-15 media containing 15 mM HEPES at pH 7.2. Individual follicles were obtained according to an established procedure (26,27). At each timepoint, a total of 25 □a or IVb follicles (containing both germ cell and surrounding follicular cells) from 4 females were pooled as one sample. In addition, the follicular cells (∼100 follicles) or their enclosed oocytes (∼20 follicles) were collected separately from IVb follicles according to an established method (28). These samples were immediately homogenized in RNAzol solution (Molecular Research Center, Cincinnati, USA) and stored at a -80 ℃ freezer until further processing.

2.3 Hormone treatments

According to established follicle classification system (26), follicles at the different stages were collected at 14:00 from several females as described above. Intact follicles with no damage were transferred into a 24-well culture plate containing 60 % L-15 medium. These follicles were incubated at 28 □ with various doses of hormones or various exposure times. Several progestin concentrations (DHP, 1-1,000 nM) with a 2 hrs incubation or several exposure times (0.5-6 hrs) with 100 nM DHP were examined. Based on the result of dose- and time-dependent experiments, follicles at different stages were incubated with DHP (100 nM) for 2 hrs. Besides, stage IVa follicles were incubated with hCG (50 IU/mL) (27) or DHP (100 nM) alone or in combination for 2 hrs. Stage IVa follicles collected from $pgr^{-/-}$ were incubated with DHP (100 nM) for 2 hrs. After incubation, minimal 25 follicles with no damage were collected and homogenized in RNAzol, stored in a -80 □ freezer until RNA purification. Minimal three replicates were collected at each timepoint.

2.4 Real-time quantitative PCR (qPCR)

RevertAid cDNA synthesis kit (Thermo Scientific, Waltham, USA) was used for reverse transcription (500 ng total RNA/sample). All qPCRs were carried out in a 20 μL reaction on the qTOWER 2.2 real-time PCR system (Analytik Jena AG, Jena, Germany) using the PowerUp SYBR green master mix (Applied Biosystems, USA). Each PCR primer of a pair was designed to target different exons of the gene (Supplemental Table

172 3) (25). Expression of the target genes was determined as relative values using the

173 comparative C_t method (29) with the *ef1α* gene serving as an internal control.

2.5 Whole-mount *in situ* hybridization (WISH)

A DNA fragment corresponding to partial *f5* or *f3a* cDNAs was amplified from

ovarian cDNAs by PCR using the gene-specific primers (see Supplemental Table 3 for

detail) (25) and cloned into the pGEM-T Easy vector (Promega, USA). Correct sequence

and orientations of inserts were validated by Sanger sequencing. Antisense or sense

digoxigenin-labeled cRNA probes for *f3a* or *f5* were synthesized by PCR using the

plasmid cDNA templates, SP6 or T7 RNA polymerase depending on the insert

orientation. IVa or IVb follicles were collected at 20:00 and 06:40 (immediately prior to

ovulation), respectively, and fixed in 4 % PFA overnight at 4 ℃. Following washing with

PBST, the follicles were treated with proteinase K (5 µg/mL) for 2 min at 37 °C, then

washed and subsequently hybridized with digoxigenin-labeled *f5* or *f3a* sense or

antisense probes at 65 °C overnight using a previously established protocol (30).

Following WISH, follicles were photographed using an M165FC dissecting microscope

and DFC 550 digital camera (Leica, Wetzlar, Germany).

2.6 Dual-luciferase reporter assay

The Pgr expression and reporter assay plasmid constructs were listed in

Supplemental Table 4 (25). Site-directed mutagenesis of four potential Pgr response

elements (PRE, i.e., Pgr binding sites) in zebrafish *f5* proximal promoter sequences (Fig.

4E, Supplemental Table 6) (25)was conducted using a Q5 site-directed mutagenesis kit

(NEB, Ipswich, MA, USA) according to the manufacturer's protocol. Human embryonic

kidney cells (HEK 293T) were used for the transactivation assays (31,32). Briefly, cells

were maintained at 37 °C with 5 % CO_2 in phenol red-free Dulbecco's modified eagle's

medium (Hyclone, USA) containing penicillin/streptomycin (Meilunbio, Dalian, China)

and supplemented with 10 % fetal bovine serum (FBS, Gibco, Brazil). Lipofectamine

3000 (Life Technologies) was used for transient transfection according to manufactures'

protocols. Cells were transiently transfected with a firefly luciferase reporter plasmid (3

μg/vector/60mm petri dish), the pRL-TK vector (150 ng/60mm petri dish) containing the

Renilla luciferase reporter gene (as an internal control for transfection efficiency), and a

Pgr expression vector or a control vector with no insert (1.5 μg/vector/60mm petri dish).

After an overnight incubation at 37 °C, the medium was replaced with luciferase assay

medium (DMEM without phenol red, supplemented with 10 % charcoal-stripped FBS)

containing various concentrations of progestins (100 pM-1 μM DHP for zebrafish Pgr, P4

or human PGRB, respectively) or vehicle. Following additional 24 hrs of incubation at 37

°C, the cells were washed with PBS and harvested to determine firefly and Renilla

luciferase activities according to the Dual-Luciferase reporter assay system (Promega,

USA) on a luminometer (GloMax 20/20, Promega, USA). Firefly luciferase activities

were normalized using Renilla luciferase data. After normalization for transfection

efficiency, induction factors were calculated as the ratios of the average luciferase value

of the steroid-treated samples versus vehicle-treated samples.

2.7 Blood capillaries in stage ☐ follicles

IVa immature follicles were obtained from paired female fish (*Tg(pgr:eGFP/*

fli1:DsRed)). In this line, the blood capillaries are labeled with DsRed, while the

follicular cells are labeled with eGFP. The nucleus of follicles was co-stained with

Hoechst 33342 in a 29-mm glass-bottom dish (Cellvis, Hangzhou, China). Images were

captured using an SP8 confocal laser scanning microscope (Leica, Germany). IVb mature

follicles were obtained from ovaries undergoing ovulation from faired female fish

(*Tg(fli1:eGFP)*) at 7:00. A series of images during ovulation was recorded using a Leica

digital camera and fluorescent dissecting microscope described previously.

2.8 Erythrocyte numbers inside the capillaries on the surface of preovulatory

follicles.

Erythrocytes inside the capillaries on the surface of ovarian follicles were stained

with *o*-dianisidine according to an established protocol (33). Ovaries with IVb follicles

were collected at 6:40 or 7:00 from siblings of $pgr^{+/-}$ in-cross, producing both wildtype

($pgr^{+/+}$) and $pgr^{-/-}$. As sampling moves closely toward 7:00, IVb follicles will be further

in the ovulation process in $pgr^{+/+}$ females. In contrast, IVb follicles in $pgr^{-/-}$ females will

never initiate ovulation (13,15), which serves as an excellent control. After collection, the

whole ovaries were immediately stained for 15 min at room temperature in the dark in 0.6

mg/ml *o*-dianisidine, 0.01 M sodium acetate (pH 4.5), 0.65 % H_2O_2, and 40 % (vol/vol)

ethanol. Stained ovaries were washed 3 times for 2 min each with PBST. At least 3 ovary

samples were collected at each timepoint for each genotype. Minimal 10 IVb follicles

were randomly chosen from each sample for image capture using DM 2500 LED

microscope and DFC 7000T digital camera (Leica), then the number of erythrocytes

under the field of view were counted by a reviewer who was blinded from sampling

conditions and genotypes.

2.9 *In vitro* ovulation assay

Ovaries with IVb mature follicles from *Tg(fli1:eGFP)* zebrafish were collected at

06:40, 20 min before ovulation, and were placed in 60 % L-15 medium. Each ovary was

immediately cut into several small pieces and then transferred into a 12-well plate with

medium containing 2 mM EDTA, 100 µg/mL heparin, or 50 µg/mL warfarin, to examine

their inhibitory effect on ovulation. The dose of each drug was chosen based on previous

reports (34,35). Follicles were immediately dispersed by pipetting using a glass pipette (~

1 mm in diameter). At least 50 IVb preovulatory follicles per well were used in each

experiment. The follicles were cultured at 28 °C in a 2 mL culture medium for 2 hrs.

Ovulation was continuously monitored until 09:00 under a fluorescent dissecting

microscope (Leica M165FC), and the numbers of oocytes that had successfully ovulated

were recorded. Minimal replicates for each treatment were eight.

2.10 Consecutive spawning test

The spawning of zebrafish was optimized under enhanced feeding conditions as

described previously (26). Multiple pairs and spawning tanks were set up. Ten to twenty

mature females (three months old) of $f5^{+/-}$ or wildtype ($f5^{+/+}$) female siblings were used.

Each female was individually paired and housed with a fertile wildtype male. Around

21:00 (1 hrs prior to lights off), water in spawning tanks was replaced, and a mesh inner

tank was inserted into each spawning tank to prevent the eating of eggs by the adults. The

number of fertilized eggs was recorded for each pair of fish around noon of the next day.

Then, water would be replaced, and fish would be fed at least three times a day. The $f5^{+/-}$

and wildtype ($f5^{+/+}$) females were continuously monitored for at least 3 weeks. Data from

the last two weeks were used for analyses.

2.11 Statistical analyses

Data analyses were performed using GraphPad Prism version 8. Depending on the

experimental setup, either independent (unpaired two-tailed) Student's t-test or one-way

ANOVA (one-tailed) followed by the Dunnett's post hoc test was used to assess statistical

differences in comparison to the control group. Each experiment (e.g. qPCR, reporter

assay, WISH, drug exposure, reporter assay, erythrocytes count, *in vitro* ovulation,

fertility test) was repeated at least once.

3.   Results

3.1 Dramatic increase of *f5* an*d f3a* transcripts in follicular cells of preovulatory

follicles during a natural spawning cycle *in vivo*

Expression of *f5* and *f3a* mRNA was exclusively restricted to the follicular cells

of mature follicles (IVb) and was not found in the germ cells (oocytes) (Fig. 1B). In

contrast, *f3b* mRNA was expressed in both follicular cells and oocytes (Fig. 1B). The

expression of *f5, f3a*, and *f3b* in mature follicles (IVb), just prior to ovulation, collected at

06:40 were ~3,145-fold, ~120-fold, and ~2-fold higher than those in immature but fully-

grown follicles (IVa, collected at 12:00, Fig. 1C). Since the increase in *f3b* expression

was modest, we focused the rest of our efforts on *f5* and *f3a*. Results from whole mount

in situ hybridization (WISH) showed that both *f5* and *f3a* were nearly undetectable in

immature follicles (IVa, Fig. 1D, E). In contrast, strong signals of both *f5* and *f3a*

transcripts were detected in the follicular cells of the mature follicles (IVb) (Fig. 1D, E),

while no signal was detectable using sense probes (data not shown).

3.2 Progestin upregulates *f5* and *f3a* transcripts through Pgr

3.2a Transcripts of *f5* and *f3a* were primarily upregulated by progestin in
preovulatory follicles

The progestin (DHP) is a well-established native ligand for Pgr and an upstream

hormonal inducer for ovulation in zebrafish (13,31,36). As predicted, the increase of *f5*

and *f3a* transcripts were dose- and time-dependent in fully grown immature follicles

(IVa) exposed to external progestin (Fig. 2A-D).

Progestin increased *f5* expression in vitellogenesis follicles (III, ~7 fold) and

fully-grown immature follicles (IVa, ~337 fold) (Fig. 3A). In comparison, only fully-

grown immature follicles (IVa) responded to progestin with increase in *f3a* transcripts,

but not the early developing follicles (stage I-III, Fig. 3B). LH is the primary endocrine

factor for inducing ovulation in all animals including zebrafish (37). As expected,

increased *f5* expression (~4 fold) was observed in IVa immature follicles exposed to hCG

(50 IU/mL, Fig. 3C), but not for *f3a* (Fig. 3D). In comparison, progestin (DHP) had much

stronger stimulatory effects on the expression of *f5* (~318 fold) and *f3a* (~10 fold) than

hCG (Fig. 3C, D).

Significantly reduced *f5 and f3a* expression was found in mature preovulatory

follicles (IVb) collected from $pgr^{-/-}$ fish compared to those in wildtype siblings (Fig. 3E).

Importantly, DHP-induced *f5* and *f3a* expression were blocked in fully-grown immature

follicles (IVa) from $pgr^{-/-}$ fish (Fig. 3F).

3.2b Progestin modulates promoter activity of *f5* and *f3a* via Pgr

As expected (31), progestin induced dose-dependent, Pgr-mediated activation of

the MMTV promoters (Fig. 4A). In the presence of Pgr, progestin significantly increased

the *f5 and f3a* promoter activities, and this was dose-dependent (Fig. 4B, C). No change

of promoter activity was observed when pcDNA3.1(+)-Pgr was replaced by a control

vector (pcDNA3.1(+), Fig. 4B, C). The increase for *f5* was much bigger than that of the

f3a. In the presence of human PGRB, P4 also significantly increased human *F5* promoter

activity in a dose-dependent manner. No change in promoter activity was observed when

pCMV-HA-hPGRB was replaced by control (pCMV-HA, Fig. 4D). Moreover, site-

directed mutagenesis of four predicted progesterone response elements (PRE)

significantly decreased zebrafish *f5* promoter activity (Fig. 4E, Supplemental Table 5)

(25). A PRE (5'-AGGACATTTTTA-3') localizing at -1438 to -1425 of zebrafish *f5*

appears to be the primary mediator of Pgr regulation (Fig. 4E).

3.3 Coagulation factor contribution to ovulation

3.3a Follicular cells with strong Pgr expression localize adjacent to blood vessels

Intriguingly, we found almost all the follicular cells with strong Pgr expression were located adjacent to capillary vessel networks on the surface of the IVa follicles in a dual transgenic zebrafish line *Tg(pgr:eGFP/ fli1:DsRed)* (Fig. 5).

3.3b Rupture of blood vessels during ovulation

The ovulation process started with the appearance of a small hole in follicular cells and a ruptured blood vessel (indicated by a red arrow in Fig. 6A, supplemental video clip A) (25) with a concomitant appearance of the ovulating oocyte on the surface of the ovaries. When the hole of the broken follicular cell layer reached a certain size, the oocyte was quickly released from the follicular layer (Fig. 6B-E, supplemental video clip A) (25).

3.3c Increased numbers of erythrocytes appear in the lumen of blood vessels of mature preovulatory follicles as ovulation proceeds

The numbers of erythrocytes in capillaries of mature preovulatory follicles (IVb) increased as they proceeded to follicular rupture, from ~06:40, (Fig. 7A, D, C) to ~07:00 (Fig. 7B, E, G). This effect was absent in the same mature follicles (IVb) collected from anovulatory *pgr*$^{-/-}$ female siblings around 07:00 (Fig. 7C, F, G).

3.3d Heparin inhibits ovulation *in vitro*

The intact IVb follicles or ovulated follicles from *Tg(fli1:eGFP)* were easily recognizable by the interconnecting blood vessels with strong GFP expression and dramatic changes in size (Fig. 8A-D). As expected, the ovulation rate decreased to 41 %

when follicles were exposed to EDTA (2 mM), an inhibitor of metalloproteases that are

required for ovulation (35). Intriguingly, heparin (100 µg/mL), an anticoagulant,

significantly reduced the rate of ovulation to 66% (Fig. 8E). Another anticoagulant,

warfarin (50 µg/mL), had a reduced but not significant adverse effect on ovulation rate

(83 %) (Fig. 8E).

3.3e Reduced fertility in *f5*$^{+/-}$ zebrafish females *in vivo*

Although *f5* homozygous mutants live to early adulthood, most die by 4-6 months

of age and do not breed well (38). Therefore, we examined female fertility in *f5*$^{+/-}$

mutants in a consecutive spawning test. We observed significantly reduced numbers of

embryos per spawning and total embryos over two weeks in comparison to wildtype

sibling females. (Fig. 9A, B). The spawning frequency of *f5*$^{+/-}$ female fish also

significantly decreased compared to that of wildtype siblings (Fig. 9C). One *f5*$^{+/-}$ female

only spawned once in two weeks (Fig. 9C). The spawning intervals of *f5*$^{+/-}$ females were

significantly longer in comparison to that of wildtype siblings (Fig. 9D).

4.   Discussion

We have provided comprehensive evidence demonstrating that the expression of

coagulation factors *f5* and *f3a* in preovulatory follicles is controlled by progestin and its

cognate receptor Pgr. Increased expression of these coagulation factors is likely critical

for the regulation of ovulation and female fertility in addition to their well-established

roles in hemostasis. Our conclusions are based on the following evidence: 1) Coagulation

factors, *f5* and *f3a*, are expressed in the follicular cells and increase dramatically prior to

ovulation during natural spawning cycles in zebrafish; 2) Expression of these cofactors in

preovulatory follicles is controlled by progestin via Pgr, which are essential and upstream

regulators for ovulation; 3) Follicular cells with strong Pgr expression are adjacent to

blood capillaries; 4). As follicular cells near rupture, a significant increase in the numbers

of erythrocytes in blood vessels suggests blood clotting which is correlated with a

dramatic increase of *f5* and *f3a* in preovulatory follicular cells; 5) Inhibition of

coagulation leads to reduced ovulation; 6) Low fertility is found in $f5^{+/-}$ zebrafish female

mutants.

A major finding of the present study is the regulation of *f5* and *f3a* by progestin

via Pgr in the preovulatory follicles during ovulation (Fig.10). F3 is expressed in a wide

array of tissues (39). It has been demonstrated that physiological concentrations of P4

could increase both F3 mRNA and protein in breast cancer cell lines ZR-75 and T47D

(40). Increased F3 expression was correlated with an increase of procoagulant activity,

which was hypothesized to be responsible for the increased risk of thrombosis in breast

cancer patients (41). Studies of the uterus have also shown that RU486, an antagonist of

PGR, not only blocks but also reverses progestin-enhanced expression of F3 mRNA and

protein in stromal cells (42). In line with these studies, PGR binding sites were identified

in the promoter region of the human F3 gene (Supplemental Table 5) (25,43). The

increase of *f3a* expression in the preovulatory follicles is primarily controlled by Pgr

(current study). A difference in the weak activity of *f3a* promoters observed in a human

immortal cell line (HEK293T) *in vitro* in comparison to progestin-induced strong *f3a*

promoter activities found in zebrafish follicles both *in vivo* and *in vitro* may be due to

missing core *cis*-regulatory elements in the cloned promoter sequence of zebrafish *f3a*

(Supplemental Table 5) (25). Nevertheless, our results demonstrate that like *f5*, *f3a*

expression is also controlled by progestin via Pgr (Supplemental Table 5, Fig.10) (25).

Until now, no information was available for progestin regulation of F5. Estrogen

and progesterone receptors regulate *F5* expression in mouse bone marrow/bone (44,45).

The present study demonstrates that a new extrahepatic site of *f5* expression is regulated

by progestin through Pgr in zebrafish. Moreover, progestin enhances the promoter

activities of *f5* in zebrafish and F5 in humans via PGR. Several putative PREs were

predicted in the promoter regions of human, mouse, and zebrafish *F5* (Supplemental

Table 6) (25). Our site-direct mutation analyses of these PREs in the zebrafish *f5*

promoter suggests direct regulation of progestin/Pgr via binding to and activating these a

PREs. One PRE site (5'- TGACTCA-3') in the *f5* promoter is also conserved between

zebrafish, mice and humans (Supplemental Table 6) (25). Whether P4 and/or PGR

regulate *F5* expression in human ovaries *in vivo* will be an interesting topic for further

investigation.

The functions of coagulation factors in hemostasis are well established (46-48).

However, the role of coagulation factors during ovulation is significantly less studied.

Formation of blood clots at apical blood vessels was found shortly before follicular

rupture in several mammalian species (1,3-6). In addition, the permeability of blood

vessels increased 3-fold prior to ovulation in rats (49), which likely facilitates material

exchange between the follicles and blood vessels. The increase of material exchange

might be important for anticipating local but avoiding systemic activation of blood

clotting. Our observation of ruptured blood vessels on the surface of ovulating follicles

further supports this hypothesis. In addition, we also found a significant increase of

erythrocytes in capillaries on the surface of ovulating follicles as they progress to

rupture/ovulation, suggestive of blood clotting. Lack of functional F5 or F8 caused

various bleeding symptoms, including recurrent and serious ovulation-related blood loss

in women (50). Occurrence of blood clotting in advance of follicle rupture could

preemptively stop excessive bleeding caused by ovulation-induced capillary rupture. The

granulosa cells appear to be the primary cells for synthesizing F5 and F3a, as these

account for most that express Pgr (this study & (10,36)). Intriguingly, we also found that

strong Pgr expressing theca cells were localized near capillaries on the follicles. Whether

these theca cells are key switchers of F3a-initiated hemostasis in the capillaries around

ovarian follicles will be an interesting topic to examine in the future.

The dramatic induction of *f5* and *f3a* expression in preovulatory follicles via Pgr,

an essential mediator for activation of various ovulation-related genes, suggests that

locally produced F5 and F3a may be required for ovulation. Activation of multiple

metalloproteases within the ovulating follicles at the right times is a critical step for

sequential digestion of extracellular matrix proteins of the follicular cells that are required

for ovulation (1,37). Their activation and substrate binding are precisely regulated by the

timing of prodomain cleavage of the enzyme and removal of endogenous tissue inhibitors

of metalloproteases (Timps) (51). Protease inhibitors (Timps and other inhibitors) are

typically present at high concentrations in the serum to prevent metalloproteases from

acting accidentally (52). Vasoconstriction of blood vessels within the thecal layer at the

apex of the follicles was suggested to be critical for the removal of protease inhibition

from the circulation (53). We hypothesize that coagulation that occurs shortly before

ovulation in the follicles may have a similar function. In addition to the removal of

protease inhibitors from serum, the cleavage of propeptide from latent form of

metalloproteinase is required for activation and thus ovulation (Fig.10). An F3-dependent

pathway of thrombin generation was also shown in bovine, equine, and human follicular

fluid (7,54,55). Granulosa cells were shown to express and secrete fibrinogen, which is

catalyzed by thrombin to become fibrin (56). Fibrin formation could stimulate activities

of plasmin/plasminogen activators (plat and plau) signal, which is well-established for

plasminogen (plg) activation (57). Furthermore, the function of Plat, Plau, and Plg in

metalloproteinase activations during ovulation is also recognized (58,59). Therefore, we

hypothesize that a dramatic increase of F3a and F5 expression in the follicular cells may

facilitate the formation of fibrin, followed by the activation of Plat, Plau, and Plg, which

further activate metalloproteinases. Taken together, as blood coagulation occurs, blood

flow decreases, then stops completely, effectively terminating inhibitory effects of

protease inhibitors from the serum, triggering metalloprotease activations to start the

degradation of the follicular extracellular matrix for preparing the release of a mature

oocyte (Fig.10). A patient diagnosed with severe F5 deficiency had to undergo bilateral

ovariectomies because of the presence of hemorrhagic anovulatory follicles and life-

threatening hemorrhage in the pouch of Douglas (60). Circulating F3 has been shown to

be elevated in women with polycystic ovary syndrome, the most common form of

anovulatory infertility (9). Consistent with findings from medaka (35), the rate of

ovulation was suppressed by EDTA treatment in the current study likely via inhibition of

metalloproteases, or possibly inhibition of coagulation. EDTA is a known anticoagulant,

acting through chelation of calcium, as many coagulation factors depend on this ion for

function. Indeed, the known anticoagulant heparin also suppressed ovulation. The

anticoagulant warfarin did not, but this was unsurprising since its mechanism of action

usually requires days of treatment. Another finding in the present study is that reduced

fecundity and impaired ovulation were observed in $f5^{+/-}$ female zebrafish *in vivo*.

Together, these results argue for conserved functions of coagulation factors in the

regulation of ovulation in vertebrate ovaries. Long-term efforts in development of new

instruments and new techniques/methods, with suitable *in vivo* imaging and examining

the association of blood flow and timing of metalloproteases activation, are required for

further elucidation of underlying mechanisms and testing our hypothetical model

(Fig.10).

In summary, our results demonstrate that *f5* and *f3a* expression are dramatically

induced by progestin via a Pgr-dependent mechanism in preovulatory follicles (Fig. 10).

Moreover, these coagulation factors may regulate ovulation and female fertility in

addition to their roles in hemostasis. Future studies on the regulation and functions of

coagulation factors in other organisms including mammalian follicles or ovarian fluid

during ovulation will likely lead to developments of diagnostic and therapeutic modalities

for patients with gynecological problems associated with mutations and/or dysregulation

of coagulation.

Acknowledgments

We want to thank Dr. Mingyu Li (Xiamen University, China) and Dr. Yu Xue (Minnan Normal University, China) for providing zebrafish transgenic lines (*Tg(fli1:eGFP)*, *Tg(fli1:DsRed)*), Dr. Shuang-Bo Kong (Xiamen University, Xiamen, China) for sharing human PGRB expression vector (pCMV-HA-hPGRB). This work was supported by Fundamental Research Funds for the Central Universities (No. 20720200114), NSFC 31672628, and NSFC 41976092 to SXC, NIH GM100461 to YZ, and NIH R35 HL150784 to JAS.

Data Availability Statement(s): Some or all data generated or analyzed during this study
are included in this published article or in the data repositories listed in References 25.

References

1.   Duffy DM, Ko C, Jo M, Brannstrom M, Curry TE. Ovulation: Parallels With Inflammatory
Processes. *Endocr Rev*. 2019;40(2):369-416.
2.   Esmon CT. The interactions between inflammation and coagulation. *British Journal of*
Haematology. 2005;131(4):417-430.
3.   Kanzaki H, Okamura H, Okuda Y, Takenaka A, Morimoto K, Nishimura T. Scanning electron-
microscopic study of rabbit ovarian follicle microvasculature using resin injection corrosion
casts. *Journal of Anatomy*. 1982;134(4):697-704.
4.   Cavender JL, Murdoch WJ. Morphological-studies of the microcirculatory system of
periovulatory ovine follicles. *Biol Reprod*. 1988;39(4):989-997.
5.   Zackrisson U, Lofman CO, Janson PO, Wallin A, Mikuni M, Brannstrom M. Alterations of
follicular microcirculation and apex structure during ovulation in the rat. *Eur J Obstet Gyn R B*.
2011;157(2):169-174.
6.   Brannstrom M, Zackrisson U, Hagstrom HG, Josefsson B, Hellberg P, Granberg S, Collins WP,
Bourne T. Preovulatory changes of blood flow in different regions of the human follicle. *Fertil*
Steril. 1998;69(3):435-442.
7.   Gentry PA, Plante L, Schroeder MOB, LaMarre J, Young JE, Dodds WG. Human ovarian
follicular fluid has functional systems for the generation and modulation of thrombin. *Fertil*
Steril. 2000;73(4):848-854.
8.   Kim YS, Kim MS, Lee SH, Choi BC, Lim JM, Cha KY, Baek KH. Proteomic analysis of recurrent
spontaneous abortion: Identification of an inadequately expressed set of proteins in human
follicular fluid. *Proteomics*. 2006;6(11):3445-3454.
9.   Gonzalez F, Kirwan JP, Rote NS, Minium J. Elevated circulating levels of tissue factor in
polycystic ovary syndrome. *Clin Appl Thromb-Hem*. 2013;19(1):66-72.
10.   Natraj U, Richards JS. Hormonal-regulation, localization, and functional-activity of the
progesterone-receptor in granulosa-cells of rat preovulatory follicles. *Endocrinology*.
1993;133(2):761-769.
11.   Lydon JP, Demayo FJ, Funk CR, Mani SK, Hughes AR, Montgomery CA, Shyamala G, Conneely
OM, Omalley BW. Mice Lacking Progesterone-Receptor Exhibit Pleiotropic Reproductive
Abnormalities. *Gene Dev*. 1995;9(18):2266-2278.
12.   Hagiwara A, Ogiwara K, Katsu Y, Takahashi T. Luteinizing hormone-induced expression of
ptger4b, a Prostaglandin E-2 receptor indispensable for ovulation of the medaka Oryzias
latipes, is regulated by a genomic mechanism involving nuclear progestin receptor. *Biol*
Reprod. 2014;90(6):1-14.
13.   Zhu Y, Liu DT, Shaner ZC, Chen SX, Hong WS, Stellwag EJ. Nuclear progestin receptor (Pgr)
knockouts in zebrafish demonstrate role for Pgr in ovulation but not in rapid non-genomic
steroid mediated meiosis resumption. *Front Endocrinol*. 2015;6:37.
14.   Kubota K, Cui W, Dhakal P, Wolfe MW, Rumi MAK, Vivian JL, Roby KF, Soares MJ. Rethinking
progesterone regulation of female reproductive cyclicity. *P Natl Acad Sci USA*.
2016;113(15):4212-4217.
15.   Tang HP, Liu Y, Li JZ, Yin YK, Li GF, Chen Y, Li SS, Zhang Y, Lin HR, Liu XC, Cheng CHK. Gene
knockout of nuclear progesterone receptor provides insights into the regulation of ovulation
by LH signaling in zebrafish. *Sci Rep*. 2016;6:28545.
16.   Liu DT, Brewer MS, Chen SX, Hong WS, Zhu Y. Transcriptomic signatures for ovulation in
vertebrates. *Gen Comp Endocr*. 2017;247:74-86.
17.   Wissing ML, Kristensen SG, Andersen CY, Mikkelsen AL, Hst T, Borup R, Grondahl ML.
Identification of new ovulation-related genes in humans by comparing the transcriptome of
granulosa cells before and after ovulation triggering in the same controlled ovarian
stimulation cycle. *Hum Reprod*. 2014;29(5):997-1010.
18.   Carletti MZ, Christenson LK. Rapid effects of LH on gene expression in the mural granulosa
cells of mouse periovulatory follicles. *Reproduction*. 2009;137(5):843-855.
19.   Runic R, Schatz F, Wan L, Demopoulos R, Krikun G, Lockwood CJ. Effects of norplant on

endometrial tissue factor expression and blood vessel structure. *J Clin Endocr Metab*.
2000;85(10):3853-3859.
20.   Haas DM, Hathaway TJ, Ramsey PS. Progestogen for preventing miscarriage in women with
recurrent miscarriage of unclear etiology. *Cochrane Db Syst Rev*. 2019(11):1-50.
21.   Jick H, Jick SS, Gurewich V, Myers MW, Vasilakis C. Risk of Idiopathic Cardiovascular Death
and Nonfatal Venous Thromboembolism in Women Using Oral-Contraceptives with Differing
Progestagen Components. *Lancet*. 1995;346(8990):1589-1593.
22.   Farley TMM, Meirik O, Chang CL, Marmot MG, Poulter NR. Effect of different progestogens in
low estrogen oral contraceptives on venous thromboembolic disease. *Lancet*.
1995;346(8990):1582-1588.
23.   Zeberg H, Kelso J, Paabo S. The neandertal progesterone receptor. *Mol Biol Evol*.
2020;37(9):2655-2660.
24.   Stein C, Caccamo M, Laird G, Leptin M. Conservation and divergence of gene families
encoding components of innate immune response systems in zebrafish. *Genome Biol*.
2007;8(11):1-23.
25.   Huang J, Sun C, Liu DT, Zhao NN, Shavit JA, Zhu Y, Chen SX. Data from: Nuclear Progestin
Receptor Mediated Linkage of Blood Coagulation and Ovulation. *Figureshare*. 2021:
https://doi.org/10.6084/m9.figshare.17427326.
26.   Liu DT, Carter NJ, Wu XJ, Hong WS, Chen SX, Zhu Y. Progestin and nuclear progestin receptor
are essential for upregulation of metalloproteinase in zebrafish preovulatory follicles. *Front*
Endocrinol. 2018;9:517.
27.   Liu DT, Hong WS, Chen SX, Zhu Y. Upregulation of adamts9 by gonadotropin in preovulatory
follicles of zebrafish. *Mol Cell Endocrinol*. 2020;499:110608.
28.   Hanna RN, Zhu Y. Controls of meiotic signaling by membrane or nuclear progestin receptor in
zebrafish follicle-enclosed oocytes. *Molecular and Cellular Endocrinology*. 2011;337(1-2):80-
88.
29.   Schmittgen TD, Livak KJ. Analyzing real-time PCR data by the comparative C-T method. *Nat*
Protoc. 2008;3(6):1101-1108.
30.   Hauptmann G, Gerster T. Multicolor whole-mount in situ hybridization. *Methods Mol Biol*.
2000;137:139-148.
31.   Chen SX, Bogerd J, Garcia-Lopez A, de Jonge H, de Waal PP, Hong WS, Schulz RW. Molecular
Cloning and Functional Characterization of a Zebrafish Nuclear Progesterone Receptor. *Biol*
Reprod. 2010;82(1):171-181.
32.   Zhang YT, Hong WS, Liu DT, Qiu HT, Zhu Y, Chen SX. Involvement of membrane progestin
receptor Beta (mPRbeta/Paqr8) in sex pheromone progestin-induced expression of
luteinizing hormone in the pituitary of male chinese black sleeper (Bostrychus Sinensis).
Front Endocrinol. 2018;9:397.
33.   Detrich HW, Kieran MW, Chan FY, Barone LM, Yee K, Rundstadler JA, Pratt S, Ransom D, Zon LI.
Intraembryonic hematopoietic-cell migration during vertebrate development. *P Natl Acad Sci*
USA. 1995;92(23):10713-10717.
34.   Hanumanthaiah R, Thankavel B, Day K, Gregory M, Jagadeeswaran P. Developmental
expression of vitamin K-dependent gamma-carboxylase activity in zebrafish embryos: effect
of warfarin. *Blood Cells Mol Dis*. 2001;27(6):992-999.
35.   Ogiwara K, Takano N, Shinohara M, Murakami M, Takahashi T. Gelatinase A and membrane-
type matrix metalloproteinases 1 and 2 are responsible for follicle rupture during ovulation
in the medaka. *Proc Natl Acad Sci U S A*. 2005;102(24):8442-8447.
36.   Hanna RN, Daly SCJ, Pang YF, Anglade I, Kah O, Thomas P, Zhu Y. Characterization and
expression of the nuclear progestin receptor in zebrafish gonads and brain. *Biol Reprod*.
2010;82(1):112-122.
37.   Takahashi T, Hagiwara A, Ogiwara K. Follicle rupture during ovulation with an emphasis on
recent progress in fish models. *Reproduction*. 2019;157(1):R1-R13.
38.   Weyand AC, Grzegorski SJ, Rost MS, Lavik KI, Ferguson AC, Menegatti M, Richter CE, Asselta R,
Duga S, Peyvandi F, Shavit JA. Analysis of factor V in zebrafish demonstrates minimal levels

591   needed for early hemostasis. *Blood Adv*. 2019;3(11):1670-1680.

592   39.   Osterud B, Bjorklid E. Sources of tissue factor. *Semin Thromb Hemost*. 2006;32(1):11-23.

593   40.   Kato S, Pinto M, Carvajal A, Espinoza N, Monso C, Sadarangani A, Villalon M, Brosens JJ, White
594   JO, Richer JK, Horwitz KB, Owen GI. Progesterone increases tissue factor gene expression,
595   procoagulant activity, and invasion in the breast cancer cell line ZR-75-1. *J Clin Endocr Metab*.
596   2005;90(2):1181-1188.

597   41.   Richer JK, Jacobsen BM, Manning NG, Abel MG, Wolf DM, Horwitz KB. Differential gene
598   regulation by the two progesterone receptor isoforms in human breast cancer cells. *J Biol*
599   *Chem*. 2002;277(7):5209-5218.

600   42.   Lockwood CJ, Krikun G, Papp C, Aigner S, Nemerson Y, Schatz F. Biological mechanisms
601   underlying ru-486 clinical effects - inhibition of endometrial stromal cell tissue factor
602   content. *J Clin Endocr Metab*. 1994;79(3):786-790.

603   43.   Bravo ML, Pinto MP, Gonzalez I, Oliva B, Kato S, Cuello MA, Lange CA, Owen GI. Progesterone
604   regulation of tissue factor depends on MEK1/2 activation and requires the proline-rich site
605   on progesterone receptor. *Endocrine*. 2015;48(1):309-320.

606   44.   Moverare S, Skrtic S, Lindberg MK, Dahlman-Wright K, Ohlsson C. Estrogen increases
607   coagulation factor V mRNA levels via both estrogen receptor-alpha and -beta in murine bone
608   marrow/bone. *Eur J Endocrinol*. 2004;151(2):259-263.

609   45.   Cleuren AC, Van der Linden IK, De Visser YP, Wagenaar GT, Reitsma PH, Van Vlijmen BJ.
610   17alpha-Ethinylestradiol rapidly alters transcript levels of murine coagulation genes via
611   estrogen receptor alpha. *J Thromb Haemost*. 2010;8(8):1838-1846.

612   46.   Spronk HM, Govers-Riemslag JW, ten Cate H. The blood coagulation system as a molecular
613   machine. *BioEssays : news and reviews in molecular, cellular and developmental biology*.
614   2003;25(12):1220-1228.

615   47.   Esmon CT. The impact of the inflammatory response on coagulation. *Thromb Res*.
616   2004;114(5-6):321-327.

617   48.   Hathcock JJ. Flow effects on coagulation and thrombosis. *Arterioscler Thromb Vasc Biol*.
618   2006;26(8):1729-1737.

619   49.   Mitsube K, Brannstrom M, Haraldsson B. Modulation of microvascular permeability in the
620   preovulatory rat ovary by an ovulatory gonadotropin stimulus. *Fertil Steril*. 2013;99(3):903-
621   909.

622   50.   Spiliopoulos D, Kadir RA. Congenital factor V and VIII deficiency in women: a systematic
623   review of literature and report of two new cases. *Blood Coagul Fibrin*. 2016;27(3):237-241.

624   51.   Zhu Y. Metalloproteases in gonad formation and ovulation. *Gen Comp Endocr*.
625   2021;314:113924.

626   52.   Travis J, Salvesen GS. Human plasma proteinase inhibitors. *Annu Rev Biochem*. 1983;52:655-
627   709.

628   53.   Migone FF, Cowan RG, Williams RM, Gorse KJ, Zipfel WR, Quirk SM. In vivo imaging reveals an
629   essential role of vasoconstriction in rupture of the ovarian follicle at ovulation. *Proc Natl*
630   *Acad Sci U S A*. 2016;113(8):2294-2299.

631   54.   Yamada M, Gentry PA. The hemostatic profile of equine ovarian follicular fluid. *Thrombosis*
632   *Research*. 1995;77(1):45-54.

633   55.   Roach LE, Petrik JJ, Plante L, LaMarre J, Gentry PA. Thrombin generation and presence of
634   thrombin receptor in ovarian follicles. *Biol Reprod*. 2002;66(5):1350-1358.

635   56.   Parrott JA, Whaley PD, Skinner MK. Extrahepatic expression of fibrinogen by granulosa cells:
636   potential role in ovulation. *Endocrinology*. 1993;133(4):1645-1649.

637   57.   Medved L, Nieuwenhuizen W. Molecular mechanisms of initiation of fibrinolysis by fibrin.
638   *Thromb Haemost*. 2003;89(3):409-419.

639   58.   Beers WH, Strickland S, Reich E. Ovarian plasminogen activator: Relationship to ovulation
640   and hormonal regulation. *Cell*. 1975;6(3):387-394.

641   59.   Ogiwara K, Minagawa K, Takano N, Kageyama T, Takahashi T. Apparent involvement of
642   plasmin in early-stage follicle rupture during ovulation in medaka. *Biol Reprod*.
643   2012;86(4):113.

644   60.   Montefusco MC, Duga S, Asselta R, Malcovati M, Peyvandi F, Santagostino E, Mannucci PM,
645   Tenchini ML. Clinical and molecular characterization of 6 patients affected by severe
646   deficiency of coagulation factor V: broadening of the mutational spectrum of factor V gene
647   and in vitro analysis of the newly identified missense mutations. *Blood*. 2003;102(9):3210-
648   3216.
649

Figure

Figure 1. Dramatic increase of *f5* **and** *f3a* **transcripts in follicular cells of preovulatory follicles during**

a daily spawning cycle *in vivo***.**

 (**A**) Schematic illustration of daily spawning cycle in zebrafish synchronized with lights on. Oocyte

maturation in zebrafish typically occurs around 06:40, while ovulation occurs around 07:00.

(**B**) Cell-specific expression of *f5*, *f3a, and f3b* in mature follicles examined by RT-PCR. Transcripts of *f5*

and *f3a* were expressed in the follicular cells but not in the oocytes, while the transcript of *f3b* was

expressed in both follicular cells and oocytes of mature preovulatory follicles. MK: molecular makers; I:

intact follicles including both surrounding follicular cells and germ cell; O: denuded oocyte (without

follicular cells); F: follicular cells (with no germ cell); N: no template control.

(**C**) Dramatic increase of *f5, f3a,* and *f3b* transcripts in preovulatory follicles during a daily spawning cycle

in vivo. Fully-grown ovarian follicles (stage IVa or IVb) were collected from wildtype mature female

zebrafish at four representative timepoints during a daily spawning cycle (see Fig. 1A). Expression of *f5,*

f3a, and *f3b* mRNA were determined by qPCR and normalized to the internal housekeeping gene (*ef1α*).

Data are expressed as the mean ± SEM (N = 4) relative to the respective transcript levels of *f5* in IVa

follicles collected at 12:00. P-values were calculated by one-tailed one-way ANOVA followed by Dunnett's

test against respective controls (samples collected at 12:00).

(**D-G**) Dramatic increase of *f5* and *f3a* transcripts in the follicular cells of mature follicles. Fully-grown but

immature (stage IVa) and mature follicles but prior to ovulation (stage IVb) were collected at 20:00 and

06:40, respectively. Strong signals of *f5* and *f3a* transcripts were only observed in mature follicles (IVb) by

whole-mount *in situ* hybridization (WISH), while no signals were observed in the immature follicles (IVa,

D, E). The positive ISH signals were localized in the follicular cells (F, G). Scale bar, 250 µm (D, E) and

50 µm (F, G).

Figure 2. Transient increase of *f5* and *f3a* expression in preovulatory follicles in response to

stimulation of progestin *in vitro*.

(**A, C**) Effects of various doses of progestin (DHP) (2 hours exposure) on *f5* and *f3a* expression in fully

grown immature preovulatory follicles (stage IVa) *in vitro*. (**B, D**) Effects of DHP (100 nM) on *f5* and *f3a*

expression in preovulatory follicles (IVa) at various timepoints *in vitro*. P-values were calculated by one-

tailed one-way ANOVA followed by Dunnett's test against respective controls (0 timepoint or vehicle

control).

Figure 3. Increase of *f5* and *f3a* transcripts in late-stage follicles is mediated by Pgr.

Follicles were incubated with progestin (DHP, 100nM), or LH (hCG, 50 IU/ml) alone, or in combination

for 2 hrs at 28 ℃ *in vitro* before qPCR analysis. (**A**) An increase of *f5* transcript was observed in late-stage

(III and IVa) follicles from wildtype (*pgr*$^{+/+}$) zebrafish, but not in early-stage follicles (I or II) from

wildtype zebrafish, in response to progestin stimulation. (**B**) An increase of *f3a* expression was observed

only in fully-grown immature follicles (IVa) from wildtype (*pgr*$^{+/+}$) zebrafish, but not in developing

follicles (I-III) from wildtype fish, in response to progestin stimulation. (**C**) Synergistic increase of *f5*

mRNA by hCG and DHP in fully grown immature follicles (IVa) from wildtype zebrafish. (**D**) Progestin

but not hCG stimulated the expression of *f3a* mRNA in full-grown immature follicles (IVa) from wildtype

zebrafish. (**E**) Both *f3a* and *f5* transcripts in preovulatory follicles (stage IVb) of *pgr*$^{-/-}$ were significantly

lower than that of *pgr*$^{+/+}$ (WT). (**F**) Stimulatory effect of progestin (DHP) for *f3a* and *f5* expression was

abolished in the stage IVa follicles of *pgr*$^{-/-}$ mutants. The expression of *f5* and *f3a* transcripts were

determined by qPCR and normalized to an internal control (*ef1α*). Data are expressed as the mean ± SEM

(N = 3) relative to the respective transcript levels of target gene measured in wildtype, or vehicle treatment

group. P-values were calculated by two-tailed Students' t-test.

Figure 4. Progestin enhances promoter activity of *f5* and *f3a* via Pgr.

HEK293T cells were transiently co-transfected with a firefly luciferase reporter vector containing putative

Pgr binding elements (3 μg/vector/60mm petri dish), a pRL-TK vector (150 ng/60mm petri dish)

containing the Renilla luciferase reporter gene (as a control for transfection efficiency), and a Pgr

expression vector (1.5 µg/vector/60mm petri dish) or an empty control expression vector (1.5

µg/vector/60mm petri dish). In the presence of zebrafish Pgr (zPgr), progestin significantly increased

MMTV (panel **A**), *zf5* (panel **B**), and *zf3a* (panel **C**) promoter activities. This increase was dose-dependent.

In the presence of human PGRB (hPGRB), P4 also significantly increased human *hF5* promoter activity,

and the increase was dose-dependent (Panel **D**). Site-directed mutagenesis of four likely progestin receptor

response elements (PRE) decreased the *zf5* promoter activity (Panel **E**). HEK293T cells were incubated for

24 hrs with increasing concentrations of progestins (DHP or P4: 100 pM to 1 µM). Extracts of the treated

cells were assayed for luciferase activity. Values are shown relative to the luciferase activity of the vehicle

treatment group. Data are expressed as mean ± SEM (N = 3). P-values were calculated by one-tailed one-

way ANOVA followed by Dunnett's test against respective controls (vehicle controls, zero dose).

Figure 5. Follicular cells with strong Pgr expression are adjacent to blood capillaries.

Images of a representative stage IVa follicle from a double transgenic zebrafish line *Tg(pgr:eGFP/*

fli1:DsRed) are shown. (**A**) Nucleus staining of follicular cells by Hoechst 33342. (**B**) Pgr promoter drives

GFP expression in follicular cells. (**C**) Red fluorescence-labeled vascular endothelial cells on the follicles.

(**D**) Merged images to show that follicular cells with strong GFP expression driven by the Pgr promoters

are adjacent to blood capillaries. Red arrowheads indicate follicular cells with strong GFP expression

driven by the Pgr promoters. Scale bars: 100 μm.

Figure 6. Rupture of blood capillaries and follicular cells during ovulation.

Ovaries undergoing ovulation were collected from *Tg(fli1:eGFP)* females at 07:00. (**A-F**), a series of

images were taken when a mature (stage IVb) ovarian follicle started ovulation. Interconnected blood

vessels labeled with GFP were observed easily on the surface of ovarian follicles. A red arrow indicates a

ruptured blood vessel on the surface of an ovulating follicle. Red arrowheads and red dotted lines indicate a

hole left in the remaining follicular cells after ovulation. The white dotted line represents the remaining

730 follicular cells. Scale bars: 250 μm. Oc, oocyte. (Please also see supplemental video clip A for additional

731 details) (25).

Figure 7. Increase of erythrocytes in capillaries on the surface of mature follicles during progression but prior to rupture.

(**A, D, B, E**) Erythrocytes within the surface capillaries of mature follicles (IVb) from wildtype ($pgr^{+/+}$) female zebrafish collected at 6:40 or 7:00 were visualized using *o*-dianisidine staining as follicles progressing towards rupture and ovulation. (**C, F**) Erythrocytes were also visualized in similar mature follicles that were collected at 7:00 from $pgr^{-/-}$ females, in which ovulation never occurs (13). (**G**) At least 30 mature follicles were randomly chosen from at least 3 different females of each condition, and the number of erythrocytes in capillaries were counted by a blinded observer. Data are expressed as mean ± SEM (N ≥ 30). P-values were calculated by two-tailed Student's t-test. Scale bars: 500 μm (A-C) or 100 μm (D-F).

Figure 8. Heparin inhibits ovulation *in vitro*.

Ovulation rate was determined using mature follicles (IVb) isolated from transgenic female zebrafish (*Tg(fli1:eGFP)*). The intact IVb follicles were easily recognizable by the interconnecting blood vessels with strong GFP expression (**A, C**). After ovulation, oocytes were easily recognizable by the absence of interconnecting GFP-positive vessels and expansion of the vitelline envelope (chorion) (**B, D**). Follicular layers separated from ovulated oocytes could also be easily distinguished due to interconnecting blood vessels and a dramatic size reduction (**B, D**). (**A, C**) Mature follicles (IVb) were collected from *Tg(fli1:eGFP)* female zebrafish. (**B, D**) Representative images of ovulated mature follicles were taken 2 hrs after incubation at 28°C. Arrows indicate the remaining follicular layer following ovulation. Arrowheads indicate expanded vitelline envelopes. Oc, oocyte. (**E**) mature follicles were incubated in a medium containing EDTA (2 mM), heparin (100 µg/mL), warfarin (50 µg/mL), or vehicle control, and rates of ovulation were determined 2 hrs following incubation. Data are expressed as the mean ± SEM (N ≥ 7). P-values were calculated by two-tailed Student's t-test. Scale bars: 1 mm (A, B) or 250 µm (C, D).

Figure 9. Reduced fecundity in *f5*⁺/⁻ **female zebrafish.**

(**A**) Reduced numbers of live embryos per spawning for each *f5*⁺/⁻ female in comparison to that of wildtype

(*f5*⁺/⁺) females. (**B**) Reduced total numbers of embryos for each *f5*⁺/⁻ female fish in comparison to wildtype

(*f5*⁺/⁺) females. (**C**) Reduced spawning times in *f5*⁺/⁻ females in comparison to that of wildtype (*f5*⁺/⁺)

females. (**D**) Increased spawning intervals in *f5*⁺/⁻ females in comparison to wildtype (*f5*⁺/⁺) females. Data

are expressed as the mean ± SEM (N ≥ 14). P-values were calculated by two-tailed Student's t-test.

Figure 10. A proposed model for nuclear progestin receptor (Pgr) mediated upregulation of *f5* and *f3a* in blood coagulation that facilitates ovulation in zebrafish.

Luteinizing hormone (LH) induces ovulation via luteinizing hormone receptor (Lhcgr) in preovulatory follicles, namely in stage IV follicles in zebrafish. LH signaling is mainly mediated by progestin (DHP, i.e., 17α,20β-dihydroxy-progesterone, a native ovulation inducing steroid in zebrafish) and its receptor (Pgr) in the production and activation of metalloproteases. The illustration on the left side is showing the upregulation of metalloproteases in preovulatory follicles, to be more specific, during the transition from stage IVa to IVb follicles, which is an important preparation step for ovulation (16,26,27). However, these proteases are produced in a latent form, i.e., inactive form of metalloproteases (iMPs, see Yong Zhu, 2021 for detail), and inhibited by tissue inhibitors of metalloproteases (TIMP) presented at high levels in and delivered by blood capillaries to the follicles. The illustration on the right side is showing a dramatic increase of coagulation factors *f5* and *f3a* shortly before ovulation, specifically in the late stage of IVb follicles, which are also mainly upregulated by progestin via Pgr. These ovarian follicles derived, and locally produced coagulation factors (F5 and F3a) are imported into inside blood vessels messed in the follicles and activated via yet to be defined mechanisms. Once F5 and F3a are activated, these coagulation factors accelerate blood coagulation, which in turn stops blood flow and therefore effectively terminates TIMP inhibition of metalloproteases from the bloodstream. In addition, the production of fibrin (not

782   shown) due to the formation of blood clots stimulates activities of plasmin/plasminogen activators, which

783   are well-known inducers for metalloprotease activation. The combined force of these two pathways triggers

784   rapid activation of metalloproteases, breakdown of extracellular matrix and basal membrane, rupture of

785   follicular cell layers, the final release of a mature oocyte.