1

The *C. elegans* gonadal sheath Sh1 cells extend asymmetrically over a differentiating germ cell population in the proliferative zone

Xin Li[1], Noor Singh[1], Camille Miller[1], India Washington[1], Bintou Sosseh[1], Kacy Lynn Gordon[1,*]

[1]Department of Biology, The University of North Carolina at Chapel Hill, Chapel Hill, NC, 27599, USA
*Correspondence: kacy.gordon@unc.edu
Department of Biology, CB #3280, University of North Carolina at Chapel Hill, Chapel Hill, NC, 27599, USA

Abstract

The *C. elegans* adult hermaphrodite germ line is surrounded by a thin tube formed by somatic sheath cells that support germ cells as they mature from the stem-like mitotic state through meiosis, gametogenesis and ovulation. Recently, we discovered that the distal Sh1 sheath cells associate with mitotic germ cells as they exit the niche (Gordon et al., 2020). Here we report that these sheath-associated germ cells differentiate first in animals with temperature-sensitive mutations affecting germ cell state, and stem-like germ cells are maintained distal to the Sh1 boundary. We analyze several markers of the distal sheath, which is best visualized with endogenously tagged membrane proteins, as overexpressed fluorescent proteins fail to localize to distal membrane processes and can cause gonad morphology defects. However, such reagents with highly variable expression can be used to determine the relative positions of the two Sh1 cells, one of which often extends further distal than the other.

Introduction

The *C. elegans* hermaphrodite gonad is a fruitful system in which to study organogenesis, meiosis, and stem cell niche biology. Recent work from our group (Gordon et al., 2020), used two endogenously tagged alleles of genetically redundant innexin genes *inx-8* and *inx-9* to visualize the somatic gonadal sheath of the *C. elegans* hermaphrodite. We discovered that the distal most pair of sheath cells, called Sh1, lies immediately adjacent to the distal tip cell (DTC), which is the stem cell niche of the germ line stem cells. Previously (based on electron microscopy and on cytoplasmic GFP overexpression from transgenes active in the sheath (*lim-7p::GFP)* (Hall et al., 1999) or its progenitor cells (*lag-2p::GFP*) (Killian and Hubbard, 2005)), Sh1 cells were thought to associate only with germ cells well into the meiotic cell cycle, so our finding required a reimagining of the anatomy of the distal gonad.

Here, we confirm that the Sh1 cells fall at the boundary of a population of germ cells in a stem-like state, report other markers that label the Sh1 cells, and verify that these markers can be used to assess gonad anatomy without unduly impacting the gonad itself. We also discuss reagents that are not suitable markers of Sh1 cells, including an overexpressed, functional cell death receptor that is used to mark Sh1 in a recent study (Tolkin et al., 2021). Finally, we consider best practices for using endogenously tagged proteins for cell and developmental studies.

Results

Distal Sh1 associates with the population of germ cells that differentiate first when progression through mitosis is halted or Notch signaling is lost

Our first experiment addresses in a new way the question of what type of germ cells associate with the distal Sh1 cell. The DTC expresses the Notch ligand LAG-2, which is necessary to maintain the germ line stem cell pool (Henderson et al., 1994). Recent work has shown that the active transcription of Notch targets *sygl-1* and *lst-1* (Lee et al., 2019) and the accumulation of their proteins (Shin et al., 2017) is restricted to the distal-most germ cells, describing a population of stem-like germ cells ~6-8 germ cell diameters (~25-35 μm) from the distal end of the gonad. Our recent work (Gordon et al., 2020) reported that the position of Sh1 coincides with *sygl-1* promoter's activation boundary on one side and the accumulation of the meiotic entry protein GLD-1 on the other, consistent with the hypothesis that the distal edge of Sh1 falls at the boundary of that stem-like cell population, ~30 μm from the distal end of the gonad.

A similarly positioned stem-like germ cell population was found in earlier work that used temperature sensitive alleles to perturb germ cell fate or progress through the cell cycle. The readout was germ cell fate as determined in one of two ways. Anti-phosphohistone H3 staining of proliferative cells and GLD-1 antibody costaining for cells accumulating meiotic factors shows where cells are dividing and beginning to differentiate, respectively. Alternatively, the "transition zone" in germ cell nuclear morphology between "mitotic" and "meiotic" zones can be visualized by the presence of crescent-shaped nuclei of meiotic prophase observed by DAPI staining. We used this latter method of visualizing nuclear morphology.

We repeated these experiments in strains that have tagged innexins to mark the distal sheath Sh1 cells to ask which population(s) of germ cells are associated with Sh1. Here, we describe the original findings and their interpretations, and then our new findings.

First, an *emb-30* temperature-sensitive allele is known to cause germ cell division to arrest at the metaphase-anaphase transition, thus halting the distal-to-proximal movement of germ cells that is driven by the proliferation of more distal cells (Cinquin et al., 2010). In a wild-type gonad, germ cells differentiate (enter and progress through the meiotic cell cycle) as they move from distal to proximal (Figure 1A, left). In *emb-30(ts)* gonads, a shift to the restrictive temperature causes proliferation to halt and germ cells to remain stationary within the gonad (Figure 1A, right). These cells can now differentiate in place—or remain in the undifferentiated state—depending on their exposure to the stemness cue. Germ cells that remain in the niche at the distal end of the gonad do not differentiate after 15 hours at the restrictive temperature, while more proximal germ cells do differentiate. Nuclear morphology differs between these two regions of the germ line.

We hypothesized that the transition in nuclear morphology in *emb-30(tn377)* animals would shift proximally after 15 hours at the restrictive temperature (as had previously been observed), to ultimately fall at the distal position of the Sh1 cell as visualized by mKate::INX-8; we hypothesize that the position of the Sh1 cell would itself not be affected by the temperature shift. Indeed, this is what we found (Figure 1B, E), supporting our prior conclusion that there is germ cell fate asymmetry across the DTC-Sh1 boundary. These results are independent of culture time of controls (Figure 1—Figure Supplement 1). The Sh1 cell covers proliferative germ cells outside the niche that are closer to differentiating than those under the DTC.

The second set of experiments using temperature sensitive alleles to reveal differences in germ cell fate along the distal-proximal axis uses *glp-1(ts)* alleles to stop Notch signal transduction and observe where and when germ cells acquire features of differentiation (Figure 1C). The same study (Cinquin et al., 2010) found that the *glp-1(q224)* temperature sensitive allele reared at the restrictive temperature over a 9-hour time course showed a progressively shrinking mitotic zone (as assessed by nuclear morphology of DAPI stained gonads) until hour ~5.5, at which time the remaining distal most ~5 rows of germ cells differentiate as a pool. A subsequent study (Fox and Schedl, 2015) used a similar approach (but with the *glp-1(bn18)* temperature sensitive allele, slightly different timing, and antibody staining to determine cell fate) and found a similar result, and additionally discovered that progress through the cell cycle influenced the precise timing of germ cell differentiation.

We used the *glp-1(bn18)* allele to deactivate Notch signaling in a strain with GFP::INX-9 to visualize the Sh1 cells and the fluorescent histone marker *naSi2(mex-5p::H2B::mCherry)* to visualize germ cell nuclei (Figure 1D, E). We hypothesized that after six hours at the restrictive temperature, only the distalmost pool of stem-like cells will not have taken on the crescent-shaped nuclear morphology of meiotic germ cells, while the germ cells under Sh1 will have entered meiosis. We predicted that the Sh1 cell would not change its position across this time interval. Indeed, this is what we found, further supporting our hypothesis that the Sh1-associated germ cells are closer to differentiation than those under the DTC, which are the last to differentiate.

Results from these temperature sensitive mutants confirm what the markers of germ cell fate revealed in Gordon et al. (2020), which is that the Sh1 cell associates with germ cells in the

111 proliferative zone that have left the stem cell niche and are on the path to differentiation, while the
112 stem-like germ cells lie immediately distal to the Sh1 cell at its interface with the DTC.
113
Different endogenously tagged membrane proteins reveal a distal position of Sh1
These experiments made use of the endogenously N-terminal tagged *inx-8(qy78[mKate::inx-8])* and
inx-9(qy79[GFP::inx-9]) alleles (Figure 2A and B) generated by (Gordon et al., 2020). Both tagged
proteins are highly specific for the somatic gonad throughout development; in the adult, their
expression differentiates, with INX-8 signal diminishing from the DTC and INX-9 signal persisting (see
white DTC outline in Figure 1D). We have since identified additional endogenous fluorescent-protein-
tagged alleles that show expression in the gonadal sheath cells and localize in or near the cell
membrane. One of these, *ina-1(qy23[ina-1::mNeonGreen])* (Figure 2C) was briefly reported in
(Gordon et al., 2020). We found another that marks the sheath, *cam-1(cp243[cam-*
1::mNeonGreen])(Heppert et al., 2018) (Figure 2D). For both tagged innexins as well as *ina-1::mNG*
and *cam-1::mNG*, we find that the Sh1 cell has a distal boundary that either displays a measurable
interface with the DTC or is so located as to be consistent with such a boundary (where the DTC is
not marked by the endogenous protein). The position of this boundary (< 40 µm, or ~ 8 germ cell
diameters) coincides with the domain in which germ cells leave the stem cell niche (Lee et al., 2019)
(Figure 2E). We have not yet found a counterexample of an endogenously tagged, membrane-
associated protein in Sh1 that demarcates an apparent Sh1 cell boundary at a great distance from
the distal end of the gonad in young adults.
131
Overexpressed transgenic markers vary in distal position and expression levels
Three integrated array transgene markers that drive overexpression of fluorescent proteins in the
sheath were also analyzed. The first is a *lim-7* promoter-driven cytoplasmic GFP that was used to
label the Sh1 cell in a foundational study of the *C. elegans* hermaphrodite gonad, *tnIs6[lim-7::GFP]*
(Hall et al., 1999) (Figure 2F). The second is a *lim-7* promoter-driven functional cell death receptor
tagged with GFP, *bcIs39[lim-7p::ced-1::GFP]* (Zhou et al., 2001), which is the basis of a recent study
that reports a more proximal boundary of Sh1 (Figure 2G, strain DG5020 (Tolkin et al., 2021)). The
third is a *lim-7* promoter-driven membrane-localized GFP made by us to mark the sheath cell
membrane without tagging an endogenous protein, *rlmIs5[lim-7p::GFP::CAAX]* (Figure 2H). The
ranges of the distal edge of GFP localization for all three strains overlaps with what we observed for
the four endogenously tagged proteins, but are far more variable, as overexpressed transgenes are
known to be (Evans, 2006) (Figure 2E-H, and Figure 2—Figure Supplement 1). Patterns are more
similar at earlier developmental stages (Figure 2—Figure Supplement 2).
145
To untangle this variance, we examined individual worms for evidence of a DTC-Sh1 interface. About
half of the scoreable *lim-7p::ced-1::GFP* gonads (strain DG5020) show a DTC-Sh1 interface, and half
show a bare region (Figure 3A). We further broke down this dataset by fluorescence intensity of distal
CED-1::GFP signal. Strikingly, among animals under a threshold of expression intensity of ~400 A.U.
(less than 1/3 as bright as the brightest GFP samples), the incidence of a DTC-Sh1 interface was
100% (10/10, as opposed to 15/30 for the whole dataset, Figure 3A). On the other extreme, gonads
with stronger CED-1::GFP signal were more likely to have a farther proximal boundary of CED-
1::GFP localization. In samples for which CED-1::GFP signal terminates at a great distance from the
distal end of the gonad, there are two possible explanations. Either in those animals, the Sh1 position
is farther proximal than in animals with other markers, or else CED-1::GFP fails to localize to the edge
of the Sh1 cell pair.
157
Expression differences between Sh1 cells in a pair can conceal distal extent of the sheath
We observed a pattern in a subset of gonads where the two Sh1 cells of a pair had dramatically
different levels of CED-1::GFP signal, and these cells had different terminal positions on the distal-
proximal axis (Figure 3B and 3B'). Exposure time and excitation laser power during image acquisition
and subsequent scaling of the resulting image determine whether or not the signal in the lowly
expressing cell is readily apparent (Figure 3B vs 3B'). In some cases, the brightness of the other Sh1
cell and the nearby proximal gonad makes the dimmer Sh1 cell nearly impossible to detect. Variable
expression levels and even complete silencing of *C. elegans* transgenes are well-known phenomena

(Evans, 2006). It was not known, however, that the two Sh1 cells of a pair could assume such
different configurations over the distal germline (Figure 3C, and see Figure 2H for the same pattern in
the *lim-7p::GFP::CAAX* transgenic strain).
169
The Sh1 positions become even more clear when *lim-7p::ced-1::GFP* is coexpressed with the mKate-
tagged innexin *inx-8(qy78)* in strain DG5131 (Figure 3D and 3E). These markers colocalize in a
substantial fraction of animals, as has been reported recently ((Tolkin et al., 2021), see Figure 2
Supplement 2 therein). In the animals that have a discrepancy between GFP and mKate localization
in Sh1, the difference in expression reveals an unexpected cell boundary between the two Sh1 cells.
We imaged 19 gonads from the coexpressing strain DG5131 through their full thickness. Of those,
4/19 had severe gonad morphology defects (see next section). Of the 15 morphologically normal
gonads, 6/15 had discrepancies in CED-1::GFP and mKate::INX-8 signal. In 3/6 such cases, one Sh1
cell makes up the entire DTC-Sh1 interface, with the other terminating at a greater distance from the
distal end. In the other 3/6 of cases, both Sh1 cells border the DTC. Fluorescence signal from
mKate::INX-8 alone does not allow these cell borders to be detected because that marker is more
consistently expressed across the Sh1 cells (Figure 3F).
182
The variability of the *lim-7p::ced-1::GFP* transgene allowed us to perform something like a mosaic
analysis when the two Sh1 cells have very different expression levels but the dimmer cell is still
visible (N=31/53 morphologically normal DG5020 gonads imaged to full depth, Figure 3 Supplement
1A-D). Where the borders of the two Sh1 cells can be distinguished, one cell extends at least 20 μm
farther distal than the other in 23/31 cases; five additional gonads have expression in only one Sh1
cell that terminates at a great distance (>70 μm) from the distal end. The edges of dimly expressing
Sh1 cells can be difficult to resolve. A similar phenomenon was observed when the cytoplasmic GFP
of *tnIs6[lim-7p::GFP]* was coexpressed with *qy78[mKate::inx-8]* ((Gordon et al., 2020) Figure 1
Supplement 1 therein). Of note, the N-terminal mKate::INX-8 and GFP::INX-9 tags are most likely
extracellular based on the innexin-6 channel structure determined by cryo-EM (Oshima et al., 2016),
so there is reason to suspect their localization at the cell membrane will be regulated differently than
that of intracellular GFPs.
195
Additionally, we noticed that in DG5131 gonads where the two Sh1 cells have very different CED-
1::GFP expression levels, sometimes mKate::INX-8 is missing from the membrane in Sh1 cells with
strong CED-1::GFP signal (Figure 3 Supplement 1E and 1F). Subtracting background, we find that
there is a 50% reduction in tagged INX-8 in such membrane regions. Since mKate::INX-8 is a
genomically encoded, functional protein, such disruption likely impacts endogenous protein function.
This observation hints at a synthetic interaction between the two fluorescent sheath membrane
proteins.
203
Overexpression of CED-1::GFP transgene is correlated with gonad abnormalities
We therefore asked whether there was further evidence of a synthetic interaction between *lim-
7p::ced-1::GFP* and *inx-8(qy78).* First, we found evidence that suggests that *lim-7p::ced-1::GFP* is
damaging to the animals with or without *qy78*. In the strain that expresses *lim-7p::ced-1::GFP* and not
qy78 (strain DG5020)*,* roughly 20% of the animals had profound gonad migration defects in one
gonad arm (Figure 4A, 4C). We also observe such defects in the DG5131 strain that combines
qy78[mKate::inx-8] with the *lim-7p::ced-1::GFP* transgene (Figure 4B, 4/19 or 21% of animals), so we
cannot attribute this defect to a spontaneous mutation arising in a single population in transit. We
have not observed such morphological defects in the original strain bearing *qy78*, nor in any other
strain we have studied. The *lim-7p::ced-1::GFP* transgene seems to cause incompletely penetrant
gonad morphology defects.
215
Whether or not overexpressed CED-1::GFP also disrupts the localization of untagged innexin proteins
or other endogenous sheath membrane proteins as it does the tagged mKate::INX-8, and whether
such disruption explains the gonad migration defects we observe for this allele, we currently cannot
say. In many of these *qy78; lim-7p::ced-1::GFP* coexpressing animals (strain DG5131), the intensity
of CED-1::GFP is notably low (Figure 4D). Lower expression levels of the CED-1::GFP fusion protein,

with or without *qy78* in the background, appear more likely to reveal the distal Sh1 cell (Figure 3A and
Figure 4D). This could either be because the absence of competing bright signal makes it easier to
detect dimly expressing distal Sh1, or because high levels of the transgene product are not tolerated
in the distal Sh1 cell. The overexpression of the functional cell death receptor CED-1, and not just the
overexpressed membrane-localized GFP, could contribute to the defects observed in this strain. We
sometimes observe abnormal sheath membrane protrusions that may result from aberrant
engulfment of distal germ cells by the sheath (Figure 4E).

The discrepancy in apparent Sh1 position when two Sh1 cells express different amounts of CED-
1::GFP and when CED-1::GFP is coexpressed with mKate::INX-8 provides definitive evidence that
CED-1::GFP sometimes fails to label the entire distal sheath (the same phenomenon is reported in
Figure 2 Figure Supplement 2B in the recent study (Tolkin et al., 2021)). Furthermore, the defects
caused in gonads overexpressing this functional cell death receptor suggests its localization to the
distal Sh1 membrane at high levels is not well-tolerated. We therefore conclude that *lim-7p::ced-*
1::GFP is an unacceptable marker of distal Sh1.

Assessing sheath markers for evidence of gonad disruption—Brood size
Just because *lim-7p::ced-1::GFP* is a poor marker of the distal sheath does not, however, relieve
concerns that the endogenously tagged innexins mKate::INX-8 and GFP::INX-9 are altering the
gonad. A control for tagged innexin function was originally carried out (Gordon et al., 2020). Briefly, a
careful genetic analysis (Starich et al., 2014) reported that the single mutant *inx-9(ok1502)* is fertile,
but the *inx-8(tn1474); inx-9(ok1502)* double mutant is sterile. Therefore, attempts to use
CRISPR/Cas9 to introduce a fluorescent tag in the *inx-8* locus were first performed in the *inx-*
9(ok1502) background, and only once a fertile edited strain was recovered was the same edit
introduced into the otherwise wild-type genetic background. We conducted brood size assays for
strains discussed in this study, including the DG5131 strain containing both *lim-7p::ced-1::GFP* and
the tagged innexin *qy78[mKate::inx-8]* that was imaged and analyzed by (Tolkin et al., 2021) but not
assayed for brood size (Table 1).

Table 1. Brood size assays.

Strain name	Full genotype	Live brood[a]	Reduction vs wt %	Unhatched eggs	Embryonic Lethality %
N2	wild type	295 ± 39 (n=57)	NA	NA	NA [b]
KLG019	*qy79[GFP::inx-9];nasi2[c]**	226 ± 22 (n=13)	23%	165±58	41 ± 8%
NK2571	*qy78[mKate::inx-8;cpIs122**	220 ± 41 (n=15)	25%	20±14	9 ± 5%
DG5020	*bcIs39[lim-7p::ced-1::GFP];naIs37**	202 ± 29 (n=12)	32%	62±47	20 ± 14%
DG5131	*qy78[mKate::inx-8];bcIs39[lim-7p::ced-1::GFP];naIs37**	187 ± 45 (n=14)	37%	40±25	18 ± 11%
LP530	*cp243[cam-1::mNG]*	260 ± 31 (n=10)	12%	NA	NA
NK2324	*qy23[ina-1::mNG]*	237 ± 37 (n=8)	20%	NA	NA

[a]Viable offspring that hatch from a single parent
[b]N2 numbers come from multiple trials, not all of which counted negligible numbers of dead embryos,
including the trial in which *ina-1(qy23)* and *cam-1(cp243)* were counted.
[C]*qy79[GFP::inx-9]* allele in strains NK2572 and NK2573 from (Gordon et al., 2020) with germ cell
nuclear marker *naSi2*; this combination of alleles was used in the cross to *glp-1(bn18)* in Figure 1D.

We find reductions in brood size for all of the strains under investigation, including a reduced brood
size and notable embryonic lethality in two strains (DG5020 and DG5131) carrying the *lim-7p::ced-*
1::GFP transgene. Interestingly, despite being genetically redundant genes (Starich et al., 2014)
tagged in highly similar ways, and having similar live brood sizes, our endogenously tagged *inx-*
8(qy78) and inx-9(qy79) strains had dramatically different degrees of embryonic lethality, with *qy79*
producing over 150 unhatched eggs per worm. All of the fluorescently marked strains have mildly to
moderately reduced brood sizes. On the basis of brood size alone, there is not a strong reason to
prefer one of these markers over another.

Assessing sheath markers for evidence of gonad disruption—Proliferative zone
Because brood size is an emergent property of many gonad, germline, embryonic, and systemic
processes (including gonadogenesis, stem cell maintenance, regulation of meiosis, spermatogenesis,
oogenesis, metabolism, ovulation, and embryogenesis), defects in brood size are not a direct proxy
for dysregulation of the germ line proliferative zone. We therefore turned our attention back to the
distal gonad and asked whether the strains with fluorescent sheath markers have abnormalities in
several metrics (Figure 5A). The length of the proliferative zones differ among strains (as measured
by DAPI staining of germ cell nuclei to detect and measure the length of the germ line distal to
crescent shaped nuclei of meiosis I, (Hubbard, 2007)) (Figure 5A, 5C, 5D). The NK2571 strain with
the tagged innexin *inx-8(qy78)* and DTC marker has a normally patterned distal germ line (average
proliferative zone length of 106 µm, or ~26 germ cell diameters) that is indistinguishable from wild
type N2 (average of 109 µm or ~27 germ cell diameters Figure 5C, 5C', and 5D). Excluding worms
with gross morphology defects, the DG5020 strain bearing a DTC marker and *lim-7p::ced-1::GFP* has
a measurably shorter distal germ line (average of 91 µm, or ~23 germ cell diameters, Figure 5C" and
5D). In the DG5131 strain that combines these alleles, the distal germ line is notably shortened
(average of 79 µm or ~20 germ cell diameters, Figure 5C''' and 5D). This is comparable to the defect
caused by the *glp-1(bn18)* allele at the permissive temperature shown in Figure 1E. Abnormal distal
gonad patterning provides further evidence that a synthetic interaction between the *lim-7p::ced-*
1::GFP transgene and the *qy78* allele—not the *qy78* allele alone—is responsible for the shorter
proliferative zone observed for strain DG5131 (Figure 4 from Tolkin et al., 2021).

We also counted the number of mitotic figures made by dividing cells in metaphase and anaphase in
these strains (Figure 5B) and the total length of the gonad from vulva to tip (Figure 5E). Wild type N2
had an average of 4.6 dividing cells per gonad; NK2571 had an average of 5.1 dividing cells per
gonad (these two were not significantly different); DG5020 had an average of 3.4 dividing cells per
gonad; DG5131 had an average of 3.2 dividing cells per gonad (these last two strains were
significantly different from wild type, see Figure 5B and legend). Gonad lengths were not significantly
different between N2 (average length of 670 µm), NK2571 (639 µm) or DG5020 (631 µm), but were
significantly shorter in DG5131 (575 µm).

In the end, we find that only the strain combining *inx-8(qy78)* and *lim-7p::ced-1::GFP* has a
dramatically smaller gonad that differs from the wild-type in three key measures. Expression of the
qy78 allele alone with a DTC marker, on the other hand, does not cause any of these quantitative
gonad phenotypes. The moderate brood size defects shown by all strains could be caused by
numerous processes outside of stem cell regulation. For example, we find the hypothesis of (Tolkin et
al., 2021) based on the findings of (Starich et al., 2020, 2014), that a major role of *inx-8*/*9* is in the
proximal gonad regulating the provisioning of oocytes with essential metabolites, to be compelling.
This hypothesis also has support from the large number of unhatched eggs observed for *inx-*
9(qy79[GFP::inx-9]). Thus, we conclude with the observation that endogenous, fluorescently tagged
sheath membrane proteins consistently mark both of the distal Sh1 cells without measurably
impairing distal gonad function and should be the reagents of choice for live-imaging in this cell type.
They also consistently report a distal Sh1 position adjacent to the stem cell zone, as we previously
found (Gordon et al., 2020).
Discussion

We discovered that the distal position of Sh1 is much closer to the distal end of the young adult
hermaphrodite gonad then than was previously observed, where it forms an interface with the DTC's
proximal projections and overlaps substantially with the proliferative zone of the germline where
mitotic cell divisions occur (Gordon et al., 2020). We have now confirmed this finding with functional
manipulations of germ cell cycling and cell fate. We observed a distal Sh1 position in other strains
with endogenously tagged sheath cell membrane proteins that act in molecular pathways outside of
gap junctional coupling, and in a substantial fraction of traditional transgenic animals expressing *lim-7*
promoter-driven CED-1::GFP, GFP::CAAX, and cytoplasmic GFP (though these strains have high
variability in fluorescence intensity and localization). Therefore, we consider the results presented
here to be confirmatory of the foundational finding of (Gordon et al., 2020), which is that almost all
mitotic germ cells in the adult hermaphrodite contact the DTC or Sh1, with a noteworthy population in
contact with both. Other recent work suggests a role for the sheath cells in promoting adult germ cell
proliferation, specifically through modulation of Notch receptor *glp-1* expression (Gopal et al., 2020).
We focus especially on young adults in these studies (less than 24 hours post mid-L4, see Methods).
An important caveat to the work is that the gonad is dynamic and cell shapes and positions change
over time. Indeed, dynamic processes could lead to the surprising difference in position often seen
between the two Sh1 cells in a single gonad arm, with one Sh1 cell growing more actively over germ
cells as they leave the niche. The high variability of an overexpressed *lim-7p::ced-1::GFP* transgene
has allowed for this surprising discovery, though that variability makes it a poor marker of the
absolute position of the Sh1 cells, it sometimes causes gonad defects, and it interacts synthetically
with *qy78* to cause gonad defects.

In physics, the observer effect states that it is impossible to observe a system without changing it. In
biological imaging in *C. elegans*, this means that we can either observe wild-type animals that are
dead, dissected and/or fixed and coated or stained, or we can observe genetically modified animals
that are alive. Some fine, membranous cellular structures do not survive fixation (Gerdes et al., 2013;
Kornberg and Roy, 2014). On the other hand, any genomic modification runs the risk of altering an
animal's physiology.

We feel most confident examining endogenously tagged gene products in Sh1 for several reasons.
First, proteins expressed at physiological levels are less likely to directly damage a cell vs.
overexpressed fluorescent proteins (Kintaka et al., 2016). Second the ability to cross-reference
among strains with different tagged proteins that act in different molecular pathways allows us to use
concordant results in reconstructing cell positions; any single marker may or may not localize to the
region of interest, but concordant results among independent experiments help construct an accurate
picture of the cell. One factor to consider, however, is that not every endogenously expressed protein
is likely to localize evenly across all regions of a cell. We would expect in a large cell like Sh1 that
interacts with germ cells in many stages of maturation that some cell-surface proteins would be
regionalized. Along those lines, it seems possible that the Sh1 cells might have mechanisms to
exclude the cell death receptor CED-1 from the cell membrane domain that contacts proliferating
germ cells. The *bcIs39* transgene is typically used to study engulfment of apoptotic germ cell corpses
at the bend of the gonad and rescues *ced-1* loss of function mutants for apoptotic germ cell corpse
engulfment (Zhou et al., 2001). We find this marker to be unreliable in the distal region of the cell, and
to cause gonad defects especially but not only when combined with endogenously tagged *inx-*
8(qy78). A recent study (Tolkin et al., 2021) uses this transgene in most of the backgrounds analyzed
(sometimes detecting the CED-1::GFP by anti-GFP antibody staining, which appears to amplify the
variability of the marker), so we find this problematic reagent to undermine that study's conclusions.

The need for caution when observing and interpreting endogenously tagged fluorescent proteins is
noted. Several steps can and should be taken to increase confidence that a tagged protein is not
causing cryptic or unwanted phenotypes. First, multiple edited lines should be recovered and
outcrossed, thereby reducing the likelihood that a phenotype is caused by off-target Cas9 cutting
creating lesions in any individual edited genome. Second, brood size should be estimated either by
timed food depletion (less rigorous) or formal brood size assays (more rigorous). Third, edited lines
should be examined for known phenotypes caused by loss of function of the targeted genes. This can

be done, in order of least to most rigorous, by consulting the literature, by comparing to RNAi
treatments or known mutants, and finally by introducing AID tags and using the degron strategy to
deplete the gene product under the lab's exact experimental conditions of choice (Zhang et al., 2015),
however this step will not work for extracellular tags (because extracellular AID tags are not
accessible to TIR1 ubiquitin ligase). Finally, any "markers" used should be assessed on their own for
the phenotype of interest. Even with these controls in place, synthetic interactions can emerge
between "markers" and alleles, including tagged proteins of interest. These interactions can
themselves reveal biologically relevant phenomena, but only if they are recognized.
374
In the end, no transgenic or genome-edited strain is truly wild type, and it should be our expectation
that such strains might be somewhat sensitized as a result. Indeed, the synthetic interaction we
document between *lim-7p::ced-1::gfp* and *inx-8(qy78)* suggests that the *qy78* is sensitized for gonad
defects caused by other genetic elements affecting the gonadal sheath. However, the perfect reagent
does not exit. We can only look for congruent results among a set of independent reagents with non-
overlapping weaknesses. Finally, we can formulate questions narrowly enough that, despite their
shortcomings, our imperfect reagents are adequate to help answer them. In the future, new
endogenously tagged alleles that are expressed in the sheath, single-copy, membrane-localized
transgenes that do not affect distal gonad patterning, and different imaging modalities like electron
microscopy will shed more light on the complex relationship between the gonadal sheath and the
germ line. At the present time, however, we consider the existence of an interface between the DTC
and Sh1 cells that coincides with the boundary of the distal-most stem-like germ cells to be supported
by the preponderance of evidence.
388

Acknowledgements
We thank T. Tolkin, A. Mohammed, T. Starich, T. Schedl, J.A. Hubbard, and D. Greenstein for
sharing their manuscript and strains DG5020 (combining published alleles *bcIs39* (Zhou et al., 2001)*,*
and *naIs37* (Pekar et al., 2017) and DG5131 (combining published alleles *qy78* (Gordon et al., 2020)*,*
bcIs39 (Zhou et al., 2001)*,* and *naIs37* (Pekar et al., 2017)). We thank D. Greenstein and the CGC for
the temperature sensitive mutant strains and B. Goldstein and A. Pani for LP530. We thank R. Dowen
and P. Breen for anonymized replication experiments. We are grateful for helpful conversations with
D. Sherwood and other colleagues.

397
References

Cinquin O, Crittenden SL, Morgan DE, Kimble J. 2010. Progression from a stem cell-like state to early differentiation in the C. elegans germ line. *Proc Natl Acad Sci U S A* **107**:2048–53. doi:10.1073/pnas.0912704107

Evans TC. 2006. Transformation and Microinjection In: WormBook, editor. The C. Elegans Research Community, WormBook.

Fox PM, Schedl T. 2015. Analysis of Germline Stem Cell Differentiation Following Loss of GLP-1 Notch Activity in Caenorhabditis elegans. *Genetics* **201**:167–84. doi:10.1534/genetics.115.178061

Francis R, Nayack S. n.d. Nayak Lab's DAPI Staining Protocol. https://owd.tcnj.edu/~nayaklab/protocols/DAPIstaining.htm

Gerdes H-H, Rustom A, Wang X. 2013. Tunneling nanotubes, an emerging intercellular communication route in development. *Mech Dev* **130**:381–387. doi:10.1016/J.MOD.2012.11.006

Gopal S, Amran A, Elton A, Ng L, Pocock R. 2020. Notch-Directed Germ Cell Proliferation Is Mediated by Proteoglycan-Dependent Transcription. *bioRxiv* 2020.07.30.229997. doi:10.1101/2020.07.30.229997

Gordon KL, Zussman JW, Li X, Martin CM, Sherwood DR. 2020. Stem cell niche exit in C. elegans via orientation and segregation of daughter cells by a cryptic cell outside the niche. *Elife* **9**:e56383. doi:DOI: 10.7554/eLife.56383

Hall DH, Winfrey VP, Blaeuer G, Hoffman LH, Furuta T, Rose KL, Hobert O, Greenstein D. 1999. Ultrastructural features of the adult hermaphrodite gonad of Caenorhabditis elegans: Relations between the germ line and soma. *Dev Biol* **212**:101–123.

Henderson ST, Gao D, Lambie EJ, Kimble J. 1994. *lag-2* may encode a signaling ligand for the GLP-1 and LIN-12 receptors of *C. elegans*. *Development* **120**:2913–24.

Heppert JK, Pani AM, Roberts AM, Dickinson DJ, Goldstein B. 2018. A CRISPR Tagging-Based Screen Reveals Localized Players in Wnt-Directed Asymmetric Cell Division. *Genetics* **208**:1147–1164. doi:10.1534/GENETICS.117.300487

Hubbard EJA. 2007. Caenorhabditis elegans germ line: a model for stem cell biology. *Dev Dyn* **236**:3343–57. doi:10.1002/dvdy.21335

Killian DJ, Hubbard EJA. 2005. Caenorhabditis elegans germline patterning requires coordinated development of the somatic gonadal sheath and the germ line. *Dev Biol* **279**:322–335. doi:10.1016/j.ydbio.2004.12.021

Kintaka R, Makanae K, Moriya H. 2016. Cellular growth defects triggered by an overload of protein localization processes. *Sci Reports 2016 61* **6**:1–11. doi:10.1038/srep31774

Kornberg TB, Roy S. 2014. Cytonemes as specialized signaling filopodia. *Development* **141**:729–36. doi:10.1242/dev.086223

Lee CH, Shin H, Kimble J. 2019. Dynamics of Notch-Dependent Transcriptional Bursting in Its Native Context. *Dev Cell* **50**:426-435.e4. doi:10.1016/j.devcel.2019.07.001

Oshima A, Tani K, Fujiyoshi Y. 2016. Atomic structure of the innexin-6 gap junction channel determined by cryo-EM. *Nat Commun* **7**:13681. doi:10.1038/ncomms13681

Pekar O, Ow MC, Hui KY, Noyes MB, Hall SE, Jane Albert Hubbard E. 2017. Linking the environment, DAF-7/TGFβ signaling and LAG-2/DSL ligand expression in the germline stem cell niche. *Dev* **144**:2896–2906. doi:10.1242/dev.147660

Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S, Rueden C, Saalfeld S, Schmid B, Tinevez J-Y, White DJ, Hartenstein V, Eliceiri K, Tomancak P, Cardona A. 2012. Fiji: an open-source platform for biological-image analysis. *Nat Methods* **9**:676–682. doi:10.1038/nmeth.2019

Shin H, Haupt KA, Kershner AM, Kroll-Conner P, Wickens M, Kimble J. 2017. SYGL-1 and LST-1 link niche signaling to PUF RNA repression for stem cell maintenance in Caenorhabditis elegans. *PLoS Genet* **13**. doi:10.1371/journal.pgen.1007121

Starich TA, Bai X, Greenstein D. 2020. Gap junctions deliver malonyl-CoA from soma to germline to support embryogenesis in Caenorhabditis elegans. *Elife* **9**. doi:10.7554/eLife.58619

Starich TA, Hall DH, Greenstein D. 2014. Two classes of gap junction channels mediate soma-germline interactions essential for germline proliferation and gametogenesis in Caenorhabditis elegans. *Genetics* **198**:1127–1153. doi:10.1534/genetics.114.168815

Team RDC. 2020. R: A language and environment for statistical computing.

Tolkin T, Mohammed A, Starich T, Schedl T, Hubbard EJA, Greenstein D. 2021. Innexin function
dictates the spatial relationship between distal somatic cells in the Caenorhabditis elegans gonad
without impacting the germline stem cell pool. *bioRxiv* 2021.10.22.465523.
doi:10.1101/2021.10.22.465523

Zhang L, Ward JD, Cheng Z, Dernburg AF. 2015. The auxin-inducible degradation (AID) system
enables versatile conditional protein depletion in C. elegans. *Dev* **142**:4374–4384.
doi:10.1242/dev.129635

Zhou Z, Hartwieg E, Horvitz HR. 2001. CED-1 Is a Transmembrane Receptor that Mediates Cell
Corpse Engulfment in C. elegans. *Cell* **104**:43–56. doi:10.1016/S0092-8674(01)00190-8

463
464 **Methods**

Appendix 1. Key Resources Table				
Reagent type (species) or resource	**Designation**	**Source or reference**	**Identifiers**	**Additional information**
gene (*C. elegans*)	*inx-8*	wormbase.org	Sequence CELE_ZK792. 2	Encodes gap junction hemichannel subunit
gene (*C. elegans*)	*inx-9*	wormbase.org	Sequence CELE_ZK792. 3	Encodes gap junction hemichannel subunit
gene (*C. elegans*)	*glp-1*	wormbase.org	Sequence CELE_F02A9. 6	Encodes Notch receptor
gene (*C. elegans*)	*emb-30*	wormbase.org	Sequence CELE_F54C8. 3	Encodes putative member of APC
gene (*C. elegans*)	*ina-1*	wormbase.org	Sequence CELE_F54G8. 3	Encodes worm alpha integrin ortholog
gene (*C. elegans*)	*cam-1*	wormbase.org	Sequence CELE_C01G6. 8	Encodes worm Wnt receptor
gene (*C. elegans*)	*ced-1*	wormbase.org	Sequence CELE_Y47H9 C.4	Encodes worm cell death receptor
genetic reagent (*C. elegans*)	*inx-8(qy78(mKate: :inx-8)) IV; cpIs122(lag-2p::mNeonGreen:: PLC$^{\delta PH}$)*	Gordon et al., 2020	NK2571	Can be obtained from K. Gordon lab
genetic reagent (*C. elegans*)	*inx-9(qy79(GFP::inx-9)) IV; naSi2(mex-5p::H2B::mCherry::nos-2 3'UTR) II*	*nasi2* transgene from (Roy et al., 2018) doi: 10.1534/g3.1 18.200511; *qy79* from	KLG019	Can be obtained from K. Gordon lab

genetic reagent (*C. elegans*)	*rlmIs5[lim-7p::GFP::CAAX]*	This study	KLG020	Can be obtained from K. Gordon lab
		Gordon et al., 2020		
genetic reagent (*C. elegans*)	*qy78(mKate::inx-8) IV*	This study	KLG021	2x outcross of NK2571 to N2
genetic reagent (*C. elegans*)	*inx-9(qy79(GFP::inx-9)) IV; naSi2(mex-5p::H2B::mCherry::nos-2 3'UTR) II; glp-1(bn18) III*	*glp-1(bn18)* from Kodoyianni et al., 1992 doi: 10.1091/mbc.3.11.1199	KLG022	mutant obtained from CGC, crossed to KLG019
genetic reagent (*C. elegans*)	*inx-8(qy78(mKate::inx-8)) IV; cpIs122(lag-2p::mNeonGreen:: PLCᵟPH); emb-30(tn377) III*	*emb-30(tn377)* from Cinquin et al., (2010) doi: 10.1073/pnas.0912704107	KLG023	mutant obtained from CGC, crossed to NK2571
genetic reagent (*C. elegans*)	*cp243[cam-1::mNG]*	Heppert et al., 2018 doi: doi:10.1534/GENETICS.117.300487	LP530	Can be obtained from B. Goldstein lab
genetic reagent (*C. elegans*)	*qy23[ina-1::mNG]*	Jayadev et al., 2019 doi: 10.1083/jcb.201903124	NK2324	Can be obtained from D. Sherwood lab
genetic reagent (*C. elegans*)	*tnIs6[lim-7p::GFP]*	Hall et al., 1999	DG1575	obtained from CGC
genetic reagent (*C. elegans*)	*bcIs39[lim-7p::ced-1::GFP];naIs37*	Tolkin et al., 2021	DG5020	See Tolkin et al., 2021

genetic reagent (*C. elegans*)	*qy78[mKate::inx-8];bcIs39[lim-7p::ced-1::GFP];naIs37*	Tolkin et al., 2021	DG5131	See Tolkin et al., 2021
Software, algorithm	µManager software v1.4.18	(Edelstein et al., 2010) doi: 10.1002/0471142727.mb1420s92	RRID:SCR_016865	https://micro-manager.org/
Software, algorithm	FIJI 2.0	(Schindelin et al., 2012) doi: 10.1038/nmeth.2019	RRID:SCR_002285	https://fiji.sc/
Software, algorithm	GraphPad Prism version 9.20 (283) for macOS	GraphPad Software, San Diego, California USA	RRID:SCR_002798	www.graphpad.com
Software, algorithm	Adobe Illustrator CC	Adobe Systems Inc.	RRID:SCR_010279	

465
Strains
In strain descriptions, we designate linkage to a promoter with a *p* following the gene name and
designate promoter fusions and in-frame fusions with a double semicolon (::). Some integrated strains
(*xxIs* designation) may still contain for example the *unc-119(ed4)* mutation and/or the *unc-119* rescue
transgene in their genetic background, but these are not listed in the strain description for the sake of
concision, nor are most transgene 3' UTR sequences.
472
Staging of animals for comparisons among sheath markers
We focused on young adult animals around the time egg laying commences, as in (Gordon et al.,
2020). Mid L4 animals are picked from healthy, unstarved populations (which are maintained without
starving for the duration of the experiment). These animals are kept at 20° C for 16-18 hours, until
adulthood is reached and ovulation begins. We prefer not to age the animals much farther into
adulthood for routine imaging (though we did this for the temperature shift experiments to follow
previously published experimental regimes), as once a full row of embryos is present in the uterus,
the distal gonads can become compressed or obscured by embryos. For strains in which a gonad
migration defect is observed (DG5020, DG5131), picking animals in the L4 stage prevents bias for or
against normal-looking adults (as the defects are profound enough to be visible on the dissecting
scope in adults).
484
Temperature-sensitive mutant analysis
Worms from the *emb-30(tn377)* mutant genotype were grown at the permissive temperature (16° C)
for 24h past L4. Plates were shifted to the restrictive temperature (25° C) for 15 h before DAPI

staining, while permissive-temperature controls were maintained at 16° C for 18 h before staining
(because development is proportionally slower at 16°C than at 25° C, permissive-temperature
controls were cultured longer). Two replicates of this experiment were performed with the results
combined in Figure 1E.
492
493 Worms from the *glp-1(bn18)* mutant genotype were grown at the permissive temperature of 16° C for
494 24 h past L4. Plates were shifted to the restrictive temperature (25° C) for 6 h (Fox and Schedl,
495 2015). Permissive-temperature controls were maintained at 16° C for 6 h. Worms were imaged live
496 (see *Confocal imaging*, below).
497
498 *DAPI staining*
499 DAPI staining was by modifying standard protocols (Francis and Nayack, n.d.), with the cold
500 methanol fixation done for a shorter time (2.5 minutes) and the concentration of DAPI higher at 1
501 ug/ml in 0.01% Tween in PBS in the dark for five minutes, washed once with 0.1% Tween in PBS.
502 Samples were briefly stored at 4° C in 75% glycerol and imaged directly in glycerol solution.
503
Confocal imaging
All images were acquired on a Leica DMI8 with an xLIGHT V3 confocal spinning disk head (89 North)
with a 63x Plan-Apochromat (1.4 NA) objective and an ORCA-Fusion Gen-III sCMOS camera
(Hamamatsu Photonics). RFPs were excited with a 555 nm laser, GFPs were excited with a 488 nm
laser, and DAPI was excited with a 405 nm laser. Worms were mounted on agar pads with 0.01M
sodium azide (live) or in 75% glycerol (DAPI stained).
510
Fluorescence intensity of lim-7p::CED-1::GFP and mKate::INX-8
For quantitative comparisons of fluorescence intensity shown in Figure 3 and Figure 4, gonads were
imaged with uniform laser power and exposure times with 1 micron Z-steps. Images were opened in
FIJI (Schindelin et al., 2012) and z-projections were made through the depth of the superficial half of
the gonad (not including signal from the deep Sh1 cell if it was present). Images without any
detectable Sh1 expression were discarded (2/32 images from the analysis in Figure 3A). A line ~20
µm long parallel to long axis of the gonad, terminating near the distal boundary of GFP expression,
and not crossing any gaps in Sh1 revealing background was drawn, and average fluorescence
intensity was measured along its length in arbitrary units.
520
Measurements of DTC and Sh1 positions
The distal tip of the gonad was identified in the fluorescence images if the DTC was marked or in a
DIC image if the DTC was not marked in a given strain. The distance from the gonad tip to the
longest DTC process (when marked), and from the gonad tip to the most distal extent of Sh1 was
measured in FIJI (Schindelin et al., 2012). A DTC-Sh1 interface is detected by subtracting the first
value from the second value—negative numbers reflect the amount of overlap of these cellular
domains across the germ line, positive numbers reflect a gap. This is a conservative estimate, as a
gap of less than one germ cell diameter (~5 µm) would still allow germ cells to contact both the DTC
and Sh1 at the same time. Min/max settings on the fluorescence images are adjusted to allow the
faintest signal to be detected when measuring.
531
Analysis of mosaic expression
The variability of the *lim-7p::ced-1::gfp* transgene allowed us to distinguish the two Sh1 cells in a pair,
especially when coexpressed with *qy78[mKate::inx-8]*. For this experiment, we imaged animals
through the full thickness of the distal gonad (40 µm instead of our usual 20 µm that captures just the
superficial half of the gonad that can be imaged more clearly). Animals in which two distinct Sh1 cells
had different levels of GFP signal were analyzed further for relative cell position. For DG5131, this
was 6/19 samples. For DG5020, this was 31/53 samples.
539
Brood size assays
DG5020 and DG5131 were shipped overnight on 9/23, passaged off the starved shipment plate onto

fresh NGM+OP50 plates and maintained by passaging unstarved animals for 3 generations before
beginning the brood size assay. For each strain, 10-15 L4 animals were singled onto NGM plates
seeded with OP50 and kept in the same incubator, on the same shelf, at 20° C. The singled animals
were passaged once per day on each of the following 5 days to fresh plates, with all plates
maintained at 20° C. Two days after removing the parent, the plates with larval offspring were moved
to 4° C for 20 minutes to cause worm motion to cease, and all larvae (and unhatched eggs when
noted) were counted on a dissecting scope with a clicker by the same team of worm counters, with
internal controls. Plates with unhatched eggs were examined and recounted one day later to see if
any hatched. Offspring from parent worms that died or burrowed in the process were not counted.
Total sample sizes and results reported in Table 1.
552
Distal germ line patterning and mitotic figures
Measurements were made in FIJI from the distal end of the gonad to the transition zone, which is the
distal-most row of germ cells with more than one crescent-shaped nucleus. Mitotic figures were
counted manually as metaphase or anaphase DAPI bodies. Observations of 0 mitotic figures were
counted in the analysis. For Figure 1—Figure Supplement 1E, measurements were made by
manually counting cell diameters. In the distalmost region of the restrictive temperature samples,
germ cell nuclei are abnormal, so absolute distances in microns were divided by the diameter of a
normal-looking germ cell from the distal end to calculate germ cell diameters in this region.
561
Gonad length measurements
Strains were synchronized by bleaching and L1 larvae transferred to OP50 seeded NGM plates. At
16C for 48 h. L4 worms were picked to fresh plates and cultured at 16C for an additional 24 hours.
These Day 1 adult worms were mounted on agar pads with 0.01M sodium azide and imaged live.
Images were analyzed in FIJI using the segmented line tool from vulva to distal gonad tip (usually in 2
tiled images to cover the whole gonad length).
568
569 *Statistical analyses*
570 Tests, test statistics, and p values given for each analysis in the accompanying figure legends. One-
571 way ANOVA followed by Tukey's multiple comparisons test were conducted in R (R Team, 2020) or
572 Prism (GraphPad Prism version 9.20 (283) for macOS, GraphPad Software, San Diego, California
573 USA, www.graphpad.com.
574
575 **Figures and Figure Legends**
576
577 **Table 1. Brood size assays.**

Figure 1. The Sh1 cells associate with proliferative germ cells that are on the path to
differentiation. (A) Schematic of hypothesis for *emb-30(tn377)* experiment. Germ cell (gc) nuclei
shown in magenta, somatic gonad cells shown in green (DTC) and transparent green (Sh1). (B)
Gonads from KLG023 *emb-30(tn377);qy78;cpIs122* worms reared at permissive (left column) and
restrictive (right column) temperatures. Top, merged image. Middle, mKate::INX-8 labeling Sh1 (edge
outlined with yellow dashed line). Bottom, DAPI staining labeling all nuclei with pink arrow marking gc
transition and same yellow dashed line as in middle image showing Sh1 edge. (C) Schematic of
hypothesis for *glp-1(ts)* experiment. (D) Gonads from KLG022 *glp-1(bn18);qy79;naSi2* worms reared
at permissive (left column) and restrictive (right column) temperatures. Top, merged image. Middle,
GFP::INX-9 labeling DTC (outlined in white) and Sh1 (edge outlined with yellow dashed line). Bottom,
germ cell histone mCherry (*naSi2[mex-5p::H2B::mCherry]*) with pink arrow showing gc transition and
same yellow dashed line as in middle image showing Sh1 edge. Note that the *glp-1(bn18)* allele is not
fully wild type at permissive temperatures and is known to have a shortened proliferative zone (Fox
and Schedl 2015). (E) Box plots overlaid with all datapoints measuring the distal position of Sh1 and
the position of the transition zone in germ cell nuclear morphology. Permissive temperature shown in
blue; restrictive temperature shown in red. Permissive *emb-30* N=30; restrictive *emb-30* N=34.
Permissive *glp-1* N=18; restrictive *glp-1* N=21. A one-way ANOVA to assess the effect of temperature
on proximodistal position of gonad features was performed, and was significant for *emb-30*:
$F_{3,124}=134.5$, p<0.0001. Tukey's multiple comparison test found that the mean values of the positions
of Sh1 and the germ cell transition zone were significantly different at the permissive temperature
(mean difference of -45.25 µm, 95% CI -52.38 µm to -38.12 µm, p<0.0001), but not at the restrictive
temperature (mean difference of -2.30 µm, 95% CI -9.00 µm to 4.40 µm, p=0.808). The position of the
germ cell transition zone differed at the permissive vs. restrictive temperatures (mean difference of
42.87 µm, 95% CI 35.95 µm to 49.79 µm, p<0.0001), but the Sh1 position did not (mean difference of
-0.078 µm, 95% CI -7.00 µm to 6.84 µm, p>0.9999). This pattern is observed across replicates and
various controls (Figure 1—Figure Supplement 1). Similar results were obtained for *glp-1*:
$F_{3,74}=52.84$, p<0.0001. Tukey's multiple comparison test found that the mean values of the positions
of Sh1 and the germ cell transition zone were significantly different at the permissive temperature
(mean difference of -35.51 µm, 95% CI -44.59 µm to -26.43 µm p<0.0001) but not at the restrictive
temperature (mean difference of 2.514 µm, 95% CI -5.892 µm to 10.92 µm, p=0.861). The position of
the germ cell transition zone differed at permissive vs. restrictive temperatures (mean difference of
36.02 µm, 95% CI 27.27 µm to 44.77 µm, p<0.0001), but the Sh1 position did not (mean difference of
-1.997 µm, 95% CI -10.75 µm to 6.753 µm, p=0.9318). All scale bars 10 µm.

Figure 2. Sheath-expressed fluorescent proteins show consistency among endogenously
tagged membrane proteins and greater variability in overexpressed transgenes. (A) NK2571
qy78[mKate::inx-8]; cpIs122[lag-2p::mNeonGreen:: PLC$^{\delta PH}$] N=21 (B) KLG019 *qy79[GFP::inx9];naSi2*
(channel not shown) N=16 (C) NK 2324 *qy23[ina-1::mNG]* N=26 (D) LP530 *cp243[cam-1::mNG]*
N=21 (E) Box plots of Sh1 positions for all strains listed above and below, with fluorescent protein
listed on the graph, including transgenes (F) DG1575 *tnIs6[lim-7p::GFP]* N=20 (G) Strain DG5020
bcIs39[lim-7p::CED-1::GFP]; naIs37[lag-2p::mCherry-PH] N=52 (note that mean and range agree
with those reported in (Tolkin et al., 2021)) (H) KLG020 *rlmIs5[lim-7p::GFP::CAAX];cpIs122* N=21.
Purple gradient marks approximate extent of stem cell zone (Lee et al., 2019; Shin et al., 2017).See
Figure 2—Figure Supplement 1 for images of minimum and maximum observed distances for all
markers. Figure 2—Figure Supplement 2 shows comparisons across development of NK2571 and
DG5020. All scale bars 10 μm.

**Figure 3. *lim-7p::*CED-1::GFP has variable expression intensity that conceals distal position of
Sh1.** (A) Plot of distal position vs. fluorescence intensity in arbitrary units of CED-1::GFP at the distal
limit of its domain in N=30 DG5020 *bcIs39[lim-7p::CED-1::GFP]; naIs37[lag-2p::mCherry-PH]*
animals. Dashed black line: all of the lowly-expressing gonads (under ~400 A.U., or <30% maximum
brightness of brightest sample) have a DTC-Sh1 interface detected. (B) DG5020 sample in which
disparate expression levels in the two Sh1 cells of a single gonad arm obscure detection of the DTC-
Sh1 interface. The GFP channel is scaled automatically in B; B' is scaled to saturate the brightest
pixels and reveal the dim second Sh1 cell. Dashed yellow link marks the edge of the bright Sh1 cell.
(C) Schematic showing Sh1 pair configuration over distal germ line, with the distal extent of Sh1p
uncertain in superficial projection. The two Sh1 cells of a pair descend from the anterior and posterior
daughters of Z1 and Z4, so the two Sh1 cells are here labeled Sh1a and Sh1p (arbitrarily). Top,
superficial view. Bottom, side view. (D) DG5131 *qy78[mKate::inx-8]; bcIs39[lim-7p::CED-1::GFP];
naIs37[lag-2p::mCherry-PH]* sample in which one Sh1 cell contacts the DTC around the
circumference of the germ line and the other Sh1 cell lies at some distance from the distal end. Gray
boxes and numbers mark planes and landmarks shown in (E). (E) Five cross sections through gonad
in (E) made by projecting through two 1 µm re-slices at the positions shown by gray boxes in (D).
Same analysis for DG5020 shown in Figure 3 Supplement 1. (F) Same worm as in (D,E); signal from
endogenously tagged allele *qy78[mKate::inx-8]* more uniformly labels the Sh1 cells, obscuring their
individual shapes. All scale bars 10 µm.

645
Figure 4. *lim-7p::ced-1::GFP* **is correlated with gonad defects.**
(A) Example of gonad morphology defect in DG5020 *bcIs39[lim-7p::CED-1::GFP]; naIs37[lag-2p::mCherry-PH]* strain, in which the gonad failed to turn. Gut outlined in dashed shape; magenta puncta in that domain are autofluorescent gut granules. (B) Example of gonad morphology defect in DG5131 *qy78[mKate::inx-8]; bcIs39[lim-7p::CED-1::GFP]; naIs37[lag-2p::mCherry-PH]* strain, in which gonad turned once and arrested without elongating along the dorsal body wall. Schematics in A and B show wild-type gonad morphology with two turns and a DTC that arrives at the dorsal midbody, left, beside schematics of defective gonad migration shown in micrographs. (C) Relative proportions of phenotypes observed in DG5020 animals (N=72). (D) Boxplot comparing fluorescence intensity for coexpressing strain DG5131 in addition to data shown in Figure 3 for DG5020. Fluorescence intensity of the *lim-7p::ced-1::GFP* transgene in this background is statistically indistinguishable from expression levels of this transgene in an otherwise wild type background in the subset of samples that display a DTC-Sh1 interface, shown here segregated from samples from this strain that show a gap between the DTC and Sh1 cells. DG5131 N=17. DG5020 gap N=13. DG5020 interface N=17. A one-way ANOVA to determine the effect of category (genotype or presence of an interface) and fluorescence intensity was performed and was significant, $F_{2,44}=7.70$, p=0.001). Tukey's multiple comparison test finds that the mean fluorescence intensity of DG5020 gonads with a DTC-Sh1 interface differs from DG5020 gonads with a gap between Sh1 and the distal end (p=0.002) and does not differ from DG5131 worms (p=0.908). (E) Gonad from DG5131 strain with white arrowheads indicating aberrant engulfment of germ cells in the distal gonad. Closer to the distal end, a large mass of germ cells showing substantial localization of the CED-1::GFP protein may also reflect ectopic engulfment. Scale bars in A and B, 10 µm; scale bar in E, 25 µm.

Figure 5. A synthetic interaction between *lim-7p::ced-1::GFP* and the tagged innexin *qy78*
shortens the proliferative zone. (A) Illustration of measurements made for Figure 5. (B) Number of
mitotic figures observed in DAPI stained animals of the four strains. Numbers of dividing cells and
gonads examined are as follows: wild type N2 (N = 311 dividing cells/67 gonads, the NK2571 strain
with the tagged innexin *qy78* (N = 184 dividing cells/36 gonads), the DG5020 strain with *lim-7p::ced-
1::gfp* (N = 105 dividing cells/31 gonads), the DG5131 strain combining these sheath markers (N= 136
dividing cells/42 gonads). A one-way ANOVA to determine the effect of genotype on number of mitotic
figures was significant $F_{3,\ 172}$ = 7.081, p=0.0002. Tukey's multiple comparison test revealed that NK2571
did not differ from wild type (mean difference -0.47 cells per gonad, 95% CI -1.65 to 0.71, p=0.7291),
DG5020 differed from wild type by 1.26 germ cells per gonad, 95% CI 0.02 to 2.50, p=0.0453), and
DG5131 differed from wild type (mean difference of 1.40 cells per gonad, 95% 0.28 to 2.52, p=0.0075).
(C-C"") DAPI stained distal gonads for measurement of proliferative zone for the four strains. Asterisk
marks tip of gonad, dashed line marks example of lengths measured. (C) Wild type N2 (N=68), (C') the
NK2571 strain with the tagged innexin *qy78* (N=49), (C") the DG5020 strain with *lim-7p::ced-1::gfp*
(N=40), (C"') and the DG5131 strain combining these sheath markers (N=45). Asterisk marks tip of
gonad. (D) Plots of proliferative zone length (left) and whole gonad length (right) for the four strains. A
one-way ANOVA to determine the effect of genotype on length of proliferative zone was significant
$F_{3,198}$=49.15, p<0.0001. Tukey's multiple comparison test revealed that NK2571 did not differ from wild
type (mean difference 2.69 µm, 95% CI -4.283 µm to 9.663 µm, p=0.750), DG5020 differed from wild
type by ~2-5 germ cell diameters (mean difference of 17.94 µm, 95% CI 10.53 µm to 25.36 µm,
p<0.0001), and DG5131 dramatically differed from wild type (mean difference of 30.36 µm, 95% CI
23.21 µm to 37.51 µm, p<0.0001). The proliferative zone length of DG5131 was also significantly
different from both of its parent strains (NK2571 vs. DG5131 mean difference of 27.67 µm, 95% CI
19.99 µm to 35.35 µm, p<0.0001; DG5020 vs. DG5131 mean difference of 12.42 µm, 95% CI 4.331 µm
to 20.50 µm, p=0.0006). (E) Plots of length of entire gonad from tip to vulva. N2 (N=12), NK2571
(N=17), DG5020 (N=19), DG5131 (N=20). A one-way ANOVA to determine the effect of genotype on
gonad length was significant $F_{3,64}$=6.27, p=0.0009. Tukey's multiple comparison test revealed that
NK2571 did not differ from wild type (mean difference 31.16 µm, 95% CI -32.99 µm to 95.32 µm,
p=0.578), DG5020 also did not differ from wild type (mean difference of 39.42 µm, 95% CI -23.32 µm to
102.2 µm, p=0.3546), and DG5131 did differ from wild type (mean difference of 95.15 µm, 95% CI
33.02 µm to 157.3 µm, p=0.0008). All scale bars 10 µm.

**Figure 1—Figure Supplement 1. Robustness of *emb-30* temperature shift experimental results to
timing of control population.** (A) Schematic of hypothesis for *emb-30(tn377)* experiment. Germ cell
(gc) nuclei shown in magenta, somatic gonad cells shown in green (DTC) and transparent green (Sh1).
(B-F) Results from analysis of strain KLG023 *emb-30(tn377);qy78;cpIs122* under different permissive
temperature control culture times. Box plots overlaid with all datapoints measuring the distal position of
Sh1 and the position of the transition zone in germ cell nuclear morphology. Permissive temperature
shown in blue; restrictive temperature shown in red. (B) Controls fixed at starting point, 36 hours after
L4 at permissive temperature as in (Cinquin et al., 2010). Permissive, N=23; restrictive N=23. (C)
Controls fixed after 21 additional hours of culture at the permissive temperature. Permissive N=18;
restrictive N=17. (D) Two replicates in which controls were cultured an additional 18 hours at the
permissive temperature. Replicate 1, permissive N=9; restrictive N=10. Replicate 2, permissive N=21;
restrictive N=24. (E) Same experiment as shown in C, but with distances measured in cell diameters
instead of microns. (F) Table of relevant ANOVA values for the significance indicators shown in B, C,
and D. For (E), a one-way ANOVA to assess the effect of temperature on proximodistal position of
gonad features was performed, and was significant $F_{3,66}=58.44$, p<0.0001. Tukey's multiple comparison
test found that the mean values of the positions of Sh1 and the germ cell transition zone were
significantly different at the permissive temperature (mean difference of -10.78 cell diameters, 95% CI -
13.09 cd to -8.47 cd, p<0.0001), but not at the restrictive temperature (mean difference of -1.24 cd,
95% CI -3.61 cd to 1.14 cd, p=0.52). The position of the germ cell transition zone differed at the
permissive vs. restrictive temperatures (mean difference of 7.73 cd, 95% CI 5.39 cd to 10,08 cd,
p<0.0001), but the Sh1 position did not (mean difference of -1.81 cd, 95% CI -4.15 cd to 0.53 cd,
p>0.19).

723
724
Figure 2—Figure Supplement 1. Endogenously tagged fluorescent proteins in the Sh1
membrane are less variable than overexpressed integrated transgenes. Minimum (left column)
and maximum (right column) measurements of the distance between distal Sh1 and the distal end of
the gonad for (A, A') *qy78[mKate::inx-8],* (B, B') *qy79[GFP::inx9],* (C, C') *qy23[ina-1::mNG],* (D, D')
cp243[cam-1::mNG] (E, E') *tnIs6[lim-7::GFP],* (F, F') *bcIs39[lim-7p::ced-1::GFP],* (G, G') *rlmIs5[lim-*
7p::GFP::CAAX]. Arrowheads in G and G' mark non-sheath cells positive for *lim-7p*::GFP::CAAX
expression in the plane of the gonad and are unavoidably included in Z-projections that capture the
gonadal cell surface. Note especially in G how dim the Sh1 expression is at the distal extent,
resembling what is sometimes seen for CED-1::GFP expression as reported by Figure 2 Figure
Supplement 2B of (Tolkin et al., 2021). In some cases, the selected images are near-minimum or near-
maximum due to imaging artifacts like low illumination or sample movement in the true minimum or
maximum images. All scale bars 10 µm.

Figure 2—Figure Supplement 2. Differences between *qy78(mKate::inx-8)* and *bcIs39(lim-7p::ced-
***1::GFP)* expression in the sheath appear at the L4-young adult transition.** Side by side
developmental comparisons between the two favored marker strains. Left column DG5020 (*bcIs39[lim-*
7p::CED-1::GFP]; naIs37[lag-2p::mCherry-PH]); right column NK2571 (*qy78[mKate::inx-8]; cpIs122[lag-*
2p::mNeonGreen:: PLC$^{\delta PH}$]). Number of worms examined for each strain at each stage given in the
figure annotations. Note that both strains show a close association of the DTC and sheath cells during
larval development; fluorescence signal from of *lim-7p::ced-1::GFP* appears later in development than
mKate::inx-8 signal (top row). During dorsal DTC migration, highly variable gaps often appear between
the DTC and Sh1 in both marker strains (not pictured). Rapid Sh1 growth during this stage has been
observed (Gordon et al., 2020), so we attribute these variable gaps to our taking snapshots of a
dynamic process. By the end of gonad elongation, most Sh1 cells come to rest within a germ cell
diameter of the DTC for both strains. The large gap between the DTC and Sh1 only appears in the *lim-*
7p::ced-1::GFP strain after larval gonad migration is complete. This stage is also when the *lim-7p::ced-*
1::GFP expressing sheath develops two other unique characteristics—holes over the germ cell bodies,
and a frilly distal edge with small, thin distal projections.

Figure 3 Supplement 1. The Sh1 cells of a pair can take two distinct configurations over the distal germ line. (A) Example of a gonad from DG5020 *lim-7p::ced-1::GFP* animal with dramatically different CED-1::GFP signals revealing the shapes of the two Sh1 cells of the pair. Gray boxes show planes depicted in (B). (B) Three cross sections through gonad in (A) made by maximum projection through two 1 μm re-slices (FIJI) at the positions shown by gray boxes in (A). Dashed yellow and white lines mark the two Sh1 cells. Depending on the proximodistal position of the gonad, one or the other Sh1 cell may surround more of the germ line. (C) Example of another gonad from DG5020. (D) Three cross sections through gonad in (C) made by projecting through two 1 μm re-slices at the positions shown by gray boxes in (C). (E,F) Gonads from strain DG5131 with merged images on top, CED-1::GFP channel in the middle, and mKate::INX-8 and distal tip marker channel on the bottom. Yellow outlines show regions of interest in which fluorescence intensity was measured. bg = background, subtracted from fluorescence intensity measured in each of the two Sh1 cells, which express CED-1::GFP at disparate levels. (E' and F') Insets from E and F. In both cases, mKate::INX-8 signal is half as strong in the Sh1 cells with more CED-1::GFP. Note also in E and F that mKate::INX-8 and bright CED-1::GFP mark a different distal extent of Sh1. All scale bars 10 μm.

Figure 1

Figure 2

Figure 3

A

Fluorescence intensity at distal limit of GFP in arbitrary units (y-axis, 0–1600)

Distal edge of CED-1::GFP signal from distal end in microns, strain DG5020 (x-axis, 0–80)

Legend:
- Shows bare region (green diamond)
- Shows DTC-Sh1 interface (purple circle)

B

DG5020 *bcls39;naIs37*
CED-1::GFP auto scaled

B'

DG5020 *bcls39;naIs37*
CED-1::GFP, saturate brightest

C

Sh1p
Sh1a
Sh1p

Sh1a
?
Sh1p

D

DG5131

1
2
3

E

1
2
3

F

DG5131 *qy78;bcls39;naIs37*

Figure 4

A DG5020 *bcIs39;naIs37*

B DG5131 *qy78;bcIs39;naIs37*

C

Gonad morphology defects 22%

DTC-Sh1 interface 42%

Gap 36%

D

Fluorescence intenisty at distal limit of GFP in arbitrary units

DG5131   DG5020 gap   DG5020 interface

E DG5131 *qy78;bcIs39;naIs37*

Figure 5

F   ANOVA table for comparisons shown in Figure 1--Figure Supplement 1

Replicate	$F_{DFn,\ DFd}$	P Value	Comparison for Tukey's	Mean Diff. μm	95.00% CI of diff.	Adjusted P Value
Starting point control (B)	$F_{3,\ 88}$ = 184.9	P<0.0001	--	--	--	--
	--	--	$Sh1_{permissive}$ to $gc_{permissive}$	-50.03	-56.31 to -43.76	<0.0001
	--	--	$Sh1_{restrictive}$ to $gc_{restrictive}$	-0.5304	-6.808 to 5.747	0.9961
	--	--	$gc_{permissive}$ to $gc_{restrictive}$	43.23	36.96 to 49.51	<0.0001
	--	--	$Sh1_{permissive}$ to $Sh1_{restrictive}$	-6.271	-12.55 to 0.005802	0.0503
18 hour control (Fig.1)	$F_{3,124}$=134.5	P<0.0001	--	--	--	--
	--	--	$Sh1_{permissive}$ to $gc_{permissive}$	-45.25	-52.38 to -38.12	<0.0001
	--	--	$Sh1_{restrictive}$ to $gc_{restrictive}$	-2.3	-9.00 to 4.40	0.808
	--	--	$gc_{permissive}$ to $gc_{restrictive}$	42.87	35.95 to 49.79	<0.0001
	--	--	$Sh1_{permissive}$ to $Sh1_{restrictive}$	-0.078	-7.00 to 6.84	0.9999
21 hour control (C)	$F_{3,\ 66}$ = 87.92	P<0.0001	--	--	--	--
	--	--	$Sh1_{permissive}$ to $gc_{permissive}$	-41.1	-48.94 to -33.25	<0.0001
	--	--	$Sh1_{restrictive}$ to $gc_{restrictive}$	-3.92	-11.99 to 4.152	0.5788
	--	--	$gc_{permissive}$ to $gc_{restrictive}$	36.63	28.67 to 44.59	<0.0001
	--	--	$Sh1_{permissive}$ to $Sh1_{restrictive}$	-0.545	-8.504 to 7.414	0.9979
18 hour, rep 1 vs. 2 (D)	--	--	--	--	--	--
Permissive	--	--	$Sh1_{replicate\ 1}$ to $Sh1_{replicate\ 2}$	7.332	-5.429 to 20.09	0.6396
Permissive	--	--	$gc_{replicate\ 1}$ to $gc_{replicate\ 2}$	4.685	-8.076 to 17.45	0.9483
Restrictive	--	--	$Sh1_{replicate\ 1}$ to $Sh1_{replicate\ 2}$	3.017	-9.038 to 15.07	0.9942
Restrictive	--	--	$gc_{replicate\ 1}$ to $gc_{replicate\ 2}$	-8.233	-20.29 to 3.823	0.4166

Minimum Sh1 distance from tip   Maximum Sh1 distance from tip   Figure 2--Figure Supplement 1

A   *inx-8(qy78)*

A'   *inx-8(qy78)*

B   *inx-9(qy79)*

B'   *inx-9(qy79)*

C   *ina-1(qy23)*

C'   *ina-1(qy23)*

D   *cam-1(cp243)*

D'   *cam-1(cp243)*

E   *tnIs6(lim-7p::GFP)*

E'   *tnIs6(lim-7p::GFP)*

F   *bcIs39(lim-7p::ced-1::GFP)*

F'   *bcIs39(lim-7p::ced-1::GFP)*

G   *lim-7p::GFP::CAAX*

G'   *lim-7p::GFP::CAAX*

DG5020 *(bcls39[lim-7p::CED-1::GFP];*
naIs37[lag-2p::mCherry-PH])

NK2571 *(qy78[mKate::inx-8];*
cpIs122[lag-2p::mNeonGreen:: PLC^δPH])

ventral migration — 5/5 worms — 5/6 worms

first turn — 4/5 worms; fifth lacks GFP — 5/5 worms

early dorsal migration — 3/4 worms; fourth lacks GFP — 5/5 worms

end of DTC migration — 6/8 worms; 2 have gap — 9/9 worms

beginning of DTC elaboration — 7/8 worms; 1 has interface — 8/8 worms

A

B

C

D

E

bg

CED-1::GFP

mKate::INX-8

F

bg

CED-1::GFP

mKate::INX-8

E'

F'

50% mKate::INX-8
signal reduction in
bright CED-1::GFP
vs. dim CED-1::GFP
Sh1 cell