Fascicles Split or Merge Every ~560 Microns Within the Human Cervical Vagus Nerve

Aniruddha R. Upadhye[1, 2], Chaitanya Kolluru[1], Lindsey Druschel[1, 2], Luna Al Lababidi[1], Sami S. Ahmad[1], Dhariyat M. Menendez[1,2], Ozge N. Buyukcelik[1], Megan L. Settell[3], Michael W. Jenkins[1], David L. Wilson[1], Jing Zhang[4], Curtis Tatsuoka[4,5], Warren M. Grill[6,7,8,9], Nicole A. Pelot[6], Kip A. Ludwig[3,10], Kenneth J. Gustafson[1,5], Andrew J. Shoffstall[1,2,*]

[1] Department of Biomedical Engineering, Case Western Reserve University, Cleveland, OH, United States of America   [2] APT Center, Louis Stokes Cleveland VA Medical Center, Cleveland, OH, United States of America   [3] Department of Biomedical Engineering, University of Wisconsin-Madison, Madison, WI, United States of America   [4] Department of Population and Quantitative Health Sciences, Case Western Reserve University, Cleveland, OH, United States of America   [5] FES Center, Louis Stokes Cleveland VA Medical Center, Cleveland, OH, United States of America   [6] Department of Biomedical Engineering, Duke University, Durham, NC, United States of America   [7] Department of Electrical and Computer Engineering, Duke University, Durham, NC, United States of America   [8] Department of Neurobiology, Duke University, Durham, NC, United States of America   [9] Department of Neurosurgery, Duke University, Durham, NC, United States of America   [10] Department of Neurosurgery, University of Wisconsin-Madison, Madison, WI, United States of America

* Corresponding Author
Andrew Shoffstall
Assistant Professor, Dept. of Biomedical Engineering
Case Western Reserve University
Cleveland, OH 44106
Ajs215@case.edu
+1 216-368-1213

1 Abstract

Vagus nerve stimulation (VNS) is FDA approved for stroke rehabilitation, epilepsy and depression; however, the underlying vagus functional anatomy underlying the implant is poorly understood. We used microCT to quantify fascicular structure and neuroanatomy within human cervical vagus nerves. Fascicles split or merged every ~560 μm (17.8 ± 6.1 events/cm). The high degree of splitting and merging of fascicles in humans may explain the clinical heterogeneity in patient responses.

2 Main

Electrical stimulation of the cervical vagus nerve (cVN) using implanted electrodes, more commonly known as cervical vagus nerve stimulation (cVNS), is an existing clinical therapy with an estimated market size of over $500 million dollars in 2020, and projected to grow exponentially at a compound annual growth rate of 11.4% to a market size of nearly 1.2 billion dollars by 2026.[1] Implanted vagus nerve stimulators are currently approved by the Food and Drug Administration (FDA) to treat epilepsy, depression, obesity and for stroke rehabilitation[2-5], and are in clinical trials to treat diverse conditions including heart failure, diabetes, and rheumatoid arthritis.[6-8]

The vagus nerve at the cervical/neck level is an attractive target for neuromodulation therapies as it is easily identifiable under ultrasound and can be instrumented with a well-established and relatively simple surgical procedure.[9] In humans, the cervical vagus consists of over 100,000 fibers; these include efferent fibers originating from the brainstem and innervating multiple visceral organs, including the lungs, heart, diaphragm, liver, and intestines, and their sensory fibers returning to the brainstem, which ultimately influence noradrenergic, serotonergic, and cholinergic inputs to the cortex.[9-11] As such, intervening at the cervical vagus

presents the opportunity to modify function both within the brain and the majority of organs

within the viscera.[12-21]

Several recent studies in animal models have suggested that smaller, multi-contact

electrodes may more selectively stimulate specific portions of the cervical vagus to take

advantage of underlying functional organization to better isolate intended activation of

therapeutic fibers from unwanted activation of off-target fibers.[22,23] The activation of low-

threshold large-diameter motor efferent fibers of the vagus that innervate the deep muscle of the

necks putatively drives the most common side effects, causing cough, throat pain, voice

alteration, and dyspnea reported in up to 66% of patients.[24-29] In a study of human patients

implanted to treat heart failure, desired heart rate responses were achieved in only 13 of 106

measurements taken at the 6- and 12-month end points, with stimulation thresholds

predominantly limited by side effects attributable to concurrent activation of the neck muscles.[24]

The vagus nerve is known to have distinct functional organization at specific points along its

path connecting the brainstem to the visceral organs.[30,31] Motor efferents responsible for deep

neck muscle activation originate within the nucleus ambiguus in the medulla oblongata and

eventually coalesce into the pharyngeal, superior laryngeal, and recurrent laryngeal branch,

which innervate the pharyngeal, cricothyroid muscle, and cricoarytenoid muscles of the deep

neck, respectively. Parasympathetic efferents originate from the dorsal motor nucleus of the

vagus within the medullar oblongata and travel down the cervical vagus and eventually join

vagal branches leading to and from the visceral organs. In contrast, sensory afferents leading

from the visceral organs follow these same branches back to the main trunk that eventually

becomes the cervical vagus.

Seminal studies by Sunderland have previously demonstrated that although the fascicles of

major peripheral nerves divide and unite to form fascicular plexuses, there is substantial

uniformity of fascicular arrangement of major nerves in the extremities.[32,33] For example, the

palmar cutaneous and motor branches of the median nerve can be dissected proximally for

several centimeters without significant cross branching.[32,33] It is unknown, however if the

human vagus at the cervical level has well maintained functional organization that may account

for the high degree of heterogeneous results across patients clinically. Prior studies in human

cadavers have focused on sparse sampling of the cervical vagus and subsequent 2D

sectioning, which has yielded highly variable results with respect to number of fascicles from

study to study with little information about the underlying functional somatotopy relevant to

VNS.[34-37]

In this study, we collected 8 mid-cervical VNs from 5 human cadavers; each nerve was 5 cm

long, and we focused our quantitative analyses on the middle 1 cm where the clinical VNS cuff

would be surgically placed.[38] We stained the nerves with osmium tetroxide, and we imaged the

nerves' morphology in three dimensions using microCT. We visualized and quantified the

merging and splitting of fascicles along the length of the vagus nerve (**Figures 1, 2**). Merging

and splitting events were detected manually by an impartial observer (**Figure 1 A, C**), noting

delineation by perineurium boundaries (**Figure 1 B**). We measured the distance over which the

events occurred; merges spanned 430 ± 117 (µm ± SD, n = 70) and splits spanned 461 ± 108 (n

= 72) (**Figure 1 D**).

Over the middle 1 cm of all 8 nerves, there were 17.8 ± 6.1 merging and splitting events

(**Figure 2 B, C**), meaning that on average, each fascicle split or merged every ~560 µm. This

number of events is much larger than expected from prior studies using histological

techniques.[34,35,37] For the standard clinical VNS cuff electrode (LivaNova, London, UK) and our

prototypical nerve (with an mean of ~6.6 fascicles), one would expect to observe ~14.2 split or

merge events over the 8 mm between the centers of the bipolar contact pair. These rapid shifts

in fascicular organization would be challenging to observe using standard histological or TEM

methods—typically using a single transverse cross section per nerve—and thus, prior studies

on vagal morphology have not observed this phenomenon.[35,37]

Merging and splitting events increased proportionally with the number of fascicles: more

fascicles provided more opportunity for split/merge events (**Figure 2 A**, β = 1.76, *p* = 0.032). We

used a two-level linear mixed model considering subject and spatial correlation between

samples to evaluate for association. This degree of fascicular reorganization has substantial

implications for VNS due to changing perineurium boundaries, which dramatically influences the

distribution of the electric field.[39] The locations of fibers—and therefore proximity of fibers to the

electrode contacts—also directly influences activation thresholds. Fascicles of a wide range of

diameters participated in splitting and merging events; reorganization was not limited to a sub-

population of small or large diameter fascicles (**Figure 2 D, E**).

Additionally, the cross-sectional areas of parent ("ab") and summed children ("a" + "b")

fascicles before and after merging or splitting events (**Figure 2 F, G)** were calculated and

compared (i.e., "ab" vs "a + b"). The parent areas were consistently larger than the sum of the

children areas (β = 0.87, *p* <0.001 and β = 1.14, *p* <0.001, for splitting and merging,

respectively, where β refers to the slope of the mixed model).

Using the microCT images, we generated a 3D model (**Figure 3 A**) and quantified the

fascicular morphology: number of fascicles, effective circular diameter, and cross-sectional area

(**Figure 3 B-G**). Overall, there was a net increase in mean fascicle diameter (*p*=0.0139) in the

rostral to caudal direction **(Figure 3 B, C)** with negligible change in overall fascicular area

(*p*=0.8399, **Figure 3 F, G**), suggesting a consolidation of the fascicles toward the distal end of

the cervical region, near the clavicle. We did not observe any major branches, although

branches may occur in this region in some individuals.[34] While there was a trend toward a

concomitant decrease in fascicle count with longitudinal distance (**Figure 3 D, E)**, the result was

statistically insignificant using a mixed-effects regression model (*p*=0.1672), likely owing to low

 sample number and the substantial variation between subjects for all three morphological

 metrics.

Figure 1: Representative example of splitting and merging of fascicles along the rostral-to-caudal direction within a 1.1 mm length of the human cVN (Specimen "2R") imaged with microCT. A) The initiation of merging "M" and splitting "S" events are annotated with arrows: 4 merges (M1-M4) and 1 split (S1). Frames are read from left-to-right, top-to-bottom, as if reading text. Frame-to-frame spacing is 100 μm (12 frames = 1.1 mm total longitudinal span). Transverse-plane scale bar shown in bottom right of the figure is 500 μm. B) Example merging event "M2", spanning 6 frames (500 μm). C) A representative line graph depicting event frequency (Split-positive, Merge- negative) along the middle 1 cm length of nerve. D) Table of mean distances

Figure 2: Graphical representation of fascicular dynamics within the central 1 cm of the surgical
window for VNS implantation across 8 nerves. *The quantification of these events was possible*
due to the high resolution along the longitudinal axis of the microCT dataset. A) Correlation
between the number of fascicles and the number of split/merge events along the 1 cm length of
nerve: subject number (color-coded, 2 – 6), left (square), right (circle). B) Box plot showing the
distribution of the number of split/merge events across all samples. C) 3D visualization of a
representative 1 cm window within the cVN (Specimen "4R"). D, E) Box plot showing the

distribution of the diameters of parent fascicles and children (a, b or a+b, respectively) for all merge and split events. F) Association plot of splitting fascicular summed areas of the children (a+b, y-axis) with the areas of the parent (ab, x-axis), mixed model slope β=0.87, p<0.001. G) Association plot of merging fascicular areas of the parent (ab, y-axis) with the summed areas of the children (a+b, x-axis), mixed model slope β=1.14, p<0.001. Note that summed areas of the children are consistently less than the area of the parent fascicle.

***Figure* 3: *Fascicle morphometry assessment within the central 5 cm of the human cVN (Specimen "4R").** A) Representative 3D visualization of segmented microCT images. B, D, E) Fascicle count, diameter, and area at 0.5 cm increments along the 5 cm surgical window for each sample, where x = –2.5 cm is the rostral end and x = +2.5 cm is the caudal end. We also averaged the data across the surgical window for each sample (C, E, G). Bars represent the mean ± SD across the sampled regions of the surgical window. Black horizontal lines represent the mean ± SD across all nerve samples.*

MicroCT enables unique three-dimensional visualization and quantification of vagal

fascicular morphology over arbitrarily long lengths of nerve, enabling new insights into the

spatial organization of the nerve that are essential for the design and analysis of effective and

selective electrical stimulation therapies to treat diseases. MicroCT has been used extensively

in orthopedic studies and other fields, but the imaging technique has only recently been applied

to neural tissues. For example, one study reported a protocol for staining rat sciatic and pig

vagus nerves, optimization of computational methods for high resolution three-dimensional

images of nerve fascicles, and development of image analysis techniques to facilitate

segmentation and tracing of the fascicles.[40] The fascicle morphology measurements obtained

from our microCT data was similar to the measurements obtained by other groups.[37] Here, we

demonstrated the unique value of microCT to quantify fascicular splitting and merging of the

human cVN.

Given the magnitude of fascicular reorganization demonstrated by our data, current VNS

cuff designs are not optimized to provide spatial selectivity. The current clinical standard

involves surgical implantation of a cuff electrode that wraps helically around the entire nerve

trunk, with bipolar contacts spanning ~270°, separated by 8 mm center-to-center. For a

representative nerve from our study, this 8 mm span would traverse over a dozen fascicle

splitting and merging events (min = 9.6, max = 22.4 events, from our limited size dataset).

Further, the fascicular reorganization varies substantially between individuals. Given this intra-

and inter-individual morphological heterogeneity of fascicles, these electrode designs are

unlikely to allow selective activation of spatially localized target fibers within the cVN.

Computational modeling of the vagus nerve can be used for engineering design of neural

cuff[41]; the basis for these models requires anatomically accurate features that reflect the

diversity observed across multiple human subjects. Currently, computational modeling of VNS

relies on longitudinal extrusion of segmented histological cross sections or simplified mock

morphologies which do not represent precise fascicle boundaries or longitudinal spatial variation

[42-44 45 46,47] Autonomic stimulation therapies will be advanced by *a priori* personalized surgical

planning, device designs, and device programming for autonomic stimulation therapies informed

by computational models as used in other neural stimulation treatments.[48] However, to make

personalized decisions and improve the accuracy of the computational predictions, better *in vivo*

imaging modalities are needed to visualize and map the fascicular morphology with higher

precision and resolution in both the transverse and the longitudinal planes.[49]

In conclusion, the fascicular anatomy of VN is extremely complex and dynamic and mapping

nerves using microCT is an effective technique to visualize and quantify fascicle reorganization.

We measured a mean of 17.8 split-or-merge events along 1 cm of the cervical vagus nerve (n=8

samples), implying that there would be ~14 events along the bipolar electrode of current clinical

VNS devices. These data on the fascicle dynamics of the human VN provide insights into the

morphological impact of morphology on VNS efficacy and provide the foundation for

computational models with increased accuracy for analysis and design of therapies with

improved selectivity.

3 Methods

3.1 <u>Tissue Acquisition and Dissection</u>

We collected 8 mid-cervical vagus nerve samples from 5 formaldehyde fixed cadavers (3

left nerves, 5 right nerves), secondary to use in medical school cadaver lab training. Since all

the specimens were harvested from de-identified donor sources, and no protected personal

health information collected, a letter of IRB exemption (non-human-subjects determination) was

sought and approved by the Case Western Reserve University Institutional Review Board.

Cadavers were already disarticulated prior to our dissection; we performed gross and fine

dissection with standard tools to isolate the vagus nerve from surrounding tissues. We made a

rostral cut directly beneath the skull (jugular foramen) approximately at the nodose ganglion.

The caudal/distal cut was made at the level of clavicle. The harvested nerves were stored in 4%

formalin solution until ready for staining. The VNS cuff electrode is clinically placed midway

between the clavicle and the mastoid process, and the surgical incision is 3-4 cm long[38]; we

therefore collected 5 cm of length for each nerve, centered around the approximate location of

VNS cuff placement, which we refer to as the "surgical window" throughout the paper.

3.2   <u>Sample Preparation: Osmium Staining & Paraffin Embedding</u>

The vagus nerves were washed three times with 1X phosphate buffered saline (PBS),

letting the sample shake on an orbital shaker for five minutes after each wash. Osmium

tetroxide (1% v/v) was prepared with deionized water, and the nerves were left fully submerged

in this solution for three days. The samples were then dehydrated with 70% and 95% ethanol

with a deionized water solvent. The dehydration included two quick rinses of the samples with

70% ethanol followed by a full wash and 30 minutes on the orbital shaker. This process was

repeated twice with 70% ethanol, then three additional times with 95% ethanol. The nerves

were stored in 70% ethanol for up to one week prior to embedding in paraffin.

The nerve samples were embedded in paraffin, mounted on a 3D printed plastic mold that fit

the nerve. At the base of the mold, there were grooves every 5 mm, and these grooves were

painted with a marking solution doped with barium sulfate to enhance sample navigation under

X-ray.

3.3   <u>MicroCT and Image Sub-Volume Reconstruction</u>

For the imaging studies, we utilized a Quantum GX2 microCT Imaging System (Perkin

Elmer, Waltham, MA, USA). The embedded nerve was placed in a 36 mm bed. Scans were

warmed up as recommended by the manufacturer, and the nerve was scanned and

reconstructed at 36 mm field of view (FOV). The resultant image block was 72 µm in voxel

resolution (isotropic). Each scan spanned 1.8 cm of nerve length, with 0.3 cm overlap (i.e.,

16.67%) between adjacent blocks to serve image reconstruction.

Post-hoc sub-block reconstruction was performed with Rigaku software provided by

Perkin Elmer. Each sub-block reconstruction was a 5.12 x 5.12 x 5.12 mm^3 cube and each

adjacent sub-blocks overlapped by 0.1 mm (20% overlap); the resolution of final reconstruction

was 10 µm voxel size (isotropic). Images were exported as DICOM files for further processing.

After down-sampling frames along the longitudinal axis by 10-fold, blocks were co-registered

and stitched using ImageJ (FIJI, Version 2.1.0/1.53c).[50] The final image dataset consisted of

pairwise stitched, evenly spaced (100 µm inter-frame spacing) TIFF images. 3D visualizations

were generated by Simpleware™ ScanIP software (Synopsys, Mountain View, CA).

3.4   Fascicle Morphometric Analysis

VN samples were analyzed using ImageJ (FIJI, Version 2.1.0/1.53c) to select, outline, and

measure individual fascicles, using the elliptical selection tool. Fascicle boundaries were

manually estimated based on visual inspection. For morphometric analysis, the operators

evaluated fascicle parameters at 0.5 cm intervals along the length of the 5 cm cervical window

for each nerve. While manual outlining potentially introduces subjective differences between

operators, the magnitude of these differences was deemed negligible based on prior inter-

operator analyses. Image scaling was set according to the microCT manufacturer provided

calibration factor: 1 pixel = 10 µm ,1.0-pixel aspect ratio. Area, minimum & maximum gray

intensity values, shape descriptors, mean intensity value, centroid coordinates, and ellipse-fit

measurements (including major and minor axes, and effective diameter – the average of major

and minor axis) were calculated.

3.5   Merging and Splitting Analysis

The splitting and merging analyses were conducted for the central 1 cm of the cervical

vagus nerve, within the 5 cm of the surgical window that we defined in this paper. The frames in

this region were isolated and loaded as an image sequence on ImageJ and analyzed from the

rostral end to caudal end. All split/merge analyses were conducted manually.

3.5.1   Defining an Event

During our analysis, we defined the start and completion of a split or merge event based on

the fascicle boundaries. We characterized an event as a start of a split when a parent fascicle,

coined "ab", appeared to create a bud or partition within the center of the otherwise consistently

shaded fascicle (e.g., **Figure 1B**). The event was marked as complete when parent fascicle "ab"

completely formed independent circular/ellipsoidal independent children fascicles "a" and "b"

with their own perineurium sheath around the fascicles. In most cases, the perineurium sheathe

is well defined and visible within the microCT. In some cases, the perineurium is inferred when

there is separation of two circular/ellipsoidal geometries. Conversely, we characterized an event

as a merge when fascicle "a" merged into another fascicle "b", resulting in a combined fascicle

"ab", applying the same logic as described above. When multiple events occurred

simultaneously (e.g., one fascicle splitting into three fascicles), we considered it as two different

splitting events. We did not observe any event where three fascicles merged to become one

fascicle in the exact same frame.

3.5.2   Measurements and Analysis

To measure the distance over which the event was taking place, the starting and the ending

frames were recorded. With the total number of frames, we calculate the distance over which

the event takes place. Using ImageJ, the fascicles were measured at the starting and the

ending frames (as mentioned in the morphometric analysis section).

We recorded the number of splitting and merging events across the central 1 cm of each

sample and calculated the average number of events across n = 8 samples. We counted the

number of fascicles in the first, middle, and last frames of the 1 cm window and calculated the

mean fascicle count in the sample. We then determined the number of events/fascicle/cm using

the values calculated as mentioned previously.

3.6   Statistics

Our primary quantitative metric was focused on fascicle splitting and merging events across

our human cadaver nerve specimens (n = 8). Descriptive statistics presented in the text include

mean and standard deviations unless otherwise denoted. Box plots presented in **Figure 2**

display individual data points (colored according to the associated legends), median values

(horizontal center line), mean values (black box), interquartile range (upper and lower box

edge), and outliers (whiskers). Bar plots presented in **Figure 3** display mean values (bar height)

and standard deviation (error bars), with horizontal lines in the background representing the

whole sample mean and standard deviations.

For all statistical tests described below, Two-sided Type I error = 0.05 was adopted.

Analysis was performed using R version 4.0.2.

Specifically, we were interested in evaluating the relationship between the number of

fascicles contained within nerve specimens and the number of splitting or merging events

observed (**Figure 2 A**). The association between the average number of fascicles at the surgical

window and the number of events along the window was investigated with a two-level linear

mixed model with subject and (left or right) side-level random intercepts.

We were also interested in evaluating the conservation of fascicular area before-and-after

splitting and merging events (**Figure 2 F, G**). In order to study the association between

fascicular area of the parent (ab) and summed areas of the children (a+b), we adopted a three-

level hierarchical linear mixed model with subject-level and side-level random intercept with

exponential spatial correlation structure for same side windows.

Similar 3-level models, as described above, were respectively used to explore the spatial

trend of outcomes along the surgical window (rostral-to-caudal) for fascicular area, mean

diameter, and fascicle count (results shared in text).

3.7   <u>**Methodological Limitations**</u>

As with standard histological processes, the staining and fixative reagents can cause

dehydration and shrinkage to tissues. Per prior publications, we anticipate shrinkage could

contribute up to 20% reduction in apparent diameters. However, we did not directly estimate this

in our study, and therefore did not apply any correction factors in our dataset. Further, we

sampled nerves from 5 cadavers, but due to the source of cadaver donation, we were unable to

acquire any demographics. This study can be expanded in the future to greater population

sample size to estimate population variability drive by demographic differences.

4   Acknowledgements

325   This work has been supported by NIH SPARC Program 1OT2OD025340, US Dept. of

326   Veterans Affairs 1IS1BX004384, the Cleveland VA APT Center, and Case Western Reserve

327   University. The opinions expressed in this article are the author's own and do not reflect the view

328   of the National Institutes of Health, the Department of Health and Human Services, or the United

329   States government.

5   Competing Interests Statement

331   KAL is a scientific board members and have stock interests in NeuroOne Medical Inc., a

332   company developing next generation epilepsy monitoring devices. KAL is also paid member of

333   the scientific advisory board of Cala Health, Blackfynn, Abbott and Battelle. KAL also is a paid

6 References

Insights, F. B. *Vagus Nerve Stimulation Market Size, Trends | Report, 2026*, <https://www.fortunebusinessinsights.com/industry-reports/vagus-nerve-stimulation-vns-market-101184> (2019).

Administration, U. S. F. D. *FDA 2021 PMA approval P2100007 for Vivistim System*, <http://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfpma/pma.cfm?id=P210007> (2021).

Administration, U. S. F. D. *FDA 2015 PMA approval P130019 for Maestro Rechargeable System*, <http://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfpma/pma.cfm?id=P130019> (2015).

Administration, U. S. F. D. *FDA 2005 PMA approval P970003 for VNS Therapy System*, <http://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfpma/pma.cfm?id=P970003S050> (2005).

Administration, U. S. F. D. *FDA 1997 PMA approval P970003 for VNS Therapy System*, <http://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfpma/pma.cfm?id=P970003> (1997).

Drewes. *Treatment of Complications to Diabetic Autonomic Neuropathy With Vagus Nerve Stimulation*, <https://clinicaltrials.gov/ct2/show/NCT04143269> (2021).

Corporation, S. M. *Long Term Extension Study of the Safety and Efficacy of Neurostimulation Using a Vagus Nerve Stimulation Device in Patients With Rheumatoid Arthritis*, <https://clinicaltrials.gov/ct2/show/NCT04862117> (2021).

Corporation, B. S. *Neural Cardiac Therapy for Heart Failure Study*, <https://clinicaltrials.gov/ct2/show/NCT01385176> (2021).

Groves, D. A. & Brown, V. J. Vagal nerve stimulation: a review of its applications and potential mechanisms that mediate its clinical effects. *Neurosci Biobehav Rev* **29**, 493-500, doi:10.1016/j.neubiorev.2005.01.004 (2005).

Pavlov, V. A. & Tracey, K. J. The vagus nerve and the inflammatory reflex--linking immunity and metabolism. *Nat Rev Endocrinol* **8**, 743-754, doi:10.1038/nrendo.2012.189 (2012).

Dorr, A. E. & Debonnel, G. Effect of vagus nerve stimulation on serotonergic and noradrenergic transmission. *J Pharmacol Exp Ther* **318**, 890-898, doi:10.1124/jpet.106.104166 (2006).

Beekwilder, J. P. & Beems, T. Overview of the clinical applications of vagus nerve stimulation. *J Clin Neurophysiol* **27**, 130-138, doi:10.1097/WNP.0b013e3181d64d8a (2010).

Bonaz, B., Sinniger, V. & Pellissier, S. The Vagus Nerve in the Neuro-Immune Axis: Implications in the Pathology of the Gastrointestinal Tract. *Front Immunol* **8**, 1452, doi:10.3389/fimmu.2017.01452 (2017).

Bottomley, J. M., LeReun, C., Diamantopoulos, A., Mitchell, S. & Gaynes, B. N. Vagus nerve stimulation (VNS) therapy in patients with treatment resistant depression: A systematic review and meta-analysis. *Comprehensive Psychiatry* **98**, 152156, doi:https://doi.org/10.1016/j.comppsych.2019.152156 (2020).

Chakravarthy, K., Chaudhry, H., Williams, K. & Christo, P. J. Review of the Uses of Vagal Nerve Stimulation in Chronic Pain Management. *Curr Pain Headache Rep* **19**, 54, doi:10.1007/s11916-015-0528-6 (2015).

16   Kin, I. *et al.* Vagus Nerve Stimulation with Mild Stimulation Intensity Exerts Anti-
Inflammatory and Neuroprotective Effects in Parkinson's Disease Model Rats.
Biomedicines **9**, 789, doi:10.3390/biomedicines9070789 (2021).

17   Koopman, F. A., van Maanen, M. A., Vervoordeldonk, M. J. & Tak, P. P. Balancing the
autonomic nervous system to reduce inflammation in rheumatoid arthritis. *J Intern Med*
282, 64-75, doi:10.1111/joim.12626 (2017).

18   Marangell, L. B. *et al.* A 1-year pilot study of vagus nerve stimulation in treatment-
resistant rapid-cycling bipolar disorder. *J Clin Psychiatry* **69**, 183-189,
doi:10.4088/jcp.v69n0203 (2008).

19   Pruitt, D. T. *et al.* Vagus Nerve Stimulation Delivered with Motor Training Enhances
Recovery of Function after Traumatic Brain Injury. *J Neurotrauma* **33**, 871-879,
doi:10.1089/neu.2015.3972 (2016).

20   Sarr, M. G. *et al.* The EMPOWER study: randomized, prospective, double-blind,
multicenter trial of vagal blockade to induce weight loss in morbid obesity. *Obes Surg* **22**,
1771-1782, doi:10.1007/s11695-012-0751-8 (2012).

21   Yin, J., Ji, F., Gharibani, P. & Chen, J. D. Vagal Nerve Stimulation for Glycemic Control
in a Rodent Model of Type 2 Diabetes. *Obes Surg* **29**, 2869-2877, doi:10.1007/s11695-
019-03901-9 (2019).

22   Fitchett, A., Mastitskaya, S. & Aristovich, K. Selective Neuromodulation of the Vagus
Nerve. *Front Neurosci* **15**, 685872, doi:10.3389/fnins.2021.685872 (2021).

23   Ordelman, S. C., Kornet, L., Cornelussen, R., Buschman, H. P. & Veltink, P. H.
Selectivity for specific cardiovascular effects of vagal nerve stimulation with a multi-
contact electrode cuff. *IEEE Trans Neural Syst Rehabil Eng* **21**, 32-36,
doi:10.1109/TNSRE.2012.2214058 (2013).

24   De Ferrari, G. M. *et al.* Long-term vagal stimulation for heart failure: Eighteen month
results from the NEural Cardiac TherApy foR Heart Failure (NECTAR-HF) trial. *Int J*
Cardiol **244**, 229-234, doi:10.1016/j.ijcard.2017.06.036 (2017).

25   Group, T. V. N. S. S. A randomized controlled trial of chronic vagus nerve stimulation for
treatment of medically intractable seizures. *Neurology* **45**, 224-230,
doi:10.1212/wnl.45.2.224 (1995).

26   Handforth, A. *et al.* Vagus nerve stimulation therapy for partial-onset seizures: a
randomized active-control trial. *Neurology* **51**, 48-55, doi:10.1212/wnl.51.1.48 (1998).

27   Nicolai, E. N. *et al.* Sources of off-target effects of vagus nerve stimulation using the
helical clinical lead in domestic pigs. *J Neural Eng* **17**, 046017, doi:10.1088/1741-
2552/ab9db8 (2020).

28   Morris, G. L., 3rd *et al.* Evidence-based guideline update: vagus nerve stimulation for the
treatment of epilepsy: report of the guideline development subcommittee of the american
academy of neurology. *Epilepsy Curr* **13**, 297-303, doi:10.5698/1535-7597-13.6.297
(2013).

29   Sackeim, H. A. *et al.* Vagus nerve stimulation (VNS) for treatment-resistant depression:
efficacy, side effects, and predictors of outcome. *Neuropsychopharmacology* **25**, 713-
728, doi:10.1016/S0893-133X(01)00271-8 (2001).

30   Câmara, R. & Griessenauer, C. J. in *Nerves and Nerve Injuries*   (eds R. Shane Tubbs *et*
al.) 385-397 (Academic Press, 2015).

31   Ellis, H. CLINICAL ANATOMY: A REVISION AND APPLIED ANATOMY FOR CLINICAL
STUDENTS. *Annals of Surgery* **159**, 31-35 (1964).

32 Sunderland Sydney, S. *Nerve and Nerve Injuries*. Pages 841–854, (Baltimore, Williams
and Wilkins Co., 1968).

33 Lee, S. K. & Wolfe, S. W. Peripheral nerve injury and repair. *J Am Acad Orthop Surg* **8**,
243-252, doi:10.5435/00124635-200007000-00005 (2000).

34 Hammer, N. *et al.* Human vagus nerve branching in the cervical region. *PLoS One* **10**,
e0118006, doi:10.1371/journal.pone.0118006 (2015).

35 Hammer, N. *et al.* Cervical vagus nerve morphometry and vascularity in the context of
nerve stimulation - A cadaveric study. *Sci Rep* **8**, 7997, doi:10.1038/s41598-018-26135-
8 (2018).

36 Kawagishi, K. *et al.* Tyrosine hydroxylase-immunoreactive fibers in the human vagus
nerve. *J Clin Neurosci* **15**, 1023-1026, doi:10.1016/j.jocn.2007.08.032 (2008).

37 Pelot, N. A. *et al.* Quantified Morphology of the Cervical and Subdiaphragmatic Vagus
Nerves of Human, Pig, and Rat. *Frontiers in Neuroscience* **14**,
doi:10.3389/fnins.2020.601479 (2020).

38 Giordano, F., Zicca, A., Barba, C., Guerrini, R. & Genitori, L. Vagus nerve stimulation:
Surgical technique of implantation and revision and related morbidity. *Epilepsia* **58**, 85-
90, doi:https://doi.org/10.1111/epi.13678 (2017).

39 Grinberg, Y., Schiefer, M. A., Tyler, D. J. & Gustafson, K. J. Fascicular perineurium
thickness, size, and position affect model predictions of neural excitation. *IEEE Trans
Neural Syst Rehabil Eng* **16**, 572-581, doi:10.1109/TNSRE.2008.2010348 (2008).

40 Thompson, N. *et al.* MicroCT optimisation for imaging fascicular anatomy in peripheral
nerves. *Journal of neuroscience methods* **338**, 108652-108652,
doi:10.1016/j.jneumeth.2020.108652 (2020).

41 Musselman, E. D., Cariello, J. E., Grill, W. M. & Pelot, N. A. ASCENT (Automated
Simulations to Characterize Electrical Nerve Thresholds): A pipeline for sample-specific
computational modeling of electrical stimulation of peripheral nerves. *PLoS Comput Biol*
17, e1009285, doi:10.1371/journal.pcbi.1009285 (2021).

42 Aristovich, K. *et al.* Model-based geometrical optimisation and in vivo validation of a
spatially selective multielectrode cuff array for vagus nerve neuromodulation. *J Neurosci
Methods* **352**, 109079, doi:10.1016/j.jneumeth.2021.109079 (2021).

43 Dali, M. *et al.* Model based optimal multipolar stimulation without a priori knowledge of
nerve structure: application to vagus nerve stimulation. *Journal of neural engineering* **15**,
046018, doi:10.1088/1741-2552/aabeb9 (2018).

44 Bucksot, J. E. *et al.* Flat electrode contacts for vagus nerve stimulation. *PLoS One* **14**,
e0215191, doi:10.1371/journal.pone.0215191 (2019).

45 Pelot, N. A., Behrend, C. E. & Grill, W. M. Modeling the response of small myelinated
axons in a compound nerve to kilohertz frequency signals. *J Neural Eng* **14**, 046022,
doi:10.1088/1741-2552/aa6a5f (2017).

46 Helmers, S. L. *et al.* Application of a computational model of vagus nerve stimulation.
Acta Neurol Scand **126**, 336-343, doi:10.1111/j.1600-0404.2012.01656.x (2012).

47 Arle, J. E., Carlson, K. W. & Mei, L. Investigation of mechanisms of vagus nerve
stimulation for seizure using finite element modeling. *Epilepsy Research* **126**, 109-118,
doi:https://doi.org/10.1016/j.eplepsyres.2016.07.009 (2016).

48 McIntyre, C. C. & Foutz, T. J. Computational modeling of deep brain stimulation. *Handb
Clin Neurol* **116**, 55-61, doi:10.1016/B978-0-444-53497-2.00005-X (2013).

49 Settell, M. L. *et al.* In vivo visualization of pig vagus nerve 'vagotopy' using ultrasound.
bioRxiv, 2020.2012.2024.424256, doi:10.1101/2020.12.24.424256 (2021).

50   Preibisch, S., Saalfeld, S. & Tomancak, P. Globally optimal stitching of tiled 3D
microscopic image acquisitions. *Bioinformatics* **25**, 1463-1465,
doi:10.1093/bioinformatics/btp184 (2009).
479