Host-associated phages disperse across the extraterrestrial analogue

Antarctica

Running title: Dispersal of Antarctic phage

Janina Rahlff[a, #], Till L.V. Bornemann[a], Anna Lopatina[b],

Konstantin Severinov[b], Alexander J. Probst[a,c]

[a]Group for Aquatic Microbial Ecology, University of Duisburg-Essen, Department of

Chemistry, Environmental Microbiology and Biotechnology (EMB), Universitätsstraße 5,

45141 Essen, Germany

[b]Institutes of Molecular Genetics and Gene Biology of the Russian Academy of Sciences,

Moscow, Russia

[c]Centre of Water and Environmental Research (ZWU), University of Duisburg-Essen,

Universitätsstraße 5, 45141, Essen, Germany

#Corresponding author: Janina Rahlff, Janina.rahlff@uol.de

Present address: Centre for Ecology and Evolution in Microbial Model Systems (EEMiS),

Department of Biology and Environmental Science, Linnaeus University, SE-39182, Kalmar,

Sweden

Key words: viruses, extreme environments, astrovirology, metagenomics, cryosphere,

Ralstonia

Abstract

Extreme Antarctic conditions provide one of the closest analogues of extraterrestrial

environments. Since air and snow samples, especially from polar regions, yield DNA amounts

in the lower picogram range, binning of prokaryotic genomes is challenging and renders

studying the dispersal of biological entities across these environments difficult. Here, we

hypothesized that dispersal of host-associated bacteriophages (adsorbed, replicating or

prophages) across the Antarctic continent can be tracked via their genetic signatures aiding our

understanding of virus and host dispersal across long distances. Phage genome fragments

(PGFs) reconstructed from surface snow metagenomes of three Antarctic stations were assigned

to four host genomes, mainly Betaproteobacteria including *Ralstonia* spp. We reconstructed the

complete genome of a temperate phage with near-complete alignment to a prophage in the

reference genome of *Ralstonia pickettii* 12D. PGFs from different stations were related to each

other at the genus level and matched similar hosts. Metagenomic read mapping and nucleotide

polymorphism analysis revealed a wide dispersal of highly identical PGFs, 13 of which were

detected in seawater from the Western Antarctic Peninsula in distance of 5538 km to the snow

sampling stations. Our results suggest that host-associated phages, especially of *Ralstonia* sp.,

disperse over long distances despite harsh conditions of the Antarctic continent. Given that 14

phages associated with two *R. pickettii* draft genomes isolated from space equipment were

identified, we conclude that *Ralstonia* phages are ideal mobile genetic elements to track

dispersal and contamination in ecosystems relevant for Astrobiology.

Importance

Host-associated phages of the bacterium *Ralstonia* identified in snow samples can be used to

track microbial dispersal over thousands of kilometers across the Antarctic continent, which

functions as an extraterrestrial analogue because of its harsh environmental conditions. Due to

presence of these bacteria carrying genome-integrated prophages on space-related equipment,

and the here demonstrated potential for dispersal of host-associated phages, our work has

implications for Planetary Protection, a discipline in Astrobiology interested in preventing

contamination of celestial bodies with alien biomolecules or forms of life.

Introduction

Due to harsh environmental conditions and isolation by the surrounding Southern Ocean's

Circumpolar Current, Antarctica is considered an analogue for multipurpose space exploration

(1, 2). For example, its McMurdo Dry Valleys are regarded as a close terrestrial analogue to

Mars (3). Astrobiology model organisms found on Antarctica are highly adapted to stressful

conditions and comprise prokaryotes such as spore-forming *Bacilli* (4, 5), but also microfungi

(3, 6). Understanding endurance and dispersal of microorganisms under conditions that mimic

those on extraterrestrial planets, i.e., high UV radiation, low temperature, and low nutrient

availability, has important implications for Planetary Protection. For instance, the dispersal of

microbes that hitchhike to a celestial body is currently not considered in Planetary Protection,

a discipline in Astrobiology set out with the aim of preventing contamination of celestial bodies

with foreign biomolecules or forms of life.

Among the potential candidates hitchhiking spacecraft are Betaproteobacteria of the genus

Ralstonia (order *Burkholderiales*). These bacteria are able to thrive under oligotrophic

conditions (7) and were reported to be ubiquitously present on space-related equipment

including water systems of the International Space Station (ISS) (8, 9), and the Mir space station

(10, 11). Likewise, they belonged to the microbial inventory of Mars Odyssey and Mars

Phoenix lander facilities and can thus prevail under strict Planetary Protection regulations (12,

13). *Ralstonia pickettii* strains were found to thrive in simulated microgravity compared to

normal gravity (11) and demonstrated high resistance against different metal ions and UV-C

radiation (8).

Ralstonia spp., (mainly *R. pickettii*), were previously found Antarctic soils (14), Antarctic snow

(15-17), in snow over Tibetan Plateau Glaciers (18), and in the air of the Antarctic base

Concordia (19). Interestingly, this genus has been regarded as an atmospheric traveler rather

than being part of true snow microflora in Antarctica (15). Despite a report on bacterial activity

at subzero temperatures in South pole snow (20), *Ralstonia* was not among active bacterial

communities as inferred from cDNA-based 16S rRNA amplicon sequencing (15). This view

was supported by a comprehensive Antarctic surface snow microbiome study that did not detect

this bacterial genus (21), and by the fact that spatial variability of snow microbiomes in

Antarctica is high (22). Although aerial dispersal is probably the major contributor to (micro-

)biological input to remote regions (23), the role of bioaerosol transport to microbial ecology

of isolated systems such as the Antarctic continent is poorly understood (24). Applying high

throughput sequencing approaches to study bacterial dispersal over the Antarctic continent

remains a considerable challenge due to the low microbial biomass of atmosphere-derived

samples regarding their DNA content (25), and resulting issues of recovering high quality

assemblies from metagenomic reads (26).

In addition to the limited knowledge about how transport via aerosols and snow across

Antarctica shapes microbial dispersal patterns, another open question relates to the distances

that microbes can cover within the atmosphere of extreme environments. A study on the

dispersal of airborne faecal coliforms showed a distribution over about just 175 m from a

sewage outfall at Rothera Research Station (Antarctic Peninsula) and thus prolonged survival

was considered unlikely (27). More stress-resistant microorganisms could, however, endure for

much longer periods. Recently, L. A. Malard et al. (22) found high abundances of spore-

forming *Bacilli* and suggested that long-term dispersal may seed continental Antarctic snow

ecosystems. However, to date, the role of aerial dispersal in shaping patterns of microbial

biogeography is supported by little empirical evidence (23).

Here, we follow the hypothesis that geographically widespread (28) *Ralstonia* spp. prophages

and/or replicating and adsorbed bacteriophages of this genus (all types further referred to as

"host-associated" phages) can be used to study host bacterium dispersal across the Antarctic

continent. Reconstructing prokaryotic genomes from samples containing low amounts of DNA,

including air or precipitation is challenging, as low input libraries (~1 pg) can result in problems

of genome binning (26). However, (pro)phage genomes or their fragments are much smaller

than prokaryotic genomes and thus easier to identify, track and compare. In this study, genome-

resolved metagenomics was applied to demonstrate dispersal of host-associated phage genome

fragments (PGFs) from surface snow across the Antarctic continent over hundreds to thousands

of kilometers. We detected PGFs belonging to the orders *Caudovirales* and *Tubulavirales* in

this extraterrestrial analogue and additionally show that similar genome-integrated phages are

frequent colonizers of space equipment. Therefore, we suggest that host-associated PGFs

represent a useful tool to study spatial dispersal of bacteria and their phages in extreme

environmental settings and further envision implications for the dispersal of microbiological

contaminations on spacecraft and celestial bodies that previously escaped Planetary Protection

measures.

Results

Reconstruction of low coverage MAGs from Antarctic snow metagenomic data

The microbial community composition of surface snow samples collected close to three Russian

Antarctic stations, Druzhnaja, Mirnii and Progress, based on 16S rRNA gene sequencing was

described earlier (16) and showed that *Ralstonia* was the most dominant organism in snow

collected at the Mirnii station, the second most dominant after *Janthinobacterium* at Druzhnaja,

and the third most dominant (after *Flavobacteria* and *Hydrogenophaga*) around Progress

station. We reconstructed four prokaryotic and one eukaryotic metagenome assembled genomes

(MAG) from the three low-biomass snow metagenomes (Figure 1A). Two MAGs related to

Ralstonia sp. and *R. pickettii* were recovered from Druzhnaja and Mirnii samples and had

86%/10% and 55%/0% completeness/contamination scores, respectively. MAGs of

Janthinobacterium lividum with 92%/6% scores and of *Flavobacterium micromati* with

96%/0% scores were recovered from Druzhnaja and Progress, respectively. We also identified

a MAG of a diatom (likely *Thalassiosira* sp.) in the Mirnii metagenome, which we used for

normalizing some of our viral analysis (see below) but was not further characterized in this

work. Read mapping revealed coverage scores of 5.9, 7.1, 6.7 and 14.6 for the MAGs of

Ralstonia sp*., R. pickettii, J. lividum* and *F. micromati*, respectively.

Prevalent absence of CRISPR-Cas systems suggest low adaptive immunity

We searched for clustered regularly interspaced short palindromic repeats (CRISPR) spacers in

the snow metagenome reads to link them to potential protospacers on PGFs. CRISPR arrays

and *cas* genes were absent from both *Ralstonia* and the *J. lividum* MAGs as determined by

CRISPRcasFinder (29). BLASTing of direct repeat (DR) sequences to NCBI's NR database

also did not indicate *Ralstonia* or *Janthinobacterium* to be the host of the CRISPR array. Solely

the genome of *F. micromati* contained two CRISPR arrays with four spacers each (both

evidence level 3 = highly likely candidates). CRISPR arrays were not detected on *R. pickettii*

draft genomes SSH4 and CW2 obtained from space equipment.

PGF-host analyses suggest shared hosts of Antarctic phages

A total of 26 predicted PGFs, eight of them being putative PGFs, was found in Antarctic snow

metagenomic data (Table S1). VIBRANT (30) additionally detected 52 PGFs, resulting in a

total of 78 PGFs (Figure 1A). PGFs with minimum 75% of the genome covered with reads had

coverages ranging between 1.9 and 17.5. Most PGFs were partial sequences according to

viralComplete (31)   and   CheckV   (32),   only   Antarcphage10_Mi_4716   and

Antarcphage49_Dr_7823_circ were estimated to be of full length (Table S1), although

annotations of Antarcphage10_Mi_4716 in comparison to *Ralstonia* phage p12J revealed

missing genes and let us question its completeness (Figure 2). Out of 78 PGFs, 77 were

categorized as lytic by VirSorter (33) and 75 were found to be of viral or unclassified origin by

CheckV (Table S1). A proportion of 43.6%, 46.1% and 10.3% identified PGFs originated from

the Druzhnaja, Mirnii and Progress, respectively.

All 78 PGFs were analyzed in conjunction with reconstructed MAGs using VirHostMatcher

(34). We defined a dissimilarity threshold of d2* value = 0.436, which corresponds to the lowest

dissimilarity value (= highest similarity) for PGFs matching the MAG of *Thalassiosira* sp. from

the snow environment (Figure 1A) by assuming that the eukaryotic MAG does not match any

of our extracted prokaryote infecting PGFs. In total, 50 of the 78 PGFs matched a host MAG

below the defined dissimilarity threshold, based on their shared k-mer patterns. Most PGFs (41)

matched both MAGs of *Ralstonia* sp. and *R. pickettii* (Figure 1A). In total, 38 and 14 PGFs

matched *J. lividum* and *F. micromati* MAGs, respectively. Five out of eight PGFs extracted

from the Progress station metagenomic snow sequences matched the *F. micromati* MAG

recovered from this station. Seven PGF matches were shared between all prokaryotic MAGs

(Figure S1). However, with 30 shared matches, the three MAGs belonging to the order

Burkholderiales shared the most overlap (Figure 1B, Figure S1).

Matching of CRISPR spacers derived from CRISPR loci of unknown hosts (reconstructed from

DR sequences) revealed that spacers from Mirnii matched 19, 14 and three PGFs from

Druzhnaja, Mirnii and Progress, respectively, and one *Flavobacterium* sp. spacer from Progress

matched a PGF from that station (Figure 1C, Table S2, ≥80% similarity). Out of these 37 spacer

matches, three matched protospacers of PGFs of unknown hosts, and 26 were assigned to a

Ralstonia host according to VirHostMatcher or other predictions (Table S1). Since p12D (see

below) was among the PGF spacer targets and is a certain *Ralstonia* phage, this could indicate

that an unknown host belonging to the order *Burkholderiales* uses adaptive immunity against

viruses. We compiled evidence of host prediction for the 78 PGFs in Table S1.

Figure 1: Host-phage pairings based on shared k-mer frequency patterns. A) Heatmap representing dissimilarity value d2* for matches between

five host MAGs and 78 PGFs derived from three Antarctic snow metagenomes from the stations Progress, Mirnii, and Druzhnaja. The PGF number

corresponds to the number in the phage name, for instance PGF#78 refers to Antarcphage78_Dr_3477 (Table S1). The table shows MAG

characteristics (phylogenetic affiliation, completeness, contamination, and coverage). n/a = not available **B)** Network of phage-host interactions based

190　　on k-mer frequency pattern reveals strong overlap between *Ralstonia* and *Janthinobacterium lividum* infecting PGFs. Here, only PGFs matching the

191　　host MAG below the defined threshold (see text) are shown. **C)** CRISPR spacer matches to PGFs. Different spacers matching to the same PGFs are

192　　shown by multiple arrows. The two hosts of CRISPR arrays for Mirnii-derived spacers remain unidentified according to their direct repeat sequence,

193　　whereas the Progress spacer is derived from a *Flavobacterium* sp. (Table S2).

The temperate phage p12D forms a distinct, monophyletic clade excluding most known

***Ralstonia* phages**

Among the 78 PGFs, we detected a circular (and thus complete) 7.8 kb PGF termed

Antarcphage49_Dr_7823_circ (Figure 2A, Figure S2). Since the Antarcphage49_Dr_7823_circ

PGF from the Druzhnaja station was found in the chromosome of a *R. pickettii* 12D strain

isolated from copper-contaminated sediment from a lake in Michigan with 99.9% identity

(accession number: CP001644.1- 2272699-2280518, Figure 2A), we here propose the name

p12D phage equivalent to *Ralstonia* p12J, a known phage infecting the *R. pickettii* 12J strain.

Antarcphage49_Dr_7823_circ (from now on referred to as p12D) only matched the MAG of

Ralstonia sp. and *R. pickettii* based on shared k-mer frequency patterns. Functional protein

annotations provided evidence that p12D contained a gene for a resolvase domain containing

protein/site-specific recombinase, likely used for integration into the host genome, as well as

for the zonular occludens toxin (Zot, PF05707) (Figure 2A, Figure S2, Table S3). The non-

toxic component of Zot at the N-terminus represents a characteristic protein in filamentous

phages that has been used for phage classification (35, 36). Reconstruction of a phylogenetic

relationships of Zot proteins from this study and respective references (Figure 2B, Figure S3)

confirmed a close identity of p12D to *Ralstonia* phage 1 NP-2014 and

Antarcphage79_WAP_18.3, all belonging to the same monophyletic clade. *Zot* of

Antarcphage10_Mi_4716 was phylogenetically related to filamentous *Ralstonia* phage p12J.

The tree shows four distinctive clusters for the Zot protein, reflecting the overall synteny of the

gene order of the different *Ralstonia* phages very well.

Annotations against UniRef100 revealed that 68.8% of all annotated Antarctic PGF proteins

remain hypothetical. The annotations taxonomically assigned 34.4% of all Antarctic PGF

proteins to either *Ralstonia* or *R. pickettii,* and 11.6 % to phages of *Janthinobacterium* or *J.*

219   *lividum*, potentially indicating lateral gene transfer between virus and host in their respective

220   evolution (37, 38) or supporting that these PGF indeed represent prophages. Three of the

221   Antarctic PGFs carried a site-specific integrase or resolvase domain, and 4.2% of genes were

222   related to phage structural proteins, e.g., head, tail, or capsid proteins (Table S4, Table S5).

223
224

Figure 2: Phage genome comparisons, functional annotations and phylogenetic

relationship based on zonular occludes toxin (Zot, Pfam-ID: PF05707). A) Synteny of

known *Ralstonia* sp. phages from NCBI and Antarctic PGFs from this study with similar coding

sequences (CDS) being in the same color if they occur on more than one phage genome.

Functional annotations performed by DRAM-v (115) are given for all colored CDS where

available. Vertical lines between sequences indicate regions of shared similarity shaded

according to tBLASTx (orange gradient for matches in the same direction or blue gradient for

inverted matches). Figure was created using Easyfig (119). **B)** The phylogenetic tree was built

using FastTree 2.1.11 (122) in Geneious 11.1.5 (112) under default settings and shows four

distinct clusters of Zot proteins based on their amino acid sequences (aligned with MUSCLE

(121)).

237   Cross-mapping and nucleotide variations reveal dispersal patterns of MAGs and PGFs

238   across Antarctica

Mapping of 100% identical reads from the three snow samples and the Western Antarctic

Peninsula (WAP, Figure 3A) seawater sample to the prokaryotic MAGs revealed that both

Ralstonia MAGs were detected at all sites (89-100% genome coverage). *J. lividum* was

considered absent from the Progress station (genome coverage of 62%) and detected at other

stations (minimum 95% covered genome). *F. micromati* was only detected in Progress (100%)

and Druzhnaja (96%) but considered absent from Mirnii (58.7%) and the WAP (1.8%). Cross-

mapping on the 78 PGFs demonstrated that PGFs derived from the Progress station could not

be found in the other two snow metagenomes (Figure 3B&C). By contrast, 16 different PGFs

from Druzhnaja were detected at Mirnii, and 27 Mirnii PGFs were found at Druzhnaja (Figure

3C). In addition, 4 and 9 PGFs from Mirnii and Druzhnaja were detected in the WAP dataset,

respectively (Figure 3 B&C), all of them having minimum 97% of their lengths covered with

reads from the snow metagenomes (Table S1). Of 43 PGFs that were found at both stations

(Mirnii and Druzhnaja) based on read mapping, 26 PGFs shared a viral cluster (VC), and 34

matched prokaryotic hosts based on k-mer frequencies between stations (Table S1).

Interestingly, nine of the 13 PGFs occurring in WAP and Mirnii/Druzhnaja samples contained

identical single nucleotide polymorphisms (SNPs, Figure 3D&E, Table S6) or were missing

common SNPs pointing towards a common phage population before dispersal led to separation.

Clustering of PGFs in VICTOR (39) and vConTACT2 (40) revealed that groups of two to four

PGFs could be assigned to twelve distinct viral genera-forming viral clusters (Figure 3E, Figure

S4 & S5, Table S1). Members of the same cluster were often recovered from different stations,

for example, Druzhnaja and Mirnii (Figure 3B), which are located 923 km apart (Figure 3C).

The phylogenomic Genome-BLAST Distance Phylogeny (GBDP) tree yielded average support

of 47% and 80% in the nucleic acid and amino acid-based analysis, respectively. OPTSIL

clustering yielded 78 and 64 species and genus clusters at the nucleic acid level and 78 and 70

at the amino acid level, respectively.

Figure 3: Evidence for dispersal of PGFs across the Antarctic continent. A) A map showing

the different snow sampling stations in the East and the sampling station for seawater at the

Western Antarctic Peninsula. Map was built using Ocean Data View (123). **B)** Pie charts

summarizing the number of PGFs considered present at each station based on cross mapping of

reads and the site of PGF assembly (phage origin). **C)** The heat map depicts the normalized

coverages based on cross-mapping of reads against the 78 PGFs and separation distances

between stations. White areas indicate PGFs that were not called present because they had less

than 75% of scaffold length covered in mapping (with least 1-fold coverage). **D)** Single-

nucleotide polymorphism (SNP) analysis of p12D/Antarcphage49_Dr_7823_circ based on

reads from Druzhnaja, Mirnii and the WAP. This PGF was absent from the metagenome of the

Progress station. Further SNP analysis data can be found in Table S6. **E)** Summary figure for

the major dispersal route and supporting evidence. The majority of PGF disperse between

Mirnii and Druzhnaja, and 13 PGFs additionally occurred at the WAP based on cross-mapping

(CM) and SNP analysis. Values refer to the number of tested features, which include number

of virus clusters (VC), number of shared viruses based on CM, virus-host matches based on k-

mer links (VHM), CRISPR-spacer matches (CSM). A VHM=1 indicates that one PGF has

infected host MAGs from two stations. No VC and only one CM were observed for Progress-

Druzhnaja and Progress-Mirnii. Therefore, this station was excluded from the summary.

Many PGFs formed VCs with no genomic relatedness to any known phages from the

ProkaryoticViralRefSeq v94 database. Others were, however, similar to known *Ralstonia* PGFs

(current family *Inoviridae,* order *Tubulavirales*) and hence clustered with those, such as p12D

with *Ralstonia* phage 1 NP-2014, or Antarcphage10_Mi_4716 with *Ralstonia* phage PE226 and

Ralstonia phage p12J (Figure 4, Table S1). Other Antarctic PGFs shared protein clusters with

Ralstonia phages from space equipment based on vConTACT2 or were related to phages of the

Caudovirales order from the RefSeq database (Figure 4).

Figure 4: Phage network of Antarctic PGFs derived from Druzhnaja, Mirnii, Western

Antarctic Peninsula (WAP) and space equipment (ISS/MarsOrb) clustered with viruses

of the viral RefSeq database. Based on shared protein clusters, Antarctic PGFs from this study

group with known *Ralstonia* phage of the family *Inoviridae* (A), with phages obtained from

Ralstonia isolates from space equipment (B) or with known phages of the order *Caudovirales*

(C). Black nodes refer prokaryotic viruses other than *Ralstonia* and PGFs from this study.

Interactions show relatedness of genomes on viral genus or higher taxonomic level. "Others"

refer to other known viral families not listed in the legend. Visualization was done using

Cytoscape 3.8.2. (109). For details about viral clusters, please see Table S1.

In summary, our results support the idea of dispersal of host-associated phages between widely

separated stations because a) cross-mapping revealed presence of Mirnii PGFs at Druzhnaja

and vice versa, presence of a temperate phage from snow in the WAP seawater dataset, and

presence of the hosts *Ralstonia* and *J. lividum* in metagenomes of most stations; b) PGFs in

widely separated locations carry identical SNPs or lack nucleotide variations; c) some PGFs

from Mirnii and Druzhnaja belong to the same genus/VC; and d) Mirnii and Druzhnaja PGFs

share host MAGs according to their k-mer frequency patterns. These observations were mainly

true for PGFs affiliated to *Ralstonia* and *J. lividum* MAGs and less so for *F. micromati* PGFs,

which could indicate that certain bacterial species or their phages in Antarctic snow are more

prone to atmospheric dispersal than others. Finally, we found CRISPR spacers from two

unidentified hosts (represented by two distinct CRISPR DR sequences) at the Mirnii station to

match PGFs from Druzhnaja, Mirnii and Progress. Mapping revealed little to no abundance of

Progress PGFs at other stations, and no *F. micromati* MAG was detected in Mirnii and WAP

samples. This could be related to the low biomass and/or or insufficient sequencing coverage.

However, spacers from Mirnii suggest an infection history with PGF from Progress, thus

reflecting (past) dispersal of host and/or phage.

***Ralstonia* phages occur across diverse ecosystems and on space equipment**

By BLASTing all recovered PGFs against the IMG/VR viral database (41), we found that p12D

shared high identity (85.9 %) with a *R. pickettii* prophage (IMG/VR v.2 scaffold ID

Ga0075447_10000781, genome ID: 3300006191) from a seawater metagenome of the WAP

(Figure 2), which was further validated by cross-mapping of WAP reads to p12D delivering

100% scaffold coverage with 90% identical reads. The WAP is 4710 km away from the

Druzhnaja station (Figure 3C). The PGF Antarcphage79_WAP_18.3 was obtained by

BLASTing the p12D scaffold against the WAP assembly but was assembled containing

genomic regions extending the actual PGF. Based on vConTACT2, it was affiliated to the same

viral genus as p12D. Other PGFs recovered from the Antarctic snow metagenome showed hits

to phages deposited at IMG/VR. Hits were related to phages originating from freshwater,

wastewater, groundwater, or phages that were associated with plant root microbial

communities, which is in accordance with the fact that *Ralstonia* sp. is a frequent

phytopathogen (42). Many of the matching entries in IMG/VR were identified as phages of

Ralstonia due to matching with CRISPR spacers (mostly of *R. solanacearum,* Table S7).

Altogether, 14 PGFs were found on space equipment, hereafter referred to as ISSphage (7) and

MarsOrbphage (7), for PGFs extracted from the *R. pickettii* strains CW2 and SSH4 draft

genomes from the ISS cooling system and Mars Odyssey Orbiter, respectively. Seven of these

PGFs were identified as lysogenic and according to protein annotations (Table S4 & S5) rely

on proteins of the integrase family for integration into the host chromosome. vConTACT2

revealed that some of the ISSphage4 formed a common genus cluster with four Antarctic PGFs,

MarsOrbphage5 shared protein clusters with ISSphage3, and MarsOrbphage7 formed a genus

cluster with several *Burkholderia* phages among other *Caudovirales* members (Figure 4, Table

S1).

Discussion

CRISPR-Cas systems equip bacteria and archaea with a powerful defense system against

invading mobile genetic elements including plasmids, phages and viruses. Only ca. 50% of

bacteria rely on this adaptive immune system, and only 31% of genomes from public databases

belonging to plant-pathogenic *Ralstonia solanacearum* contain CRISPR-Cas arrays (43). Upon

exposure to a virulent phage under laboratory conditions, the CRISPR array of *R. solanacearum*

strain CFBP2957 did not acquire new spacers from viral protospacers (43). O. S. Gonçalves et

al. (28) reported that in the presence of CRISPR arrays, 27.9% of CRISPR spacers from

Ralstonia genomes targeted prophage elements. In our study, two *Ralstonia* MAGs from a low

temperature environment were devoid of CRISPR systems, implying that representatives of this

genus have alternative strategies for defense against mobile genetic elements (43). However,

analyses on more *Ralstonia* genomes are necessary to corroborate this statement.

Based on shared protein clusters, some Antarctic PGFs such as Antarcphage10_Mi_4716,

Antarcphage48_Dr_1123,   Antarcphage17_Mi_3026,   and p12D clustered with known

Ralstonia phages including *Ralstonia* phage p12J (NC_005131.2), *Ralstonia* phage PE226

(NC_015297.1) or *Ralstonia* phage 1 NP-2014 (NC_023586.1). These are filamentous phages

of the *Inoviridae*, a family that has been recently called for reclassification to higher taxonomic

ranks (36, 44). Inoviruses (order *Tubulavirales*) typically feature circular, single stranded DNA

genomes of ~5-15 kb length, lead to chronic infections, and are globally abundant (36). As our

protocol should have only revealed double-stranded DNA, detection of ssDNA inoviruses

sequences shows that these must be either replicating phages or genome-integrated prophages.

Zot, which was detected on p12D, Antarcphage10_Mi_4716 and Antarcphage79_WAP_18.3 is

a typical gene in filamentous *Ralstonia* phages (45, 46). This phage type confers little burden

to its host or can even serve it, e.g., by increasing its hosts' virulence and evolutionary fitness,

and because virions can leave the host in a non-destructible way (35, 36). *Ralstonia* phages

detected in Arctic viromes were shown to transduce genomic information of cold-shock

proteins to their hosts (47), a clear asset for microorganisms in polar environments. However,

we did not find transduction of beneficial genes, which seems to be common in extreme

environments (48).

Some *Ralstonia* phages occur as non-integrative, episomal forms, e.g., RS603, a hybrid of

RSM1/3 infecting the phytopathogen *R. solanacearum* (49)*,* whose genome lacks a resolvase

domain (Figure 2A), but many mesophilic *Ralstonia* also occur as lysogens (50). Lysogeny, a

lifestyle during which the phage genome becomes integrated into the host chromosome, is a

widespread phenomenon in low temperature environments (51-53) and likely attributed to

prolonged starvation and low activity of host cells under harsh conditions, the latter being

previously reported for *Ralstonia* (15). Since p12D and its counterpart from the WAP have a

resolvase-domain containing protein likely functioning in integration/excision during

lysogenisation (54, 55), we conclude that they must be temperate phages of *Ralstonia*.

Many, i.e., 75 of the 78 Antarctic PGFs found in this study shared little protein clusters with

known phages from public databases. This is certainly related to the often high diversity of

viruses, limited accessibility to Antarctic environments as well as a stronger focus on

sequencing metaviromes and viral isolates of direct human interest (56). The missing

relatedness is known for ssDNA viruses originating from Antarctic cryoconite holes and was

attributed to the isolation and extreme environmental conditions at the Antarctic continent (57).

In total, 45 of the 78 PGFs could be assigned to a host, which in 71% of cases was *Ralstonia*

(summarized in Table S1, column "host prediction").

Aeolian transport of viruses over polar environments, especially attached to snowflakes, has

been barely investigated to date. Former investigations, mainly conducted at low latitudes,

demonstrated intercontinental transport of microbes by winds (58), and that highly identical

phages can be found in distantly related areas and in various ecosystems around the globe (59-

62). C. M. Bellas et al. (63) recently reported the presence of near-identical phage genomes

being spread by up to 4000 km in cryoconite holes of Svalbard, Greenland, and the Alps. Our

data show that despite the isolation of the Antarctic continent, and under no consideration of

anthropogenic dispersal (64), bacteria and phage distribution via snow over extensive distances

across Antarctica is possible. Man-made dispersal seems unlikely for our samples, due to the

relatedness of Antarctic phages to phages from environmental sources according to database

hits. We cannot be certain about the nature of the PGFs (prophage or lytic phage) by

metagenomic predictions alone. While the *Inoviridae* fraction likely occurs as prophages or

episomal forms, detection of PGFs assigned to lytic categories by VIBRANT and VirSorter

(category 1-3) suggests that host-associated, lytic phages captured at the adsorption or infection

stages were present as well. The degree of uncertainty about the category of a virus presumably

results from many fragmented scaffolds with relatively low coverages, which were however

sufficient to identify shared elements of viruses between stations. Conclusions about the

presence/absence of hallmark genes should be drawn from more complete datasets based on

greater sequencing depth of samples containing more biomass. We further commend

experiments that involve cultivation attempts and sequencing of metaviromes (free phages) to

reveal the extent of lysogenic or chronic compared to lytic infection styles in Antarctic snow.

Snow PGFs carried identical SNPs with those from the seawater metagenome from the WAP

located 4710 km and 5338 km apart from Druzhnaja and Mirnii, respectively (Figure 3C),

implying long distance transport. This is further supported by the detection of their hosts,

Ralstonia and *J. lividum,* in the WAP. From seawater, a transmission route via sea spray

aerosols to snow that is blown over ice surfaces (65) can be assumed. Aerosolization of bacteria

from the sea surface is highly taxon-specific but seems to work well for *R. pickettii* (66) and

also viruses (67). Alternatively, a transport via aerosols to clouds and precipitating snow is

possible. In the latter scenario, bioaerosols including microbial cells would act as ice nucleation

particles (68), transferring microbes to ice clouds where they might induce their own

precipitation (69, 70). The abundance data support the isolation of Progress-derived PGFs and

their potential host *F. micromati* and point towards a decreasing gradient of abundances from

Druzhnaja to Mirnii to the WAP for many PGFs (Table S1). Thus, general dispersal patterns of

microbes across Antarctica seem governed by westward drift and are probably mediated by the

prevailing Southern Hemisphere westerly winds (71). On short spatial-temporal scales,

dispersal seems more complex and is probably shaped by multifactorial dependencies such as

the different potential of a species to become airborne (66), meteorological conditions or the

local geography. For instance, while Druzhnaja is located ~50 km into the continent, the stations

Mirnii and Progress are near the coast and more exposed to the sea in summer when ice breaks

occur.

We assume that the transferability of the use of PGFs to study microbial dispersal in space

analogues such as the Antarctic continent is likely applicable to other celestial bodies like Mars.

Three celestial bodies in our solar system (Mars, Europa, and Enceladus) have environmental

conditions that could favor microbial life (reviewed by M. G. Netea et al. (72)). Most microbial

isolates (85-95%) obtained from spacecraft and assembly facilities are associated with humans

(73). Since *Ralstonia* spp. can be human and plant pathogens (42, 74) and are able to thrive

under harsh and oligotrophic conditions (7), they might contribute to the transmission of viruses

to extraterrestrial environments, particularly via manned missions.

We found evidence for viral signatures associated with two *R. pickettii* strain draft genomes

previously obtained from space equipment of a spacecraft assembly clean room and from water

systems of the ISS. This result in conjunction with the result that temperate *Ralstonia* phages

(and other viruses) can undergo long-range dispersal in association with their hosts across the

extraterrestrial analogue Antarctica suggest that contaminations of space equipment with

particularly persistent microbes such as *Ralstonia* should receive more focus during

microbiological monitoring in the framework of Planetary Protection. However, the field of

astrovirology has so far generally found little attention (75). The contribution of lysogenic and

episomal phage ('hidden hitchhikers') to overall viral loads on spacecraft and associated

equipment has been overlooked despite early work reporting on alterations in prophage

induction during spaceflight (76-78), tobacco mosaic virus to survive space flight equivalent

proton irradiation (79), the occurrence of phages and human-related circoviruses in clean rooms

(80) and inoviruses on the ISS (81, 82).

Planetary Protection aims to prevent the spread of biological contaminants (forward

contamination) to space shuttles and stations as well as extraterrestrial environments of the solar

system. However, this policy largely ignores the potential of escaped biological contaminants

to heavily disperse on foreign celestial bodies once being released, for instance after crash

landings as happened for the Schiaparelli module of the ExoMars program in 2016 (83). Our

results show that host-associated PGFs are not only suitable indicators for tracking long-

distance dispersal in space analogues but also demonstrate that the release of contaminants that

previously escaped Planetary Protection measures could spread far across extraterrestrial

ecosystems and, in the worst-case scenario, confound future life detection missions.

Material and Methods

Metagenomic and genomic data processing

We analyzed publicly available metagenomic data sets, which correspond to Antarctic surface

snow collected around three Russian stations (Druzhnaja, Mirnii, Progress) and are deposited

at NCBI's Sequence Read Archive (SRA) as Bioproject PRJNA674475 (Table S8) and MG-

RAST under project accession mgp13052 including taxon abundance data. Snow sampling was

conducted in December 2008 and 2009 as described previously (15); in brief, a sterile plastic

scoop was used to sample ~10 kg corresponding to a 2-3 cm layer of surface snow across several

1 m^2 areas. Snow was melted over a period of 12 hours and concentrated using Pellicon

tangential flow filters (Millipore, Burlington, MA, USA) to a final volume of ~ 10 mL. DNA

extraction for these metagenomes was carried out with the DNA Blood & Tissue kit (Qiagen,

Hilden, Germany), whose protocol causes removal of free viruses, and resulted in DNA

amounts of 170 – 490 ng. Sequencing libraries were prepared with MiSeq reagent kit v.2

(Illumina, USA), targeting mainly dsDNA viruses. Metagenomic data of a seawater sample

from the WAP was also obtained from SRA (accession #SRR5591034). Raw shotgun

sequencing reads of the three snow metagenomes and the WAP dataset were quality-trimmed

using BBDuk (https://github.com/BioInfoTools/BBMap/blob/master/sh/bbduk.sh) from the

BBTools package (84) and Sickle (85) resulting in read counts between 1.31 – 2.65 Mio. for

the three snow samples. Assembly of reads was done using MetaSPAdes version 3.13 (86), and

scaffolds <1 kbps were removed. Draft genome sequences of *R. pickettii* strains SSH4 and CW2

isolated from space equipment were obtained from Genbank accession #JFZG00000000 and

#JFZH00000000. Strains SSH4 and CW2 were isolated pre-flight from the surface of the Mars

Odyssey Orbiter during assembly and from a water sample taken in-flight from the ISS cooling

system, respectively (9).

Reconstruction of microbial genomes from metagenomes and CRISPR prediction

Binning of MAGs from the three stations Druzhnaja, Mirnii and Progress was performed using

Emergent   self-organizing   maps   (ESOMs,   (87)),   ABAWACA

(https://github.com/CK7/abawaca) and MaxBin 2.0 (88). Aggregation of bins was performed

using DASTool (89) and curation of bins was done in uBin (90), also delivering contamination

and completeness scores (91). Read coverage of recovered MAGs was obtained from read

mapping using Bowtie2 (92) in sensitive mode to the individual bins, followed by mismatch

filtering (2% mismatch allowance, depending on read length). To investigate the dispersal of

MAGs, mismatch criteria were set to 0% mismatches (100% similarity) and calcopo.rb

(https://github.com/ProbstLab/viromics/tree/master/calcopo) was used to calculate the

coverage per nucleotide (breadth) and the percentage of positions in the genome covered by

reads. Only genomes with a least 70% breadth were considered present in the respective

metagenome.

CRISPR arrays, which represent the prokaryotic adaptive immunity, were searched in host

MAGs using CRISPRcasFinder and considering evidence level 3 or 4 (29). CRISPR loci consist

of repeats interspaced with short DNA sequences (spacers) obtained from invading mobile

genetic elements such as phages and thus provide a record of past infections. Since CRISPR

arrays might get lost during the binning processes, e.g., due to fragmentation in assembly of

strain variants, absence of CRISPR arrays in CRISPRcasFinder was further investigated by

reconstructing CRISPR systems from raw reads using Crass (93) and BLASTing the obtained

direct repeat (DR) sequences, which are phylogenetically well-conserved (94), against the

NCBI non-redundant database (release 1st March 2021) using BLASTn --short algorithm (95)

with subsequent filtering at 80% similarity and e-value threshold of 10e-05. DR sequences were

BLASTed against the MAGs and used to extract CRISPR spacers from reads using

MetaCRAST with settings -d 3 -l 60 -c 0.99 -a 0.99 -r (96). Spacers were matched against PGFs

as mentioned above for DRs. In most cases, BLASTing the DR sequences against the NR

database did not reveal the host's identity. Nevertheless, spacers derived from unknown hosts

were considered, as their matches show that targeted PGFs represent true mobile genetic

elements, and matches can be used to infer infection patterns between stations.

PGF detection, host allocation and viral clustering

PGFs were identified from metagenome assemblies using a combination of bioinformatic tools,

namely Virsorter v1 (33), VirFinder (97), CircMG (98), renamed to VRCA

(https://github.com/alexcritschristoph/VRCA),   VOGdb   (version   VOG93)   (99),   and

Endmatcher   (https://github.com/ProbstLab/viromics/tree/master/Endmatcher).   The

classification of predicted PGFs as "putative viruses" and "viruses" was done as previously

described in the Supplementary Figure 3 of (100). VIBRANT v.1.2.1 (30) with default settings

was used to find additional PGFs. No length cut-off was set (101), since many known *Ralstonia*

phages and inoviruses have genome sizes <10 kb (36, 50), and because of the low biomass of

Antarctic snow samples little PGFs compared to other ecosystems were expected. VirSorter

and VIBRANT aided in (pro)phage detection in draft genomes of *R. pickettii* from space

equipment. CheckV (32) was used to determine the type of identified PGF and completeness,

and viralComplete (31) was applied to predict closely related phages. PGFs associated with

MAGs were identified by grepping the scaffold ID on the bins. Pairwise comparisons of

Antarctic PGFs and clustering was done using nucleic acid- and amino acid-based VICTOR

(39). The resulting intergenomic distances were used to infer a balanced minimum evolution

tree with branch support via FASTME including Subtree Pruning and Regrafting (SPR)

postprocessing (102) using the distance formula D0. Branch support was inferred from 100

pseudo-bootstrap replicates each. Trees were rooted at the midpoint (103) and visualized with

FigTree v1.4.4 (104). Taxon boundaries at the species and genus level were estimated with the

OPTSIL program (105), using the recommended clustering thresholds (39) and an F value

(fraction of links required for cluster fusion) of 0.5 (106).

Clustering of PGFs from Antarctic snow and space equipment was further substantiated via

vConTACT2 v.0.9.19 (40, 107) in combination with the ProkaryoticViralRefSeq database

(v94, (108)) followed by visualization of viral clusters (VC) in Cytoscape v. 3.8.2 (109). Virus-

host matches were determined using the tool VirHostMatcher (34) with d_2* oligonucleotide

frequency dissimilarity measures for a k-mer length of 6. Viral genus clusters were determined

using VIRIDIC (110). Venn diagrams were calculated using the VIB-ugent webtool

(http://bioinformatics.psb.ugent.be/webtools/Venn/).

Mapping of reads to assembled PGFs and nucleotide polymorphism analysis

We assume that presence of a PGF at two locations confirmed by read mapping represents

dispersal. To determine if an assembled PGF from a single sample occurred in other

metagenomes even if not being assembled, we performed read mapping following the

previously published guidelines (101): that reads should map to a PGF with at least 90% identity

(Bowtie2 settings as in (111)), and more than 75% of scaffold should have a coverage of at least

1x. To detect breadth of a PGF, we again used calcopo.rb (see above). Mean coverage of PGFs

was   calculated   using   calc_coverage_v3 (https://github.com/ProbstLab/uBin-

helperscripts/blob/master/bin/04_01calc_coverage_v3.rb) and normalized to sequencing depth.

Analysis of nucleotide polymorphisms was conducted for the 13 PGFs that underwent long

range dispersal, i.e., PGF being present in the WAP sample and at least one of the snow

metagenomes based on read mapping (Table S1). Variant analysis was performed in Geneious

11.1.5 (112) by applying default settings to the read-mappings generated as explained above.

Gene prediction and annotations

Open reading frames on PGFs were detected using Prodigal in meta mode (113). Functional

and taxonomic annotations of predicted proteins of PGFs were performed by DIAMOND

searches with a e-value of 10e-05 (114) against FunTaxDB (90) and by using DRAM-v (115).

For the full-length genome of the PGF p12D, annotations were improved using HHpred against

PDB, Pfam, UniProt-SwissProt-viral and NCBI_Conserved Domains ((116, 117),

https://toolkit.tuebingen.mpg.de/tools/hhpred) with a probability threshold of 70%. Sequences

of PGFs were BLASTed against IMG/VR 2.0 (118) with an e-value cut-off of 10e-05 to find

related phages from other metagenomic datasets.

Synteny of *Ralstonia* phage and phylogenetic comparison of the *zot* gene

Synteny of known *Ralstonia* sp. phages from NCBI and all Antarctic PGFs that were identified

herein and carried the zonular occludens toxin (*zot*, Pfam-ID: PF05707) was performed with

tBLASTx comparisons using Easyfig v.2.2.5 (119) on .gbk files generated by Prokka (120). A

phylogenetic tree for the MUSCLE-aligned (121) amino acid and nucleic acid sequences of *zot*

was constructed using the FastTree (122) algorithm in Geneious 11.1.5 (112).

Acknowledgements

We acknowledge Ken Dreger for server administration and maintenance as well as Cristina

Moraru for sharing insights on virus taxonomy.

Author Contribution statement

J.R. designed the study, wrote the manuscript, and carried out the analyses with input from

T.L.V.B.; A.L. and K.S. generated the raw data and performed the sampling; A.J.P

conceptualized the project, provided supervision, resources, and was involved in data

interpretation; All authors edited drafts of the manuscript.

Author Disclosure Statement

For all authors, no competing financial interests exist.

Funding statement

We acknowledge funding from the German Aerospace Center (DLR) for the project DISPERS

(50WB1922). J.R. was partially supported by the German Science Foundation for the project

VIBOCAT (grant number DFG RA3432/1-1). A.J.P. and T.L.V.B. were supported by the

Ministerium   für   Kultur   und   Wissenschaft   des   Landes   Nordrhein-Westfalen

("Nachwuchsgruppe Dr. Alexander Probst").

References:

1.   Pyne SJ. 2007. The extraterrestrial Earth: Antarctica as analogue for space

exploration. Space Policy 23:147-149.

2.   Lugg D, Shepanek M. 1999. Space analogue studies in Antarctica. Acta Astronaut

44:693-9.

3.   Onofri S, Selbmann L, Zucconi L, Pagano S. 2004. Antarctic microfungi as models for

exobiology. Planet Space Sci 52:229-237.

4.   Puskeppeleit M, Quintern LE, El Naggar S, Schott JU, Eschweiler U, Horneck G,

Bucker H. 1992. Long-term dosimetry of solar UV radiation in Antarctica with spores

of *Bacillus subtilis*. Appl Environ Microbiol 58:2355-9.

5.   Nicholson WL, Munakata N, Horneck G, Melosh HJ, Setlow P. 2000. Resistance of

Bacillus endospores to extreme terrestrial and extraterrestrial environments. Microbiol

Mol Biol Rev 64:548-72.

6.   Onofri S, Barreca D, Selbmann L, Isola D, Rabbow E, Horneck G, De Vera J, Hatton

J, Zucconi L. 2008. Resistance of Antarctic black fungi and cryptoendolithic

communities to simulated space and Martian conditions. Stud Mycol 61:99-109.

7.   McAlister MB, Kulakov LA, O'Hanlon JF, Larkin MJ, Ogden KL. 2002. Survival and

nutritional requirements of three bacteria isolated from ultrapure water. J Ind

Microbiol Biotechnol 29:75-82.

8.   Mijnendonckx K, Provoost A, Ott CM, Venkateswaran K, Mahillon J, Leys N, Van

Houdt R. 2013. Characterization of the survival ability of *Cupriavidus metallidurans*

and *Ralstonia pickettii* from space-related environments. Microb Ecol 65:347-60.

9.   Monsieurs P, Mijnendonckx K, Provoost A, Venkateswaran K, Ott CM, Leys N, Van

Houdt R. 2014. Draft genome sequences of *Ralstonia pickettii* strains SSH4 and CW2,

isolated from space equipment. Genome Announc 2.

10.   Ott CM, Bruce RJ, Pierson DL. 2004. Microbial characterization of free floating

condensate aboard the Mir space station. Microb Ecol 47:133-6.

11.   Baker PW, Leff L. 2004. The effect of simulated microgravity on bacteria from the

Mir space station. Microgravity Sci Technol 15:35-41.

12.   Vaishampayan P, Osman S, Andersen G, Venkateswaran K. 2010. High-density 16S

microarray and clone library-based microbial community composition of the Phoenix

spacecraft assembly clean room. Astrobiology 10:499-508.

13.   La Duc MT, Nicholson W, Kern R, Venkateswaran K. 2003. Microbial

characterization of the Mars Odyssey spacecraft and its encapsulation facility. Environ

Microbiol 5:977-85.

14.   Lysak V, Maksimova IA, Nikitin DA, Ivanova AE, Kudinova AG, Soina VS,

Marfenina OE. 2018. Soil microbial communities of Eastern Antarctica. Moscow

Univ Biol Sci Bull 73:104-112.

15.   Lopatina A, Krylenkov V, Severinov K. 2013. Activity and bacterial diversity of snow

639   around Russian Antarctic stations. Res Microbiol 164:949-58.

640   16.   Lopatina A, Medvedeva S, Shmakov S, Logacheva MD, Krylenkov V, Severinov K.

641   2016. Metagenomic analysis of bacterial communities of Antarctic surface snow.

642   Front Microbiol 7:398.

643   17.   Antony R, Mahalinganathan K, Krishnan KP, Thamban M. 2012. Microbial

644   preference for different size classes of organic carbon: a study from Antarctic snow.

645   Environ Monit Assess 184:5929-43.

646   18.   Liu Y, Yao T, Jiao N, Kang S, Xu B, Zeng Y, Huang S, Liu X. 2009. Bacterial

647   diversity in the snow over Tibetan Plateau Glaciers. Extremophiles 13:411-23.

648   19.   Van Houdt R, De Boever P, Coninx I, Le Calvez C, Dicasillati R, Mahillon J,

649   Mergeay M, Leys N. 2009. Evaluation of the airborne bacterial population in the

650   periodically confined Antarctic base Concordia. Microb Ecol 57:640-8.

651   20.   Carpenter EJ, Lin S, Capone DG. 2000. Bacterial activity in South Pole snow. Appl

652   Environ Microbiol 66:4514-7.

653   21.   Michaud L, Lo Giudice A, Mysara M, Monsieurs P, Raffa C, Leys N, Amalfitano S,

654   Van Houdt R. 2014. Snow surface microbiome on the High Antarctic Plateau (DOME

655   C). PLoS One 9:e104505.

656   22.   Malard LA, Sabacka M, Magiopoulos I, Mowlem M, Hodson A, Tranter M, Siegert

657   MJ, Pearce DA. 2019. Spatial variability of Antarctic surface snow bacterial

658   communities. Front Microbiol 10:461.

659   23.   Pearce DA, Alekhina IA, Terauds A, Wilmotte A, Quesada A, Edwards A,

660   Dommergue A, Sattler B, Adams BJ, Magalhaes C, Chu WL, Lau MC, Cary C, Smith

661   DJ, Wall DH, Eguren G, Matcher G, Bradley JA, de Vera JP, Elster J, Hughes KA,

662   Cuthbertson L, Benning LG, Gunde-Cimerman N, Convey P, Hong SG, Pointing SB,

Pellizari VH, Vincent WF. 2016. Aerobiology over Antarctica - A new initiative for

atmospheric ecology. Front Microbiol 7:16.

24. Bottos EM, Woo AC, Zawar-Reza P, Pointing SB, Cary SC. 2014. Airborne bacterial

populations above desert soils of the McMurdo Dry Valleys, Antarctica. Microb Ecol

67:120-8.

25. Behzad H, Gojobori T, Mineta K. 2015. Challenges and opportunities of airborne

metagenomics. Genome Biol Evol 7:1216-26.

26. Bowers RM, Clum A, Tice H, Lim J, Singh K, Ciobanu D, Ngan CY, Cheng JF,

Tringe SG, Woyke T. 2015. Impact of library preparation protocols and template

quantity on the metagenomic reconstruction of a mock microbial community. BMC

Genomics 16:856.

27. Hughes KA. 2003. Aerial dispersal and survival of sewage-derived faecal coliforms in

Antarctica. Atmos Environ 37:3147-3155.

28. Gonçalves OS, de Oliveira Souza F, Bruckner FP, Santana MF, Alfenas-Zerbini P.

2021. Widespread distribution of prophages signaling the potential for adaptability

and pathogenicity evolution of *Ralstonia solanacearum* species complex. Genomics

113:992-1000.

29. Couvin D, Bernheim A, Toffano-Nioche C, Touchon M, Michalik J, Neron B, Rocha

EPC, Vergnaud G, Gautheret D, Pourcel C. 2018. CRISPRCasFinder, an update of

CRISRFinder, includes a portable version, enhanced performance and integrates

search for Cas proteins. Nucleic Acids Res 46:W246-W251.

30. Kieft K, Zhou Z, Anantharaman K. 2020. VIBRANT: automated recovery, annotation

and curation of microbial viruses, and evaluation of viral community function from

genomic sequences. Microbiome 8:90.

31. Antipov D, Raiko M, Lapidus A, Pevzner PA. 2020. Metaviral SPAdes: assembly of

viruses from metagenomic data. Bioinformatics 36:4126-4129.

32.   Nayfach S, Camargo AP, Schulz F, Eloe-Fadrosh E, Roux S, Kyrpides NC. 2020.

690   CheckV assesses the quality and completeness of metagenome-assembled viral

691   genomes. Nat Biotechnol doi:10.1038/s41587-020-00774-7.

692   33.   Roux S, Enault F, Hurwitz BL, Sullivan MB. 2015. VirSorter: mining viral signal

693   from microbial genomic data. PeerJ 3:e985.

694   34.   Ahlgren NA, Ren J, Lu YY, Fuhrman JA, Sun F. 2017. Alignment-free d_2^{*}

695   oligonucleotide frequency dissimilarity measure improves prediction of hosts from

696   metagenomically-derived viral sequences. Nucleic Acids Res 45:39-53.

697   35.   Hay ID, Lithgow T. 2019. Filamentous phages: masters of a microbial sharing

698   economy. EMBO Rep 20.

699   36.   Roux S, Krupovic M, Daly RA, Borges AL, Nayfach S, Schulz F, Sharrar A, Matheus

700   Carnevali PB, Cheng JF, Ivanova NN, Bondy-Denomy J, Wrighton KC, Woyke T,

701   Visel A, Kyrpides NC, Eloe-Fadrosh EA. 2019. Cryptic inoviruses revealed as

702   pervasive in bacteria and archaea across Earth's biomes. Nat Microbiol 4:1895-1906.

703   37.   Filée J, Forterre P, Laurent J. 2003. The role played by viruses in the evolution of their

704   hosts: a view based on informational protein phylogenies. Res Microbiol 154:237-243.

705   38.   Moreira D. 2000. Multiple independent horizontal transfers of informational genes

706   from bacteria to plasmids and phages: implications for the origin of bacterial

707   replication machinery. Mol Microbiol 35:1-5.

708   39.   Meier-Kolthoff JP, Göker M. 2017. VICTOR: genome-based phylogeny and

709   classification of prokaryotic viruses. Bioinformatics 33:3396-3404.

710   40.   Bolduc B, Jang HB, Doulcier G, You ZQ, Roux S, Sullivan MB. 2017. vConTACT:

711   an iVirus tool to classify double-stranded DNA viruses that infect Archaea and

712   Bacteria. PeerJ 5:e3243.

41.   Prospero JM, Blades E, Mathison G, Naidu R. 2005. Interhemispheric transport of

viable fungi and bacteria from Africa to the Caribbean with soil dust. Aerobiologia

21:1-19.

42.   Askora A, Kawasaki T, Usami S, Fujie M, Yamada T. 2009. Host recognition and

integration of filamentous phage phiRSM in the phytopathogen, *Ralstonia*

solanacearum. Virology 384:69-76.

43.   da Silva Xavier A, de Almeida JCF, de Melo AG, Rousseau GM, Tremblay DM, de

Rezende RR, Moineau S, Alfenas-Zerbini P. 2019. Characterization of CRISPR-Cas

systems in the *Ralstonia solanacearum* species complex. Mol Plant Pathol 20:223-

239.

44.   Koonin EV, Dolja VV, Krupovic M, Varsani A, Wolf YI, Yutin N, Zerbini FM, Kuhn

JH. 2020. Global organization and proposed megataxonomy of the virus world.

Microbiol Mol Biol Rev 84.

45.   Murugaiyan S, Bae JY, Wu J, Lee SD, Um HY, Choi HK, Chung E, Lee JH, Lee SW.

2011. Characterization of filamentous bacteriophage PE226 infecting *Ralstonia*

solanacearum strains. J Appl Microbiol 110:296-303.

46.   Mai-Prochnow A, Hui JG, Kjelleberg S, Rakonjac J, McDougald D, Rice SA. 2015.

'Big things in small packages: the genetics of filamentous phage and effects on fitness

of their host'. FEMS Microbiol Rev 39:465-87.

47.   Sanguino L, Franqueville L, Vogel TM, Larose C. 2015. Linking environmental

prokaryotic viruses and their host through CRISPRs. FEMS Microbiol Ecol 91.

48.   Hwang Y, Rahlff J, Schulze-Makuch D, Schloter M, Probst AJ. 2021. Diverse Viruses

Carrying Genes for Microbial Extremotolerance in the Atacama Desert Hyperarid

Soil. mSystems 6:e00385-21.

49.   Van TTB, Yoshida S, Miki K, Kondo A, Kamei K. 2014. Genomic characterization of

φRS603, a filamentous bacteriophage that is infectious to the phytopathogen Ralstonia

solanacearum. Microbiology and immunology 58:697-700.

50.   Askora A, Yamada T. 2015. Two different evolutionary lines of filamentous phages in

Ralstonia solanacearum: their effects on bacterial virulence. Front Genet 6:217.

51.   Filippova SN, Surgucheva NA, Sorokin VV, Akimov VN, Karnysheva EA, Brushkov

AV, Andersen D, Gal'chenko VF. 2016. Bacteriophages in Arctic and Antarctic low-

temperature systems. Microbiology 85:359-366.

52.   Dziewit L, Radlinska M. 2016. Two inducible prophages of an Antarctic

Pseudomonas sp. ANT_H14 use the same capsid for packaging their genomes -

Characterization of a novel phage helper-satellite system. PLoS One 11:e0158889.

53.   Filippova SN, Surgucheva NA, Kulikov EE, Sorokin VV, Akimov VN, Bej AK,

McKay C, Andersen D, Galchenko VF. 2013. Detection of phage infection in the

bacterial population of Lake Untersee (Antarctica). Microbiology 82:383-386.

54.   Askora A, Kawasaki T, Fujie M, Yamada T. 2011. Resolvase-like serine recombinase

mediates integration/excision in the bacteriophage phiRSM. J Biosci Bioeng 111:109-

16.

55.   Ahmad AA, Stulberg MJ, Mershon JP, Mollov DS, Huang Q. 2017. Molecular and

biological characterization of varphiRs551, a filamentous bacteriophage isolated from

a race 3 biovar 2 strain of *Ralstonia solanacearum*. PLoS One 12:e0185034.

56.   Rodrigues RAL, Andrade A, Boratto PVM, Trindade GS, Kroon EG, Abrahao JS.

2017. An anthropocentric view of the virosphere-host relationship. Front Microbiol

8:1673.

57.   Sommers P, Fontenele RS, Kringen T, Kraberger S, Porazinska DL, Darcy JL,

Schmidt SK, Varsani A. 2019. Single-stranded DNA viruses in Antarctic cryoconite

holes. Viruses 11.

763 58. Smith DJ, Timonen HJ, Jaffe DA, Griffin DW, Birmele MN, Perry KD, Ward PD,

764   Roberts MS. 2013. Intercontinental dispersal of bacteria and archaea by transpacific

765   winds. Appl Environ Microbiol 79:1134-9.

766 59. Short CM, Suttle CA. 2005. Nearly identical bacteriophage structural gene sequences

767   are widely distributed in both marine and freshwater environments. Appl Environ

768   Microb 71:480-486.

769 60. Breitbart M, Rohwer F. 2005. Here a virus, there a virus, everywhere the same virus?

770   Trends Microbiol 13:278-84.

771 61. Breitbart M, Miyake JH, Rohwer F. 2004. Global distribution of nearly identical

772   phage-encoded DNA sequences. FEMS Microbiol Lett 236:249-56.

773 62. Li Y, Endo H, Gotoh Y, Watai H, Ogawa N, Blanc-Mathieu R, Yoshida T, Ogata H.

774   2019. The Earth is small for "Leviathans": Long distance dispersal of giant viruses

775   across aquatic environments. Microbes Environ 34:334-339.

776 63. Bellas CM, Schroeder DC, Edwards A, Barker G, Anesio AM. 2020. Flexible genes

777   establish widespread bacteriophage pan-genomes in cryoconite hole ecosystems. Nat

778   Commun 11:4403.

779 64. Hughes KA, Convey P, Pertierra LR, Vega GC, Aragon P, Olalla-Tarraga MA. 2019.

780   Human-mediated dispersal of terrestrial species between Antarctic biogeographic

781   regions: A preliminary risk assessment. J Environ Manage 232:73-89.

782 65. Benninghoff W, Benninghoff A. 1985. Wind transport of electrostatically charged

783   particles and minute organisms in Antarctica, p 592-596, Antarctic nutrient cycles and

784   food webs. Springer.

785 66. Michaud JM, Thompson LR, Kaul D, Espinoza JL, Richter RA, Xu ZZ, Lee C, Pham

786   KM, Beall CM, Malfatti F, Azam F, Knight R, Burkart MD, Dupont CL, Prather KA.

787   2018. Taxon-specific aerosolization of bacteria and viruses in an experimental ocean-

788   atmosphere mesocosm. Nat Commun 9:2017.

67.   Aller JY, Kuznetsova MR, Jahns CJ, Kemp PF. 2005. The sea surface microlayer as a

source of viral and bacterial enrichment in marine aerosols. J Aerosol Sci 36:801-812.

68.   Wilbourn EK, Thornton DCO, Ott C, Graff J, Quinn PK, Bates TS, Betha R, Russell

LM, Behrenfeld MJ, Brooks SD. 2020. Ice nucleation by marine aerosols over the

North Atlantic Ocean in late spring. J Geophys Res Atmos 125:e2019JD030913.

69.   Amato P. 2012. Clouds provide atmospheric oases for microbes. Microbe 7:119-123.

70.   Christner BC, Morris CE, Foreman CM, Cai R, Sands DC. 2008. Ubiquity of

biological ice nucleators in snowfall. Science 319:1214.

71.   Strother SL, Salzmann U, Roberts SJ, Hodgson DA, Woodward J, Van Nieuwenhuyze

W, Verleyen E, Vyverman W, Moreton SG. 2015. Changes in Holocene climate and

the intensity of Southern Hemisphere Westerly Winds based on a high-resolution

palynological record from sub-Antarctic South Georgia. Holocene 25:263-279.

72.   Netea MG, Dominguez-Andres J, Eleveld M, Op den Camp HJM, van der Meer JWM,

Gow NAR, de Jonge MI. 2020. Immune recognition of putative alien microbial

structures: Host-pathogen interactions in the age of space travel. PLoS Pathog

16:e1008153.

73.   Nicholson WL, Schuerger AC, Race MS. 2009. Migrating microbes and planetary

protection. Trends Microbiol 17:389-92.

74.   Coenye T, Vandamme P, LiPuma JJ. 2002. Infection by *Ralstonia* species in cystic

fibrosis patients: identification of *R. pickettii* and *R. mannitolilytica* by polymerase

chain reaction. Emerg Infect Dis 8:692-6.

75.   Berliner AJ, Mochizuki T, Stedman KM. 2018. Astrovirology: Viruses at large in the

universe. Astrobiology 18:207-223.

76.   Mattoni RHT, Keller Jr EC. 1972. Induction of lysogenic bacteria in the space. The

Experiments of Biosatellite II 204:309.

77.   Mattoni RHT. 1968. Influence of spaceflight and radiation on induction of prophage

P-22 in *Salmonella typhimurium*. Jpn J Genet 43:465-465.

78.   Mattoni RHT. 1968. Space-flight effects and gamma radiation interaction on growth

and induction of lysogenic bacteria, a preliminary report. BioScience 18:602-608.

79.   Koike J. 1991. Fundamental questions concerning the contamination of other planets

with terrestrial microorganisms carried by space-probes. The Journal of Space

Technology and Science 7:9-14.

80.   Weinmaier T, Probst AJ, La Duc MT, Ciobanu D, Cheng JF, Ivanova N, Rattei T,

Vaishampayan P. 2015. A viability-linked metagenomic analysis of cleanroom

environments: eukarya, prokaryotes, and viruses. Microbiome 3:62.

81.   Mora M, Wink L, Kogler I, Mahnert A, Rettberg P, Schwendner P, Demets R, Cockell

C, Alekhova T, Klingl A, Krause R, Zolotariof A, Alexandrova A, Moissl-Eichinger

C. 2019. Space Station conditions are selective but do not alter microbial

characteristics relevant to human health. Nat Commun 10:3990.

82.   Pavletić B, Runzheimer K, Siems K, Koch S, Cortesão M, Ramos-Nascimento A,

Moeller R. 2022. Spaceflight virology: What do we know about viral threats in the

spaceflight environment? Astrobiology doi:10.1089/ast.2021.0009.

83.   Aboudan A, Colombatti G, Bettanini C, Ferri F, Lewis S, Van Hove B, Karatekin O,

Debei S. 2018. ExoMars 2016 Schiaparelli module trajectory and atmospheric profiles

reconstruction. Space Science Reviews 214:1-31.

84.   Bushnell B. 2014. BBTools software package. URL http://sourceforge

net/projects/bbmap 578:579.

85.   Joshi N, Fass J. 2011. Sickle: A sliding-window, adaptive, quality-based trimming

tool for FastQ files (Version 1.33)[Software].

86.   Nurk S, Meleshko D, Korobeynikov A, Pevzner PA. 2017. metaSPAdes: a new

versatile metagenomic assembler. Genome Res 27:824-834.

87.   Dick GJ, Andersson AF, Baker BJ, Simmons SL, Thomas BC, Yelton AP, Banfield

841   JF. 2009. Community-wide analysis of microbial genome sequence signatures.

842   Genome Biol 10:R85.

843   88.   Wu YW, Simmons BA, Singer SW. 2016. MaxBin 2.0: an automated binning

844   algorithm to recover genomes from multiple metagenomic datasets. Bioinformatics

845   32:605-7.

846   89.   Sieber CMK, Probst AJ, Sharrar A, Thomas BC, Hess M, Tringe SG, Banfield JF.

847   2018. Recovery of genomes from metagenomes via a dereplication, aggregation and

848   scoring strategy. Nat Microbiol 3:836-843.

849   90.   Bornemann TLV, Esser SP, Stach L, Burg T, Probst AJ. 2020. uBin – a manual

850   refining tool for metagenomic bins designed for educational purposes. bioRxiv

851   doi:10.1101/2020.07.15.204776

852   91.   Probst AJ, Castelle CJ, Singh A, Brown CT, Anantharaman K, Sharon I, Hug LA,

853   Burstein D, Emerson JB, Thomas BC, Banfield JF. 2017. Genomic resolution of a

854   cold subsurface aquifer community provides metabolic insights for novel microbes

855   adapted to high CO_2 concentrations. Environ Microbiol 19:459-474.

856   92.   Langmead B, Salzberg SL. 2012. Fast gapped-read alignment with Bowtie 2. Nat

857   Methods 9:357-9.

858   93.   Skennerton CT, Imelfort M, Tyson GW. 2013. Crass: identification and reconstruction

859   of CRISPR from unassembled metagenomic data. Nucleic Acids Res 41:e105.

860   94.   Mojica FJ, Diez-Villasenor C, Soria E, Juez G. 2000. Biological significance of a

861   family of regularly spaced repeats in the genomes of Archaea, Bacteria and

862   mitochondria. Mol Microbiol 36:244-6.

863   95.   Altschul SF. 2001. BLAST algorithm. e LS.

864   96.   Moller AG, Liang C. 2017. MetaCRAST: reference-guided extraction of CRISPR

865   spacers from unassembled metagenomes. PeerJ 5:e3788.

866   97.   Ren J, Ahlgren NA, Lu YY, Fuhrman JA, Sun F. 2017. VirFinder: a novel k-mer

867   based tool for identifying viral sequences from assembled metagenomic data.

868   Microbiome 5:69.

869   98.   Crits-Christoph A, Gelsinger DR, Ma B, Wierzchos J, Ravel J, Davila A, Casero MC,

870   DiRuggiero J. 2016. Functional interactions of archaea, bacteria and viruses in a

871   hypersaline endolithic community. Environ Microbiol 18:2064-77.

872   99.   Marz M, Beerenwinkel N, Drosten C, Fricke M, Frishman D, Hofacker IL, Hoffmann

873   D, Middendorf M, Rattei T, Stadler PF, Topfer A. 2014. Challenges in RNA virus

874   bioinformatics. Bioinformatics 30:1793-9.

875   100.   Rahlff J, Turzynski V, Esser SP, Monsees I, Bornemann TLV, Figueroa-Gonzalez PA,

876   Schulz F, Woyke T, Klingl A, Moraru C, Probst AJ. 2021. Lytic archaeal viruses

877   infect abundant primary producers in Earth's crust. Nat Commun 12:4642.

878   101.   Roux S, Emerson JB, Eloe-Fadrosh EA, Sullivan MB. 2017. Benchmarking viromics:

879   an in silico evaluation of metagenome-enabled estimates of viral community

880   composition and diversity. PeerJ 5:e3817.

881   102.   Lefort V, Desper R, Gascuel O. 2015. FastME 2.0: A comprehensive, accurate, and

882   fast distance-based phylogeny inference program. Mol Biol Evol 32:2798-800.

883   103.   Farris JS. 1972. Estimating phylogenetic trees from distance matrices. The American

884   Naturalist 106:645-668.

885   104.   Rambaut A. 2006. FigTree, a graphical viewer of phylogenetic trees and as a program

886   for producing publication-ready figures.

887   105.   Göker M, Garcia-Blazquez G, Voglmayr H, Telleria MT, Martin MP. 2009. Molecular

888   taxonomy of phytopathogenic fungi: a case study in *Peronospora*. PLoS One 4:e6319.

889   106.   Meier-Kolthoff JP, Hahnke RL, Petersen J, Scheuner C, Michael V, Fiebig A, Rohde

890   C, Rohde M, Fartmann B, Goodwin LA, Chertkov O, Reddy T, Pati A, Ivanova NN,

891   Markowitz V, Kyrpides NC, Woyke T, Göker M, Klenk HP. 2014. Complete genome

sequence of DSM 30083(T), the type strain (U5/41(T)) of *Escherichia coli*, and a

proposal for delineating subspecies in microbial taxonomy. Stand Genomic Sci 9:2.

107.   Bin Jang H, Bolduc B, Zablocki O, Kuhn JH, Roux S, Adriaenssens EM, Brister JR,

Kropinski AM, Krupovic M, Lavigne R, Turner D, Sullivan MB. 2019. Taxonomic

assignment of uncultivated prokaryotic virus genomes is enabled by gene-sharing

networks. Nat Biotechnol 37:632-639.

108.   Brister JR, Ako-Adjei D, Bao Y, Blinkova O. 2015. NCBI viral genomes resource.

Nucleic Acids Res 43:D571-7.

109.   Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D, Amin N,

Schwikowski B, Ideker T. 2003. Cytoscape: a software environment for integrated

models of biomolecular interaction networks. Genome Res 13:2498-504.

110.   Moraru C, Varsani A, Kropinski AM. 2020. VIRIDIC-A novel tool to calculate the

intergenomic similarities of prokaryote-infecting viruses. Viruses 12.

111.   Nilsson E, Bayfield OW, Lundin D, Antson AA, Holmfeldt K. 2020. Diversity and

host interactions among virulent and temperate Baltic Sea *Flavobacterium* phages.

Viruses 12:158.

112.   Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, Buxton S,

Cooper A, Markowitz S, Duran C, Thierer T, Ashton B, Meintjes P, Drummond A.

2012. Geneious Basic: an integrated and extendable desktop software platform for the

organization and analysis of sequence data. Bioinformatics 28:1647-9.

113.   Hyatt D, Chen GL, Locascio PF, Land ML, Larimer FW, Hauser LJ. 2010. Prodigal:

prokaryotic gene recognition and translation initiation site identification. BMC

Bioinformatics 11:119.

114.   Buchfink B, Xie C, Huson DH. 2015. Fast and sensitive protein alignment using

DIAMOND. Nature Methods 12:59-60.

115. Shaffer M, Borton MA, McGivern BB, Zayed AA, La Rosa SL, Solden LM, Liu P,

Narrowe AB, Rodriguez-Ramos J, Bolduc B, Gazitua MC, Daly RA, Smith GJ, Vik

DR, Pope PB, Sullivan MB, Roux S, Wrighton KC. 2020. DRAM for distilling

microbial metabolism to automate the curation of microbiome function. Nucleic Acids

Res 48:8883-8900.

116. Zimmermann L, Stephens A, Nam SZ, Rau D, Kubler J, Lozajic M, Gabler F, Soding

J, Lupas AN, Alva V. 2018. A completely reimplemented MPI bioinformatics toolkit

with a new HHpred server at its core. J Mol Biol 430:2237-2243.

117. Söding J, Biegert A, Lupas AN. 2005. The HHpred interactive server for protein

homology detection and structure prediction. Nucleic Acids Res 33:W244-8.

118. Paez-Espino D, Roux S, Chen IA, Palaniappan K, Ratner A, Chu K, Huntemann M,

Reddy TBK, Pons JC, Llabres M, Eloe-Fadrosh EA, Ivanova NN, Kyrpides NC. 2019.

IMG/VR v.2.0: an integrated data management and analysis system for cultivated and

environmental viral genomes. Nucleic Acids Res 47:D678-D686.

119. Sullivan MJ, Petty NK, Beatson SA. 2011. Easyfig: a genome comparison visualizer.

Bioinformatics 27:1009-10.

120. Seemann T. 2014. Prokka: rapid prokaryotic genome annotation. Bioinformatics

30:2068-9.

121. Edgar RC. 2004. MUSCLE: a multiple sequence alignment method with reduced time

and space complexity. BMC Bioinformatics 5:113.

122. Price MN, Dehal PS, Arkin AP. 2010. FastTree 2--approximately maximum-

likelihood trees for large alignments. PLoS One 5:e9490.

123. Schlitzer R. 2015. Ocean Data View. https://odv.awi.de/.