1 **Host-associated phages disperse across the extraterrestrial analogue**

2 **Antarctica**

3 **Running title: Dispersal of Antarctic phage**

6 **Janina Rahlff[a, #], Till L.V. Bornemann[a], Anna Lopatina[b],**

7 **Konstantin Severinov[b], Alexander J. Probst[a,c]**

9 [a]Group for Aquatic Microbial Ecology, University of Duisburg-Essen, Department of

10 Chemistry, Environmental Microbiology and Biotechnology (EMB), Universitätsstraße 5,

11 45141 Essen, Germany

12 [b]Institutes of Molecular Genetics and Gene Biology of the Russian Academy of Sciences,

13 Moscow, Russia

14 [c]Centre of Water and Environmental Research (ZWU), University of Duisburg-Essen,

15 Universitätsstraße 5, 45141, Essen, Germany

17 #Corresponding author: Janina Rahlff, Janina.rahlff@uol.de

18 Present address: Centre for Ecology and Evolution in Microbial Model Systems (EEMiS),

19 Department of Biology and Environmental Science, Linnaeus University, SE-39182, Kalmar,

20 Sweden

22 **Key words:** viruses, extreme environments, astrovirology, metagenomics, cryosphere,

23 *Ralstonia*

Abstract

Extreme Antarctic conditions provide one of the closest analogues of extraterrestrial

environments. Since air and snow samples especially from polar regions yield DNA amounts

in the lower picogram range, binning of prokaryotic genomes is challenging and renders

studying the dispersal of biological entities across these environments difficult. Here, we

hypothesized that dispersal of host-associated bacteriophages (adsorbed, replicating or

prophages) across the Antarctic continent can be tracked via their genetic signatures and

benefits our understanding of virus and host dispersal across long distances. Phage genome

fragments (PGFs) reconstructed from surface snow metagenomes of three Antarctic stations

were assigned to four host genomes, mainly Betaproteobacteria including *Ralstonia* spp.

Betaproteobacteria of this genus have been found in Antarctic snow as well as on space-

related equipment. We reconstructed the complete genome of a temperate phage with near-

complete alignment to a prophage in the reference genome of *Ralstonia pickettii* 12D. PGFs

from different stations were related to each other at the genus level and matched similar hosts.

Metagenomic read mapping and nucleotide polymorphism analysis revealed a wide dispersal

of highly identical PGFs, 13 of which appeared in seawater from the Western Antarctic

Peninsula with up to 5538 km to the snow sampling stations. Our results suggest that host-

associated phages, especially of *Ralstonia* sp. disperse over long distances despite harsh

conditions of the Antarctic continent. Due to the additional identification of 14 phages

associated with two *R. pickettii* draft genomes isolated from space equipment, we conclude

implications for the spread of biological contaminants in extraterrestrial settings.

Importance

Host-associated phages of the bacterium *Ralstonia* identified in snow samples can be used to

track microbial dispersal over thousands of kilometers across the Antarctic continent, which

functions as an extraterrestrial analogue because of its harsh environmental conditions. Due to

presence of this bacterial strain including genome-integrated prophages on space-related

equipment, and the here demonstrated potential for dispersal of host-associated phages, our

work has implications for Planetary Protection, a discipline in Astrobiology interested in

preventing contamination of celestial bodies with alien biomolecules or forms of life.

Introduction

Due to harsh environmental conditions and isolation of Antarctica by the surrounding

Southern Ocean's Circumpolar Current, this continent is considered an analogue for

multipurpose space exploration (1, 2). For example, its McMurdo Dry Valleys are regarded as

a close terrestrial analogue to Mars (3). Astrobiology model organisms found on Antarctica

are highly adapted to stressful conditions and comprise prokaryotes such as spore-forming

Bacilli (4, 5), but also microfungi (3, 6). Understanding endurance and dispersal of

microorganisms under conditions that mimic those on extraterrestrial planets, i.e., high UV

radiation, low temperature, and low nutrient availability, has important implications for

Planetary Protection. For instance, the dispersal of microbes that hitchhike to a celestial body

is currently not considered in Planetary Protection, a discipline in Astrobiology set out with

the aim of preventing contamination of celestial bodies with foreign biomolecules or forms of

life.

Among the potential candidates hitchhiking spacecraft are Betaproteobacteria of the genus

Ralstonia (order *Burkholderiales*). These bacteria are able to thrive under oligotrophic

conditions (7) and were reported to be ubiquitously present on space-related equipment

including water systems of the International Space Station (ISS) (8, 9), and the Mir space

station (10, 11). Likewise, they belonged to the microbial inventory of Mars Odyssey and

Mars Phoenix lander facilities and can thus prevail under strict Planetary Protection

regulations (12, 13). *Ralstonia picketii* strains were found to thrive in simulated microgravity

compared to normal gravity (11) and demonstrated high resistance against different metal ions

and UV-C radiation (8).

Ralstonia spp., (mainly *Ralstonia pickettii*), were previously found in in Antarctic soils (14),

Antarctic snow (15-17), in snow over Tibetan Plateau Glaciers (18), and in the air of the

Antarctic base Concordia (19). Interestingly, this genus has been regarded as an atmospheric

traveler rather than being part of true snow microflora in Antarctica (15). Despite a report on

bacterial activity at subzero temperatures in South pole snow (20), *Ralstonia* was not among

active bacterial communities as inferred from cDNA-based 16S rRNA amplicon sequencing

(15). This view was supported by a comprehensive Antarctic surface snow microbiome study

that did not detect this bacterial genus (21), and by the fact that spatial variability of snow

microbiomes in Antarctica is high (22). Although aerial dispersal is probably the major

contributor to (micro-)biological input to remote regions (23), the role of bioaerosol transport

to microbial ecology of isolated systems such as the Antarctic continent is poorly understood

(24). Applying high throughput sequencing approaches to study bacterial dispersal over the

Antarctic continent remains considerable challenging due to the low microbial biomass of

atmosphere-derived samples regarding their DNA content (25), and resulting issues of

recovering high quality assemblies from metagenomic reads (26).

In addition to the limited knowledge about how transport via aerosols and snow across

Antarctica shapes microbial dispersal patterns, another open question relates to the distances

that microbes can cover within the atmosphere of extreme environments. A study on the

dispersal of airborne faecal coliforms showed a distribution over about just 175 m from a

sewage outfall at Rothera Research Station (Antarctic Peninsula) and thus prolonged survival

was considered unlikely (27). More stress-resistant microorganisms could, however, endure

for much longer periods. Recently, L. A. Malard et al. (22) found high abundances of spore-

forming *Bacilli* and suggested that long-term dispersal may seed continental Antarctic snow

ecosystems. However, to date, the role of aerial dispersal in shaping patterns of microbial

biogeography is supported by little empirical evidence (23).

Here, we follow the hypothesis that *Ralstonia* spp. prophages, which are geographically

widespread entities (28), as well as replicating and adsorbed bacteriophages (all types further

referred to as "host-associated" phages) can be used to study dispersal across the Antarctic

continent. Extracting prokaryotic genomes from low DNA samples including air or

precipitation is challenging, as low input libraries (1 pg) can result in the absence of bins (26).

However, (pro)phage genomes or their fragments are much smaller than prokaryotic genomes

and thus easier to identify, track and compare. In this study, genome-resolved metagenomics

was applied to demonstrate dispersal of host-associated phage genome fragments (PGFs)

from surface snow across the Antarctic continent over hundreds to thousands of kilometers.

We detected PGFs belonging to the orders *Caudovirales* and *Tubulavirales* in this

extraterrestrial analogue and additionally show that similar genome-integrated phages are

frequent colonizers of space equipment. Therefore, we suggest that host-associated PGFs

represent a useful tool to study spatial dispersal of bacteria and their phages in extreme

environmental settings and further envision implications for the dispersal of microbiological

contaminations on spacecraft and celestial bodies that previously escaped Planetary Protection

measures.

Results

Reconstruction of MAGs from Antarctic snow metagenomic data with little evidence for

adaptive immunity

The microbial community composition of surface snow samples collected close to three

Russian Antarctic stations, Druzhnaja, Mirnii and Progress, based on 16S rRNA gene

sequencing was described earlier (16) and showed that *Ralstonia* was the most dominant

organism in snow collected at the Mirnii station, the second most dominant after

Janthinobacterium at Druzhnaja, and the third most dominant (after *Flavobacteria* and

Hydrogenophaga) around Progress station. We reconstructed four prokaryotic and one

eukaryotic metagenome assembled genomes (MAG) from the three low-biomass snow

metagenomes (Figure 1A). Two MAGs related to *Ralstonia* sp. and *Ralstonia pickettii* were

recovered from Druzhnaja and Mirnii samples and had 86%/10% and 55%/0%

completeness/contamination scores, respectively. MAGs of *Janthinobacterium lividum* with

92%/6% scores and of *Flavobacterium micromati* with 96%/0% scores were recovered from

Druzhnaja and Progress, respectively. We also identified a MAG of a diatom (likely

Thalassiosira sp.) in the Mirnii metagenome, which we used for normalizing some of our

viral analysis (see below) but was not further characterized in this manuscript. Coverage of

bacterial MAGs as performed by read mapping revealed coverage scores of 5.9, 7.1, 6.7 and

14.6 for the MAGs of *Ralstonia* sp.*, Ralstonia pickettii, Janthinobacterium lividum* and

Flavobacterium micromati, respectively.

We searched for clustered regularly interspaced short palindromic repeats (CRISPR) spacers

in the snow metagenome reads to link them to potential protospacers on PGFs. CRISPR

arrays and *cas* genes were absent from both *Ralstonia* and the *Janthinobacterium lividum*

MAGs as determined by CRISPRcasFinder. BLASTing of direct repeat (DR) sequences to

NCBI's NR database also did not indicate *Ralstonia* or *Janthinobacterium* to be the host of

the CRISPR array. Solely the genome of *Flavobacterium micromati* contained two CRISPR

arrays with four spacers each (both evidence level 3= highly likely candidates). No CRISPR

arrays were found on the *Ralstonia pickettii* draft genomes SSH4 and CW2 obtained from

space equipment.

PGF-host analyses suggest shared hosts of Antarctic phages and a newly discovered

temperate phage p12D

A total of 26 predicted PGFs, eight of them being putative PGFs, was found in Antarctic snow

metagenomic data (Table S1). VIBRANT (29) additionally detected 52 PGFs, resulting in a

total of 78 PGFs (Figure 1A). PGFs with minimum 75% of the genome covered with reads

had coverages ranging between 1.9 and 17.5. Most PGFs were partial sequences according to

viralComplete, only Antarcphage10_Mi_4716 and Antarcphage49_Dr_7823_circ were

estimated to be of full length (Table S1), although annotations of Antarcphage10_Mi_4716 in

comparison to *Ralstonia* phage p12J revealed missing genes and let us question its

completeness (Figure 2). Out of 78 PGFs, 77 were categorized as lytic by VirSorter (30) and

75 were found to be of viral or unclassified source by CheckV (Table S1). 43.6%, 46.1% and

10.3% of identified PGFs originated from the Druzhnaja, Mirnii and Progress, respectively.

All 78 PGFs were analyzed in conjunction with reconstructed MAGs using VirHostMatcher.

We defined a dissimilarity threshold of d2* value= 0.436, which corresponds to the lowest

dissimilarity value (=highest similarity) for PGFs matching the MAG of *Thalassiosira* sp.

from the snow environment (Figure 1A) by assuming that the eukaryotic MAG does not

match any of our extracted presumably prokaryote infecting PGFs. In total, 50 of the 78 PGFs

matched a host MAG below the defined dissimilarity threshold, based on their shared k-mer

patterns. Most PGFs (41) matched both MAGs of *Ralstonia* sp. and *Ralstonia pickettii* with

dissimilarity d2* values <0.436 (Figure 1A). 38 and 14 PGFs matched *Janthinobacterium*

lividum and *Flavobacterium micromati* MAGs, respectively. Five out of eight PGFs extracted

from the Progress Station metagenomic snow sequences matched the MAG of

Flavobacterium micromati recovered from this station. Seven PGF matches were shared

between all prokaryotic MAGs (Figure S1). However, with 30 shared matches, the three

MAGs belonging to the order *Burkholderiales* shared the most overlap (Figure 1B, Figure

S1).

182
Figure 1: Host-phage pairings based on shared k-mer frequency patterns. A) Heatmap representing dissimilarity value d2* for matches

between five host MAGs and 78 PGFs derived from three Antarctic snow metagenomes from the stations Progress, Mirnii, and Druzhnaja. The

table shows MAG characteristics (phylogenetic affiliation, completeness, contamination, and coverage). **B)** Network of phage-host interactions

based on k-mer frequency pattern reveals strong overlap between *Ralstonia* and *Janthinobacterium lividum* infecting PGFs. Here, only PGFs

matching the host MAG below the defined threshold (see text) are shown. n/a= not available

Among the 78 PGFs, we detected a circular and thus complete 7.8 kb PGF termed

Antarcphage49_Dr_7823_circ   (Figure   2A,   Figure   S2).   Since   the

Antarcphage49_Dr_7823_circ PGF from the Druzhnaja station was found in the chromosome

of a *Ralstonia pickettii* 12D strain isolated from copper-contaminated sediment from a lake in

Michigan with 99.88% identity (accession number: CP001644.1- 2272699-2280518, Figure

2A), we here propose the name p12D phage equivalent to *Ralstonia* p12J, a known phage

infecting the strain *Ralstonia pickettii* 12J. Antarcphage49_Dr_7823_circ (from now on

referred to as p12D) only matched the MAG of *Ralstonia* sp. and *Ralstonia pickettii* based on

shared k-mer frequency patterns. Functional protein annotations provided evidence that p12D

contained a gene for a resolvase domain containing protein/ site-specific recombinase, likely

used for integration into the host genome, as well as for the zonular occludens toxin (Zot,

PF05707) (Figure 2A, Figure S2, Table S2). The non-toxic component of Zot at the N-

terminus represents a well-conserved protein in filamentous phages and has been used for

phylogenetic analysis (31, 32). Consequent construction of a phylogenetic tree for Zot

proteins from this study and respective references (Figure 2B, Figure S3) confirmed close

identity of p12D to *Ralstonia* phage 1 NP-2014 and Antarcphage79_WAP_18.3. *Zot* of

Antarcphage10_Mi_4716 was phylogenetically related to filamentous *Ralstonia* phage p12J.

The tree shows four distinctive clusters for the Zot protein, reflecting the overall synteny of

the gene order of the different *Ralstonia* phages very well.

Annotations against UniRef100 revealed that 68.8% of all annotated Antarctic PGF proteins

remain hypothetical. The annotations taxonomically assigned 34.4% of all Antarctic PGF

proteins to either *Ralstonia* or *Ralstonia pickettii,* and 11.6 % to phages of *Janthinobacterium*

or *Janthinobacterium lividum*, potentially indicating lateral gene transfer between virus and

host in their respective evolution (33, 34) or supporting that these PGF indeed represent

prophages. Three of the Antarctic PGFs carried a site-specific integrase or resolvase domain,

and 4.2% of genes were related to phage structural proteins, e.g., head, tail, or capsid proteins

(Table S3, Table S4).

Figure 2: Phage genome comparisons, functional annotations and phylogenetic

relationship based on zonular occludes toxin (Zot, Pfam-ID: PF05707). A) Synteny of

known *Ralstonia* sp. phages from NCBI and Antarctic PGFs from this study with similar

coding sequences (CDS) being in the same color if they occur on more than one phage

genome. Functional annotations performed by DRAM-v (35) are given for all colored CDS

where available. Vertical lines between sequences indicate regions of shared similarity shaded

according to tBLASTx (orange gradient for matches in the same direction or blue gradient for

inverted matches). Figure was created using Easyfig (36). **B)** The phylogenetic tree was built

using FastTree 2.1.11 (37) in Geneious 11.1.5 (38) under default settings and shows four

distinct clusters of Zot proteins based on their amino acid sequences (aligned with MUSCLE

(39)).

Cross-mapping and nucleotide variations reveal dispersal patterns of PGFs across

Antarctica

Cross-mapping on the 78 PGFs with reads from the three snow samples and the WAP

seawater sample (Figure 3A) demonstrated that PGFs derived from the Progress station could

not be found in the other two snow metagenomes. By contrast, 16 different PGFs from

Druzhnaja were present at Mirnii, and 27 Mirnii PGFs were present at Druzhnaja (Figure 3B).

In addition, 4 and 9 PGFs from Mirnii and Druzhnaja were detected in the WAP dataset,

respectively (Figure 3B), all of them having minimum 97% of their lengths covered with

reads from the snow metagenomes (Table S1). Of 43 PGFs that were found at both stations

(Mirnii and Druzhnaja) based on read mapping, 26 PGFs shared a viral cluster (VC), and 34

matched prokaryotic hosts based on k-mer frequencies between stations (Table S1).

Interestingly, nine of the 13 PGFs occurring in WAP and Mirnii/Druzhnaja samples,

contained identical single nucleotide polymorphisms (Table S5) or were even devoid of them

pointing towards a common phage population before dispersal led to separation.

Only one of the DR sequences matched back to the MAG assigned *Flavobacterium micromati*

from Progress station (Table S6). Matching of CRISPR spacers derived from CRISPR loci of

unknown hosts (reconstructed from DR sequences) revealed that spacers from Mirnii matched

19, 14 and three PGFs from Druzhnaja, Mirnii and Progress, respectively, and one

Flavobacterium sp. spacer from Progress matched a PGF from that station (Table S6, Figure

S4). Out of these 37 spacer matches, ten matched protospacers of PGFs of unknown hosts and

the remaining 27 were associated with PGFs (including p12D) that infect one or more of the

four recovered prokaryotic MAGs according to VirHostMatcher. Since p12D was among the

PGF spacer targets and is a certain *Ralstonia* phage, this could indicate that the unknown host

from which the CRISPR spacer were derived also belongs to the order *Burkholderiales* and

uses adaptive immunity against its viruses.

Clustering of PGFs in VICTOR and vConTACT2 revealed that groups of two to four PGFs

could be assigned to twelve distinct viral genera-forming viral clusters (Figure S5 & S6,

Table S1, Figure 3A). Members of the same cluster were often recovered from different

stations, for example, Druzhnaja and Mirnii, which are located 923 km apart (Figure 3B). The

phylogenomic Genome-BLAST Distance Phylogeny (GBDP) tree yielded average support of

47% and 80% in the nucleic acid and amino acid-based analysis, respectively. OPTSIL

clustering yielded 78 and 64 species and genus clusters at the nucleic acid level and 78 and 70

at the amino acid level, respectively.

Figure 3: Dispersal distances of Antarctic PGFs. A) The map shows different stations and

PGF whose presence was determined by cross mapping (CM) of reads and shared between

two locations. Dispersal is only shown if at least one PGF occurred with equal or more than

75% of scaffold length of at least 1-fold coverage at two stations. Number of virus clusters

(VC) and virus-host cross matches (VHM) was determined for assembled PGFs from snow

metagenomes (Druzhnaja, Mirnii, Progress). A VHM indicates that one PGF infects host

MAGs from two stations. No VC and only one CM were observed for Progress-Druzhnaja

and Progress-Mirnii. The Western Antarctic Peninsula (WAP) sample is derived from

seawater (zigzag/wave symbol), the remaining samples are from snow (snowflake symbol).

Map was built using Ocean Data View (40). **B)** Pictures of two sampling sites (photos: Anna

Lopatina). **C)** Heatmap of read-normalized coverages based on CM of reads against 78

Antarctic PGFs and separation distances between stations. White areas indicate PGFs that

were not present because they had less than 75% of scaffold length with least 1-fold coverage.

Many PGFs formed VCs with no genomic relatedness to any known phages from the

ProkaryoticViralRefSeq v94 database. Others were, however, like known *Ralstonia* PGFs

(current family *Inoviridae,* order *Tubulavirales*) and hence clustered with those, such as p12D

with *Ralstonia* phage 1 NP-2014, or Antarcphage10_Mi_4716 with *Ralstonia* phage PE226

and *Ralstonia* phage p12J (Figure 4, Table S2). Other Antarctic PGFs shared protein clusters

with *Ralstonia* phages from space equipment based on vCONTACT2 or were related to

phages of the *Caudovirales* order from the RefSeq database (Figure 4).

Figure 4: Phage network of Antarctic PGFs derived from Druzhnaja, Mirnii, Western Antarctic Peninsula (WAP) and space equipment (ISS/MarsOrb) clustered with viruses of the viral RefSeq database. Based on shared protein clusters, Antarctic PGFs from this study group with known *Ralstonia* phage of the family *Inoviridae* (A), with phages obtained from *Ralstonia* isolates from space equipment (B) or with known phages of the order *Caudovirales* (C). Black nodes refer prokaryotic viruses other than *Ralstonia* and PGFs from this study. Interactions show relatedness of genomes on viral genus or higher taxonomic level. "Others" refer to other known viral families not listed in the legend. Visualization was done using Cytoscape 3.8.2. (41). For details about viral clusters, please see Table S1.

In summary, our results support the idea of dispersal of host-associated phages between widely separated stations because a) cross-mapping revealed presence of Mirnii PGFs at Druzhnaja and vice versa, and presence of a temperate phage from snow in the WAP seawater dataset, b) PGFs in widely separated locations carry identical single nucleotide polymorphisms or lack nucleotide variations, c) some PGFs from Mirnii and Druzhnaja belong to the same genus/VC, and d) Mirnii and Druzhnaja PGFs share host MAGs according

to their k-mer frequency patterns. This was mainly true for PGFs affiliated to the host MAGs

Ralstonia and *Janthinobacterium lividum* but rarely for PGFs of *Flavobacterium micromati,*

which could indicate that certain bacterial species or their phage in Antarctic snow are more

prone to atmospheric dispersal than others. And e) we found CRISPR spacers from two

unidentified hosts (represented by two distinct CRISPR DR sequences) at the Mirnii station to

match PGFs from Druzhnaja, Mirnii and Progress (Figure S4). This is particularly interesting

because cross-mapping revealed little to no abundance of Progress PGFs at other stations,

which could be related to the low biomass and/or little coverage of PGFs, but spacers from

Mirnii suggest an infection history with PGF from that station, thus reflecting (past) dispersal

of host and/or phage.

Ralstonia phages occur across diverse ecosystems and on space equipment

By BLASTing all recovered PGFs against the IMG/VR viral database, we found that p12D

shared high identity (85.9 %) with a *Ralstonia pickettii* prophage (IMG/VR v.2 scaffold ID

Ga0075447_10000781, genome ID: 3300006191) from a seawater metagenome of the WAP

(Figure 2, Table S7), which was further validated by cross-mapping of WAP reads to p12D

delivering 100% scaffold coverage with 90% identical reads. The WAP has 4710 km to the

station Druzhnaja (Figure 3). The PGF Antarcphage79_WAP_18.3 was obtained by

BLASTing the p12D scaffold against the WAP assembly but was assembled containing

genomic regions extending the actual PGF. Based on vConTACT2, it was affiliated to the

same viral genus as p12D, fostering the genomic relatedness between these two phages. Other

PGFs recovered from the Antarctic snow metagenome showed hits to phages deposited at

IMG/VR. Hits were related to phages originating from freshwater, wastewater, groundwater,

or phages that were associated with plant root microbial communities, which is in accordance

with the fact that *Ralstonia* sp. is a frequent phytopathogen (42). Many of the matching

entries in IMG/VR were identified as phages of *Ralstonia* due to matching CRISPR spacers of

this host (mostly *R. solanacearum,* Table S7).

Altogether, 14 PGFs were found on space equipment, hereafter referred to as ISSphage (7)

and MarsOrbphage (7) for PGFs extracted from the *Ralstonia pickettii* strains CW2 and SSH4

draft genomes from the ISS cooling system and Mars Odyssey Orbiter, respectively. Seven of

these PGFs were identified as lysogenic and according to protein annotations (Table S3 & S4)

mostly use proteins of the integrase family for integration into the host chromosome.

VCONTACT revealed that some of the ISSphage4 formed a common genus cluster with 4

Antarctic PGFs, MarsOrbphage5 shared protein clusters with ISSphage3, and MarsOrbphage7

formed a genus cluster with several *Burkholderia* phages among other *Caudovirales* members

(Table S1, Figure 4).

Discussion

Some Antarctic PGFs are temperate phages of the order *Tubulavirales*

CRISPR-Cas systems equip bacteria and archaea with a powerful defense system against

invading mobile genetic elements including plasmids, phages and viruses. Only ca. 50% of

bacteria rely on this adaptive immune system, and only 31% of genomes from public

databases belonging to plant-pathogenic *Ralstonia solanacearum* contain CRISPR-Cas arrays

(43). Upon exposure to a virulent phage under laboratory conditions, the CRISPR array of

Ralstonia solanacearum strain CFBP2957 did not acquire new spacers from viral

protospacers. This demonstrates the CRISPR system's inefficiency in viral eradication and

suggests alternative antiviral mechanisms in this *Ralstonia* strain (43). More recently, a

publication reported that in the presence of CRISPR arrays, 27.9% of spacers from *Ralstonia*

genomes targeted prophage elements (28). Overall, our study demonstrated the lack of

CRISPR systems in *Ralstonia* from a low temperature environment, and further underpins

that this genus might employ other strategies for defense against mobile genetic elements

(43). J. L. Weissman et al. (44) suggested that psychrophilic and psychrotolerant microbes

would disfavor CRISPR systems more than mesophilic species, due to lower viral decay rates,

higher viral diversity, and greater dispersal in colder compared to warmer environments. In

addition, frequent targeting of temperate phages by CRISPR spacers can be another reason for

the loss of CRISPR systems (45). We assume that not all sequenced PGFs are prophages but

could also have been sequenced during their adsorption or replicating stage in the lytic cycle.

Based on shared protein clusters, some Antarctic PGFs such as Antarcphage10_Mi_4716,

Antarcphage48_Dr_1123, Antarcphage17_Mi_3026, and p12D clustered with known

Ralstonia phages including *Ralstonia* phage p12J (NC_005131.2), *Ralstonia* phage PE226

(NC_015297.1) or *Ralstonia* phage 1 NP-2014 (NC_023586.1). These are filamentous phages

of the *Inoviridae*, a family that has been recently called for re-classification to higher

taxonomic ranks (32, 46). Inoviruses (order *Tubulavirales*) typically feature circular, single

stranded DNA genomes of ~5-15 kb length, partake in chronic infection cycles and are

globally abundant (32). As we sequenced dsDNA viruses, detection of ssDNA inoviruses

shows that these must be either replicating phages or genome-integrated prophages. *Zot*,

which was detected on p12D, Antarcphage10_Mi_4716 and Antarcphage79_WAP_18.3 is a

typical gene in filamentous *Ralstonia* phages (47, 48). Filamentous phages confer little burden

to their hosts or even serve them, e.g., by increasing their hosts' virulence and evolutionary

fitness, and because virions can leave the host in a non-destructible way (31, 32). *Ralstonia*

phages detected in Arctic viromes were shown to transduce genomic information of cold-

shock proteins to their hosts (49), which would be an asset for microorganisms in polar

environments. Although we did not find cold shock genes on PGFs in this study, transduction

of beneficial genes, which is common in extreme environments (50), could theoretically be

another reason why *Ralstonia* hosts do not apply CRISPR defenses against their phages.

Some *Ralstonia* phages occur as non-integrative, episomal forms, e.g., RS603, a hybrid of

RSM1/3 infecting the phytopathogen *Ralstonia solanacearum* (51)*,* whose genome lacks a

resolvase domain (Figure 2A), but many mesophilic *Ralstonia* also occur as lysogens (52).

Lysogeny, a lifestyle during which the phage genome becomes integrated into the host

chromosome, is a widespread phenomenon in low temperature environments (53-55) and

likely attributed to prolonged starvation and low activity of host cells under harsh conditions.

Low activity was previously also reported for *Ralstonia* (15). Since p12D and its counterpart

from the Western Antarctic Peninsula (WAP) have a resolvase-domain containing protein

functioning in integration/excision during lysogenisation (56, 57), we conclude that they must

be temperate phages of *Ralstonia*.

PGFs travel long distances across the Antarctic continent

Many, i.e., 75 of the 78 Antarctic PGFs found in this study shared little protein clusters with

known phages from public databases. This missing relatedness was previously reported for

ssDNA viruses originating from Antarctic cryoconite holes and attributed to the isolation and

extreme environmental conditions at the Antarctic continent (58). In fact, 28 of the 78 PGFs

did not match any host MAG below a 0.436 threshold (Table S1), meaning that they were

either false-positives PGFs or PGFs of non-identified host MAGs (ten of them had at least

one spacer match).

Aeolian transport of viruses over polar environments, especially attached to snowflakes, has

been scarcely investigated to date. Former investigations, mainly conducted at low latitudes,

demonstrated intercontinental transport of microbes by winds (59), and that highly identical

phages can be found in distantly related areas and in various ecosystems around the globe

(60-63). C. M. Bellas et al. (64) recently reported the presence of near-identical phage

genomes being spread by up to 4000 km in cryoconite holes of Svalbard, Greenland, and the

Alps. Our data show that despite the isolation of the Antarctic continent, and under no

consideration of anthropogenic dispersal (65), bacteria and phage distribution via snow over

extensive distances across Antarctica is possible. We cannot be certain about the nature of the

PGFs (prophage or lytic phage) by metagenomic predictions alone. While the *Inoviridae*

fraction of the dataset is likely to occur as prophages or episomal forms, frequent CRISPR

spacer matches against PGFs and PGFs assigned to lytic categories by VIBRANT and

VirSorter (category 1-3) support that host-associated, lytic phages during adsorption stage

were present as well. We thus commend further experiments that involve cultivation attempts

and sequencing of metaviromes (free phages) to reveal the extent of lysogenic or chronic

compared to lytic infection styles in Antarctic snow.

Snow PGFs were additionally found and carried identical single nucleotide polymorphisms in

the seawater metagenome from the WAP being 4710 km or even 5338 km apart from

Druzhnaja and Mirnii, respectively (Figure 3B), implying long distance transport. From

seawater, a transmission route via sea spray aerosols to snow that is blown over ice surfaces

(66) can be assumed. Aerosolization of bacteria from the sea surface is highly taxon-specific

but seems to work well for *Ralstonia pickettii* (67) and also viruses (68). Alternatively, a

transport via aerosols to clouds and precipitating snow is possible. In the latter scenario,

bioaerosols including microbial cells would act as ice nucleation particles (69), transferring

microbes to ice clouds where they might induce their own precipitation (70, 71). The

abundance results from this study support isolation of Progress-derived PGFs and point

towards a decreasing gradient of abundances from Druzhnaja to Mirnii to the WAP for many

PGFs (Table S1). Thus, general dispersal patterns of microbes across Antarctica seem

governed by westward drift and are probably mediated by the prevailing Southern

Hemisphere westerly winds (72). On short spatial-temporal scales, dispersal seems more

complex and probably shaped by multifactorial dependencies such as the different potential of

a species to become airborne (67), meteorological conditions or the local geography. For

instance, while Druzhnaja is located ~50 km into the continent, the stations Mirnii and

Progress are near the coast and more exposed to the sea in summer when ice breaks occur.

Implications for dispersal of viruses and microbes in extraterrestrial settings

We assume that the transferability of using PGFs to study microbial dispersal in space

analogues such as the Antarctic continent is likely applicable to other celestial bodies like

Mars. Three celestial bodies in our solar system (Mars, Europa, and Enceladus) have

environmental conditions that could favor microbial life (reviewed by M. G. Netea et al.

(73)). Most microbial isolates (85-95%) obtained from spacecraft and assembly facilities are

associated with humans and consequently unmanned spacecraft contain fewer and less diverse

microbes than their manned counterparts (74). Since *Ralstonia* spp. including *Ralstonia*

pickettii can be human pathogens (75) and are able to thrive under harsh and oligotrophic

conditions (7), they might contribute to the transmission of viruses to extraterrestrial

environments, particularly via manned missions.

We found evidence for viral signatures associated with two *Ralstonia pickettii* strain draft

genomes previously obtained from space equipment of a spacecraft assembly clean room and

from water systems of the ISS. This result in conjunction with the result that temperate

Ralstonia phages (and other viruses) can undergo long-range dispersal in association with

their hosts across the extraterrestrial analogue Antarctica suggests that contaminations of

space equipment with particularly persistent microbes such as *Ralstonia* should receive more

focus during microbiological monitoring in the framework of Planetary Protection. However,

the field of astrovirology has so far generally found little attention (76). The contribution of

lysogenic and episomal phage ('hidden hitchhikers') to overall viral loads on spacecraft and

associated equipment has been overlooked despite early work reporting on alterations in

prophage induction during spaceflight (77-79), tobacco mosaic virus to survive space flight

equivalent proton irradiation (80), and the occurrence of phages and human-related

circoviruses in clean rooms (81).

Planetary Protection aims preventing the spread of biological contaminants (forward

contamination) to space shuttles and stations as well as extraterrestrial environments of the

solar system. However, this policy largely ignores the potential of escaped biological

contaminants to heavily disperse on foreign celestial bodies once being released, for instance

after crash landings as happened for the Schiaparelli module of the ExoMars program in 2016

(82). Our results show that host-associated PGFs are not only suitable indicators for tracking

long-distance dispersal in space analogues but also demonstrate that the release of

contaminants that previously escaped Planetary Protection measures could spread far across

extraterrestrial ecosystems and, in the worst-case scenario, confound future life detection

missions.

Material and Methods

Metagenomic and genomic data processing

We analyzed publicly available metagenomic data sets, which correspond to Antarctic surface

snow collected around three Russian stations (Druzhnaja, Mirnii, Progress) and are deposited

at NCBI's Sequence Read Archive (SRA) as Bioproject PRJNA674475 (Table S8) and MG-

RAST under project accession mgp13052 including taxon abundance data. DNA extraction

for these metagenomes was carried out with DNA Blood & Tissue kit (Qiagen, Hilden,

Germany), whose protocol causes removal of free viruses. Sequencing libraries were prepared

with MiSeq reagent kit v.2 (Illumina, USA), targeting mainly dsDNA viruses. Metagenomic

data of a seawater sample from the Western Antarctic Peninsula (WAP) was also obtained

from SRA (accession number SRR5591034). Raw shotgun sequencing reads of the three

snow   metagenomes   and   the   WAP   dataset   were   quality-trimmed   using   BBDuk

(https://github.com/BioInfoTools/BBMap/blob/master/sh/bbduk.sh)   from   the   BBTools

package (83) and Sickle (84) resulting in read counts between 1.31 – 2.65 Mio. for the three

snow samples. Assembly of reads was done using MetaSPAdes version 3.13 (85). Scaffolds

<1 kbps in length were removed. Draft genome sequences of *Ralstonia pickettii* strains SSH4

and CW2 isolated from space equipment were obtained from Genbank accession numbers

JFZG00000000 and JFZH00000000 (9). Strains SSH4 and CW2 were isolated pre-flight from

the surface of the Mars Odyssey Orbiter during assembly and from a water sample taken in-

flight from the ISS cooling system, respectively.

Reconstruction of microbial genomes from metagenomes and CRISPR prediction

Binning of metagenome-assembled genomes (MAGs) from the three stations Druzhnaja,

Mirnii and Progress was performed using Emergent self-organizing maps (ESOMs, (86)),

ABAWACA (https://github.com/CK7/abawaca) and MaxBin 2.0 (87). Cross-mapping for use

of MaxBin 2.0 was conducted using Bowtie2 in --sensitive mode (88). Aggregation of bins

was performed using DASTool (89) and bins were curated using GC base content, coverage

and taxonomy in uBin (90), which also uses an universal marker gene set for bacteria and

archaea to estimate genome quality (contamination and completeness) (91). Read coverage of

recovered MAGs was obtained from read mapping using Bowtie2 in sensitive mode to the

individual bins, followed by mismatch filtering (2% mismatch allowance, depending on read

length).

Clustered regularly interspaced short palindromic repeats (CRISPR) arrays, which represent

the prokaryotic adaptive immunity, were searched in host MAGs using CRISPRcasFinder and

only those with evidence level 3 or 4 were considered (92). CRISPR loci consist of repeats

interspaced with short DNA sequences (spacers) obtained from invading mobile genetic

elements such as phages and thus provide a record of past infection events. Since CRISPR

arrays might get lost during the binning processes, e.g., due to fragmentation in assembly of

strain variants, absence of CRISPR arrays in CRISPRcasFinder was further investigated by

reconstructing CRISPR systems from raw reads using Crass (93) and BLASTing the obtained

direct repeat (DR) sequences, which are phylogenetically well-conserved (94), against the

NCBI non-redundant database (release 1st March 2021) using a BLASTn --short algorithm

(95) with subsequent filtering at 80% similarity and e-value threshold of 10e-05 for hits with

potential hosts. DR sequences were also BLASTed against the MAGs and were used to

extract CRISPR spacers from reads using MetaCRAST with settings -d 3 -l 60 -c 0.99 -a 0.99

-r (96), which were in turn matched against PGFs applying the same BLASTn and filtering

step as mentioned above. In most cases, BLASTing the DR sequences against the NR

database did not reveal the host's identity. Nevertheless, spacers derived from unknown hosts

were considered as well, as their matches to PGFs increased our confidence that these

represent true mobile genetic elements, and matches can be used to infer infection patterns

between stations.

PGF detection, host allocation and viral clustering

Phage genome fragments (PGFs) were identified from metagenome assemblies using a

combination of bioinformatic tools and the classification of predicted viral genome

(fragments) as "putative viruses" and "viruses" was done as previously described (97).

VIBRANT v.1.2.1 with default settings was used to find additional PGFs (29). We did not set

the usual cut-off length of 10 kb for PGFs (98), since many known *Ralstonia* phages and

inoviruses have genome sizes <10 kb (32, 52), and because the low biomass Antarctic snow

samples were overall very limited in PGFs compared to other ecosystems. VirSorter v1 (30)

and VIBRANT aided in (pro)phage detection in draft genomes of *R. pickettii* from space

equipment, and VirSorter contigs were taken for downstream analyses. CheckV (99) was used

to determine the type of identified PGF, and viralComplete (100) applied to predict

completeness and closely related phage. Pairwise comparisons of Antarctic PGFs and

clustering was done using nucleic acid- and amino acid-based VICTOR (101). The resulting

intergenomic distances were used to infer a balanced minimum evolution tree with branch

support via FASTME including Subtree Pruning and Regrafting (SPR) postprocessing (102)

using the distance formula D0. Branch support was inferred from 100 pseudo-bootstrap

replicates each. Trees were rooted at the midpoint (103) and visualized with FigTree v1.4.4

(104). Taxon boundaries at the species and genus level were estimated with the OPTSIL

program (105), using the recommended clustering thresholds (101) and an F value (fraction of

links required for cluster fusion) of 0.5 (106).

Clustering of PGFs from Antarctic snow and space equipment was further substantiated via

vConTACT2 v.0.9.19 (107, 108) in combination with the ProkaryoticViralRefSeq database

(v94, (109)) followed by visualization of viral clusters (VC) in Cytoscape v. 3.8.2 (41). Virus-

host matches were determined using the tool VirHostMatcher (110) with d_{2*} oligonucleotide

frequency dissimilarity measures for a k-mer length of six as these settings should result in

highest host prediction accuracy (110). Viral genus clusters were further determined using

VIRIDIC (111). Venn diagrams for overlapping PGF-host matches between MAGs were

calculated using the VIB-ugent webtool (http://bioinformatics.psb.ugent.be/webtools/Venn/).

Mapping of reads to assembled PGFs and nucleotide polymorphism analysis

We assume that presence of a PGF at two locations confirmed by read mapping represents

dispersal. To determine if an assembled PGF from a single sample occurred in other

metagenomes even if not being assembled, we performed read mapping (Bowtie2) following

the previously published guidelines: that reads should map to a PGF with at least 90% identity

and more than 75% of scaffold should have a coverage of at least 1x (98). Bowtie2 (88)

settings were: --ignore-quals –mp = 1,1 –np = 1 –rdg = 0,1 –rfg = 0,1 --score-min = L,0,-0.1

as previously suggested (112). To detect the coverage of a PGF by reads after mapping we

used an inhouse script *Calcopo* (https://github.com/ProbstLab/viromics/blob/master/calpoco)

to calculate the coverage per nucleotide and then the percentage of positions in the genome

covered.   Mean   coverage   of   PGFs   was   calculated   using

calc_coverage_v3 (https://github.com/ProbstLab/uBin-

helperscripts/blob/master/bin/04_01calc_coverage_v3.rb) and normalized to sequencing depth

to compare between samples. Analysis of nucleotide polymorphisms was conducted for the

13 PGFs that underwent long range dispersal, i.e., PGF being present in the WAP sample and

at least one of the snow metagenomes according to read mapping (Table S1). Variation

analysis was performed in Geneious 11.1.5 (38) by applying default settings to the read-

mappings generated as explained above.

Gene prediction and annotations

Open reading frames on PGFs were detected using Prodigal in meta mode (113). Functional

and taxonomic annotations of predicted proteins of PGFs were performed by DIAMOND

searches with a e-value of 10e-05 (114) against FunTaxDB (90) and by using DRAM-v (35).

For the full-length genome of the PGF p12D, annotations were improved using HHpred

against PDB, Pfam, UniProt-SwissProt-viral and NCBI_Conserved Domains ((115, 116),

https://toolkit.tuebingen.mpg.de/tools/hhpred) with a probability threshold of 70%. Sequences

of PGFs were BLASTed against IMG/VR 2.0 (117) with an e-value cut-off of 10e-05 to find

related phages from other metagenomic datasets.

Synteny of *Ralstonia* phage and phylogenetic comparison of the *zonular occludens toxin*

gene

586   Synteny of known *Ralstonia* sp. phages from NCBI and all Antarctic PGFs that were

587   identified herein and carried the zonular occludens toxin (*zot*, Pfam-ID: PF05707) was

588   performed with tBLASTx comparisons using Easyfig v.2.2.5 (36) on .gbk files generated by

589   Prokka (118). A phylogenetic tree for the MUSCLE-aligned (39) amino acid and nucleic acid

590   sequences of *zot* was constructed using the FastTree (37) algorithm in Geneious 11.1.5 (38).

Acknowledgements

We acknowledge Ken Dreger for server administration and maintenance as well as Cristina

Moraru for sharing insights on virus taxonomy.

Author Contribution statement

J.R. designed the study, wrote the manuscript, and carried out the analyses with input from

T.L.V.B.; A.L. and K.S. generated the raw data and performed the sampling; A.J.P

conceptualized the project, provided supervision, resources, and was involved in data

interpretation; All authors edited drafts of the manuscript.

Author Disclosure Statement

For all authors, no competing financial interests exist.

Funding statement

We acknowledge funding from the German Aerospace Center (DLR) for the project

DISPERS (50WB1922). J.R. was partially supported by the German Science Foundation for

the project VIBOCAT (grant number DFG RA3432/1-1). A.J.P. and T.L.V.B. were supported

by the Ministerium für Kultur und Wissenschaft des Landes Nordrhein-Westfalen

("Nachwuchsgruppe Dr. Alexander Probst").

References:

1.   Pyne SJ. 2007. The extraterrestrial Earth: Antarctica as analogue for space

614   exploration. Space Policy 23:147-149.

2.   Lugg D, Shepanek M. 1999. Space analogue studies in Antarctica. Acta Astronaut

44:693-9.

3.   Onofri S, Selbmann L, Zucconi L, Pagano S. 2004. Antarctic microfungi as models for

exobiology. Planet Space Sci 52:229-237.

4.   Puskeppeleit M, Quintern LE, El Naggar S, Schott JU, Eschweiler U, Horneck G,

Bucker H. 1992. Long-term dosimetry of solar UV radiation in Antarctica with spores

of *Bacillus subtilis*. Appl Environ Microbiol 58:2355-9.

5.   Nicholson WL, Munakata N, Horneck G, Melosh HJ, Setlow P. 2000. Resistance of

Bacillus endospores to extreme terrestrial and extraterrestrial environments. Microbiol

Mol Biol Rev 64:548-72.

6.   Onofri S, Barreca D, Selbmann L, Isola D, Rabbow E, Horneck G, De Vera J, Hatton

J, Zucconi L. 2008. Resistance of Antarctic black fungi and cryptoendolithic

communities to simulated space and Martian conditions. Stud Mycol 61:99-109.

7.   McAlister MB, Kulakov LA, O'Hanlon JF, Larkin MJ, Ogden KL. 2002. Survival and

nutritional requirements of three bacteria isolated from ultrapure water. J Ind

Microbiol Biotechnol 29:75-82.

8.   Mijnendonckx K, Provoost A, Ott CM, Venkateswaran K, Mahillon J, Leys N, Van

Houdt R. 2013. Characterization of the survival ability of *Cupriavidus metallidurans*

and *Ralstonia pickettii* from space-related environments. Microb Ecol 65:347-60.

9.   Monsieurs P, Mijnendonckx K, Provoost A, Venkateswaran K, Ott CM, Leys N, Van

Houdt R. 2014. Draft genome sequences of *Ralstonia pickettii* strains SSH4 and CW2,

isolated from space equipment. Genome Announc 2.

10.   Ott CM, Bruce RJ, Pierson DL. 2004. Microbial characterization of free floating

condensate aboard the Mir space station. Microb Ecol 47:133-6.

11.   Baker PW, Leff L. 2004. The effect of simulated microgravity on bacteria from the

Mir space station. Microgravity Sci Technol 15:35-41.

12.   Vaishampayan P, Osman S, Andersen G, Venkateswaran K. 2010. High-density 16S

642   microarray and clone library-based microbial community composition of the Phoenix

643   spacecraft assembly clean room. Astrobiology 10:499-508.

644   13.   La Duc MT, Nicholson W, Kern R, Venkateswaran K. 2003. Microbial

645   characterization of the Mars Odyssey spacecraft and its encapsulation facility. Environ

646   Microbiol 5:977-85.

647   14.   Lysak V, Maksimova IA, Nikitin DA, Ivanova AE, Kudinova AG, Soina VS,

648   Marfenina OE. 2018. Soil microbial communities of Eastern Antarctica. Moscow

649   Univ Biol Sci Bull 73:104-112.

650   15.   Lopatina A, Krylenkov V, Severinov K. 2013. Activity and bacterial diversity of snow

651   around Russian Antarctic stations. Res Microbiol 164:949-58.

652   16.   Lopatina A, Medvedeva S, Shmakov S, Logacheva MD, Krylenkov V, Severinov K.

653   2016. Metagenomic analysis of bacterial communities of Antarctic surface snow.

654   Front Microbiol 7:398.

655   17.   Antony R, Mahalinganathan K, Krishnan KP, Thamban M. 2012. Microbial

656   preference for different size classes of organic carbon: a study from Antarctic snow.

657   Environ Monit Assess 184:5929-43.

658   18.   Liu Y, Yao T, Jiao N, Kang S, Xu B, Zeng Y, Huang S, Liu X. 2009. Bacterial

659   diversity in the snow over Tibetan Plateau Glaciers. Extremophiles 13:411-23.

660   19.   Van Houdt R, De Boever P, Coninx I, Le Calvez C, Dicasillati R, Mahillon J,

661   Mergeay M, Leys N. 2009. Evaluation of the airborne bacterial population in the

662   periodically confined Antarctic base Concordia. Microb Ecol 57:640-8.

663   20.   Carpenter EJ, Lin S, Capone DG. 2000. Bacterial activity in South Pole snow. Appl

664   Environ Microbiol 66:4514-7.

665   21.   Michaud L, Lo Giudice A, Mysara M, Monsieurs P, Raffa C, Leys N, Amalfitano S,

666   Van Houdt R. 2014. Snow surface microbiome on the High Antarctic Plateau (DOME

667   C). PLoS One 9:e104505.

668   22.   Malard LA, Sabacka M, Magiopoulos I, Mowlem M, Hodson A, Tranter M, Siegert

669   MJ, Pearce DA. 2019. Spatial variability of Antarctic surface snow bacterial

670   communities. Front Microbiol 10:461.

671   23.   Pearce DA, Alekhina IA, Terauds A, Wilmotte A, Quesada A, Edwards A,

672   Dommergue A, Sattler B, Adams BJ, Magalhaes C, Chu WL, Lau MC, Cary C, Smith

673   DJ, Wall DH, Eguren G, Matcher G, Bradley JA, de Vera JP, Elster J, Hughes KA,

674   Cuthbertson L, Benning LG, Gunde-Cimerman N, Convey P, Hong SG, Pointing SB,

675   Pellizari VH, Vincent WF. 2016. Aerobiology over Antarctica - A new initiative for

676   atmospheric ecology. Front Microbiol 7:16.

677   24.   Bottos EM, Woo AC, Zawar-Reza P, Pointing SB, Cary SC. 2014. Airborne bacterial

678   populations above desert soils of the McMurdo Dry Valleys, Antarctica. Microb Ecol

679   67:120-8.

680   25.   Behzad H, Gojobori T, Mineta K. 2015. Challenges and opportunities of airborne

681   metagenomics. Genome Biol Evol 7:1216-26.

682   26.   Bowers RM, Clum A, Tice H, Lim J, Singh K, Ciobanu D, Ngan CY, Cheng JF,

683   Tringe SG, Woyke T. 2015. Impact of library preparation protocols and template

684   quantity on the metagenomic reconstruction of a mock microbial community. BMC

685   Genomics 16:856.

686   27.   Hughes KA. 2003. Aerial dispersal and survival of sewage-derived faecal coliforms in

687   Antarctica. Atmos Environ 37:3147-3155.

688   28.   Gonçalves OS, de Oliveira Souza F, Bruckner FP, Santana MF, Alfenas-Zerbini P.

689   2021. Widespread distribution of prophages signaling the potential for adaptability

and pathogenicity evolution of *Ralstonia solanacearum* species complex. Genomics

113:992-1000.

29.   Kieft K, Zhou Z, Anantharaman K. 2020. VIBRANT: automated recovery, annotation

and curation of microbial viruses, and evaluation of viral community function from

genomic sequences. Microbiome 8:90.

30.   Roux S, Enault F, Hurwitz BL, Sullivan MB. 2015. VirSorter: mining viral signal

from microbial genomic data. PeerJ 3:e985.

31.   Hay ID, Lithgow T. 2019. Filamentous phages: masters of a microbial sharing

economy. EMBO Rep 20.

32.   Roux S, Krupovic M, Daly RA, Borges AL, Nayfach S, Schulz F, Sharrar A, Matheus

Carnevali PB, Cheng JF, Ivanova NN, Bondy-Denomy J, Wrighton KC, Woyke T,

Visel A, Kyrpides NC, Eloe-Fadrosh EA. 2019. Cryptic inoviruses revealed as

pervasive in bacteria and archaea across Earth's biomes. Nat Microbiol 4:1895-1906.

33.   Filée J, Forterre P, Laurent J. 2003. The role played by viruses in the evolution of their

hosts: a view based on informational protein phylogenies. Res Microbiol 154:237-243.

34.   Moreira D. 2000. Multiple independent horizontal transfers of informational genes

from bacteria to plasmids and phages: implications for the origin of bacterial

replication machinery. Mol Microbiol 35:1-5.

35.   Shaffer M, Borton MA, McGivern BB, Zayed AA, La Rosa SL, Solden LM, Liu P,

Narrowe AB, Rodriguez-Ramos J, Bolduc B, Gazitua MC, Daly RA, Smith GJ, Vik

DR, Pope PB, Sullivan MB, Roux S, Wrighton KC. 2020. DRAM for distilling

microbial metabolism to automate the curation of microbiome function. Nucleic Acids

Res 48:8883-8900.

36.   Sullivan MJ, Petty NK, Beatson SA. 2011. Easyfig: a genome comparison visualizer.

Bioinformatics 27:1009-10.

37.   Price MN, Dehal PS, Arkin AP. 2010. FastTree 2--approximately maximum-

716   likelihood trees for large alignments. PLoS One 5:e9490.

717   38.   Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, Buxton S,

718   Cooper A, Markowitz S, Duran C, Thierer T, Ashton B, Meintjes P, Drummond A.

719   2012. Geneious Basic: an integrated and extendable desktop software platform for the

720   organization and analysis of sequence data. Bioinformatics 28:1647-9.

721   39.   Edgar RC. 2004. MUSCLE: a multiple sequence alignment method with reduced time

722   and space complexity. BMC Bioinformatics 5:113.

723   40.   Schlitzer R. 2015. Ocean Data View. https://odv.awi.de/.

724   41.   Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D, Amin N,

725   Schwikowski B, Ideker T. 2003. Cytoscape: a software environment for integrated

726   models of biomolecular interaction networks. Genome Res 13:2498-504.

727   42.   Askora A, Kawasaki T, Usami S, Fujie M, Yamada T. 2009. Host recognition and

728   integration of filamentous phage phiRSM in the phytopathogen, *Ralstonia*

729   *solanacearum*. Virology 384:69-76.

730   43.   da Silva Xavier A, de Almeida JCF, de Melo AG, Rousseau GM, Tremblay DM, de

731   Rezende RR, Moineau S, Alfenas-Zerbini P. 2019. Characterization of CRISPR-Cas

732   systems in the *Ralstonia solanacearum* species complex. Mol Plant Pathol 20:223-

733   239.

734   44.   Weissman JL, Laljani RMR, Fagan WF, Johnson PLF. 2019. Visualization and

735   prediction of CRISPR incidence in microbial trait-space to identify drivers of antiviral

736   immune strategy. ISME J 13:2589-2602.

737   45.   Rollie C, Chevallereau A, Watson BN, Chyou T-y, Fradet O, McLeod I, Fineran PC,

738   Brown CM, Gandon S, Westra ER. 2020. Targeting of temperate phages drives loss of

739   type I CRISPR–Cas systems. Nature 578:149-153.

740   46.   Koonin EV, Dolja VV, Krupovic M, Varsani A, Wolf YI, Yutin N, Zerbini FM, Kuhn

741   JH. 2020. Global organization and proposed megataxonomy of the virus world.

742   Microbiol Mol Biol Rev 84.

743   47.   Murugaiyan S, Bae JY, Wu J, Lee SD, Um HY, Choi HK, Chung E, Lee JH, Lee SW.

744   2011. Characterization of filamentous bacteriophage PE226 infecting *Ralstonia*

745   *solanacearum* strains. J Appl Microbiol 110:296-303.

746   48.   Mai-Prochnow A, Hui JG, Kjelleberg S, Rakonjac J, McDougald D, Rice SA. 2015.

747   'Big things in small packages: the genetics of filamentous phage and effects on fitness

748   of their host'. FEMS Microbiol Rev 39:465-87.

749   49.   Sanguino L, Franqueville L, Vogel TM, Larose C. 2015. Linking environmental

750   prokaryotic viruses and their host through CRISPRs. FEMS Microbiol Ecol 91.

751   50.   Hwang Y, Rahlff J, Schulze-Makuch D, Schloter M, Probst AJ. 2021. Diverse viruses

752   carrying genes for microbial extremotolerance in the Atacama Desert hyperarid soil.

753   mSystems 6:e00385-21.

754   51.   Van TTB, Yoshida S, Miki K, Kondo A, Kamei K. 2014. Genomic characterization of

755   ⬜RS603, a filamentous bacteriophage that is infectious to the phytopathogen

756   *Ralstonia solanacearum*. Microbiology and immunology 58:697-700.

757   52.   Askora A, Yamada T. 2015. Two different evolutionary lines of filamentous phages in

758   *Ralstonia solanacearum:* their effects on bacterial virulence. Front Genet 6:217.

759   53.   Filippova SN, Surgucheva NA, Sorokin VV, Akimov VN, Karnysheva EA, Brushkov

760   AV, Andersen D, Gal'chenko VF. 2016. Bacteriophages in Arctic and Antarctic low-

761   temperature systems. Microbiology 85:359-366.

762   54.   Dziewit L, Radlinska M. 2016. Two inducible prophages of an Antarctic

763   *Pseudomonas* sp. ANT_H14 use the same capsid for packaging their genomes -

764   Characterization of a novel phage helper-satellite system. PLoS One 11:e0158889.

55.   Filippova SN, Surgucheva NA, Kulikov EE, Sorokin VV, Akimov VN, Bej AK,

McKay C, Andersen D, Galchenko VF. 2013. Detection of phage infection in the

bacterial population of Lake Untersee (Antarctica). Microbiology 82:383-386.

56.   Askora A, Kawasaki T, Fujie M, Yamada T. 2011. Resolvase-like serine recombinase

mediates integration/excision in the bacteriophage phiRSM. J Biosci Bioeng 111:109-

16.

57.   Ahmad AA, Stulberg MJ, Mershon JP, Mollov DS, Huang Q. 2017. Molecular and

biological characterization of varphiRs551, a filamentous bacteriophage isolated from

a race 3 biovar 2 strain of *Ralstonia solanacearum*. PLoS One 12:e0185034.

58.   Sommers P, Fontenele RS, Kringen T, Kraberger S, Porazinska DL, Darcy JL,

Schmidt SK, Varsani A. 2019. Single-stranded DNA viruses in Antarctic cryoconite

holes. Viruses 11.

59.   Smith DJ, Timonen HJ, Jaffe DA, Griffin DW, Birmele MN, Perry KD, Ward PD,

Roberts MS. 2013. Intercontinental dispersal of bacteria and archaea by transpacific

winds. Appl Environ Microbiol 79:1134-9.

60.   Short CM, Suttle CA. 2005. Nearly identical bacteriophage structural gene sequences

are widely distributed in both marine and freshwater environments. Appl Environ

Microb 71:480-486.

61.   Breitbart M, Rohwer F. 2005. Here a virus, there a virus, everywhere the same virus?

Trends Microbiol 13:278-84.

62.   Breitbart M, Miyake JH, Rohwer F. 2004. Global distribution of nearly identical

phage-encoded DNA sequences. FEMS Microbiol Lett 236:249-56.

63.   Li Y, Endo H, Gotoh Y, Watai H, Ogawa N, Blanc-Mathieu R, Yoshida T, Ogata H.

2019. The Earth is small for "Leviathans": Long distance dispersal of giant viruses

across aquatic environments. Microbes Environ 34:334-339.

64.   Bellas CM, Schroeder DC, Edwards A, Barker G, Anesio AM. 2020. Flexible genes

establish widespread bacteriophage pan-genomes in cryoconite hole ecosystems. Nat

Commun 11:4403.

65.   Hughes KA, Convey P, Pertierra LR, Vega GC, Aragon P, Olalla-Tarraga MA. 2019.

Human-mediated dispersal of terrestrial species between Antarctic biogeographic

regions: A preliminary risk assessment. J Environ Manage 232:73-89.

66.   Benninghoff W, Benninghoff A. 1985. Wind transport of electrostatically charged

particles and minute organisms in Antarctica, p 592-596, Antarctic nutrient cycles and

food webs. Springer.

67.   Michaud JM, Thompson LR, Kaul D, Espinoza JL, Richter RA, Xu ZZ, Lee C, Pham

KM, Beall CM, Malfatti F, Azam F, Knight R, Burkart MD, Dupont CL, Prather KA.

2018. Taxon-specific aerosolization of bacteria and viruses in an experimental ocean-

atmosphere mesocosm. Nat Commun 9:2017.

68.   Aller JY, Kuznetsova MR, Jahns CJ, Kemp PF. 2005. The sea surface microlayer as a

source of viral and bacterial enrichment in marine aerosols. J Aerosol Sci 36:801-812.

69.   Wilbourn EK, Thornton DCO, Ott C, Graff J, Quinn PK, Bates TS, Betha R, Russell

LM, Behrenfeld MJ, Brooks SD. 2020. Ice nucleation by marine aerosols over the

North Atlantic Ocean in late spring. J Geophys Res Atmos 125:e2019JD030913.

70.   Amato P. 2012. Clouds provide atmospheric oases for microbes. Microbe 7:119-123.

71.   Christner BC, Morris CE, Foreman CM, Cai R, Sands DC. 2008. Ubiquity of

biological ice nucleators in snowfall. Science 319:1214.

72.   Strother SL, Salzmann U, Roberts SJ, Hodgson DA, Woodward J, Van Nieuwenhuyze

W, Verleyen E, Vyverman W, Moreton SG. 2015. Changes in Holocene climate and

the intensity of Southern Hemisphere Westerly Winds based on a high-resolution

palynological record from sub-Antarctic South Georgia. Holocene 25:263-279.

73.   Netea MG, Dominguez-Andres J, Eleveld M, Op den Camp HJM, van der Meer JWM,

816   Gow NAR, de Jonge MI. 2020. Immune recognition of putative alien microbial

817   structures: Host-pathogen interactions in the age of space travel. PLoS Pathog

818   16:e1008153.

819   74.   Nicholson WL, Schuerger AC, Race MS. 2009. Migrating microbes and planetary

820   protection. Trends Microbiol 17:389-92.

821   75.   Coenye T, Vandamme P, LiPuma JJ. 2002. Infection by *Ralstonia* species in cystic

822   fibrosis patients: identification of *R. pickettii* and *R. mannitolilytica* by polymerase

823   chain reaction. Emerg Infect Dis 8:692-6.

824   76.   Berliner AJ, Mochizuki T, Stedman KM. 2018. Astrovirology: Viruses at large in the

825   universe. Astrobiology 18:207-223.

826   77.   Mattoni RHT, Keller Jr EC. 1972. Induction of lysogenic bacteria in the space. The

827   Experiments of Biosatellite II 204:309.

828   78.   Mattoni RHT. 1968. Influence of spaceflight and radiation on induction of prophage

829   P-22 in *Salmonella typhimurium*. Jpn J Genet 43:465-465.

830   79.   Mattoni RHT. 1968. Space-flight effects and gamma radiation interaction on growth

831   and induction of lysogenic bacteria, a preliminary report. BioScience 18:602-608.

832   80.   Koike J. 1991. Fundamental questions concerning the contamination of other planets

833   with terrestrial microorganisms carried by space-probes. The Journal of Space

834   Technology and Science 7:9-14.

835   81.   Weinmaier T, Probst AJ, La Duc MT, Ciobanu D, Cheng JF, Ivanova N, Rattei T,

836   Vaishampayan P. 2015. A viability-linked metagenomic analysis of cleanroom

837   environments: eukarya, prokaryotes, and viruses. Microbiome 3:62.

838   82.   Aboudan A, Colombatti G, Bettanini C, Ferri F, Lewis S, Van Hove B, Karatekin O,

839   Debei S. 2018. ExoMars 2016 Schiaparelli Module Trajectory and Atmospheric

840   Profiles Reconstruction. Space Science Reviews 214:1-31.

841   83.   Bushnell B. 2014. BBTools software package. URL http://sourceforge

842   net/projects/bbmap 578:579.

843   84.   Joshi N, Fass J. 2011. Sickle: A sliding-window, adaptive, quality-based trimming

844   tool for FastQ files (Version 1.33)[Software].

845   85.   Nurk S, Meleshko D, Korobeynikov A, Pevzner PA. 2017. metaSPAdes: a new

846   versatile metagenomic assembler. Genome Res 27:824-834.

847   86.   Dick GJ, Andersson AF, Baker BJ, Simmons SL, Thomas BC, Yelton AP, Banfield

848   JF. 2009. Community-wide analysis of microbial genome sequence signatures.

849   Genome Biol 10:R85.

850   87.   Wu YW, Simmons BA, Singer SW. 2016. MaxBin 2.0: an automated binning

851   algorithm to recover genomes from multiple metagenomic datasets. Bioinformatics

852   32:605-7.

853   88.   Langmead B, Salzberg SL. 2012. Fast gapped-read alignment with Bowtie 2. Nat

854   Methods 9:357-9.

855   89.   Sieber CMK, Probst AJ, Sharrar A, Thomas BC, Hess M, Tringe SG, Banfield JF.

856   2018. Recovery of genomes from metagenomes via a dereplication, aggregation and

857   scoring strategy. Nat Microbiol 3:836-843.

858   90.   Bornemann TLV, Esser SP, Stach L, Burg T, Probst AJ. 2020. uBin – a manual

859   refining tool for metagenomic bins designed for educational purposes. bioRxiv

860   doi:10.1101/2020.07.15.204776

861   91.   Probst AJ, Castelle CJ, Singh A, Brown CT, Anantharaman K, Sharon I, Hug LA,

862   Burstein D, Emerson JB, Thomas BC, Banfield JF. 2017. Genomic resolution of a

863   cold subsurface aquifer community provides metabolic insights for novel microbes

864   adapted to high CO_2 concentrations. Environ Microbiol 19:459-474.

865   92.   Couvin D, Bernheim A, Toffano-Nioche C, Touchon M, Michalik J, Neron B, Rocha

866   EPC, Vergnaud G, Gautheret D, Pourcel C. 2018. CRISPRCasFinder, an update of

CRISPRFinder, includes a portable version, enhanced performance and integrates

search for Cas proteins. Nucleic Acids Res 46:W246-W251.

93.   Skennerton CT, Imelfort M, Tyson GW. 2013. Crass: identification and reconstruction

of CRISPR from unassembled metagenomic data. Nucleic Acids Res 41:e105.

94.   Mojica FJ, Diez-Villasenor C, Soria E, Juez G. 2000. Biological significance of a

family of regularly spaced repeats in the genomes of Archaea, Bacteria and

mitochondria. Mol Microbiol 36:244-6.

95.   Altschul SF. 2001. BLAST algorithm. e LS.

96.   Moller AG, Liang C. 2017. MetaCRAST: reference-guided extraction of CRISPR

spacers from unassembled metagenomes. PeerJ 5:e3788.

97.   Rahlff J, Turzynski V, Esser SP, Monsees I, Bornemann TLV, Figueroa-Gonzalez PA,

Schulz F, Woyke T, Klingl A, Moraru C, Probst AJ. 2021. Lytic archaeal viruses

infect abundant primary producers in Earth's crust. Nat Commun 12:4642.

98.   Roux S, Emerson JB, Eloe-Fadrosh EA, Sullivan MB. 2017. Benchmarking viromics:

an in silico evaluation of metagenome-enabled estimates of viral community

composition and diversity. PeerJ 5:e3817.

99.   Nayfach S, Camargo AP, Schulz F, Eloe-Fadrosh E, Roux S, Kyrpides NC. 2020.

CheckV assesses the quality and completeness of metagenome-assembled viral

genomes. Nat Biotechnol doi:10.1038/s41587-020-00774-7.

100. Antipov D, Raiko M, Lapidus A, Pevzner PA. 2020. Metaviral SPAdes: assembly of

viruses from metagenomic data. Bioinformatics 36:4126-4129.

101. Meier-Kolthoff JP, Göker M. 2017. VICTOR: genome-based phylogeny and

classification of prokaryotic viruses. Bioinformatics 33:3396-3404.

102. Lefort V, Desper R, Gascuel O. 2015. FastME 2.0: A comprehensive, accurate, and

fast distance-based phylogeny inference program. Mol Biol Evol 32:2798-800.

103.   Farris JS. 1972. Estimating phylogenetic trees from distance matrices. The American

893   Naturalist 106:645-668.

894   104.   Rambaut A. 2006. FigTree, a graphical viewer of phylogenetic trees and as a program

895   for producing publication-ready figures.

896   105.   Göker M, Garcia-Blazquez G, Voglmayr H, Telleria MT, Martin MP. 2009. Molecular

897   taxonomy of phytopathogenic fungi: a case study in *Peronospora*. PLoS One 4:e6319.

898   106.   Meier-Kolthoff JP, Hahnke RL, Petersen J, Scheuner C, Michael V, Fiebig A, Rohde

899   C, Rohde M, Fartmann B, Goodwin LA, Chertkov O, Reddy T, Pati A, Ivanova NN,

900   Markowitz V, Kyrpides NC, Woyke T, Göker M, Klenk HP. 2014. Complete genome

901   sequence of DSM 30083(T), the type strain (U5/41(T)) of *Escherichia coli*, and a

902   proposal for delineating subspecies in microbial taxonomy. Stand Genomic Sci 9:2.

903   107.   Bolduc B, Jang HB, Doulcier G, You ZQ, Roux S, Sullivan MB. 2017. vConTACT:

904   an iVirus tool to classify double-stranded DNA viruses that infect Archaea and

905   Bacteria. PeerJ 5:e3243.

906   108.   Bin Jang H, Bolduc B, Zablocki O, Kuhn JH, Roux S, Adriaenssens EM, Brister JR,

907   Kropinski AM, Krupovic M, Lavigne R, Turner D, Sullivan MB. 2019. Taxonomic

908   assignment of uncultivated prokaryotic virus genomes is enabled by gene-sharing

909   networks. Nat Biotechnol 37:632-639.

910   109.   Brister JR, Ako-Adjei D, Bao Y, Blinkova O. 2015. NCBI viral genomes resource.

911   Nucleic Acids Res 43:D571-7.

912   110.   Ahlgren NA, Ren J, Lu YY, Fuhrman JA, Sun F. 2017. Alignment-free d_2^{*}

913   oligonucleotide frequency dissimilarity measure improves prediction of hosts from

914   metagenomically-derived viral sequences. Nucleic Acids Res 45:39-53.

915   111.   Moraru C, Varsani A, Kropinski AM. 2020. VIRIDIC-A novel tool to calculate the

916   intergenomic similarities of prokaryote-infecting viruses. Viruses 12.

917   112.   Nilsson E, Bayfield OW, Lundin D, Antson AA, Holmfeldt K. 2020. Diversity and

918   host interactions among virulent and temperate Baltic Sea *Flavobacterium* phages.

919   Viruses 12:158.

920   113.   Hyatt D, Chen GL, Locascio PF, Land ML, Larimer FW, Hauser LJ. 2010. Prodigal:

921   prokaryotic gene recognition and translation initiation site identification. BMC

922   Bioinformatics 11:119.

923   114.   Buchfink B, Xie C, Huson DH. 2015. Fast and sensitive protein alignment using

924   DIAMOND. Nature Methods 12:59-60.

925   115.   Zimmermann L, Stephens A, Nam SZ, Rau D, Kubler J, Lozajic M, Gabler F, Soding

926   J, Lupas AN, Alva V. 2018. A completely reimplemented MPI bioinformatics toolkit

927   with a new HHpred server at its core. J Mol Biol 430:2237-2243.

928   116.   Söding J, Biegert A, Lupas AN. 2005. The HHpred interactive server for protein

929   homology detection and structure prediction. Nucleic Acids Res 33:W244-8.

930   117.   Paez-Espino D, Roux S, Chen IA, Palaniappan K, Ratner A, Chu K, Huntemann M,

931   Reddy TBK, Pons JC, Llabres M, Eloe-Fadrosh EA, Ivanova NN, Kyrpides NC. 2019.

932   IMG/VR v.2.0: an integrated data management and analysis system for cultivated and

933   environmental viral genomes. Nucleic Acids Res 47:D678-D686.

934   118.   Seemann T. 2014. Prokka: rapid prokaryotic genome annotation. Bioinformatics

935   30:2068-9.