1

LuxT is a Global Regulator of Low-Cell Density Behaviors Including Type III

Secretion, Siderophore Production, and Aerolysin Secretion in *Vibrio harveyi*

Michaela J. Eickhoff[a]*, Chenyi Fei[a,b], Jian-Ping Cong[a,c]^, and Bonnie L. Bassler[a,c]#

[a]Department of Molecular Biology, Princeton University, Princeton, New Jersey, USA.

[b]Lewis-Sigler Institute for Integrative Genomics, Princeton University, Princeton, New

Jersey, USA.

[c]Howard Hughes Medical Institute, Chevy Chase, Maryland, USA.

Running Title: LuxT is a global regulator in *Vibrio harveyi*

20 #Address correspondence to Bonnie L. Bassler, bbassler@princeton.edu.

21 *Present address: New Jersey Department of Health, Public Health and Environmental

22 Laboratories, Ewing, New Jersey, USA.

23 ^Deceased.

Abstract

Quorum sensing (QS) is a chemical communication process in which bacteria produce, release, and detect extracellular signaling molecules called autoinducers. Via combined transcriptional and post-transcriptional regulatory mechanisms, QS allows bacteria to collectively alter gene expression on a population-wide scale. Recently, the LuxT transcription factor was shown to control *V. harveyi qrr*1, encoding the Qrr1 small RNA that functions at the core of the QS regulatory cascade. Here, we use RNA-Sequencing to reveal that, beyond control of *qrr*1, LuxT is a global regulator of 414 *V. harveyi* genes including those involved in type III secretion, siderophore production, and aerolysin toxin biosynthesis. Importantly, LuxT directly represses *swrZ*, encoding a transcription factor, and LuxT control of type III secretion, siderophore, and aerolysin genes occurs by two mechanisms, one that is SwrZ-dependent and one that is SwrZ-independent. All of these target genes specify QS-controlled behaviors that are enacted when *V. harveyi* is at low cell density. Thus, LuxT and SwrZ function in parallel with QS to drive particular low cell density behaviors. Phylogenetic analyses reveal that *luxT* is highly conserved among *Vibrionaceae,* but *swrZ* is less well conserved. In a test case to examine the relationship between LuxT and SwrZ, we find that in *Aliivibrio fischeri,* LuxT also functions as a *swrZ* repressor, and LuxT activates *A. fischeri* siderophore production via *swrZ* repression. Our results indicate that LuxT is a major regulator among *Vibrionaceae,* and, in the species that also possess *swrZ*, LuxT functions with SwrZ to control gene expression.

Importance

Bacteria precisely tune gene expression patterns to successfully react to changes that occur in the environment. Defining the mechanisms that enable bacteria to thrive in diverse and fluctuating habitats, including in host organisms, is crucial for a deep understanding of the microbial world and also for development of effective applications to promote or to combat particular bacteria. In this study, we show that a regulator called LuxT controls over 400 genes in the marine bacterium *Vibrio harveyi* and, moreover, that LuxT is highly conserved among *Vibrionaceae* species, ubiquitous marine bacteria that often cause disease. We characterize the mechanisms by which LuxT controls genes involved in virulence and nutrient acquisition. We show that LuxT functions in parallel with a set of regulators of the bacterial cell-to-cell communication process called quorum sensing to promote *V. harveyi* behaviors at low cell density.

Introduction

Bacteria have the remarkable ability to rapidly detect and adapt to environmental fluctuations. Often, bacteria employ transcriptional and post-transcriptional regulatory mechanisms to tune gene expression patterns that enhance survival under varying conditions (1, 2). Such combined regulatory mechanisms are used in a process called quorum sensing (QS) to monitor and react to changes in the cell density and the species composition of the vicinal community. QS involves the production, release, accumulation, and group-wide detection of signaling molecules called autoinducers (AIs). QS fosters the synchronous execution of collective behaviors, typically, ones that are unproductive for

an individual bacterium to carry out alone but that become effective when enacted by the

group, e.g., bioluminescence, virulence factor production, and biofilm formation (3, 4).

Vibrio harveyi, the focus of the current work, is a model QS bacterium that uses

three AIs that act in parallel to control bioluminescence, type III secretion, siderophore

production, and hundreds of other traits (5-8). Each AI is detected by a cognate

membrane-bound receptor. At low-cell density (LCD), AI concentrations are low, and the

three unliganded QS receptors act as kinases that drive phosphorylation of the response

regulator LuxO. LuxO-P activates transcription of genes encoding five small regulatory

RNAs (sRNAs) called the Qrr sRNAs that function post-transcriptionally to activate and

repress, respectively, the translation of the master QS regulators AphA and LuxR (9-12).

Thus, at LCD, AphA is made and LuxR is not. AphA is responsible for executing the LCD

QS program (Fig S1A). At LCD, the Qrr sRNAs also directly control 16 other target

mRNAs, and they operate as feedback regulators within the QS signaling pathway (13-

15). At high-cell density (HCD), accumulated AIs bind to their cognate QS receptors. In

the liganded state, the QS receptors acts as phosphatases, and phosphate is stripped

from LuxO, which inactivates it (16, 17). Thus, at HCD, Qrr sRNA production terminates,

AphA is not made, and, by contrast, LuxR is produced (8, 10, 12). LuxR drives the HCD

QS regulon (Fig S1B).

The five *V. harveyi* Qrr sRNAs possess high sequence identity, and Qrr2-5

regulate an identical set of mRNA targets (10, 13). Qrr1 is the outlier. Because it lacks

nine nucleotides that are present in Qrr2-5, Qrr1 fails to regulate *aphA* and two additional

mRNA targets that are controlled by Qrr2-5 (10, 11, 13). The genes encoding the Qrr

sRNAs differ in their LCD expression levels: *qrr*4 is the most highly transcribed of the set,

whereas only low-level transcription of *qrr*1 and *qrr*5 occurs (10). Recently, we discovered

that a transcription factor called LuxT represses expression of *qrr*1. LuxT does not

regulate *qrr*2-5. Thus, the exclusive involvement of LuxT in *qrr*1 regulation provides a

mechanism enabling Qrr1 to uniquely control downstream targets (18). Indeed, via

repression of *qrr*1, LuxT indirectly controls the translation of Qrr1 target mRNAs including

those encoding a secreted protease, an aerolysin toxin, a chitin deacetylase, and a

component involved in capsule polysaccharide secretion. LuxT also activates

transcription of this same set of genes by a Qrr1-independent mechanism (18). LuxT

repression of *qrr*1 does not significantly alter LuxR translation, and as mentioned, Qrr1

does not regulate AphA (11, 18). These findings indicate that LuxT repression of *qrr*1

tunes the expression of select Qrr1-controlled mRNAs without altering the entire QS

regulon.

To date, *qrr*1 is the only gene identified to be directly controlled by LuxT (18, 19).

However, LuxT has been linked to the regulation of additional phenotypes in *V. harveyi*

and other *Vibrionaceae* including bioluminescence, siderophore production, virulence

factor production, and motility (18, 20-23). These findings, together with our initial

knowledge that LuxT can regulate transcription of target genes independently of Qrr1 in

V. harveyi (18), inspired us to investigate the role of LuxT beyond its control of *qrr*1. Here,

we use RNA-Sequencing (RNA-Seq) to identify the LuxT regulon, revealing LuxT to be a

global regulator of ~414 genes in *V. harveyi*. We find that LuxT directly represses the *V.*

harveyi swrZ gene encoding a GntR-family transcriptional regulator. We use genetic and

molecular analyses to show that, in *V. harveyi*, LuxT activates genes required for type III

secretion, siderophore production, aerolysin toxin production, and activation occurs by

two mechanisms. One mechanism depends on SwrZ: LuxT represses *swrZ* and SwrZ

represses target gene expression. The second mechanism is SwrZ independent. Finally,

the LuxT-controlled traits identified here are also QS-controlled and are enacted primarily

at LCD. Therefore, LuxT functions in parallel with QS to establish *V. harveyi* LCD

behaviors. Finally, we analyze *luxT* and *swrZ* conservation among *Vibrionaceae* bacteria,

and we demonstrate that, via *swrZ* repression, LuxT also activates genes required for

siderophore production in *Aliivibrio fischeri.*

Results

LuxT is a global regulator that directly represses *swrZ* transcription.

To define the set of genes regulated by LuxT*,* we used RNA-Seq to compare the

transcriptomes of WT and Δ*luxT V. harveyi*. We considered transcripts with 2-fold or

higher changes in expression ($p < 0.01$) to be LuxT-regulated, revealing a total of 414

genes: 243 activated and 171 repressed (Fig 1A and Data Set S1). One gene, *swrZ*

(*VIBHAR_RS03920*), stood out due to its dramatic repression by LuxT. *swrZ* was

previously shown to be repressed by the LuxT homolog SwrT in *Vibrio parahaemolyticus*

(22). Specifically, SwrT repressed transcription of *swrZ*, and SwrZ repressed the *laf* genes

encoding the lateral flagellar machinery required for *V. parahaemolyticus* swarming

locomotion. Thus, through this cascade – repression of a repressor – SwrT activates *V.*

parahaemolyticus swarming (22). The *V. harveyi* genome does not encode *laf* genes, and

V. harveyi does not swarm. Nonetheless, our data demonstrate that the regulatory

arrangement in which LuxT represses *swrZ* exists in *V. harveyi*: qRT-PCR verified that

LuxT represses *swrZ* expression by ~100-fold (Fig 1B).

To determine if LuxT repression of *swrZ* is direct, we used recombinant

Escherichia coli to isolate LuxT and the *swrZ* promoter from other *V. harveyi* regulatory

components. Two plasmids were introduced into *E. coli.* One plasmid harbored a P*swrZ*-

lux transcriptional reporter, and the second plasmid carried arabinose-inducible *luxT*.

Induction of *luxT* expression drove repression of P*swrZ*-*lux* in the *E. coli* strain carrying the

p*luxT* vector, whereas no repression occurred in *E. coli* carrying the empty control vector

(Fig 1C). We conclude that LuxT directly represses *swrZ*. Consistent with this supposition,

purified LuxT-6xHis protein bound to the *swrZ* promoter in an *in vitro* electrophoretic

mobility shift assay (EMSA) whereas LuxT-6xHis did not bind to control DNA (Fig S2A).

The laddering present in the P*swrZ* EMSA may indicate that multiple LuxT-binding sites

exist in the *swrZ* promoter (Fig S2A).

Both *luxT* and *swrZ* are subject to feedback regulation, and neither *luxT* nor *swrZ*

is controlled by QS.

To further explore connections between LuxT and SwrZ in *V. harveyi*, we assessed

whether feedback regulatory loops exist. First, regarding SwrZ regulation of *luxT*: We find

no evidence for SwrZ-mediated feedback onto *luxT* because *luxT* transcript levels were

similar in WT and Δ*swrZ V. harveyi* (Fig 1D), and, moreover, overexpression of *swrZ* in

WT *V. harveyi* did not alter *luxT* transcription as measured by qRT-PCR (Fig S3A). qRT-

PCR validated that *swrZ* was indeed overexpressed from the p*swrZ* plasmid (Fig S3A).

Second, we investigated autoregulation of *luxT*. Activity of a P*luxT*-*lux* reporter was

measured in Δ*luxA* and Δ*luxA* Δ*luxT V. harveyi* strains. *V. harveyi* is naturally

bioluminescent and *luxA* encodes a luciferase subunit. Thus, deletion of *luxA* was

required to ensure that all light produced by the test strains came from the P*luxT*-*lux*

reporter. Fig 1E shows that, in strains harboring an empty control vector, P*luxT*-*lux*

expression is 6-fold higher in the Δ*luxA V. harveyi* strain than in the Δ*luxA* Δ*luxT* strain.

Introduction of a vector expressing *luxT* restored light production (Fig 1E). Thus, LuxT

activates its own transcription. Unfortunately, autoregulation of *luxT* could not be tested

in recombinant *E. coli* because the P*luxT*-*lux* reporter was not expressed to any detectable

level. However, in an *in vitro* EMSA, LuxT-6xHis bound the *luxT* promoter, providing

evidence for a direct auto-regulatory role (Fig S2B). Finally, we investigated direct

autoregulation of *swrZ*. In recombinant *E. coli,* induction of *swrZ* expression repressed a

P*swrZ*-*lux* reporter 30-fold indicating direct feedback repression (Fig 1F). We conclude that

LuxT and SwrZ can be placed into a regulatory pathway in which LuxT directly represses

swrZ. Positive and negative feedback loops exist for LuxT and SwrZ, respectively (Fig

1G). Below, we speculate on the implications of this regulatory arrangement.

Analysis of the LuxT regulon revealed that a subset of LuxT-controlled genes are

also regulated by QS ((8) and Data Set S1). One possible explanation for this finding is

that regulatory interconnections exist between LuxT and QS. We know that LuxT does

not regulate *aphA* or *luxR,* so LuxT cannot reside upstream of those two components in

the QS cascade (18). Alternatively, QS could control *luxT* and/or *swrZ* expression. To test

this possibility, we measured *luxT* and *swrZ* expression in WT, Δ*luxO,* and *luxO* D61E *V.*

harveyi strains. The logic is as follows: The WT strain undergoes the normal LCD to HCD

QS transition. The Δ*luxO* strain exhibits "HCD-locked" behavior because in the absence

of LuxO activity, no Qrr sRNAs are produced ((9, 10) and Fig S1B). *luxO* D61E encodes

a LuxO-P mimetic, so the strain harboring this mutation exhibits "LCD-locked" behavior

in which the Qrr sRNAs are constitutively produced ((10, 17) and Fig S1A). There were

no significant differences in *luxT* or *swrZ* expression in these strains (Fig S3B). Consistent

with this finding, no dramatic changes in *luxT* expression occurred over growth as would

be expected for a QS-regulated gene (Fig S3C). As expected from our above findings

(Fig 1B), *swrZ* transcript levels were > 100-fold higher in the Δ*luxT* strain than in WT *V.*

harveyi, but again, cell-density-dependent changes in *swrZ* expression did not occur in

either strain, arguing against QS control (Fig S3D). We conclude that expression of *luxT*

and *swrZ* are not QS regulated. Thus, we infer that QS and LuxT/SwrZ function

independently and in parallel in the regulation of a subset of genes in each of their

regulons.

LuxT activates expression of type III secretion system genes via repression of

swrZ.

As mentioned above, LuxT regulates ~413 genes in addition to *swrZ* (Fig 1A and

Data Set S1). To gain insight into the LuxT regulon, as examples, we characterize the

mechanisms by which LuxT controls genes specifying three *V. harveyi* functions: type III

secretion, siderophore production, and aerolysin secretion. We begin with type III

secretion genes. Type III secretion systems (T3SSs) are syringe-like molecular machines

that ferry toxic effector proteins across bacterial inner and outer membranes and into

target eukaryotic cells (24-26). In *V. harveyi* and other vibrios, type III secretion is crucial

for virulence (27-29). The structural proteins comprising the *V. harveyi* T3SS are encoded

in four operons (T3SS.1, T3SS.2, T3SS.3, and T3SS.4; Fig 2A) (5, 8, 30). Type III

secretion is QS-regulated in *V. harveyi* (5). Specifically, T3SS genes are repressed by

both AphA and LuxR. The consequence is that T3SS gene expression peaks at low- to

mid-cell density when AphA levels have decreased and LuxR levels have not risen (8,

12). QS regulation of type III secretion occurs indirectly through ExsA, the master

transcriptional activator of T3SS genes (Fig 2A). Both AphA and LuxR directly repress

exsA (8, 30).

The genes in all four T3SS operons and *exsA* were revealed by RNA-Seq as

activated by LuxT (Data Set S1 and Fig 2A). To understand dual control of T3SS genes

by QS and LuxT, we engineered a set of strains and developed a companion strategy to

probe regulatory mechanisms. First, we verified LuxT involvement from the RNA-Seq

analysis, and we determined if *qrr*1 is required for LuxT control of T3SS genes. Using

qRT-PCR, we quantified transcript levels of four T3SS genes: *hyp, vopN, vscO,* and *exsD*,

representing the four T3SS operons (underlined in Fig 2A). Expression was compared

between WT and Δ*luxT V. harveyi* and between Δ*qrr*1 and Δ*qrr*1 Δ*luxT V. harveyi*.

Expression was, respectively, ~10, 14, 8, and 4-fold lower in the Δ*luxT* strain than in WT,

confirming that LuxT is an activator of T3SS genes (Fig S4A). Nearly identical results to

those shown in Fig S4A were obtained for the Δ*qrr*1 and Δ*qrr*1 Δ*luxT* strains, indicating

that LuxT activates T3SS genes independently of Qrr1 (Fig S4B).

To verify the mechanism by which QS regulates T3SS genes, we measured

transcript levels of the four representative genes in WT, Δ*luxO* (HCD-locked), and *luxO*

D61E (LCD-locked) *V. harveyi* strains. Expression of *hyp, vopN, vscO,* and *exsD* were >

100-fold higher in the *luxO* D61E strain than in the WT and Δ*luxO* strains at HCD (Fig

2B). Consistent with previously published results, high level expression of T3SS genes

occurs at LCD due to LuxR-mediated repression of them at HCD (5, 8). Specifically,

compared to WT, a Δ*luxR V. harveyi* strain exhibited > 100-fold higher expression of the

four genes (Fig S5A). AphA is known to repress T3SS.1 and T3SS.4 in *V. harveyi* (8).

Expression of *hyp* (T3SS.1) and *exsD* (TS33.4) were ~1.6-fold higher in the *luxO* D61E

Δ*aphA* strain compared to the *luxO* D61E strain, however, the differences were not

statistically significant (Fig S5B). We presume that we did not observe a larger role for

AphA here because of differences in our experimental growth conditions compared to

those used previously (8).

To assess the mechanism by which LuxT controls T3SS genes expression, we

first determined whether LuxT-mediated activation of T3SS genes requires SwrZ by

measuring transcript levels of the four representative T3SS genes in WT, Δ*luxT,* Δ*swrZ,*

and Δ*swrZ* Δ*luxT V. harveyi* strains at LCD. Comparison of expression levels between the

WT and Δ*luxT* strains validates LuxT activation of the genes. Comparison of expression

levels between the Δ*swrZ* and Δ*swrZ* Δ*luxT* strains tests whether LuxT activation of genes

requires SwrZ. As above (Fig S4A), expression of *hyp, vopN, vscO,* and *exsD* was lower

in the Δ*luxT* strain than in WT (Fig 2C). However, elimination of *luxT* did not alter transcript

levels in the Δ*swrZ* background (Fig 2C). Thus, LuxT activates expression of all four T3SS

operons via a SwrZ-dependent mechanism. Most likely, LuxT represses *swrZ,* and SwrZ

represses T3SS genes. We test this assumption below.

LuxT regulates *exsA* by SwrZ-dependent and SwrZ-independent mechanisms to

activate type III secretion.

To quantify the contributions of QS and LuxT/SwrZ to regulation of type III

secretion, we used Western blot assessment of VopD, a T3SS.1 protein that accumulates

in and is secreted from the cytoplasm following activation of T3SS gene expression (5).

We measured VopD levels in WT, ∆*luxT*, ∆*swrZ,* and ∆*swrZ* ∆*luxT V. harveyi* strains at

HCD and in the LCD-locked (*luxO* D61E) strain. No VopD was detected in the strains at

HCD, consistent with LuxR repression of T3SS genes (Fig 3A, first four lanes) (5, 30). By

contrast, VopD was made in the *luxO* D61E strain, and deletion of *luxT* in the *luxO* D61E

strain reduced VopD levels to below our detection limit (Fig 3A, fifth and sixth lanes).

These data mirror our qRT-PCR results from Fig 2C and confirm the role of LuxT as an

activator of type III secretion. To assess whether *swrZ* is required for LuxT activation of

type III secretion, VopD levels were measured in *luxO* D61E ∆*swrZ* and *luxO* D61E ∆*swrZ*

∆*luxT* strains. Lower levels of VopD were present in the *luxO* D61E ∆*swrZ* ∆*luxT* strain

than in the *luxO* D61E ∆*swrZ* strain (Fig 3A, right-most two lanes). These results differ

from the qRT-PCR results in Fig 2C, likely because of the different growth conditions

required for the qRT-PCR and the VopD Western blot analyses. We summarize the data

in Fig 3A as follows: Deletion of *luxT* reduces VopD production in the presence and

absence of *swrZ,* however a more dramatic reduction occurs when *swrZ* is present. To

explain these data, we developed and tested the following model for LuxT/SwrZ

regulation of T3SS genes:

We propose that LuxT activates T3SS genes by two mechanisms: One mechanism

is SwrZ-dependent, and one mechanism is SwrZ-independent. This model is depicted in

Fig 3B-E and is used to explain the data in the LCD-locked strains of the Western blot

shown in Fig 3A (right-most four lanes). In WT *V. harveyi* (Fig 3B), LuxT is present. It

represses *swrZ,* and it also activates T3SS genes by a SwrZ-independent mechanism.

The consequence is LuxT activation of T3SS genes. Therefore, high-level VopD

production occurs (fifth lane in Fig 3A). In Δ*luxT V. harveyi* (Fig 3C), *swrZ* is derepressed.

The consequence is that SwrZ is produced and available to repress T3SS genes.

Therefore, no VopD is produced (sixth lane in Fig 3A). In the Δ*swrZ* strain (Fig 3D), the

situation is essentially identical to WT in which *swrZ* is repressed by LuxT, but in this

case, there simply is no *swrZ.* The consequence is LuxT activation of T3SS genes.

Therefore, high-level VopD production occurs (seventh lane in Fig 3A). Finally, in the case

of the Δ*swrZ* Δ*luxT* double mutant (Fig 3E), there is no repression of T3SS genes by SwrZ

and there is no activation by LuxT. Thus, basal level expression of T3SS genes ensues,

and an intermediate amount of VopD is made (eighth lane in Fig 3A). Key to this model

is that VopD levels in the LCD-locked Δ*swrZ* and Δ*swrZ* Δ*luxT* strains are not identical

because of the SwrZ-independent mechanism by which LuxT activates T3SS genes (Fig

3D and 3E).

To test our model, we assessed whether LuxT activates T3SS gene expression by

a SwrZ-independent mechanism. To do this, we introduced a p*luxT* overexpression vector

into a Δ*luxR* Δ*swrZ* Δ*luxT V. harveyi* strain. The Δ*luxR* mutation was included to relieve

HCD repression of T3SS genes by LuxR. In this setup, we measured *hyp, vopN, vscO,*

and *exsD* transcript levels*.* We also measured expression of *exsA,* which is located

immediately upstream of T3SS.4 and, as mentioned, encodes the master transcriptional

activator of T3SS genes. Compared to the vector control, overexpression of *luxT* caused

~122, 74, 34, 57, and 19-fold increases in *hyp, vopN, vscO, exsD,* and *exsA* expression,

respectively (Fig 4A). These data demonstrate that LuxT activates the four T3SS operons

and *exsA* by a mechanism that does not require SwrZ. We also assessed *swrZ*

overexpression in this experimental setup. Repression by SwrZ was not observed (Fig

4A) presumably because in the absence of LuxT activation, T3SS gene expression is

minimal so no detectable repression by SwrZ can occur. To test this supposition, we

introduced the p*swrZ* overexpression construct into a Δ*luxR* Δ*swrZ V. harveyi* strain.

Indeed, in the *luxT*+ background, overexpression of *swrZ* caused an ~10-fold reduction in

hyp, vopN, and *vscO* transcript levels and ~9 and 3-fold reductions in *exsD* and *exsA*

levels, respectively (Fig 4B). Thus, SwrZ is a repressor of the four T3SS operons and

exsA.

Previously, it was discovered that T3SS gene expression does not occur in the

absence of *exsA* irrespective of the presence or absence of LuxR. Thus, ExsA is epistatic

to QS in the control of T3SS genes (30). Consistent with this finding, overexpression of

exsA overrode repression by LuxR to activate T3SS gene expression at HCD (30). Based

on these earlier data and our identification of LuxT and SwrZ as regulators of T3SS genes,

we wondered whether LuxT and SwrZ also exert control over T3SS genes via regulation

of *exsA*. To test this possibility, we measured *hyp, vopN, vscO,* and *exsD* expression in

WT, Δ*luxT*, Δ*exsA,* and Δ*exsA* Δ*luxT V. harveyi* strains. Fig 4C shows that, as expected,

expression of all four genes was lower in the Δ*luxT* strain than in WT. However, there

were no significant differences in *hyp, vopN,* and *vscO* transcript levels between the

Δ*exsA* and Δ*exsA* Δ*luxT* strains (Fig 4C). We conclude that *exsA* is required for LuxT

activation of T3SS.1, T3SS.2, and T3SS.3, encompassing *hyp, vopN,* and *vscO,*

respectively. Because LuxT activation of *hyp, vopN,* and *vscO* requires *swrZ* (see Fig

2C), we can also conclude that regulation by SwrZ requires *exsA*. Regarding T3SS.4,

there was ~3-fold less *exsD* transcript present in the Δ*exsA* Δ*luxT* strain than the Δ*exsA*

strain. It is possible that LuxT activates the T3SS.4 operon encompassing *exsD*

independently of ExsA or, alternatively, based on their sequential orientation in the

genome (Fig 2A), LuxT only activates the *exsA* promoter and read-through transcription

occurs for T3SS.4.

Together, our results show that LuxT activates T3SS genes by two mechanisms;

one is SwrZ-dependent, and one is SwrZ-independent. Also, LuxT functions

independently of QS to control these genes. QS and LuxT/SwrZ all modulate T3SS.1,

T3SS.2, and T3SS.3 via control of expression of *exsA* encoding a transcriptional activator

of T3SS genes. Previous results showed that *exsA* is repressed by both AphA and LuxR,

which constrains its expression to low- to mid-cell densities. Our scheme for QS and

LuxT/SwrZ regulation of T3SS genes is depicted in Fig 4D.

QS and LuxT/SwrZ regulate *V. harveyi* siderophore production.

Similar to type III secretion, the RNA-Seq analysis revealed LuxT to be an activator

of siderophore production genes in *V. harveyi.* Siderophores are small molecule iron

chelators that bacteria produce and secrete to scavenge extracellular iron (31).

Companion siderophore uptake systems import siderophore-Fe^{3+} complexes, facilitating

iron acquisition. *V. harveyi* encodes the genes required to produce, secrete, and import

two siderophores: amphi-enterobactin and anguibactin (32, 33), and both sets of genes

are regulated by LuxT (Fig 5A and 5B, respectively, and Data Set S1). *V. harveyi*

produces siderophores at LCD suggesting QS regulation, however, the mechanism

connecting QS to siderophore genes is not defined (6, 34). Using the steps laid out above

for characterization of T3SS gene regulation, we determine how QS, LuxT, and SwrZ

control siderophore production.

First, qRT-PCR validated the RNA-Seq data. Transcript levels of *aebC* encoding

isochorismate synthase were measured as a representative amphi-enterobactin

biosynthetic gene (32). Regarding anguibactin, expression of *angM* and *fatD* were

measured, and they encode a non-ribosomal peptide synthetase and an anguibactin-Fe^{3+}

transporter, respectively (35, 36). Transcript levels of *aebC, angM,* and *fatD* were ~3-fold

lower in Δ*luxT V. harveyi* than in WT (Fig S4C). Analogous results were obtained for the

three genes when measured in Δ*qrr*1 and Δ*qrr*1 Δ*luxT* strains, showing that LuxT

activates siderophore genes independently of Qrr1 (Fig S4D).

Second, to determine the mechanism by which QS controls siderophore

production, we used a chrome azurol S (CAS) dye to quantify siderophore levels in cell-

free culture fluids prepared from WT *V. harveyi* and QS mutant strains. Approximately 24-

fold more siderophore was present in fluids from the LCD-locked *luxO* D61E strain than

in fluids from the WT and Δ*luxO* strains, consistent with the known LCD production pattern

(Fig 5C) (6, 34). We used mutants defective in each siderophore biosynthesis pathway to

show that the siderophore detected by the CAS assay was amphi-enterobactin, not

anguibactin as follows: AebF is required for amphi-enterobactin production (32), and the

luxO D61E Δ*aebF* strain had almost no siderophore in its cell-free culture fluids (Fig S6A).

In contrast, the *luxO* D61E Δ*angN* strain, which lacks a gene required for anguibactin

production (37), did not exhibit reduced siderophore production compared to the *luxO*

D61E strain (Fig S6A). *V. harveyi* apparently does not produce anguibactin when grown

under standard laboratory conditions as reported previously (34).

With respect to *V. harveyi* QS, three mechanisms can explain how a gene acquires

a LCD expression pattern (Fig S1): It is activated at LCD by AphA, it is activated, directly

or indirectly, at LCD by the Qrr sRNAs, or it is repressed at HCD by LuxR. Regarding

AphA: Siderophore levels were similarly high in fluids from the *luxO* D61E and *luxO* D61E

Δ*aphA V. harveyi* strains, so AphA has no role in regulation of siderophore production

(Fig S6B). Regarding LuxR: The Δ*luxR* strain possessed ~16-fold more amphi-

enterobactin in its culture fluids than did WT at HCD (Fig S6C). Thus, QS control of amphi-

enterobactin production occurs via LuxR repression at HCD. Regarding the Qrr sRNAs:

At HCD somewhat less amphi-enterobactin was produced by the Δ*luxR* strain than by the

luxO D61E strain (Fig S6B and S6C), hinting that, in addition to the primary repressive

role of LuxR at HCD, at LCD, the Qrr sRNAs could activate siderophore biosynthesis or

export. Possibilities include direct post-transcriptional activation by the Qrr sRNAs or

alternatively, another QS-regulated factor may exist that is involved in amphi-enterobactin

gene regulation. We did not investigate this mechanism further.

To position LuxT and SwrZ in the siderophore regulatory pathway, we used qRT-

PCR quantitation of *aebC, angM,* and *fatD* in WT, Δ*luxT,* Δ*swrZ,* and Δ*swrZ* Δ*luxT V.*

harveyi strains (Fig 5D). As in Fig S4C, there were lower mRNA levels in the Δ*luxT* strain

than in the WT showing again that LuxT activates siderophore genes. Modestly increased

expression of the three genes occurred in the Δ*swrZ* strain compared to the WT, indicating

that SwrZ is a repressor of siderophore genes. Finally, the Δ*luxT* Δ*swrZ* double mutant

expressed siderophore genes at a level intermediate between those of the Δ*luxT* and

Δ*swrZ* strains. These data resemble the findings in Fig 3A for LuxT/SwrZ regulation of

T3SS genes, suggesting an identical model: LuxT activates siderophore production by

both a SwrZ-dependent and a SwrZ-independent mechanism. We validate this model

below.

Fur represses *V. harveyi* siderophore production in iron-rich conditions.

The above qRT-PCR measurements of siderophore transcripts showed only modest differences between strains (Fig 5D) possibly as a consequence of the generally low siderophore gene expression that occurs in iron-replete conditions due to repression by Fur, the major transcriptional regulator of iron transport. Fur repression is relieved in iron-limiting conditions (38). Indeed, culture fluids used for the CAS assays in Fig 5C and S6A-C were prepared from *V. harveyi* grown in low-iron minimal medium, and significant siderophore was produced. To assess the role of Fur in regulation of *V. harveyi* siderophore production, *aebC, angM,* and *fatD* transcript levels were compared between WT and Δ*fur V. harveyi*. In iron-rich medium, in the absence of *fur*, expression of the three test genes was ~284, 119, and 66-fold higher than when *fur* was present (Fig S6D)*.* Deletion of *luxT* in the Δ*fur* strain caused 3, 43, and 52-fold decreases in *aebC, angM,* and *fatD* transcript levels, respectively (Fig S6D). Thus, Fur is a repressor of amphi-enterobactin and anguibactin genes, which diminished our ability to observe their regulation by LuxT/SwrZ in iron-replete conditions, particularly for *angM* and *fatD.* We note that *fur* expression is not controlled by LuxT or SwrZ (Fig S6E). Thus, Fur and LuxT/SwrZ regulate siderophore genes independently. The siderophore experiments in the next section were conducted following growth of *V. harveyi* under iron-limited conditions.

LuxT activates *V. harveyi* siderophore production by SwrZ-dependent and SwrZ-independent mechanisms.

To validate the roles of and relationship between LuxT and SwrZ in regulation of siderophore production, we quantified siderophore produced by the LCD-locked *luxO* D61E, *luxO* D61E Δ*luxT*, *luxO* D61E Δ*swrZ*, and *luxO* D61E Δ*swrZ* Δ*luxT* mutant strains using the CAS assay. Consistent with the qRT-PCR results (Fig 5D), the *luxO* D61E Δ*swrZ* Δ*luxT* mutant produced intermediate siderophore levels between those of the *luxO* D61E Δ*luxT* strain and the *luxO* D61E Δ*swrZ* strain (Fig 5E). These data support a model in which LuxT activates siderophore production by two mechanisms – one dependent on SwrZ and one independent of SwrZ. We performed complementation analyses in a *V. harveyi* Δ*luxR* Δ*luxT* Δ*swrZ* strain as a final test of our model. Even in the absence of *swrZ,* overexpression of *luxT* caused increased expression of *aebC, angM,* and *fatD,* supporting the SwrZ-independent role for LuxT (Fig 5F). Following the reasoning described above for T3SS genes, repression due to *swrZ* overexpression could only be observed in the *luxT*+ background (Fig 5F and 5G). Our model for regulation of siderophore production in *V. harveyi* is depicted in Fig 5H: Fur represses siderophore genes in iron-replete conditions, and LuxR represses siderophore genes at HCD. LuxT activates siderophore genes by both a SwrZ-dependent and a SwrZ-independent mechanism.

QS and LuxT/SwrZ regulate production of the aerolysin toxin.

The final *V. harveyi* LuxT-regulated behavior we investigated from the RNA-Seq was production of the pore-forming toxin aerolysin. Aerolysin-like proteins, originally

discovered in the bacterium *Aeromonas hydrophila,* are a family of secreted proteins that

form transmembrane β-barrel pores in eukaryotic target cells, causing cell death (39-41).

Aerolysins are virulence factors in *Aeromonas* spp. and in *Vibrio splendidus* (42, 43).

We previously reported that LuxT is an activator of *VIBHAR_RS11620* encoding

an aerolysin toxin. For clarity, we now call *VIBHAR_RS11620 aerB.* We determined that

LuxT activates *aerB* expression by two mechanisms. First, LuxT activates *aerB*

transcription by a Qrr1-independent mechanism. Second, LuxT indirectly activates *aerB*

translation by repressing expression of *qrr*1 encoding a post-transcriptional repressor of

aerB (13, 18)*. V. harveyi* aerolysin production can be measured by monitoring hemolysis

of blood cells in liquid or on blood agar plates. Activation by LuxT is required for *V. harveyi*

hemolytic activity in both assays. Observable repression by Qrr1 occurs only in the plate

assay (18).

In agreement with our earlier findings, the RNA-Seq revealed LuxT as an activator

of the *aerB* aerolysin toxin gene. Three additional genes upstream of *aerB* were also

activated (Fig 6A and Data Set S1): *VIBHAR_RS11600*, which we call *aerR*, encodes a

transcriptional regulator, *VIBHAR_RS11605*, which we do not name, encodes a protein

of unknown function, and *VIBHAR_RS11610*, which we call *aerA*, encodes an additional

aerolysin family toxin. The protein sequences of AerA and AerB are similar (50% pairwise

identity, 66% pairwise positive), indicating a possible gene duplication event. The

VIBHAR_RS11615 gene that resides between *aerA* and *aerB* was not identified as

regulated by LuxT (Fig 6A).

As above, we first confirmed LuxT regulation of *aerR, aerA,* and *aerB*, and

determined if Qrr1 is required. Deletion of *luxT* in WT *V. harveyi* resulted in ~5, 28, and

15-fold lower expression of *aerR, aerA,* and *aerB,* respectively (Fig S4E). In a strain

lacking *qrr*1, deletion of *luxT* caused similar reductions in expression (Fig S4F). These

data show that Qrr1 is not required for LuxT activation of aerolysin genes. We know from

our earlier work that Qrr1 is a post-transcriptional regulator of *aerB.* Regulation by Qrr1

can only be detected when measuring *aerB* translation, not transcription (13, 18). For the

remainder of the present work, we focus on the transcriptional, Qrr1-independent,

mechanism by which LuxT activates aerolysin production.

QS regulation of aerolysin secretion is more complicated than we anticipated,

nonetheless, we lay out what we know here. In the liquid hemolysis assay, culture fluids

from the LCD-locked *luxO* D61E strain possessed ~17-fold more hemolysis activity than

fluids from HCD WT *V. harveyi* (Fig 6B). Fluids from the HCD-locked Δ*luxO* strain had

activity similar to that of the WT strain (Fig 6B). Thus, aerolysin secretion is QS-controlled

and produced at LCD. Deletion of *aphA* in the *luxO* D61E LCD-locked strain did not

reduce hemolysis (Fig S7A), and deletion of *luxR* in the WT did not increase hemolysis

at HCD (Fig S7B). These data were unexpected and show that, while aerolysin is QS-

controlled, neither AphA nor LuxR regulates its production. We confirmed the results

using blood agar plates. The results on the plates were identical to those in liquid with the

exception that the Δ*luxR* strain caused less hemolysis on plates than did the WT (Fig

S7C). Given the LCD production pattern for aerolysin, we conclude that aerolysin genes

must be activated at LCD by the QS Qrr sRNAs, either directly or indirectly. We note that

this finding seemingly contradicts our earlier result showing that Qrr sRNAs post-

transcriptionally repress *aerB* (13, 18)*.* We preliminarily predict that the Qrr sRNAs exert

both positive and negative effects on aerolysin secretion. Activation by the Qrr sRNAs is

likely indirect through an unknown regulator, while post-transcriptional repression of *aerB*

occurs directly (13). Further investigation to characterize the mechanism of QS control of

aerolysin genes is required.

While the mechanism underlying QS regulation of aerolysin genes is not fully

defined, what is clear is that *aerR*, *aerA*, and *aerB* are all regulated by LuxT (Data Set S1

and Fig S4E). Thus, we next explored which of the three genes is required for *V. harveyi*

hemolysis activity. We deleted each gene individually in the LCD-locked *luxO* D61E *V.*

harveyi strain. Deletion of *aerR* and *aerA* eliminated hemolysis activity in the liquid and

plate assays (Fig S7D and S7E, respectively). In contrast, deletion of *aerB* did not reduce

hemolytic activity, so *aerB* is not required (Fig S7D and S7E). We speculate that AerR is

an activator of *aerA*, and AerA is the primary secreted aerolysin toxin. Because *aerB* is

repressed by the Qrr sRNAs, *aerB* expression may remain low at LCD.

To understand the mechanism underlying LuxT activation of aerolysin gene

expression, we measured *aerR, aerA,* and *aerB* transcript levels in WT, Δ*luxT,* Δ*swrZ,*

and Δ*swrZ* Δ*luxT V. harveyi* strains at LCD. Expression of all three genes was high in the

WT and Δ*swrZ* strains, low in the Δ*luxT* strain, and intermediate in the Δ*swrZ* Δ*luxT* strain

(Fig 6C). The corresponding hemolysis patterns in the LCD-locked *luxO* D61E

background exactly mirrored the transcription patterns both in the liquid assay (Fig 6D)

and on the blood agar plates (Fig S7F). Exactly analogous to what we have explained

above, these data indicate that LuxT activates aerolysin production by both a SwrZ-

dependent and a SwrZ-independent mechanism. We verified this assumption using

complementation analyses (Fig 6E and 6F). The results were as expected according to

our putative mechanism except that *swrZ* overexpression did not repress *aerB* (Fig 6F).

However, the qRT-PCR results in Fig 6C show that SwrZ is a repressor of *aerB.* We

presume *swrZ* complementation did not repress *aerB* because *aerB* exhibits quite low

basal expression. Our model for LuxT/SwrZ regulation of aerolysin production is depicted

in Fig 6G.

luxT is and swrZ is not conserved among Vibrionaceae family members.

As mentioned, SwrT (LuxT) repression of *swrZ* was originally discovered in *V.*

parahaemolyticus and shown to be relevant for regulation of swarming motility (22). Here,

we have shown that LuxT also regulates *swrZ* in *V. harveyi,* and LuxT and SwrZ both

regulate type III secretion, siderophore production, and aerolysin secretion. LuxT also

regulates genes in *V. harveyi* that SwrZ does not control. For example, LuxT activates

luxCDABE, encoding luciferase, and SwrZ plays no role ((18, 20) and Fig S8). We

assessed SwrZ involvement in the regulation of 5 additional LuxT-activated and 3

additional LuxT-repressed genes. All of the additional LuxT-activated and none of the

LuxT-repressed genes exhibited regulation by SwrZ (Fig S8). We conclude that within the

LuxT regulon, a subset of genes does not employ SwrZ in regulation.

In *V. harveyi* and *V. parahaemolyticus,* the *luxT* (*swrT*) and *swrZ* genes are

encoded on the two different chromosomes, and so, while they co-regulate many genes,

it is unlikely they were inherited as a pair. Knowing their interconnected roles in *V. harveyi*

and *V. parahaemolyticus*, we wondered whether LuxT and SwrZ jointly regulate functions

in other *Vibrionaceae* species. To garner preliminary evidence for or against this

possibility, we examined the conservation of both genes within the *Vibrionaceae* family.

Among the 418 sequenced *Vibrionaceae,* the *luxT* gene is present in all but 16 species

(96%) (Fig 7A). None of the 16 species lacking *luxT* encodes a *swrZ* gene (Fig 7A).

Among the 402 *Vibrionaceae* that possess *luxT,* 227 species (56%) also have *swrZ* (Fig

7A); these genera include *Aliivibrio, Photobacterium,* and *Vibrio* (Fig 7A). Thus, in cases

in which both *luxT* and *swrZ* are present, the possibility for co-regulation of target genes

exists. In the species that have *luxT* but lack *swrZ*, LuxT must regulate genes

independently of SwrZ. Together, these observations may explain the evolution of SwrZ-

dependent and SwrZ-independent functions for LuxT in *V. harveyi.*

To probe the possibility that LuxT also represses *swrZ* in other species, we

performed a phylogenetic analysis by comparing the *swrZ* promoter regions of all *luxT*⁺

swrZ⁺ species. Our prediction was that *swrZ* promoters would be more similar in species

in which LuxT represses *swrZ* than in species in which LuxT does not regulate *swrZ*. In

support of this idea, the *V. harveyi* and *V. parahaemolyticus swrZ* promoter sequences

are indeed similar. Thus, when we constructed a phylogenetic tree based on *swrZ*

promoter sequences, *V. harveyi* and *V. parahaemolyticus* reside in close proximity (Fig

7B). As an initial test of LuxT regulation of *swrZ* in a species beyond *V. harveyi* and *V.*

parahaemolyticus, we analyzed LuxT repression of *swrZ* in *Aliivibrio fischeri,* a species

that neighbors *V. harveyi* and *V. parahaemolyticus* on the *swrZ* promoter phylogenetic

tree (Fig 7B). Indeed, *swrZ* transcript levels were ~10-fold higher in Δ*luxT A. fischeri* than

in WT showing that LuxT is a repressor of *swrZ* in *A. fischeri* (Fig 7C). We previously

reported that LuxT is an indirect activator of *iucABCD,* encoding the aerobactin

siderophore biosynthesis genes in *A. fischeri* (21). At that time, we did not know the

identity of the regulator connecting LuxT to *iucABCD*. Our findings here point to SwrZ

fulfilling that function. To test this idea, we used the CAS assay to measure siderophore

levels in culture fluids from WT, Δ*luxT,* Δ*swrZ,* and Δ*swrZ* Δ*luxT A. fischeri* strains. Indeed,

LuxT activates *A. fischeri* siderophore production only in the *swrZ*⁺ background as

deletion of *luxT* in the Δ*swrZ* strain caused no change in siderophore production (Fig 7D).

Thus, LuxT activates *A. fischeri* siderophore production via a SwrZ-dependent

mechanism. There is no evidence that a second SwrZ-independent mechanism occurs

(Fig 7D). Our model for LuxT and SwrZ regulation of *A. fischeri* siderophore production is

depicted in Fig 7E.

Discussion

QS controls over 600 genes in *V. harveyi* (8). Integral to the *V. harveyi* QS process

are five, largely redundant, Qrr sRNAs that post-transcriptionally regulate target gene

expression. LuxT is a TetR family transcriptional regulator that was previously identified

to repress expression of *V. harveyi qrr*1 encoding the Qrr1 sRNA (18). When LuxT acts

as a *qrr*1 repressor, it tunes the expression of select Qrr1-controlled genes that are

members of the much larger QS regulon. Here, we use RNA-Seq to discover that beyond

controlling *qrr*1 expression, LuxT is a global regulator in *V. harveyi* that controls 414

genes. Analogous to what has been reported for *V. parahaemolyticus* (22), LuxT directly

represses *swrZ,* encoding an additional transcriptional regulator. We show that, in *V.*

harveyi, LuxT activates type III secretion, siderophore production, and aerolysin secretion

genes independently of Qrr1, by both SwrZ-dependent and SwrZ-independent

mechanisms.

Two feedback loops exist in the *V. harveyi* LuxT/SwrZ regulatory pathway: LuxT

and SwrZ activate and repress, respectively, their own transcription. Generally, positive

feedback is thought to amplify responses to a stimuli, whereas negative feedback buffers

pathway output against fluctuations in stimuli (44, 45). Thus, including both positive and

negative feedback in a single system can alter both input sensitivity and output dynamics

(45, 46). We speculate that the LuxT positive feedback loop enables *V. harveyi* to rapidly

commit to the LuxT "ON" state following initial activation of *luxT* expression. By contrast,

SwrZ-directed negative feedback on *swrZ,* which occurs when LuxT is in the "OFF" state,

should prevent runaway *swrZ* expression. Moreover, LuxT and SwrZ have opposing

regulatory functions, so genes activated by LuxT are repressed by SwrZ. Negative SwrZ

autoregulation could, by tamping down SwrZ production, accelerate the expression of

LuxT-activated genes when *V. harveyi* transitions to the LuxT "ON" state.

Our analyses revealed that LuxT activates T3SS, siderophore, and aerolysin

genes by two mechanisms, one SwrZ-dependent and one SwrZ-independent. We

propose two advantages to this regulatory arrangement. First, this dual mechanism may

promote a "switch-like" response. Because SwrZ represses genes that are activated by

LuxT, when LuxT is in the "ON" state, stronger target gene activation will occur if *swrZ* is

repressed by LuxT than if not. Second, if expression and/or activity of either or both LuxT

and SwrZ respond to additional regulatory inputs, linking the two nodes could promote

more nuanced control of gene expression than if LuxT and *swrZ* were not connected in

the circuit.

For *V. harveyi* to transition between LuxT "ON" and LuxT "OFF" states, a cue(s)

must activate *luxT* expression or increase LuxT activity. We do not know the identity of

this putative cue. We do know that LuxT is not regulated by QS and, consistent with this

finding, *luxT* expression remains steady over the *V. harveyi* growth curve (Fig S3B and

S3C). Curiously, however, genes in the LuxT-controlled regulon overlap with those in the

QS-controlled regulon. In particular, LuxT activates expression of type III secretion,

siderophore production, and aerolysin secretion genes, all of which are also QS-

controlled and expressed at LCD, which is the typical pattern for QS-controlled virulence

genes in vibrios (47). The above mentioned QS- and LuxT-regulated traits are all

virulence factors that contribute to the success of vibrios as pathogens in host organisms

(42, 48, 49). For example, in shrimp larvae, *V. harveyi* T3SS gene expression is 1,000-

fold higher than that in laboratory culture (48). Possibly, *luxT* expression or LuxT activity

is modulated by a host factor. Alternatively or in addition, TetR-family members frequently

bind small-molecule ligands (50). It could be that a *V. harveyi-,* host-, or environmentally-

derived ligand promotes heightened LuxT activity. Future studies investigating regulation

of *luxT* expression or LuxT activity should define the connection between the LCD state

and LuxT function.

Our phylogenetic analyses revealed that *luxT,* but not *swrZ,* is conserved among

Vibrionaceae. And, LuxT repression of *swrZ* is conserved at least in *V. parahaemolyticus,*

V. harveyi, and our *A. fischeri* test case (22). Going forward, it will be interesting to learn

how LuxT function has diverged in species that harbor *luxT* but lack *swrZ,* such as in

Vibrio cholerae C6706. RNA-Seq experiments similar to those described here could be

used to begin to parse the roles LuxT plays and, based on the findings, paint a coherent

evolutionary picture of LuxT control of *Vibrionaceae* biology*.*

Materials and Methods

Bacterial strains, culture conditions, and standard methods

All strains are listed in Table S1A. *V. harveyi* strains were derived from *V. harveyi* BB120

(BAA-1116) (51). *A. fischeri* strains were derived from *A. fischeri* ES114 (52). *E. coli* S17–

1 λ*pir* was used for cloning. *E. coli* MG1655 was used for heterologous gene expression.

V. harveyi and *A. fischeri* strains were grown in either Luria Marine (LM) medium or

minimal Autoinducer Bioassay (AB) medium at 30˚C with shaking (53, 54). AB medium

contained 0.4% vitamin-free casamino acids (Difco). *E. coli* strains were grown in LB

medium at 37˚C for cloning or at 30˚C for heterologous gene expression. When

necessary, kanamycin, chloramphenicol, ampicillin, and polymyxin B were added at 100

μg mL^{-1}, 10 μg mL^{-1}, 100 μg mL^{-1}, and 50 μg mL^{-1}, respectively. Gene expression from

P$_{BAD}$ and P$_{tac}$ promoters was induced following addition of 0.01% arabinose and 0.5 mM

isopropyl β-D-1-thiogalactopyranoside (IPTG), respectively, at the time of inoculation.

Procedures for LuxT-6xHis purification, EMSA analyses, qRT-PCR measurements, and

hemolysis assays have been described (18), as have CAS siderophore detection assays

(21). Primers used to amplify DNA probes for EMSA analyses and for qRT-PCR

measurements are listed in Table S1B. Growth conditions for qRT-PCR experiments are

provided in the figure legends.

DNA manipulation and strain construction

Oligonucleotide primers were purchased from Integrated DNA Technologies (IDT) and

are listed in Table S1B. PCR reactions used either KOD Hot Start DNA Polymerase

(Sigma) or iProof DNA Polymerase (Bio-Rad). All cloning was completed using Gibson

Assembly Master Mix (New England Biolabs) for isothermal DNA assembly (55).

Plasmids were verified by sequencing (Genewiz) and are listed in Table S1C. Regarding

nomenclature of our constructs, a capital P designates the promoter driving transcription

(e.g. P*swrZ*-*lux*). Plasmids that promote overexpression of genes are designated with a

lowercase p (e.g. p*swrZ*). The P*swrZ*-*lux* and P*luxT*-*lux* transcriptional reporters included

114 and 531 bp promoter regions, respectively, to drive transcription of *luxCDABEG*. A

consensus ribosome-binding site was included in both reporters to drive translation.

Plasmids were introduced into *E. coli* by electroporation using a Bio-Rad Micro Pulser.

Plasmids were conjugated into *V. harveyi* and *A. fischeri* from *E. coli* S17–1 λ*pir*. *V.*

harveyi exconjugants were selected on agar plates containing polymyxin B, and *A.*

fischeri exconjugants were selected on agar plates containing ampicillin. Chromosomal

alterations in *V. harveyi* and *A. fischeri* were introduced using the pRE112 vector

harboring the *sacB* counter-selectable marker as previously described (18, 56, 57).

Mutations were verified by PCR and/or sequencing.

RNA-Sequencing

Cells from overnight cultures of WT and Δ*luxT V. harveyi* strains grown in LM medium

were pelleted by centrifugation at 21,100 x *g* (Eppendorf 5424) and resuspended in fresh

LM medium. Fresh cultures containing 25 mL of LM medium were inoculated with the

washed cells at OD_{600} = 0.005. The cultures were incubated at 30˚C with shaking, and

RNA was isolated from 3 biological replicates of each strain when the cultures had

reached OD_{600} = 0.1 using the RNeasy mini kit (Qiagen #74106). RNA-Seq was

performed at the Genomics Core Facility at Princeton University, as previously described

(58). Reads were mapped to the *V. harveyi* BB120 (BAA-1116) genome using TopHat

(59). Genes with differential expression were identified using DESeq2 (60) and those

exhibiting a $\log_2$ fold-change in expression > 1, along with a *p*-value < 0.01, in the $\Delta luxT$

strain compared to WT *V. harveyi,* were designated as members of the LuxT regulon.

Bioluminescence assays

Cells in overnight cultures of *E. coli* harboring the the P_{swrZ}-*lux* reporter or *V. harveyi*

harboring the P_{luxT}-*lux* reporter were pelleted by centrifugation at 21,100 x *g* (Eppendorf

5424) and resuspended in LB or LM medium, respectively. The *E. coli* and *V. harveyi*

cells were inoculated into fresh LB and LM, respectively, with normalization to a starting

OD_{600} of 0.005. 150 µL of each culture was transferred to wells of a 96-well plate (Corning)

in quadruplicate technical replicates and overlayed with 50 µL of mineral oil (Sigma). The

plates were incubated with shaking at 30°C, and bioluminescence and OD_{600} were

measured every 15 min for 24 h using a BioTek Synergy Neo2 Multi-Mode Reader

(BioTek, Winooski, VT, USA). Relative light units (RLU: bioluminescence/OD_{600})

represent the values when each sample was at OD_{600} = 1.

VopD Western blot analyses

Cytoplasmic VopD levels were detected in *V. harveyi* cultures that had been grown in AB

medium supplemented with 5 mM EGTA as previously described (5). Cells equivalent to

1 OD_{600} were pelleted by centrifugation at 21,100 x *g* (Eppendorf 5424), resuspended in

100 µL of SDS-PAGE sample buffer (Bio-Rad), and boiled for 10 min. Samples were

loaded onto 4-20% Mini-PROTEAN TGX Gels (Bio-Rad) and subjected to electrophoresis

for 30 min at 50 mA. Following Western transfer, nitrocellulose membranes were cut in

half. One portion was probed with an antibody against VopD, and the other portion was

probed with an antibody against the LuxS control, as previously described (5). Anti-rabbit

IgG (H+L) HRP Conjugate (Promega) was used as the secondary antibody. Proteins were

visualized using SuperSignal West Femto Maximum Sensitivity Substrate (Thermo

Fisher) and an ImageQuant LAS 4000 imager.

Phylogenetic analyses

Genomic DNA sequences of 418 *Vibrionaceae* species were downloaded from the

GenBank database (61). Genes encoding *luxT* among *Vibrionaceae* species were

identified previously (18). A custom MATLAB (Mathworks, 2020) search algorithm based

on protein sequence similarity was used to identify genes encoding *swrZ*. Briefly, protein

sequences of SwrZ from *V. harveyi* ATCC BAA-1116 and *V. parahaemolyticus* RIMD

2210633 were used as queries. Both yielded similar search results. The chromosomes or

contigs of species under consideration were first converted to amino acid sequences, and

subsequently scanned for regions similar to the query sequences. To identify regions of

highest similarity between protein sequences, local sequence alignments were performed

using the Smith-Waterman (SW) algorithm (62). The standard substitution matrix

BLOSUM62 (https://ftp.ncbi.nih.gov/blast/matrices/) was used to compute similarity

scores S, which considers both the length and sequence similarity of the alignment. A

cut-off value $S > 150$ was used to identify putative genes encoding SwrZ. Genes identified

as *swrZ* homologs were verified to encode GntR family transcriptional regulators. Species

lacking *swrZ* were excluded from further phylogenetic analyses. The protocols used to

perform phylogenetic analyses and tree building have been described previously (18).

Acknowledgments

RNA-Sequencing was performed at the Genomics Core Facility in the Lewis-Sigler

Institute for Integrative Genomics at Princeton University. This work was supported by the

Howard Hughes Medical Institute, National Institutes of Health (NIH) Grant

5R37GM065859, and National Science Foundation Grant MCB-1713731 (to BLB). MJE

was supported by the NIH graduate training grant (NIGMS T32GM007388). The funders

had no role in study design, data collection and interpretation, or the decision to submit

the work for publication. The authors have no competing interests to declare. We thank

members of the Bassler laboratory for insightful discussions and ideas. We lovingly

acknowledge Jian-Ping Cong and her fundamental contributions to the work reported

here. We dedicate this manuscript to her memory. We are grateful for her decades of

devotion to advancing research in the bacterial cell-cell communication field and for her

unflagging kindness to and support of former and current Bassler laboratory members.

Main Figures

Figure 1

Fig 1. LuxT is a global transcriptional regulator in *V. harveyi* that directly represses *swrZ*. (A) Volcano plot depicting RNA-Seq data comparing the transcriptomes of WT and Δ*luxT V. harveyi*. Each data point represents one *V. harveyi* gene. FC designates fold-change. 243 genes were significantly activated (black) and 171 genes were significantly repressed by LuxT (dark gray). **(B)** qRT-PCR measurements of *swrZ* transcript levels in WT (black) and Δ*luxT* (blue) *V. harveyi*. RNA was isolated from strains grown in LM to OD_{600} = 0.1 **(C)** Activity of a plasmid-borne P_{swrZ}-*lux* transcriptional reporter in recombinant *E. coli*. pControl denotes that the second introduced plasmid is the empty parent vector and p*luxT* designates that the second vector harbors arabinose-inducible *luxT*. Strains were grown in LB to OD_{600} = 1 in the absence (black) or presence (gray) of 0.01% arabinose. **(D)** qRT-PCR measurements of *luxT* transcript levels in WT (black) and Δ*swrZ* (red) *V. harveyi*. Growth conditions as in B. **(E)** Activity of a plasmid-borne P_{luxT}-

lux transcriptional reporter in Δ*luxA* (black) and Δ*luxA* Δ*luxT* (blue) *V. harveyi* strains. pControl and p*luxT* denote the empty parent vector and a vector harboring IPTG-inducible *luxT,* respectively. Strains were grown in LM supplemented with 0.5 mM IPTG to OD_{600} = 1. **(F)** Activity of a plasmid-borne P$_{swrZ}$-*lux* transcriptional reporter in recombinant *E. coli.* pControl denotes that the second introduced plasmid is the empty parent vector and p*swrZ* designates that the second vector harbors IPTG-inducible *swrZ.* Strains were grown in LB to OD_{600} = 1 in the absence (black) or presence (gray) of 0.5 mM IPTG. **(G)** Model for the LuxT/SwrZ regulatory circuit. In C, E, and F, RLU denotes relative light units. In B-F, error bars represent standard deviations of the means of n = 3 biological replicates. Unpaired two-tailed *t* tests with Welch's correction were performed comparing the indicated conditions. *p*-values: ns ≥ 0.05, ** < 0.01, **** < 0.0001.

Figure 2

Fig 2. QS and LuxT/SwrZ regulate *V. harveyi* T3SS gene expression. (A) Schematic of T3SS gene organization in *V. harveyi*. There are four major T3SS operons, as labeled. All genes shown in blue were identified by RNA-Seq as members of the LuxT regulon. The numbers shown within select genes designate the fold-change differences in transcript levels between WT and Δ*luxT V. harveyi*, as measured by RNA-Seq (Data Set S1). Expression of underlined genes was measured in panels B and C. **(B)** qRT-PCR measurements of mRNA levels of the indicated genes in WT (black), Δ*luxO* (green), and *luxO* D61E (orange) *V. harveyi* strains. RNA was isolated from strains grown in AB to $OD_{600} = 1$. **(C)** qRT-PCR measurements of mRNA levels of the indicated genes in WT (black), Δ*luxT* (blue), Δ*swrZ* (red), and Δ*swrZ* Δ*luxT* (purple) *V. harveyi* strains. RNA was isolated from strains grown in LM to $OD_{600} = 0.1$. In B and C, error bars represent standard deviations of the means of $n = 3$ biological replicates. Different letters indicate significant differences between strains, $p < 0.05$ (two-way analysis of variation (ANOVA) followed by Tukey's multiple comparisons test).

Figure 3

Fig 3. LuxT activates *V. harveyi* type III secretion gene expression by two mechanisms. (A) Western blot of cytoplasmic VopD levels in the indicated *V. harveyi* strains at HCD. "LCD-locked" denotes that the parent strain for the samples in the right four lanes harbors the *luxO* D61E mutation. Detection of LuxS protein serves as the loading control, as previously described (5). **(B-E)** Schematics depicting the outcomes for LuxT regulation of T3SS genes by SwrZ-dependent and SwrZ-independent mechanisms for WT **(B),** Δ*luxT* **(C),** Δ*swrZ* **(D)**, and Δ*swrZ* Δ*luxT* **(E)** scenarios. See text for details.

Figure 4

Fig 4. LuxT activates *V. harveyi exsA*, **and in turn, T3SS genes, by SwrZ-dependent and SwrZ-independent mechanisms. (A)** qRT-PCR measurements of transcript levels of the indicated genes in Δ*luxR* Δ*swrZ* Δ*luxT V. harveyi* harboring the indicated plasmid. pControl (black) denotes the empty parent vector, and p*luxT* (blue) and p*swrZ* (red) denote vectors harboring IPTG-inducible *luxT* and *swrZ,* respectively. RNA was isolated from strains grown for 16 h in AB supplemented with 0.5 mM IPTG. **(B)** As in A, for Δ*luxR* Δ*swrZ V. harveyi* harboring the indicated plasmid. **(C)** qRT-PCR measurements of transcript levels of the indicated genes in WT (black), Δ*luxT* (blue), Δ*exsA* (gray), and Δ*exsA* Δ*luxT* (white) *V. harveyi* strains. RNA was isolated from *V. harveyi* strains grown in LM to OD_{600} = 0.1. **(D)** Model for QS and LuxT/SwrZ regulation of T3SS genes. In A-C, error bars represent standard deviations of the means of n = 3 biological replicates. In A and C, different letters indicate significant differences between strains, $p < 0.05$ (two-way ANOVA followed by Tukey's multiple comparisons test). In B, unpaired two-tailed t tests with Welch's correction were performed comparing the indicated pControl and p*swrZ* conditions. p-value: * < 0.05.

Fig 5. QS, LuxT, and SwrZ regulate *V. harveyi* siderophore production. (A-B) Schematic of siderophore gene organization in *V. harveyi* for amphi-enterobactin (A) and anguibactin (B). All genes labeled in blue were identified by RNA-Seq as members of the LuxT regulon. The numbers shown within or below the genes designate the fold-change differences in transcript levels between WT and Δ*luxT V. harveyi*, as measured by RNA-Seq. Expression of underlined genes was measured in panels D, F, and G. **(C)** CAS assay quantitation of siderophore levels in cell-free culture fluids isolated from the indicated *V. harveyi* strains. Strains were grown for 16 h in AB medium. **(D)** qRT-PCR measurements of transcript levels of the indicated genes in WT (black), Δ*luxT* (blue), Δ*swrZ* (red), and Δ*swrZ* Δ*luxT* (purple) *V. harveyi* strains. RNA was isolated from strains

grown in LM to OD_{600} = 0.1. **(E)** As in C. "LCD-locked" denotes that the strains harbor the *luxO* D61E mutation. **(F)** qRT-PCR measurements of transcript levels of the indicated genes in Δ*luxR* Δ*swrZ* Δ*luxT V. harveyi* harboring the indicated plasmid. pControl (black) denotes the empty parent vector, and p*luxT* (blue) and p*swrZ* (red) denote vectors harboring IPTG-inducible *luxT* and *swrZ,* respectively. RNA was isolated from strains grown for 16 h in AB supplemented with 0.5 mM IPTG. **(G)** As in F, for Δ*luxR* Δ*swrZ V. harveyi* harboring the indicated plasmid. **(H)** Model for Fur, QS, and LuxT/SwrZ regulation of *V. harveyi* siderophore production. In C-G, error bars represent standard deviations of the means of *n* = 3 biological replicates. In C-F, different letters indicate significant differences between strains, *p* < 0.05 (two-way ANOVA followed by Tukey's multiple comparisons test). In G, unpaired two-tailed *t* tests with Welch's correction were performed comparing the indicated pControl and p*swrZ* conditions. *p*-values: * < 0.05, ** < 0.01.

Figure 6

Fig 6. QS, LuxT, and SwrZ regulate *V. harveyi* aerolysin secretion. (A) Schematic of aerolysin gene organization in *V. harveyi*. All genes labeled in blue were identified by RNA-Seq as members of the LuxT regulon. The numbers shown within the genes designate the fold-change differences in transcript levels between WT and Δ*luxT V. harveyi*, as measured by RNA-Seq. Expression of underlined genes was measured in panels C, E, and F. **(B)** Hemolytic activity in the indicated *V. harveyi* cell-free culture fluids as judged by lysis of defibrinated sheep's blood cells. Culture fluids were collected after 24 h of growth in AB medium. **(C)** qRT-PCR measurements of transcript levels of the indicated genes in WT (black), Δ*luxT* (blue), Δ*swrZ* (red), and Δ*swrZ* Δ*luxT* (purple) *V. harveyi* strains. RNA was isolated from strains grown in LM to $OD_{600} = 0.1$. **(D)** As in B for the designated strains. **(E)** qRT-PCR measurements of transcript levels of the indicated genes in Δ*luxR* Δ*swrZ* Δ*luxT V. harveyi* harboring the indicated plasmid. pControl (black) denotes the empty parent vector, and p*luxT* (blue) and p*swrZ* (red) denote vectors harboring IPTG-inducible *luxT* and *swrZ,* respectively. RNA was isolated from strains grown for 16 h in AB supplemented with 0.5 mM IPTG. **(F)** As in E, for Δ*luxR* Δ*swrZ V. harveyi* harboring the indicated plasmid. **(G)** Model for LuxT/SwrZ regulation of aerolysin

production. In B-F, error bars represent standard deviations of the means of $n = 3$ biological replicates. In B-E, different letters indicate significant differences between strains, $p < 0.05$ (two-way ANOVA followed by Tukey's multiple comparisons test). In F, unpaired two-tailed t tests with Welch's correction were performed comparing the indicated pControl and p*swrZ* conditions. *p*-values: ns ≥ 0.05, * < 0.05, ** < 0.01.

Figure 7

Fig 7. Co-occurrence of *luxT* and *swrZ* genes among *Vibrionaceae* species. **(A)** Highest similarity in protein sequences (BLOSUM62) in the indicated genera to *V. parahaemolyticus* SwrZ. Species are divided into two groups, one possessing *luxT* (*luxT*+) and one lacking *luxT* (*luxT*−). The dashed line indicates the cutoff used for the similarity score to differentiate *swrZ*+ from *swrZ*− strains. Gray indicates groups of species that lack *swrZ*. The numbers in parentheses indicate the number of species in each group. Boxes show the means ± SDs for groups with > 5 species. Circles represent individual species in groups with ≤ 5 members. Purple, green, and orange denote, respectively, *Aliivibrio*, *Photobacterium*, and *Vibrio* species that possess *swrZ*. **(B)** Phylogenetic tree based on the *swrZ* promoter regions in *Vibrionaceae*. Phylogenetic distances were computed based on sequences spanning -122 to -8, relative to the *swrZ* start codons. Colors as in panel A. **(C)** qRT-PCR measurements of *swrZ* transcript levels in WT (black) and Δ*luxT* (blue) *A. fischeri*. RNA was isolated from strains grown in LM to OD_{600} = 0.1. **(D)** CAS assay quantitation of siderophore levels in cell-free culture fluids prepared from the indicated *A. fischeri* strains. Strains were grown for 16 h in AB medium. **(E)** Model for LuxT/SwrZ regulation of *A. fischeri* aerobactin siderophore production. In C and D, error bars represent standard deviations of the means of n = 3 biological replicates. In C, an unpaired two-tailed t test with Welch's correction was performed comparing the indicated WT and Δ*luxT* conditions. p-value: ** < 0.01. In D, different letters indicate significant differences between strains, p < 0.05 (two-way ANOVA followed by Tukey's multiple comparisons test).

References

1. Bervoets I, Charlier D. 2019. Diversity, versatility and complexity of bacterial gene regulation mechanisms: opportunities and drawbacks for applications in synthetic biology. FEMS Microbiol Rev 43:304–339.

2. Gottesman S. 2019. Trouble is coming: Signaling pathways that regulate general stress responses in bacteria. J Biol Chem 294:11685–11700.

3. Papenfort K, Bassler BL. 2016. Quorum sensing signal-response systems in Gram-negative bacteria. Nat Rev Microbiol 14:576–588.

4. Waters CM, Bassler BL. 2005. Quorum sensing: cell-to-cell communication in bacteria. Annu Rev Cell Dev Biol 21:319–346.

5. Henke JM, Bassler BL. 2004. Quorum sensing regulates type III secretion in *Vibrio harveyi* and *Vibrio parahaemolyticus*. J Bacteriol 186:3794–3805.

6. Lilley BN, Bassler BL. 2000. Regulation of quorum sensing in *Vibrio harveyi* by LuxO and Sigma-54. Mol Microbiol 36:940–954.

7. Henke JM, Bassler BL. 2004. Three parallel quorum-sensing systems regulate gene expression in *Vibrio harveyi*. J Bacteriol 186:6902–6914.

8. van Kessel JC, Rutherford ST, Shao Y, Utria AF, Bassler BL. 2013. Individual and combined roles of the master regulators AphA and LuxR in control of the *Vibrio harveyi* quorum-sensing regulon. J Bacteriol 195:436–443.

9.   Lenz DH, Mok KC, Lilley BN, Kulkarni RV, Wingreen NS, Bassler BL. 2004. The small RNA chaperone Hfq and multiple small RNAs control quorum sensing in *Vibrio harveyi* and *Vibrio cholerae*. Cell 118:69–82.

10. Tu KC, Bassler BL. 2007. Multiple small RNAs act additively to integrate sensory information and control quorum sensing in *Vibrio harveyi*. Genes Dev 21:221–233.

11. Shao Y, Bassler BL. 2012. Quorum-sensing non-coding small RNAs use unique pairing regions to differentially control mRNA targets. Mol Microbiol 83:599–611.

12. Rutherford ST, van Kessel JC, Shao Y, Bassler BL. 2011. AphA and LuxR/HapR reciprocally control quorum sensing in vibrios. Genes Dev 25:397–408.

13. Shao Y, Feng L, Rutherford ST, Papenfort K, Bassler BL. 2013. Functional determinants of the quorum-sensing non-coding RNAs and their roles in target regulation. EMBO J 32:2158–2171.

14. Tu KC, Long T, Svenningsen SL, Wingreen NS, Bassler BL. 2010. Negative feedback loops involving small regulatory RNAs precisely control the *Vibrio harveyi* quorum-sensing response. Mol Cell 37:567–579.

15. Teng S-W, Schaffer JN, Tu KC, Mehta P, Lu W, Ong NP, Bassler BL, Wingreen NS. 2011. Active regulation of receptor ratios controls integration of quorum-sensing signals in *Vibrio harveyi*. Mol Syst Biol 7:491.

16.   Freeman JA, Lilley BN, Bassler BL. 2000. A genetic analysis of the functions of LuxN: a two-component hybrid sensor kinase that regulates quorum sensing in *Vibrio harveyi*. Mol Microbiol 35:139–149.

17.   Freeman JA, Bassler BL. 1999. A genetic analysis of the function of LuxO, a two-component response regulator involved in quorum sensing in *Vibrio harveyi*. Mol Microbiol 31:665–677.

18.   Eickhoff MJ, Fei C, Huang X, Bassler BL. 2021. LuxT controls specific quorum-sensing-regulated behaviors in *Vibrionaceae* spp. via repression of *qrr*1, encoding a small regulatory RNA. PLoS Genet 17:e1009336.

19.   Lin YH, Miyamoto C, Meighen EA. 2000. Purification and characterization of a luxO promoter binding protein LuxT from *Vibrio harveyi*. Protein Expr Purif 20:87–94.

20.   Lin YH, Miyamoto C, Meighen EA. 2000. Cloning and functional studies of a luxO regulator LuxT from *Vibrio harveyi*. Biochim Biophys Acta 1494:226–235.

21.   Eickhoff MJ, Bassler BL. 2020. *Vibrio fischeri* siderophore production drives competitive exclusion during dual-species growth. Mol Microbiol 114:244–261.

22.   Jaques S, McCarter LL. 2006. Three new regulators of swarming in *Vibrio parahaemolyticus*. J Bacteriol 188:2625–2635.

23.   Liu H, Gu D, Cao X, Liu Q, Wang Q, Zhang Y. 2012. Characterization of a new quorum sensing regulator *luxT* and its roles in the extracellular protease production,

motility, and virulence in fish pathogen *Vibrio alginolyticus*. Arch Microbiol 194:439–452.

24. Green ER, Mecsas J. 2016. Bacterial Secretion Systems: An Overview. Microbiol Spectr 4.

25. Deng W, Marshall NC, Rowland JL, McCoy JM, Worrall LJ, Santos AS, Strynadka NCJ, Finlay BB. 2017. Assembly, structure, function and regulation of type III secretion systems. Nat Rev Microbiol 15:323–337.

26. Galán JE, Lara-Tejero M, Marlovits TC, Wagner S. 2014. Bacterial type III secretion systems: specialized nanomachines for protein delivery into target cells. Annu Rev Microbiol 68:415–438.

27. Morot A, Fekih El S, Bidault A, Le Ferrand A, Jouault A, Kavousi J, Bazire A, Pichereau V, Dufour A, Paillard C, Delavat F. 2021. Virulence of *Vibrio harveyi* ORM4 towards the European abalone *Haliotis tuberculata* involves both quorum sensing and a type III secretion system. Environ Microbiol 23:5273–5288.

28. Park K-S, Ono T, Rokuda M, Jang M-H, Okada K, Iida T, Honda T. 2004. Functional characterization of two type III secretion systems of *Vibrio parahaemolyticus*. Infect Immun 72:6659–6665.

29. Zeb S, Shah MA, Yasir M, Awan HM, Prommeenate P, Klanchui A, Wren BW, Thomson N, Bokhari H. 2019. Type III secretion system confers enhanced virulence in clinical non-O1/non-O139 *Vibrio cholerae*. Microb Pathog 135:103645.

30. Waters CM, Wu JT, Ramsey ME, Harris RC, Bassler BL. 2010. Control of the type 3 secretion system in *Vibrio harveyi* by quorum sensing through repression of ExsA. Appl Environ Microbiol 76:4996–5004.

31. Neilands JB. 1995. Siderophores: structure and function of microbial iron transport compounds. J Biol Chem 270:26723–26726.

32. Zane HK, Naka H, Rosconi F, Sandy M, Haygood MG, Butler A. 2014. Biosynthesis of amphi-enterobactin siderophores by *Vibrio harveyi* BAA-1116: identification of a bifunctional nonribosomal peptide synthetase condensation domain. J Am Chem Soc 136:5615–5618.

33. Naka H, Actis LA, Crosa JH. 2013. The anguibactin biosynthesis and transport genes are encoded in the chromosome of *Vibrio harveyi*: a possible evolutionary origin for the pJM1 plasmid–encoded system of *Vibrio anguillarum*? MicrobiologyOpen 2:182–194.

34. McRose DL, Baars O, Seyedsayamdost MR, Morel FMM. 2018. Quorum sensing and iron regulate a two-for-one siderophore gene cluster in *Vibrio harveyi*. Proc Natl Acad Sci USA 115:7581–7586.

35. Di Lorenzo M, Poppelaars S, Stork M, Nagasawa M, Tolmasky ME, Crosa JH. 2004. A nonribosomal peptide synthetase with a novel domain organization is essential for siderophore biosynthesis in *Vibrio anguillarum*. J Bacteriol 186:7327–7336.

36.   Naka H, López CS, Crosa JH. 2010. Role of the pJM1 plasmid-encoded transport proteins FatB, C and D in ferric anguibactin uptake in the fish pathogen *Vibrio anguillarum*. Environ Microbiol Rep 2:104–111.

37.   Tolmasky ME, Actis LA, Crosa JH. 1988. Genetic analysis of the iron uptake region of the *Vibrio anguillarum* plasmid pJM1: molecular cloning of genetic determinants encoding a novel trans activator of siderophore biosynthesis. J Bacteriol 170:1913–1919.

38.   Bagg A, Neilands JB. 1987. Ferric uptake regulation protein acts as a repressor, employing iron (II) as a cofactor to bind the operator of an iron transport operon in *Escherichia coli*. Biochemistry 26:5471–5477.

39.   Wretlind B, Möllby R, Wadström T. 1971. Separation of two hemolysins from *Aeromonas hydrophila* by isoelectric focusing. Infect Immun 4:503–505.

40.   Bernheimer AW, Avigad LS, Avigad G. 1975. Interactions between aerolysin, erythrocytes, and erythrocyte membranes. Infect Immun 11:1312–1319.

41.   Podobnik M, Kisovec M, Anderluh G. 2017. Molecular mechanism of pore formation by aerolysin-like proteins. Philos Trans R Soc Lond, B, Biol Sci 372.

42.   Macpherson HL, Bergh Ø, Birkbeck TH. 2012. An aerolysin-like enterotoxin from *Vibrio splendidus* may be involved in intestinal tract damage and mortalities in turbot, *Scophthalmus maximus* (L.), and cod, *Gadus morhua* L., larvae. J Fish Dis 35:153–167.

43. Rasmussen-Ivey CR, Figueras MJ, McGarey D, Liles MR. 2016. Virulence Factors of *Aeromonas hydrophila*: In the Wake of Reclassification. Front Microbiol 7:1337.

44. Mitrophanov AY, Groisman EA. 2008. Positive feedback in cellular control systems. Bioessays 30:542–555.

45. Rao SD, Igoshin OA. 2021. Overlaid positive and negative feedback loops shape dynamical properties of PhoPQ two-component system. PLoS Comput Biol 17:e1008130.

46. Mitarai N, Jensen MH, Semsey S. 2015. Coupled positive and negative feedbacks produce diverse gene expression patterns in colonies. mBio 6: e00059-15.

47. Miller MB, Skorupski K, Lenz DH, Taylor RK, Bassler BL. 2002. Parallel quorum sensing systems converge to regulate virulence in *Vibrio cholerae*. Cell 110:303–314.

48. Ruwandeepika HAD, Karunasagar I, Bossier P, Defoirdt T. 2015. Expression and quorum sensing regulation of type III secretion system genes of *Vibrio harveyi* during infection of gnotobiotic brine shrimp. PLoS ONE 10:e0143935.

49. Balado M, Lages MA, Fuentes-Monteverde JC, Martínez-Matamoros D, Rodríguez J, Jiménez C, Lemos ML. 2018. The siderophore piscibactin Is a relevant virulence factor for *Vibrio anguillarum* favored at low temperatures. Front Microbiol 9:1766.

50. Cuthbertson L, Nodwell JR. 2013. The TetR family of regulators. Microbiol Mol Biol Rev 77:440–475.

51. Bassler BL, Greenberg EP, Stevens AM. 1997. Cross-species induction of luminescence in the quorum-sensing bacterium *Vibrio harveyi*. J Bacteriol 179:4043–4045.

52. Boettcher KJ, Ruby EG. 1990. Depressed light emission by symbiotic *Vibrio fischeri* of the sepiolid squid *Euprymna scolopes*. J Bacteriol 172:3701–3706.

53. Bassler BL, Wright M, Silverman MR. 1994. Multiple signalling systems controlling expression of luminescence in *Vibrio harveyi*: sequence and function of genes encoding a second sensory pathway. Mol Microbiol 13:273–286.

54. Greenberg EP, Hastings JW, Ulitzur S. 1979. Induction of luciferase synthesis in *Beneckea harveyi* by other marine bacteria. Arch Microbiol 120:87–91.

55. Gibson DG, Young L, Chuang RY, Venter JC, Hutchison CA, Smith HO. 2009. Enzymatic assembly of DNA molecules up to several hundred kilobases. Nat Methods 6:343–345.

56. Edwards RA, Keller LH, Schifferli DM. 1998. Improved allelic exchange vectors and their use to analyze 987P fimbria gene expression. Gene 207:149–157.

57. Chaparian RR, Olney SG, Hustmyer CM, Rowe-Magnus DA, van Kessel JC. 2016. Integration host factor and LuxR synergistically bind DNA to coactivate quorum-sensing genes in *Vibrio harveyi*. Mol Microbiol 101:823–840.

58. Mukherjee S, Moustafa D, Smith CD, Goldberg JB, Bassler BL. 2017. The RhlR quorum-sensing receptor controls *Pseudomonas aeruginosa* pathogenesis and

biofilm development independently of its canonical homoserine lactone autoinducer. PLoS Pathog 13:e1006504.

59. Kim D, Pertea G, Trapnell C, Pimentel H, Kelley R, Salzberg SL. 2013. TopHat2: accurate alignment of transcriptomes in the presence of insertions, deletions and gene fusions. Genome Biol 14:R36–13.

60. Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. Genome Biol 15:550–21.

61. Benson DA, Karsch-Mizrachi I, Lipman DJ, Ostell J, Rapp BA, Wheeler DL. 2000. GenBank. Nucleic Acids Res 28:15–18.

62. Smith TF, Waterman MS. 1981. Identification of common molecular subsequences. J Mol Biol 147:195–197.

Supplemental Material

Figure S1

Fig S1. Simplified _V. harveyi_ QS pathway. (A) LCD and **(B)** HCD _V. harveyi_ QS circuit. Circles, triangles, and squares represent AIs. See Introduction in the main text for details.

Figure S2.

A

LuxT-6xHis
(Dimer):   - 0.5x 1x 2x 4x 8x 16x - 0.5x 1x 2x 4x 8x 16x

Unbound

P_{swrZ} Control

B

LuxT-6xHis
(Dimer):   - 0.5x 1x 2x 4x 8x 16x - 0.5x 1x 2x 4x 8x 16x

Unbound

P_{luxT} Control

Fig S2. LuxT binds to the *V. harveyi swrZ* and *luxT* promoters. (A) EMSA showing binding of LuxT-6xHis to a 113 bp DNA fragment containing the *swrZ* promoter (left) and a 107 bp control fragment containing the *V. harveyi luxC* promoter (right). Previously, it was shown that LuxT does not bind the *luxC* promoter (18). 20 nM DNA probe was incubated with the indicated relative concentrations of the LuxT-6xHis dimer: - indicates no protein, 1x indicates 20 nM, and 16x indicates 320 nM. **(B)** As in A, for a 99 bp DNA fragment containing the *luxT* promoter (left) and a 107 bp control fragment containing the *V. harveyi luxC* promoter (right).

Figure S3

Fig S3. SwrZ does not regulate *luxT,* and QS does not regulate *luxT* or *swrZ* in *V. harveyi.* (A) qRT-PCR measurements of *luxT* and *swrZ* transcript levels in WT *V. harveyi* harboring the following plasmids: pControl (black) denotes the empty parent vector and p*swrZ* (red) denotes a vector carrying IPTG-inducible *swrZ.* RNA was isolated from strains grown in LM containing 0.5 mM IPTG to OD_{600} = 1. **(B)** qRT-PCR measurements of *luxT* and *swrZ* transcript levels in WT (black), Δ*luxO* (green), and *luxO* D61E (orange) *V. harveyi* strains. RNA was isolated from strains grown in AB to OD_{600} = 1. **(C)** qRT-PCR measurements of *luxT* transcript levels over growth. RNA was isolated from WT *V. harveyi* grown in LM to different cell densities, as indicated by the OD_{600} values on the x-axis. **(D)** As in C, except *swrZ* transcript levels were measured in WT (black) and Δ*luxT* (blue) *V. harveyi* strains. In all panels, error bars represent standard deviations of the means of *n* = 3 biological replicates. Standard deviations that are smaller than the symbols are not shown. In A, unpaired two-tailed *t* tests with Welch's correction were performed comparing the pControl and p*swrZ* conditions for each gene. *p*-values: ns ≥ 0.05, *** < 0.001. In B, different letters indicate significant differences between strains, $p < 0.05$ (two-way analysis of variation (ANOVA) followed by Tukey's multiple comparisons test).

Figure S4

Fig S4. LuxT activates T3SS, siderophore, and aerolysin genes by Qrr1-independent mechanisms in *V. harveyi*. qRT-PCR measurements of transcript levels of the indicated genes in WT (black) and Δ*luxT* (blue) *V. harveyi* **(A, C,** and **E)** or in Δ*qrr*1 (black) and Δ*qrr*1 Δ*luxT* (blue) *V. harveyi* strains **(B, D,** and **F)**. RNA was isolated from strains grown in LM to OD_{600} = 0.1. In all panels, error bars represent standard deviations of the means of *n* = 3 biological replicates. Unpaired two-tailed *t* tests with Welch's correction were performed comparing two conditions, as indicated. *p*-values: * <0.05, ** < 0.01, *** < 0.001, **** < 0.0001.

Figure S5

Fig S5. *V. harveyi* **T3SS genes are repressed at HCD by LuxR. (A)** qRT-PCR measurements of transcript levels of the indicated T3SS genes in WT (black) and Δ*luxR* (gray) *V. harveyi* strains. RNA was isolated from strains grown in AB to OD$_{600}$ = 1. **(B)** As in A, except transcript levels were measured in *luxO* D61E (black) and *luxO* D61E Δ*aphA* (gray) *V. harveyi* strains. In both panels, error bars represent standard deviations of the means of *n* = 3 biological replicates. Unpaired two-tailed *t* tests with Welch's correction were performed comparing two conditions, as indicated. *p*-values: ns ≥ 0.05, * < 0.05, ** < 0.01.

Figure S6

Fig S6. LuxR and Fur repress production of the *V. harveyi* amphi-enterobactin siderophore. (A-C) CAS assay quantitation of siderophore levels in cell-free culture fluids isolated from the indicated *V. harveyi* strains. Strains were grown for 16 h in AB. **(D)** qRT-PCR measurements of transcript levels of the indicated genes in WT (black), Δ*fur* (gray), and Δ*fur* Δ*luxT* (blue) *V. harveyi* strains. Strains were grown in LM to OD_{600} = 0.1. **(E)** As in D for WT (black), Δ*luxT* (blue), Δ*swrZ* (red), and Δ*swrZ* Δ*luxT* (purple) *V. harveyi* strains. In all panels, error bars represent standard deviations of the means of n = 3 biological replicates. In A, D, and E, different letters indicate significant differences between strains, $p < 0.05$ (two-way ANOVA followed by Tukey's multiple comparisons test). In B and C, unpaired two-tailed t tests with Welch's correction were performed comparing the two conditions, as indicated. p-values: ns ≥ 0.05, ** < 0.01.

Figure S7

Fig S7. Assessment of the requirements for *V. harveyi* **aerolysin secretion. (A, B, and D)** Hemolytic activity present in the indicated *V. harveyi* cell-free culture fluids as judged by lysis of defibrinated sheep's blood cells. Culture fluids were collected after 24 h of growth in AB medium. **(C, E,** and **F)** Hemolytic activity produced by the indicated *V. harveyi* strains as judged by halo formation on TSA plates containing 5% sheep's blood. Strains were grown for 72 h at 30°C. In A, B, and D, error bars represent standard deviations of the means of $n = 3$ biological replicates. In A and B, unpaired two-tailed *t* tests with Welch's correction were performed comparing two conditions, as indicated. *p*-values: ns $\geq$ 0.05. In D, different letters indicate significant differences between strains, $p < 0.05$ (two-way ANOVA followed by Tukey's multiple comparisons test).

Figure S8

Fig S8. LuxT regulates genes by SwrZ-dependent and SwrZ-independent mechanisms in *V. harveyi.* qRT-PCR measurements of transcript levels of the indicated genes in WT (black), Δ*luxT* (blue), Δ*swrZ* (red), and Δ*swrZ* Δ*luxT* (purple) *V. harveyi* strains. Strains were grown in LM to $OD_{600} = 0.1$. Error bars represent standard deviations of the means of $n = 3$ biological replicates. Different letters indicate significant differences between strains, $p < 0.05$ (two-way ANOVA followed by Tukey's multiple comparisons test).

Data Set S1. The *V. harveyi* LuxT regulon. Genes activated (Sheet 1) and repressed (Sheet 2) by LuxT in *V. harveyi.* FC designates fold-change in gene expression in the Δ*luxT* strain compared to WT.

Table S1. Strains, primers, and plasmids used in this study. (A) Strain list, **(B)** primer list, and **(C)** plasmid list.

Table S1

A. Strains used in this study

(WT strains are depicted in bold with variants listed below them.)

Strain	Relevant genotype or feature	Reference
V. harveyi **BB120**	WT	(1)
ME287	Δ*luxT*	(2)
ME291	Δ*swrZ*	(2)
ME1057	Δ*luxA*	This study
ME1056	Δ*luxA* Δ*luxT*	This study
BB721	Δ*luxO*	(3)
JSV780	*luxO* D61E	(4)
KT39	Δ*qrr*1	(5)
ME431	Δ*qrr*1 Δ*luxT*	(2)
KM669	Δ*luxR*	(6)
STR416	*luxO* D61E Δ*aphA*	(7)
ME302	Δ*swrZ* Δ*luxT*	(2)
ME181	*luxO* D61E Δ*luxT*	(2)
ME805	*luxO* D61E Δ*swrZ*	This study
ME807	*luxO* D61E Δ*swrZ* Δ*luxT*	This study
ME944	Δ*luxR* Δ*swrZ* Δ*luxT*	This study
ME942	Δ*luxR* Δ*swrZ*	This study
ME1044	Δ*exsA*	This study
ME1032	Δ*exsA* Δ*luxT*	This study
ME132	*luxO* D61E Δ*aebF*	This study
ME133	*luxO* D61E Δ*angN*	This study
ME1009	Δ*fur*	This study
ME1010	Δ*fur* Δ*luxT*	This study
ME1042	*luxO* D61E Δ*aerR*	This study
ME1019	*luxO* D61E Δ*aerA*	This study
ME1017	*luxO* D61E Δ*aerB*	This study
A. fischeri **ES114**	WT	(8)
ME226	Δ*luxT*	(9)
ME1030	Δ*swrZ*	This study
ME1050	Δ*swrZ* Δ*luxT*	This study
E. coli **S17-1 λ*pir***	WT	(10)
E. coli **MG1655**	WT	(11)

B. Primers used in this study

Sequences are provided in the 5' → 3' direction

Name	Sequence	Description
STR-0040	CGTGAGCGTATCCCGGTATCTAT	qRT-PCR, *hfq* (*V. harveyi*)
STR-0041	TTGCAGTTTGATACCGTTCACAAG	qRT-PCR, *hfq* (*V. harveyi*)
ME-405	TGAAACAGAAATACAACGTTGGTGC	qRT-PCR, *swrZ* (*V. harveyi*)
ME-501	CACAGGTGCAACCCAACAGC	qRT-PCR, *swrZ* (*V. harveyi*)
ME-605	TCTAGAGGCATCAAATAAAACGAAAG	Plasmid construction, pSC101Kan
ME-606	AAGACGGGTAAGCCTGTTGATG	Plasmid construction, pSC101Kan
ME-607	AGCGGTATCATCAACAGGCTTACCCGTCTTCCTGTTAAGTATCTTCCTGGCATC	Plasmid construction, pSC101Kan
ME-608	GAGCCTTTCGTTTTATTTGATGCCTCTAGACACATGGTCCTTCTTGAGTTTGTAAC	Plasmid construction, pSC101Kan
ME-611	ACATGGCATGGATGAACTGTACAAC	Plasmid construction, P$_{swrZ}$-*lux* and P$_{luxT}$-*lux* (pSC101Kan vector)
ME-612	GTTACAAACTCAAGAAGGACCATGTG	Plasmid construction, P$_{swrZ}$-*lux* and P$_{luxT}$-*lux* (pSC101Kan vector)
ME-1121	TAGGAATTCAATTAGGAGGTAATTAAGCATGGAAAAACACTTACCTTTAATAATAAATGG	Plasmid construction, P$_{swrZ}$-*lux* and P$_{luxT}$-*lux* (*luxCDABE*)
ME-1122	TAGACACATGGTCCTTCTTGAGTTTGTAACTTATTACAAATAAGCGAACGCGTCC	Plasmid construction, P$_{swrZ}$-*lux* and P$_{luxT}$-*lux* (*luxCDABE*)
ME-714	CCTCAGTTGTACAGTTCATCCATGCCATGTAAAAAGAGGGCTTCAGCCCTC	Plasmid construction, P$_{swrZ}$-*lux* (P$_{swrZ}$)
ME-604	CCTAGGCCTGTCGAGGCTGTTTCCTGTGTGATTATCGTGACGACAACTAAACAGTAC	Plasmid construction, P$_{swrZ}$-*lux* (P$_{swrZ}$)
ME-1032	GTTTGATCTGCTTAATAAATCTGGTTAAAA	P$_{luxC}$ Control EMSA probe forward primer
ME-1033	TCCATATCAAGAGCTTCTCCTTTG	P$_{luxC}$ Control EMSA probe reverse primer
ME-1190	AATCATCAGTTCAATAAAAAGAGGGC	P$_{swrZ}$ EMSA probe forward primer
ME-1174	TAAACAGTACGTAGTTCGGTTGTAATTG	P$_{swrZ}$ EMSA probe reverse primer
ME-443	GCTTAATTACCTCCTCTTCCTTAGCTCCTGAATTCCTAG	Plasmid construction, p*swrZ* (pFED343 vector)

ME-444	TCACTACTCTGTGCTATGGTGTTC	Plasmid construction, p*swrZ* (pFED343 vector)
ME-839	CAGGAGCTAAGGAAGAGGAGGTAATTAAGCAT GAGGTTGCTTGTGTCTAGTCC	Plasmid construction, p*swrZ* (*swrZ*)
ME-840	AGCATTGAACACCATAGCACAGAGTAGTGATTA AGCAGGCTCTAGCTCTAGTAC	Plasmid construction, p*swrZ* (*swrZ*)
ME-128	GAAGATCATGGATGCCGTTGTTG	qRT-PCR, *luxT* (*V. harveyi*)
ME-415	AATGGTGGCTAATACCTGTACGC	qRT-PCR, *luxT* (*V. harveyi*)
ME-1130	CCTCAGTTGTACAGTTCATCCATGCCATGTCGA ATCCTCTTGTAAATACTGTGGG	Plasmid construction, P$_{luxT}$-*lux* (P$_{luxT}$)
ME-1131	CCTAGGCCTGTCGAGGCTGTTTCCTGTGTGTT ATGGCATATTTTAAGCTCTTCTCTTTG	Plasmid construction, P$_{luxT}$-*lux* (P$_{luxT}$)
pRE112-F	ATGCAGTTCACTTACACCGCTTC	Plasmid construction, pRE112 mediated chromosomal alterations
pRE112-R	GGGATCGGGCCCTATCACTT	Plasmid construction, pRE112 mediated chromosomal alterations
ME-1191	GGGTTGAGAAGCGGTGTAAGTGAACTGCATGT CGGTGGTGCACATGATGC	Plasmid construction, *luxA* deletion (*V. harveyi*)
ME-1192	AGGTGGCTGATAAGTGAGAAGG	Plasmid construction, *luxA* deletion (*V. harveyi*)
ME-1193	GGAAACTTCCTTCTCACTTATCAGCCACCTATT ATCGCATCTATGAAGCTATTCC	Plasmid construction, *luxA* deletion (*V. harveyi*)
ME-1194	ACGCCTGAATAAGTGATAGGGCCCGATCCCTG CGCTTTATCTTCCATCGAC	Plasmid construction, *luxA* deletion (*V. harveyi*)
ME-1172	CAAGGACATACAATACATAATGCTGAG	P$_{luxT}$ EMSA probe forward primer
ME-1173	TGGCATATTTTAAGCTCTTCTCTTTG	P$_{luxT}$ EMSA probe reverse primer
ME-1184	GGAGTTCTCAATACAACAGCCAATC	qRT-PCR, *hyp* (*V. harveyi*)
ME-1185	GATTCGGGTCGATATGGGCATC	qRT-PCR, *hyp* (*V. harveyi*)
ME-817	TCCGCGTACACAATGCAACTC	qRT-PCR, *vopN* (*V. harveyi*)
ME-818	CGTCTTTCACTTTGCGTTTCG	qRT-PCR, *vopN* (*V. harveyi*)
ME-506	GGTTCAGCAACAAGAGTATCGAG	qRT-PCR, *vscO* (*V. harveyi*)
ME-507	CGCCGTTCTTCTTCCTCTTC	qRT-PCR, *vscO* (*V. harveyi*)
ME-1186	TAACCCAATCATCGCAAGTGCC	qRT-PCR, *exsD* (*V. harveyi*)

ME-1187	CGTTGCAGCAAAGACAATTGTTC	qRT-PCR, *exsD* (*V. harveyi)*
ME-1175	CAGTCTCACATTGTTGTCGTTCAC	qRT-PCR, *exsA* (*V. harveyi)*
ME-1176	CTTGAGAGAGGAACACACCACAAC	qRT-PCR, *exsA* (*V. harveyi)*
ME-1177	GGGTTGAGAAGCGGTGTAAGTGAACTGCATAT AAGTAAATAGAGTACGCAGTGAAATC	Plasmid construction, *exsA* deletion (*V. harveyi)*
ME-1178	CACATCCATTTTCTACCCTTCATAATCC	Plasmid construction, *exsA* deletion (*V. harveyi)*
ME-1179	TAGGATTATGAAGGGTAGAAAATGGATGTGGT CGCTATTGCTAAGTGAAATGTC	Plasmid construction, *exsA* deletion (*V. harveyi)*
ME-1180	ACGCCTGAATAAGTGATAGGGCCCGATCCCCG CATTTAAGTCGCTATACACATG	Plasmid construction, *exsA* deletion (*V. harveyi)*
ME-823	TTACCCAAGGCATTACCGCAG	qRT-PCR, *aebC* (*V. harveyi)*
ME-824	TTTGGTTTCCCGCATCCTTTGC	qRT-PCR, *aebC* (*V. harveyi)*
ME-510	GAACCAGAAAGCATCTCATGGC	qRT-PCR, *angM* (*V. harveyi)*
ME-511	CATAGCGTAAATTGAGCCCATAATC	qRT-PCR, *angM* (*V. harveyi)*
ME-805	GGTCACTACTGCTGAAGATGGAC	qRT-PCR, *fatD* (*V. harveyi)*
ME-806	AGCCGGCCCCTGTTAGAATC	qRT-PCR, *fatD* (*V. harveyi)*
ME-10	GGGTTGAGAAGCGGTGTAAGTGAACTGCATAT GTACTGCCGAAAACTCCCGTC	Plasmid construction, *aebF* deletion (*V. harveyi)*
ME-11	TCCTTTGTTTTTCTTATTCTTTGGCTG	Plasmid construction, *aebF* deletion (*V. harveyi)*
ME-12	TAGCAGCCAAAGAATAAGAAAAACAAAGGATTC GTCAACACAATCTAGGTCGAAC	Plasmid construction, *aebF* deletion (*V. harveyi)*
ME-13	ACGCCTGAATAAGTGATAGGGCCCGATCCCAA TCTCTCATTTCATGGACACTACTG	Plasmid construction, *aebF* deletion (*V. harveyi)*
ME-20	GGGTTGAGAAGCGGTGTAAGTGAACTGCATCT TGCGCGATGGGATGTTAAG	Plasmid construction, *angN* deletion (*V. harveyi)*
ME-21	CTGCTTTCACTTTGTTGACAATCAATC	Plasmid construction, *angN* deletion (*V. harveyi)*
ME-22	ACCGATTGATTGTCAACAAAGTGAAAGCAGCA CATCATTGATTTAGTAGCATATTTTC	Plasmid construction, *angN* deletion (*V. harveyi)*
ME-23	ACGCCTGAATAAGTGATAGGGCCCGATCCC CCTCCTATCGAACGCTCATAAAG	Plasmid construction, *angN* deletion (*V. harveyi)*
ME-60	GGGTTGAGAAGCGGTGTAAGTGAACTGCATGA CCAAGAAGATTACGCAGAGTAC	Plasmid construction, *fur* deletion (*V. harveyi)*

ME-61	ATACTTTCCCGTTGGATCATCTGC	Plasmid construction, *fur* deletion (*V. harveyi*)
ME-62	ATCGCTGCAGATGATCCAACGGGAAAGTATAA AGTAACGCTTTTTGTTAAGAAAGAC	Plasmid construction, *fur* deletion (*V. harveyi*)
ME-63	ACGCCTGAATAAGTGATAGGGCCCGATCCCCC GTAAAGTGAAAGGCGTAATTCAC	Plasmid construction, *fur* deletion (*V. harveyi*)
ME-1128	GCCAGACTGCCAGCACATC	qRT-PCR, *fur* (*V. harveyi*)
ME-1129	CATCATCGAATTGGTTTAGTACACG	qRT-PCR, *fur* (*V. harveyi*)
ME-813	TCTTACCCTTTTACTGGAACGCTC	qRT-PCR, *aerR* (*V. harveyi*)
ME-814	GCTAGCGAAGTTTCACCGATG	qRT-PCR, *aerR* (*V. harveyi*)
ME-512	CCTATCTACATTCGCAACTGGC	qRT-PCR, *aerA* (*V. harveyi*)
ME-513	TCAAGTGGACGATAGCCTGAAC	qRT-PCR, *aerA* (*V. harveyi*)
ME-782	GGCTATCATGGGGAGATCAAGTC	qRT-PCR, *aerB* (*V. harveyi*)
ME-783	GGTGATGGGCATTGAGACGTTAC	qRT-PCR, *aerB* (*V. harveyi*)
ME-1198	GGGTTGAGAAGCGGTGTAAGTGAACTGCATAA CGGGTGGTTTCTTGGGTGC	Plasmid construction, *aerR* deletion (*V. harveyi*)
ME-1199	TTTAATAGGGTTATATAAGTTCATACGACAG	Plasmid construction, *aerR* deletion (*V. harveyi*)
ME-1200	TGTCGTATGAACTTATATAACCCTATTAAATGAA ACCCTATATCGCTGTTGCC	Plasmid construction, *aerR* deletion (*V. harveyi*)
ME-1201	ACGCCTGAATAAGTGATAGGGCCCGATCCCGC CAGTTGCGAATGTAGATAGG	Plasmid construction, *aerR* deletion (*V. harveyi*)
ME-1153	GGGTTGAGAAGCGGTGTAAGTGAACTGCATAC ACCAGACCAGTTTGAGCAAG	Plasmid construction, *aerA* deletion (*V. harveyi*)
ME-1154	AAGGCTGATGTTGATGTTTCGC	Plasmid construction, *aerA* deletion (*V. harveyi*)
ME-1155	ACGTTCATGCGAAACATCAACATCAGCCTTGCT CTTGGTTTTGAAGGTGCTC	Plasmid construction, *aerA* deletion (*V. harveyi*)
ME-1156	ACGCCTGAATAAGTGATAGGGCCCGATCCCGA GGGTTAGCCTGATGAGAGTG	Plasmid construction, *aerA* deletion (*V. harveyi*)
ME-1141	GGGTTGAGAAGCGGTGTAAGTGAACTGCATAG CATACAAGCAGCAGACTAAG	Plasmid construction, *aerB* deletion (*V. harveyi*)
ME-1142	CATACTGAGCTGCGTACTGAC	Plasmid construction, *aerB* deletion (*V. harveyi*)
ME-1143	ATGCCTAACGTCAGTACGCAGCTCAGTATGCG CCTACTCATTCAACCTCTAG	Plasmid construction, *aerB* deletion (*V. harveyi*)

ME-1144	ACGCCTGAATAAGTGATAGGGCCCGATCCCCG CATTGATGAAGTCAAGCAG	Plasmid construction, *aerB* deletion (*V. harveyi*)
ME-416	TATACAACAGGGCAGCGTTGG	qRT-PCR, *luxC* (*V. harveyi*)
ME-417	TCCAATTTGCTTCGAGTTTCGC	qRT-PCR, *luxC* (*V. harveyi*)
ME-809	TGCTAAGCGAATACTTTGAAGGAC	qRT-PCR, *VIBHAR_RS19875* (*V. harveyi*)
ME-810	AAGAACGAGATCTGGCTGGC	qRT-PCR, *VIBHAR_RS19875* (*V. harveyi*)
ME-780	AGACAACAGCTCCAATACGGC	qRT-PCR, *VIBHAR_RS25670* (*V. harveyi*)
ME-781	ATTTGGGTTGGCTTTGGTCTCTAC	qRT-PCR, *VIBHAR_RS25670* (*V. harveyi*)
ME-1118	AGTAATCCAAGCGGCTCAAACC	qRT-PCR, *VIBHAR_RS22290* (*V. harveyi*)
ME-1119	TTTTCCATGATACTCGACCAGTCAC	qRT-PCR, *VIBHAR_RS22290* (*V. harveyi*)
ME-516	CGCTGGTAATGGTAATTCCACTAG	qRT-PCR, *VIBHAR_RS27345* (*V. harveyi*)
ME-517	TAGATCGACAAACACTAAACCTCTG	qRT-PCR, *VIBHAR_RS27345* (*V. harveyi*)
ME-514	TACAACAAGCACGCAAAGCG	qRT-PCR, *VIBHAR_RS08855* (*V. harveyi*)
ME-515	ATCTGCACGGTTACGTTCTGG	qRT-PCR, *VIBHAR_RS08855* (*V. harveyi*)
ME-831	CCACGCAACAAACGATGCAG	qRT-PCR, *VIBHAR_RS09515* (*V. harveyi*)
ME-832	TCTCTCTCAAATCACAGTTCAACAC	qRT-PCR, *VIBHAR_RS09515* (*V. harveyi*)
ME-835	TACAGTGCCCTAAGCTGACTTC	qRT-PCR, *cheV* (*V. harveyi*)
ME-836	CAATATCTGTGGTTGGACGGCC	qRT-PCR, *cheV* (*V. harveyi*)
ME-502	GTAGCCATTCACAAGCTGAGC	qRT-PCR, *csgG* (*V. harveyi*)
ME-503	ATTCGACCTAATGCGACTTTGC	qRT-PCR, *csgG* (*V. harveyi*)
ME-93	GTTAACGGGATCAAACTACAGGGAC	qRT-PCR, *hfq* (*A. fischeri*)
ME-94	AGTAGAAATCGCATGCTTGTATACC	qRT-PCR, *hfq* (*A. fischeri*)
ME-1239	TGATCCACCAAGATATTCTCTCTGG	qRT-PCR, *swrZ* (*A. fischeri*)

ME-1240	AGCTTCACGAATAGGGGATGC	qRT-PCR, *swrZ* (*A. fischeri*)
ME-1223	GGGTTGAGAAGCGGTGTAAGTGAACTGCATCG TTTCTGTTCAACTTCCAGCTC	Plasmid construction, *swrZ* deletion (*A. fischeri*)
ME-1224	TTTTCCAGAGAGAATATCTTGGTGG	Plasmid construction, *swrZ* deletion (*A. fischeri*)
ME-1225	ATGATCCACCAAGATATTCTCTCTGGAAAATAC TCGAAAGCCAGTGATCTTATATC	Plasmid construction, *swrZ* deletion (*A. fischeri*)
ME-1226	ACGCCTGAATAAGTGATAGGGCCCGATCCCCT TTCGATGTATTGCTATCTGAGAG	Plasmid construction, *swrZ* deletion (*A. fischeri*)

C. Plasmids used in this study

Plasmid Name	Stock Name	Description	Origin, marker	Reference
pSC101Kan	pME94	Low-copy vector. pSC101 origin from pJES-24 (12) and Kan^R cassette from pEVS143 (13)	pSC101, Kan^R	This study
P*swrZ*-*lux*	pME161	*swrZ-luxCDABE* transcriptional reporter in pME94	pSC101, Kan^R	This study
pKP8-35	pKP8-35	P*BAD* overexpression vector	pBR322, Amp^R	(14)
p*luxT*	pME109	*V. harveyi luxT* overexpression vector, cloned in pKP8-35	pBR322, Amp^R	(2)
pFED343	pFED343	P*tac* overexpression vector	P15A, Cam^R	(15)
p*swrZ*	pME192	*V. harveyi swrZ* overexpression vector, cloned in pFED343	P15A, Cam^R	This study
P*luxT*-*lux*	pME186	*luxT-luxCDABE* transcriptional reporter in pME94	pSC101, Kan^R	This study
pRE112	pRE112	allelic exchange vector harboring *sacB* as a counter-selectable marker	R6Kγ, Cam^R	(16)
pRE112-Δ*luxA*	pME213	*V. harveyi luxA* deletion construct in pRE112	R6Kγ, Cam^R	This study
pRE112-Δ*swrZ*	pME64	*V. harveyi swrZ* deletion construct in pRE112	R6Kγ, Cam^R	(2)
p*luxT*	pME69	*V. harveyi luxT* overexpression vector, cloned in pFED343	P15A, Cam^R	(2)
pRE112-Δ*luxT*	pME12	*V. harveyi luxT* deletion construct in pRE112	R6Kγ, Cam^R	(2)

pRE112-ΔexsA	pME212	*V. harveyi exsA* deletion construct in pRE112	R6Kγ, Cam^R	This study
pRE112-ΔaebF	pME1	*V. harveyi aebF* deletion construct in pRE112	R6Kγ, Cam^R	This study
pRE112-ΔangN	pME3	*V. harveyi angN* deletion construct in pRE112	R6Kγ, Cam^R	This study
pRE112-Δfur	pME9	*V. harveyi fur* deletion construct in pRE112	R6Kγ, Cam^R	This study
pRE112-ΔaerR	pME219	*V. harveyi aerR* deletion construct in pRE112	R6Kγ, Cam^R	This study
pRE112-ΔaerA	pME215	*V. harveyi aerA* deletion construct in pRE112	R6Kγ, Cam^R	This study
pRE112-ΔaerB	pME214	*V. harveyi aerB* deletion construct in pRE112	R6Kγ, Cam^R	This study
pRE112-ΔswrZ (*A. fischeri*)	pME220	*A. fischeri swrZ* deletion construct in pRE112	R6Kγ, Cam^R	This study

Table S1 References

1.	Bassler BL, Greenberg EP, Stevens AM. 1997. Cross-species induction of luminescence in the quorum-sensing bacterium *Vibrio harveyi*. J Bacteriol 179:4043–4045.

2.	Eickhoff MJ, Fei C, Huang X, Bassler BL. 2021. LuxT controls specific quorum-sensing-regulated behaviors in *Vibrionaceae* spp. via repression of *qrr*1, encoding a small regulatory RNA. PLoS Genet 17:e1009336.

3.	Bassler BL, Wright M, Silverman MR. 1994. Sequence and function of LuxO, a negative regulator of luminescence in *Vibrio harveyi*. Mol Microbiol 12:403–412.

4.	Freeman JA, Bassler BL. 1999. A genetic analysis of the function of LuxO, a two-component response regulator involved in quorum sensing in *Vibrio harveyi*. Mol Microbiol 31:665–677.

5.	Tu KC, Bassler BL. 2007. Multiple small RNAs act additively to integrate sensory information and control quorum sensing in *Vibrio harveyi*. Genes Dev 21:221–233.

6.	Pompeani AJ, Irgon JJ, Berger MF, Bulyk ML, Wingreen NS, Bassler BL. 2008. The *Vibrio harveyi* master quorum-sensing regulator, LuxR, a TetR-type protein is both an activator and a repressor: DNA recognition and binding specificity at target promoters. Mol Microbiol 70:76–88.

7.   van Kessel JC, Rutherford ST, Shao Y, Utria AF, Bassler BL. 2013. Individual and combined roles of the master regulators AphA and LuxR in control of the *Vibrio harveyi* quorum-sensing regulon. J Bacteriol 195:436–443.

8.   Boettcher KJ, Ruby EG. 1990. Depressed light emission by symbiotic *Vibrio fischeri* of the sepiolid squid *Euprymna scolopes*. J Bacteriol 172:3701–3706.

9.   Eickhoff MJ, Bassler BL. 2020. *Vibrio fischeri* siderophore production drives competitive exclusion during dual-species growth. Mol Microbiol 114:244–261.

10.   de Lorenzo V, Timmis KN. 1994. Analysis and construction of stable phenotypes in gram-negative bacteria with Tn*5*- and Tn*10*-derived minitransposons. Methods Enzymol 235:386–405.

11.   Blattner FR, Plunkett G, Bloch CA, Perna NT, Burland V, Riley M, Collado-Vides J, Glasner JD, Rode CK, Mayhew GF, Gregor J, Davis NW, Kirkpatrick HA, Goeden MA, Rose DJ, Mau B, Shao Y. 1997. The complete genome sequence of *Escherichia coli* K-12. Science 277:1453–1462.

12.   Papenfort K, Silpe JE, Schramma KR, Cong J-P, Seyedsayamdost MR, Bassler BL. 2017. A *Vibrio cholerae* autoinducer-receptor pair that controls biofilm formation. Nat Chem Biol 13:551–557.

13.   Dunn AK, Millikan DS, Adin DM, Bose JL, Stabb EV. 2006. New rfp- and pES213-derived tools for analyzing symbiotic *Vibrio fischeri* reveal patterns of infection and *lux* expression in situ. Appl Environ Microbiol 72:802–810.

14.   Papenfort K, Pfeiffer V, Mika F, Lucchini S, Hinton JCD, Vogel J. 2006. SigmaE-dependent small RNAs of *Salmonella* respond to membrane stress by accelerating global *omp* mRNA decay. Mol Microbiol 62:1674–1688.

15.   Swem LR, Swem DL, Wingreen NS, Bassler BL. 2008. Deducing receptor signaling parameters from in vivo analysis: LuxN/AI-1 quorum sensing in *Vibrio harveyi*. Cell 134:461–473.

16.   Edwards RA, Keller LH, Schifferli DM. 1998. Improved allelic exchange vectors and their use to analyze 987P fimbria gene expression. Gene 207:149–157.