Global membrane tension is independent of polyacrylamide substrate stiffness

Eva Kreysing[1,*], Jeffrey Mc Hugh[2,3,*], Sarah K. Foster[1,4], Kurt Andresen[5], Ryan D. Greenhalgh[1], Eva K. Pillai[1], Andrea Dimitracopoulos[1], Ulrich F. Keyser[2,#], Kristian Franze[1,6,7,#]

1) Department of Physiology, Development and Neuroscience, University of Cambridge, Cambridge CB2 3DY, UK

2) Biological and Soft Systems, Maxwell Centre, Cavendish Laboratory, Cambridge CB3 0HE, UK

3) Neuroglial Interactions in Cerebral Physiopathology, CIRB, CNRS UMR 7241/INSERM U1050, Collège de France, Paris 75005, France

4) Systems Biology of Microbial Communities, Cluster of Excellence – CMFI, University of Tübingen, 72076 Tübingen, Germany

5) Department of Physics, Gettysburg College, Gettysburg, PA 17325, United States of America

6) Institute of Medical Physics, Friedrich-Alexander-Universität Erlangen-Nürnberg, 91052 Erlangen, Germany.

7) Max-Planck-Zentrum für Physik und Medizin, 91054 Erlangen, Germany

* These authors contributed equally.

\# To whom correspondence should be addressed: ufk20@cam.ac.uk or kf284@cam.ac.uk

Abstract

Most animal cells are surrounded by a cell membrane and an underlying actomyosin cortex.

Both structures are linked, and they are under tension. In-plane membrane tension and cortical

tension both influence many cellular processes, including cell migration, division, and

endocytosis. However, while actomyosin tension is regulated by substrate stiffness, how

membrane tension responds to mechanical substrate properties is currently poorly understood.

Here, we probed the effective membrane tension of neurons and fibroblasts cultured on glass

and polyacrylamide substrates of varying stiffness using optical tweezers. In contrast to

actomyosin-based traction forces, both peak forces and steady state tether forces of cells

cultured on hydrogels were independent of substrate stiffness and did not change after

blocking myosin II activity using blebbistatin, indicating that tether and traction forces are not

directly linked. Peak forces in fibroblasts on hydrogels were about twice as high as those in

neurons, indicating stronger membrane-cortex adhesion in fibroblasts. Steady state tether

forces were generally higher in cells cultured on hydrogels than on glass, which we explain by a

mechanical model. Our results provide new insights into the complex regulation of effective

membrane tension and pave the way for a deeper understanding of the biological processes it

instructs.

Introduction

The cell membrane is supported by the underlying actomyosin cortex, and molecular linkers

form tight connections between the two layers. Osmotic pressure resulting from ion gradients

across the membrane and interactions of the membrane with the force-generating actomyosin

cytoskeleton contribute to a tension across the membrane. This membrane tension regulates

many cellular processes, such as cell migration, division, stem cell fate choice, and endo- and

exocytosis[1–6]. Despite its biological importance[7–9], membrane tension is still difficult to

quantify. In-plane membrane tension is defined as the force per unit length acting tangentially

on the plasma membrane. Because of the membrane's coupling to cytoskeletal elements,

measurements of the 'effective' membrane tension usually contain contributions of the

underlying actomyosin cortex and other force-generating elements[10,11].

The actomyosin cortex is coupled to the extracellular environment through membrane proteins

such as integrins and cadherins. It generates tensile forces which are exerted on the

environment. These forces, and hence cytoskeletal tension, increase with substrate stiffness

for most cell types[12,13], suggesting that the measured effective membrane tension should also

increase on stiffer substrates. However, the regulation of membrane tension by substrate

mechanics is yet to be probed experimentally.

Results

Here, we used optical tweezers (OT) to probe membrane properties of cells cultured on

polyacrylamide substrates of different stiffnesses as well as on glass. Using a membrane-

adherent bead held in an optical trap, we pulled membrane tethers from cells, and the resulting

tether forces served as an indicator for effective membrane tension [2,14,15] (Fig. 1a, b; Fig. S1,

Video S1). Throughout a pull, the force on the tether can be continuously monitored, resulting

in a characteristic force-time curve (Fig. 1c). The force rises quickly after initiation of the pull

and reaches a peak before decreasing sharply. In agreement with Sheetz[14], we attribute this

peak force (PF) to the initial local detachment of plasma membrane from the actomyosin cortex

(Fig. 1a).

Subsequently, the tether is pulled until it reaches a set length. During this elongation process,

the plasma membrane slides over the actin cortex while bonds between the membrane and the

lipid-binding proteins located in or at the cortex are broken and quickly re-established[16].

Following the elongation phase, the bead is held in a stable position and the steady-state force

(SSF) is measured (Fig. 1b). The SSF is thought to reflect a combination of the in-plane plasma

membrane tension and the cortical tension, with the strength of the membrane-cortex

adhesion determining the coupling between these two elements[16,17].

To test the effect of substrate mechanics and cortical tension on effective membrane tension,

we performed tether pulls in fibroblasts, which possess a dense actomyosin cortex, and in

neuronal axons, whose sub-membranous cytoskeleton is characterized by actin rings spaced by

spectrin tetramers[18]. Cells were cultured on glass as well as on hydrogels fabricated within a

stiffness range adjusted to match their natural environment. For fibroblasts, hydrogel

stiffnesses ranged from 100Pa (very soft; similar to subcutaneous adipose tissue), to 10kPa

(stiff; already in the stiffness range of many fibrotic tissues)[19]. For neurons, the highest

substrate stiffness used was 1kPa, corresponding to the highest stiffness experienced by this

type of neuron *in vivo*[20].

Peak forces (PFs) measured in fibroblasts were about twice as high as those measured in

neuronal axons (Fig. 2b, f). Within each cell type, PFs were similar across all tested hydrogel

substrates. Steady state forces (SSFs), which were comparable between fibroblasts and

neurons, were also similar across all polyacrylamide substrate stiffnesses (Fig. 2c, g).

In contrast, actomyosin-based traction forces are known to scale with substrate stiffnesses for

both neurons and fibroblasts[21,22]. We explicitly measured traction forces in our culture

conditions and confirmed that traction forces exhibited by both cell types increased

significantly with increasing substrate stiffness within the investigated stiffness range (Fig. 2d,

Fig. S3). Similarly, net forces, which are obtained by integrating traction forces across neuronal

growth cones, and which provide an approximation for the tension exerted by a growth cone

on its axon[21,23], significantly increased on stiffer substrates (Fig 2h). These data indicated that

traction forces and net forces assessed by 2D traction force microscopy and tether forces

measured by OT are not tightly coupled.

Cellular cortical tension is mediated by actomyosin contractility[24]. However, cortical tension

further away from adhesion sites, where traction forces are generated and transmitted to the

environment, was recently suggested to be largely independent of traction forces[22]. In order to

further examine the interplay between cortical tension and effective membrane tension, we

measured tether forces in fibroblasts cultured on 10 kPa hydrogels, where we would expect the

highest cytoskeletal tension[22], following treatment with the myosin II (and thus contractility)

inhibitor blebbistatin. Blebbistatin treatment had no significant effect on either PFs or SSFs (Fig.

3), suggesting that there is indeed no strong coupling between actomyosin-based cortical

tension and global effective membrane tension in cells cultured on hydrogels.

As numerous *in vitro* studies use glass-bottomed culture dishes, and the surface properties of

glass differ considerably from those of hydrogels, we also cultured fibroblasts and neurons on

glass to test whether effective membrane tension is comparable in both culture environments.

In neurons, PFs measured in axons cultured on glass were indeed like those observed on

hydrogels (Fig. 2f). However, PFs for fibroblasts cultured on glass were significantly lower than

those for fibroblasts grown on hydrogels (Fig. 2b). Similarly, SSFs were significantly lower for

fibroblasts and neurons cultured on glass compared to those cultured on hydrogels (Fig. 2c, g).

Unlike on hydrogels, tether forces measured on glass were similar for fibroblasts and neurons

(Fig. 2).

While PFs did not change significantly after blocking actomyosin contractility in fibroblasts

cultured on glass, SSFs increased on glass following blebbistatin treatment[25] (Fig 3b), reaching

values similar to SSFs seen in fibroblasts cultured on hydrogels. These data showed that

effective membrane tension depends on the type of culture substrate, and they suggested that

the coupling between the cell cortex and the membrane also differs between glass and

hydrogel substrates.

Discussion

Here we found that the steady-state tether forces, which are thought to scale with effective

membrane tension, do not change as a function of hydrogel substrate stiffness within a

physiologically relevant stiffness range in both fibroblasts and neurons (Fig. 2c, g). In contrast,

actomyosin-based traction and net forces increased on stiffer substrates (Fig. 2d, h), as shown

previously[22], suggesting that the effective membrane tension measured by OT is not directly

linked to cellular contractility on hydrogels[26]. In line with this interpretation, the application of

the myosin II blocker, blebbistatin, did not alter the effective membrane tension of fibroblasts

cultured on hydrogels (Fig. 3). Alternatively, membrane tension might be influenced by cortical

tension on short time scales but is then actively regulated to resume original values quickly[27].

Another possibility is that traction forces transmitted from focal adhesions to the cortex might

dissipate within a finite distance, so that overall cortical tension would not scale with local

traction forces[22]. If cortical and membrane tension are coupled, membrane tension would only

increase locally[28] near focal adhesions, which would not be seen when probed further away.

Such local membrane tension fluctuations would explain, for example, why the

mechanosensitive ion channel Piezo1, which is thought to be activated by membrane tension, is

activated only near focal adhesion sites despite being distributed across the whole cell

surface[29].

The effective membrane tension was higher in cells cultured on hydrogels than in those

cultured on glass (Fig. 2). Fibroblasts developed much stronger stress fibres on glass than on

hydrogels (Fig. S2)[30]. The strong contraction of stress fibres on glass might deform the cortex

between focal adhesions, this way releasing cortical tension and causing the membrane to

relax, as schematically indicated in Figure 4. This could not only explain why SSFs in fibroblasts

were lower on glass than on hydrogels (Fig. 2c, Fig. 4b) but also why blebbistatin treatment,

which relaxes stress fibres, led to a significant increase in SSFs on glass but not on hydrogels

(Figs. 3b, 4c-d). This hypothesis agrees with recent experiments showing that perturbations of

stress fibres lead to an increase in tension in the nearby actin cortex[31]. The tension exerted by

growth cones on the distal axon, which increased on stiffer substrates (Fig. 2h), could similarly

lead to membrane relaxation in neurons cultured on glass (Fig. 2g). Alternatively, or

additionally, differences in physical and chemical properties between glass and hydrogels could

also activate different signalling pathways involved in the regulation of membrane tension

homeostasis[27].

The PF, which is the initial force required to detach the membrane from the cortex, is related to

membrane-cortex adhesion[14] (in analogy to attempting to open Velcro from the middle).

Fibroblasts have a dense actomyosin cortex, and proteins such as ezrin/radixin/moesin (ERM)

can link the membrane to the cortex across the cell surface. Neuronal axons, on the other hand,

have a periodic actin-spectrin network underneath their membrane[18] and much less area for

ERM proteins to connect the membrane to actin rings. This morphological difference may lead

to lower membrane-cortex adhesion in axons compared to fibroblasts, explaining why PFs are

about twice as high in fibroblasts compared to axons cultured on physiologically stiff hydrogels.

A low membrane-cortex coupling could also explain why tethers pulled from axons can freely

"slide" along the axon as observed in a previous[15] and the current study (video S2). To our

knowledge, tether sliding has not been reported in fibroblasts.

Our results confirm that membrane tension measurements need to be interpreted carefully.

Effective membrane tension did not depend on substrate stiffness within a physiological

stiffness range, yet it varies between hydrogels and glass, and between cell types. Future

technological developments enabling highly resolved quantitative membrane tension

measurements will shed light on the molecular control of local membrane tension, an

important regulator of many cellular functions.

Acknowledgments

The authors would like to thank Alex Winkel and Joy Thompson for AFM measurements of hydrogels, Liz Williams for preparation of hydrogels, and Ewa Paluch, Ruby Peters and Aki Stubb for discussions. J. Mc H. acknowledges funding from AFOSR (Grant No. FA9550-17-1-0118). S. K. F. acknowledges funding from the Herchel Smith Foundation. U. F. K. was supported by an ERC Consolidator Grant DesignerPores No. 647144, K. F. was supported by the European Research Council (Consolidator Grant 772426) and the Alexander von Humboldt Foundation (Alexander von Humboldt Professorship).

Author Contributions

J. Mc H., E. K., and K. A. performed tether pulling experiments. J. Mc H. analysed the results from OT experiments. E. K. cultured 3T3 fibroblasts and prepared hydrogels. S. K. F. dissected and cultured Xenopus laevis RGCs and prepared hydrogels. A.D. performed TFM experiments with 3T3 fibroblasts. R. G. performed the TFM experiments with neurons and all TFM analyses. E. K. and E. P. performed the immunostainings. J. Mc H., E. K., K. F., and U. F. K. designed experiments with 3T3 fibroblasts. J. Mc H., S. K. F., K. A., and U. F. K. designed pulling experiments with X. laevis RGCs. S. K. F., Mc H., E. K., U.F.K. and K. F. conceived the project. E. K., J. Mc H., S. K. F., U.F.K. and K.F. wrote the manuscript.

Competing interests

The authors declare no competing interests.

References

1. Raucher, D. & Sheetz, M. P. Cell Spreading and Lamellipodial Extension Rate Is Regulated by Membrane Tension. *J. Cell Biol.* **148**, 127–136 (2000).

2. Nussenzveig, H. M. Cell membrane biophysics with optical tweezers. *Eur. Biophys. J.* **47**, 499–514 (2018).

3. Gauthier, N. C., Fardin, M. A., Roca-Cusachs, P. & Sheetz, M. P. Temporary increase in plasma membrane tension coordinates the activation of exocytosis and contraction during cell spreading. *Proc. Natl. Acad. Sci. U. S. A.* **108**, 14467–14472 (2011).

4. Keren, K. *et al.* Mechanism of shape determination in motile cells. *Nature* **453**, 475–480 (2008).

5. De Belly, H. *et al.* Membrane Tension Gates ERK-Mediated Regulation of Pluripotent Cell Fate. *Cell Stem Cell* **28**, 273-284.e6 (2021).

6. Dai, J. & Sheetz, M. P. Regulation of endocytosis, exocytosis, and shape by membrane tension. *Cold Spring Harb. Symp. Quant. Biol.* **60**, 567–571 (1995).

7. Sheetz, M. P. Cell control by membrane–cytoskeleton adhesion. *Nat. Rev. Mol. Cell Biol.* **2**, 392–396 (2001).

8. Pontes, B., Monzo, P. & Gauthier, N. C. Membrane tension: A challenging but universal physical parameter in cell biology. *Semin. Cell Dev. Biol.* **71**, 30–41 (2017).

9. Pontes, B. *et al.* Membrane tension controls adhesion positioning at the leading edge of cells. *J. Cell Biol.* **216**, 2959–2977 (2017).

10. Salbreux, G., Charras, G. & Paluch, E. Actin cortex mechanics and cellular morphogenesis. *Trends Cell Biol.* **22**, 536–545 (2012).

11.   Clark, A. G., Wartlick, O., Salbreux, G. & Paluch, E. K. Stresses at the Cell Surface during

Animal Cell Morphogenesis. *Curr. Biol.* **24**, R484–R494 (2014).

12.   Califano, J. P. & Reinhart-King, C. A. Substrate Stiffness and Cell Area Predict Cellular

Traction Stresses in Single Cells and Cells in Contact. *Cell. Mol. Bioeng.* **3**, 68–75 (2010).

13.   Sabass, B., Gardel, M. L., Waterman, C. M. & Schwarz, U. S. High Resolution Traction

Force Microscopy Based on Experimental and Computational Advances. *Biophys. J.* **94**, 207–

220 (2008).

14.   Sheetz, M. P. Cell control by membrane–cytoskeleton adhesion. *Nat. Rev. Mol. Cell Biol.*

2, 392–396 (2001).

15.   Datar, A., Bornschlögl, T., Bassereau, P., Prost, J. & Pullarkat, P. A. Dynamics of membrane

tethers reveal novel aspects of cytoskeleton-membrane interactions in axons. *Biophys. J.* **108**,

489–497 (2015).

16.   Sens, P. & Plastino, J. Membrane tension and cytoskeleton organization in cell motility. *J.*

Phys. Condens. Matter **27**, (2015).

17.   Gauthier, N. C., Masters, T. A. & Sheetz, M. P. Mechanical feedback between membrane

tension and dynamics. *Trends Cell Biol.* **22**, 527–535 (2012).

18.   Xu, K., Zhong, G. & Zhuang, X. Actin, spectrin, and associated proteins form a periodic

cytoskeletal structure in axons. *Science* **339**, 452–456 (2013).

19.   Wells, R. G. Tissue Mechanics and Fibrosis. *Biochim. Biophys. Acta* **1832**, 884–890 (2013).

20.   Franze, K., Janmey, P. A. & Guck, J. Mechanics in Neuronal Development and Repair.

Annu. Rev. Biomed. Eng. **15**, 227–251 (2013).

21.   Koch, D., Rosoff, W. J., Jiang, J., Geller, H. M. & Urbach, J. S. Strength in the Periphery:

Growth Cone Biomechanics and Substrate Rigidity Response in Peripheral and Central

Nervous System Neurons. *Biophys. J.* **102**, 452–460 (2012).

22.   Rheinlaender, J. *et al.* Cortical cell stiffness is independent of substrate mechanics. *Nat.*

Mater. **19**, 1019–1025 (2020).

23.   Franze, K. Integrating Chemistry and Mechanics: The Forces Driving Axon Growth. *Annu.*

Rev. Cell Dev. Biol. **36**, 61–83 (2020).

24.   Chugh, P. *et al.* Actin cortex architecture regulates cell surface tension. *Nat. Cell Biol.* **19**,

689–697 (2017).

25.   Lieber, A. D., Yehudai-Resheff, S., Barnhart, E. L., Theriot, J. A. & Keren, K. Membrane

Tension in Rapidly Moving Cells Is Determined by Cytoskeletal Forces. *Curr. Biol.* **23**, 1409–

1417 (2013).

26.   Mandal, K. *et al.* Soft hyaluronic gel promotes cell spreading, stress fibers, focal adhesion,

membrane tension by phosphoinositide signaling, not traction force. *ACS Nano* **13**, 203–214

(2019).

27.   Thottacherry, J. J. *et al.* Mechanochemical feedback control of dynamin independent

endocytosis modulates membrane tension in adherent cells. *Nat. Commun.* **9**, 1–14 (2018).

28.   Shi, Z., Graber, Z. T., Baumgart, T., Stone, H. A. & Cohen, A. E. Cell membranes resist flow.

Cell **175**, 1769-1779.e13 (2018).

29.   Ellefsen, K. L. *et al.* Myosin-II mediated traction forces evoke localized Piezo1-dependent

Ca 2+ flickers. *Commun. Biol.* **2**, 1–13 (2019).

30.   Walcott, S. & Sun, S. X. A mechanical model of actin stress fiber formation and substrate

elasticity sensing in adherent cells. *Proc. Natl. Acad. Sci.* **107**, 7757–7762 (2010).

31.   Vignaud, T. *et al.* Stress fibres are embedded in a contractile cortical network. *Nat.*

Mater. 1–11 (2020) doi:10.1038/s41563-020-00825-z.

32.   Leung, K.-M. *et al.* Asymmetrical β-actin mRNA translation in growth cones mediates

attractive turning to netrin-1. *Nat. Neurosci.* **9**, 1247–1256 (2006).

33.   Nieuwkoop, P. D. & Faber, J. *Normal table of xenopus laevis (daudin)*. (Routledge, 1994).

34.   Mc Hugh, J., Andresen, K. & Keyser, U. F. Cation dependent electroosmotic flow in glass

nanopores. *Appl. Phys. Lett.* **115**, 113702 (2019).

35.   Gittes, F. & Schmidt, C. F. Chapter 8 signals and noise in micromechanical measurements.

in *Methods in cell biology.* vol. 55 129–156 (Academic Press, 1997).

36.   Buer, C. S., Weathers, P. J. & Swartzlander, G. A. Changes in Hechtian strands in cold-

hardened cells measured by optical microsurgery. *Plant Physiol.* **122**, 1365–1377 (2000).

37.   Kucik, D. F., Kuo, S. C., Elson, E. L. & Sheetz, M. P. Preferential attachment of membrane

glycoproteins to the cytoskeleton at the leading edge of lamella. *J. Cell Biol.* **114**, 1029–1036

(1991).

38.   Hochmuth, R. M., Shao, J. Y., Dai, J. & Sheetz, M. P. Deformation and flow of membrane

into tethers extracted from neuronal growth cones. *Biophys. J.* **70**, 358–369 (1996).

39.   Pontes, B. *et al.* Cell cytoskeleton and tether extraction. *Biophys. J.* **101**, 43–52 (2011).

40.   Moshayedi, P. *et al.* The relationship between glial cell mechanosensitivity and foreign

body reactions in the central nervous system. *Biomaterials* **35**, 3919–3925 (2014).

41.   Koser, D. E. *et al.* Mechanosensing is critical for axon growth in the developing brain. *Nat.*

Neurosci. **19**, 1592–1598 (2016).

42.   Moshayedi, P. *et al.* Mechanosensitivity of astrocytes on optimized polyacrylamide gels

analyzed by quantitative morphometry. *J. Phys. Condens. Matter* **22**, (2010).

Figures

Figure 1: Illustration of an OT measurement. Peak and steady state tether forces provide

different insights into cell membrane properties. (a) When an OT pull is initiated, the local

membrane-cortex adhesion first needs to be overcome, resulting in a peak force (PF) measured

at the beginning of the experiment (see (c)). (b) After tether extension, the bead is held

stationary, and the steady state force (SSF), which scales with effective membrane tension, is

recorded. (c) A typical force-time curve recorded during a tether pull, showing a peak (PF) at

the beginning of the pull and a plateau (SSF) when the tether is held at its maximum extension.

Figure 2: Tether and traction forces of fibroblasts and neurons on different substrates. (a-d)

3T3 fibroblasts and (e-h) *Xenopus* retinal ganglion cell axons; shear moduli of polyacrylamide

substrates are provided in Pa. (a, e) Fluorescence images of membrane tethers pulled from a

3T3 fibroblast and an axon. Fibroblast membranes were labelled with CellMask membrane

stain, neurons with Tetramethylrhodamine Dextran. Scale bars: 5µm. (b) Peak tether forces

(PFs) and (c) steady state tether forces (SSFs) were similar on all hydrogels irrespective of their

stiffness (Kruskal-Wallis tests followed by Tukey posthoc tests), but significantly larger than on

glass. (d) 3T3 fibroblasts exerted higher traction forces on stiffer hydrogels compared to on

softer ones (two-tailed Mann-Whitney test). (f) PFs in axons were similar on all hydrogel and

glass substrates. (g) SSFs in neurons were similar on different hydrogels but higher than those

measured on glass. (h) Net forces (growth cone forces pulling on the axon) were higher on

stiffer hydrogels than on softer ones (two-tailed Mann-Whitney test).

Figure 3: Inhibition of actomyosin contractility increases SSFs on glass but not on hydrogels.

(a) The application of 20 µM Blebbistatin did not significantly alter PFs on hydrogels or on glass

(Mann-Whitney test). (b) SSFs of fibroblasts grown on glass but not on hydrogels increased

significantly following blebbistatin treatment. The untreated conditions correspond to the data

shown in Figure 2.

Figure 4: Toy model of substrate type-dependent effective membrane tension. Schematic

representation of the cell cortex and stress fibres in fibroblasts on glass and hydrogel substrates

without and with blebbistatin treatment. (a) In fibroblasts on glass substrates without

blebbistatin treatment, the strong contraction of stress fibres might relax the tension in the

cortex and thus reduce membrane tension. (b) In fibroblasts on hydrogel substrates without

blebbistatin treatment, we observe fewer stress fibres, most likely resulting in lower

contractility and less relaxation of the membrane. The resulting membrane tension is higher

compared to that of cells on glass. (c) In fibroblasts on glass substrates with blebbistatin

treatment, the stress fibres relax. This might increase membrane tension. (d) In fibroblasts on

hydrogel substrates with blebbistatin treatment, fewer stress fibres relax, and the effect on

membrane tension is too low to be detected by OT.

Materials and methods

NIH 3T3 fibroblast cell culture

3T3 cells were cultured in plastic cell culture flasks in culture media (90 ml DMEM with glutamine (Gibco 22320-022), 10 ml Fetal Bovine Serum (Gibco 25300054), 1 ml Penicillin-Streptomycin-Amphotericin B Mixture (PSF, Lonza 17-745E)). After reaching confluency, cells were enzymatically detached. For this step, cells were washed twice with prewarmed (37°C) PBS (Sigma D8537) and then incubated with 1ml 10% TrypLE (Thermo Fisher Scientific A1285901) in PBS at 37°C for 5 min. 5 ml culture media was then added, and the cells were cultured in fresh culture flasks or plated on glass dishes and polyacrylamide (PAA) hydrogels for 4-48 h prior to experiments.

Hydrogels of 3 different elastic moduli (shear moduli = 100 Pa,1 kPa, and 10 kPa) were prepared and functionalized as described below. PAA gels as well as glass bottom culture dishes were coated with 0.01 mg/ml PDL in PBS for at least 4 h (typically overnight) followed by 0.02 mg/ml fibronectin in PBS for at least 4 h. Culture media was exchanged with prewarmed (37°C) imaging media (Live Cell Imaging Solution, henceforth 'imaging media' (Thermo Fisher Scientific A14291DJ)) supplemented with 1% PSF 15 min prior to measurements.

For the blebbistatin treatment, cells were incubated with 20 μM blebbistatin (Sigma-Aldrich, B0560-1MG) for 15 min prior to measurement. For visualisation of membrane tethers, fibroblasts were stained with CellMask Orange Plasma Membrane Stain (ThermoFisher Scientific, USA, C10045-100 μl). The dye was added to the imaging media in the culture dish at 1:1000 dilution 15 min prior to the tether pulls.

Xenopus laevis retinal ganglion cell culture

All animal experiments were conducted in compliance with the Ethical Review Committee of the

University of Cambridge and Home office Guidelines.

Xenopus embryos were derived from *in vitro* fertilization of eggs kindly provided by the Gurden

Institute Frog Facility (University of Cambridge, UK). For fluorescent labelling of *Xenopus*

neurons, 5 nL of 0.2% w/v tetramethylrhodamine-labeled dextran (Thermo Fisher, D3308)

diluted in nuclease-free water was injected in each dorsal blastomere of a 4-cell stage *Xenopus*

embryo. Injections were performed as previously described[32].

Embryos were grown at 14-16°C with daily cleaning in 0.1X Modified Barth's Saline (10X MBS:

88mM NaCL, 2.4mM NaHCO3, 1mM HEPES, 0.82mM MgSO4, 0.33mM Ca(NO3)2, 0.41 mM

CaCl2, pH 7.6).

All dissections took place at stage 35/36 (stages according to Nieuwkoop and Faber[33]). In the

case of dextran-injected embryos, embryos were screened using a fluorescence

stereomicroscope prior to dissection to ensure the presence of dextran in the eye primordia

(note that dextran-injected embryos are morphologically normal). Embryos were anesthetized

prior to dissection in 20% tricaine methanesulfonate solution (MS222, pH 7.6-7.8, with 1x PSF)

and subsequently immobilized in a Sylgard®184-lined petri dish using a bent 0.2 mm insect bin

(Austerlitz). Eye primordia were carefully explanted using a 0.15 mm insect pin (Austerlitz) held

in a pin holder. Explants were immediately transferred to 60% L15 media (Sigma-Aldrich, L4386,

pH 7.6-7.8, with 1x PSF, henceforth referred to as 'Xenopus culture media').

Following dissection, explants were placed with the lens facing up onto PAA gels or glass bottom

culture dishes. The hydrogels were prepared as described below.

Hydrogels and glass dishes were coated overnight (hydrogels) or for 30 min (glass) with 0.01

mg/ml PDL (Sigma-Aldrich, P6407, diluted in PBS) followed by 2 h in 5 µg/mL laminin (Sigma-

Aldrich, L2020, diluted in PBS). Following explant plating, dishes were allowed to sit for 1-2 h on

the benchtop to allow explants to adhere, and then were moved to a 20°C incubator overnight.

Immunostainings

Cells were washed twice with prewarmed (37°C) PBS. Afterwards, they were fixed and

permeabilized with 4% PFA (Thermo Fisher 28908) and 0.2% tritonX (Sigma-Aldrich T8787) in PBS

for 10 min at room temperature (RT). The cells were washed once with PBS at RT. Cells were

incubated with DAPI (1.7 µg/ml Sigma-Aldrich D9542) and Phalloidin (Life Technologies A12379,

diluted 1:300) in 1% Bovine Serum Albumin (Sigma-Aldrich A3294) in PBS for 1 h at RT. In the

end, the dishes were washed twice with PBS. Cells were imaged with a Leica DMi8

epifluorescence setup with a ×63 oil objective (NA 1.4, Leica).

Optical tweezers and force measurement

OT measurements took place on a custom-built inverted microscope[34]. Briefly, a 5 W 1064 nm

ytterbium fibre laser (YLM-5-LP, IPG Laser, Germany) was used at a high power (> 1 W) to

minimize intensity fluctuations. Excess power was redirected to a beam dump and the power at

the optical trap itself was 600 mW. The trap was generated by overfilling the back aperture of an

NA 1.2 UPlanSApo water immersion objective (Olympus, Japan) with the laser beam.

To enable controlled positioning of samples, the sample stage was mounted on top of a precision

piezoelectric stage (P-517.43 and E-710.3, Physik Instrumente, Germany) with a positioning

resolution of 1 nm. The sample stage was illuminated by a white fibre light source (DC-950 Fiber-

Lite, Edmund Optics, USA). White light entered the objective and then passed through a dichroic

mirror (DM) to a 550 nm short pass DM (DMSP550, Thorlabs, USA). This passed light below 550

nm to the 532 nm notch DM. This mirror coupled the fluorescence laser into the microscope,

and in the process blocked the portion of incoming white light ranging from 515 to 540 nm. Light

that passed through the notch DM was focused onto a CMOS camera (MC1362, Mikrotron,

Germany), which was used for tracking particles in the optical trap. Light above 550 nm was

passed to the fluorescence imaging camera (optiMOS Scientific CMOS, Teledyne Imaging, USA),

which was used to find beads to trap, locate cells, and record fluorescence images of tethers.

Forces were measured with the OT by modelling a particle in the trap potential as a Hookean

mass and spring system. By tracking the position of a trapped particle and multiplying this by a

calibrated trap stiffness constant, the force experienced by the particle was determined. Trap

calibration was performed using the power spectrum method[35]. Postprocessing of the data was

done using a custom script in Python 3.6, which is available on GitHub

https://github.com/mchughj33/mem-tension.

OT bead functionalisation

As reported previously[36,37], beads with 2 μm diameter (Kisker PC-S-2.0) were functionalized with

Concanavalin A (ConA, from Canavalia, Sigma-Aldrich, C2272-10MG) as follows.

10 µL bead solution were added into 100 µL of 2 mg/mL ConA in a tube (low-protein binding)

and incubated for 30 min at 4°C. Beads were washed twice via centrifugation in 100µL of

5mg/mL bovine serum albumin (BSA, Sigma-Aldrich, A3294) in 200mM Tris (Sigma, GE17-1321-

01) in ddH2O. All centrifugation steps were conducted at 4°C, 10,000 rcf. After the final wash, 1

ml imaging media supplemented with 1% PSF was added and gently mixed by pipetting.

Membrane tether pulling experiments

15 min prior to each fibroblast measurement, the culture media was exchanged with prewarmed

(37°C) imaging media supplemented with 1% PSF. For Xenopus experiments, Xenopus media was

exchanged with RT imaging media ~15 minutes prior to imaging.

In each OT experiment, 2 µl functionalised beads (see above) were added to the culture media

and a bead captured in the trap. During the entire process, the position of the bead relative to

the laser trap was recorded in order to measure the force produced by the tether pull. Before

attaching the bead to the cell, a short recording of the Brownian motion of the trapped bead was

taken. The mean force from each recording was used to define the point of zero force in the

subsequent analysis of each respective pull. Afterwards, the bead was carefully brought into

contact with the cell membrane. The image of the bead was monitored for a change in the

interference pattern around the bead, indicating contact between the bead and membrane. At

this point the beads was held stationary for 5 s. Then it was moved approximately 0.25 µm away

from the cell while monitoring the particle tracking system to see if the mean displacement of

the bead from the trap centre was reduced, which signified that the bead was attached to the

membrane.

Beads were then moved away from the cell at 0.1 µm s^{-1} in the case of 3T3 fibroblasts and 0.5

µm s^{-1} in the case of neurons, leading to the formation of a membrane tether between the cell

and the bead. After reaching the end of the pull, 10 µm for 3T3 fibroblasts and 20 µm for

neurons, the bead was held stationary for 30 s. The protocol for pulling tethers from neurons

was based on previous work[38]. Due to high rate of tether loss during preliminary pulls with

fibroblasts, to the pulling velocity was reduced, leading to a subsequent decrease in the pull

distance to avoid reducing experimental throughput. Pontes et al. showed that varying the

velocities for tether pulls in 3T3 fibroblasts within the range of 0.07 µm/s to 1µm/s, did not

affect the force results[39]. We therefore do not expect this velocity change to affect our

observations. For each pull, the peak force was defined as the maximum force value reached,

and the steady state force was determined as the mean of the 30 s period at the end of the pull

during which the bead was held stationary. Analysis was performed using custom analysis scripts

written in python 3.6.

Hydrogels for fibroblasts

Deformable hydroxy-polyacryamide substrates for fibroblast cultures were prepared as

described previously[22,40]. Briefly, for each gel a top coverslip and a glass-bottomed culture dishe

(MatTek P35G-0.20-C and Ibidi 81158 were used for OT and TFM, respectively) were prepared.

Both coverslips were first washed in 70% ethanol and distilled water, and then dried. Bottom

coverslips were wiped with a cotton bud soaked in 1M NaOH, dried, and then incubated in (3-

aminopropyl)trimethoxysilane (APTMS, Sigma Aldrich, 281778) for 3 min. APTMS was washed off

thoroughly in ddH20 and coverslips were then incubated in a 0.5% solution of glutaraldehyde

(Sigma Aldrich, G6257) for 30 min and then rinsed 3x with distilled water. Top coverslips were

treated for 10 min with Rain-X (Shell Car Care International). Acrylamide (AA, Sigma Aldrich,

A4058) stock solution was made by mixing 500 µL of 40% Acrylamide + 65 µL of 100%

Hydroxyacrylamide (Sigma Aldrich, 697931). Afterwards the gel premix was prepared by mixing

500 µL Acrylamide stock solution with 250 µL 2% Bisacrylamide (BA, Fisher Scientific, BP1404-

250). The premix was diluted in PBS to achieve the different shear moduli according to the **table**

1. For TFM gels, 10 µL of PBS was replaced by 10 µl bead solution (FluoSpheres™ Carboxylate-

Modified Microspheres, 0.2 µm, dark red fluorescent)) and the gel mix was sonicated for 30 s to

separate the beads.

Volume of gel premix	Volume of PBS	Shear modulus (G')
53 µL	447 µL	0.1 kPa
75 µL	425 µL	1 kPa
150 µL	350 µL	10 kPa

Table 1: Hydroxy-polyacrylamide gel composition used for fibroblasts

Freshly-made premixes were de-gassed under vacuum for 4-6 min. To each 500 µL of final gel

solution, 1.5 µL 0.3% (v/v) N, N, N', N'-tetramethylethylenediamine (TEMED, Thermo Fisher,

15524–010) and 5 µL 0.1% (w/v) ammonium persulfate (APS, Sigma, 215589) solution were

added and the premix was gently mixed. 14 µL or 10 µL of this mixture were pipetted on the

bottom coverslip and 18 mm or 16 mm top coverslips (for 14 µL and 10 µL volume, respectively)

were placed on top (with RainX-treated side facing down). The gels were soaked in PBS for at

least 20 min before the top coverslips were removed. Afterwards, they were washed twice with

PBS and then treated with ultraviolet light for 30 min. Hydrogels were coated as described

above.

Hydrogels for neurons

Deformable polyacrylamide gel substrates were prepared as described previously[41,42] for neuron

culture. The coverslips were prepared as described in the hydroxy-acryamide gel protocol using

MatTek dishes as the bottom coverslip. The gel mix was prepared according to **table 2** to achieve

the different shear moduli. For TFM gels, the PBS volume was reduced by 10 µL and replaced by

10 µL bead solution and the gel mix was sonicated for 30 s to separate the beads.

Volume of AA	Volume of BA	Volume of PBS	Shear modulus (G')
63 µL	10 µL	427 µL	0.1 kPa
63 µL	17.5 µL	419.5 µL	0.3 kPa
94 µL	15 µL	391 µL	1 kPa

471
472   Table 2: Polyacryamide gel composition used for neurons
473

Freshly-made gel mixes were briefly de-gassed under vacuum for 4-6 min. To each 500 µL of final

gel solution, 1.5 µL 0.3% (v/v) TEMED and 5 µL 0.1% (w/v) APS solution were added and gently

mixed. 7 µL, 14 µL of this mixture were pipetted onto the bottom coverslip and 18 mm top

coverslips were placed on top (with RainX treated side facing down) for OT experiments and TFM

experiments, respectively. The gels were soaked in PBS for at least 20 min before the top

coverslips were removed. The gels were subsequently functionalized via a 4h treatment with

hydrazine hydrate (Sigma Aldrich, 225819), followed by 1 h in 5% acetic acid (Fisher Scientific,

10171460), and then washed thoroughly in PBS. Afterwards, they were treated with ultraviolet

light for 30 min. Hydrogels were coated as described above.

TFM

24 h after plating, fibroblasts were imaged with an inverted microscope (Leica DMi8) at 37°C and

5% CO_2, equipped with a digital sCMOS camera (ORCA-Flash4.0, Hamamatsu Photonics), an

EL6000 illuminator (Leica) and a ×63 oil objective (NA = 1.4, Leica). Images were acquired with

Leica LAS X software. Fluorescence images of beads and wide-field images of cells were taken

every 2 min for 10 minutes. After image acquisition, the culture media was exchanged with

Trypsin-EDTA (Gibco) to detach cells from the gel. Reference images of fluorescent beads were

taken 15 min after trypsinization.

16 - 24 h after plating, *Xenopus* eye primordia explants were imaged using an inverted Nikon

microscope (Ti-E at room temperature) equipped with an sCMOS camera (Prime BSI Scientific,

Photometrics), a CoolLED pE-4000, and ×60 oil objective (NA = 1.4, Nikon). Fluorescence images

of beads and wide-field images of cells were taken every 5 min for 1 h. After image acquisition,

the culture media was exchanged with Trypsin-EDTA (Gibco) or 20% sodium dodecyl sulfate

solution (Severn Biotech Ltd., 2 0-4002-01) to detach the explants from the gel. Reference

images of fluorescent beads were taken 15 min after trypsinization or immediately after sodium

dodecyl sulfate treatment.

Traction stress maps were calculated for each frame. Traction stresses were averaged over time

(12 frames for neurons, 5 frames for fibroblasts) for each cell. Postprocessing of the data and

statistical analyses were done with a custom script in Python 3.8 and is available on GitHub

https://github.com/rg314/pytraction.

Statistics

OT data are shown as violin plots, TFM data as box plots. Violin shapes are kernel density

estimations, boxes indicate the range of data that falls between the first and third quartile,

whiskers show the range of data within 1.5 times the interquartile range. Density estimator

width is based on the number of samples at a given value. Median values for each dataset are

marked by a white circle. Lines between violins indicate significant differences in the samples.

The boxplot diagrams were generated using python seaborn. The borders of the box indicate the

quartiles of the dataset, the median is indicated by a line within the box, and the whiskers

include 1.5*interquartile range. Points that were determined to be outliers were plotted

individually. Kruskal-Wallis one-way analysis of variance (ANOVA) tests followed by a Tukey

posthoc test were used for statistical analysis when comparing multiple groups. When

comparing two groups, a Mann-Whitney test was used.