Effective cell membrane tension is independent of polyacrylamide substrate

stiffness

Eva Kreysing[1,*], Jeffrey Mc Hugh[2,3,*], Sarah K. Foster[1,4], Kurt Andresen[5], Ryan D.

Greenhalgh[1], Eva K. Pillai[1], Andrea Dimitracopoulos[1], Ulrich F. Keyser[2,#], Kristian

Franze[1,6,7,#]

1) Department of Physiology, Development and Neuroscience, University of

Cambridge, Cambridge CB2 3DY, UK

2) Biological and Soft Systems, Maxwell Centre, Cavendish Laboratory, Cambridge

CB3 0HE, UK

3) Neuroglial Interactions in Cerebral Physiopathology, CIRB, CNRS UMR

7241/INSERM U1050, Collège de France, Paris 75005, France

4) Systems Biology of Microbial Communities, Cluster of Excellence – CMFI,

University of Tübingen, 72076 Tübingen, Germany

5) Department of Physics, Gettysburg College, Gettysburg, PA 17325, United States

of America

6) Institute of Medical Physics, Friedrich-Alexander-Universität Erlangen-Nürnberg,

91052 Erlangen, Germany.

7) Max-Planck-Zentrum für Physik und Medizin, 91054 Erlangen, Germany

*   These authors contributed equally.

To whom correspondence should be addressed: ufk20@cam.ac.uk or

kf284@cam.ac.uk

Abstract

Most animal cells are surrounded by a cell membrane and an underlying actomyosin cortex. Both structures are linked, and they are under tension. In-plane membrane tension and cortical tension both influence many cellular processes, including cell migration, division, and endocytosis. However, while actomyosin tension is regulated by substrate stiffness, how membrane tension responds to mechanical substrate properties is currently poorly understood. Here, we probed the effective membrane tension of neurons and fibroblasts cultured on glass and polyacrylamide substrates of varying stiffness using optical tweezers. In contrast to actomyosin-based traction forces, both peak forces and steady state tether forces of cells cultured on hydrogels were independent of substrate stiffness and did not change after blocking myosin II activity using blebbistatin, indicating that tether and traction forces are not directly linked. Peak forces in fibroblasts on hydrogels were about twice as high as those in neurons, indicating stronger membrane-cortex adhesion in fibroblasts. Steady state tether forces were generally higher in cells cultured on hydrogels than on glass, which we explain by a mechanical model. Our results provide new insights into the complex regulation of effective membrane tension and pave the way for a deeper understanding of the biological processes it instructs.

Introduction

The cell membrane is supported by the underlying actomyosin cortex, and molecular linkers form tight connections between the two layers. Osmotic pressure resulting from ion gradients across the membrane and interactions of the membrane with the force-generating actomyosin cytoskeleton contribute to a tension across the membrane. This membrane tension regulates many cellular processes, such as cell migration, division, stem cell fate choice, and endo- and exocytosis[1–6]. Despite its biological importance[7–9], membrane tension is still difficult to quantify. In-plane membrane tension is defined as the force per unit length acting tangentially on the plasma membrane. Because of the membrane's coupling to cytoskeletal elements, measurements of the 'effective' membrane tension usually contain contributions of the underlying actomyosin cortex and other force-generating elements[10,11].

The actomyosin cortex is coupled to the extracellular environment through transmembrane proteins such as integrins and cadherins, which transmit tensile forces generated by acto-myosin interaction to the environment. The resulting traction forces, and hence cytoskeletal tension, increase with substrate stiffness for most cell types[12,13], suggesting that the measured effective membrane tension should also increase on stiffer substrates. However, the regulation of membrane tension by substrate mechanics is yet to be probed experimentally.

Results

Here, we used optical tweezers (OT) to probe membrane properties of cells cultured on polyacrylamide substrates of different stiffnesses as well as on glass. Using a membrane-

adherent bead held in an optical trap, we pulled membrane tethers from cells, and the resulting

tether forces served as an indicator for effective membrane tension [2,14,15] (Fig. 1a, b; Fig. S1,

Video S1). Throughout a pull, the force on the tether can be continuously monitored, resulting

in a characteristic force-time curve (Fig. 1c). The force rises quickly after initiation of the pull

and reaches a peak before decreasing sharply. In agreement with Sheetz[14], we attribute this

peak force (PF) to the initial local detachment of plasma membrane from the actomyosin cortex

(Fig. 1a), although it may also be influenced by measurement parameters such as the adhesion

area between the bead and the cell membrane[16].

Subsequently, the tether is pulled until it reaches a set length. During this elongation process,

the plasma membrane slides over the actin cortex while bonds between the membrane and the

lipid-binding proteins located in or at the cortex are broken and quickly re-established[17]. At the

same time, membrane proteins may accumulate at the base of the tether and influence the

pulling force[15]. Overall, the measured forces then decrease during the elongation phase.

Afterwards, the bead is held in a stable position and the steady-state force (SSF) is measured

(Fig. 1b). The SSF is thought to reflect a combination of the in-plane plasma membrane tension

and the cortical tension, with the strength of the membrane-cortex adhesion determining the

coupling between these two elements[17,18]. As such, the SSF depends in a non-linear manner on

the number of cortex-plasma membrane linkers such a ERM proteins[19].

To test the effect of substrate mechanics and cortical tension on effective membrane tension,

we performed tether pulls in fibroblasts, which possess a dense actomyosin cortex, and in

neuronal axons, whose sub-membranous cytoskeleton is characterized by actin rings spaced by

spectrin tetramers[20]. Cells were cultured on glass as well as on hydrogels fabricated within a

stiffness range adjusted to match their natural environment. For fibroblasts, hydrogel

stiffnesses ranged from 100Pa (very soft; similar to subcutaneous adipose tissue), to 10kPa

(stiff; already in the stiffness range of many fibrotic tissues)[21]. For neurons, the highest

substrate stiffness used was 1kPa, corresponding to the highest stiffness experienced by this

type of neuron *in vivo*[22].

Peak forces (PFs) measured in fibroblasts were about twice as high as those measured in

neuronal axons (Fig. 2b, f). Within each cell type, PFs were similar across all tested hydrogel

substrates. Steady state forces (SSFs), which were comparable between fibroblasts and

neurons, were also similar across all polyacrylamide substrate stiffnesses (Fig. 2c, g).

In contrast, actomyosin-based traction forces, i.e., contractile cellular forces exerted on the

substrate, are known to scale with substrate stiffnesses for both neurons and fibroblasts[23,24].

We measured these forces in our culture conditions using traction force microscopy (TFM),

where cells are cultured on soft, elastic hydrogels with embedded fluorescent nanoparticles

(see Methods)[25]. Forces exerted by the cells lead to local substrate deformations, which can be

tracked using fluorescence imaging of the nanoparticles, whose displacement can be used to

calculate traction forces[26].

We confirmed that traction forces exhibited by both cell types increased significantly with

increasing substrate stiffness within the investigated stiffness range (Fig. 2d, Fig. S2). Similarly,

net forces, which are obtained by integrating traction forces across neuronal growth cones, and

which provide an approximation for the tension exerted by a growth cone on its axon[23,27],

significantly increased on stiffer substrates (Fig 2h). These data indicated that traction forces

and net forces assessed by 2D traction force microscopy and tether forces measured by OT are

not tightly coupled.

In order to further examine the interplay between cortical tension and effective membrane

tension, we measured tether forces in fibroblasts cultured on 10 kPa hydrogels, where we

would expect the highest cytoskeletal tension[24], following treatment with the myosin II (and

thus contractility) inhibitor blebbistatin. Blebbistatin treatment had no significant effect on

either PFs or SSFs (Fig. 3), suggesting that there is indeed no strong coupling between

actomyosin-based cortical tension and average effective membrane tension in cells cultured on

hydrogels.

As numerous *in vitro* studies use glass-bottomed culture dishes, and the surface properties of

glass differ considerably from those of hydrogels, we also cultured fibroblasts and neurons on

glass to test whether effective membrane tension is comparable in both culture environments.

In neurons, PFs measured in axons cultured on glass were indeed like those observed on

hydrogels (Fig. 2f). However, PFs for fibroblasts cultured on glass were significantly lower than

those for fibroblasts grown on hydrogels (Fig. 2b). Similarly, SSFs were significantly lower for

fibroblasts and neurons cultured on glass compared to those cultured on hydrogels (Fig. 2c, g).

Unlike on hydrogels, tether forces measured on glass were similar for fibroblasts and neurons

(Fig. 2).

While PFs did not change significantly after blocking actomyosin contractility in fibroblasts

cultured on glass, SSFs increased on glass following blebbistatin treatment[28] (Fig 3b), reaching

values similar to SSFs seen in fibroblasts cultured on hydrogels. These data showed that

effective membrane tension depends on the type of culture substrate, and they suggested that

the coupling between the cell cortex and the membrane also differs between glass and

hydrogel substrates.

Discussion

Here we found that the steady-state tether forces, which are thought to scale with effective

membrane tension, do not change as a function of hydrogel substrate stiffness within a

physiologically relevant stiffness range in both fibroblasts and neurons (Fig. 2c, g). In contrast,

actomyosin-based traction and net forces increased on stiffer substrates (Fig. 2d, h), as shown

previously[24], suggesting that the effective membrane tension measured by OT dorsolaterally to

adhesion sites is not directly linked to cellular contractility on hydrogels[29]. In line with this

interpretation, the application of the myosin II blocker, blebbistatin, did not alter the effective

membrane tension of fibroblasts cultured on hydrogels (Fig. 3).

Alternatively, membrane tension might be influenced by cortical tension on short time scales

but is then actively regulated to resume original values quickly[30]. Another possibility is that

traction forces and membrane tension are coupled also on longer time scales but traction

forces transmitted from focal adhesions and ventral stress fibres to the actin cortex[31] dissipate

with increasing distance from the stress fibres. Thus, on hydrogels enhanced cortical tension

might be limited to ventral areas of the cell surface not accessible by OT. This would be in line

with a recent study suggesting that cortical tension in dorsolateral parts of the cell are largely

independent of traction forces[24].

If cortical and membrane tension are coupled, membrane tension would then only change

locally[32] near focal adhesions and stress fibres[31], which would not be seen when probed further

away. Such local membrane tension heterogeneities would explain, for example, why the

mechanosensitive ion channel Piezo1, which is thought to be activated by membrane tension, is

activated only near focal adhesion sites despite being distributed across the whole cell

surface[33].

The effective membrane tension was higher in cells cultured on hydrogels than in those

cultured on glass (Fig. 2). Fibroblasts developed much stronger stress fibres on glass than on

hydrogels (Fig. S3)[34], indicating considerably larger traction forces. The strong contraction of

stress fibres and the embedding actin cortex[33] on glass could propagate further within the actin

cortex than on hydrogels. The concomitant decrease in cortex area could cause the membrane

to relax also in areas assessed by OT in cells cultured on glass, as schematically indicated in

Figure 4, explaining why tether forces were lower on glass than on hydrogels (Fig. 2).

This could not only explain why SSFs in fibroblasts were lower on glass than on hydrogels (Fig.

2c, Fig. 4b) but also why blebbistatin treatment, which relaxes both stress fibres, led to a

significant increase in SSFs on glass but not on hydrogels (Figs. 3b, 4c-d). The tension exerted

by growth cones on the distal axon, which increased on stiffer substrates (Fig. 2h), could

similarly lead to membrane relaxation in neurons cultured on glass (Fig. 2g). Alternatively, or

additionally, differences in physical and chemical properties between glass and hydrogels could

also activate different signalling pathways involved in the regulation of membrane tension

homeostasis[30].

The PF, which is the initial force required to detach the membrane from the cortex, is related to

membrane-cortex adhesion[14] (in analogy to attempting to open Velcro from the middle).

Fibroblasts have a dense actomyosin cortex, and proteins such as ezrin/radixin/moesin (ERM

proteins) can link the membrane to the cortex across the cell surface. Neuronal axons, on the

other hand, have a periodic actin-spectrin network underneath their membrane[20] and much

less area for ERM proteins to connect the membrane to actin rings. This morphological

difference may lead to lower membrane-cortex adhesion in axons compared to fibroblasts,

explaining why PFs are about twice as high in fibroblasts compared to axons cultured on

physiologically stiff hydrogels. A low membrane-cortex coupling could also explain why tethers

pulled from axons can freely "slide" along the axon as observed in a previous[15] and the current

study (video S2). To our knowledge, tether sliding has not been reported in fibroblasts.

Our results confirm that membrane tension measurements need to be interpreted carefully.

Effective membrane tension dorsolaterally of adhesion sites did not depend on substrate

stiffness within a physiological stiffness range, yet it varied between hydrogels and glass, and

between cell types. Future technological developments enabling highly resolved quantitative

membrane tension measurements will shed light on the molecular control of local membrane

tension, an important regulator of many cellular functions.

Acknowledgments

The authors would like to thank Alex Winkel and Joy Thompson for AFM measurements of hydrogels, Liz Williams for preparation of hydrogels, and Ewa Paluch, Ruby Peters and Aki Stubb for discussions. J. Mc H. acknowledges funding from AFOSR (Grant No. FA9550-17-1-0118). S. K. F. acknowledges funding from the Herchel Smith Foundation. U. F. K. was supported by an ERC Consolidator Grant DesignerPores No. 647144, K. F. was supported by the European Research Council (Consolidator Grant 772426) and the Alexander von Humboldt Foundation (Alexander von Humboldt Professorship).

Author Contributions

J. Mc H., E. K., and K. A. performed tether pulling experiments. J. Mc H. analysed the results from OT experiments. E. K. cultured 3T3 fibroblasts and prepared hydrogels. S. K. F. dissected and cultured Xenopus laevis RGCs and prepared hydrogels. A.D. performed TFM experiments with 3T3 fibroblasts. R. G. performed the TFM experiments with neurons and all TFM analyses. E. K. and E. P. performed the immunostainings. J. Mc H., E. K., K. F., and U. F. K. designed experiments with 3T3 fibroblasts. J. Mc H., S. K. F., K. A., and U. F. K. designed pulling experiments with X. laevis RGCs. S. K. F., J. Mc H., E. K., U.F.K. and K. F. conceived the project. E. K., J. Mc H., S. K. F., U.F.K. and K.F. wrote the manuscript.

Competing interests

The authors declare no competing interests.

References

References

1. Raucher, D. & Sheetz, M. P. Cell Spreading and Lamellipodial Extension Rate Is

Regulated by Membrane Tension. *J. Cell Biol.* **148**, 127–136 (2000).

2. Nussenzveig, H. M. Cell membrane biophysics with optical tweezers. *Eur. Biophys. J.*

47, 499–514 (2018).

3. Gauthier, N. C., Fardin, M. A., Roca-Cusachs, P. & Sheetz, M. P. Temporary

increase in plasma membrane tension coordinates the activation of exocytosis and

contraction during cell spreading. *Proc. Natl. Acad. Sci. U. S. A.* **108**, 14467–14472

(2011).

4. Keren, K. *et al.* Mechanism of shape determination in motile cells. *Nature* **453**, 475–

480 (2008).

5. De Belly, H. *et al.* Membrane Tension Gates ERK-Mediated Regulation of Pluripotent

Cell Fate. *Cell Stem Cell* **28**, 273-284.e6 (2021).

6. Dai, J. & Sheetz, M. P. Regulation of endocytosis, exocytosis, and shape by

membrane tension. *Cold Spring Harb. Symp. Quant. Biol.* **60**, 567–571 (1995).

7. Sheetz, M. P. Cell control by membrane–cytoskeleton adhesion. *Nat. Rev. Mol. Cell*

Biol. **2**, 392–396 (2001).

8. Pontes, B., Monzo, P. & Gauthier, N. C. Membrane tension: A challenging but

universal physical parameter in cell biology. *Semin. Cell Dev. Biol.* **71**, 30–41 (2017).

9. Pontes, B. *et al.* Membrane tension controls adhesion positioning at the leading edge

of cells. *J. Cell Biol.* **216**, 2959–2977 (2017).

10.   Salbreux, G., Charras, G. & Paluch, E. Actin cortex mechanics and cellular

morphogenesis. *Trends Cell Biol.* **22**, 536–545 (2012).

11.   Clark, A. G., Wartlick, O., Salbreux, G. & Paluch, E. K. Stresses at the Cell

Surface during Animal Cell Morphogenesis. *Curr. Biol.* **24**, R484–R494 (2014).

12.   Califano, J. P. & Reinhart-King, C. A. Substrate Stiffness and Cell Area Predict

Cellular Traction Stresses in Single Cells and Cells in Contact. *Cell. Mol. Bioeng.* **3**,

68–75 (2010).

13.   Sabass, B., Gardel, M. L., Waterman, C. M. & Schwarz, U. S. High Resolution

Traction Force Microscopy Based on Experimental and Computational Advances.

Biophys. J. **94**, 207–220 (2008).

14.   Sheetz, M. P. Cell control by membrane–cytoskeleton adhesion. *Nat. Rev. Mol.*

Cell Biol. **2**, 392–396 (2001).

15.   Datar, A., Bornschlögl, T., Bassereau, P., Prost, J. & Pullarkat, P. A. Dynamics of

membrane tethers reveal novel aspects of cytoskeleton-membrane interactions in

axons. *Biophys. J.* **108**, 489–497 (2015).

16.   Koster, G., Cacciuto, A., Derényi, I., Frenkel, D. & Dogterom, M. Force barriers

for membrane tube formation. *Phys. Rev. Lett.* **94**, 16–19 (2005).

17.   Sens, P. & Plastino, J. Membrane tension and cytoskeleton organization in cell

motility. *J. Phys. Condens. Matter* **27**, (2015).

18.   Gauthier, N. C., Masters, T. A. & Sheetz, M. P. Mechanical feedback between

membrane tension and dynamics. *Trends Cell Biol.* **22**, 527–535 (2012).

19.   Paraschiv, A. *et al.* Influence of membrane-cortex linkers on the extrusion of

membrane tubes. *Biophys. J.* **120**, 598–606 (2021).

20.   Xu, K., Zhong, G. & Zhuang, X. Actin, spectrin, and associated proteins form a

periodic cytoskeletal structure in axons. *Science* **339**, 452–456 (2013).

21.   Wells, R. G. Tissue Mechanics and Fibrosis. *Biochim. Biophys. Acta* **1832**, 884–

890 (2013).

22.   Franze, K., Janmey, P. A. & Guck, J. Mechanics in Neuronal Development and

Repair. *Annu. Rev. Biomed. Eng.* **15**, 227–251 (2013).

23.   Koch, D., Rosoff, W. J., Jiang, J., Geller, H. M. & Urbach, J. S. Strength in the

Periphery: Growth Cone Biomechanics and Substrate Rigidity Response in

Peripheral and Central Nervous System Neurons. *Biophys. J.* **102**, 452–460 (2012).

24.   Rheinlaender, J. *et al.* Cortical cell stiffness is independent of substrate

mechanics. *Nat. Mater.* **19**, 1019–1025 (2020).

25.   Schwarz, U. S. & Soiné, J. R. D. Traction force microscopy on soft elastic

substrates: A guide to recent computational advances. *Biochim. Biophys. Acta BBA -*

Mol. Cell Res. **1853**, 3095–3104 (2015).

26.   Huang, Y. *et al.* Traction force microscopy with optimized regularization and

automated Bayesian parameter selection for comparing cells. *Sci. Rep.* **9**, 539

(2019).

27.   Franze, K. Integrating Chemistry and Mechanics: The Forces Driving Axon

Growth. *Annu. Rev. Cell Dev. Biol.* **36**, 61–83 (2020).

28.   Lieber, A. D., Yehudai-Resheff, S., Barnhart, E. L., Theriot, J. A. & Keren, K.

Membrane Tension in Rapidly Moving Cells Is Determined by Cytoskeletal Forces.

Curr. Biol. **23**, 1409–1417 (2013).

29.   Mandal, K. *et al.* Soft hyaluronic gel promotes cell spreading, stress fibers, focal

adhesion, membrane tension by phosphoinositide signaling, not traction force. *ACS*

Nano **13**, 203–214 (2019).

30.   Thottacherry, J. J. *et al.* Mechanochemical feedback control of dynamin

independent endocytosis modulates membrane tension in adherent cells. *Nat.*

Commun. **9**, 1–14 (2018).

31.   Vignaud, T. *et al.* Stress fibres are embedded in a contractile cortical network.

Nat. Mater. 1–11 (2020) doi:10.1038/s41563-020-00825-z.

32.   Shi, Z., Graber, Z. T., Baumgart, T., Stone, H. A. & Cohen, A. E. Cell membranes

resist flow. *Cell* **175**, 1769-1779.e13 (2018).

33.   Ellefsen, K. L. *et al.* Myosin-II mediated traction forces evoke localized Piezo1-

dependent Ca 2+ flickers. *Commun. Biol.* **2**, 1–13 (2019).

34.   Walcott, S. & Sun, S. X. A mechanical model of actin stress fiber formation and

substrate elasticity sensing in adherent cells. *Proc. Natl. Acad. Sci.* **107**, 7757–7762

(2010).

35.   Leung, K.-M. *et al.* Asymmetrical β-actin mRNA translation in growth cones

mediates attractive turning to netrin-1. *Nat. Neurosci.* **9**, 1247–1256 (2006).

36.   Nieuwkoop, P. D. & Faber, J. *Normal table of xenopus laevis (daudin).*

(Routledge, 1994).

37.   Mc Hugh, J., Andresen, K. & Keyser, U. F. Cation dependent electroosmotic flow

in glass nanopores. *Appl. Phys. Lett.* **115**, 113702 (2019).

38.   Gittes, F. & Schmidt, C. F. Chapter 8 signals and noise in micromechanical

measurements. in *Methods in cell biology.* vol. 55 129–156 (Academic Press, 1997).

39.   Buer, C. S., Weathers, P. J. & Swartzlander, G. A. Changes in Hechtian strands

in cold-hardened cells measured by optical microsurgery. *Plant Physiol.* **122**, 1365–

1377 (2000).

40.   Kucik, D. F., Kuo, S. C., Elson, E. L. & Sheetz, M. P. Preferential attachment of

membrane glycoproteins to the cytoskeleton at the leading edge of lamella. *J. Cell*

Biol. **114**, 1029–1036 (1991).

41.   Hochmuth, R. M., Shao, J. Y., Dai, J. & Sheetz, M. P. Deformation and flow of

membrane into tethers extracted from neuronal growth cones. *Biophys. J.* **70**, 358–

369 (1996).

42.   Pontes, B. *et al.* Cell cytoskeleton and tether extraction. *Biophys. J.* **101**, 43–52

(2011).

43.   Moshayedi, P. *et al.* The relationship between glial cell mechanosensitivity and

foreign body reactions in the central nervous system. *Biomaterials* **35**, 3919–3925

(2014).

44.   Koser, D. E. *et al.* Mechanosensing is critical for axon growth in the developing

brain. *Nat. Neurosci.* **19**, 1592–1598 (2016).

45.   Moshayedi, P. *et al.* Mechanosensitivity of astrocytes on optimized

polyacrylamide gels analyzed by quantitative morphometry. *J. Phys. Condens. Matter*

22, (2010).

Figures

Figure 1: Illustration of an OT measurement. Peak and steady state tether forces provide

different insights into cell membrane properties. (a) When an OT pull is initiated, the local

membrane-cortex adhesion first needs to be overcome, resulting in a peak force (PF) measured

at the beginning of the experiment (see (c)). (b) After tether extension, the bead is held

stationary, and the steady state force (SSF), which scales with effective membrane tension, is

recorded. (c) A typical force-time curve recorded during a tether pull, showing a peak (PF) at

the beginning of the pull and a plateau (SSF) when the tether is held at its maximum extension.

Figure 2: Tether and traction forces of fibroblasts and neurons on different substrates. (a-d)

3T3 fibroblasts and (e-h) *Xenopus* retinal ganglion cell axons; shear moduli of polyacrylamide

substrates are provided in Pa. (a, e) Fluorescence images of membrane tethers pulled from a

3T3 fibroblast and an axon. Fibroblast membranes were labelled with CellMask membrane

stain, neurons with Tetramethylrhodamine Dextran. Scale bars: 5μm. (b) Peak tether forces

(PFs) and (c) steady state tether forces (SSFs) were similar on all hydrogels irrespective of their

stiffness (Kruskal-Wallis tests followed by Tukey posthoc tests), but significantly larger than on

glass. (d) 3T3 fibroblasts exerted higher traction forces on stiffer hydrogels compared to on

softer ones (two-tailed Mann-Whitney test). (f) PFs in axons were similar on all hydrogel and

glass substrates. (g) SSFs in neurons were similar on different hydrogels but higher than those

measured on glass. (h) Net forces (growth cone forces pulling on the axon) were higher on

stiffer hydrogels than on softer ones (two-tailed Mann-Whitney test).

Figure 3: Inhibition of actomyosin contractility increases SSFs on glass but not on hydrogels.

(a) The application of 20 µM Blebbistatin did not significantly alter PFs on hydrogels or on glass

(Mann-Whitney test). (b) SSFs of fibroblasts grown on glass but not on hydrogels increased

significantly following blebbistatin treatment. The untreated conditions correspond to the data

shown in Figure 2.

Figure 4: Toy model of substrate type-dependent effective membrane tension. Schematic

representation of the cell cortex, stress fibres and cell membrane in fibroblasts on glass and

hydrogel substrates without and with blebbistatin treatment. (a) In fibroblasts on glass

substrates without blebbistatin treatment, the strong contraction of stress fibres, which is

transmitted to the nearby actin cortex[31], might lead to the relaxation of the membrane and

thus reduce membrane tension. (b) In fibroblasts on hydrogel substrates without blebbistatin

treatment, we observed fewer stress fibres, most likely resulting in lower contractility and less

relaxation of the membrane. The resulting membrane tension is higher compared to that of

cells on glass. (c) In fibroblasts on glass substrates with blebbistatin treatment, the stress fibres

relax. This might increase membrane tension. (d) In fibroblasts on hydrogel substrates with

blebbistatin treatment, fewer stress fibres relax, and the effect on membrane tension is too low

to be detected by OT.

Materials and methods

NIH 3T3 fibroblast cell culture

3T3 cells were cultured in plastic cell culture flasks in culture media (90 ml DMEM with glutamine (Gibco 22320-022), 10 ml Fetal Bovine Serum (Gibco 25300054), 1 ml Penicillin-Streptomycin-Amphotericin B Mixture (PSF, Lonza 17-745E)). After reaching confluency, cells were enzymatically detached. For this step, cells were washed twice with prewarmed (37°C) PBS (Sigma D8537) and then incubated with 1ml 10% TrypLE (Thermo Fisher Scientific A1285901) in PBS at 37°C for 5 min. 5 ml culture media was then added, and the cells were cultured in fresh culture flasks or plated on glass dishes and polyacrylamide (PAA) hydrogels for 4-48 h prior to experiments.

Hydrogels of 3 different elastic moduli (shear moduli = 100 Pa,1 kPa, and 10 kPa) were prepared and functionalized as described below. PAA gels as well as glass bottom culture dishes were coated with 0.01 mg/ml PDL in PBS for at least 4 h (typically overnight) followed by 0.02 mg/ml fibronectin in PBS for at least 4 h. Culture media was exchanged with prewarmed (37°C) imaging media (Live Cell Imaging Solution, henceforth 'imaging media' (Thermo Fisher Scientific A14291DJ)) supplemented with 1% PSF 15 min prior to measurements.

For the blebbistatin treatment, cells were incubated with 20 µM blebbistatin (Sigma-Aldrich, B0560-1MG) for 15 min prior to measurement. For visualisation of membrane tethers, fibroblasts were stained with CellMask Orange Plasma Membrane Stain (ThermoFisher Scientific, USA, C10045-100 µl). The dye was added to the imaging media in the culture dish at 1:1000 dilution 15 min prior to the tether pulls.

***Xenopus laevis* retinal ganglion cell culture**

All animal experiments were conducted in compliance with the Ethical Review

Committee of the University of Cambridge and Home office Guidelines.

Xenopus embryos were derived from *in vitro* fertilization of eggs kindly provided by the

Gurden Institute Frog Facility (University of Cambridge, UK). For fluorescent labelling of

Xenopus neurons, 5 nL of 0.2% w/v tetramethylrhodamine-labeled dextran (Thermo

Fisher, D3308) diluted in nuclease-free water was injected in each dorsal blastomere of

a 4-cell stage *Xenopus* embryo. Injections were performed as previously described[35].

Embryos were grown at 14-16°C with daily cleaning in 0.1X Modified Barth's Saline

(10X MBS: 88mM NaCL, 2.4mM NaHCO3, 1mM HEPES, 0.82mM MgSO4, 0.33mM

Ca(NO3)2, 0.41 mM CaCl2, pH 7.6).

All dissections took place at stage 35/36 (stages according to Nieuwkoop and Faber[36]).

In the case of dextran-injected embryos, embryos were screened using a fluorescence

stereomicroscope prior to dissection to ensure the presence of dextran in the eye

primordia (note that dextran-injected embryos are morphologically normal). Embryos

were anesthetized prior to dissection in 20% tricaine methanesulfonate solution

(MS222, pH 7.6-7.8, with 1x PSF) and subsequently immobilized in a Sylgard®184-

lined petri dish using a bent 0.2 mm insect bin (Austerlitz). Eye primordia were carefully

explanted using a 0.15 mm insect pin (Austerlitz) held in a pin holder. Explants were

immediately transferred to 60% L15 media (Sigma-Aldrich, L4386, pH 7.6-7.8, with 1x

PSF, henceforth referred to as 'Xenopus culture media').

Following dissection, explants were placed with the lens facing up onto PAA gels or

glass bottom culture dishes. The hydrogels were prepared as described below.

Hydrogels and glass dishes were coated overnight (hydrogels) or for 30 min (glass) with

0.01 mg/ml PDL (Sigma-Aldrich, P6407, diluted in PBS) followed by 2 h in 5 µg/mL

laminin (Sigma-Aldrich, L2020, diluted in PBS). Following explant plating, dishes were

allowed to sit for 1-2 h on the benchtop to allow explants to adhere, and then were

moved to a 20°C incubator overnight.

Immunostainings

Cells were washed twice with prewarmed (37°C) PBS. Afterwards, they were fixed and

permeabilized with 4% PFA (Thermo Fisher 28908) and 0.2% tritonX (Sigma-Aldrich

T8787) in PBS for 10 min at room temperature (RT). The cells were washed once with

PBS at RT. Cells were incubated with DAPI (1.7 µg/ml Sigma-Aldrich D9542) and

Phalloidin (Life Technologies A12379, diluted 1:300) in 1% Bovine Serum Albumin

(Sigma-Aldrich A3294) in PBS for 1 h at RT. In the end, the dishes were washed twice

with PBS. Cells were imaged with a Leica DMi8 epifluorescence setup with a ×63 oil

objective (NA 1.4, Leica).

Optical tweezers and force measurement

OT measurements took place on a custom-built inverted microscope[37]. Briefly, a 5 W

1064 nm ytterbium fibre laser (YLM-5-LP, IPG Laser, Germany) was used at a high

power (> 1 W) to minimize intensity fluctuations. Excess power was redirected to a

beam dump and the power at the optical trap itself was 600 mW. The trap was

generated by overfilling the back aperture of an NA 1.2 UPlanSApo water immersion

objective (Olympus, Japan) with the laser beam.

To enable controlled positioning of samples, the sample stage was mounted on top of a

precision piezoelectric stage (P-517.43 and E-710.3, Physik Instrumente, Germany)

with a positioning resolution of 1 nm. The sample stage was illuminated by a white fibre

light source (DC-950 Fiber-Lite, Edmund Optics, USA). White light entered the objective

and then passed through a dichroic mirror (DM) to a 550 nm short pass DM (DMSP550,

Thorlabs, USA). This passed light below 550 nm to the 532 nm notch DM. This mirror

coupled the fluorescence laser into the microscope, and in the process blocked the

portion of incoming white light ranging from 515 to 540 nm. Light that passed through

the notch DM was focused onto a CMOS camera (MC1362, Mikrotron, Germany), which

was used for tracking particles in the optical trap. Light above 550 nm was passed to

the fluorescence imaging camera (optiMOS Scientific CMOS, Teledyne Imaging, USA),

which was used to find beads to trap, locate cells, and record fluorescence images of

tethers.

Forces were measured with the OT by modelling a particle in the trap potential as a

Hookean mass and spring system. By tracking the position of a trapped particle and

multiplying this by a calibrated trap stiffness constant, the force experienced by the

particle was determined. Trap calibration was performed using the power spectrum

method[38]. Postprocessing of the data was done using a custom script in Python 3.6,

which is available on GitHub https://github.com/mchughj33/mem-tension.

OT bead functionalisation

As reported previously[39,40], beads with 2 µm diameter (Kisker PC-S-2.0) were

functionalized with Concanavalin A (ConA, from Canavalia, Sigma-Aldrich, C2272-

10MG) as follows.

10 µL bead solution were added into 100 µL of 2 mg/mL ConA in a tube (low-protein

binding) and incubated for 30 min at 4°C. Beads were washed twice via centrifugation

in 100µL of 5mg/mL bovine serum albumin (BSA, Sigma-Aldrich, A3294) in 200mM Tris

(Sigma, GE17-1321-01) in ddH2O. All centrifugation steps were conducted at 4°C,

10,000 rcf. After the final wash, 1 ml imaging media supplemented with 1% PSF was

added and gently mixed by pipetting.

Membrane tether pulling experiments

15 min prior to each fibroblast measurement, the culture medium was exchanged with

pre-heated (37°C) imaging medium supplemented with 1% PSF. For Xenopus

experiments, Xenopus medium was exchanged with RT imaging medium ~15 minutes

prior to imaging.

In each OT experiment, 2 µl functionalised beads (see above) were added to the culture

media and a bead captured in the trap. To reduce variations in the contact area

between bead and cell[16], all relevant measurement parameters were tightly controlled.

Beads of one size (2.16 µm) were used, ConA coated beads were prepared freshly for

each experiment under temperature-controlled conditions, and the beads were attached

by bringing them into contact with the cells for the same amount of time (5 seconds, see

below).

During the entire process, the position of the bead relative to the laser trap was

recorded in order to measure the force produced by the tether pull. Before attaching the

bead to the cell, a short recording of the Brownian motion of the trapped bead was

taken. The mean force from each recording was used to define the point of zero force in

the subsequent analysis of each respective pull. Afterwards, the bead was carefully

brought into contact with the cell membrane. The image of the bead was monitored for a

change in the interference pattern around the bead, indicating contact between the

bead and membrane. At this point the beads was held stationary for 5 s. Then it was

moved approximately 0.25 μm away from the cell while monitoring the particle tracking

system to see if the mean displacement of the bead from the trap centre was reduced,

which signified that the bead was attached to the membrane.

Beads were then moved away from the cell at 0.1 μm s^{-1} in the case of 3T3 fibroblasts

and 0.5 μm s^{-1} in the case of neurons, leading to the formation of a membrane tether

between the cell and the bead. After reaching the end of the pull, 10 μm for 3T3

fibroblasts and 20 μm for neurons, the bead was held stationary for 30 s. The protocol

for pulling tethers from neurons was based on previous work[41]. Due to high rate of

tether loss during preliminary pulls with fibroblasts, to the pulling velocity was reduced,

leading to a subsequent decrease in the pull distance to avoid reducing experimental

throughput. Pontes et al. showed that varying the velocities for tether pulls in 3T3

fibroblasts within the range of 0.07 μm/s to 1μm/s, did not affect the force results[42]. We

therefore do not expect this velocity change to affect our observations. For each pull,

the peak force was defined as the maximum force value reached, and the steady state

force was determined as the mean of the 30 s period at the end of the pull during which

the bead was held stationary. Analysis was performed using custom analysis scripts

written in python 3.6.

Hydrogels for fibroblasts

Deformable hydroxy-polyacryamide substrates for fibroblast cultures were prepared as

described previously[24,43]. Briefly, for each gel a top coverslip and a glass-bottomed

culture dishe (MatTek P35G-0.20-C and Ibidi 81158 were used for OT and TFM,

respectively) were prepared. Both coverslips were first washed in 70% ethanol and

distilled water, and then dried. Bottom coverslips were wiped with a cotton bud soaked

in 1M NaOH, dried, and then incubated in (3-aminopropyl)trimethoxysilane (APTMS,

Sigma Aldrich, 281778) for 3 min. APTMS was washed off thoroughly in ddH20 and

coverslips were then incubated in a 0.5% solution of glutaraldehyde (Sigma Aldrich,

G6257) for 30 min and then rinsed 3x with distilled water. Top coverslips were treated

for 10 min with Rain-X (Shell Car Care International). Acrylamide (AA, Sigma Aldrich,

A4058) stock solution was made by mixing 500 µL of 40% Acrylamide + 65 µL of 100%

Hydroxyacrylamide (Sigma Aldrich, 697931). Afterwards the gel premix was prepared

by mixing 500 µL Acrylamide stock solution with 250 µL 2% Bisacrylamide (BA, Fisher

Scientific, BP1404-250). The premix was diluted in PBS to achieve the different shear

moduli according to the **table 1.** For TFM gels, 10 µL of PBS was replaced by 10 µl

bead solution (FluoSpheres™ Carboxylate-Modified Microspheres, 0.2 µm, dark red

fluorescent)) and the gel mix was sonicated for 30 s to separate the beads.

Volume of gel	Volume of PBS	Shear modulus (G')

premix		
53 µL	447 µL	0.1 kPa
75 µL	425 µL	1 kPa
150 µL	350 µL	10 kPa

Table 1: Hydroxy-polyacrylamide gel composition used for fibroblasts

Freshly-made premixes were de-gassed under vacuum for 4-6 min. To each 500 µL of

final gel solution, 1.5 µL 0.3% (v/v) N, N, N', N'-tetramethylethylenediamine (TEMED,

Thermo Fisher, 15524–010) and 5 µL 0.1% (w/v) ammonium persulfate (APS, Sigma,

215589) solution were added and the premix was gently mixed. 14 µL or 10 µL of this

mixture were pipetted on the bottom coverslip and 18 mm or 16 mm top coverslips (for

14 µL and 10 µL volume, respectively) were placed on top (with RainX-treated side

facing down). The gels were soaked in PBS for at least 20 min before the top coverslips

were removed. Afterwards, they were washed twice with PBS and then treated with

ultraviolet light for 30 min. Hydrogels were coated as described above.

Hydrogels for neurons

Deformable polyacrylamide gel substrates were prepared as described previously[44,45]

for neuron culture. The coverslips were prepared as described in the hydroxy-acryamide

gel protocol using MatTek dishes as the bottom coverslip. The gel mix was prepared

according to **table 2** to achieve the different shear moduli**.** For TFM gels, the PBS

volume was reduced by 10 µL and replaced by 10 µL bead solution and the gel mix was

sonicated for 30 s to separate the beads.

Volume of AA	Volume of BA	Volume of PBS	Shear modulus (G')
63 µL	10 µL	427 µL	0.1 kPa
63 µL	17.5 µL	419.5 µL	0.3 kPa
94 µL	15 µL	391 µL	1 kPa

529
Table 2: Polyacryamide gel composition used for neurons

Freshly-made gel mixes were briefly de-gassed under vacuum for 4-6 min. To each 500

µL of final gel solution, 1.5 µL 0.3% (v/v) TEMED and 5 µL 0.1% (w/v) APS solution

were added and gently mixed. 7 µL, 14 µL of this mixture were pipetted onto the bottom

coverslip and 18 mm top coverslips were placed on top (with RainX treated side facing

down) for OT experiments and TFM experiments, respectively. The gels were soaked in

PBS for at least 20 min before the top coverslips were removed. The gels were

subsequently functionalized via a 4h treatment with hydrazine hydrate (Sigma Aldrich,

225819), followed by 1 h in 5% acetic acid (Fisher Scientific, 10171460), and then

washed thoroughly in PBS. Afterwards, they were treated with ultraviolet light for 30

min. Hydrogels were coated as described above.

TFM

24 h after plating, fibroblasts were imaged with an inverted microscope (Leica DMi8) at

37°C and 5% CO_2, equipped with a digital sCMOS camera (ORCA-Flash4.0,

Hamamatsu Photonics), an EL6000 illuminator (Leica) and a ×63 oil objective (NA =

1.4, Leica). Images were acquired with Leica LAS X software. Fluorescence images of

beads and wide-field images of cells were taken every 2 min for 10 minutes. After image

acquisition, the culture media was exchanged with Trypsin-EDTA (Gibco) to detach

cells from the gel. Reference images of fluorescent beads were taken 15 min after

trypsinization.

16 - 24 h after plating, *Xenopus* eye primordia explants were imaged using an inverted

Nikon microscope (Ti-E at room temperature) equipped with an sCMOS camera (Prime

BSI Scientific, Photometrics), a CoolLED pE-4000, and ×60 oil objective (NA = 1.4,

Nikon). Fluorescence images of beads and wide-field images of cells were taken every

5 min for 1 h. After image acquisition, the culture media was exchanged with Trypsin-

EDTA (Gibco) or 20% sodium dodecyl sulfate solution (Severn Biotech Ltd., 2 0-4002-

01) to detach the explants from the gel. Reference images of fluorescent beads were

taken 15 min after trypsinization or immediately after sodium dodecyl sulfate treatment.

Traction stress maps were calculated for each frame. Traction stresses were averaged

over time (12 frames for neurons*, 5* frames for fibroblasts) for each cell. Postprocessing

of the data and statistical analyses were done with a custom script in Python 3.8 and is

available on GitHub https://github.com/rg314/pytraction.

Statistics

OT data are shown as violin plots, TFM data as box plots. Violin shapes are kernel

density estimations, boxes indicate the range of data that falls between the first and

third quartile, whiskers show the range of data within 1.5 times the interquartile range.

Density estimator width is based on the number of samples at a given value. Median

values for each dataset are marked by a white circle. Lines between violins indicate

significant differences in the samples. The boxplot diagrams were generated using

python seaborn. The borders of the box indicate the quartiles of the dataset, the median

is indicated by a line within the box, and the whiskers include 1.5*interquartile range.

Points that were determined to be outliers were plotted individually. Kruskal-Wallis one-

way analysis of variance (ANOVA) tests followed by a Tukey posthoc test were used for

statistical analysis when comparing multiple groups. When comparing two groups, a

Mann-Whitney test was used.