Regulatory T cells suppress the formation of potent KLRK1 and IL-7R expressing effector CD8 T cells by limiting IL-2

Oksana Tsyklauri[1, 2], Tereza Chadimova[1,2], Veronika Niederlova[1], Jirina Kovarova[3], Juraj Michalik[1], Iva Malatova[3], Sarka Janusova[1], Helene Rossez[4], Ales Drobek[1], Hana Vecerova[1], Virginie Galati[4], Marek Kovar[3], Ondrej Stepanek[1,4*]

[1] Laboratory of Adaptive Immunity, Institute of Molecular Genetics of the Czech Academy of Sciences, Prague, Czech Republic

[2] Faculty of Science, Charles University, Prague, Czech Republic

[3] Institute of Microbiology of the Czech Academy of Sciences, Prague, Czech Republic

[4] Department of Biomedicine, University Hospital and University of Basel, Basel, Switzerland

*Correspondence should be addressed to Ondrej Stepanek:
ondrej.stepanek@img.cas.cz

Abstract

Regulatory T cells (Tregs) are indispensable for maintaining self-tolerance by suppressing conventional T cells. On the other hand, Tregs may promote tumor growth by inhibiting anti-cancer immunity. In this study, we identified that Tregs increase the quorum of self-reactive CD8$^+$ T cells required for the induction of experimental autoimmune diabetes. Their major suppression mechanism is limiting available IL-2, an essential T-cell cytokine. Specifically, Tregs inhibit the formation of a previously uncharacterized subset of antigen-stimulated KLRK1$^+$ IL7R$^+$ (KILR) CD8$^+$ effector T cells, which are distinct from conventional effector CD8$^+$ T cells. KILR CD8$^+$ T cells show a superior cell killing abilities in vivo. The administration of agonistic IL-2 immunocomplexes phenocopies the absence of Tregs, i.e., it induces KILR CD8$^+$ T cells, promotes autoimmunity, and enhances anti-tumor responses. Counterparts of KILR CD8$^+$ T cells were found in the human blood, revealing them as a potential target for immunotherapy.

Introduction

Physiological immune responses aim at invading pathogens, but not at healthy tissues. There are two major principles of controlling self-reactive lymphocytes: clonal deletion (central tolerance) and suppression (peripheral tolerance). FOXP3[+] regulatory T cells (Tregs) represent a major force of peripheral tolerance as they regulate homeostasis and immune responses of conventional T cells [1]. The absence of Tregs in FOXP3-deficient individuals leads to a severe autoimmune condition called immunodysregulation polyendocrinopathy enteropathy X-linked syndrome [2-4]. On the other hand, Tregs inhibit anti-tumor immune responses.

Although most studies have focused on how Tregs suppress conventional CD4[+] T cells, CD8[+] T cells are also regulated by Tregs. At steady-state, Tregs suppress the proliferation of CD8[+] T cells [5] and prevent the spontaneous differentiation of memory CD8[+] T cells into effector cells [6, 7]. Moreover, Tregs suppress antigenic responses of CD8[+] T cells [8-10]. Multiple mechanisms of Treg-mediated suppression of CD8[+] T cells have been proposed, such as reducing the expression of co-stimulatory molecules on antigen-presenting cells (APC) via CTLA4 [7, 9, 11], limiting the availability of IL-2 [8, 9], production of anti-inflammatory cytokines IL-10 [6] and TGFβ [12], and limiting the production of CCL3, CCL4, and CCL5 chemokines by APCs [10]. However, none of these studies investigated the interaction between Tregs and CD8[+] T cells in an autoimmunity model. Thus, it is still unclear which of these mechanisms is the most important in preventing self-reactive CD8[+] T cells from inducing an autoimmune disease.

A single study addressed how Tregs suppress CD8[+] T-cell responses with various affinity to the cognate antigen [11]. The authors concluded that Tregs inhibit CD8[+] T-cell activation by low-affinity, but not high-affinity, antigens. In such a case, Tregs would not correct errors of central tolerance by suppressing highly self-reactive CD8[+] T cells that escape negative selection in the thymus. Their tolerogenic role would be limited to increasing the antigen-affinity threshold in the periphery [13] by suppressing positively selected CD8[+] T cells with intermediate to low affinity to self-antigens.

In this study, we focused on Treg-mediated peripheral tolerance, which prevents self-reactive CD8[+] T cells from inducing an autoimmune pathology. We found that Tregs increase the number of self-reactive CD8[+] T cells required to trigger experimental diabetes across a wide range of their affinities to self-antigens. In addition, we observed that excessive IL-2 prevents Tregs from suppressing CD8[+] T cells in vitro as well as during autoimmune and anti-tumor responses in vivo. Moreover, Treg depletion, as well as high IL-2 levels, induce the expansion of an unusual population of potent cytotoxic KLRK1[+] IL7R[+] CD8[+] T cells. Altogether, our data provide strong evidence that the major mechanism of Treg suppression of CD8[+] T cells is limiting the availability of IL-2.

Results

Tregs increase the quorum of self-reactive CD8[+] T cells for inducing an autoimmune pathology

To study the potential role of FOXP3[+] Tregs in the autoimmune response of self-reactive T cells, we employed a model of experimental autoimmune diabetes based on the transfer of ovalbumin (OVA)-specific OT-I T cells into double-transgenic mice expressing ovalbumin in pancreatic β-cells [14] and diphtheria toxin receptor (DTR) in FOXP3[+] Tregs. Most of the experiments were performed using RIP.OVA DEREG mice carrying a random insertion of a transgene encoding for DTR-GFP fusion protein under the control of *Foxp3* promoter [15], but some experiments were performed using *Foxp3*[DTR] RIP.OVA strain with *DTR-GFP* knocked-in into the 3' untranslated region of the *Foxp3* locus [16]. Subsequent priming of OT-I T cells induced the formation of effector T cells and eventual destruction of pancreatic β-cells. Administration of diphtheria toxin (DT) selectively depleted Tregs uncovering their role in this autoimmune pathology. The major advantages of this model are: i) known etiology – the pathology is triggered by self-reactive CD8[+] T cells, ii) control of key parameters (number of transferred OT-I T cells, affinity of the priming antigen, presence of Tregs etc.), and iii) the bypass of central tolerance, allowing us to study the mechanisms of peripheral tolerance separately from the central tolerance.

The induction of diabetes in the majority of Treg-replete mice required 10^6 adoptively transferred OT-I T cells followed by priming with OVA peptide and LPS (Fig. 1A-C). This revealed efficient mechanisms of peripheral tolerance preventing the self-reactive T cells from destroying the insulin-producing cells. After the depletion of Tregs, the number of transferred OT-I T cells, sufficient to induce diabetes in the vast majority of animals, dropped dramatically (Fig. 1B-C). Accordingly, the expansion of OT-I T cells was enhanced in Treg-deficient mice compared with Treg-replete mice (Fig. 1D, Fig. S1A-B). When 10^6 OT-I T cells were transferred, 50 % of Treg-depleted mice developed diabetes, even when they were primed with LPS alone without the cognate peptide (Fig. 1E, Fig. S1C). This was possible because the small amount of endogenous OVA was sufficient to prime OT-I T cells, as revealed by comparing their expansion and CD44[+] CD62L[-] effector T-cell formation in RIP.OVA, *Foxp3*[DTR] RIP.OVA, and *Foxp3*[DTR] mice (Fig. 1F, Fig. S1D). Overall, these results revealed that Tregs represent an important mechanism of peripheral tolerance that prevents the induction of diabetes by self-reactive CD8[+] T cells.

To investigate how Tregs suppress priming of self-reactive T cells by antigens with various affinities, we applied a priming protocol based on the adoptive transfer of bone marrow-derived dendritic cells (DCs) loaded with OVA peptide ($K_D \sim 50$ μM) or its variants recognized by OT-I T cells with lower affinity; Q4R7 ($K_D \sim 300$ μM) or Q4H7 ($K_D \sim 850$ μM) [17]. In contrast to OVA peptide and LPS priming, as few as 10^3 transferred OT-I T cells were able to induce autoimmune diabetes in RIP.OVA mice upon DC-OVA

priming (Fig. 2A). When 3×10^2 OT-I T cells were transferred, all Treg-depleted mice, but only one third of Treg-replete mice, were diabetic (Fig. 2A). Accordingly, the absence of Tregs increased the susceptibility to diabetes upon priming with DC-Q4R7, although as many as 3×10^4 OT-I T cells were required to induce diabetes in most Treg-depleted hosts (Fig. 2B). Along this line, Treg-replete RIP.OVA mice were resistant to 10^6 OT-I T cells primed with DC-Q4H7, whereas most Treg-depleted mice manifested diabetes (Fig. 2C). The expansion of OT-I T cells, their expression of IL-2Rα (CD25; a subunit of the high-affinity IL-2 receptor) and KLRG1 (a marker of short-lived effector cells), and the absolute number of KLRG1$^+$ OT-I T cells in the spleen were greater in Treg-depleted mice than in Treg-replete RIP.OVA mice on day six post-immunization with DC-OVA, -Q4R7, or -Q4H7 (Fig. 2D-F, EV2A-C). Please, note that different numbers of OT-I cells were transferred for immunizations with different peptides in these experiments. DCs not loaded with any peptide did not induce diabetes with one exceptional mouse (Fig. S2D-E). When we titrated the amount of transferred OT-I T cells, we observed that the hierarchy of the biological potencies of high-, intermediate-, and low-affinity antigens was preserved in Treg-deficient and Treg-replete hosts (Fig. S2F). Collectively, these results showed that Tregs increase the quorum of self-reactive CD8$^+$ T cells required for inducing autoimmune diabetes, irrespective of the affinity of the priming antigen.

To elucidate, whether the physiological activity of Tregs is a limiting factor of self-tolerance, we selectively expanded the Treg compartment prior to the diabetes induction using complexes of IL-2 and anti-IL-2 mAb JES6-1A12 (JES6) [18, 19] (Fig. 2G). We observed that the expanded Tregs prevented DC-OVA-induced diabetes in most animals (Fig. 2H), showing that enhancing the overall Treg activity increases self-tolerance and prevents the induction of the experimental diabetes.

Tregs suppress self-reactive CD8$^+$ T cells in the absence of conventional CD4$^+$ T cells

Tregs could inhibit self-reactive CD8$^+$ T cells directly or via the suppressing a by-stander help of CD4$^+$ T cells. To address this question, we used T-cell deficient $Cd3\varepsilon^{-/-}$ RIP.OVA mice as hosts. First, we adoptively transferred 10^6 polyclonal OVA-tolerant CD8$^+$ T cells isolated from RIP.OVA mice to replenish the CD8$^+$ T-cell compartment in the $Cd3\varepsilon^{-/-}$ RIP.OVA mice. One week later, we transferred conventional (*Foxp3-GFP$^-$*) or Treg (*Foxp3-GFP$^+$*) CD4$^+$ T cells from DEREG RIP.OVA mice. A control group of mice did not receive any CD4$^+$ T cells. Finally, we transferred OT-I T cells and subsequently primed them with DC-OVA (Fig. 3A). We observed that conventional CD4$^+$ T cells, albeit presumably tolerant to OVA, increased the incidence of experimental diabetes and accelerated its onset (Fig. 3B-C, EV3A). In contrast, Tregs reduced the incidence of diabetes compared with mice receiving no CD4$^+$ T cells (Fig. 3B-C, EV3A). Conventional CD4$^+$ T cells enhanced the expansion of OT-I T cells and increased the number of KLRG1$^+$ OT-I T cells, whereas Tregs elicited the opposite effect (Fig. 3D-E, Fig. S3B-C), as revealed on day eight post-immunization. The role of Tregs was even more apparent on day 14 post-immunization, which

corresponded to the typical onset of diabetes in the absence of conventional CD4[+] T cells (Fig. 3F-G, Fig. S3D-E).

Overall, these experiments showed that conventional CD4[+] T cells provide a by-stander help to self-reactive CD8[+] T cells, whereas Tregs directly suppress priming of CD8[+] T cells.

Tregs suppress self-reactive CD8[+] T cells via limiting IL-2

One of the Treg-mediated effects observed in the previous experiments was the downregulation of IL-2Rα in CD8[+] T cells. IL-2Rα expression on T cells can be induced by antigen stimulation or by IL-2 itself in a positive feedback loop [20]. Accordingly, downregulation of IL-2Rα in OT-I T cells in the presence of Tregs might be a consequence of the limited availability of IL-2. We observed significantly higher levels of IL-2Rα and an IL-2 signaling intermediate, pSTAT5 in OT-I T cells in Treg-deficient mice early after priming (Fig. 4A-B, Fig EV4A-B). These results were reproduced when congenic Ly5.1 OT-I T cells were used (Fig. S4C-E). These results showed that Tregs limit IL-2 signaling in self-reactive CD8[+] T cells in vivo.

Tregs efficiently suppressed proliferation and IL-2Rα expression in CD8[+] T cells ex vivo (Fig. 4C). An addition of excessive IL-2 abolished this suppression (Fig. 4C). To address whether Tregs can suppress self-reactive CD8[+] T cells in the excess of IL-2 in vivo, we performed the diabetic assay with IL-2/anti-IL-2 mAb complexes (IL-2ic) provided during OT-I priming (Fig. 4D). IL-2ic have markedly increased biological activity in comparison to IL-2 alone [21]. We used IL-2ic with JES6-1A12 antibody clone, which selectively promotes IL-2 signaling in T cells expressing high-affinity IL-2Rαβγ trimeric IL-2 receptor, or with the antibody clone S4B6, which promotes IL-2 signaling irrespective of IL-2Rα expression [21, 22]. Both clones of IL-2ic broke the peripheral tolerance and induced the onset of autoimmune diabetes similarly to the Treg-depletion (Fig. 4E-F). IL-2ic promoted the formation of OT-I KLRG1[+] effector cells (Fig. 4G), despite a dramatic IL-2-dependent expansion of Tregs at the same time (Fig. 4H). Overall, these results showed that Tregs are unable to suppress CD8[+] T cells in the excess of IL-2 signal, although Tregs themselves expand dramatically, which should augment their overall suppressive capacity with the exception of the limiting IL-2 mechanism. This supports the hypothesis that the major mechanism of Treg-mediated suppression of self-reactive CD8[+] T cells is the reduction of IL-2 availability. This IL-2-dependent mechanism might be specific for the suppression of CD8[+] T cells, since antigen-activated CD8[+] T cells were much more responsive to IL-2 than activated CD4[+] T cells (Fig. S4G).

Tregs suppress formation of a unique subset of CD8[+] effector T cells

To study how Treg-mediated suppression alters the gene expression profiles of activated CD8$^+$ T cells, we analyzed the transcriptomes of OT-I T cells primed in the presence or absence of Tregs on a single-cell level. IL-2 responsive genes were upregulated in OT-I T cells primed in the Treg-depleted mice, including those encoding the key cytotoxic molecules granzyme B and perforin, as well as IL-2Rα (Fig. 5A, Fig. S5A-B). Unsupervised clustering high-lighted five distinct activated OT-I T cell subsets (Fig. 5B). Based on their gene expression profiles (Fig. S5C, Table S1) and comparison with CD8$^+$ T-cell subsets formed during an antiviral response (Fig. S5D), we identified clusters 0 and 1 as early effector T cells, cluster 2 as memory T cells precursors, and cluster 3 as precursors of exhausted cells. Cluster 4, on the other hand, did not match to any established CD8$^+$ T-cell subset. It was characterized by the expression of a natural killer cell activation receptor gene *Klrk1* (alias *Nkg2d*), *Gzmb* (Fig. 5C, EV5C) and by the overall similarity to the NK cell gene expression profile (Fig. S5E). Interestingly, these cells occurred almost exclusively in the absence of Tregs (Fig. 5D-E). This cluster could be further divided into two subsets. The larger of them expressed *Ilr7a*, *Cd103*, and natural killer cell markers *Ifitm1-3* and *Cd7*, but not a typical effector T-cell marker *Cd49d (Itga4)*, which encodes a subunit of the integrin VLA4 (Fig. 5F-H, Fig. S5C). We refer to these cells as <u>K</u>LRK1$^+$ <u>IL</u>-7<u>R</u>α$^+$ (KILR) effector CD8$^+$ T cells in this study, reflecting their co-expression of typical cytotoxic effector genes (*Gzmb, Prf1*) together with markers of NK cells (*Klrk1*) or T-cell memory (*Il7r*) genes. In the next step, we validated the scRNAseq results using flow cytometry. This confirmed that KLRK1$^+$ IL-7R$^+$ CD49d$^-$ CD8$^+$ T cells are ~20-fold more frequent in the absence of Tregs during the priming than in Treg-replete conditions (Fig. 5I). This experiment also confirmed that part of these KLRK1$^+$ IL-7R$^+$ CD49d$^-$ cells expressed CD103 (Fig. S5F). Moreover, the gene expression profile of FACS-sorted populations of OT-I T cells primed in the presence or absence of Tregs was consistent with the original scRNAseq data (Fig. S5G).

KILR effector CD8$^+$ T cells have strong cytotoxic properties

In the next step, we tested whether Treg depletion induces KILR effector CD8$^+$ T cells in polyclonal mice. Although Treg deficiency increased the frequency of KILR effector CD8$^+$ T cells, they were present at low numbers (Fig. 6A, Fig. S6A-B). We hypothesized that only the combination of antigenic and IL-2 signals can effectively induce KILR effector CD8$^+$ T cells. To address this hypothesis, we directly treated the OT-I Rag2$^{-/-}$ mice with OVA and/or IL-2ic. Indeed, the combination of IL-2 administration and antigenic stimulation efficiently induced KILR effector CD8$^+$ T cells in OT-I mice (Fig. 6B-E), which were characterized by high GZMB and IL-7Rα expression, whereas the stimulation with IL-2 or antigen alone failed to induce the complete KILR phenotype.

To assess the cytotoxic properties of these cells, we compared the ability of KLRK1$^+$ and conventional KLRK1$^-$ effector T cells from OT-I mice treated with OVA + IL-2ic, and KLRK1$^+$ T cells from mice treated

only with OVA to kill splenocytes loaded with their cognate antigen in vivo (Fig. S6C). KLRK1[+] T cells induced by the combination of the antigen and IL-2ic showed the most potent cytotoxic activity on per cell basis (Fig. 6F, Fig. S6D-E).

To identify putative human counterparts of KILR effector CD8[+] T cells, we generated a human blood CD8[+] T-cell atlas by integrating publicly available single cell transcriptomic datasets from healthy donors, because CD8[+] T cell data from Treg-deficient patients are not available. After removing MAIT cells (Fig. S6F, cluster 6), we identified naïve, memory, and effector T cells, and the population expressing some KILR effector CD8[+] T-cell signature genes such as NK markers (*KLRD1*, *IFITM3, CD7)* and *IL7R* (Fig. 6G-H, Fig. S6G, Table S2). Similarly to mouse KILR effector CD8[+] T cells, this human T cell subset was enriched for NK signature genes (Fig. S6H). Human KILR-like T cells constituted for ~1-10% of all CD8[+] T cells (Fig. 6I). These cells showed no signs of clonal expansion, but expressed rearranged αβTCR genes, ruling out the possibility that these cells were natural killer cells or another non-T cell subset (Fig. 6J). Because these cells expressed high levels of *IL2RB* (Fig. 6H), a subunit of IL-2 and IL-15 receptors, their gene expression profile was probably modulated by strong IL-2/IL-15 signals, which is in line with the origin of mouse KILR effector T cells. Although human KILR-like T cells expressed cytotoxic genes *GNLY* and *GZMK*, the expression of *GZMA* and *GZMB* was lower in these cells than in the conventional effector cells (Fig. 6H). This probably reflects the fact that these T cells have not been stimulated by IL-2 together with their cognate antigen recently, which is required for the formation of the full KILR phenotype in mice (Fig. 6B-E). Nevertheless, this previously uncharacterized population of human CD8[+] T cells shows apparent similarities to mouse KILR effector CD8[+] T cells induced by supra-physiological IL-2 levels.

Strong IL-2 signal promotes anti-tumor CD8[+] T-cell responses

So far, we have documented that Tregs suppress self-reactive CD8[+] T cells by limiting IL-2 signal in the context of the autoimmune pathology. We hypothesized that excessive IL-2 signaling may override Treg-mediated suppression of tumor-reactive CD8[+] T cells as well. Therefore, we investigated the effect of IL-2/JES6 immunocomplexes which were previously considered tolerogenic, in a BCL1 leukemia model (Fig. 7A). This IL-2ic slowed down the progression of the disease, when co-administrated with a chemotherapeutic drug doxorubicin (Fig. 7B-C, EV7A). This effect was dependent on CD8[+] but not on CD4[+] T cells, as revealed by antibody-mediated depletion of these subsets (Fig. 7B). Depletion of CD4[+] T cells even improved the anti-tumor effect of doxorubicin plus IL-2ic combinational treatment, presumably due to the depletion of Tregs, which could still limit IL-2 availability between the IL-2ic injections. The administration of IL-2ic without the chemotherapy showed no therapeutic effect (Fig. 7B, EV7A), suggesting that the combination of the IL-2 and antigens released from dead tumor cells was required.

We observed very similar effects of combinational therapy of Doxorubicin with IL-2/JES6 in a B16F10 melanoma model (Fig. 7D-E, EV7B). Since IL-2/JES6 highly selectively stimulates IL-2Rα^+ cells, represented mostly by Tregs in naïve unprimed mice, it was considered as an immunotherapeutic tool for the specific expansion of Tregs in vivo with a potential application in the treatment of autoimmune diseases and pre-transplantation care. On the contrary, our data showed that it promotes anti-tumor response of CD8$^+$ T cells, particularly in the combination with immunogenic chemotherapy [23], and that this anti-tumor activity was not counteracted by Tregs.

Discussion

In this study, we investigated how Tregs prevent CD8[+] T-cell mediated autoimmune pathology. Using a well-controlled system based on the transfer of CD8[+] T cells specific for a pancreatic neo-self-antigen, we revealed that regulatory T cells substantially increase the quorum of self-reactive T cells [24] required to induce the autoimmune pathology.

We observed that Tregs do not alter the antigen-affinity discrimination as they suppress both high-affinity and low-affinity CD8[+] T cells, which is in an apparent contrast with a previous study [10] concluding that Tregs suppress low-affinity T-cell responses exclusively. The most possible explanation is that the previous study focused on the role of Tregs in the CD8[+] T-cell response to *Listeria monocytogenes*, where the presence or depletion of Tregs might regulate the kinetics of pathogen clearance and thus the dose of bacterial antigens. Contrary to that, the dose of the priming antigen was constant in our experimental diabetes.

We have identified the reduction of available IL-2 as the major mechanism of Treg-mediated tolerance of CD8[+] T cells. The evidence is that Treg depletion increases IL-2 signaling in these cells, i.e., pSTAT5 levels and expression of IL-2-responsive genes, and that the administration of IL-2 and IL-2ic mimics the absence of Tregs in vitro and in vivo, respectively. Strong exogenous IL-2 signals stimulate the expansion of Tregs and the expression of their effector molecules [25], which should enhance all their potential suppression mechanisms with the notable exception of IL-2 sequestration. However, as this exogenous IL-2 signal mimics the effect of Treg depletion in our model, mechanisms non-targeting IL-2 do not seem to contribute substantially to the suppression of self-reactive CD8[+] T cells. This conclusion is in line with some previous studies of Treg-mediated suppression of effector CD8[+] T cells [8, 9] and with the importance of IL-2Rα expression in self-reactive CD8[+] T cells for the infiltration of pancreatic islets [26].

It has been found that the biological activity of IL-2ic is dictated by the anti-IL-2 mAb clone [21, 22]. IL-2/JES6 immunocomplexes selectively stimulate cells expressing the high-affinity trimeric IL-2 receptor and were considered as an immunosuppressive agent acting via the expansion of Tregs [25]. In contrast, our conclusion that IL-2 restriction is the major mechanism of Treg-mediated suppression of CD8[+] T cells implies that the administration of IL-2/JES6 should release antigen-activated CD8[+] T cells from the Treg control. Indeed, IL-2/JES6 in combination with chemotherapy significantly prolonged the survival of tumor-bearing mice in two different tumor models in a CD8[+] T-cell-dependent manner. Accordingly, the administration of IL-2/JES6 after priming, but not before the transfer of self-reactive T cells, decreased the T-cell quorum for the induction of experimental diabetes. Collectively, these results reveal that a strong sustained IL-2 signal, even if selective for IL-2Rα[+] cells, potentiates antigen-induced CD8[+] T-cell anti-

tumor immunity and autoimmunity despite its concomitant stimulatory effects on Tregs. Recently, a combinational therapy of in vitro expanded polyclonal Tregs and low-dose IL-2 was tested in patients with type I diabetes in a phase I clinical trial [27]. The authors of the study did not observe a benefit of this intervention for the patients but they reported an expansion of Tregs along with activated NK and cytotoxic CD8[+] T cells, which corresponds to our conclusions.

Our finding that IL-2 restriction is the major mechanism of Treg-mediated regulation of self- and tumor-reactive CD8[+] T cells does not exclude the involvement of additional mechanisms regulating other immune cell types, such as conventional CD4[+] T cells. On the contrary, there is substantial evidence that Tregs suppress conventional CD4[+] T cells largely mechanisms not depending on IL-2 sequestration. First, we show that antigen-stimulated CD4[+] T cells are less sensitive to IL-2 signals than CD8[+] T cells in vivo, which limits the potential impact of the IL-2 restriction on them. Second, it has been shown that Tregs lacking the high-affinity IL-2 receptor, implied in IL-2 sequestration, can still control the homeostasis of CD4[+] T cells but not CD8[+] T cells [5]. Third, it has been shown that Tregs suppress conventional CD4[+] T cells by depleting their cognate peptide-MHCII from APC, which is not applicable for the suppression of MHCI-restricted CD8[+] T cells [28]. In contrast, a recent study proposed that self-reactive CD4[+] T cells produce IL-2 locally attracting Tregs to their close proximity, which leads to the suppression of later phases of self-reactive T-cell activation at least partially via limiting IL-2 [29].

Antigenic stimulation of CD8[+] T cells in the presence of excessive IL-2, induced by depletion of Tregs or by administration of exogenous IL-2R agonists, leads to the formation of a previously uncharacterized subset of KILR effector CD8[+] T cells. They express high levels of cytotoxic molecules, such as granzymes, but unlike conventional effector CD8[+] T cells, they also express high levels of IL-7R. Moreover, they express several natural killer receptors, including KLRK1/NKG2D, which is involved in target cell killing [30]. Since these cells show superior cytotoxic properties in vivo, suppression of their formation is probably a major mechanism of Treg-mediated self-tolerance. On the other hand, the existence of the KILR effector gene expression program suggests the possibility that these cells might arise naturally in specific immunological conditions, such as some type of infection. Indeed, it has been shown that the infection with lymphocytic choriomeningitis virus induces a transient drop in Treg numbers [31], which might alleviate the Treg-mediated suppression of virus-specific CD8[+] T cells in the early phase of the immune response. However, we could not find KILR effector CD8[+] T cells in any scRNAseq dataset from the course of mouse infection, suggesting that such KILR effector-inducing conditions would be rather rare.

The expression of IL-7Rα in KILR effector CD8[+] T cells is striking as conventional effector T cells are characterized as IL-7Rα[-], which is associated with their short life span [32, 33]. A previous study showed that antigenic stimulation together with activating antibodies to OX40 and 4-1BB TNF-family receptors

induced IL-7Rα[+] effector CD8[+] T cells [34]. These cells produced high levels of proinflammatory cytokines in an IL-7-dependent manner [34]. Since 4-1BB signaling has been shown to enhance IL-2 production and IL-2Rα expression in T cells [35, 36], it is possible that those IL-7Rα[+] effector cells were induced via a strong IL-2 signal. However, as the gene expression profile of OX40/4-1BB-induced IL-7Rα[+] effector CD8[+] T cells has not been analyzed [34], it is not clear whether or not these cells have other similarities to KILR effector CD8[+] T cells.

Our reanalysis of publicly available datasets of CD8[+] T cells revealed KILR-like CD8[+] T cells in the human peripheral blood. These cells show patterns of KILR T-cell gene expression, but they do not express high levels of cytotoxic molecules, probably because they lack recent antigenic/IL-2 stimulation. Therefore, future studies are needed to resolve their phenotype and function. In any case, the potent cytotoxic capacity and their expansion by IL-2 agonists make KILR effector CD8[+] T cells a promising clinical target in cancer immunotherapy.

Materials and methods

Antibodies, peptides and dyes

Antibodies to the following antigens were purchased from BioLegend and used for flow cytometry: CD4 BV650 (RM4-5, #100545), CD4 Alexa Flour 700 (RM4-5, #100536), CD4 APC-Cy7 (GK15, #100414), CD4 APC (RM4-5, #100516), CD8a BV421 (53-6.7, #100738), CD8a PE (53-6.7, #100708), CD11c Alexa Flour 700 (N418, #117319), CD25 Alexa Fluor 700 (PC61, #102024), CD25 BV605 (PC61, #102036), CD25 BV650 (PC61, #102038), CD25 Pe-Cy7 (PC61, #102016), CD44 BV 650 (IM7, #103049), CD45.1 Alexa Fluor 700 (A20, # 110723), CD45.1 BV650 (A20, #110735), CD45.1 FITC (A20, #110706), CD45.1 PerCP-Cy5.5 (A20, #110728), CD45.2 Alexa Fluor 700 (104, #109822), CD45.2 APC (104, #109814), CD49d APC (R1-2, #103622), CD49d Pe-Cy7 (R1-2, #103618), CD62L FITC (MEL-14, #104406), CD80 PerCP-Cy5.5 (16-10A1, #104722), CD86 Alexa Fluor 700 (GL-1, #105024), CD103 PerCP-Cy5.5 (2 E7, #121416), CD127 PE-Cy7 (A7R34, #135013), TCRβ APC (H57-597, #109212). Antibodies to TCR Vα2 FITC (B20.1, #553288), CD45.1 APC (A20, #558701), CD45.1 PE (A20, #553776), CD49d PE (R1-2, #553157) were purchased from BD Pharmingen. Antibodies to Granzyme B eFluor660 (NGZB, #50-8898-82), FOXP3 PE-Cy7 (FJK-16s, #25-5773-82), FR4 PE/Dazzle594 (12A5, #125016), KLRG1 BV510 (2F1/KLRG1, #138421), KLRK1 PE-eFluor610 (CX5, #61-5882-82), MHC class II (I-A/I-E) FITC (M5/114.15.2, #11-5321-82) were purchased from eBioscience. Anti-TCRβ PE antibody (H57-597, #1146040) was purchased from SONY.

Goat anti-rabbit IgG (H+L) secondary antibody conjugated to Alexa Fluor 647 from Invitrogen (#21245) was used following rabbit anti-mouse phospho-Stat5 (Tyr694) (D47E7) from Cell Signaling (#4322S). Anti-mouse CD3ε antibody (clone 145-2C11, #100302, BioLegend) was used for plate coating.

Anti-CD4 (GK1.5, #BE0003-1) and anti-CD8α (53-6.7, #BE0004-1) depletion antibodies were purchased from Bioxcell (USA).

Anti-mouse IL-2 mAb S4B6 (#BE0043-1) and anti-mouse IL-2 mAb JES6-1A12 (#BE0043) used for preparation of IL-2/S4B6 and IL-2/JES6 complexes, respectively, were purchased from Bioxcell (USA).

OVA (SIINFEKL), Q4R7 (SIIRFERL), and Q4H7 (SIIRFEHL) peptides were purchased from Eurogentec or Peptides&Elephants.

CFSE (#65-0850-84) and Cell Trace Violet (CTV) (#C34557), LIVE/DEAD near-IR (#L10119), and Hoechst 33258 (#H3569) dyes were purchased from Invitrogen.

IL-2/S4B6 and IL-2/JES6 complexes

IL-2/S4B6 and IL-2/JES6 immunocomplexes were described previously [25]. Complexes were prepared by mixing recombinant mouse IL-2 (100 µg/ml; Peprotech) with anti-IL-2 mAb at a molar ratio of 2:1 in PBS. After 15 min incubation at room temperature, the complexes were diluted in PBS into the desired concentration, frozen at -20 °C, and thawed shortly before application.

Mice

All the mice had C57Bl/6J or BALB/C background. DEREG [15], $FOXP3^{DTR}$ [16], RIP.OVA [14], OT-I $Rag2^{-/-}$ [37], OT-II $Rag2^{-/-}$ [38], Ly5.1 [39] strains were described previously. Mice were bred in specific-pathogen-free facilities (Institute of Molecular Genetics of the Czech Academy of Sciences, Prague; Department of Biomedicine, University Hospital, Basel) or in a conventional facility (Institute of Microbiology of the Czech Academy of Sciences, Prague). Animal protocols were performed in accordance with the laws of the Czech Republic and Cantonal and Federal laws of Switzerland, and approved by the Czech Academy of Sciences or the Cantonal Veterinary Office of Baselstadt, Switzerland, respectively.

Males and females were used for the experiments. At the start of the experiment, all mice were 6-11 weeks old. If possible, age- and sex-matched pairs of animals were used in the experimental groups. If possible, littermates were equally divided into the experimental groups. No randomization was performed when the experimental groups were based on the genotype of the mice; otherwise mice were assigned to experimental groups randomly (defined by their ID numbers) prior to the contact between the experimenter and the mice. The experiments were not blinded since no subjective scoring method was used. The estimation of the sample size was based on our previous experience. We typically aimed at having 12 animals per group in three independent experiments. The final number of mice per group depended on the number of available mice with required genotype, number of isolated T-cells for adoptive transfers, and unpredicTable Sents (e.g., mouse death). In cancer experiments, we aimed at 16 mice per group in two independent experiments.

RT-qPCR

2-10 × 10⁴ OT-I T lymphocytes were FACS sorted as CD8⁺ CD45.1⁺ CD49d⁻ KLRK1⁻ cells (naïve), CD8⁺ CD45.1⁺ CD49d⁺ KLRK1⁻ cells (antigen experienced), CD8⁺ CD45.1⁺ CD49d⁺ KLRK1⁺ cells (double positive), or CD8⁺ CD45.1⁺ CD49d⁻ KLRK1⁺ cells (SE). Total RNA was isolated by TRIzol LS (Invitrogen, #10296010) and in-column DNase digestion using RNA Clean & Concentrator Kit (Zymo Research), according to manufacturers' instructions. RNA was stored at -80 ºC or transcribed immediately using RevertAid reverse transcriptase (Thermo Fisher Scientific, #EP0442) with oligo(dT)18 primers according to the manufacturer's instructions. RT-qPCR was carried out using LightCycler 480 SYBR green I master

chemistry and a LightCycler 480 machine (Roche). All samples were measured in triplicates. Median CT values were normalized to a reference gene, Glyceraldehyde-3-Phosphate Dehydrogenase (*Gapdh*). The sequences of used primers are:

Gapdh: F TGCACCACCAACTGCTTAGC, R GGCATGGACTGTGGTCATGAG

mCd7: F TGGATGCCCAAGACGTACA, R TAAGATCCCTTCCAGGTGCC

mIfitm1: F ATGCCTACTCCGTGAAGTCTAGG, R GACAACGATGACGACGATGGC

mGzma: F AAAGGACTCCTGCAATGGGG, R ATCGGCGATCTCCACACTTC

mGzmb: F GGGGCCCACAACATCAAAGA, R GGCCTTACTCTTCAGCTTTAGCA

mGzmk: F AAGGATTCCTGCAAGGGTGA, R ATTCCAGGCTTTTTGGCGATG

mKlrd1: F TCGGTGGAGACTGATGTCTG, R AACACAGCATTCAGAAACTTCC

mIl7r: F AAAGCCAGAGCGCCTGGGTG, R CTGGGCAGGGCAGTTCAGGC

mIl2ra: F AGAACACCACCGATTTCTGG, R GGCAGGAAGTCTCACTCTCG

mCxcr6: F ACTGGGCTTCTCTTCTGATGC, R AAGCGTTTGTTCTCCTGGCT

Enrichment of T lymphocytes

T lymphocytes were enriched by negative selection using the Dynabeads Biotin Binder kit (Invitrogen, #11047), and biotinylated anti-CD19 (clone 1D3, ATCC# HB305), anti-CD4 (clone YTS177), or anti-CD8 antibodies (clone 2.43, ATCC# TIB-210), depending on the experimental setup. Antibodies were produced and biotinylated in house using (+)-Biotin N-hydroxysuccinimide ester (Sigma Aldrich) in bicarbonate buffer. The excessive biotin was separated from the antibody using Sephadex G-25 (Sigma Aldrich). For RT-qPCR experiments, CD8$^+$ T cells were enriched using Dynabeads Untouched Mouse CD8 Cells Kit (Invitrogen, #11417D) according to manufacturer's instructions.

Flow cytometry and cell sorting

Live cells were stained with relevant antibodies on ice. LIVE/DEAD near-IR dye or Hoechst 33258 were used for discrimination of viable and dead cells.

For intracellular staining of FOXP3 and Granzyme B, cells were fixed and permeabilized using Foxp3 / Transcription Factor Staining Buffer Set (#00-5523-00, Invitrogen) according to the manufacturer's instructions. Fixed cells were stained at room temperature for 1 h.

For intracellular staining of pSTAT5, splenic cells were fixed using Fixation/ Permeabilization buffer (Foxp3 / Transcription Factor Staining Buffer Set, #00-5523-00, Invitrogen) at room temperature for 15 min, washed twice with Permeabilization buffer (Foxp3 / Transcription Factor Staining Buffer Set, #00-5523-00, Invitrogen), washed with PBS and stained with anti-pSTAT5 antibody at room temperature overnight. The next day, cells were stained with secondary antibody at room temperature for 1 h.

Flow cytometry was carried out using an LSRII (BD Bioscience) or an Aurora (Cytek). Data were analyzed using FlowJo software (BD Bioscience).

Cell sorting was performed on an Influx or an Aria machines (both BD Bioscience).

ELISA

The anti-idiotypic B1 mAb was produced by the use of B1 hybridoma via the conventional ascites producing approach in parafine oil pre-treated BALB/C mice. It was purified by 45% supercritical antisolvent precipitation (ammonium sulphate) followed by extensive dialysis against distilled water, centrifuged at 12000 g for removal of IgM, and purified by a protein A affinity chromatography. Next, it was biotinylated with Sulpho NHS-biotin reagent (Pierce) according to the manufacturer´s protocol. Blood of mice was taken on day 54 of B-cell leukemia/lymphoma experiment. Blood serum was separated and serially diluted (1:40 – 1:640) in PBS. Plate wells (Costar) were coated with 50 µl of diluted samples and incubated overnight at 4 °C, followed by blocking with 1% gelatin (200 µl per well, 2 h at room temperature). Biotinylated anti-idiotypic B1 mAb was added (20 ng/ml) in a buffer containing 0.5% gelatin, 3% PEG, and 0.1% tween, and plate was incubated for 2 h at room temperature. Next, samples were conjugated with ExtrAvidin−Peroxidase (Sigma-Aldrich) for 1 h at room temperature, and 3,3′,5,5′-tetramethylbenzidine substrate (Sigma-Aldrich) was added for 10 min in dark. Reaction was stopped with 50 µl of 2M H_2SO_4, and absorbance at 450 nm was measured by a Biolisa spectrometer (Bioclin).

Bone marrow-derived dendritic cells

Bone marrow cells were seeded on 100 mm plates (tissue culture untreated) and maintained in DMEM (Sigma Aldrich) containing 10% FBS (GIBCO), 100 U/ml penicillin (BB Pharma), 100 mg/ml streptomycin (Sigma Aldrich), 40 mg/ml gentamicin (Sandoz), and 2% of supernatant from J558 cells producing GM-CSF for 10 days at 5% CO_2 at 37 °C [40]. The cells were split every 2-3 days. On day 10, cells were incubated in the presence of 100 ng/ml LPS (Sigma Aldrich) and 200 nM of indicated peptide for 3 h at 5% CO_2 at 37 °C. Next, plates were incubated with 0.02% EDTA in PBS for 5 min at 5% CO_2 at 37 °C, and harvested. Washed and filtered cells were used for adoptive transfers.

In vitro proliferation assay

CD8$^+$ T lymphocytes from WT mice were FACS sorted, labeled with CTV, and plated into an anti-CD3ε-antibody-coated 48-well plate in IMDM (10% FBS (GIBCO), 100 U/ml penicillin (BB Pharma), 100 mg/ml streptomycin (Sigma Aldrich), 40 mg/ml gentamicin (Sandoz)). Tregs, sorted as CD4$^+$ GFP$^+$ T lymphocytes from DEREG$^+$ mice were added to corresponding wells of the plate in 1:1 ratio with conventional CD8$^+$ T lymphocytes. Recombinant IL-2 (2 ng/ml) was added or not. Cells were incubated at 37°C, 5% CO2 for 72 h and analyzed by flow cytometry.

In vivo proliferation assay

OT-I CD8$^+$ and OT-II CD4$^+$ T cells were isolated from OT-I *Rag2$^{-/-}$* and OT-II *Rag2$^{-/-}$* mice, respectively, using MACS negative selection kits. On day 0, mixture of OT-I CD8$^+$ (0.75 × 10^6 cells) and OT-II CD4$^+$ (1.5 × 10^6 cells) CFSE labeled cells was adoptively transferred into recipient Ly5.1 mice. The next day, recipient mice were immunized with ovalbumin protein i.p. (75 μg/mouse), and after 6 h, they received the first dose of IL-2ic (1.5 μg IL-2 equivalent/dose, i.p). On days 2, 3, and 4 mice received additional IL-2ic doses. On day 5 post-immunization, spleens of Ly5.1 mice were collected and used for flow cytometry analysis.

Treg depletion

In order to deplete Tregs, 0.25 μg of DT (#D0564, Sigma-Aldric) was administered i.p. to DEREG$^+$ RIP.OVA mice and control DEREG$^-$ RIP.OVA mice for two consecutive days.

Model of autoimmune diabetes

The model of autoimmune diabetes has been described previously [13]. Briefly, an indicated number of OT-I T cells isolated from OT-I *Rag2$^{-/-}$* mice were adoptively transferred into a recipient RIP.OVA mouse i.v. On the following day, the recipient mice were immunized with 10^6 of bone marrow-derived dendritic cells loaded with indicated peptide (i.v.) or with 25 μg LPS + 50 μg OVA peptide in 200 μl PBS (i.p.).

Alternatively, *Cd3ε$^{-/-}$* RIP.OVA recipient mice received 10^6 of polyclonal CD8$^+$ T lymphocytes derived from RIP.OVA Ly5.1 donors 8 days prior to immunization. One day prior to immunization, *Cd3ε$^{-/-}$* RIP.OVA mice received 4-8 × 10^5 Tregs (sorted as CD4$^+$ GFP$^+$ TCRβ$^+$) or 10^6 conventional CD4$^+$ T lymphocytes (sorted as CD4$^+$ GFP$^-$ TCRβ$^+$) derived from DEREG RIP.OVA donors.

If IL-2ic were used, 1.5 μg IL-2 equivalent/dose of IL-2/S4B6 or IL-2/JES6 were injected i.p. on day 0, 1, and 2 post-immunization. Alternatively, mice received 5 doses of IL-2/JES6 (2.5 μg IL-2 equivalent) on days -7, -6, -5, -4, -3 prior to the immunization.

Urine glucose was monitored on a daily basis using test strips (GLUKOPHAN, Erba Lachema). Blood glucose was measured using Contour blood glucose meter (Bayer) on a day 7, 8, or 14 post-immunization, depending on the experimental design. The animal was considered diabetic when the concentration of glucose in the urine reached ≥ 1000 mg/dl for two consecutive days.

Induction of KILR effector T cells using IL-2ic

On day 0, recipient mice (OT-I Rag2$^{-/-}$ or RIP.OVA) received 25 µg OVA peptide in 200 µl of PBS and/or 0.75 µg IL-2 equivalent/dose of IL-2/S4B6 in 250 µl PBS i.p. On days 1 and 2, mice received 2 more doses of IL-2/S4B6. On day 3, spleens were collected, and used for flow cytometry analysis or FACS sort.

In vivo killing assay

In vivo CD8$^+$ T cell killing assay was performed as described previously [41] with minor modifications. In short, OT-I *Rag2$^{-/-}$* mice were immunized or not with OVA peptide (25 µg, single dose on day 0, i.p.) and/or IL-2/S4B6 (0.75 µg IL-2 equivalent/dose of IL-2/S4B6, 3 doses, days 0-2, i.p.). On day 3, FACS sorted KLRK1$^+$ CD8$^+$ or KLRK1$^-$ CD8$^+$ cells were injected i.v. to recipient RIP.OVA mice, which had received a mixture of target cells (10^7 OVA-pulsed cells and 10^7 unpulsed target cells) earlier the same day. Target cells were prepared from spleens of Ly5.1 mice. OVA-pulsed cells were prepared via 1 h incubation in RPMI-1640 containing 10% FBS (GIBCO), 100 U/ml penicillin (BB Pharma), 100 mg/ml streptomycin (Sigma Aldrich), 40 mg/ml gentamicin (Sandoz), and 2 µM OVA peptide at 37°C, 5% CO_2, followed by loading with CTV. Unpulsed target cells were incubated in parallel in the medium without OVA peptide, and subsequently loaded with CFSE. On day 4, splenocytes of the recipient mice were analyzed by flow cytometry. Target cells were identified as CD45.1$^+$ cells. Ratio of unpulsed (CFSE$^+$) to OVA pulsed target cells (CTV$^+$) was determined and normalized to those of recipients which did not receive OT-I T cells.

Murine B-cell leukemia

On day 0, BALB/C female mice were injected i.p. with 5×10^5 BCL1 cells (a BALB/c-derived leukemia cell line) [42] in PBS. On days 11 and 24 post-inoculation, mice received doxorubicin (Adriblastina) (5 mg/kg in 250 µl PBS, i.v.), followed or not by anti-CD4 or anti-CD8α depletion mAbs (200 µg in 250 µl of PBS for both, i.p.). On days 12, 13, 14, 25, 26, and 27 IL-2/JES6 was administrated to mice (5 µg IL-2 equivalent/dose in 250 µl PBS, i.p.). Survival of mice was monitored from day 30 to day 100. On days 28, 54, and 64-96 blood serum of mice was used for ELISA for antibody against the idiotype of IgM expressed on the BCL1 cells.

B16F10 melanoma

Female C57Bl/6J mice were inoculated s.c. with 5×10^5 B16F10 melanoma cells (day 0). On day 6, mice received doxorubicin (Adriblastina) (8 mg/kg in 250 µl PBS, i.v.) followed or not by anti-CD8α or anti-CD4 depletion mAbs (200 ug in 250 µl of PBS, i.p.). On days 7, 8, and 9 IL-2/JES6 was injected to mice (5 µg IL-2 equivalent/dose in 250 µl PBS, i.p.). Survival of mice was monitored on a daily basis.

ScRNA sequencing

Six weeks old female DEREG⁻ RIP.OVA (n=3) and DEREG⁺ RIP.OVA (n=3) littermate mice were treated with DT (0.250 µg per mouse in 0.5 ml of PBS, i.p.) on days -2 and -1 prior to the immunization, followed by i.v. injection of 5×10^4 OT-I T cells in 200 µl of PBS on day -1. OT-I T cells were obtained from spleens and lymph nodes of Ly5.1 OT-I *Rag2⁻ᐟ⁻* mouse. On day 0, mice received Ly5.1 dendritic cells loaded with OVA (10^6 cells per mouse in 200 µl of PBS). On day 3 post-immunization, mice were sacrificed and spleens were collected. Erythrocytes were lysed with ACK buffer (2 min, on ice), cells were washed and resuspended in PBS/ 2% FBS. Next, cells were stained with LIVE/DEAD near-IR dye, anti-CD8a BV421, anti-CD45.1 PE, and anti-CD45.2 APC antibodies together with TotalSeq-C anti-mouse hashtag antibodies (anti-CD45 clone 30-F11, anti-H-2 clone M1/42, Biolegend, #155869 (MH5), #155871 (MH6), #155873 (MH7), #155875 (MH8), #155877 (MH9) and #155879 (MH10)). Viable CD8a⁺, CD45.1⁺, CD45.2⁻ OT-I cells were FACS sorted. The individual samples were pooled together and with cells coming from an unrelated experiment. These unrelated cells were labeled with unique hashtag antibodies and removed during the analysis. The viability and concentration of cells after sort were measured using the TC20 Automated Cell Counter (#1450102, Bio-Rad). The viability of the cells pre-loading was > 90%. Cells were loaded onto a 10x Chromium machine (10x Genomics) aiming at the yield of 1000 cells per sample and processed with Feature Barcode technology for Cell Surface Protein protocol (#CG000186 Rev D) with the Chromium Single Cell 5' Library & Gel Bead and Chromium Single Cell 5' Feature Barcode Library kits (10x Genomics, #PN-1000014, #PN-1000020, #PN-1000080, #PN-1000009, #PN-1000084). Resulting cDNA libraries were sequenced on a NovaSeq 6000 (Illumina) with the S1 Reagent Kit (100 or 300 cycles, Illumina, #20012865, #20012863).

Analysis of scRNAseq data

The raw scRNA data were mapped to Mouse Reference GRCm38 obtained from Ensembl database v102 [43] by 10x Genomics Cell Ranger 5.0.0 [44]. The same software was also used to create the employed mouse transcriptome reference. Default parameters were kept.

The hashtag sequences were mapped using the hashtag sequence references by 10x Genomics Cell Ranger 5.0.0. The data were pre-processed using Seurat R package v4.0.3 [45] on R v4.0.4 (https://www.r-project.org/). The cells with less than 200 transcripts were removed. A histogram of cell counts having a

specified number of hashtag reads was computed for each hashtag. To detect the end of an initial slope of a histogram, a function based on descent along the gradient – moving to the neighboring point in the histogram as long as its associated value is lower - was used. Before its application, each histogram was averaged using the sliding window of size n = 5 points. The resulting limit was used to associate or not each cell with the respective sample. Cells marked by multiple hashtags were considered as doublets and excluded from further analysis. Read limits (lim) and the number of recovered cells (#) for individual hashtags were: MH5 (lim 19, #744), MH6 (lim 19, #778), MH7 (lim 24, #789), MH8 (lim 21, #849), MH9 (lim 35, #836), MH10 (lim 25, #878).

The lists of barcodes of each uniquely marked cell were created and used to separate read pairs where the start of first read R1 matches one of barcodes exactly from those with different barcodes and consequently either originating from the cells from unrelated experiment, from cell doublets or insufficiently marked cells. These reads were mapped again using 10x Genomics Cell Ranger 5.0.0 to generate data used further for downstream analysis. Cells with less than 200 transcripts and/or more than 15 % of transcripts mapping to mitochondrial genes were removed. Mitochondrial genes, TCRα and TCRβ V(D)J-genes, ribosomal genes and genes whose transcripts were detected in less than three cells were excluded. Log normalization (scale factor = 1×10^4), scaling, identification of variable features (2000 variable features), dimensional reduction (PCA and UMAP with top 50 and 15 principal components respectively, 40 nearest neighbors for UMAP), identification of nearest neighbors in the reduced space ('rann' algorithm) and Louvain clustering (resolution = 0.8) were performed using the Seurat R package v4.0.3 [45] on R v4.0.4 (https://www.r-project.org/). These steps allowed to identify the low-quality cells and contaminating cell types that were removed along with cells with more than 7.5% of transcripts mapping to mitochondrial genes. In total, 4043 cells passed the QC steps. Afterwards, all steps starting from and including log-normalization were repeated using slightly different parameters (800 variable features, 12 top principal components for both UMAP and PCA dimensional reductions and 30 nearest neighbors for UMAP, resolution = 0.3 for Louvain clustering; other parameters stayed the same). Cell cycle scores for S phase and G2/M phases were regressed out in the scaling step. Projection of clusters on a reference dataset of acute and chronic viral infection CD8+ T cells was done using the ProjecTILs R package v0.6.0 [46]. Natural killer cell signature was calculated for each cluster using the AddModuleScore function from the Seurat package with the default parameters. Signature genes were selected from the Molecular Signatures Database v7.5.1 (Systematic name M4838 and M5669 for murine and human datasets, respectively). Signature genes that were not detected in our datasets were filtered out. For the separate analysis of KILR effector CD8+ T cells, cluster 4 was extracted from the original data and re-clustered again with adjusted parameters.

The code for the cell filtration to hashtags, barcode extraction and whole downstream analysis is accessible on GitHub (https://github.com/Lab-of-Adaptive-Immunity/Supereffectors_scRNAseq).

Gene set enrichment analysis

Lists of IL-2 responsive genes were obtained from literature [47, 48]. Fold changes between DEREG[+] and DEREG[-] samples were calculated with the Seurat R package v4.0.3 [45]. Gene set enrichment analysis (GSEA) analysis was performed using the fgsea R package v1.16.0 [49]. Genes with similar fold changes were ranked in a random order.

Building and analysis of Human CD8[+] T-cell atlas

The Human CD8[+] Atlas was built from 14 different data sets previously mapped to human genome reference GRCh38 by 10x Genomics and downloaded from their support site (https://support.10xgenomics.com/) either as raw feature matrices or, if not available for given data set, as filtered feature matrices. For data sets with available V(D)J information their list was downloaded either as list of filtered annotated contigs or, if not available for given data set or it was originally procesed by Cell Ranger 5.0.0 or higher, as list of all annotated contigs. The following data sets were used: *CD8[+] T cells of Healthy Donor 1*, Single Cell Immune Profiling Dataset by Cell Ranger 3.0.2, 10x Genomics, (2019, May 9); *CD8[+] T cells of Healthy Donor 2*, Single Cell Immune Profiling Dataset by Cell Ranger 3.0.2, 10x Genomics, (2019, May 9); *CD8[+] T cells of Healthy Donor 3*, Single Cell Immune Profiling Dataset by Cell Ranger 3.0.2, 10x Genomics, (2019, May 9); *CD8[+] T cells of Healthy Donor 4*, Single Cell Immune Profiling Dataset by Cell Ranger 3.0.2, 10x Genomics, (2019, May 9); *10k Human PBMCs with TotalSeq-B Human TBNK Antibody Cocktail, 3' v3.1*, Single Cell Gene Expression Dataset by Cell Ranger 6.0.0, 10x Genomics, (2021, March 31); *10k Human PBMCs Multiplexed, 2 CMOs - Inputs/Library*, Single Cell Gene Expression Dataset by Cell Ranger 6.0.0, 10x Genomics, (2021,March 2); *5k Peripheral blood mononuclear cells (PBMCs) from a healthy donor (v3 chemistry),* Single Cell Gene Expression Dataset by Cell Ranger 3.0.2, 10x Genomics, (2019, May 29); *10k PBMCs from a Healthy Donor - Gene Expression and Cell Surface Protein*, Single Cell Gene Expression Dataset by Cell Ranger 3.0.0, 10x Genomics, (2018, November 19); *8k PBMCs from a Healthy Donor*, Single Cell Gene Expression Dataset by Cell Ranger 2.1.0, 10x Genomics, (2017, November 8); *PBMCs of a Healthy Donor (v1)*, Single Cell Immune Profiling Dataset by Cell Ranger 3.1.0, 10x Genomics, (2019, July 24); *Human T cells from a Healthy Donor, 1k cells - multi (v2)*, Single Cell Immune Profiling Dataset by Cell Ranger 5.0.0, 10x Genomics, (2020, November 19); *Human PBMC from a Healthy Donor, 10k cells - multi (v2),* Single Cell Immune Profiling Dataset by Cell Ranger 5.0.0, 10x Genomics, (2020, November 19); *Human PBMC from a Healthy Donor, 1k cells (v2)*, Single Cell Immune Profiling Dataset by Cell Ranger 4.0.0, 10x Genomics, (2020, August

25); and *PBMCs of a healthy donor - 5' gene expression and cell surface protein*, Single Cell Immune Profiling Dataset by Cell Ranger 3.0.0, 10x Genomics, (2018, November 19).

Cells with less than 200 transcripts, above-average transcript count plus identified as doublets by scds R package v1.4.0 [50] on R v4.0.4 (https://www.r-project.org/), and/or cells with more than 10% transcripts mapping to mitochondrial genes were removed. The genes equivalent to those removed in our mouse data set plus V(D)J genes of TCRγ and TCRδ genes were excluded. Centered log-ratio normalization of cell surface protein counts for data sets that have them, log-normalization (scale factor = 1×10^4) and identification of variable features for each data set (2500 variable features) were performed using Seurat R package v4.0.0 [45]. The same package was used for the integration of all data sets and subsequent scaling, dimensional reduction (PCA and UMAP with top 20 principal components), nearest neighbors identification and Louvain clustering (resolution = 0.4) of the resulting data set. The newly emerging cluster of cells with below-average gene count and above-average proportion of transcripts mapping to mitochondrial genes was removed. In total, 140,564 cells passed the QC steps. Afterwards, the previous steps starting from and including both normalizations were repeated using the same parameters. A cluster containing 5,160 MAIT cells was identified using differential expression analysis and V(D)J information, which was kept for some analyses, but removed for others. OPTICs method from dbscan R package v1.1-6 [51] was applied (parameters minPts = 500 and eps_cl = .55 for data both with and without MAIT cells) to generate the final clustering.

The code for building the atlas and its whole analysis is available on GitHub (https://github.com/Lab-of-Adaptive-Immunity/HS-CD8-Atlas).

Statistical analysis

Statistical analysis was performed using an appropriate test for given type of data using GraphPad Prism 5.0 or R. Quantitative data from mice were usually tested using non-parametric Mann-Whitney test or, for a comparison of more than two sample groups, Kruskal-Wallis test with Dunn's posttest, if required. Unpaired Student's t test was used only once for the statistical analysis of the size of cell clusters from the scRNAseq experiments, where the limited number of biological replicates (n=3) did not allow a non-parametric test with reasonable statistical power. However, the results were subsequently confirmed using independent assays with more biological replicates. Mann-Whitney test with Bonferroni correction (adjusted p-value) for multiple comparisons was used for the statistical analysis of differentially expressed genes in the scRNAseq data. The survival/disease-free curves were analyzed by Log-rank (Mantel-Cox) test. The statistical test is indicated for each experiment in Figure Legends. All tests were two-tailed. When applicable, the correction for multiple comparisons was applied.

Data availability

The scRNAseq data generated in this study were deposited in the Gene Expression Omnibus (GSE183940).

Author Contribution

OS conceived the study. OT, TC, JK, IM, SJ, HR, AD, HV, VG, MK, and OS performed experiments. OT, TC, JK, IM, SJ, MK, and OS analyzed experiments. VN and JM analyzed scRNAseq data. OT, VN, JM, and OS wrote the first draft. All authors reviewed and commented on the manuscript.

Conflict of Interest

All authors declare that they have no conflict of interest.

Acknowledgement

This project has received funding from the European Union's Horizon 2020 research and innovation programme under grant agreement No 802878 (ERC Starting Grant FunDiT to OS) The project was supported by the Czech Science Foundation (19-03435Y to OS and 18-12973S to MK), Purkyne Fellowship of the Czech Academy of Sciences (to OS), Research Fund for Young Scientists at the University of Basel (DMS2336 to OS), the Institute of Molecular Genetics of the Czech Academy of Sciences (RVO 68378050 to OS), the Institute of Microbiology of the Czech Academy of Sciences (RVO 61388971 to MK), and Charles University Grant Agency (1706119 to OT and TC). The animal facility of the Institute of Molecular Genetics is a part of the Czech Center for Phenogenomics supported by the Czech Ministry of Education, Youth and Sports and the European Regional Development Fund (LM2015040, LM2018126, OP RDI CZ.1.05/2.1.00/19.0395, OP RDI BIOCEV CZ.1.05/1.1.00/02.0109).

VN and SJ are students of the Faculty of Science, Charles University in Prague.

Figure Legends

Figure 1

A. Scheme of RIP.OVA diabetes model. OT-I T cells were transferred into $Foxp3^{DTR}$ RIP.OVA or DEREG RIP.OVA mice (Treg-depleted with DT) and RIP.OVA controls (Treg-replete). The next day, mice were immunized with OVA peptide and LPS. Urine glucose levels were monitored on a daily basis for 14 days.

B-D. Diabetes was induced in RIP.OVA mice as described in A. **B.** Percentage of diabetic mice is shown. Number of diabetic mice and total number of mice per group is indicated on top of each column. Number of transferred OT-I T cells is indicated for each group. **C.** Glucose concentration in blood on day 7 post-immunization is shown. RIP.OVA n=19, DEREG RIP.OVA n=9, $Foxp3^{DTR}$ RIP.OVA n=9. One mouse from $FOXP3^{DTR}$ RIP.OVA group died before the measurement (shown as "x"). **D.** Diabetes was induced in RIP.OVA (Ly5.1/Ly5.2, n=4) and $Foxp3^{DTR}$ RIP.OVA (Ly5.1, n=7) mice using 0.5×10^6 OT-I T cells. Spleens were collected on day 5 and analyzed by flow cytometry. Percentage of OT-I T cells in CD8$^+$ T-cell population is shown.

E-F. RIP.OVA (Ly5.1/Ly5.2), $Foxp3^{DTR}$ RIP.OVA (Ly5.1), DEREG RIP.OVA (Ly5.1/Ly5.2), and $Foxp3^{DTR}$ (Ly5.1) mice were treated as shown in A, with the exception that on day 0, mice were stimulated with LPS only (without OVA peptide). **E.** Percentage of diabetic mice is shown. Number of diabetic mice and total number of mice per group is indicated on top of each column. Number of transferred OT-I T cells is indicated for each group. **F.** Diabetes was induced using 0.5×10^6 OT-I T cells. Spleens were collected on day 5 and analyzed by flow cytometry. Left: percentage of OT-I T cells in CD8$^+$ T-cell population is shown. RIP.OVA n=5, $Foxp3^{DTR}$ RIP.OVA n=9, $Foxp3^{DTR}$ n=8. Right: percentage of effector cells defined as CD44$^+$ CD62L$^-$ in OT-I T-cell population is shown. RIP.OVA n=4, $Foxp3^{DTR}$ RIP.OVA n=9, $Foxp3^{DTR}$ n=8.

Statistical significance was calculated by Kruskal-Wallis test (p-value is shown in italics) with Dunn's post-test (* <0.05, **<0.01) for comparison of three groups (C, F), or two-tailed Mann-Whitney test for comparison of two groups (D, p-value shown in italics). Median is shown.

Figure 2

A-C. Treg-depleted DEREG$^+$ RIP.OVA mice and control DEREG$^-$ RIP.OVA mice received indicated numbers of OT-I T cells (10^3 or 3×10^2 OT-I T cells in **A**, 3×10^4 OT-I T cells in **B**, 10^6 OT-I T cells in **C**). The next day, mice were immunized with DC loaded with an indicated peptide (OVA in **A**, Q4R7 in **B**, Q4H7 in **C**). Urine glucose level was monitored on a daily basis for 14 days. **A.** Left: Percentage of diabetic mice is shown. Number of diabetic mice and total number of mice per group is indicated on top of each

column. Middle: Survival curve. Number of mice per group is indicated. Right: Blood glucose concentration on day 7 post-immunization. **B-C.** Left: Survival curve. Number of mice per group is indicated. Right: Blood glucose concentration on day 7 post-immunization.

D-F. Diabetes was induced in DEREG⁻ RIP.OVA and DEREG⁺ RIP.OVA mice similarly to A-C. On day 6, spleens were collected and analyzed by flow cytometry. Percentage of OT-I T cells among CD8⁺ T cells, count of KLRG1⁺ OT-I T cells, and count of IL-2Rα⁺ OT-I T cells are shown. **D.** Diabetes was induced using DC-OVA and 10^3 OT-I T cells. Left: DEREG⁻ n=9, DEREG⁺ n=10. Middle: DEREG⁻ n=7, DEREG⁺ n=9. Right: n=5 mice per group. **E.** Diabetes was induced using DC-Q4R7 and 3×10^4 OT-I T cells. Left: DEREG⁻ n=13, DEREG⁺ n=10. Middle: DEREG⁻ n=10, DEREG⁺ n=9. Right: DEREG⁻ n=6, DEREG⁺ n=7. **F.** Diabetes was induced using DC-Q4H7 and 10^6 OT-I T cells, n=8 mice per group.

G-H. RIP.OVA mice were treated or not with IL-2/JES6 for five consecutive days. Two days after the last dose, the mice received 10^3 OT-I T cells, and the next day they were immunized with DC-OVA. Urine glucose level was monitored on a daily basis for 14 days. **G.** Experimental scheme. **H.** Left: Survival curve. Right: Blood glucose concentration on day 7 post-immunization. n=12 mice per group.

Statistical significance was calculated by Log-rank (Mantel-Cox) test (survival) or two-tailed Mann-Whitney test (glucose concentration and flow cytometry analysis). P-value is shown in italics. Median is shown.

Figure 3

Cd3ε⁻ᐟ⁻ RIP.OVA mice received 10^6 polyclonal CD8⁺ T cells from Ly5.1 RIP.OVA mice. After seven days, they received either 10^6 conventional CD4⁺ T cells (GFP⁻), or $0.4-1 \times 10^6$ Tregs (GFP⁺) from DEREG⁺ RIP.OVA mice, or no CD4⁺ T cell transfer (NT). Next, 250 OT-I T cells were adoptively transferred to all the recipients. Next day, mice were immunized with DC-OVA.

A. Experimental scheme.

B-C. Urine glucose level was monitored on a daily basis until day 21 post-immunization. **B.** Survival curve. Number of mice is indicated. **C.** Blood glucose concentration on day 8 and day 14 post-immunization is shown. Day 8: NT n=28, Conv. CD4⁺ n=25, Tregs n=15. Day 14: NT n=15, Tregs n=8.

D-G. On day 8 or day 14, spleens were collected and analyzed by flow cytometry. Percentage of OT-I T cells among CD8⁺ T cells (**D, F**), and count of KLRG1⁺ OT-I T cells (**E, G**) are shown. Day 8: NT n=25, Conv. CD4⁺ n=15, Tregs n=12. Day 14: NT n=15, Tregs n=7.

Statistical significance was calculated by Kruskal-Wallis test (p-value is shown in italics) with Dunn's post-test (* <0.05, **<0.01, ***<0.001) for comparison of three groups, or two-tailed Mann-Whitney test for comparison of two groups (p-value is shown in italics). Median is shown.

Figure 4

A-B. OT-I T cells (5×10^4) were transferred into Treg-depleted DEREG$^+$ RIP.OVA mice and control DEREG$^-$ RIP.OVA mice. The next day, mice were immunized with DC-OVA. On day 3 post-immunization, spleens were collected and analyzed by flow cytometry. OT-I T cells were identified as OVA-tetramer and TCRVα2 double positive. **A.** IL-2Rα expression on OT-I T cells. Left: a representative experiment out of 3 in total. Right: n=5 mice per group. **B.** pSTAT5 expression in OT-I T cells. Left: a representative experiment out of 3 in total. Right: Geometric mean fluorescence intensity (MFI) of anti-pSTAT5-Alexa Fluor 647 in OT-I T cells. Mean of MFI values for each group per experiment are shown as a grey dots. Lines connect data from corresponding experiments. DEREG$^-$ n=7, DEREG$^+$ n=6.

C. In vitro proliferation assay. CTV-labelled CD8$^+$ T lymphocytes were stimulated with plate-bounded anti-CD3ε-antibody in the presence or absence of Tregs and/or recombinant IL-2. After 72 h, cells were analyzed by flow cytometry. Left: Proliferation is indicated by the CTV dilution. Right: IL-2Rα expression on CD8$^+$ T cells. A representative experiment out of 4 in total.

D-F. Treg-depleted DEREG$^+$ RIP.OVA mice and control DEREG$^-$ RIP.OVA mice received OT-I T cells, followed by the immunization with OVA peptide and LPS. On days 0, 1, and 2 post-immunization, RIP.OVA mice received IL-2ic (IL-2/S4B6 or IL-2/JES6). **D.** Experimental scheme. **E-F.** Urine glucose level was monitored on a daily basis for 14 days. Number of mice is indicated. Four mice from IL-2/JES6 group died before day 5 and were excluded from the analysis **E.** Survival curve is shown. **F.** Blood glucose concentration on day 7 post-immunization.

G-H. Diabetes was induced in Treg-depleted DEREG$^+$ RIP.OVA mice (with DT) or Treg-replete DEREG$^+$ RIP.OVA mice (without DT), which later received IL-2ic (IL-2/S4B6 or IL-2/JES6), or were left untreated (control) (scheme D). On day 5 post-immunization, spleens were collected and analyzed by flow cytometry. Four mice from IL-2/JES6 group died before the analysis (shown as "x"). Control n=6, IL-2/S4B6 n=6, IL-2/JES6 n=2, DEREG$^+$ DT n=4. **G.** Number of KLRG1$^+$ OT-I T cells. **H.** Number of Tregs (defined as GFP$^+$ CD4$^+$ T cells).

Statistical significance was calculated by two-tailed Mann-Whitney test (comparison of two groups), Kruskal-Wallis test (comparison of four groups), or Log-rank (Mantel-Cox) test (survival), p-value is shown in italics. Median is shown.

Figure 5

A-E. Ly5.1 OT-I T cells were adoptively transferred into Treg-depleted DEREG$^+$ RIP.OVA and control DEREG$^-$ RIP.OVA mice (n=3 mice per group). The next day, mice were immunized with DC-OVA. On day 3 post-immunization, spleens were isolated and OT-I T cells were FACS sorted as Ly5.1$^+$ CD8$^+$ cells, and analyzed via scRNAseq.

A. Heat map showing the relative expression of canonical IL-2 responsive genes**.** Each column represents one mouse.

B-D. UMAP projection of the individual OT-I T cell based on their gene expression profile. **B.** The colors indicate individual clusters revealed by unsupervised clustering. **C.** The intensity of the blue color indicates the level of *Klrk1* expression in individual cells. **D.** The origin of the cells (DEREG$^+$ or DEREG$^-$ mice) is indicated.

E. The percentage of cells assigned to specific clusters is shown for individual mice. Statistical significance was calculated by unpaired Student's t test with Bonferroni correction, p-value is shown in italics. Median is shown.

F-H. Cluster 4 was reanalyzed separately. **F.** UMAP projection showing two subclusters identified by unsupervised clustering. **G.** The intensity of the blue color indicates the level of expression of *Itgα4* or *Il7r* in individual cells. **H.** Projection of subclusters 0 and 1 on the original UMAP plot.

I. Ly5.1 OT-I T cells (5×10^4) were adoptively transferred into Treg-depleted DEREG$^+$ RIP.OVA mice (n=7) or DEREG$^-$ RIP.OVA mice (n=11). The next day, mice were immunized with DC-OVA. On day 3 post-immunization, spleens were collected and analyzed by flow cytometry. Left: A representative experiment out of three in total is shown. KLRK1$^+$ subset of Ly5.1 OT-I T cells was divided into two gates based on their expression of CD49d and IL-7R. Percentage of KILR effector T cells, defined as KLRK1$^+$ IL-7R$^+$ CD49d$^-$ cells out of OT-I T cells is shown. Statistical significance was calculated by two-tailed Mann-Whitney test for comparison, p-value shown in italics. Median is shown.

Figure 6

A. DEREG$^+$ RIP.OVA mice were treated with DT in order to deplete Tregs on days 0 and 1. On day 3, spleens were collected and analyzed by flow cytometry. Percentage of KILR effector T cells defined as KLRK1$^+$ CD49d$^-$ IL-7R$^+$ among CD8$^+$ T cells is shown. Untreated n=6, DT n=7.

B-E. OT-I *Rag2$^{-/-}$* mice were treated with OVA peptide (single dose on day 0, n=7), and/or IL-2/S4B6 (3 doses, days 0, 1 and 2, n=7 mice per group) or left untreated (n=6). Spleens were collected and analyzed by

flow cytometry on day 3. **B.** Four populations of OT-I T cells were identified based on their expression of KLRK1 and CD49d. A representative experiment out of three in total. **C.** Left: IL-7R expression on KLRK1$^+$ CD49d$^-$ OT-I T cells from mice treated with OVA, and/or IL-2/S4B6. Expression of IL-7R on naïve OT-I T cells is shown as a positive control. Representative staining. Right: Number of KILR effector T cells, defined as KLRK1$^+$ CD49d$^-$ IL-7R$^+$ OT-I T cells. **D.** IL-7R expression in CD49d$^+$ KLRK1$^-$, CD49d$^+$ KLRK1$^+$, CD49d$^-$ KLRK1$^+$, and CD49d$^-$ KLRK1$^-$ OT-I T cells from mice treated with OVA + IL-2/S4B6. Left: A representative histogram. Right: Percentage of IL-7R$^+$ cells among OT-I T cells in indicated populations. **E.** GZMB levels in OT-I T cells. Left: A representative histogram. Right: Geometric mean fluorescence intensity (MFI) of anti-GZMB-eFluor 660 antibody on OT-I T cells. Obtained values were normalized to the average of MFI of untreated samples in each experiment (=1).

F. OT-I *Rag2$^{-/-}$* mice were treated with OVA peptide (day 0), and/or IL-2/S4B6 (days 0, 1 and 2) or left untreated. Spleens were collected on day 3. KLRK1$^+$ CD8$^+$ or KLRK1$^-$ CD8$^+$ cells were sorted and adoptively transferred into recipient RIP.OVA mice, which have received a mixture of target OVA-pulsed CTV-loaded and unpulsed CFSE-loaded splenocytes from Ly5.1 mice at ~1:1 ratio earlier the same day. The next day, the spleens were analyzed for the presence of Ly5.1 donor cells by flow cytometry. Ratio of unpulsed (CFSE$^+$) to OVA pulsed (CTV$^+$) target cells was determined and normalized to control recipients which did not receive OT-I T cells (=1). KLRK1$^-$ (no treatment) n=6, KLRK1$^-$ (OVA + IL-2/JES6) n=11, KLRK1$^+$(OVA + IL-2/JES6) n=13, KLRK1$^+$(OVA peptide) n=6. Four independent experiments.

G-J. Human CD8$^+$ T-cell atlas was generated by integrating 14 scRNAseq data sets from blood of healthy donors. The gene expression data after the removal of MAIT cells were projected into a 2D UMAP plot. **G.** The assignment of individual cells to clusters identified by unsupervised clustering. Individual clusters were matched to established CD8$^+$ T cell subsets based on the expression of their signature markers (see Fig. S6H). **H.** The intensity of the blue color indicates the level of expression of indicated genes in individual cells. **I.** The percentage of CD8$^+$ T cells assigned to the KILR-like CD8$^+$ T-cell cluster in individual donors (n=14). **J.** Left: Clonally expanded T cells were identified based on their TCRαβ VDJ sequences. The intensity of the red color indicates the size of individual clones. Right: T cells with recovered complete TCRαβ VDJ information are shown in green.

Statistical significance was calculated by two-tailed Mann-Whitney test for comparison of two groups, or Kruskal-Wallis test for multiple groups comparison (p-value is shown in italics) with Dunn's post-test (* <0.05, **<0.01, ***<0.001). Median is shown.

Figure 7

A-C. On day 0, BALB/C mice were inoculated with BCL1 leukemia cells. On days 11 and 24 post inoculation, mice received doxorubicin (Dox), with or without anti-CD4 or anti-CD8 depletion mAbs. On three consecutive days, mice were treated with IL-2/JES6. n=16 mice per group in two independent experiments with the exception of "Dox + IL-2/JES6 + αCD8" conditions where three mice in the second experiment did not tolerate anti-CD8 administration for unknown reasons and were removed. **A.** Scheme of the experiment. **B.** Survival curves. **C.** The detection of the BCL1 IgM/IgD in the serum on day 54 by ELISA using an anti-idiotypic antibody. Percentage of mice with undetectable BCL1, detectable BCL1, and dead mice in each group is shown.

D-E. On day 0, C57Bl/6J mice were inoculated with B16F10 melanoma cells. On day 6, mice received doxorubicin (Dox). On three consecutive days, mice were treated with IL-2/JES6. n=16 mice per group in two independent experiments. **D**. Scheme of the experiment showing also the eventual treatment with anti-CD4 or anti-CD8 depletion mAb relevant for Figure EV7B. **E**. Survival curves.

Statistical significance was calculated by Log-rank (Mantel-Cox) test (survival) or Fisher's exact test (ELISA, two groups of the outcome: mice with undetectable BCL1 vs. mice with detectable BCL1 or dead). P-value is shown in italics. Median is shown.

Supplementary Figure Legends

Figure S1

A. Gating strategy for the experiment shown in Fig. 1D. CD4$^+$ T cells and B cells were depleted via magnetic bead separation prior to the analysis.

B. Representative dot plots for the experiment shown in Fig. 1D.

C. Blood glucose concentration (day 7) in the experiment shown in Fig. 1E (10^6 transferred OT-I T cells). RIP.OVA n=6, DEREG RIP.OVA n=8, *Foxp3^{DTR}* RIP.OVA n=8. One *Foxp3^{DTR}* RIP.OVA mouse died before the measurement (shown as "x"). Median is shown.

D. Representative dot plots for the experiment shown in Fig. 1F.

Figure S2

A. Representative dot plots for the experiment shown in Fig. 2D.

B. Representative dot plots for the experiment shown in Fig. 2E. **C.** Representative dot plots for the experiment shown in Fig. 2F.

D-E. Treg-depleted DEREG$^+$ RIP.OVA mice and control DEREG$^-$ RIP.OVA mice received 10^6 OT-I T cells. The next day, they were immunized with unloaded DC. **D.** Urine glucose level was monitored on a daily basis. Left: Survival curve. Number of mice per group is indicated. Right: Blood glucose concentration on day 7 post- immunization. **E.** On day 6 post-immunization, spleens were collected and analyzed by flow cytometry. Percentage of OT-I T cells among CD8$^+$ T cells, count of KLRG1$^+$ OT-I T cells, and count of IL-2Rα$^+$ OT-I T cells are shown. DEREG$^-$ n=6, DEREG$^+$ n=4.

F. Bubble chart illustrating diabetes incidence in DEREG$^+$ RIP.OVA and DEREG$^-$ RIP.OVA mice immunized with DC loaded with indicated peptides vs. numbers of transferred OT-I T cells. Number of diabetic mice and total number of mice per group is indicated.

Statistical significance was calculated by Log-rank (Mantel-Cox) test (survival) or two-tailed Mann-Whitney test (glucose concentration and flow cytometry analysis), p-value is shown in italics. Median is shown.

Figure S3

A. Percentage of diabetic mice in individual groups in the experiment shown in Fig. 3A. Number of diabetic mice and total number of mice per group is indicated.

B-C. In the experimental setup shown in Fig. 3A, spleens were collected on day 8 post-immunization and analyzed by flow cytometry. **B.** Left: Percentage of OT-I T cells among CD8$^+$ T cells. A representative experiment out of five in total. Right: Number of OT-I T cells is shown. NT n=25, Conv. CD4$^+$ n=15, Tregs n=12. **C.** KLRG1 expression on OT-I T cells on day 8, a representative histogram.

D-E. In the experimental setup shown in Fig. 3A, spleens were collected on day 14 post-immunization and analyzed by flow cytometry. **D.** Left: Percentage of OT-I T cells among CD8$^+$ T cells. Representative plots are shown. Right: Number of OT-I T cells is shown. NT n=15, Tregs n=7. **E.** KLRG1 expression on OT-I T cells on day 14. A representative experiment out of 3 in total.

Statistical significance was calculated using Chi-square test (p-value is shown in italics) and Fisher's exact post-test (***<0.001, ****<0.0001) for comparison of two groups of the outcomes: diabetic mice vs. diabetes-free mice (A). Statistical significance for flow cytometry analysis was calculated by Kruskal-Wallis test (p-value is shown in italics) with Dunn's post-test (* <0.05, ***<0.001, ****<0.0001) for comparison of three groups (B), or two-tailed Mann-Whitney test for comparison of two groups (D). Median is shown.

Figure S4

A. Scheme of the experiments described in Fig. 4A-C and Fig. S4B-E.

B. Gating strategy for the experiment shown in Fig. 4A.

C-E. The experiment described in Fig. 4A was modified by using congenic Ly5.1 OT-I T cells for the adoptive transfer. **C.** The gating strategy. **D.** IL-2Rα expression on OT-I Ly5.1 T cells. Left: a representative experiment out of 3 in total. Right: Percentage of IL-2Rα$^+$ cells among OT-I Ly5.1 T cells, n=7 mice per group. **E.** pSTAT5 expression in OT-I Ly5.1 T cells. Left: a representative experiment out of 3 in total. Right: Geometric mean fluorescence intensity (MFI) of anti-pSTAT5-Alexa Fluor 647 on Ly5.1 OT-I T cells. Mean of MFI values for each genotype per experiment are shown as a grey dots. Lines connect data from corresponding experiments. n=7 mice per group.

G. OT-I CD8$^+$ and OT-II CD4$^+$ T cells (both Ly5.2) were labeled with CFSE and adoptively transferred into Ly5.1 mice as a mixture (0.75 × 10^6 CD8$^+$ OT-I + 1.5 × 10^6 CD4$^+$ OT-II T cells), followed by the administration of OVA protein with or without IL-2ic (IL-2/S4B6 or IL-2/JES6). On day 5 post-immunization, spleens were collected and analyzed by flow cytometry. Top: Representative staining showing CD45.2$^+$ OT-I (CD4$^-$) and OT-II (CD4$^+$) T cells among viable cells. Bottom: Proliferation is indicated by the CFSE dilution. Representative histograms for OT-I and OT-II T cells.

Statistical significance was calculated by two-tailed Mann-Whitney test, p-value is shown in italics. Median is shown.

Figure S5

A-E. Additional analysis for the scRNAseq experiment shown in Fig. 5A-H. **A.** Gene set enrichment analyses comparing the ranked expression of IL-2 induced genes in DEREG$^+$ versus DEREG$^-$ samples. The list of IL-2-responsive genes is based on the indicated publications [40, 41]. P-value (shown in italics) is estimated based on an adaptive multi-level split Monte-Carlo scheme. **B.** Violin plot indicating the expression of *Gzmb* in OT-I T cells primed in DEREG$^-$ and DEREG$^+$ mice. **C.** The intensity of the blue color indicates the level of expression of indicated genes in individual cells. **D.** Projection of clusters of primed OT-I T cells (Fig. 5B) on the UMAP visualization of CD8$^+$ T cells subsets formed during an anti-viral response using TILPRED algorithm. **E.** Natural killer cell signature was calculated for each cluster. Top: The intensity of the blue color indicates the level of expression of natural killer signature genes. Bottom: violin plot showing natural killer cell signature score calculated for each cluster.

F. Additional results of the experiments described in Fig. 5I. CD103 expression on KLRK1$^+$ CD49d$^+$ IL-7R$^-$ (Effector) cells and KLRK1$^+$ CD49d$^-$ IL-7R$^+$ (KILR) T cells. Left: A representative histogram. Right: percentage of CD103$^+$ cells among cells falling into KILR gate. DEREG$^-$ n=11, DEREG$^+$ n=7. Median is shown. Statistical significance was calculated using two-tailed Mann-Whitney test, p-value is shown in italics.

G. Ly5.1 OT-I T cells (5 × 10^4) were transferred into Treg-depleted DEREG$^+$ RIP.OVA mice and DEREG$^-$ RIP.OVA mice. The next day, mice were immunized with DC-OVA. On day 3 post immunization, splenic CD49d$^+$ KLRK1$^-$ antigen-experienced (AE), CD49d$^+$ KLRK1$^+$ double-positive (DP), and CD49d$^-$ KLRK1$^+$ (KILR) OT-I T cells (identified as Ly5.1$^+$ CD8$^+$) were FACS-sorted as indicated and analyzed via RT-qPCR. The correspondence of the FACS gates to scRNAseq clusters is shown. The expression of indicated genes was normalized to *Gapdh*. Mean + SEM is indicated. Four independent experiments. Statistical significance was calculated by Kruskal-Wallis test, p-value is shown in italics.

Figure S6

A. Representative experiment out of three in total for the data shown in Fig. 6A.

B. GZMB levels in CD8$^+$ T cells from the experiment shown in Fig. 6A. Left: A representative histogram. Right: Geometric mean fluorescence intensity (MFI) of anti-GZMB-eFluor 660 antibody on CD8$^+$ T cells was determined. Obtained values were normalized to the average of MFI of untreated samples in each experiment (=1). DT treated n=6, Untreated n=5 mice.

C-E. Additional datafor the experiment shown in Fig. 6F. **C.** Scheme of the experiment. **D.** The percentage of OVA-pulsed target cells out of all viable splenocytes. No OT-I T cell transfer n=8, KLRK1$^-$ (no treatment) n=6, KLRK1$^-$ (OVA + IL-2/JES6) n=11, KLRK1$^+$ (OVA + IL-2/JES6) n=13, KLRK1$^+$ (OVA peptide) n=6. **E.** A representative experiment out of 4 in total.

F. Identification of MAIT cells in the Human CD8[+] T cell atlas. Top left: UMAP projection. Individual cells are assigned to individual clusters based on unsupervised clustering. Cluster 6 corresponds to MAIT cells. Top right: The intensity of the blue color indicates the level of *KLRB1* expression (a MAIT cell marker) in individual cells. Bottom: Cells expressing five most abundant semi-invariant MAIT cell-specific receptor-alpha (TCRα) are shown.

G. The intensity of the blue color indicates the expression level of selected genes based on the detection using hash-tagged antibodies (CD45RA and CD45RO) or transcripts in Human CD8[+] T cell atlas after the removal of MAIT cells.

Statistical significance was calculated by Kruskal-Wallis test (p-value is shown in italics) with Dunn's multiple comparison post-test (* <0.05, ***<0.001) (C, F), or two-tailed Mann-Whitney test (B). Median is shown.

H. Natural killer cell signature was calculated for each cluster of the Human CD8[+] T cell atlas. Left: The intensity of the blue color indicates the level of expression of natural killer signature genes. Right: violin plot showing natural killer cell signature score calculated for each cluster.

Figure S7

A. In experiments described in Fig. 7A-C, BCL1 IgM/IgD in the serum on days 28, 54, and 94-96 post tumor inoculation was detected by ELISA. n=16 mice per group in two independent experiments with two exceptions: (i) "Dox + IL-2/JES6 + αCD4" conditions on day 54 where the analysis failed for the second experiment and thus, n=8 mice and (ii) control conditions on day 54, when one mouse was not analyzed for technical reasons and thus, n=15. The data shown here include data shown in Fig. 7C. The statistical significance was calculated using Chi-square test (two groups of the outcomes: mice with undetectable BCL1 vs. mice with detectable BCL1 or dead).

B. On day 0, C57Bl/6J mice were inoculated with B16F10 melanoma cells. On day 6, mice received doxorubicin (Dox) with/without anti-CD4 or anti-CD8 depletion mAb. On three consecutive days, mice were treated with IL-2/JES6 (see Fig. 7D). Survival curves. n=8 mice per group in one experiments. The data shown here partially overlap with data shown in Fig. 7E. There were two experiments in total, but only the second experiment had the mouse groups with the depletion of CD8[+] or CD4[+] cells. Fig. 7E shows pooled data for all conditions included in both experiments. Here, results from the second experiment are shown. Statistical significance was calculated by Log-rank (Mantel-Cox) test.

Table S1

Differentially expressed genes for all clusters shown in Figure 5B. The used differential expression criteria were fold change above 2, minimum of 0.1 difference in the fraction of detection between each cluster and

the rest of the cells and adjusted p-value below 0.01. For each gene, the fractions of cells with at least one detected transcript in the tested cluster (pct.1) or among all other cells (pct.2) are listed. The columns 'p_val', 'avg_log2FC' and 'p_val_adj' show p-values (Mann Whitney U Test), log2 fold changes and adjusted p-values (Bonferroni correction).

Table S2

Differentially expressed genes between the super-effector-like cell cluster and all the remaining cells in the human CD8[+] atlas (after MAIT removal). The used differential expression criteria were average fold change above 1.5 and adjusted p-value below 0.01. For each gene, the fractions of super-effector-like cells (pct.1) or remaining cells (pct.2) with at least one detected transcript are listed. The columns 'p_val ', 'avg_log2FC' and 'p_val_adj' show p-values (Mann Whitney U Test), log2 fold changes and adjusted p-values (Bonferroni correction).

References

1. Josefowicz, S.Z. and A. Rudensky, *Control of Regulatory T Cell Lineage Commitment and Maintenance.* Immunity, 2009. **30**(5): p. 616-625.
2. Bennett, C.L., et al., *The immune dysregulation, polyendocrinopathy, enteropathy, X-linked syndrome (IPEX) is caused by mutations of FOXP3.* Nature Genetics, 2001. **27**(1): p. 20-21.
3. Brunkow, M.E., et al., *Disruption of a new forkhead/winged-helix protein, scurfin, results in the fatal lymphoproliferative disorder of the scurfy mouse.* Nature Genetics, 2001. **27**(1): p. 68-73.
4. Wildin, R.S., et al., *X-linked neonatal diabetes mellitus, enteropathy and endocrinopathy syndrome is the human equivalent of mouse scurfy.* Nature Genetics, 2001. **27**(1): p. 18-20.
5. Chinen, T., et al., *An essential role for the IL-2 receptor in T-reg cell function.* Nature Immunology, 2016. **17**(11): p. 1322-1333.
6. Laidlaw, B.J., et al., *Production of IL-10 by CD4(+) regulatory T cells during the resolution of infection promotes the maturation of memory CD8(+) T cells.* Nature Immunology, 2015. **16**(8): p. 871-+.
7. Kalia, V., et al., *Quiescence of Memory CD8(+) T Cells Is Mediated by Regulatory T Cells through Inhibitory Receptor CTLA-4.* Immunity, 2015. **42**(6): p. 1116-1129.
8. McNally, A., et al., *CD4+CD25+ regulatory T cells control CD8+ T-cell effector differentiation by modulating IL-2 homeostasis.* Proc Natl Acad Sci U S A, 2011. **108**(18): p. 7529-34.
9. Kastenmuller, W., et al., *Regulatory T cells selectively control CD8+ T cell effector pool size via IL-2 restriction.* J Immunol, 2011. **187**(6): p. 3186-97.
10. Pace, L., et al., *Regulatory T Cells Increase the Avidity of Primary CD8(+) T Cell Responses and Promote Memory.* Science, 2012. **338**(6106): p. 532-536.
11. Schildknecht, A., et al., *FoxP3+ regulatory T cells essentially contribute to peripheral CD8+ T-cell tolerance induced by steady-state dendritic cells.* Proc Natl Acad Sci U S A, 2010. **107**(1): p. 199-203.
12. Green, E.A., et al., *CD4(+)CD25(+) T regulatory cells control anti-islet CD8(+) T cells through TGF-beta-TGF-beta receptor interactions in type 1 diabetes.* Proceedings of the National Academy of Sciences of the United States of America, 2003. **100**(19): p. 10878-10883.
13. King, C.G., et al., *T cell affinity regulates asymmetric division, effector cell differentiation, and tissue pathology.* Immunity, 2012. **37**(4): p. 709-20.
14. Kurts, C., et al., *Major histocompatibility complex class I-restricted cross-presentation is biased towards high dose antigens and those released during cellular destruction.* Journal of Experimental Medicine, 1998. **188**(2): p. 409-414.
15. Lahl, K., et al., *Selective depletion of Foxp3(+) regulatory T cells induces a scurfy-like disease.* Journal of Experimental Medicine, 2007. **204**(1): p. 57-63.
16. Kim, J.M., J.P. Rasmussen, and A.Y. Rudensky, *Regulatory T cells prevent catastrophic autoimmunity throughout the lifespan of mice.* Nature Immunology, 2007. **8**(2): p. 191-197.
17. Stepanek, O., et al., *Coreceptor scanning by the T cell receptor provides a mechanism for T cell tolerance.* Cell, 2014. **159**(2): p. 333-45.
18. Polhill, T., et al., *IL-2/IL-2Ab complexes induce regulatory T cell expansion and protect against proteinuric CKD.* J Am Soc Nephrol, 2012. **23**(8): p. 1303-8.
19. Liu, R., et al., *Expansion of regulatory T cells via IL-2/anti-IL-2 mAb complexes suppresses experimental myasthenia.* Eur J Immunol, 2010. **40**(6): p. 1577-89.
20. Kim, H.P., J. Kelly, and W.J. Leonard, *The basis for IL-2-induced IL-2 receptor alpha chain gene regulation: importance of two widely separated IL-2 response elements.* Immunity, 2001. **15**(1): p. 159-72.

21. Boyman, O., et al., *Selective stimulation of T cell subsets with antibody-cytokine immune complexes.* Science, 2006. **311**(5769): p. 1924-1927.

22. Spangler, J.B., et al., *Antibodies to Interleukin-2 Elicit Selective T Cell Subset Potentiation through Distinct Conformational Mechanisms.* Immunity, 2015. **42**(5): p. 815-825.

23. Kim, R. and T. Kin, *Current and Future Therapies for Immunogenic Cell Death and Related Molecules to Potentially Cure Primary Breast Cancer.* Cancers (Basel), 2021. **13**(19).

24. Bosch, A.J.T., et al., *A minimum number of autoimmune T cells to induce autoimmunity?* Cell Immunol, 2017. **316**: p. 21-31.

25. Tomala, J. and M. Kovar, *IL-2/anti-IL-2 mAb immunocomplexes: A renascence of IL-2 in cancer immunotherapy?* Oncoimmunology, 2016. **5**(3): p. e1102829.

26. Marchingo, J.M., et al., *Antigen affinity, costimulation, and cytokine inputs sum linearly to amplify T cell expansion.* Science, 2014. **346**(6213): p. 1123-1127.

27. Dong, S., et al., *The effect of low-dose IL-2 and Treg adoptive cell therapy in patients with type 1 diabetes.* JCI Insight, 2021. **6**(18).

28. Akkaya, B., et al., *Regulatory T cells perform antigen specific suppression by depleting cognate peptide-MHC class II via trogocytosis.* Journal of Immunology, 2019. **202**(1).

29. Wong, H.S., et al., *A local regulatory T cell feedback circuit maintains immune homeostasis by pruning self-activated T cells.* Cell, 2021. **184**(15): p. 3981-3997 e22.

30. Prajapati, K., et al., *Functions of NKG2D in CD8(+) T cells: an opportunity for immunotherapy.* Cell Mol Immunol, 2018. **15**(5): p. 470-479.

31. Schorer, M., et al., *Rapid expansion of Treg cells protects from collateral colitis following a viral trigger.* Nat Commun, 2020. **11**(1): p. 1522.

32. Joshi, N.S., et al., *Inflammation directs memory precursor and short-lived effector CD8(+) T cell fates via the graded expression of T-bet transcription factor.* Immunity, 2007. **27**(2): p. 281-95.

33. Kaech, S.M., et al., *Selective expression of the interleukin 7 receptor identifies effector CD8 T cells that give rise to long-lived memory cells.* Nature Immunology, 2003. **4**(12): p. 1191-1198.

34. Lee, S.J., et al., *CD134 Costimulation Couples the CD137 Pathway to Induce Production of Supereffector CD8 T Cells That Become IL-7 Dependent.* J Immunol, 2007. **179**(4): p. 2203-14.

35. Barsoumian, H.B., E.S. Yolcu, and H. Shirwan, *4-1BB Signaling in Conventional T Cells Drives IL-2 Production That Overcomes CD4+CD25+FoxP3+ T Regulatory Cell Suppression.* PLoS One, 2016. **11**(4): p. e0153088.

36. Oh, H.S., et al., *4-1BB Signaling Enhances Primary and Secondary Population Expansion of CD8+ T Cells by Maximizing Autocrine IL-2/IL-2 Receptor Signaling.* PLoS One, 2015. **10**(5): p. e0126765.

37. Palmer, E., A. Drobek, and O. Stepanek, *Opposing effects of actin signaling and LFA-1 on establishing the affinity threshold for inducing effector T-cell responses in mice.* Eur J Immunol, 2016. **46**(8): p. 1887-901.

38. Barnden, M.J., et al., *Defective TCR expression in transgenic mice constructed using cDNA-based alpha- and beta-chain genes under the control of heterologous regulatory elements.* Immunol Cell Biol, 1998. **76**(1): p. 34-40.

39. Jang, Y., et al., *Cutting Edge: Check Your Mice-A Point Mutation in the Ncr1 Locus Identified in CD45.1 Congenic Mice with Consequences in Mouse Susceptibility to Infection.* J Immunol, 2018. **200**(6): p. 1982-1987.

40. Kralova, J., et al., *Expression of Fluorescent Fusion Proteins in Murine Bone Marrow-derived Dendritic Cells and Macrophages.* J Vis Exp, 2018(140).

41. Kim, M.V., et al., *Murine in vivo CD8(+) T Cell Killing Assay.* Bio Protoc, 2014. **4**(13).

42. Slavin, S. and S. Strober, *Spontaneous Murine B-Cell Leukemia.* Nature, 1978. **272**(5654): p. 624-626.

43. Howe, K.L., et al., *Ensembl 2021.* Nucleic Acids Res, 2021. **49**(D1): p. D884-D891.

44. Zheng, G.X., et al., *Massively parallel digital transcriptional profiling of single cells.* Nat Commun, 2017. **8**: p. 14049.

45. Haberman, Y., et al., *Pediatric Crohn disease patients exhibit specific ileal transcriptome and microbiome signature.* J Clin Invest, 2014. **124**(8): p. 3617-33.

46. Andreatta, M., et al., *Interpretation of T cell states from single-cell transcriptomics data using reference atlases.* Nat Commun, 2021. **12**(1): p. 2965.

47. Kovanen, P.E., et al., *Global analysis of IL-2 target genes: identification of chromosomal clusters of expressed genes.* Int Immunol, 2005. **17**(8): p. 1009-21.

48. Lin, J.X., et al., *Critical Role of STAT5 transcription factor tetramerization for cytokine responses and normal immune function.* Immunity, 2012. **36**(4): p. 586-99.

49. Korotkevich, G., et al., *Fast gene set enrichment analysis.* bioRxiv, 2021: p. 060012.

50. Bais, A.S. and D. Kostka, *scds: computational annotation of doublets in single-cell RNA sequencing data.* Bioinformatics, 2020. **36**(4): p. 1150-1158.

51. Hahsler, M., M. Piekenbrock, and D. Doran, *dbscan: Fast Density-Based Clustering with R.* Journal of Statistical Software, 2019. **91**(1): p. 1-30.

A

Foxp3^{DTR} or DEREG RIP.OVA

d -2 → DT

d -1 → OT-I T cells / DT

d 0 → LPS + OVA

d 5 - d 14 → Glucose monitoring

B Diabetes incidence

% diabetic

11/12 · 0/14 · 6/6 · 10/11 · 2/4

OT-I #: 1M · 10k · 10k · 10k · 3k

■ RIP.OVA
■ DEREG RIP.OVA
■ *Foxp3*^{DTR}RIP.OVA

C Day 7 (10k OT-I T cells)

Blood glucose [mmol/l]

<0.0001

RIP-OVA · DEREG RIP.OVA · *Foxp3*^{DTR} RIP.OVA

D OT-I T cells

% out of CD8$^+$ T cells

0.012

RIP-OVA · *Foxp3*^{DTR} RIP.OVA

E Priming with LPS only

% diabetic

0/9 · 8/16 · 4/8 · 4/13 · 2/12

OT-I #: 1M · 1M · 1M · 100k · 10k

■ RIP.OVA
■ DEREG RIP.OVA
■ *Foxp3*^{DTR} RIP.OVA

F OT-I T cells

% out of CD8$^+$ T cells

0.0026 **

RIP-OVA · *Foxp3*^{DTR} RIP.OVA · *Foxp3*^{DTR}

Effector cells (CD44$^+$ CD62L$^-$)

% out of OT-I T cells

0.0045 * **

RIP-OVA · *Foxp3*^{DTR} RIP.OVA · *Foxp3*^{DTR}

A

$Cd3\varepsilon^{-/-}$ RIP.OVA

CD8⁺ T cells

CD4⁺ T cells (FOXP3⁺ or FOXP3⁻)
OT-I T cells

DC-OVA

Glucose monitoring

d -8   d -1   d 0   d 5 - d 21

B Diabetes induction in $Cd3\varepsilon^{-/-}$ RIP.OVA

<0.0001

% diabetes free

Days post immunization

— NT, n=28
— Conv. CD4⁺ T cells, n=28
— Tregs, n=14

C Day 8   Day 14

Blood glucose [mmol/l]

*   <0.0001   **

NT   Conv. CD4⁺   Tregs

0.0029

NT   Tregs

D OT-I T cells (day 8)

0.0099

% out of CD8⁺ T cells

*   **

NT   Conv. CD4⁺   Tregs

E KLRG1⁺ OT-I cells (day 8)

0.0004

Cell number × 10⁴

*   ***

NT   Conv. CD4⁺   Tregs

F OT-I T cells (day 14)

0.0019

% out of CD8⁺ T cells

NT   Tregs

G KLRG1⁺ OT-I cells (day 14)

0.0024

Cell number

NT   Tregs

A IL-2Rα⁺ OT-I T cells

B pSTAT5 (OT-I T cells)

C Anti-CD3ε antibody | Anti-CD3ε + IL-2 | Anti-CD3ε antibody | Anti-CD3ε + IL-2

D DEREG RIP.OVA

E Diabetes induction with IL-2ic

F Day 7

G KLRG1⁺ OT-I T cells

H Tregs (GFP⁺ CD4⁺ T cells)

A — BALB/C: BCL1 (d 0); Dox, αCD4 or αCD8 (d 11); IL-2/JES6 (d 12, 13, 14); Dox, αCD4 or αCD8 (d 24); IL-2/JES6 (d 25, 26, 27); Survival monitoring (d 120)

B — BCL1 leukemia; *<0.0001*; Percent survival vs Time (days). Control, Dox, IL-2/JES6, Dox + IL2/JES6, Dox + IL2/JES6 + αCD8, Dox + IL2/JES6 + αCD4

C — BCL1-derived IgM (day 54); *0.011*; % of mice in group; Dox, Dox + IL-2/JES6; BCL1 undetected, BCL1 positive, Dead

D — C57Bl/6J: B16F10 (d 0); Dox, αCD4 or αCD8 (d 6); IL-2/JES6 (d 7, 8, 9); Survival monitoring (d 70)

E — B16F10 melanoma; *0.0048*; Percent survival vs Time (days). Control, Dox, IL-2/JES6, Dox + IL2/JES6

FIGURE S1

A *Cd3ε⁻ʹ⁻*RIP.OVA

B

OT-I T cells (day 8)

C OT-I T cells

Conv. CD4⁺ T cells
No CD4⁺ T cell transfer
Tregs

D No CD4⁺ T cell transfer / Tregs

OT-I T cells (day 14)

E OT-I T cells

No CD4⁺ T cell transfer
Tregs

FIGURE S5

FIGURE S7