What happened when? Brain and behavioral responses to violated expectations about the structure and content of episodic memories

Sophie Siestrup[1,2*], Benjamin Jainta[1], Nadiya El-Sourani[1], Ima Trempler[1,2], Oliver T. Wolf[c], Sen Cheng[d], Ricarda I. Schubotz[1,2]

[1] Department of Psychology, University of Münster, Münster, Germany,

[2] Otto Creutzfeldt Center for Cognitive and Behavioral Neuroscience, University of Münster, Münster, Germany

[3] Department of Cognitive Psychology, Faculty of Psychology, Ruhr University Bochum, Bochum, Germany

[4] Institute for Neural Computation, Ruhr University Bochum, Bochum, Germany

*Corresponding author

E-mail: s.siestrup@uni-muenster.de (SS)

Abstract

Episodic memories are not static but can be modified on the basis of new experiences, potentially

allowing us to make valid predictions in the face of an ever-changing environment. Recent research

has identified mnemonic prediction errors as a possible trigger for such modifications. In the present

study, we investigated the influence of different types of mnemonic prediction errors on brain

activity and subsequent memory performance using a novel paradigm for episodic modification.

Participants encoded different episodes which consisted of short toy stories. During a subsequent

functional magnetic resonance imaging (fMRI) session, episodic retrieval was cued by presenting

videos showing the original episodes, or modified versions thereof. In modified videos either the

order of two subsequent action steps was changed (violating structure expectancy) or an object was

exchanged for another (violating content expectancy). While brain responses to structure expectancy

violations were only subtle, content expectancy violations recruited brain areas relevant for

processing of new object information. In a post-fMRI memory test, the participants' tendency to

accept modified episodes as originally encoded increased significantly when they had experienced

expectancy violations during the fMRI session. Our study provides valuable initial insights into the

neural processing of different types of mnemonic prediction errors and their influence on

subsequent memory.

Introduction

Episodic memories are our window to the past. They enable us to vividly re-live events that we experienced at some point in our personal life, allowing us to re-locate them in space and time [1]. But there is evidence that episodic memories are not always an exact, veridical reconstruction of our past. On the contrary, memories are flexible and prone to change [2, 3, 4, 5]. It is often believed that the modification of episodic memories is undesirable and occurs due to erroneous processes during memory encoding, consolidation or retrieval [6, 7]. However, a valuable new perspective is that such modifications serve an important function for our predictive capabilities [8, 9, 10]. Prediction is thought to be based on internal models [11] derived from memories. What is crucial for predictive success is not only that we establish and recall appropriate internal models of the world, but also attune them to persistent change to maintain valid predictions in the long run. This is accomplished via prediction errors that serve as bottom-up learning signals for model updating [12, 13, 14]. Accordingly, it is likely adaptive that largely faithful memories are modified in favor of valid internal models informed and updated by later experiences [10, 15]. Such prediction errors could arise in everyday life if the episodic memory recalled in a situation only partially resembles this new situation, which is probably the normal case. Accordingly, these differences could become effective as prediction errors in the process of memory reconstruction. Although some have suggested that prediction errors can also lead to an improvement of memories under certain circumstances [16, 17, 18], potentially as they can trigger one-shot learning [17], evidence is also accumulating that violated expectations are an important driver of memory change, consistent with the theoretical framework mentioned above [19]. In addition to behavioral studies, researchers have begun to investigate how the human brain processes mnemonic prediction errors. For example, some have found evidence that the hippocampus responds to unexpected outcomes that contradict previously established memories, and thus may play an important role in memory updating based on new relevant information [20, 21].

Memory traces include different types of information about an experienced episode, i.e., "what"

happened "when" and "where" [21]. To our knowledge, previous studies have not yet specifically

targeted different types of mnemonic prediction errors based on separate components of the

memory trace. Moreover, reported memory tasks are often not very naturalistic, employing simple

stimuli (e.g. word or picture lists) and encoding periods with little participant involvement (e.g.

learning a list or watching a video; e.g. [22, 23]). In the present study, we therefore aimed to extend

the understanding of mnemonic prediction errors at the neural and behavioral level by selectively

targeting predictions about two different aspects of episodic memories, either their content ("what")

or their structure ("when"). To do so, we employed a new naturalistic paradigm for encoding

episodic memories in the laboratory, then violated expectations based on reactivated episodes in

two different ways and measured neural responses using functional magnetic resonance imaging

(fMRI). Episodes were reactivated by videos reminiscent of episodes that each participant had

encoded in a training session one week before entering the fMRI session. A subset of these videos

was selectively manipulated with regard to either the occurrence of an object (content modification)

or the order of two consecutive action steps (structure modification). Importantly, these

modifications were rather subliminal as they neither challenged general world knowledge nor the

overall outcome of the episodes. Following the paradigm implemented by Schiffer et al. [24, 25],

modified episodes were repeatedly presented to simulate the natural circumstances which

potentially foster memory modification, i.e. updating of internal models due to increasing evidence

for the validity of an alternative. In the present study, episodes were comprised of short stories

played with *PLAYMOBIL*® toys. We chose this approach to allow for complex but controlled,

laboratory-encoded episodes which are largely unbiased by previous everyday experiences.

In a post-fMRI memory test, participants were presented videos with episodes reminiscent of the

original training session or modified versions thereof. Participants were asked to indicate on a four-

point Likert scale whether the presented video shows an episode from the training session or not.

We expected that mnemonic prediction errors experienced during scanning should trigger internal

model updating [24, 25] and thus lead to a decrease in memory performance afterwards (H1), which

we examined using the discrimination index P_r. More specifically, we might find that repeated

expectancy violation during the fMRI session leads to an increased acceptance of modified videos as

original episodes (H1a), similar to previous behavioral findings reporting misattribution of new items

to old memories [20, 22]. Moreover, we might even see a reduced acceptance of unmodified videos

as a result of updating (H1b) [26, 27]. We tested these assumptions by analyzing rating scores from

the memory test. Since we expected to find differential brain activation between the two types of

episodic modifications (as described below), we might also see that the type of modification further

modulates memory performance which we also wanted to investigate in a more exploratory manner.

Concerning brain activation, we expected that some regions would react to both types of expectancy

violations (H2). Common activation was investigated by first calculating the separate contrasts of the

two types of expectancy violation against original videos, i.e. the no-expectancy violation condition

(structure expectancy violation vs. original, content expectancy violation vs. original), and combining

them through conjunction analysis. Several brain networks have been described to underlie the

retrieval of declarative memories regardless of their specific content [28]. Particularly, the

frontomedian cortex (mPFC) is a candidate area that may serve more general control over

consolidation and retrieval of long-term memories [29, 30]. Furthermore, the hippocampus is

regarded a core structure of episodic memory [31, 32, 33] and memory modification is believed to be

accomplished by the hippocampus linking reactivated content to new experience during reactivation,

and delivering updated memories to the frontomedian cortex [34]. Additionally, as outlined above,

the hippocampus has already been identified as a region that responds to mnemonic prediction

errors [20, 21].

The episodes employed in the present study could be classified in two different ways, as stories

showing events experienced by toy figure actors/actresses, but also as playing actions performed by

an actor/actress. As to the latter interpretation, a hypothesis was that brain areas related to the

observation of actions may also be engaged for videos showing modified episodes, including lateral

premotor cortex, the intraparietal sulcus (IPS), the ventrolateral prefrontal cortex, and the lateral

occipitotemporal cortex (OTC), together coined 'AON' for "action-observation network" [35].

Accordingly, we found previously that activity in all of these four areas attenuated when subjects

adapted to an expectancy violation affecting both objects and sequential structure *combined* [24,

25]. Taken together, the AON, the mPFC and the hippocampal formation were expected to respond

to both types of expectancy violations.

Additionally, we expected to also find specific activations for structure and content episodic

expectancy violations, which we investigated by calculating the above-mentioned separate contrasts.

First, we hypothesized that structure-expectancy violation leads to elevated activity in the lateral

premotor cortex [36] and mesial premotor cortex (called supplementary motor area (SMA) [37]) due

to their role in sequential association of single manipulations. We further expected to find elevated

brain activity in the area of dorsal premotor cortex and adjacent prefrontal sites along the superior

frontal sulcus (PMd/SFS) for structure expectancy violation. This region is characteristically active in

stepwise ordinal linking of individual action steps, as required in quite different, but especially

predictive, tasks [38, 39, 40, 41, 42](H3).

Content expectancy violation was expected to lead to a particularly strong response in the lateral

OTC [43], in the anterior IPS [44, 45], and, in addition, in the fusiform gyrus (FG [46]), due to their

complex object-action-focused functional profile (H4).

Apart from the expected common activation in the ventrolateral prefrontal cortex, we might find

different response patterns in this area for the two types of episodic expectancy violation. For

example, it has been suggested that within the inferior frontal gyrus (IFG), Brodmann area 44 (BA 44)

may be especially involved in processing the sequential structure of actions [47, 48] whereas BA 45

may be more relevant for semantic content [49, 50].

Materials and Methods

Participants

Forty female participants took part in the study. Only females participated to guarantee a high level of self-identification with our stimulus material which showed female hands. Two participants did not return for the fMRI session after training. Additionally, data from one participant had to be excluded from analysis due to nausea and dizziness during the fMRI session. In another case, an unknown technical problem led to the incomplete acquisition of functional data. Thus, the final sample for fMRI analyses consisted of 36 participants, age 18 – 28 years (M = 22.67 years, SD = 2.40 years). Participants of the final sample were all right-handed as assessed by the Edinburgh Handedness Inventory [51]. Handedness scores varied from +60 to +100 (M = 92.17, SD = 10.95). Data from one participant was not used for the analysis of the post-fMRI memory test due to misunderstood instructions. All participants had normal or corrected-to-normal vision, were native German speakers, and reported no history of neurological or psychiatric disorders or substance abuse. Participants signed a written informed consent and received monetary compensation or course credits for their participation. The study was conducted in accordance with the Declaration of Helsinki and approved by the Local Ethics Committee of the University of Münster.

Stimuli

Even though episodic memory can potentially never be completely independent of semantic knowledge [52] we wanted to reduce influences of everyday experiences as far as possible in our study. To this end we chose to use abstract but complex toy-based stories so that we could ensure controlled encoding of unique episodes in the laboratory. At the end of the study, participants were asked whether the stories used in the study reminded them of experiences they made in everyday life and 61.1 % answered no.

Stimuli were 78 short videos (duration = 8.80 – 17.88 s, M = 12.71 s) of stories that were played with PLAYMOBIL® toys, showing only the toys and hands and underarms of an actress. Stories comprised

six to nine action steps (M = 7.4 steps) and four to 14 separable objects (M = 6.93 objects), such as

characters, animals, vehicles and tools. The exact same object appeared in only one of the stories.

Most participants were familiar with *PLAYMOBIL®* toys, as 78 % of them indicated in the first session

that they had played with these toys during their childhood. Those that had not played with

PLAYMOBIL® toys had all played with similar toys.

Stories were filmed from above with a digital single-lens reflex camera (Nikon D5300) which was

centrally mounted above the table and faced straight down. Matt white paper served as a base. A

frame of 47.5 cm x 28 cm was taped on the paper, congruent with the section captured by the

camera (in the following referred to as camera frame). Objects that were needed for a particular

story were positioned next to the camera frame and were only moved into view in the moment at

which they appeared in the story. During filming, the actress wore a black pullover and black rubber

gloves to mask individual physical characteristics. To facilitate future imitation from demo videos, the

back of the right hand was marked with a yellow dot [53]. For a schematic overview of the filming

setup, see Fig 1A. Video material was edited using Adobe Premiere Pro CC (Adobe Systems Software,

Dublin, Ireland, Version 12.1.2). All videos had a frame of size 1920 x 1080 pixels and a frame rate of

25 frames per second. Videos started with seven frames showing only background and ended after

seven frames showing the final toy constellation. The original perspective of the videos was the third-

person (observer) perspective (3pp) and they were rotated 180° to create their first-person (field)

perspective (1pp) counterparts using Adobe Premiere Pro CC [54] (Fig 1B). Throughout the

experiment, videos were presented at a visual angle of approximately 7.3° x 13° using the stimulus

presentation software Presentation (version 20.3 02.01.19, NeuroBehavioral Systems, Berkeley, CA,

USA).

Fig 1. Creation of stimulus material. (A) An actress wearing a black pullover and black gloves (right

hand marked with a yellow dot) was filmed from above while playing stories with toy objects within a

marked frame. (B) Videos were rotated by 180° to create videos in the first-person perspective (1pp)

using Adobe Premiere Pro CC.

On the basis of two pilot studies, we chose 24 out of originally 30 stories for our stimulus set. Stories

were excluded when they were particularly difficult to imitate or describe. We made sure that all

chosen stories were comparably memorable. We did this by presenting a questionnaire that included

the correct description of each story and additionally one lure description (similar objects and

context, but different story) to a separate sample of 16 participants and checked for outliers in the

discrimination index P_r for each story. One of the 30 stories was excluded due to low memorability.

The 24 final stories existed in three different versions each: (1) an original version (*ori*) as encoded by

the participants, (2) a version in which two adjacent action steps were switched (structure

modification) to elicit structure-based expectancy violations (*str*) and (3) another variation of the

original video in which one object was exchanged (content modification) to elicit content-based

expectancy violations (*con*). Story scripts were created by five experimenters who all had to agree

that the original story as well as modifications thereof were semantically valid (within a toy world).

This was important since we did not want to violate general semantic expectations. For creating

videos with modifications, the respective stories were played and filmed again exactly the way as for

the original video. The only aspect that differed between original and modified versions was a single

change of either the order of two action steps (i.e., one transition out of 7.33 transitions on average,

for structure modifications) or one object (i.e., one object out of 6.95 objects on average, for content

modifications). Modifications were never introduced in the first two action steps to allow for

expectation build-up prior to modifying the videos and an associated prediction error. Furthermore,

no modifications were introduced in the last two action steps either. For examples of different story

versions, see Fig 2 and S1 Table. The exact time point of the modification in each video (in the

following referred to as time of violation) was determined by identifying the video frame which

diverged from the original version.

Fig 2. Different story versions used in the study. Twenty-four stories existed in three different versions each: an original, a structure modification, and a content modification. For the structure modifications, two adjacent action steps were switched compared to the original (here: step 4 and 5). For the content modifications, an object was exchanged compared to the original (here: tortoise instead of guinea pig on the sales counter in step 5). Note that in the fMRI experiment, each participant was only presented with one of the three versions of a story.

Six additional stories existed in only one version. Four of those served as so-called novels in the fMRI session to provide an additional control condition, which will be discussed in the paper by Jainta et al. (in preparation). The two remaining videos were used for practice trials and did not appear in the fMRI experiment. The stimulus material will be made available at the Action Video Corpus Muenster (https://www.uni-muenster.de/IVV5PSY/AvicomSrv/). Some example videos are available in an OSF repository (https://osf.io/34knw/?view_only=3194f4ad2fa54a0fa47f4189b2aa1f77).

Procedure

Training

Training sessions were conducted in a computer lab at the Department of Psychology at the University of Münster. The training consisted of two sessions that took place on two consecutive days and lasted about 2 h and 1.5 h, respectively. During each of the two sessions, participants encoded half of the episodes. We chose to split the training over two days to avoid fatigue or a decrease in motivation due to the relatively long duration of the task. For encoding episodes, participants imitated half of the 24 stories from demo videos (i.e., the original versions) and only observed the other half. Please note that this experimental factor (*agency*) will not be further addressed in this paper, but in a companion paper (Jainta et al., in preparation).

The 24 demo videos were organized in four subsets, containing six videos each, balanced for the

number of action steps (*A1, A2, B1, B2*). Participants either imitated videos belonging to *A* subsets

and observed videos belonging to *B* subsets or vice versa. In each training session, participants went

through one imitation block and one observation block, each consisting of one subset of videos. This

means that each participant trained each video either during session one or during session two, the

same video was not trained on both days. The order of videos within these blocks was randomized

for each participant. All stories were imitated and only observed equally often over the course of the

study. The first training session started with a short practice phase of the task. Then, half of the

participants started with imitation, the other half with observation. The same applied to the second

training session, except that participants who had started with imitation on the first day now started

with observation, and vice versa. The order of blocks and the order in which video subsets were

trained was balanced between participants. Participants always went through each story eight times,

including four times in 3pp and another four times in 1pp. If the story was to be imitated,

participants first watched the story four times in 3pp, then once in 1pp and then performed the last

three repetitions of the story themselves. If the story was only observed, the video was presented

three more times in 1pp. The optimal number of video presentations for successful imitation was

determined in a pilot study. Fig 3 depicts the structure of the training.

Fig 3. Training procedure. Training consisted of two sessions on two consecutive days (middle row).

The 24 videos to be encoded were organized in four subsets with 6 videos each (*A1, A2, B1, B2*).

Participants either imitated videos belonging to *A* subsets and observed videos belonging to *B*

subsets or vice versa. In each training session, participants went through one imitation block and one

observation block, each consisting of one subset of videos. Hence, Participants imitated and

observed half of the stories each (balanced between participants), equally divided between the two

sessions. In an imitation block (top row), each demo video was shown four times in 3pp and once in

1pp, followed by three correct imitations by the participant. In an observation block (bottom row),

participants watched each demo video four times in both perspectives. After each video, the

participant had to give a detailed description of the story (not depicted here).

During training, the participant sat at the same setup which had been used for filming the stimulus

material and likewise wore a black pullover and gloves with a yellow dot on the right hand. The

experimenter sat opposite of the participant, supervising the performance (Fig 4). For each story, the

toys were positioned next to the camera frame, following the same arrangement as used while

creating the stimulus material. After imitation or observation, participants had to deliver a detailed

description of the story to ensure that they understood it correctly and had paid attention to all

objects involved. If participants made a mistake during an imitation or description trial, they were

immediately interrupted by the experimenter to avoid encoding of incorrect scripts. They would then

start over with a new imitation/description attempt.

Fig 4. Training setup. During training participants observed and imitated toy stories, while sitting at

the filming setup. Their performance was monitored by the experimenter.

Cover-story

Participants were told that videos of themselves playing the stories would be presented in the fMRI

session. Even though participants were actually filmed, these videos were not used during the fMRI

experiment. This was only a cover story to elevate personal identification with the videos to benefit

episode reactivation. In a previous pilot study, we examined whether participants could be convinced

that they are shown in the videos one week after training. Participants were asked for all videos

(presented in pseudo-random order) whether they believed to be shown themselves, rated on a

scale from 0 – 4 (0 = no, 1 = rather no, 2 = rather yes, 4 = yes). We found a significant main effect of

agency (repeated measures analysis of variance, $F_{(1,15)}$ = 33.130, $p < 0.001$, ηp^2 = 0.688, n = 16), driven

by higher identification ratings for imitated than only observed videos (M_{imi} = 2.89, SD_{imi} = 0.44; M_{obs}

= 2.12, SD_{obs} = 0.59), which showed that participants were convinced by our cover story. In our

current study, participants were asked to indicate in percent in how many videos they had been

depicted themselves during the fMRI session. On average, they believed that 40.5 % percent of the

videos showed themselves (SD = 21.8 %). Please note that this cover story was most relevant for the

experimental factor *agency* which is the subject of a different paper. Participants were fully

debriefed after completion of the study.

fMRI session

The fMRI session took place approximately one week after the second training session (range = 7 - 12

days; M = 7.42 days, SD = 0.90 days). Participants were presented with original and modified videos

reminiscent of the previously encoded episodes. Importantly, each video was only shown in the

original or one divergent version. Thus, eight videos were repeatedly presented in the original

version, eight included a structure modification and eight a content modification. Which stories

belonged to which conditions was counterbalanced between participants, so that over the course of

the study, all stories equally often belonged to all experimental conditions. Half of the stories in the

original and modification conditions were shown in 1pp, the other half in 3pp. The perspective of

videos was counterbalanced between participants. This experimental factor (*perspective*) is not the

subject of the present paper, but of another paper by Jainta et al. (in preparation). Additionally, four

novel stories that had not been part of the training were included in the fMRI session to serve as an

additional control.

The fMRI experiment consisted of six blocks, each containing the 24 videos reminiscent of the

previously encoded episodes. Consequently, each video was presented six times over the course of

the session. Within blocks, videos were presented in pseudo-random order. Additionally, each block

contained three null events during which only a fixation cross was presented (duration: 7 – 10 s).

Furthermore, each novel video was presented once per block. Therefore, the whole experiment

contained 18 null events and 24 novel video trials. Participants were not informed about the block-

structure of the experiment.

Participants were instructed to attentively watch the presented videos. They were told that after

some videos a short description would be presented (e.g. "Rescuing princess") that either matched

or did not match the story shown in the video (question trials). The task was to either accept or reject

the description by pressing one of two buttons on a response box with the right index or middle

finger, respectively. This type of task has been used successfully before to focus participants'

attention on complex video stimuli [55]. The descriptions represented the stories on a gist-level, so

specific details were irrelevant to the correct answer (Fig 5). Throughout the entire experiment, each

story was once followed by a matching description and once by a non-matching description, resulting

in a total number of 56 question trials in the experiment. Each block contained nine to ten question

trials and per block, approximately 50 % of descriptions were to be accepted, 50 % were to be

rejected. The question was presented for a maximum of 3 s or until participants responded. Upon

response delivery, participants received 1 s written feedback whether they answered correctly,

incorrectly, or too late, in case no response was given. Participants were naïve with regard to this

distribution of question trials.

Between trials, a fixation cross was presented for a duration of 2 s (1 s after question trials) to serve

as an interstimulus interval. Before each trial, a variable jitter of 0, 0.5, 1 or 1.5 s of fixation was

added for enhancement of the temporal resolution of the BOLD response (Fig 5).

Fig 5. Schematic depiction of task during fMRI session. Video trials consisted of a variable jitter (0,

0.5, 1, or 1.5 s of fixation), a video showing a toy story (ca. 9 – 18 s) and a 2 s interstimulus interval

(fixation). Question trials included a variable jitter, a question regarding the story shown in the

preceding video (maximally 3 s long or terminated by response), and a 2 s interstimulus interval. The

interstimulus interval after question trials was divided into 1 s feedback ("correct", "incorrect", "too

late") and 1 s fixation. Aside from the question, it was depicted which button should be pressed to

accept (left, green) or reject (right, red) the description.

In total, the fMRI task had a duration of approximately 48 min. The task had previously been

practiced at the end of the second training session (four video trials, four question trials, one null

event). Practice videos were shown twice each and not re-used in the fMRI session.

Memory test

Immediately after the fMRI session, participants completed an explicit memory test. Importantly,

encoding occurred incidentally, as participants were not informed beforehand that their memory for

episodes would be tested.

Participants were seated in a separate room in front of a laptop and instructed to remember their

training sessions one week prior. They were presented all stories that they had seen in the fMRI

session in two different versions. More precisely, when expectations had been violated during the

fMRI session by the presentation of episode videos containing a structure modification, these

modified videos were presented again during the memory test and *additionally* each story was

shown in the original version. The same applied when expectations had been violated by content

modifications. When expectations had not been violated, i.e., when original episode videos had been

presented during the fMRI scan, these original videos were presented again in the memory test and

additionally, each story was shown in a modified version, *either* containing a structure modification

in half of the cases *or* a content modification.

The participants' task was to rate after each video whether they knew this exact episode from the

training, using a Likert scale including 1 (*"yes"*), 2 (*"rather yes"*), 3 (*"rather no"*) and 4 (*"no"*), by

pressing one out of four marked keys on the laptop's keyboard. Response time was not restricted but

participants were instructed to respond quickly and intuitively. Videos were always presented in the

same perspective (1pp or 3pp) as during the fMRI session and in a pseudo-randomized order, so that

half of the stories (of each experimental condition) were first presented in their original version

followed by a modified version and vice versa. Novels were shown twice in the same version, so that,

in total, the memory test comprised 56 video trials. The completion of the task took approximately

15 min.

MRI data acquisition and preprocessing

MRI scans were conducted with a 3-Tesla Siemens Magnetom Prisma MR tomograph (Siemens,

Erlangen, Germany) using a 20-channel head coil. Participants lay supine on the scanner bed with

their right index and middle finger positioned on the two appropriate buttons on a response box.

Head, arm and hand movements were minimized by tight fixation with form-fitting cushions.

Participants were provided with earplugs and headphones to attenuate scanner noise. Stimuli were

projected on a screen that the participants saw via an individually adjusted mirror which was

mounted on the head coil.

High resolution T1 weighted anatomical images were obtained with a 3D-multiplanar rapidly

acquired gradient-echo (MPRAGE) sequence prior to functional imaging. 192 slices with a thickness

of 1 mm were acquired, using a repetition time (TR) of 2130 ms, an echo time (TE) of 2.28 ms, a flip

angle of 8° and a field of view (FoV) of 256 x 256 mm². Functional images of the whole brain were

acquired in interleaved order along the AC–PC plane using a gradient-echo echoplanar imaging (EPI)

sequence to measure blood-oxygen-level-dependent (BOLD) contrast. Thirty-three axial slices with a

thickness of 3 mm were obtained in an interleaved order, using a TR of 2000 ms, a TE of 30 ms, a FoV

of 192 x 192 mm² and a flip angle of 90°.

Processing of imaging data was conducted with SPM12 (Wellcome Trust, London, England)

implemented in MATLAB (version R2018b, The MathWorks Inc., Natick, MA, USA). Data were

preprocessed by slice time correction to the middle slice, movement correction and realignment to

the mean image, co-registration of the functional data to individual structural scans, normalization of

functional and structural images into the standard MNI space (Montreal Neurological Institute,

Montreal, QC, Canada) on the basis of segmentation parameters and spatial smoothing using a

Gaussian kernel of full-width at half maximum (FWHM) of 8 mm. Further, a 128 s high-pass temporal

filter was applied.

Statistical data analysis

Behavioral data analysis

The behavioral data analysis was conducted using RStudio ([56]; version 1.3.1073).

After completion of the memory test task, one participant reported that she had not understood it

correctly, so data from this participant was excluded from the analysis (n = 35).

For the analysis of P_r values (general memory performance; unmodified videos = targets, modified

videos = distractors) and ratings/reaction times for modified videos in the memory test, we applied a

2 x 2 within-subject factorial design with the factors *violation* (*yes* = expectancy violation during fMRI

session; *no* = no expectancy violation) and *alternative* (*str* = structure modification included in

memory test; *con* = content modification included), giving rise to the four factorial combinations

no,str (1), *no,con* (2), *yes,str* (3) and *yes,con* (4). Please note that due to the described allocation of

videos to factorial combinations, there were always twice as many trials belonging to factor level *yes*

(8 trials) than *no* (4 trials). For each factorial combination, we calculated the discrimination index P_r

(hit rate minus false alarm rate [57]), the mean rating score, and the mean reaction time for correct

responses. In cued memory tasks, reaction time has traditionally been used as a measure for how

long it takes to retrieve information from memory [58]. Longer reaction times are thought to indicate

increased difficulty of retrieval due to higher cognitive processing demands [59, 60].

For analyzing responses to original (unmodified) videos in the memory test, videos were allocated to

the three different conditions *ori* (no expectancy violation during fMRI session), *str* (structure

expectancy violation during fMRI session) and *con* (content expectancy violation during fMRI

session). Mean rating scores as well as mean reaction times were calculated. Additionally, we

calculated the mean rating scores for novel videos. Please note that low mean rating scores indicate

high acceptance, while high mean rating scores mean low acceptance.

As a next step, we conducted an explorative analysis on behavioral data from the fMRI session. We

calculated the error rate and mean reaction time for each condition (*ori, str, con*) per participant. In

only 0.7 % of all question trials no answer was given, so that such trials were not further considered

in the analysis.

Data were inspected for normal distribution using the Shapiro Wilk Test. Further, data were checked

for extreme outliers as defined as values above quartile 3 + 3 x interquartile range (*IQR*) or lower

than quartile 1 - 3 x *IQR*. When data were normally distributed or could be transformed to fit normal

distribution (reaction times for original videos, reaction times during fMRI task; logarithmic

transformation) and showed no extreme outliers, we used parametric analysis approaches (one-way

repeated measures analyses of variance (rmANOVA)). Post-hoc pair-wise comparisons were

conducted with paired *t*-tests (one-tailed when comparing *ori* and *str* and *ori* and *con* conditions,

two-tailed when comparing *str* and *con* conditions). When the prerequisites for parametric analysis

were not met, we used a non-parametric two-way rmANOVA based on aligned rank data (package

ARTool [61]; P_r values, ratings of modified videos) or the Friedman test for one-way approaches

(ratings of original videos, error rates in fMRI task). Several participants did not give any correct

answers (i.e. rejection) in response to modified videos in the post-fMRI memory test for one or more

factorial combinations. To avoid the listwise exclusion of many datapoints for reaction times when

using a rmANOVA approach, we decided to use one-tailed Wilcoxon signed rank tests when

comparing reaction time between levels of the factor *violation*, separately for each modification

type. This way, we could include 16 datapoints for structure and 29 datapoints for content modified

videos. We additionally excluded another datapoint for content modifications due to a very long

mean reaction time in the *yes* condition (8.85 s) which likely does not represent purely mnemonic

processes. Please note that this did not influence the overall outcome of the analysis.

When parametric statistical approaches were used, we report (untransformed) mean values and

standard errors of the mean as descriptive statistics, when non-parametric approaches were applied,

we report median values and *IQR*. For all behavioral analyses, we applied a significance level of $\alpha =$

0.05. *P*-values were adjusted according to the Bonferroni correction for multiple comparisons [62].

Raw and processed behavioral data have been deposited in an OSF repository

(https://osf.io/34knw/?view_only=3194f4ad2fa54a0fa47f4189b2aa1f77).

fMRI design specifications

Statistical analyses of the fMRI data were conducted with SPM12. We used a general linear model

(GLM) for serially autocorrelated observations [63, 64] and convolved regressors with the canonical

hemodynamic response function. Regressors were original videos (*ori*), videos containing a structure

modification (*str*) and videos containing a content modification (*con*), each comprising 48 trials. For

str and *con* trials, the onsets of events were time-locked to the point in the video at which the

modification occurred (time of violation). For *ori* trials, we calculated a hypothetical time of violation

(mean of times that corresponded to points of structure and content modification in the non-

modified video) to serve as a comparable onset. These conditions were modeled as events as we

were interested in the phasic effect of the expectancy violation at the precise moment it occurred.

To each of those regressors, we added a parametric modulator to model the repeated presentation

of each video. Two additional regressors modeled the 18 null events and the 56 question trials. The

modeled activation of null events and questions was time-locked to their respective onsets. Null

events were modeled as epochs, containing their full presentation time (7 – 10 s), while questions

were modeled as events. The six subject-specific rigid-body transformations obtained from

realignment were included as regressors of no interest. Therefore, the GLM comprised 14 regressors

in total. Please note that factors which were not considered in the present paper (*agency*,

perspective) were perfectly balanced over the factors of interest so that we can rule out any

confounding effects (fully-crossed design).

We calculated first-level-*t*-contrasts for *str > ori* and *con > ori* to analyze brain activity in response to

the specific expectation violation types. To exploratively examine whether the BOLD response to the

violations attenuated over repetitions, we additionally considered *t*-contrasts for the parametric

regressors. Grey matter masking was applied on the first level of the analysis. For masking, we used

the smoothed individual normalized grey matter image (8 mm FWHM) which was thresholded at 0.2

using ImCalc in SPM12 to create a binary mask. Second-level group analyses were performed by

using one-sample *t*-tests across participants. A conjunction of *str > ori* and *con > ori* contrasts was

calculated to detect shared effects of both expectancy violations [65]. We applied false discovery

rate (FDR) correction and a threshold of $p < 0.05$ (peak level). In case no significant activation was

detected using this threshold, we applied no correction for multiple comparison with $p < 0.001$. This

is a common procedure when specific neuroanatomical hypotheses are examined [66, 67, 68, 69].

Brain activation patterns were visualized with the software MRIcroGL (version 1.2.20200331,

McCausland Center for Brain Imaging, University of South Carolina, USA). Unthresholded statistical

maps have been uploaded to NeuroVault.org ([70], https://neurovault.org/collections/SKPCADYC/).

To explore whether IFG activity is modulated by the two types of expectancy violation, we performed

region of interest (ROI) analyses for two different IFG areas (pars opercularis, BA 44; pars triangularis,

BA 45). Separate structural ROIs were generated per IFG area and hemisphere using the automated

anatomical labeling atlas with the WFU PickAtlas toolbox [71] in SPM12. Mean beta values for the *str*

> ori and *con > ori* contrasts were extracted from each ROI using the MarsBaR toolbox [72]. The

conditions for parametric analysis were confirmed as described above for behavioral data. One-tailed

one-sample *t*-tests were used to test whether mean beta values per ROI and contrast were greater

than zero. For each ROI, mean beta values were compared between *str > ori* and *con > ori* using two-

tailed paired *t*-tests. There were no specific hypotheses regarding differences between left and right

hemispheres, so we did not compare them directly. Because we analyzed four ROIs per contrast and

performed multiple pair-wise comparisons, we applied the Bonferroni correction [62] and report

adjusted *p*-values, for both one-sample and paired *t*-tests. The extracted mean beta values were

deposited in an OSF repository

(https://osf.io/34knw/?view_only=3194f4ad2fa54a0fa47f4189b2aa1f77).

Results

Post-fMRI memory test

Discrimination index P_r

We found a significant main effect of *violation* ($F_{(1,34)}$ = 20.220, p < 0.001, ηp^2 = 0.373) on P_r values

which was driven by higher P_r values for *no* (*Median* = 0.50, *IQR* = 0.50) compared to *yes* (*Median* =

0.19, *IQR* = 0.50). There was also a significant main effect of *alternative* ($F_{(1,34)}$ = 85.828, p < 0.001, ηp^2

= 0.716), which was explained by higher P_r values for *con* (*Median* = 0.63, *IQR* = 0.38) than for *str*

(*Median* = 0.06, *IQR* = 0.25). There was no significant interaction of *violation* and *alternative* ($F_{(1,34)}$ =

2.074, p = 0.159, ηp^2 = 0.058) (Fig 6A).

Ratings and reaction times for modified videos

We found a significant main effect of *violation* on mean rating scores for modified videos ($F_{(1,34)}$ =

25.554, p < 0.001, ηp^2 = 0.429), which was explained by higher ratings for *no* (*Median* = 2.5, *IQR* =

1.75) than for *yes* (*Median* = 2, *IQR* = 1.75). Thus, when an expectancy violation had occurred during

the fMRI session, participants were more likely to accept modified versions afterwards. Additionally,

there was a significant main effect of *alternative* on mean rating scores for modified videos ($F_{(1,34)}$ =

122.647, p < 0.001, ηp^2 = 0.783). Higher ratings were detected for *con* (*Median* = 3.25, *IQR* = 1.375)

than for *str* (*Median* = 1.563, *IQR* = 1.094), which means that participants were generally more prone

to accept structurally modified videos than those with modified content. There was no significant

interaction of the two factors *violation* and *alternative* ($F_{(1,34)}$ = 2.010, p = 0.165, ηp^2 = 0.056; Fig 6B).

We found a near-significant trend that reaction times were longer for structurally modified videos

when this version had already been presented in the fMRI session (Z = -1.80, p = 0.072; Fig 6C). No

significant difference was found concerning reaction times for videos with content modification ($Z = -$

1.089, $p = 0.276$; Fig 6D).

Ratings and reaction times for unmodified videos

Mean rating scores for unmodified videos did not differ significantly between conditions ($\chi2_{(2)} =$

2.124, $p = 0.346$, *Kendall's W* = 0.030). Descriptively, rating scores were slightly higher when repeated

expectancy violation had occurred during the fMRI session ($Median_{ori} = 1.0$, $IQR_{ori} = 0.313$; $Median_{str} =$

1.25, $IQR_{str} = 0.375$; $Median_{con} = 1.125$, $IQR_{con} = 0.5$), which might indicate a tendency to accept

original videos slightly less after repeated expectancy violation (Fig 6E).

There was a significant main effect of condition on mean reaction times for correctly classifying

unmodified videos ($F_{(2,68)} = 3.725$, $p = 0.029$, $\eta p^2 = 0.099$). Pairwise comparisons revealed that

reaction times were significantly longer after structure expectancy violation in the scanner than

when no expectancy violation had occurred ($t_{(34)} = 3.287$, $p = 0.004$). No significant differences were

detected between *con* and *ori* ($t_{(34)} = 1.779$, $p = 0.126$) and *str* and *con* conditions ($t_{(34)} = 0.725$, $p = 1$;

Fig 6F).

Fig 6: Behavioral results from post-fMRI memory test. (A) Discrimination index P_r from post-fMRI

memory test. Original (unmodified) videos were the targets while modified videos (*str* = structure

modification, *con* = content modification) were distractors. Statistics: non-parametric rmANOVA

based on aligned rank data. a vs. b = $p < 0.001$ (main effect *violation*); *** = $p < 0.001$ (main effect

alternative). (B) Mean rating scores for modified videos for each factorial combination. Statistics:

non-parametric rmANOVA based on aligned rank data. a vs. b = $p < 0.001$ (main effect *violation*); ***

= $p < 0.001$ (main effect *alternative*. (C) Mean reaction times for structure modified videos, $n = 16$.

Statistics: Wilcoxon signed rank tests. (D) Mean reaction times for content modified videos, $n = 28$.

(E) Mean rating scores for unmodified videos for each condition (*ori* = no expectation violation during

fMRI session, *str* = structure expectancy violation, *con* = content expectancy violation). Statistics:

Friedman test. (F) Mean reaction times for unmodified videos for each condition. Statistics:

rmANOVA, post-hoc pairwise comparisons: paired t-tests, p-values were corrected for multiple

comparisons. ** = $p < 0.01$. Box plots show medians, quartiles, minima and maxima. Bar plots show

means and standard errors.

Ratings for novel videos

546 High ratings for novel videos, i.e., videos which were new during the fMRI session ($Median = 4$, $IQR =$

547 1), indicate that participants correctly remembered that these videos had not been part of the

548 training.

Behavioral performance during fMRI session

Error rates of the task during the fMRI session were generally very low ($Median_{ori} = 0$, $IQR_{ori} = 0$;

$Median_{con} = 0$, $IQR_{con} = 0.02$; $Median_{str} = 0$, $IQR_{str} = 0$) and did not differ significantly between

conditions ($\chi2_{(2)} = 0.520$, $p = 0.771$, $Kendall's\ W = 0.007$). We also did not find a significant main effect

of condition on mean reaction times during the fMRI session ($F_{(2,70)} = 0.983$, $p = 0.380$, $\eta p^2 = 0.027$).

These findings indicate that the participants' attention to (and understanding of) the stories was very

good and not deteriorated by the switch of single action steps or objects. Importantly, this shows

that any neural differences between these conditions are not simply based on confusion or generally

elevated cognitive effort in respect to expectancy violations. All questions in the fMRI session asked

for the gist of the stories. We can thus assume that neither the structure, nor the content

modification changed this gist compared to the original story.

fMRI results

First, we investigated whether structure and content expectation violations elicit common brain

activation patterns compared to original episodes. The conjunction of the whole brain contrast $str >$

ori and $con > ori$ did not yield significant activation after correction for multiple comparisons.

Without correction and at a threshold of $p < 0.001$, there was significant activation in the opercular

part of the right IFG which corresponds to BA 44 ($x = 51$, $y = 17$, $z = 20$, 44 voxels).

To identify which brain regions are active for structure expectancy violation, we calculated the whole

brain contrast *str > ori*. This contrast did not yield significant activation after correction for multiple

comparisons. Without correction and at a threshold of $p < 0.001$, activation clusters were detected in

the right hemisphere in SFS, PMd and the opercular part of the IFG (BA 44) (Fig 7 and Table 1).

To detect brain regions which respond to content expectancy violations, we calculated the whole

brain contrast *con > ori*. Significant bilateral activation was found in pre-SMA and the middle IPS.

Additionally, there were activation clusters in left and right inferior frontal sulcus (IFS) which

expanded into the triangular part of the IFG (BA 45) and the inferior frontal junction (IFJ). There was

significant bilateral activation in the anterior insula. Finally, elevated brain activity was detected in

the occipitotemporal cortex (OTC), bilaterally including FG and extending into parahippocampal gyrus

(PHG) in the right hemisphere (Fig 7 and Table 1).

Fig 7. Whole brain activation for expectancy violation effects. (A) Uncorrected *t*-map ($p < 0.001$) for

the *str > ori* contrast. (B) FDR-corrected *t*-map ($p < 0.05$) for the *con > ori* contrast. IFG: inferior

frontal gyrus; SFS: superior frontal sulcus; PMd: dorsal premotor cortex; IFS: inferior frontal sulcus;

pre-SMA: pre-supplementary motor area; IPS: intraparietal sulcus; AI: anterior insula; FG: fusiform

gyrus; PHG: parahippocampal gyrus.

Table 1. Peak activations from second-level whole-brain analyses of expectancy violation effects.

Localization	H	Cluster extent	MNI Coordinates			*t*-value
			x	y	z	
str > ori (uncorrected at $p < 0.001$)						

Superior frontal sulcus	R	16	27	23	59	3.72
Dorsal premotor cortex	R	48	39	2	56	4.74
Inferior frontal gyrus (pars opercularis, BA 44)	R	70	54	14	20	4.63

con > ori (FDR-corrected at $p < 0.05$)

Pre-supplementary motor area	L	49	-6	14	53	4.16
	R	64	9	17	47	4.11
Intraparietal sulcus	R	294	30	-61	50	6.18
	L	241	-27	-64	38	5.41
Inferior frontal junction	L	320	-39	2	32	6.38
Inferior frontal sulcus, extending into inferior frontal gyrus (pars triangularis, BA 45)	L		-42	20	26	5.51
	R	505	45	29	20	6.99
Anterior insula	R	134	33	20	-1	5.58
	L	143	-30	20	-4	5.70
Occipitotemporal cortex	L	236	-45	-55	-13	7.48
Fusiform gyrus	L		-27	-40	-22	3.87
Occipitotemporal cortex, including fusiform gyrus and extending into parahippocampal gyrus	R	259	33	-52	-13	5.10

Note: H = Hemisphere, MNI = Montreal Neurological Institute, L = Left, R = Right, *str* = structure

expectancy violation, *con* = content expectancy violation, *ori* = no expectancy violation.

For an exploratory, more detailed analysis of IFG activity, we extracted beta values from different IFG

areas (pars opercularis, BA 44; pars triangularis, BA 45) for the *str > ori* and *con > ori* contrasts. As we

did not have specific hypotheses concerning differences between left and right hemisphere, we

extracted beta values separately for both hemispheres but did not directly compare them. One-

sample *t*-tests against zero revealed significant activation for content expectancy violations in left

and right BA 45 as well as in right BA 44. For structure expectancy violations, significant activation

was only found for right BA 44. There were near-significant trends for activation ($p < 0.01$) in left BA

44 for content expectancy violations and in right BA 45 for structure expectancy violations (Fig 8). For

a detailed overview of all one-sample t-tests, see S3 Table. Paired t-tests showed that mean beta

values were significantly higher for content expectancy violations compared to structure expectancy

violations in left and right BA 45 (left: $t_{(35)}$ = 4.545, p < 0.001; right: $t_{(35)}$ = 4.005, p < 0.001). We did not

detect significant differences between conditions in BA 44 (left: $t_{(35)}$ = -2.053, p = 0.191; right: $t_{(35)}$ = -

0.771, p = 1; Fig 8).

Fig 8. Region of interest analysis of left (A) and right (B) opercular (BA 44) and triangular (BA 45)

part of the IFG. Mean beta values for the contrasts *str > ori* and *con > ori* were extracted from

subregions of the IFG as defined by the automated anatomical labeling atlas (depicted on the right).

Bar plots show means and standard errors. One-tailed one-sample t-test were used to determine

whether values were greater than zero (symbols above bars). Two-tailed paired t-tests were used to

determine differences between conditions (symbols above lines). *P*-value were corrected for multiple

comparisons. ** = p < 0.01; *** = p < 0.001.

Regarding the explorative analysis of attenuation of the BOLD response to structure expectancy

violations, we observed only sub-threshold attenuation in the IFS, occipitotemporal areas (including

FG), IPS and thalamus. In contrast, several regions showed significant attenuation in for content

expectancy violations, including posterior precuneus, OTC (including FG and PHG), pIPS, thalamus

and the right inferior frontal junction (S1 Fig and S2 Table)

Discussion

Episodic memories allow us to mentally relive our personal past. Against what we might believe, they

are not exact imprints of previous experiences, as they are prone to later modification, potentially

triggered by prediction errors. In the present study, we investigated brain and behavioral responses

to expectancy violations based on two properties which characterize episodic memories, namely

structure and content information. Participants encoded episodes in the form of toy stories. By

presenting modified videos in a subsequent fMRI session, we aimed at violating the brain's

expectations based on episode structure or content. While brain responses were largely as expected

for content expectancy violations, only subtle activation was detected following structure expectancy

violations, potentially because structure modifications were less salient. An explicit memory test

revealed that the acceptance of original videos slightly decreased after expectancy violation. Further,

we found an increased tendency to accept modified versions when the same had been presented

during the fMRI scan. Together, findings provide first evidence that different types of mnemonic

prediction errors are processed differently by the brain and may contribute to memory changes.

629 **Influence of expectancy violation on post-fMRI memory**
630 **performance**

As expected, episodes that had been reactivated in a modified version during the fMRI experiment

were generally remembered worse, as reflected by lower discrimination indexes (P_r). More

specifically, modified videos were more likely to be perceived as original stories thereafter, as

evidenced by significantly decreased rating scores. Similarly, others have reported that mnemonic

prediction errors lead to an intrusion of new information into an established memory repertoire [20,

22]. In our study, participants were only descriptively less prone to accept original stories following

expectation violation in the scanner. This finding is in line with previous reports that the inclusion of

new information into an old memory is not necessarily accompanied by a deterioration of the

original memory [73, 22]. Thus, it is likely that the prediction errors in the current study left the

original memory traces largely intact but led to the encoding of an additional, alternative

representation of the same episode. This finding contrasts with the results of a study by Kim et al.

[27], who showed that prediction violations led to decreased recognition of original memory content.

However, in their study, original memories were encoded by showing the to-be-predicted sequence

only once, whereas participants in our study had experienced and actively encoded each story

multiple times. Thus, it may be that memory traces were particularly stable in the current paradigm,

such that a greater number of repeated alternative evidences would be required to significantly

reduce memory performance for the initially encoded episodes. All memory changes we observed in

the present study were rather subtle, as median values indicated a change in the degree of certainty

rather than a complete transition from acceptance to rejection or vice versa. Participants also took

longer to correctly classify unmodified videos when they had previously experienced structure

expectancy violations. This tendency was also observed for modified videos. Longer response times

in cued-memory paradigms are interpreted as indicative of increased difficulty of retrieval due to

higher cognitive processing demands [59, 60]. Thus, it is likely that it became more difficult to

differentiate between alternative competing versions of episodes when versions diverging from the

original experience had been already encountered in the fMRI session.

Taken together, our results suggest that violations of both, structure and content expectations,

influenced subsequent memory performance for affected episodes. These findings are in line with

the idea that mnemonic prediction errors can trigger episodic memory modification [19]. We can,

however, not rule out that the repeated presentation of modified versions during the fMRI scan led

to a confusion of different versions, which could be an alternative explanation for our findings. On

the other hand, participants remembered correctly that novel videos had not been part of the

original episode repertoire even though they had repeatedly been presented during the fMRI

experiment as well. This finding suggests that our post-test results are at least not a product of

source confusion in its simplest form. Even if our results are to be attributed to confusion-effects,

this would not per se contradict the idea of memory modification. In fact, it is not yet clear exactly by

what mechanism memory modifications are brought about, and it has been suggested that they may

result, at least in part, from interference between old and new memory traces [74, 75, 22].

Another interesting behavioral finding was that participants generally had a strong tendency to

accept structurally modified versions as originals in the memory test. It is likely that structural

modifications were generally less salient than content modifications, although both types of

modifications resulted from a single change in the story (see S1 Table for examples). This result also

fits the fact that after the fMRI session, two-thirds of the participants self-reported noticing at least

one object swap, while less than a quarter of the participants had noticed a change regarding the

order of action steps. Recently, it was reported that memory performance after an expectancy

violation differed depending on whether changes were detected (and remembered) by participants

or not [18]. While undetected changes led to reduced memory performance, detected changes had

the opposite effect. Although our results suggest that structural changes were probably significantly

more difficult to detect, content expectancy violations led to an increase in the acceptance of

modified version, i.e. reduced memory performance, as well.

The current study provides evidence that different types of mnemonic prediction errors can modify

memory. Interestingly, others have found that prediction errors can also have to opposite effect, that

is improve subsequent memory [16, 17, 18]. Thus, it seems that the effect of expectation violation on

memory underlies a complex dynamic which might depend on the strength of prediction errors. It

has been suggested that strong expectancy violations lead to better memory by promoting learning

of new associations [17]. Additionally, no/weak prediction errors were also found to yield better

memory performance [76], and it has been suggested that expectancy violations need to overcome a

certain threshold to initiate memory updating [10]. Overall, it seems like the strength of prediction

errors modulates memory in a U-shaped fashion [76]. These findings are in accordance with the

theoretical framework by Ritvo et al. [77], which would predict a decrease in memory performance

following moderate memory reactivation. In conclusion, it is likely that mnemonic prediction errors

in the current study were medium-sized, i.e. large enough to decrease, but too small to increase,

memory performance. Within this range, prediction errors caused by structural deviance might have

been smaller than those elicited by content deviance, as our behavioral data suggests that the latter

were more salient. However, one must keep in mind that much of the previous research used one-

shot learning to study the effects of prediction errors on memory [17, 22]. In our paradigm, we

intentionally took a different approach, namely, repetition of the same divergent information that

often gives rise to learning processes in daily life. The aim of the current study was to characterize

brain and behavioral responses to different types of mnemonic prediction errors for this first time

with a novel paradigm. In contrast, we did not specifically aim to investigate prediction error

strengths, so this aspect is not discussed in detail here.

Neural responses to different episodic expectancy violations

Content and structure violation of expectation were expected to draw on distinct brain areas, but

also to share some common violation-induced activation in mPFC, AON, and the hippocampal

formation. Without correction for multiple comparisons, common activation was detected in right

IFG, which reflects our previous finding that IFG increases for inconsistent or highly informative detail

in observed actions [78, 79, 55]. After correction for multiple comparisons, no common violation-

related activity could be observed. At this point, it is not possible to distinguish whether this was due

to a large functional disparity between content and structural information in episodic memories or

simply to the brain's overly subtle response to violations of structural expectancy.

On the one hand, we had expected that structure expectation violations lead to activation in brain

regions involved in the temporal organization of episodes. We did not find significant activation

patterns after correction for multiple comparisons. Without such correction, we detected activation

in a hypothesized area, right PMd/SFS. The activity of a region comprising PMd/SFS is related to

linking successive action steps [38, 39, 41, 42], and could contribute to updating the current event or

action model with respect to each next segment [80, 40, 39, 42].

On the other hand, we found content expectancy violations to specifically recruit pIPS and OTC,

including FG, which were hypothesized on the basis of their role for processing of object properties in

the context of actions [81, 46, 82, 43, 55]. More specifically, pIPS encodes basic visual features of

graspable objects [44, 83], reflecting the interaction with toy objects in our paradigm. Additionally,

content expectancy violations elicited activity in the hippocampal formation (right PHG). In general,

hippocampus and PHG are important in learning contexts [84, 85, 86, 87] and there is evidence that

the PHG is involved in processing of competing memories [88]. Moreover, the hippocampal

formation is believed to generate mismatch signals when predictions do not fit perceptual inputs [69,

89, 20]. Because our post-fMRI memory test data imply that content changes were more salient than

structural changes, one could speculate that the discrepancy between prediction and perception was

strong enough to be reflected in the activation of the hippocampal formation for content expectancy

violations. This would also fit with our finding that content expectancy violation triggered activity in

the anterior insula, which is relevant for error detection [90] and salience processing [91].

Similar to our conjunction analysis, the ROI analysis in different IFG regions showed that the right BA

44 responded to both types of episodic violations. These results confirm the assumption that BA 45 is

more responsive to content than to structural expectancy violations. We take this as a first indication

that within IFG, different subregions might process different types of episodic prediction errors.

However, our study cannot rule out the possibility that this effect is merely due to quantitative

(rather than qualitative) differences between structural and content expectancy violation, as

structure modifications might have elicited weaker prediction errors in our paradigm.

In order to interpret brain activity triggered by violating expectations considering retrieved episodes,

it is informative to consider attenuation of this response when participants repeatedly encountered

modified stories, as this gives insight into adaptation processes during model updating [24, 25]. In

fact, such neuronal repetition suppression has been suggested to occur due to a decrease of

prediction error strength [92], which would be in line with idea that memory modification occurs due

to gradual updating of internal models triggered by prediction errors. We thus additionally explored

possible attenuation effects concerning modified episodes. Brain activation was found to attenuate

with repeated exposure to content modifications, indicating ongoing adaptation processes. Some

regions that showed a decrease in brain activation were also identified in the general violation effect

contrast, including PHG. Correspondingly, Schiffer et al. [24] found hippocampal attenuation with

repeated presentation of divergent episodes. They concluded that this activity decrease is a sign of

increasing validity of internal models due to updating. The attenuation effects we found for structure

expectancy violations were quite subtle, so one could speculate that for this type of expectancy

violations, adaptation might be still at an early stage. It is also possible that, at least because of the

substantial attenuation effect on content violation, some brain areas that responded to expectation

violation could no longer be detected in the main effect. For example, attenuation was found

bilaterally in FG, extending into PHG for content expectancy violations. While widespread activation

in FG was also found in response to the violation itself, only a small portion of right PHG showed

significant activation.

756 Limitations and implications for future research

One factor that might limit the generalizability of our findings is that, for practical reasons, only

women participated in the study. However, because the processing of episodic memory in the brain

seems to be broadly similar between women and men [93], we are confident that our findings are

applicable to a more general population. In the future, our paradigm could be adapted to circumvent

such practical limitations, for example, by applying virtual reality techniques so that encoding could

be detached from the true physical appearance of participants' hands.

One potential limitation of the current paradigm is that in our modified episodes, the occurrence of

objects and temporal structures were new. Thus, we cannot completely rule out that aside from

prediction errors, novelty influenced our findings. It would be valuable to find a way to keep novelty

constant and have participants make active predictions which are then either violated or not. Aside

from that, it would be interesting to compare different prediction strengths which we will target in

an upcoming study where we will introduce manipulations of different salience.

Another limitation of the current study is that we only found very subtle brain responses to

structure-expectancy violations, which may be explained by their lower salience. As discussed above,

structure modification might have caused weaker prediction errors than content modifications. The

hypothesized (subthreshold) activation in the right PMd/SFS in response to structure-expectancy

violations could be indicative for qualitative difference in the processing of the two types of

expectancy violation. Nevertheless, an important question arises: could prediction errors be less

informative for complex episode structures in general, or is this finding specific to certain, more

variable circumstances? Our results therefore represent exiting starting points for future research.

We plan to investigate whether additional consolidation of encoded memories through active

retrieval training [94] could enhance the brain response to structure violations. In addition, we will

analyze whether structural modifications of higher salience (e.g., rearrangement of multiple steps, or

rearrangements more significant to the episode) could produce a similar effect.

It is possible that by repeating the same modified versions, attenuation effects may have limited the

detectability of brain regions that process certain types of mnemonic prediction errors. To clarify

this, we will conduct a follow-up study in which we do both, repeating the same modified versions of

an episode, but also including the presentations of constantly changing versions, similar to the

paradigm of Schiffer et al. [24]. Finally, it will be interesting to reanalyze our dataset to link fMRI and

behavioral data to analyze whether specific brain activations predict good or bad memory

performance.

Conclusion

When recalling episodes, our memory can change, for instance due to prediction errors in the re-

activation process. Our results suggest that structural and content prediction errors in episode

retrieval differ in their neural processing. Future studies need to further validate whether the

observed differences are indeed qualitative, and not just quantitative, in nature. This is also

suggested by the fact that the tendency to falsely classify modified episodes as originally experienced

ones increased comparably after experiencing repeated structural as well as content-related

prediction errors. Thus, memory change seems to be able to occur as a result of different types of

prediction errors. In summary, the present findings and our newly developed paradigm provide very

good starting points for further research into the temporal dynamics, limits and conditions of

episodic memory modification.

Acknowledgments

The authors thank Monika Mertens for her help during data collection. Furthermore, we thank

Annika Garlichs, Christin Schwarzer, Helena Sydlik and Yuyi Xu for their assistance during the creation

of stimulus material, data collection and data management. Lastly, we thank Jennifer Pomp, Lena

Schliephake, Falko Mecklenbrauck and Nina Heins for advice regarding data analysis and the

members of research unit FOR 2812 for valuable discussions. This work was funded by the German

Research Foundation (Deutsche Forschungsgemeinschaft) – project numbers 419037023,

419039274, 419037518. The funders had no role in study design, data collection, analysis and

interpretation, decision to publish, or writing of the report.

References

1.   Tulving E. Episodic Memory: From Mind to Brain. Annu Rev Psychol. 2002;53:1–25. doi: 0084-
6570/02/0201-0001.

2.   Nader K, Einarsson EÖ. Memory reconsolidation: An update. Ann N Y Acad Sci. 2010;1191:27–
41. doi: 10.1111/j.1749-6632.2010.05443.x.

3.   Nader K. Reconsolidation and the Dynamic Nature of Memory. Nov Mech Mem. 2015;7(10):1–
20. doi: 10.1007/978-3-319-24364-1_1.

4.   Lee JLC, Nader K, Schiller D. An Update on Memory Reconsolidation Updating. Trends Cogn
Sci. 2017;21(7):531–45. doi: 10.1016/j.tics.2017.04.006.

5.   Scully ID, Napper LE, Hupbach A. Does reactivation trigger episodic memory change? A meta-
analysis. Neurobiol Learn Mem. 2017;142(Pt A):99–107. doi: 10.1016/j.nlm.2016.12.012.

6.   Schacter DL. The cognitive neuroscience of false memories: Introduction. Cogn Neuropsychol.
1999;16(3/4/5):193–5. doi: 10.1080/026432999380753.

7.   Straube B. An overview of the neuro-cognitive processes involved in the encoding,
consolidation, and retrieval of true and false memories. Behav Brain Funct. 2012;8(35):1–10.
doi: 10.1186/1744-9081-8-35.

8.   Suddendorf T, Corballis MC. The evolution of foresight: What is mental time travel, and is it
unique to humans? Behav Brain Sci. 2007;30(3):299–351. doi: 10.1017/S0140525X07001975.

9.   Schacter DL, Guerin SA, St. Jacques PL. Memory distortion: An adaptive perspective. Trends
Cogn Sci. 2011;15(10):467–74. doi: 10.1016/j.tics.2011.08.004.

10.   Exton-McGuinness MTJ, Lee JLC, Reichelt AC. Updating memories-The role of prediction errors
in memory reconsolidation. Behav Brain Res. 2015;278:375–84. doi:
10.1016/j.bbr.2014.10.011.

11.   Rosen R. Anticipatory systems: Philosophical, mathematical, and methodological foundations.
2nd ed. New York, Springer; 2012. doi: 10.1007/978-1-4614-1269-4.

12.   Friston K. A theory of cortical responses. Philos Trans R Soc B Biol Sci. 2005;360(1456):815–36.
doi: 10.1098/rstb.2005.1622.

13.   Friston K, Kiebel S. Predictive coding under the free-energy principle. Philos Trans R Soc B Biol
Sci. 2009;364(1521):1211–21. doi: 10.1098/rstb.2008.0300.

14.   Schubotz RI. Prediction and expectation. In: Toga AW, editor. Brain Mapping: An Encyclopedic

Reference. Cambridge: Academic Press: Elsevier; 2015. pp. 295–302. doi: 10.1016/B978-0-12-
397025-1.00205-0.

15.   Fernández RS, Boccia MM, Pedreira ME. The fate of memory: Reconsolidation and the case of
Prediction Error. Neurosci Biobehav Rev. 2016;68:423–41. doi:
10.1016/j.neubiorev.2016.06.004.

16.   Smith TA, Hasinski AE, Sederberg PB. The context repetition effect: Predicted events are
remembered better, even when they don't happen. J Exp Psychol Gen. 2013;142(4):1298–
308. doi: 10.1037/a0034067.

17.   Greve A, Cooper E, Kaula A, Anderson MC, Henson R. Does prediction error drive one-shot
declarative learning? J Mem Lang. 2017;94:149–65. doi: 10.1016/j.jml.2016.11.001.

18.   Wahlheim CN, Zacks JM. Memory guides the processing of event changes for older and
younger adults. J Exp Psychol Gen. 2019;148(1):30–50. doi: 10.1037/xge0000458.

19.   Sinclair AH, Barense MD. Prediction Error and Memory Reactivation: How Incomplete
Reminders Drive Reconsolidation. Trends Neurosci. 2019;42(10):727–39. doi:
10.1016/j.tins.2019.08.007.

20.   Long NM, Lee H, Kuhl BA. Hippocampal Mismatch Signals Are Modulated by the Strength of
Neural Predictions and Their Similarity to Outcomes. J Neurosci. 2016;36(50):12677–87. doi:
10.1523/JNEUROSCI.1850-16.2016.

21.   Bein O, Duncan K, Davachi L. Mnemonic prediction errors bias hippocampal states. Nat
Commun. 2020;11(1). doi: 10.1038/s41467-020-17287-1.

22.   Sinclair AH, Barense MD. Surprise and destabilize: Prediction error influences episodic
memory reconsolidation. Learn Mem. 2018;25(8):369–81. doi: 10.1101/lm.046912.117.

23.   Scully ID, Hupbach A. Different reactivation procedures enable or prevent episodic memory
updating. Hippocampus. 2020;30(8):806–14. doi: 10.1002/hipo.23159.

24.   Schiffer AM, Ahlheim C, Wurm MF, Schubotz RI. Surprised at all the entropy: Hippocampal,
caudate and midbrain contributions to learning from prediction errors. PLoS One.
2012;7(5):1–11. doi: 10.1371/journal.pone.0036445.

25.   Schiffer AM, Ahlheim C, Ulrichs K, Schubotz RI. Neural Changes when Actions Change:
Adaptation of Strong and Weak Expectations. Hum Brain Mapp. 2013;34(7):1713–27. doi:
10.1002/hbm.22023.

26.   Wichert S, Wolf OT, Schwabe L. Updating of episodic memories depends on the strength of

new learning after memory reactivation. Behav Neurosci. 2013;127(3):331–8. doi:
10.1037/a0032028.

27.   Kim G, Lewis-Peacock JA, Norman KA, Turk-Browne NB. Pruning of memories by context-
based prediction error. Proc Natl Acad Sci U S A. 2014;111(24):8997–9002. doi:
10.1073/pnas.1319438111.

28.   Burianova H, McIntosh AR, Grady CL. A common functional brain network for
autobiographical, episodic, and semantic memory retrieval. Neuroimage. 2010;49(1):865–74.
doi: 10.1016/j.neuroimage.2009.08.066.

29.   Euston DR, Gruber AJ, McNaughton BL. The Role of Medial Prefrontal Cortex in Memory and
Decision Making. Neuron. 2012;76(6):1057–70. doi: 10.1016/j.neuron.2012.12.002.

30.   Peters GJ, David CN, Marcus MD, Smith DM. The medial prefrontal cortex is critical for
memory retrieval and resolving interference. Learn Mem. 2013;20(4):201–9. doi:
10.1101/lm.029249.112.

31.   Maguire EA, Intraub H, Mullally SL. Scenes, Spaces, and Memory Traces: What Does the
Hippocampus Do? Neuroscientist. 2016;22(5):432–9. doi: 10.1177/1073858415600389.

32.   Horner AJ, Doeller CF. Plasticity of hippocampal memories in humans. Curr Opin Neurobiol.
2017;43:102–9. doi: 10.1016/j.conb.2017.02.004.

33.   Stachenfeld KL, Botvinick MM, Gershman SJ. The hippocampus as a predictive map. Nat
Neurosci. 2017;20(11):1643–53. doi: 10.1038/nn.4650.

34.   Van Kesteren MTR, Ruiter DJ, Fernández G, Henson RN. How schema and novelty augment
memory formation. Trends Neurosci. 2012;35(4):211–9. doi: 10.1016/j.tins.2012.02.001.

35.   Caspers S, Zilles K, Laird AR, Eickhoff SB. ALE meta-analysis of action observation and imitation
in the human brain. Neuroimage. 2010;50(3):1148–67. doi:
10.1016/j.neuroimage.2009.12.112.

36.   Schubotz RI. Human premotor cortex: Beyond motor performance. MPI Series in Human
Cognitive and Brain Sciences, vol. 50. Max Planck Institute for Human Cognitive and Brain
Sciences, Leipzig. 2004.

37.   Shima K, Tanji J. Neuronal Activity in the Supplementary and Presupplementary Motor Areas
for Temporal Organization of Multiple Movements. J Neurophysiol. 2000;84(4):2148–60. doi:
10.1152/jn.2000.84.4.2148.

38.   Stadler W, Schubotz RI, von Cramon DY, Springer A, Graf M, Prinz W. Predicting and

Memorizing Observed Action: Differential Premotor Cortex Involvement. Hum Brain Mapp.
2011;32(5):677–87. doi: 10.1002/hbm.20949.

39.   Schubotz RI, Korb FM, Schiffer AM, Stadler W, von Cramon DY. The fraction of an action is
904   more than a movement: Neural signatures of event segmentation in fMRI. Neuroimage.
905   2012;61(4):1195–205. doi: 10.1016/j.neuroimage.2012.04.008.

906   40.   Tamber-Rosenau BJ, Esterman M, Chiu YC, Yantis S. Cortical Mechanisms of Cognitive Control
907   for Shifting Attention in Vision and Working Memory. J Cogn Neurosci. 2011;23(10):2905–19.
908   doi: 10.1162/jocn.2011.21608.

909   41.   Hrkać M, Wurm MF, Schubotz RI. Action Observers Implicitly Expect Actors to Act Goal-
910   Coherently, Even if they Do Not: An fMRI Study. Hum Brain Mapp. 2014;35(5):2178–90. doi:
911   10.1002/hbm.22319.

912   42.   Pomp J, Heins N, Trempler I, Kulvicius T, Tamosiunaite M, Mecklenbrauck F, et al. Touching
913   events predict human action segmentation in brain and behavior. Neuroimage.
914   2021;243:118534. doi: 10.1016/j.neuroimage.2021.118534.

915   43.   Lingnau A, Downing PE. The lateral occipitotemporal cortex in action. Trends Cogn Sci.
916   2015;19(5):268–77. doi: 10.1016/j.tics.2015.03.006.

917   44.   Creem-Regehr SH. Sensory-motor and cognitive functions of the human posterior parietal
918   cortex involved in manual actions. Neurobiol Learn Mem. 2009;91(2):166–71. doi:
919   10.1016/j.nlm.2008.10.004.

920   45.   Schubotz RI, Wurm MF, Wittmann MK, von Cramon DY. Objects tell us what action we can
921   expect: Dissociating brain areas for retrieval and exploitation of action knowledge during
922   action observation in fMRI. Front Psychol. 2014;5(JUN):1–15. doi: 10.3389/fpsyg.2014.00636.

923   46.   Reber PJ, Gitelman DR, Parrish TB, Mesulam MM. Priming Effects in the Fusiform Gyrus:
924   Changes in Neural Activity beyond the Second Presentation. Cereb Cortex. 2005;15(6):787–
925   95. doi: 10.1093/cercor/bhh179.

926   47.   Fiebach CJ, Schubotz RI. Dynamic anticipatory processing of hierarchical sequential events: A
927   common role for Broca's area and ventral premotor cortex across domains? Cortex.
928   2006;42(4):499–502. doi: 10.1016/S0010-9452(08)70386-1.

929   48.   Clerget E, Winderickx A, Fadiga L, Olivier E. Role of Broca's area in encoding sequential human
930   actions: A   virtual   lesion   study.   Neuroreport.   2009;20(16):1496–9.   doi:
931   10.1097/WNR.0b013e3283329be8.

49. Badre D, Wagner AD. Left ventrolateral prefrontal cortex and the cognitive control of
memory.   Neuropsychologia.   2007;45(13):2883–901.   doi:
10.1016/j.neuropsychologia.2007.06.015.

50. Heim S, Eickhoff SB, Amunts K. Different roles of cytoarchitectonic BA 44 and BA 45 in
phonological and semantic verbal fluency as revealed by dynamic causal modelling.
Neuroimage. 2009;48(3):616–24. doi: 10.1016/j.neuroimage.2009.06.044.

51. Oldfield RC. The assessment and analysis of handedness: The Edinburgh inventory.
Neuropsychologia. 1971;9(1):97–113. doi: 10.1016/0028-3932(71)90067-4.

52. Cheng S, Werning M, Suddendorf T. Dissociating memory traces and scenario construction in
mental   time   travel.   Neurosci   Biobehav   Rev.   2016;60:82–9.   doi:
10.1016/j.neubiorev.2015.11.011.

53. Franz EA, Ford S, Werner S. Brain and cognitive processes of imitation in bimanual situations:
Making inferences about mirror neuron systems. Brain Res. 2007;1145:138–49. doi:
10.1016/j.brainres.2007.01.136.

54. Wurm MF, Schubotz RI. The role of the temporoparietal junction (TPJ) in action observation:
Agent detection rather than visuospatial transformation. Neuroimage. 2018;165(2017):48–55.
doi: 10.1016/j.neuroimage.2017.09.064.

55. El-Sourani N, Trempler I, Wurm MF, Fink GR, Schubotz RI. Predictive Impact of Contextual
Objects during Action Observation: Evidence from Functional Magnetic Resonance Imaging. J
Cogn Neurosci. 2019;32:326–337. doi: 10.1162/jocn_a_01480.

56. R Core Team. R: A language and environment for statistical computing. R Foundation for
Statistical Computing, Vienna, Austria. URL https://www.R-project.org/. 2020.

57. Snodgrass JG, Corwin J. Pragmatics of measuring recognition memory: Applications to
dementia and amnesia. J Exp Psychol Gen. 1988;117(1):34–50. doi: 10.1037/0096-
3445.117.1.34.

58. Collins AM, Quillian MR. Retrieval Time from Semantic Memory. J Verbal Learning Verbal
Behav. 1969;8(2):240–7. doi: 10.1016/S0022-5371(69)80069-1.

59. Larsen SF, Plunkett K. Remembering experienced and reported events. Appl Cogn Psychol.
1987;1(1):15–26. doi: 10.1002/acp.2350010104.

60. Noppeney U, Price CJ. Retrieval of abstract semantics. Neuroimage. 2004;22(1):164–70. doi:
10.1016/j.neuroimage.2003.12.010.

61. Wobbrock JO, Findlater L, Gergle D, Higgins JJ. The Aligned Rank Transform for Nonparametric
Factorial Analyses Using Only ANOVA Procedures. CHI '11 CHI Conf Hum Factors Comput Syst.
2011;143–6. doi: 10.1145/1978942.1978963.

62. Bonferroni CE. Teoria Statistica Delle Classi e Calcolo Delle Probabilità, Vol. 8. Pubblicazioni del
R Istituto Superiore di Scienze Economiche e Commerciali di Firenze, 3–62. 1936.

63. Friston KJ, Holmes AP, Worsley KJ, Poline J -P, Frith CD, Frackowiak RSJ. Statistical Parametric
Maps in Functional Imaging: A General Linear Approach. Hum Brain Mapp. 1994;2(4):189–
210. doi: 10.1002/hbm.460020402.

64. Worsley KJ, Friston KJ. Analysis of fMRI Time-Series Revesited - Again. Neuroimage.
1995;2(3):173–81. doi: 10.1006/nimg.1995.1023.

65. Nichols T, Brett M, Andersson J, Wager T, Poline JB. Valid conjunction inference with the
minimum statistic. Neuroimage. 2005;25(3):653–60. doi: 10.1016/j.neuroimage.2004.12.005.

66. Worsley KJ, Marrett S, Neelin P, Vandal AC, Friston KJ, Evans AC. A unified statistical approach
for determining significant signals in images of cerebral activation. Hum Brain Mapp.
1996;4(1):58–73. doi: 10.1002/(SICI)1097-0193(1996)4:1<58::AID-HBM4>3.0.CO;2-O.

67. Farrer C, Frith CD. Experiencing Oneself vs Another Person as Being the Cause of an Action:
The Neural Correlates of the Experience of Agency. Neuroimage. 2002;15(3):596–603. doi:
10.1006/nimg.2001.1009.

68. Mechelli A, Sartori G, Orlandi P, Price CJ. Semantic relevance explains category effects in
medial   fusiform   gyri.   Neuroimage.   2006;30(3):992–1002.   doi:
10.1016/j.neuroimage.2005.10.017.

69. Kumaran D, Maguire EA. Match–Mismatch Processes Underlie Human Hippocampal
Responses   to   Associative   Novelty.   J   Neurosci.   2007;27(32):8517–24.   doi:
10.1523/JNEUROSCI.1677-07.2007.

70. Gorgolewski KJ, Varoquaux G, Rivera G, Schwarz Y, Ghosh SS, Maumet C, et al.
NeuroVault.Org: A web-based repository for collecting and sharing unthresholded statistical
maps of the human brain. Front Neuroinform. 2015;9(8):1–9. doi: 10.3389/fninf.2015.00008.

71. Maldjian JA, Laurienti PJ, Kraft RA, Burdette JH. An automated method for neuroanatomic and
cytoarchitectonic atlas-based interrogation of fMRI data sets. Neuroimage. 2003;19(3):1233–
9. doi: 10.1016/S1053-8119(03)00169-1.

72. Brett M, Anton JL, Valabregue R, Poline J-B. Region of interest analysis using an SPM toolbox.

Present 8th Int Conferance Funct Mapp Hum Brain, June 2-6, 2002, Sendai, Japan.

73.   St. Jacques PL, Olm C, Schacter DL. Neural mechanisms of reactivation-Induced updating that
enhance and distort memory. Proc Natl Acad Sci U S A. 2013;110(49):19671–8. doi:
10.1073/pnas.1319630110.

74.   Sederberg PB, Gershman SJ, Polyn SM, Norman KA. Human memory reconsolidation can be
explained using the temporal context model. Psychon Bull Rev. 2011;18(3):455–68. doi:
10.3758/s13423-011-0086-9.

75.   Klingmüller A, Caplan JB, Sommer T. Intrusions in episodic memory: reconsolidation or
interference? Learn Mem. 2017;24(5):216–24. doi: 10.1101/lm.045047.117.

76.   Greve A, Cooper E, Tibon R, Henson RN. Knowledge Is Power: Prior Knowledge Aids Memory
for Both Congruent and Incongruent Events, but in Different Ways. J Exp Psychol Gen.
2019;148(2):325–41. doi: 10.1037/xge0000498.

77.   Ritvo VJH, Turk-Browne NB, Norman KA. Nonmonotonic Plasticity: How Memory Retrieval
Drives Learning. Trends Cogn Sci. 2019;23(9):726–42. doi: 10.1016/j.tics.2019.06.007.

78.   Wurm MF, Schubotz RI. Squeezing lemons in the bathroom: Contextual information
modulates action recognition. Neuroimage. 2012;59(2):1551–9. doi:
10.1016/j.neuroimage.2011.08.038.

79.   Hrkać M, Wurm MF, Kühn AB, Schubotz RI. Objects Mediate Goal Integration in Ventrolateral
Prefrontal Cortex during Action Observation. PLoS One. 2015;10(7):1–12. doi:
10.1371/journal.pone.0134316.

80.   Kurby CA, Zacks JM. Segmentation in the perception and memory of events. Trends Cogn Sci.
2008;12(2):72–9. doi: 10.1016/j.tics.2007.11.004.

81.   Grill-Spector K, Kourtzi Z, Kanwisher N. The lateral occipital complex and its role in object
recognition. Vision Res. 2001;41(10–11):1409–22. doi: 10.1016/S0042-6989(01)00073-6.

82.   Wiggett AJ, Downing PE. Representation of Action in Occipito-temporal Cortex. J Cogn
Neurosci. 2011;23(7):1765–80. doi: 10.1162/jocn.2010.21552.

83.   Mruczek REB, von Loga IS, Kastner S. The representation of tool and non-tool object
information in the human intraparietal sulcus. J Neurophysiol. 2013;109(12):2883–96. doi:
10.1152/jn.00658.2012.

84.   Aguirre GK, Detre JA, Alsop DC, D'Esposito M. The Parahippocampus Subserves Topographical
Learning in Man. Cereb Cortex. 1996;6(6):823–9. doi: 10.1093/cercor/6.6.823.

85. O'Reilly RC, Rudy JW. Computational Principles of Learning in the Neocortex and
Hippocampus. Hippocampus. 2000;10(4):389–97. doi: 10.1002/1098-
1063(2000)10:4<389::AID-HIPO5>3.0.CO;2-P.

86. Davachi L, Wagner AD. Hippocampal Contributions to Episodic Encoding: Insights From
Relational and Item-Based Learning. J Neurophysiol. 2002;88(2):982–90. doi:
10.1152/jn.2002.88.2.982.

87. Köhler S, Crane J, Milner B. Differential Contributions of the Parahippocampal Place Area and
the Anterior Hippocampus to Human Memory for Scenes. Hippocampus. 2002;12(6):718–23.
doi: 10.1002/hipo.10077.

88. Kuhl BA, Bainbridge WA, Chun MM. Neural Reactivation Reveals Mechanisms for Updating
Memory. J Neurosci. 2012;32(10):3453–61. doi: 10.1523/JNEUROSCI.5846-11.2012.

89. Duncan K, Curtis C, Davachi L. Distinct Memory Signatures in the Hippocampus: Intentional
States Distinguish Match and Mismatch Enhancement Signals. J Neurosci. 2009;29(1):131–9.
doi: 10.1523/JNEUROSCI.2998-08.2009.

90. Klein TA, Ullsperger M, Danielmeier C. Error awareness and the insula: links to neurological
and psychiatric diseases. Front Hum Neurosci. 2013;7(JAN):1–14. doi:
10.3389/fnhum.2013.00014.

91. Menon V, Uddin LQ. Saliency, switching, attention and control: a network model of insula
function. Brain Struct Funct. 2010;214(5–6):655–67. doi: 10.1007/s00429-010-0262-0.

92. Summerfield C, Trittschuh EH, Monti JM, Mesulam MM, Egner T. Neural repetition
suppression reflects fulfilled perceptual expectations. Nat Neurosci. 2008;11(9):1004–6. doi:
10.1038/nn.2163.

93. Nyberg L, Habib R, Herlitz A. Brain activation during episodic memory retrieval: Sex
differences. Acta Psychol (Amst). 2000;105(2–3):181–94. doi: 10.1016/s0001-6918(00)00060-
3.

94. Antony JW, Ferreira CS, Norman KA, Wimber M. Retrieval as a Fast Route to Memory
Consolidation. Trends Cogn Sci. 2017;21(8):573–6. doi: 10.1016/j.tics.2017.05.001.

Supporting information

S1 Fig. Whole brain activation for attenuation effects. (A) Uncorrected t-map ($p < 0.001$) for

structure violation attenuation effect. (B) FDR-corrected t-map ($p < 0.05$) for content expectancy

violation attenuation effect. IFJ: inferior frontal junction; IFS: inferior frontal sulcus; IPS: intraparietal

sulcus; FG: fusiform gyrus; PCUN: precuneus; THA: thalamus; PHG: parahippocampal gyrus.

S1 Table. Three examples of the 24 toy stories used in the study.

S2 Table. Peak activations from second-level whole-brain analyses of attenuation effects at the

time of violation.

S3 Table. One-sample t-tests (one-tailed) for mean beta values compared to zero.

A

Camera

Camera frame

Toy object

Actress

B

Third-person-perspective (3pp)

First-person-perspective (1pp)

Rotation

180°

Experimenter

Camera monitor

Experiment monitor

Camera

Toy objects

Camera frame

Participant

A

str > ori

Uncorrected, *p* < 0.001

IFG
SFS
PMd

B

con > ori

FDR-corrected, *p* < 0.05

IFS/IFG

pre-SMA
IPS   IPS
AI   AI
FG   FG
PHG