Title

First evidence of *in vitro* cytotoxic effects of marine microlitter on *Merluccius merluccius* and

Mullus barbatus, two Mediterranean commercial fish species

Authors

Miccoli A*[1], Mancini E[2], Saraceni PR[1], Della Ventura G[3,4], Scapigliati G[1] and Picchietti S[1]

[1] Department for Innovation in Biological, Agro-food and Forest Systems, University of

Tuscia, Viterbo, 01100, Italy

[2] Italian Fishery Research and Studies Center, Rome, 00184, Italy

[3] Department of Science, Roma 3 University, Rome, 00146, Italy

[4] INFN Laboratori Nazionali di Frascati, Via E. Fermi 54, Frascati, 00044, Italy

*Corresponding author, andrea.miccoli@unitus.it

ORCiD

AM: 0000-0002-4545-7229

SP: 0000-0002-6116-213X

GDV: 0000-0001-6277-961X

Abstract

Marine litter is composed mainly of plastics and is recognized as a serious threats to marine

ecosystems. Ecotoxicological approaches have started elucidating the potential severity of

microplastics (MPs) in controlled laboratory studies with pristine materials but no information

exist on marine environmental microlitter as a whole. Here, we characterized the litter in the

coastal Northern Tyrrhenian sea and in the stomach of two fish species of socio-economic

importance, and exposed primary cell cultures of mucosal and lymphoid organs to marine

microlitter for evaluating possible cytotoxic effects. An average of 0.30 ± 0.02 microlitter items

m^{-3} was found in water samples. µFT-IR analysis revealed that plastic particles, namely HDPE,

polyamide and polypropylene were present in 100% and 83.3% of *Merluccius merluccius* and

Mullus barbatus analyzed, which overall ingested 14.67 ± 4.10 and 5.50 ± 1.97

items/individual, respectively. Moreover, microlitter was confirmed as a vector of

microorganisms. Lastly, the apical end-point of viability was found to be significantly reduced

in splenic cells exposed *in vitro* to two microlitter conditions. Considering the role of the spleen

in the mounting of adaptive immune responses, our results warrant more in-depth

investigations for clarifying the actual susceptibility of these two species to anthropogenic

microlitter.

Keywords

Marine microlitter; Bioindicators; Cytotoxicity; *In vitro* approaches; Primary cell cultures;

Biological agents

Highlights

- 0.30 ± 0.02 microlitter items m^{-3} were found at the surface of coastal Northern Tyrrhenian sea
- 14.67 ± 4.10 and 5.50 ± 1.97 items/individual were retrieved from the stomach of hakes and mullets
- The ingested microlitter contained plastic items
- Microlitter was validated as a carrier of bacteria, fungi and flagellates
- Splenic cells exposed to two microlitter conditions for 72 hours suffered cytotoxicity

Figure/Table captions

Fig. 1 Map of sampling sites - Study area with 250 μm net tows GPS positions, haul route

and bathymetry.

Fig. 2 A-E Microlitter collected from sea surface - Some examples of litter particles collected

from surface 250 μm net tows (SUP1-SUP4).

Fig. 3 Quali-quantitative characterization of anthropogenic litter collected from the sea

surface- A) Particle density of the micro- and macrolitter fractions per net tow. Average litter

density is reported as mean ± SE. B) Cumulative microlitter density per item type per net

tow. C) Size class distribution per litter shape type. D-F) Percentage of color abundance per

microlitter shape type.

Fig. 4 Examples of microlitter items retrieved from the digestive tract of fish - A) White

fragment from *M. barbatus*, 0.58 match with polyamide (Primpke et al., 2018). B) Black

filament from *M. merluccius*, 0.46 match with HDPE (Chabuka and Kalivas, 2020). C) Green

fragment from *M. merluccius*, 0.72 match with HDPE (Chabuka and Kalivas, 2020). Scale

bars: 500 μm.

Fig. 5 Quali-quantitative characterization of microlitter retrieved from fish stomach contents -

A) Microlitter particle abundance distribution per species (log y scale). Whiskers plotted

according to the Tukey method. B) Cumulative particle abundance per type. C) Size class

distribution per microlitter shape type per species. D-G) Percentage of color abundance per

microlitter shape type per species.

Fig. 6 μFT-IR spectra - Spectra of randomly selected representative microlitter items

retrieved from fish stomach contents. A) Blue fragment from *M. barbatus*; B) blue filament

from *M. merluccius*; C) white fragment from *M. barbatus*. Matching with reference polymer

as per the Open Specy open source database. r: Pearson's correlation coefficient as

measure of linear correlation between data sets.

Fig. 7 A-D Microlitter as a vector of biological agents - Examples of microorganisms

observed in cell suspensions following a 72-hour incubation with microlitter. A) European

hake splenic cell, negative control. B-D) Grey, white and black arrowheads indicate

bacilliform bacteria, unicellular fungi and flagellates, respectively. Scale bars: 10 µm.

Fig. 8 Quantification of intracellular ATP content as a proxy of cell viability - ATP data

distribution per species, time, organ and treatment. Statistical significance as per one-way

ANOVA followed by Tukey's HSD *post hoc* test. *: $p< 0.05$

Fig. 9 Comparison of microlitter densities retrieved from representative scientific literature

with data herein presented. Boxes indicate mean ± SD items m^{-3}. Sampling locations

differentiated by color-coding.

Table 1 Sampling details - Summary of the experimental campaign with net tows details.

Time expressed as UTC/GMT +2:00.

Table 2 Full statistical details of the rank-based nonparametric Kruskal-Wallis tests

performed on microbial counts per species and organ. ns: non significant.

Table S1 Sea surface microlitter abundance and density (items m^{-3}) per color per type per

net tow and relative share of litter types.

Table S2 Sea surface litter size classes per particle type per net tow.

Table S3 Full statistical details of the one way ANOVA tests performed on cytotoxic data of

A) *Merluccius merluccius* and B) *Mullus barbatus*, per time and organ. DFn: degrees of

freedom in the numerator; DFd: degrees of freedom in the denominator; F: test statistic for

ANOVA; ges: generalized eta squared.

1. Introduction

Coastal areas are subject to an exponential increase in population density and the development of impacting human activities, e.g. industries, tourism, recreational activities, fishing and aquaculture. As a consequence, they can be affected by both sporadic and continuous pollution events, with consequences on all compartments, and are thus considered "hotspots" of contamination (Llorca et al., 2020; Lu et al., 2018).

Marine litter is defined as "any anthropogenic manufactured, or processed solid material (regardless of size) discarded, disposed of, or abandoned in the environment, including all materials discarded into the sea, on the shore, or brought indirectly to the sea by rivers, sewage, storm water, waves, or winds" (UNEP, 2016). It is one of the most serious threats to marine ecosystems and a global environmental concern. It comprises glass, metal, cardboard and textiles items (Löhr et al., 2017) as well as anthropogenic particles produced by industrial activities (e.g., coal-fired power plants) and transport emissions (Piazzolla et al., 2020), but Tekman et al. (2021) revealed that plastic accounts for the 66-79% of the global litter composition.

Annual global plastic production accounted for 368 million tons in 2019 (Association of Plastic Manufacturers, 2020): about 1.3-3.1% of these (5-12 million tons year^{-1}) reach the Oceans (Jambeck et al., 2015), and the total amount of floating plastic was estimated at 0.3 million tons (van Sebille et al., 2015). The Mediterranean Sea, in particular, is regarded as an accumulation zone for marine litter, with densies comparable to those of the five subtropical gyres (Cózar et al., 2015; UNEP/MAP, 2015; Van Sebille et al., 2020).

In addition to primary microplastics (MPs), i.e. particles that are purposefully manufactured of microscopic sizes < 5 mm, the vast majority of marine litter is subject to degradation by abiotic (UV radiation, mechanical abrasion, temperature) and biotic (microbiological depolymerization) agents, resulting into secondary MPs (Ru et al., 2020; Thompson et al., 2004). Their chemico-physical properties, such as type of polymer, density, size, shape, internal geometry and color, influence their transport, buoyancy and sinking as well as rates

of ingestion and removal by aquatic organisms (Kowalski et al., 2016; Nguyen et al., 2020; Shim et al., 2018).

Due to their small size, MPs are bioavailable to a variety of taxa (e.g. Cole et al., 2013; Fossi et al., 2018; Gomiero et al., 2018; Lusher et al., 2013; Pittura et al., 2018) and can either be mistaken with or selectively chosen instead of food (Clark et al., 2016; Moore, 2008), with demonstrated impacts. Once ingested, MPs can affect biological functions and tissue integrity of organisms (Cole et al., 2015; Pedà et al., 2016; Sussarellu et al., 2016). Moreover, MPs can be potential carriers of pollutants (Amelia et al., 2021; Guo and Wang, 2019) and can be colonized by microbial pathogens, transferring them along the trophic web (Caruso, 2019; Casabianca et al., 2019). Ecotoxicological and physiological impacts of MPs were also demonstrated in controlled laboratory studies, but the commercially-available and pristine materials employed hardly reflect the actual heterogeneity of the environmental litter.

The European hake *Merluccius merluccius* (Linnaeus, 1758) and the red mullet *Mullus barbatus* (Linnaeus, 1758) are good experimental models in MPs-related research because of their biological features, commercial relevance, abundance in the Mediterranean region and suitability as small-scale plastic pollution bioindicators. Building on research that demonstrated the presence of MPs in their gastrointestinal tracts (Atamanalp et al., 2021; Avio et al., 2020, 2015; Bellas et al., 2016; Digka et al., 2018; Giani et al., 2019; Mancuso et al., 2019), in the present study we hypothesized that microlitter could be retrieved from the marine environment and ingested by fish, and that it could affect the *in vitro* viability of immunologically-relevant cells. We therefore investigated the litter abundance in the coastal sector of Civitavecchia (Northern Tyrrhenian Sea, Latium, Italy) at the sea surface and in the stomachs of *M. merluccius* and *M. barbatus*. We also evaluated the microbiological contamination and the *in vitro* cytotoxicity of environmentally-collected microlitter particles on primary cell cultures of mucosal and lymphoid organs, with the ultimate aim of defining the actual susceptibility of these two socio-economically important fish species to anthropogenic microlitter.

2. Material & Methods

a. Study area

The research was conducted in the northern Tyrrhenian Sea (Italy), FAO's General Fisheries Commission for the Mediterranean (GFCM) Geographical Sub-Area 9 (GSA 9). The experimental campaign fell within the physiographical unit (PU) M. Argentario – Cape Linaro, and included the coastal platform that extends from Santa Severa (42.01676 N, 11.95604 E) to the Tarquinia coastal area (42.22243 N, 11.70495 E). The study area is characterized by a 120-150 m deep continental margin and a sandy to sandy-muddy seabed, as reported in Mancini et al. (2021).

b. Experimental campaign

The experimental campaign took place on October 23rd 2020 and extended over a 14-hour period (03:00 - 17:00 UTC/GMT +2:00). Four 15-minute long horizontal tows with a 250 μm mesh size net were performed at sea surface for litter sampling (Fig. 1). Table 1 reports time of start, GPS position, micro- and macrolitter abundance m^{-3} and meteo-marine conditions as per the World Meteorological Organization sea state coding.

Start time (sample #)	Start coordinates	End coordinates	Microlitter items m^{-3} (>0.25, <5 mm)	Macrolitter items m^{-3} (>0.5, < 6 cm)	Meteo-marine conditions
07:35 (SUP1)	42°10.883N 11°30.070E	42°10.570N 11°30.470E	0.29	0.008	Calm (rippled)
11:15 (SUP2)	42°06.349N 11°37.538E	42°05.789N 11°38.128E	0.23	0.01	Smooth (wavelets)
14:05 (SUP3)	42°02.957N 11°41.231E	42°02.4230N 11°41.919E	0.32	0.013	Smooth (wavelets)
16:45 (SUP4)	42°03.554N 11°43.158E	42°03.522N 11°43.884E	0.35	0.023	Sligth

c. Qualitative and quantitative characterization of litter at the sea surface

The 250 μm WP2 net was equipped with a metered line, a non-filtering cod-end and a flow

meter. Upon deployment from the starboard side of the vessel, utmost care was taken to

ensure that the net stretched out correctly. Upon retrieval, the content of the cod-end was

transferred to 500 mL containers and stored at 4 °C until transported to the laboratory. To

prevent airborne contamination in the laboratory, microlitter exposure to air was kept as short

as possible and all following steps were performed under a sterile laminar flow cabinet

previously cleaned with 100% ethanol. Each sample (SUP1-SUP4) was visually sorted 5 mL

at a time using a stereomicroscope (Leica 8APO). Microlitter was classified in terms of shape

(i.e. filament, fragment, film), color and diameter size (A: $250 < x < 500$ μm; B: $500 < x < 1000$

μm; C: $1000 < x < 3000$ μm; D: $3000 < x < 5000$ μm; E: $x > 5000$ μm). Size fractionation, as a

method for representing litter size, was used in other recent peer-reviewed publications (Digka

et al., 2018; Giani et al., 2019). Following the processing in the lab, microlitter sampled from

the marine environment was stored in sterile 50 mL Falcon tubes filled with 0.22 μm-filtered

sea water. Microlitter abundances are presented as items m^{-3} of seawater ± standard error

(SE).

d. Fish sampling and characterization of litter quantity and quality in

184   **stomach contents**

Fish specimens were collected in the same area (October 30[th] 2020, 01:11:00 PM haul start

time, haul start GPS position 41°59.669 N 11°49.214 E, haul end GPS position 42°03.0410 N

11°44.4621 E) by means of the bottom trawl net typically known as "volantina" geared with a

cod-end mesh size of 50 mm diamond and a vertical opening of 4 m (Sala et al., 2013). *M.*

merluccius and *M. barbatus* specimens had a total length (TL) of 18.6 ± 1.02 cm and 11.7 ±

0.64 and a body weight (BW) of 47.1 ± 8.7 g and 16.5 ± 1.15 g, respectively (average ± SD):

they were employed for the quali-quantitative characterization of microlitter within the stomach

contents (n=6 per species) and for cytotoxicity assays (n=3 per species). Immediately

following the opening of the net, specimens of similar within-species TL were sorted,

thoroughly rinsed with 0.22 µm-filtered Milli-Q water, wrapped in autoclaved aluminum foil and

immediately transferred to ice, where they were kept until lab processing. Stomach eversion

was not observed during sampling.

In the laboratory, fish stomachs were excised under a sterile laminar flow cabinet, previously

cleaned with 100% ethanol, and preserved in 75% ethanol until processing. Tools for organ

dissection (e.g. stainless steel tweezers and forceps, glass Petri dishes) were sterilized by

autoclave. Stomach contents were then placed in sterile Petri dishes and visually sorted with

a stereomicroscope (Leica 8APO) to classify litter particles in terms of shape, diameter size

and color, using the same classification as in section 2c. The particles retrieved from the

stomach contents were maintained in 50 mL falcon tubes with 0.22 µm-filtered sea water.

Microlitter abundances are presented as mean particles per species ± standard error (items ±

SE). Our workflow was compliant with the recommendations of Bessa et al. (2019) with

regards to species selection, sampling methods, airborne/external cross-contamination

prevention and particle classification.

e. µFT-IR analysis

To ascertain the nature of the microlitter items retrieved from fish stomach contents, forty items

were analyzed by Fourier-transform infrared microspectroscopy (µFT-IR) following visual

inspection. The subsample was representative of shape types, color and size ranges of items

both within- and between-species. The experiments were performed at the DAFNE Laboratory

of INFN (Frascati, Italy) in transmission mode, using a Bruker Hyperion 3000 FTIR microscope

equipped with a Globar IR source, a broadband beamsplitter (KBr) and a mercury-cadmium

telluride (MCT) detector; the beam size was set at 20x20 µm. Spectroscopic analysis yielded

absorbance spectra, which were analyzed using the Open Specy open source database

(Cowger et al., 2021) with the Pearson's correlation coefficient as measure of the linear

correlation between the data sets. Spectra visualization and overlay were achieved with the

SpectraGryph 1.2 software using the peak normalization method (i.e. each spectrum highest

peak within the visible area was set to 1).

f. Assessment of microlitter cytotoxic effects on fish primary cell cultures

Marine microlitter particles were obtained by surface 250 µm net tows from the same sampling

location of fish, and were stored in 50 mL falcon tubes filled with 0.22 µm-filtered sea water

following visual inspection until employed for cytotoxicity assays. Microlitter particles were

dried under laminar flow cabinet on a lint-free tissue (Kimwipes, Kimtech Science, USA) in a

sterile glass Petri dish at room temperature and there counted using a stereo microscope

(average item weight 160 µg, size ranging from 250 and 5000 µm, colors observed red, blue,

black, green, grey). Isolation and cultivation of fish primary cells were performed according to

published standard procedure (e.g. Miccoli et al., 2021b). Gills (G), head kidney (HK) and

spleen (SPL) were dissected from *M. merluccius* and *M. barbatus* (n=3) under a sterile laminar

flow cabinet previously cleaned with 100% ethanol with sterilized tools (see section 2d), and

immediately immersed in cold Hank's Balanced Salt Solution without calcium and magnesium

(HBSS), previously adjusted for appropriate sea water osmolarity (355 mOsm Kg^{-1}) with 3M

NaCl. Cells were obtained through teasing in cold HBSS using disposable strainers of 100 µm

and 40 µm mesh size, and homogenates were washed by centrifugation (10 min, 400 g, 4 °C).

Cells were then resuspended in L-15 (Leibovitz) medium containing 10% heat-inactivated fetal

calf serum (FCS, Gibco) and antibiotics (penicillin–streptomycin, Gibco). Cells were counted

in a Neubauer chamber and adjusted to a concentration of $5x10^5$ cells mL^{-1} in L-15 medium.

Both HBSS and L-15 media were 0.22 µm-filtered.

Cellular suspensions were exposed to "Low" and "High" microlitter conditions corresponding

to 4 and 20 field-collected microlitter particles/ml. Both microlitter concentrations were

consistent number-wise with the ingestion rates of microlitter herein described, and the lowest

microlitter concentration was in line with future modelled estimates weight-wise (Isobe et al.,

2019).

Cells were cultured for 2 and 72 hours at 15 °C with gentle rotary shaking to ensure a

continuous contact with microlitter particles. Because the bottom trawl net was not equipped

with any temperature sensor, the incubation temperature was selected according to the near

real-time numerical model MEDSEA_ANALYSISFORECAST_PHY_006_013 (Clementi et al.,

2021), resolving for variable "sea_water_potential_temperature_at_sea_floor (bottomT)"

using sampling location, date and depth as input. Three technical replicates per biological

sample were used in all experimental groups. Negative controls consisting of cells incubated

at the same conditions without microlitter were considered for each organ and species.

Positive controls, i.e. cells incubated as negative controls but with 0.2% NaN_3, were also tested

separately for each organ and species. Intracellular ATP value, as a proxy of cell

viability/cytotoxicity (Schoonen et al., 2005), was then quantitatively evaluated simultaneously

in all samples of both species for any given incubation time using the ATPlite assay

(PerkinElmer, catalog no. 6016943) following the manufacturer's instructions: 50 µL of cell

lysis and 50 µL substrate solutions were added to 100 µL cell suspensions per replicate and

shaken for 5 min. Resulting homogenates were transferred to disposable opaque well plates

(OptiPlate-96, PerkinElmer) and luminescence was measured using a microplate reader

(Wallac Victor2, PerkinElmer), following a 10-minute dark adaptation period.

262   **g. Validation of microlitter as a carrier of biological agents**

At the end of a 72-hour incubation of primary cultures with microlitter following an alike

experimental design as above, 10 µL of cell suspension from each experimental group were

qualitatively observed under a Zeiss microscope equipped with a colour 8 camera (AxioCam

MRC) and a software package (KS 300 and AxioVision). Multiple sets of photographs at

random frames were taken per each experimental group and microorganisms were quantified

over a 100.000 μm^2 area by an operator unaware of treatments. Sterile glass slides and cover

slips were used for these steps.

270   **h. Data analysis, visualization and statistics**

Stomach content particle abundance was tested for statistical significance between species

using an independent samples t-test with the null hypothesis of equal population means

between groups. Datasets were checked for normality with the Shapiro-Wilk test and for

homoscedasticity with the Levene test. A log-transformation was applied to meet the normal

distribution assumption.

The relation among species and ingestion of microlitter particle types by color was examined

with a chi-square test on a two-way contingency table. The null hypothesis assumed no

association between variables. Results are reported as χ^2_{df}=test statistic.

Cytotoxicity data, grouped by species, time, organ and treatment, were tested for statistical

significance using a one-way ANOVA with the null hypothesis of equal population means

among groups, followed by a Tukey's HSD *post-hoc* test in case the main effect of the models

was significant. Datasets were checked for normality with the Shapiro-Wilk test and for

homoscedasticity with the Levene test.

Microbial count was analyzed with the rank-based nonparametric Kruskal-Wallis test because

datasets, many of which were zero-inflated, did not meet the assumptions for parametric

testing. The null hypothesis was that samples were drawn from the same population or from

populations whose medians did not differ.

A comparison of microlitter densities retrieved from representative scientific literature was

visualized as mean ± SD items m^{-3} with the R "forestplot" package v 2.0.1. Data reported in

other units than items m^{-3} were excluded from the analysis. References were organized

hierarchically by location and year, and box size was set to constant.

i. Ethics statement

Animal manipulation complied with the guidelines of the European Union Directive

(2010/63/EU) and the Italian Legislative Decree 26 of 4 March 2014 "Attuazione della Direttiva

2010/63/UE sulla protezione degli animali utilizzati a fini scientifici". Ethical review and

approval was not required because animals were sampled from the natural environment and

were not subject to any experimental manipulation, in line with the Explanatory Note of the

Italian Ministry of Health's Directorate-General for Animal Health and Veterinary Medicinal

Products (DGSAF) of 26 July 2017.

3. Results

a. Qualitative and quantitative characterization of litter at the sea surface

Anthropogenic marine litter visually of plastic origin and in the form of filaments, fragments and films was found in all water samples taken in the Civitavecchia area (Fig. 2, Table 1). Both microlitter (250 µm < x < 5 mm) and macrolitter (5 mm < x < 6 cm) fractions were identified.

SUP4 and SUP2 had the highest and lowest microlitter particle densities with 0.35 and 0.23

items m^{-3}, respectively (Table 1). SUP4 and SUP 1 had the highest and lowest macrolitter

particle densities with 0.023 and 0.008 items m^{-3}, respectively (Table 1). Their average

abundance among all samples was 0.30 ± 0.02 items m^{-3} and 0.014 ± 0.003 items m^{-3} (Fig.

3A).

With regard to microlitter shapes, filaments, fragments and films were observed. In particular,

the highest and lowest concentration of filaments were found in SUP4 (0.12 items m^{-3}) and

SUP3 (0.02 items m^{-3}), respectively; the highest and lowest concentration of fragments were

found in SUP3 (0.17 items m^{-3}) and SUP1 (0.039 items m^{-3}), and the highest and lowest

concentration of films was found in SUP4 (0.18 items m^{-3}) and SUP2 (0.07 items m^{-3}) (Fig.

3B). Number and density of particles per type per color per station, total number of particles

collected and relative share of litter types are reported in Table S1.

The vast majority (51%) of filaments belonged to size class C and a smaller population to size

class A (5%). Fragments fell for the most part within size class B (38%), while size class E

was the least represented (3%). Films mostly belonged to size classes B and C (39%), while

size class E was found in the 1% of cases (Fig. 3C and Table S2).

Filaments from all tows were mostly white (38%) and black (37%); less frequent colors in terms

of abundance were blue (23%) and yellow (1%). Fragments were more chromatically

diversified: while most of them were white (34%), blue and green fragments were found with

a percentage of 24% and 21%, respectively; red (7%), black (3%) and yellow (2%) fragments

were less abundant. Films were mostly blue (60%), followed by white (17%) and green (15%);

yellow (2%) and red (2%) films were less frequent (Fig. 3D-F and Table S1).

b. Characterization of quantity, quality and chemical composition of

microlitter in fish stomach content

Microlitter items were found in 100% *M. merluccius* specimens and in 5 out of 6 (83.3%) *M.*

barbatus specimens (Fig. 4).

A higher abundance of microlitter was found in the stomach contents of hakes (14.67 ± 4.10

items/individual) than mullets (5.50 ± 1.97 items/individual), but the difference between group

means was not statistically significant (t_{10}=2.045, p=0.068) (Fig. 5A).

The retrieved microlitter was classified as filaments and fragments, and no films were found.

12.67 ± 4.27 filaments and 2 ± 0.71 fragments were retrieved from hakes, while 4.33 ± 2

filaments and 1.17 ± 0.37 fragments were found in mullets. Filaments were the most abundant

shape type, with 76/88 particles (83.36%) in hake and 22/28 (78.57%) in mullet (Fig. 5B).

M. merluccius ingested mostly filaments and fragments in the B and A size classes,

respectively. The most represented filament size class in *M. barbatus* stomach contents was

B, while fragments equally fell into the three size classes (Fig. 5C).

Filaments found in the stomach contents of *M. merluccius* were generally blue (53.95%) and

black (18.42%) followed by red, green, white and other colored types (all below 12%). Blue,

black and green filaments were found in similar percentages also in *M. barbatus* (54.55%,

18.18% and 4.55%, respectively). Microlitter fragments in hake and mullet were mostly white

(33% and 66.67%, respectively) and red (25% and 16.67%, respectively); however, green,

blue and other colored-fragments were retrieved only from hake, while black fragments were

only found in mullet (Fig. 5D-G). Differences in microlitter stomach content by color between

species was not statistically significant either for filaments ($X^2_{(5)}$=9.38, p=0.094) or fragments

($X^2_{(5)}$=5.63, p=0.34).

Forty items visually classified as plastics, equaling approximately the 35% of all items

retrieved, were further analyzed by µFT-IR for a qualitative term of reference. Samples for

which a spectrum could be obtained matched exclusively to synthetic polymers: filaments

exclusively matched with HDPE while fragments were identified as HDPE, polyamide and

polypropylene. HDPE and polypropylene, with a cumulative identification rate of 88.8%, were

the two most frequent polymers. Selected spectra obtained from a blue fragment from *M.*

barbatus, a blue filament from *M. merluccius* and a white fragment from *M. barbatus* are

presented, showing a Pearson's correlation coefficient of 0.87, 0.72 and 0.58 with reference

spectra, respectively (Fig. 6 A-C).

c. Microlitter as carrier of biological agents to _M. merluccius_ and _M. barbatus_ cells

To evaluate the presence of microorganisms and consequently validate the role of microlitter as carrier of biological agents, 10 µL cell suspension from all cultures following a 72-hour incubation were qualitatively assessed by optical microscopy. Bacilliform bacteria (grey arrowheads), unicellular fungi (white arrowheads) and flagellates (black arrowheads) were observed in microlitter-exposed cells (Fig. 7 B-D) but not in control cells (Fig. 7 A).

375 Their abundance was quantified over a 37.000 μm^2 area and normalized to 100.000 μm^2 area

376 per species, organ and microlitter concentration. A variable degree of biological contamination

377 was found in conditioned primary cultures of both species (Table 2).

| Species | Organ | Mean microbial count | | | SE microbial count | | | 95% CI microbial count [L;U] | | Statistical test | | |
		ctrl	low	high	ctrl	low	high	ctrl	low	high	H	p	
M. merluccius	Gills	0	269	241.2	0	143.8	57.95	0;0	-349.8;887.8	-8.14;490.5	5.609	0.068	ns
	Head kidney	0	14.4	23.31	0	7.00	3.91	0;0	-15.79;44.48	6.49;40.13	6.161	0.025	*
	Spleen	0	8.07	48.42	0	5.6	34.97	0;0	-16.02;32.16	-102.0;198.9	5.162	0.1	ns
M. barbatus	Gills	0	0	18.83	0	0	4.11	0;0	0;0	1.15;36.51	7.624	0.036	*
	Head kidney	0	0	17.04	0	0	5.88	0;0	0;0	-8.26;42.34	7.624	0.036	*
	Spleen	0	5.38	23.31	0	5.38	13.92	0;0	-17.77;28.53	-36.58;83.21	4.587	0.107	ns

For the hake, a significant effect of microlitter concentration on microbial counts was found

only in HK primary cultures (H statistics=6.16, p=0.025). For the mullet, statistical significance

was evidenced for microbial counts in G and HK (H statistics=7.62, p=0.036).

d. Microlitter cytotoxicity in *M. merluccius* and *M. barbatus* cells

Microlitter cytotoxicity was evaluated based on the viability of cells from G, HK and SPL

following a 2- and a 72-hour long incubation with two concentrations of microlitter sampled by

250 μm net tows (Fig. 8).

Exposure to microlitter did not induce any statistically significant decrease in intracellular ATP

following the short incubation time in either species or organs (Table S3a-b).

In *M. merluccius*, microlitter induced a decrease in median cell viability after a 72-hour

incubation in primary G cultures at the high concentration and in both HK and SPL cultures at

the low concentration. The linear model fit to splenic cell culture data following the 72-hour

exposure explained the 62.3% of ATP levels variation, even though differences among

experimental groups were slightly non-significant ($F_{(2,6)}$ = 4.96, p=0.054) (Table S3a).

In *M. barbatus* primary cultures from all organs, the median intracellular ATP levels were lower

in the Low and High groups than in corresponding controls. Such a decrease revealed a

statistically significant main effect of microlitter concentration on splenic cells ($F_{(2,6)}$ = 10.8, p

=0.01), with the overall treatment effect explaining almost 80% of ATP levels variation of the

model (η^2 = 0.783) (Table S3b). Pairwise comparisons between the control and the "Low" and

"High" groups were statistically significant (p adjusted = 0.0113 and 0.027, respectively) (Fig.

8).

4. Discussion

In this work we applied a multidisciplinary approach combining oceanographical, spectroscopical, cellular and microscopical methods to characterize the quality and quantity of microlitter particles in the coastal surface waters and in the digestive tract of two commercially-valuable Mediterranean fish species; we also preliminarily addressed the cytotoxic potential of field-collected microlitter on primary cultures of cells extracted from mucosal (i.e. gills) and lymphoid (i.e. head kidney and spleen) organs.

The marine litter causes multiple environmental, economic, social, political and cultural impacts (Barboza et al., 2019; Galgani et al., 2019; GESAMP, 2015; UNEP, 2014), especially to the health and functioning of organisms and ecosystems (Corinaldesi et al., 2021; Garcia-Vazquez et al., 2018; Rios et al., 2007). At the European level, such pollutant was included among the 11 qualitative descriptors of the Marine Strategy Framework Directive against which the quality of the marine environment is assessed (European Parliament, 2008/56/EC). Extensive research has demonstrated the ubiquity of plastic pollution in several matrices such as beaches (Fortibuoni et al., 2021; Prevenios et al., 2018), sediments (Piazzolla et al., 2020; Renzi et al., 2018) and seawater (Atwood et al., 2019; Capriotti et al., 2021) – although remote (Cincinelli et al., 2017; Lusher et al., 2015). Microlitter was retrieved from all water samples taken within the framework of the PISCES project in a much higher (~20-fold) average concentration (0.30 ± 0.02 items m^{-3}) than litter particles > 5 mm (0.014 ± 0.003 items m^{-3}). Keeping in mind the environmental and biological severity of litter < 5 mm, our results are in good agreement with microlitter concentrations reported from other areas of the Mediterranean Sea, Yellow Sea and oceanic waters (Baini et al., 2018; Cincinelli et al., 2017; Collignon et al., 2012; Constant et al., 2018; Cózar et al., 2015; de Lucia et al., 2018, 2014; Expósito et al., 2021; Fagiano et al., 2022; Fossi et al., 2012, 2016; Kazour et al., 2019; Lusher et al., 2015; Panti et al., 2015; Suaria et al., 2016; van der Hal et al., 2017; Zhang et al., 2017; Zhao et al., 2014) (Fig. 9), suggesting that also surveys that are not extensive in either duration or sample sizes can effectively capture the extent of microlitter pollution. This is desirable to

minimize the impacts of research-related anthropogenic activities. An exception was represented by the Eastern Mediterranean Sea, which appears to be much more polluted than the western basin. We must highlight that data dispersion could not be quantified from Cózar et al. (2015) and Constant et al. (2018) as only mean items m^{-3} were reported, and that data from Vasilopoulou et al. (2021) was discarded because of non-informative results (41.31 ± 112.05 mean ± SD items m^{-3} - SD could be back-calculated from standard error because a sample size was clearly indicated by authors).

The relationship between the marine biota and microlitter was so far mostly evaluated by ingestion rates (e.g. Rios-Fuster et al., 2019; Savoca et al., 2019). MPs are thought to be mistaken for or even purposefully chosen instead of food (Clark et al., 2016; Ling et al., 2017) probably also depending on their color (Du et al., 2021; Wright et al., 2013). The presence of both microlitter shape types in the stomachs of *M. merluccius* and *M. barbatus* and the lack of statistically significant differences based on the chromatic factor support the idea that microlitter may be ingested non-selectively by these two species, even though a biomagnification origin cannot be excluded. Hake and mullet were chosen as experimental

models for a variety of reasons: on one hand they are among the most targeted demersal fish
species by the Mediterranean deep-sea fisheries and the two most fished target species in
the shallow area of the coastal sector (Sabatella et al., 2017; Tiralongo et al., 2021); on the
other, they are regarded as bioindicators of coastal marine ecosystems and display a benthic
feeding behavior at a certain life stage (juvenile *M. merluccius*) or throughout the lifespan (*M.*
barbatus) (Carrozzi et al., 2019; Esposito et al., 2014). Moreover, some authors already
described the occurrence of MPs in these two species (Atamanalp et al., 2021; Avio et al.,
2020, 2015; Bellas et al., 2016; Digka et al., 2018; Giani et al., 2019; Mancuso et al., 2019).
Here we confirmed that microlitter ranging in size from 250 to 3000 μm was present in the
stomachs of both our target species in higher numbers compared to literature data (e.g. Avio
et al., 2020; Bellas et al., 2016; Giani et al., 2019), especially for *M. Merluccius*. A reason for
this may be found in an ontogenic trait of the species, for which a shift from benthic towards
pelagic and necto-benthic feeding habits was demonstrated to occur over a TL of 18 cm, and
in the fact that a greater microlitter contamination is usually observed in sediments than water
column. Benthic macrolitter in the area was recently described quali-quantitatively, with the
most abundant categories being ascribed to plastic (Mancini et al., 2021); MPs were also
demonstrated to abound in superficial sediments in the study area (Piazzolla et al., 2020). The
most commonly ingested microlitter items were confirmed by μFT-IR as high density
polyethylene, polyamide and polypropylene (Chabuka and Kalivas, 2020; Primpke et al.,
2018). On this, we must specify that a small number of spectra could be identified with certainty
due to poor signal quality caused by the saturation of the detector (i.e. large sample size), or
significant degradation of the plastic items likely caused by stomach acids. More importantly,
we call the attention on the fact that such synthetic polymers were also demonstrated in marine
sediments (Piazzolla et al., 2020) and in the atmosphere (Lucci et al., 2021) of the same area,
pointing to the high and pervasive dispersion of anthropogenic litter across multiple
environmental compartments.
472

It is known that MPs act as a carrier of biological agents (Amaral-Zettler, 2019; Kiessling et al., 2015), and our data confirmed this (Fig. 7). Because a dedicated experiment aimed at molecular taxonomy could not be set up due to limited availability of microorganisms on microlitter, flagellates were classified on phenotypic properties. Based on flagellar features and because they are extremely common in marine plankton, where they can be found free-swimming or attached to bacterial mats or other surfaces, we hypothesize they may belong to the *Paraphysomonas* or *Spumella* genus or to the aloricate Bicosoecida order. Microbial composition has the ability to condition the fate of MPs in the water column and sediments (Rogers et al., 2020). Once the microlitter is ingested, its associated microorganisms may colonize the gastrointestinal tract of the host, possibly affecting its welfare: in fact, harmful microorganisms, including potential human and animal pathogens, were found associated to litter (Zettler et al., 2013) and, according to data from Zwollo et al. (2021), serious consequences may arise due to the reduced ability to respond adequately to pathogens because of suboptimal humoral immune responses.

Research aimed at also elucidating physiological impacts of MPs have exposed fish to pristine commercially-available particles under controlled laboratory conditions. Their bioavailability was demonstrated and effects such as altered feeding behavior, metabolic disorders, energy depletion, growth impairment, delayed development, compromised immune response, reproduction and lifespan were reported (Botterell et al., 2019; Espinosa et al., 2019, 2017; Guerrera et al., 2021; Mazurais et al., 2015; Rios-Fuster et al., 2021; Sendra et al., 2021; Yong et al., 2020).

Recently, beach-sampled microlitter was employed in *in vivo* experiments on the European sea bass *Dicentrachus labrax* (Zitouni et al., 2021) and medaka *Oryzias latipes* (Pannetier et al., 2020) to investigate survival, development, uptake, oxidative stress and genotoxicity following the administration of a microlitter-spiked feed. Their results showed the ability of environmental MPs to i) accumulate in fish organs, ii) significantly affect the activity of enzymes involved in the antioxidant defense system and iii) induce DNA damages following

acute exposures. HK primary cultures were also employed to explore the impacts of non-environmental MPs on the abundance and antibody response of B cells in rainbow trout (Zwollo et al., 2021): a lower rate of B cell development together with reduced expression of Ig heavy chain genes were found, suggesting that not only innate but also adaptive immunity may be threatened by such an emerging contaminant.

Despite some similarities with the three just-mentioned studies may be perceived, we must highlight that no other research has ever investigated the apical cytotoxicity event in primary cell cultures derived from fish mucosal and lymphoid organs following their exposure to microlitter collected in the same area from where animals were sampled (literature search conducted on Web of Science on December 3 , 2021). The primary organs for examining the cytotoxic effects of microlitter would have been the stomach or intestine, but the methods for cell extraction are difficult, lengthy, prone to contamination and poor in yield, which were all incompatible factors with our experimental design. Keeping in mind that microlitter and MP impacts occur via cell internalization or chemical contamination and that the former mode of action was not expectable due to the large size of particles employed, we investigated additional immunologically-relevant organs that could be reached by the release of chemical contaminants contained by or adsorbed on the microlitter particles, and affected in their physiological status. We believe that our results, obtained in an attempt to bridge the fields of biological oceanography and experimental toxicology, are biologically significant and actual because i) microlitter particles and fish specimens originated from the same sampling site, ii) microlitter cytotoxicity was measured by the well-established, highly-sensitive and unambiguous direct luciferase-based quantification of cellular ATP (Cree and Andreotti, 1997; Mahto et al., 2010) iii) primary cultures were obtained from organs that are key in ensuring immune barrier and competency and iv) the suitability of the strategy for testing for MP toxicity was overall demonstrated and recently reviewed in details (Revel et al., 2021). In addition, fish have been increasingly established as experimental models in the fields of biomedical sciences and toxicology because they share many similarities with higher vertebrates immunology-wise (Miccoli et al., 2021a; Scapigliati et al., 2018).

Taking into account cytotoxicity data (Fig. 8) and the lack of statistically different microbial

counts observed within *M. barbatus* spleen cultures (Table 2), splenic cell subpopulations

appeared to be the most sensitive to microlitter exposures among all investigated organs. No

further reduction in ATP levels were seen in the High compared to the Low condition,

suggesting that such a pollutant can impact cell viability already at concentrations that are in

line with estimates modelled over the next three decades. These results are concerning

because spleen, together with thymus and kidney, is the major lymphoid organ of teleosts in

which adaptive immune responses are mounted (Flajnik, 2018; Zapata et al., 2006). Neither

the physiological endpoints reported in the large majority of scientific literature nor our results

herein presented provide insight into the molecular mechanisms underlying microlitter toxicity

pathways, but inform about apical events manifested either by the whole organism or primary

cell cultures, respectively. However, the novelty of our approach was to provide data on a

lower, possibly more predictive, level of biological organization (cellular vs. organismal) by

means of so-called New Approach Methodologies, which heavily rely on *in vitro* testing. This

is compliant with the 3Rs principle in animal testing, in addition to having been validated by

the latest internationally-agreed test guidelines (OECD, 2021) and supported by regulatory

toxicology roadmaps (e.g. EPA's strategic vision).

5. Conclusions

In conclusion, the present study has investigated the anthropogenic litter in the coastal

epipelagic Northern Tyrrhenian Sea and the stomach of two commercially-relevant fish

species, validated the microlitter fraction as a carrier of biological agents and, for the first time,

demonstrated that splenic cell viability is negatively affected following exposure to such a

contaminant. Future investigations with larger sample sizes, primary or continuous cell

cultures from additional organs and more in-depth methodological approaches are warranted

for clarifying the susceptibility of *Merluccius merluccius* and *Mullus barbatus* to anthropogenic

microlitter.

6. Author contribution

AM: Conceptualization, Funding acquisition, Data curation, Formal analysis, Visualization, Supervision, Project administration, Writing - original draft, Writing - review.

EM: Conceptualization, Funding acquisition, Investigation, Writing - Review & Editing.

PRS: Methodology, Investigation, Writing - original draft, Writing - Review & Editing.

GDV: Methodology, Resources, Writing - Review & Editing.

GS: Resources, Supervision, Writing - Review & Editing.

SP: Supervision, Writing - Review & Editing

7. Acknowledgements

This research was funded by the ''PISCES'' project (FEAMP PO 2014–2020 promoted by Lazio Region, Italy) and the ''Departments of Excellence-2018'' Program (Dipartimenti di Eccellenza) of the Italian Ministry of Education, University and Research awarded to the DIBAF Department of University of Tuscia. AM and EM are extremely thankful to the ''Marinai & Caratisti'' Cooperative of Civitavecchia and to Dr. Roberto Arciprete for their support in all sampling operations. Financial support to GDV was provided by the Grant to Department of Science, Roma Tre University (MIUR-Italy Dipartimenti di Eccellenza, ARTICOLO 1, COMMI 314-337 LEGGE 232/2016). Dr. Daniele Piazzolla of the Laboratory of Experimental Oceanology and Marine Ecology (University of Tuscia) is acknowledged for his contribution to the PISCES project. AM wishes to thank Prof. Pierangelo Luporini (University of Camerino) for his valuable input on flagellate taxonomy.

8. References

Amaral-Zettler, L., 2019. Plastics: Colonization and Degradation, in: Reference Module in Life Sciences. Elsevier. doi:10.1016/B978-0-12-809633-8.90685-X

Amelia, T.S.M., Khalik, W.M.A.W.M., Ong, M.C., Shao, Y.T., Pan, H.-J., Bhubalan, K., 2021. Marine microplastics as vectors of major ocean pollutants and its hazards to the marine ecosystem and humans. Prog. Earth Planet. Sci. 8, 12. doi:10.1186/s40645-020-00405-4

Association of Plastic Manufacturers, 2020. Plastics - the Facts 2020 - An analysis of European

plastics production, demand and waste data.

Atamanalp, M., Köktürk, M., Uçar, A., Duyar, H.A., Özdemir, S., Parlak, V., Esenbuğa, N., Alak, G.,

2021. Microplastics in Tissues (Brain, Gill, Muscle and Gastrointestinal) of Mullus barbatus and

Alosa immaculata. Arch. Environ. Contam. Toxicol. 81, 460–469. doi:10.1007/s00244-021-

00885-5

Atwood, E.C., Falcieri, F.M., Piehl, S., Bochow, M., Matthies, M., Franke, J., Carniel, S., Sclavo, M.,

Laforsch, C., Siegert, F., 2019. Coastal accumulation of microplastic particles emitted from the

Po River, Northern Italy: Comparing remote sensing and hydrodynamic modelling with in situ

sample collections. Mar. Pollut. Bull. 138, 561–574. doi:10.1016/j.marpolbul.2018.11.045

Avio, C.G., Gorbi, S., Regoli, F., 2015. Experimental development of a new protocol for extraction and

characterization of microplastics in fish tissues: First observations in commercial species from

Adriatic Sea. Mar. Environ. Res. 111, 18–26. doi:10.1016/j.marenvres.2015.06.014

Avio, C.G., Pittura, L., D'Errico, G., Abel, S., Amorello, S., Marino, G., Gorbi, S., Regoli, F., 2020.

Distribution and characterization of microplastic particles and textile microfibers in Adriatic food

webs: General insights for biomonitoring strategies. Environ. Pollut. 258, 113766.

doi:10.1016/j.envpol.2019.113766

Baini, M., Fossi, M.C., Galli, M., Caliani, I., Campani, T., Finoia, M.G., Panti, C., 2018. Abundance

and characterization of microplastics in the coastal waters of Tuscany (Italy): The application of

the MSFD monitoring protocol in the Mediterranean Sea. Mar. Pollut. Bull. 133, 543–552.

doi:10.1016/J.MARPOLBUL.2018.06.016

Barboza, L.G.A., Cózar, A., Gimenez, B.C.G., Barros, T.L., Kershaw, P.J., Guilhermino, L., 2019.

Macroplastics Pollution in the Marine Environment, in: Shepperd, C. (Ed.), World Seas: An

Environmental Evaluation. Elsevier, pp. 305–328. doi:10.1016/B978-0-12-805052-1.00019-X

Bellas, J., Martínez-Armental, J., Martínez-Cámara, A., Besada, V., Martínez-Gómez, C., 2016.

Ingestion of microplastics by demersal fish from the Spanish Atlantic and Mediterranean coasts.

Mar. Pollut. Bull. 109, 55–60. doi:10.1016/j.marpolbul.2016.06.026

Bessa, F., Centre, E.S., Lusher, A., 2019. Harmonized protocol for monitoring microplastics in biota.

JPI-Oceans BASEMAN project. doi:10.13140/RG.2.2.28588.72321/1

Botterell, Z.L.R., Beaumont, N., Dorrington, T., Steinke, M., Thompson, R.C., Lindeque, P.K., 2019.

Bioavailability and effects of microplastics on marine zooplankton: A review. Environ. Pollut. 245,

98–110. doi:10.1016/j.envpol.2018.10.065

Capriotti, M., Cocci, P., Bracchetti, L., Cottone, E., Scandiffio, R., Caprioli, G., Sagratini, G., Mosconi,

G., Bovolin, P., Palermo, F.A., 2021. Microplastics and their associated organic pollutants from

the coastal waters of the central Adriatic Sea (Italy): Investigation of adipogenic effects in vitro.

Chemosphere 263, 128090. doi:10.1016/j.chemosphere.2020.128090

Carrozzi, V., Di Lorenzo, M., Massi, D., Titone, A., Ardizzone, G., Colloca, F., 2019. Prey preferences

and ontogenetic diet shift of European hake Merluccius merluccius (Linnaeus, 1758) in the

central Mediterranean Sea. Reg. Stud. Mar. Sci. 25, 100440. doi:10.1016/j.rsma.2018.100440

Caruso, G., 2019. Microplastics as vectors of contaminants. Mar. Pollut. Bull. 146, 921–924.

doi:10.1016/j.marpolbul.2019.07.052

Casabianca, S., Capellacci, S., Giacobbe, M.G., Dell'Aversano, C., Tartaglione, L., Varriale, F.,

Narizzano, R., Risso, F., Moretto, P., Dagnino, A., Bertolotto, R., Barbone, E., Ungaro, N.,

Penna, A., 2019. Plastic-associated harmful microalgal assemblages in marine environment.

Environ. Pollut. 244, 617–626. doi:10.1016/j.envpol.2018.09.110

Chabuka, B.K., Kalivas, J.H., 2020. Application of a Hybrid Fusion Classification Process for

Identification of Microplastics Based on Fourier Transform Infrared Spectroscopy. Appl.

Spectrosc. 74, 1167–1183. doi:10.1177/0003702820923993

Cincinelli, A., Scopetani, C., Chelazzi, D., Lombardini, E., Martellini, T., Katsoyiannis, A., Fossi, M.C.,

Corsolini, S., 2017. Microplastic in the surface waters of the Ross Sea (Antarctica): Occurrence,

distribution and characterization by FTIR. Chemosphere 175, 391–400.

doi:10.1016/j.chemosphere.2017.02.024

Clark, J.R., Cole, M., Lindeque, P.K., Fileman, E., Blackford, J., Lewis, C., Lenton, T.M., Galloway,

T.S., 2016. Marine microplastic debris: a targeted plan for understanding and quantifying

interactions with marine life. Front. Ecol. Environ. 14, 317–324. doi:10.1002/fee.1297

Clementi, E., Aydogdu, A., Goglio, A.C., Pistoia, J., Escudier, R., Drudi, M., Grandi, A., Mariani, A.,

Lyubartsev, V., Lecci, R., Cretí, S., Coppini, G., Masina, S., Pinardi, N., 2021. Mediterranean

Sea Physical Analysis and Forecast (CMEMS MED-Currents, EAS6 system) (Version 1) [Data

set]. Copernicus Monitoring Environment Marine Service (CMEMS).

doi:10.25423/CMCC/MEDSEA_ANALYSISFORECAST_PHY_006_013_EAS6

Cole, M., Lindeque, P., Fileman, E., Halsband, C., Galloway, T.S., 2015. The Impact of Polystyrene

Microplastics on Feeding, Function and Fecundity in the Marine Copepod Calanus

helgolandicus. Environ. Sci. Technol. 49, 1130–1137. doi:10.1021/es504525u

Cole, M., Lindeque, P., Fileman, E., Halsband, C., Goodhead, R., Moger, J., Galloway, T.S., 2013.

Microplastic Ingestion by Zooplankton. Environ. Sci. Technol. 47, 6646–6655.

doi:10.1021/es400663f

Collignon, A., Hecq, J.H., Glagani, F., Voisin, P., Collard, F., Goffart, A., 2012. Neustonic microplastic

and zooplankton in the North Western Mediterranean Sea. Mar. Pollut. Bull.

doi:10.1016/j.marpolbul.2012.01.011

Constant, M., Kerherve, P., Sola, J., Sanchez-Vidal, A., Canals, M., Heussner, S., 2018. Floating

Microplastics in the Northwestern Mediterranean Sea: Temporal and Spatial Heterogeneities, in:

M., C., Di Pace, E., Errico, M., Gentile, G., Montarsolo, A., Mossotti, R. (Eds.), Proceedings of

the International Conference on Microplastic Pollution in the Mediterranean Sea. pp. 9–15.

doi:10.1007/978-3-319-71279-6_2

Corinaldesi, C., Canensi, S., Dell'Anno, A., Tangherlini, M., Di Capua, I., Varrella, S., Willis, T.J.,

Cerrano, C., Danovaro, R., 2021. Multiple impacts of microplastics can threaten marine habitat-

forming species. Commun. Biol. 4, 1–13. doi:10.1038/s42003-021-01961-1

Cowger, W., Steinmetz, Z., Gray, A., Munno, K., Lynch, J., Hapich, H., Primpke, S., De Frond, H.,

Rochman, C., Herodotou, O., 2021. Microplastic Spectral Classification Needs an Open Source

Community: Open Specy to the Rescue! Anal. Chem. 93, 7543–7548.

doi:10.1021/acs.analchem.1c00123

Cózar, A., Sanz-Martín, M., Martí, E., González-Gordillo, J.I., Ubeda, B., Gálvez, J.Á., Irigoien, X.,

Duarte, C.M., 2015. Plastic Accumulation in the Mediterranean Sea. PLoS One 10, e0121762.

doi:10.1371/journal.pone.0121762

Cree, I.A., Andreotti, P.E., 1997. Measurement of cytotoxicity by ATP-based luminescence assay in

primary cell cultures and cell lines. Toxicol. Vitr. 11, 553–556. doi:10.1016/S0887-

2333(97)00060-X

de Lucia, G.A., Caliani, I., Marra, S., Camedda, A., Coppa, S., Alcaro, L., Campani, T., Giannetti, M.,

Coppola, D., Cicero, A.M., Panti, C., Baini, M., Guerranti, C., Marsili, L., Massaro, G., Fossi,

M.C., Matiddi, M., 2014. Amount and distribution of neustonic micro-plastic off the western

674   Sardinian coast (Central-Western Mediterranean Sea). Mar. Environ. Res. 100, 10–16.

675   doi:10.1016/j.marenvres.2014.03.017

676 de Lucia, G.A., Vianello, A., Camedda, A., Vani, D., Tomassetti, P., Coppa, S., Palazzo, L., Amici, M.,

677   Romanelli, G., Zampetti, G., Cicero, A.M., Carpentieri, S., Di Vito, S., Matiddi, M., 2018. Sea

678   water contamination in the Vicinity of the Italian minor islands caused by microplastic pollution.

679   Water (Switzerland) 10. doi:10.3390/w10081108

680 Digka, N., Tsangaris, C., Torre, M., Anastasopoulou, A., Zeri, C., 2018. Microplastics in mussels and

681   fish from the Northern Ionian Sea. Mar. Pollut. Bull. 135, 30–40.

682   doi:10.1016/j.marpolbul.2018.06.063

683 Du, S., Zhu, R., Cai, Y., Xu, N., Yap, P.-S., Zhang, Yunhai, He, Y., Zhang, Yongjun, 2021.

684   Environmental fate and impacts of microplastics in aquatic ecosystems: a review. RSC Adv. 11,

685   15762–15784. doi:10.1039/D1RA00880C

686 Espinosa, C., Cuesta, A., Esteban, M.Á., 2017. Effects of dietary polyvinylchloride microparticles on

687   general health, immune status and expression of several genes related to stress in gilthead

688   seabream (Sparus aurata L.). Fish Shellfish Immunol. 68, 251–259.

689   doi:10.1016/j.fsi.2017.07.006

690 Espinosa, C., Esteban, M.Á., Cuesta, A., 2019. Dietary administration of PVC and PE microplastics

691   produces histological damage, oxidative stress and immunoregulation in European sea bass

692   (Dicentrarchus labrax L.). Fish Shellfish Immunol. 95, 574–583. doi:10.1016/j.fsi.2019.10.072

693 Esposito, V., Andaloro, F., Bianca, D., Natalotto, A., Romeo, T., Scotti, G., Castriota, L., 2014. Diet

694   and prey selectivity of the red mullet, Mullus barbatus (Pisces: Mullidae), from the southern

695   Tyrrhenian Sea: the role of the surf zone as a feeding ground. Mar. Biol. Res. 10, 167–178.

696   doi:10.1080/17451000.2013.797585

697 Expósito, N., Rovira, J., Sierra, J., Folch, J., Schuhmacher, M., 2021. Microplastics levels, size,

698   morphology and composition in marine water, sediments and sand beaches. Case study of

699   Tarragona coast (western Mediterranean). Sci. Total Environ.

700   doi:10.1016/j.scitotenv.2021.147453

701 Fagiano, V., Alomar, C., Compa, M., Soto-Navarro, J., Jordá, G., Deudero, S., 2022. Neustonic

702   microplastics and zooplankton in coastal waters of Cabrera Marine Protected Area (Western

703   Mediterranean Sea). Sci. Total Environ. 804, 150120. doi:10.1016/J.SCITOTENV.2021.150120

Flajnik, M.F., 2018. A cold-blooded view of adaptive immunity. Nat. Rev. Immunol. 18, 438–453.

doi:10.1038/s41577-018-0003-9

Fortibuoni, T., Amadesi, B., Vlachogianni, T., 2021. Composition and abundance of macrolitter along

the Italian coastline: The first baseline assessment within the european Marine Strategy

Framework Directive. Environ. Pollut. 268, 115886. doi:10.1016/j.envpol.2020.115886

Fossi, M.C., Marsili, L., Baini, M., Giannetti, M., Coppola, D., Guerranti, C., Caliani, I., Minutoli, R.,

Lauriano, G., Finoia, M.G., Rubegni, F., Panigada, S., Bérubé, M., Urbán Ramírez, J., Panti, C.,

2016. Fin whales and microplastics: The Mediterranean Sea and the Sea of Cortez scenarios.

Environ. Pollut. 209, 68–78. doi:10.1016/j.envpol.2015.11.022

Fossi, M.C., Panti, C., Baini, M., Lavers, J.L., 2018. A Review of Plastic-Associated Pressures:

Cetaceans of the Mediterranean Sea and Eastern Australian Shearwaters as Case Studies.

Front. Mar. Sci. 5. doi:10.3389/fmars.2018.00173

Fossi, M.C., Panti, C., Guerranti, C., Coppola, D., Giannetti, M., Marsili, L., Minutoli, R., 2012. Are

baleen whales exposed to the threat of microplastics? A case study of the Mediterranean fin

whale (Balaenoptera physalus). Mar. Pollut. Bull. 64, 2374–2379.

doi:10.1016/j.marpolbul.2012.08.013

Galgani, L., Beiras, R., Galgani, F., Panti, C., Borja, A., 2019. Editorial: Impacts of Marine Litter. Front.

Mar. Sci. 6. doi:10.3389/fmars.2019.00208

Garcia-Vazquez, E., Cani, A., Diem, A., Ferreira, C., Geldhof, R., Marquez, L., Molloy, E., Perché, S.,

2018. Leave no traces – Beached marine litter shelters both invasive and native species. Mar.

Pollut. Bull. 131, 314–322. doi:10.1016/j.marpolbul.2018.04.037

GESAMP, 2015. Sources, fate and effects of microplastics in the marine environment: a global

assessment, IMO/FAO/UN. ed, Rep. Stud. GESAMP.

Giani, D., Baini, M., Galli, M., Casini, S., Fossi, M.C., 2019. Microplastics occurrence in edible fish

species (Mullus barbatus and Merluccius merluccius) collected in three different geographical

sub-areas of the Mediterranean Sea. Mar. Pollut. Bull. 140, 129–137.

doi:10.1016/j.marpolbul.2019.01.005

Gomiero, A., Strafella, P., Pellini, G., Salvalaggio, V., Fabi, G., 2018. Comparative Effects of Ingested

PVC Micro Particles With and Without Adsorbed Benzo(a)pyrene vs. Spiked Sediments on the

Cellular and Sub Cellular Processes of the Benthic Organism Hediste diversicolor. Front. Mar.

Sci. 5. doi:10.3389/fmars.2018.00099

Guerrera, M.C., Aragona, M., Porcino, C., Fazio, F., Laurà, R., Levanti, M., Montalbano, G., Germanà,

G., Abbate, F., Germanà, A., 2021. Micro and Nano Plastics Distribution in Fish as Model

Organisms: Histopathology, Blood Response and Bioaccumulation in Different Organs. Appl.

Sci. 11, 5768. doi:10.3390/app11135768

Guo, X., Wang, J., 2019. The chemical behaviors of microplastics in marine environment: A review.

Mar. Pollut. Bull. 142, 1–14. doi:10.1016/j.marpolbul.2019.03.019

Isobe, A., Iwasaki, S., Uchida, K., Tokai, T., 2019. Abundance of non-conservative microplastics in the

upper ocean from 1957 to 2066. Nat. Commun. 10, 1–3. doi:10.1038/s41467-019-08316-9

Jambeck, J.R., Geyer, R., Wilcox, C., Siegler, T.R., Perryman, M., Andrady, A., Narayan, R., Law,

K.L., 2015. Plastic waste inputs from land into the ocean. Science (80-.). 347, 768–771.

doi:10.1126/science.1260352

Kazour, M., Jemaa, S., Issa, C., Khalaf, G., Amara, R., 2019. Microplastics pollution along the

Lebanese coast (Eastern Mediterranean Basin): Occurrence in surface water, sediments and

biota samples. Sci. Total Environ. doi:10.1016/j.scitotenv.2019.133933

Kiessling, T., Gutow, L., Thiel, M., 2015. Marine Litter as Habitat and Dispersal Vector, in: Marine

Anthropogenic Litter. Springer International Publishing, Cham, pp. 141–181. doi:10.1007/978-3-

319-16510-3_6

Kowalski, N., Reichardt, A.M., Waniek, J.J., 2016. Sinking rates of microplastics and potential

implications of their alteration by physical, biological, and chemical factors. Mar. Pollut. Bull. 109,

310–319. doi:10.1016/j.marpolbul.2016.05.064

Ling, S.D., Sinclair, M., Levi, C.J., Reeves, S.E., Edgar, G.J., 2017. Ubiquity of microplastics in

coastal seafloor sediments. Mar. Pollut. Bull. 121, 104–110.

doi:10.1016/j.marpolbul.2017.05.038

Llorca, M., Álvarez-Muñoz, D., Ábalos, M., Rodríguez-Mozaz, S., Santos, L.H.M.L.M., León, V.M.,

Campillo, J.A., Martínez-Gómez, C., Abad, E., Farré, M., 2020. Microplastics in Mediterranean

coastal area: toxicity and impact for the environment and human health. Trends Environ. Anal.

Chem. 27, e00090. doi:10.1016/j.teac.2020.e00090

Löhr, A., Savelli, H., Beunen, R., Kalz, M., Ragas, A., Van Belleghem, F., 2017. Solutions for global

marine litter pollution. Curr. Opin. Environ. Sustain. 28, 90–99. doi:10.1016/j.cosust.2017.08.009

Lu, Y., Yuan, J., Lu, X., Su, C., Zhang, Y., Wang, C., Cao, X., Li, Q., Su, J., Ittekkot, V., Garbutt, R.A.,

Bush, S., Fletcher, S., Wagey, T., Kachur, A., Sweijd, N., 2018. Major threats of pollution and

climate change to global coastal ecosystems and enhanced management for sustainability.

Environ. Pollut. 239, 670–680. doi:10.1016/j.envpol.2018.04.016

Lucci, F., Della Ventura, G., Piazzolla, D., Venettacci, C., Terribili, A., La Bella, C., Conte, A., Scanu,

S., Bonamano, S., Radica, F., Marcelli, M., 2021. Development and testing for a low-cost device

for airborne PM monitoring: the port of Civitavecchia (Italy) during COVID-19 lockdown. Atmos.

Environ. under Submiss.

Lusher, A.L., McHugh, M., Thompson, R.C., 2013. Occurrence of microplastics in the gastrointestinal

tract of pelagic and demersal fish from the English Channel. Mar. Pollut. Bull. 67, 94–99.

doi:10.1016/j.marpolbul.2012.11.028

Lusher, A.L., Tirelli, V., O'Connor, I., Officer, R., 2015. Microplastics in Arctic polar waters: the first

reported values of particles in surface and sub-surface samples. Sci. Rep. 5, 14947.

doi:10.1038/srep14947

Mahto, S.K., Chandra, P., Rhee, S.W., 2010. In vitro models, endpoints and assessment methods for

the measurement of cytotoxicity. Toxicol. Environ. Health Sci. 2, 87–93.

doi:10.1007/BF03216487

Mancini, E., Miccoli, A., Piazzolla, D., Saraceni, P.R., Lezzi, M., Tiralongo, F., Bonifazi, A., Picchietti,

S., Marcelli, M., 2021. Macrozoobenthic fauna associated with benthic marine litter (Northern

Tyrrhenian Sea, Italy) and first report of two bryozoan species in Italian waters. Reg. Stud. Mar.

Sci. 47, 101912. doi:10.1016/j.rsma.2021.101912

Mancuso, M., Savoca, S., Bottari, T., 2019. First record of microplastics ingestion by European hake

Merluccius merluccius from the Tyrrhenian Sicilian coast (Central Mediterranean Sea). J. Fish

Biol. 94, 517–519. doi:10.1111/jfb.13920

Mazurais, D., Ernande, B., Quazuguel, P., Severe, A., Huelvan, C., Madec, L., Mouchel, O., Soudant,

P., Robbens, J., Huvet, A., Zambonino-Infante, J., 2015. Evaluation of the impact of

polyethylene microbeads ingestion in European sea bass (Dicentrarchus labrax) larvae. Mar.

Environ. Res. 112, 78–85. doi:10.1016/j.marenvres.2015.09.009

Miccoli, A., Buonocore, F., Picchietti, S., Scapigliati, G., 2021a. The sea bass Dicentrarchus labrax as

a marine model species in immunology : Insights from basic and applied research. Aquac. Fish.

doi:10.1016/j.aaf.2021.09.003

Miccoli, A., Guerra, L., Pianese, V., Saraceni, P.R., Buonocore, F., Taddei, A.R., Couto, A., Wolf, T.

De, Fausto, A.M., Scapigliati, G., Picchietti, S., 2021b. Molecular , Cellular and Functional

Analysis of TR γ Chain along the European Sea Bass Dicentrarchus labrax Development. Int. J.

Mol. Sci. 22, 3376. doi:ijms22073376

Moore, C.J., 2008. Synthetic polymers in the marine environment: A rapidly increasing, long-term

threat. Environ. Res. 108, 131–139. doi:10.1016/j.envres.2008.07.025

Nguyen, T.H., Tang, F.H.M., Maggi, F., 2020. Sinking of microbial-associated microplastics in natural

waters. PLoS One 15, e0228209. doi:10.1371/journal.pone.0228209

OECD, 2021. Test No. 249: Fish Cell Line Acute Toxicity - The RTgill-W1 cell line assay, OECD

Guidelines for the Testing of Chemicals, Section 2. OECD, Paris. doi:10.1787/C66D5190-EN

Pannetier, P., Morin, B., Le Bihanic, F., Dubreil, L., Clérandeau, C., Chouvellon, F., Van Arkel, K.,

Danion, M., Cachot, J., 2020. Environmental samples of microplastics induce significant toxic

effects in fish larvae. Environ. Int. 134, 105047. doi:10.1016/j.envint.2019.105047

Panti, C., Giannetti, M., Baini, M., Rubegni, F., Minutoli, R., Fossi, M.C., 2015. Occurrence, relative

abundance and spatial distribution of microplastics and zooplankton NW of Sardinia in the

Pelagos Sanctuary Protected Area, Mediterranean Sea. Environ. Chem. 12, 618.

doi:10.1071/EN14234

Pedà, C., Caccamo, L., Fossi, M.C., Gai, F., Andaloro, F., Genovese, L., Perdichizzi, A., Romeo, T.,

Maricchiolo, G., 2016. Intestinal alterations in European sea bass Dicentrarchus labrax

(Linnaeus, 1758) exposed to microplastics: Preliminary results. Environ. Pollut.

doi:10.1016/j.envpol.2016.01.083

Piazzolla, D., Cafaro, V., de Lucia, G.A., Mancini, E., Scanu, S., Bonamano, S., Piermattei, V.,

Vianello, A., Della Ventura, G., Marcelli, M., 2020. Microlitter pollution in coastal sediments of the

northern Tyrrhenian Sea, Italy: microplastics and fly-ash occurrence and distribution. Estuar.

Coast. Shelf Sci. doi:10.1016/j.ecss.2020.106819

Pittura, L., Avio, C.G., Giuliani, M.E., D'Errico, G., Keiter, S.H., Cormier, B., Gorbi, S., Regoli, F.,

2018. Microplastics as Vehicles of Environmental PAHs to Marine Organisms: Combined

Chemical and Physical Hazards to the Mediterranean Mussels, Mytilus galloprovincialis. Front.

Mar. Sci. 5. doi:10.3389/fmars.2018.00103

Prevenios, M., Zeri, C., Tsangaris, C., Liubartseva, S., Fakiris, E., Papatheodorou, G., 2018. Beach

litter dynamics on Mediterranean coasts: Distinguishing sources and pathways. Mar. Pollut. Bull.

129, 448–457. doi:10.1016/j.marpolbul.2017.10.013

Primpke, S., Wirth, M., Lorenz, C., Gerdts, G., 2018. Reference database design for the automated

analysis of microplastic samples based on Fourier transform infrared (FTIR) spectroscopy. Anal.

Bioanal. Chem. 410, 5131–5141. doi:10.1007/s00216-018-1156-x

Renzi, M., Blašković, A., Bernardi, G., Russo, G.F., 2018. Plastic litter transfer from sediments

towards marine trophic webs: A case study on holothurians. Mar. Pollut. Bull. 135, 376–385.

doi:10.1016/j.marpolbul.2018.07.038

Revel, M., Roman, C., Châtel, A., 2021. Is cell culture a suitable tool for the evaluation of micro- and

nanoplastics ecotoxicity? Ecotoxicology 30, 421–430. doi:10.1007/s10646-021-02355-z

Rios-Fuster, B., Alomar, C., Compa, M., Guijarro, B., Deudero, S., 2019. Anthropogenic particles

ingestion in fish species from two areas of the western Mediterranean Sea. Mar. Pollut. Bull.

144, 325–333. doi:10.1016/j.marpolbul.2019.04.064

Rios-Fuster, B., Arechavala-Lopez, P., García-Marcos, K., Alomar, C., Compa, M., Álvarez, E., Julià,

M.M., Solomando Martí, A., Sureda, A., Deudero, S., 2021. Experimental evidence of

physiological and behavioral effects of microplastic ingestion in Sparus aurata. Aquat. Toxicol.

231, 105737. doi:10.1016/j.aquatox.2020.105737

Rios, L.M., Moore, C., Jones, P.R., 2007. Persistent organic pollutants carried by synthetic polymers

in the ocean environment. Mar. Pollut. Bull. 54, 1230–1237.

doi:10.1016/j.marpolbul.2007.03.022

Rogers, K.L., Carreres-Calabuig, J.A., Gorokhova, E., Posth, N.R., 2020. Micro-by-micro interactions:

How microorganisms influence the fate of marine microplastics. Limnol. Oceanogr. Lett. 5, 18–

36. doi:10.1002/lol2.10136

Ru, J., Huo, Y., Yang, Y., 2020. Microbial Degradation and Valorization of Plastic Wastes. Front.

Microbiol. 11. doi:10.3389/fmicb.2020.00442

Sabatella, E.C., Colloca, F., Coppola, G., Fiorentino, F., Gambino, M., Malvarosa, L., Sabatella, R.,

2017. Key Economic Characteristics of Italian Trawl Fisheries and Management Challenges.

Front. Mar. Sci. 4. doi:10.3389/fmars.2017.00371

Sala, A., Conides, A., De Carlo, F., Klaoudatos, D., Grech, D., Lucchetti, A., Mayans, A., Notti, E.,

Paci, N., Salom, S., Sartor, P., Sbrana, M., Soler, I., Spedicato, M.T., Virgili, M., 2013. Final

855   project report Technical specifications of Mediterranean trawl gears (myGears).

856   doi:10.13140/RG.2.2.24528.76804

857   Savoca, S., Capillo, G., Mancuso, M., Bottari, T., Crupi, R., Branca, C., Romano, V., Faggio, C.,

858   D'Angelo, G., Spanò, N., 2019. Microplastics occurrence in the Tyrrhenian waters and in the

859   gastrointestinal tract of two congener species of seabreams. Environ. Toxicol. Pharmacol. 67,

860   35–41. doi:10.1016/j.etap.2019.01.011

861   Scapigliati, G., Fausto, A.M., Picchietti, S., 2018. Fish Lymphocytes: An Evolutionary Equivalent of

862   Mammalian Innate-Like Lymphocytes? Front. Immunol. 9, 1–8. doi:10.3389/fimmu.2018.00971

863   Schoonen, W.G.E.J., de Roos, J.A.D.M., Westerink, W.M.A., Débiton, E., 2005. Cytotoxic effects of

864   110 reference compounds on HepG2 cells and for 60 compounds on HeLa, ECC-1 and CHO

865   cells. Toxicol. Vitr. 19, 491–503. doi:10.1016/j.tiv.2005.01.002

866   Sendra, M., Pereiro, P., Yeste, M.P., Mercado, L., Figueras, A., Novoa, B., 2021. Size matters:

867   Zebrafish (Danio rerio) as a model to study toxicity of nanoplastics from cells to the whole

868   organism. Environ. Pollut. 268, 115769. doi:10.1016/j.envpol.2020.115769

869   Shim, W.J., Hong, S.H., Eo, S., 2018. Marine Microplastics: Abundance, Distribution, and

870   Composition, in: Zeng, E. (Ed.), Microplastic Contamination in Aquatic Environments. Elsevier,

871   pp. 1–26. doi:10.1016/B978-0-12-813747-5.00001-1

872   Suaria, G., Avio, C.G., Mineo, A., Lattin, G.L., Magaldi, M.G., Belmonte, G., Moore, C.J., Regoli, F.,

873   Aliani, S., 2016. The Mediterranean Plastic Soup: Synthetic polymers in Mediterranean surface

874   waters. Sci. Rep. 6, 1–10. doi:10.1038/srep37551

875   Sussarellu, R., Suquet, M., Thomas, Y., Lambert, C., Fabioux, C., Pernet, M.E.J., Le Goïc, N.,

876   Quillien, V., Mingant, C., Epelboin, Y., Corporeau, C., Guyomarch, J., Robbens, J., Paul-Pont, I.,

877   Soudant, P., Huvet, A., 2016. Oyster reproduction is affected by exposure to polystyrene

878   microplastics. Proc. Natl. Acad. Sci. 113, 2430–2435. doi:10.1073/pnas.1519019113

879   Tekman, M.B., Gutow, L., Macario, A., Haas, A., Walter, A., Bergmann, M., 2021. LITTERBASE

880   [WWW Document]. Alfred Wegener Inst. Helmholtz Cent. Polar Mar. Res. URL

881   https://litterbase.awi.de/litter_detail

882   Thompson, R.C., Olsen, Y., Mitchell, R.C., Davis, A., Rowland, S.J., John, A.W.G., McGonigle, D.,

883   Russel, A.E., 2004. Lost at Sea: Where Is All the Plastic? Science (80-.). 304, 838–838.

doi:10.1126/science.1094559

Tiralongo, F., Mancini, E., Ventura, D., de Malherbe, S., Paladini de Mendoza, F., Sardone, M.,

Arciprete, R., Massi, D., Marcelli, M., Fiorentino, F., Minervini, R., 2021. Commercial catches

and discards composition in the central Tyrrhenian Sea: a multispecies quantitative and

qualitative analysis from shallow and deep bottom trawling. Mediterr. Mar. Sci. 22, 521–531.

doi:10.12681/mms.25753

UNEP/MAP, 2015. Marine litter assessment in the Mediterranean Sea. Athens (Greece).

UNEP, 2016. Marine plastic debris and microplastics - Global Lessons and Research to Inspire Action

and guide policy change, UN.

UNEP, 2014. Valuing Plastics: The Business Case for Measuring, Managing and Disclosing Plastic

Use in the Consumer Goods Industry, UN.

van der Hal, N., Ariel, A., Angel, D.L., 2017. Exceptionally high abundances of microplastics in the

oligotrophic Israeli Mediterranean coastal waters. Mar. Pollut. Bull.

doi:10.1016/j.marpolbul.2016.12.052

Van Sebille, E., Aliani, S., Law, K.L., Maximenko, N., Alsina, J.M., Bagaev, A., Bergmann, M.,

Chapron, B., Chubarenko, I., Cózar, A., Delandmeter, P., Egger, M., Fox-Kemper, B., Garaba,

S.P., Goddijn-Murphy, L., Hardesty, B.D., Hoffman, M.J., Isobe, A., Jongedijk, C.E., Kaandorp,

M.L.A., Khatmullina, L., Koelmans, A.A., Kukulka, T., Laufkötter, C., Lebreton, L., Lobelle, D.,

Maes, C., Martinez-Vicente, V., Morales Maqueda, M.A., Poulain-Zarcos, M., Rodríguez, E.,

Ryan, P.G., Shanks, A.L., Shim, W.J., Suaria, G., Thiel, M., Van Den Bremer, T.S., Wichmann,

D., 2020. The physical oceanography of the transport of floating marine debris. Environ. Res.

Lett. doi:10.1088/1748-9326/ab6d7d

van Sebille, E., Wilcox, C., Lebreton, L., Maximenko, N., Hardesty, B.D., van Franeker, J.A., Eriksen,

M., Siegel, D., Galgani, F., Law, K.L., 2015. A global inventory of small floating plastic debris.

Environ. Res. Lett. 10, 124006. doi:10.1088/1748-9326/10/12/124006

Vasilopoulou, G., Kehayias, G., Kletou, D., Kleitou, P., Triantafyllidis, V., Zotos, A., Antoniadis, K.,

Rousou, M., Papadopoulos, V., Polykarpou, P., Tsiamis, G., 2021. Microplastics Investigation

Using Zooplankton Samples from the Coasts of Cyprus (Eastern Mediterranean). Water 2021,

Vol. 13, Page 2272 13, 2272. doi:10.3390/W13162272

Wright, S.L., Thompson, R.C., Galloway, T.S., 2013. The physical impacts of microplastics on marine

organisms: A review. Environ. Pollut. 178, 483–492. doi:10.1016/j.envpol.2013.02.031

Yong, C., Valiyaveettil, S., Tang, B., 2020. Toxicity of Microplastics and Nanoplastics in Mammalian

Systems. Int. J. Environ. Res. Public Health 17, 1509. doi:10.3390/ijerph17051509

Zapata, A., Diez, B., Cejalvo, T., Gutiérrez-De Frías, C., Cortés, A., 2006. Ontogeny of the immune

system of fish. Fish Shellfish Immunol. 20, 126–136. doi:10.1016/j.fsi.2004.09.005

Zettler, E.R., Mincer, T.J., Amaral-Zettler, L.A., 2013. Life in the "Plastisphere": Microbial Communities

on Plastic Marine Debris. Environ. Sci. Technol. 47, 7137–7146. doi:10.1021/es401288x

Zhang, W., Zhang, S., Wang, J., Wang, Y., Mu, J., Wang, P., Lin, X., Ma, D., 2017. Microplastic

pollution in the surface waters of the Bohai Sea, China. Environ. Pollut. 231, 541–548.

doi:10.1016/j.envpol.2017.08.058

Zhao, S., Zhu, L., Wang, T., Li, D., 2014. Suspended microplastics in the surface water of the Yangtze

Estuary System, China: First observations on occurrence, distribution. Mar. Pollut. Bull. 86, 562–

568. doi:10.1016/j.marpolbul.2014.06.032

Zitouni, N., Bousserrhine, N., Missawi, O., Boughattas, I., Chèvre, N., Santos, R., Belbekhouche, S.,

Alphonse, V., Tisserand, F., Balmassiere, L., Dos Santos, S.P., Mokni, M., Guerbej, H., Banni,

M., 2021. Uptake, tissue distribution and toxicological effects of environmental microplastics in

early juvenile fish Dicentrarchus labrax. J. Hazard. Mater. 403.

doi:10.1016/j.jhazmat.2020.124055

Zwollo, P., Quddos, F., Bagdassarian, C., Seeley, M.E., Hale, R.C., Abderhalden, L., 2021.

Polystyrene microplastics reduce abundance of developing B cells in rainbow trout

(Oncorhynchus mykiss) primary cultures. Fish Shellfish Immunol. 114, 102–111.

doi:10.1016/j.fsi.2021.04.014