Research Article

Sonic hedgehog is basolaterally sorted from the TGN and transcytosed to
the apical domain involving Dispatched-1 at Rab11-ARE

Lisette Sandoval[1,#], Mariana Labarca[1,#], Claudio Retamal[1], Paula Sanchez[2],
Juan Larraín[2], Alfonso González[1,2,3]

[1]Centro de Biología Celular y Biomedicina (CEBICEM), Facultad de Medicina y
Ciencia, Universidad San Sebastián, Santiago 7510157, Chile. [2]Centro de
Envejecimiento y Regeneración (CARE), Facultad de Ciencias Biológicas, Pontificia
Universidad Católica de Chile, Santiago 8330025, Chile. [3]Fundación Ciencia y Vida,
Santiago 7780272, Chile.

[#] These authors contributed equally and are considered first authors

Keywords: Hedgehog, polarity, trafficking, dispatched, Rab11, lipidation.

17
The authors declare no conflicts of interest

Corresponding author: Alfonso González

Centro de Biología Celular y Biomedicina (CEBICEM)

Facultad de Medicina y Ciencia

Universidad San Sebastián

Santiago 7510157, Chile

E-mail: alfonso.gonzalez@uss.cl

Phone: +56 2 22606300

ABSTRACT

Hedgehog (Hh) secretion from apical and/or basolateral domains occurs in different epithelial cells impacting development and tissue homeostasis. Palmitoylation and cholestyrolation attach Hh proteins to membranes and Dispatched-1 (Disp-1) promotes their release. How these lipidated proteins are handled by the complex secretory and endocytic pathways of polarized epithelial cells remains unknown. We show that MDCK cells address newly synthesized sonic hedgehog (Shh) from the TGN to the basolateral cell surface and then to the apical domain through a transcytosis pathway that includes Rab11-apical recycling endosomes (Rab11-ARE). Both palmitoylation and cholestyrolation contribute to this sorting behavior, otherwise Shh lacking these lipid modifications is unpolarized. Disp-1 mediates first basolateral secretion from the TGN and then transcytosis from the Rab11-ARE. At steady state, Shh predominates apically and can be basolaterally transcytosed. This complex Shh trafficking provides several steps for regulation and variation in different epithelia, subordinating the apical to the basolateral secretion.

INTRODUCTION

Hedgehog proteins (Hhs) are signaling determinants of embryonic patterning,

differentiation and organogenesis and after birth contribute to tissue homeostasis and repair

(Beachy et al., 2004; Briscoe and Therond, 2013; Guerrero and Kornberg, 2014; Parchure et

al., 2018; Petrov et al., 2017). Alterations of Hhs functions are involved in developmental

defects, tissue fibrosis and cancer (Briscoe and Therond, 2013; Edeling et al., 2016; Machado

and Diehl, 2017). In most of these processes the function of Hhs depends on polarized

secretion from epithelial cells that possess apical and basolateral domains facing opposite

compartments of the organism (Briscoe and Therond, 2013; Guerrero and Kornberg, 2014; Mao

et al., 2018; Matusek et al., 2020; Parchure et al., 2018; Peng et al., 2015; Petrov et al., 2017;

Walton and Gumucio, 2021). Apical secretion sends signals to neighboring epithelial cells while

basolateral secretion targets neighboring epithelial cells and underlying stromal cells

(Chamberlain et al., 2008; Lehmann et al., 2020; Mao et al., 2018; Walton and Gumucio, 2021).

Elucidating the Hhs intracellular trafficking pathways in epithelial cells is then required to fully

understand the signaling function of these important morphogens in development, tissue

homeostasis and pathology.

Hhs are synthesized in the endoplasmic reticulum (ER) as a 45kD precursor that is

autocatalytically cleaved and processed, generating a ~20kDa N-terminal peptide modified by

the covalent addition of cholesterol to its C-terminus and palmitate to its N-terminal end (Chen

et al., 2011; Pepinsky et al., 1998; Porter et al., 1996). These lipid moieties attach the mature

and functionally competent Hhs to the membrane (Petrov et al., 2017). Multiple forms of Hhs

secretion have been described, including oligomers, lipoproteins, exosomes and a main

releasing mechanism from the plasma membrane involving the transmembrane and cholesterol

binding protein Dispatched (Disp) (Burke et al., 1999; Caspary et al., 2002; Creanga et al.,

2012; Parchure et al., 2018; Stewart et al., 2018; Tukachinsky et al., 2012). Disp functions as

Hhs/Na+ antiporter driven by the Na+ gradient (Petrov et al., 2020; Wang et al., 2021) and

becomes functionally competent after its cleavage at the cell surface by the proprotein

convertase Furin, which in epithelial cells occurs at the basolateral plasma membrane (Hall et

al., 2019; Stewart et al., 2018).

In different epithelial cells Hhs can be found localized to the apical, basolateral or both

plasma membrane domains and their polarized secretion can be inferred from target cell

responses. Sonic hedgehog (Shh), the most studied vertebrate orthologs of Hhs (Parchure et

al., 2018; Petrov et al., 2017), generates paracrine responses of stromal mesenchymal cells

underneath the epithelial cells of adult intestine (Buller et al., 2012; Shyer et al., 2015), lung

(Mao et al., 2018; Peng et al., 2015), and kidney (Ding et al., 2012; Edeling et al., 2016; Zhou et

al., 2014), indicating basolateral secretion. Instead, gastric parietal cells distribute and secrete

Shh mostly apically, showing also detectable levels at the basolateral side (Zavros et al., 2008).

Apically secreted Shh is also found in the neural tube of chick (Danesin et al., 2006) and mouse

(Chamberlain et al., 2008) embryos. In the mature airway epithelia, Shh is secreted from both

apical and basolateral domains, eliciting different signaling pathways and functional

consequences at each compartment (Mao et al., 2018).

The routes followed by Hhs along the complex vesicular trafficking pathways of

epithelial cells have been mainly explored in *Drosophila* (Guerrero and Kornberg, 2014;

Therond, 2012). Extensive studies interpreting subcellular localizations in producing cells and

responses of target cells describe both apical and basolateral pools of Hh and propose a

primary biosynthetic pathway to the apical domain, followed by endocytosis and either apical

recycling (D'Angelo et al., 2015) or transcytosis of Hh to the basolateral domain (Callejo et al.,

2011). These alternative pathways remain highly debated and would determine different

secretion and signaling characteristics (Gore et al., 2021; Matusek et al., 2020; Simon et al.,

2016). How extensive these *Drosophila* models are to mammalian epithelial cells is unknown.

The biosynthetic routes of different epithelial cells display variations reflecting plasticity

in the polarity program (Rodriguez-Boulan and Macara, 2014). Exocytic pathways emerging

from the TGN can predominate towards the basolateral domain, from where apical proteins then

undergo transcytosis, as in intestinal cells and hepatocytes (Bartles et al., 1987; Hauri and

Matter, 1991; Rindler and Traber, 1988; Tuma and Hubbard, 2003). In other epithelial cells,

epitomized by kidney epithelial cells, apical and basolateral proteins are first sorted at the TGN

into distinct polarized secretion pathways of equivalent transport capacity and endocytic

proteins can then recycle to the same or the opposite cell surface (Gonzalez and Rodriguez-

Boulan, 2009; Mellman and Nelson, 2008; Rodriguez-Boulan and Macara, 2014; Tuma and

Hubbard, 2003). Protein trafficking along polarized vesicular pathways is controlled by sorting

signals embedded in the cargo and decoded by a complex machinery coupled to the generation

of vesicular/tubular vehicles (Rodriguez-Boulan and Macara, 2014). Transmembrane proteins

can have apical sorting signals in extracellular, membrane-spanning or cytosolic domains, or

basolateral signals located in their cytosolic domains (Bonifacino, 2014; Gonzalez and

Rodriguez-Boulan, 2009; Mellman and Nelson, 2008; Rodriguez-Boulan and Macara, 2014).

Luminal proteins anchored to membranes through lipid moieties, such as Hh proteins and

glycosylphosphatidylinositol-anchored proteins (GPI-APs), associate with lipid rafts (Long et al.,

2015; Parchure et al., 2018; Petrov et al., 2017), originally proposed as platforms for apical

sorting (Simons and Ikonen, 1997). Although most GPI-APs are addressed to the apical cell

surface, there are also examples of basolateral sorting of GPI-APs associated with lipid-rafts

(Imjeti et al., 2011; Lebreton et al., 2021; Lebreton et al., 2019; Paladino et al., 2015; Sezgin et

al., 2017). Previous studies describe Hhs subcellular distributions and secretion at steady-state

conditions that do not define the trafficking pathways followed by newly synthesized Hhs

proteins in epithelial cells.

Madin-Darby canine kidney (MDCK) cells forming polarized monolayers in culture

constitute the most characterized model system for studies of protein trafficking in epithelial

cells (Rodriguez-Boulan and Macara, 2014). In MDCK cells, proteins lacking polarity sorting

signals are distributed unpolarized, reflecting TGN-derived apical and basolateral routes of

equivalent capacity (Gonzalez et al., 1987; Gottlieb et al., 1986), whereas domain-selective

secretion implies specific sorting events occurring at the TGN (Gonzalez et al., 1993; Scheiffele

et al., 1995). A unique study in transfected MDCK cells describes Shh steady-state apical and

basolateral distribution and enhanced basolateral secretion by coexpressed Disp-1 (Etheridge

et al., 2010). This study leaves uncertainty as to whether Shh basolateral secretion occurred

directly from the TGN or indirectly through transcytosis from the apical domain, as suggested in

Drosophila (Callejo et al., 2011).

A combination of experimental approaches previously used for GPI-APs (Paladino et

al., 2006; Zurzolo et al., 1993), transmembrane (Cancino et al., 2007; Gan et al., 2002; Kreitzer

et al., 2000) and secretion (Gonzalez et al., 1987) proteins can be applied to define the

biosynthetic and endocytic trafficking routes of newly synthesized Shh. Here we evaluate Shh

trafficking in MDCK cells using pulse-chase domain-specific targeting assays (Cancino et al.,

2007; Paladino et al., 2006), plasmid microinjection expression combined with synchronized

post-TGN trafficking imaging of Shh, co-expression with Disp1 in live cells (Cancino et al., 2007;

Kreitzer et al., 2003), and cell surface tagging with biotin or antibody. The results reveal that

newly synthesized Shh is first sorted from the TGN to the basolateral plasma membrane and

then undergo transcytosis to the apical cell surface involving the Rab11-apical recycling

endosome (Rab11-ARE). Shh polarized sorting requires the lipid modifications and therefore

depends on membrane association. At steady state, Shh is mainly localized at the apical cell

surface and its secretion is also mainly apical. A small proportion recycles back to the

basolateral cell surface. We also show that Disp-1 first promotes basolateral secretion of newly

synthesized Shh transported from the TGN and then is required for its transcytosis through the

Rab11-ARE to the apical domain. Such a complex trafficking behavior predicts that apical Shh

secretion reciprocally depends on previous basolateral secretion and transcytosis processes

that are both sensitive to Disp-1 cleavage and expression levels.

RESULTS

Predominant apical distribution and secretion of Sonic Hedgehog in polarized

epithelial MDCK cells

Previous studies indicate that Shh can be found distributed at and secreted from apical

and basolateral domains with variable preponderance in different epithelial cells (Briscoe and

Therond, 2013; Chamberlain et al., 2008; Guerrero and Kornberg, 2014; Mao et al., 2018;

Petrov et al., 2017; Shyer et al., 2015; Zavros et al., 2008; Zhou et al., 2014). To explore the

trafficking behavior of Shh in a widely characterized epithelial model system (Mellman and

Nelson, 2008; Rodriguez-Boulan and Macara, 2014), we generated permanently transfected

MDCK cells expressing different levels of Shh and first assessed the polarity of Shh distribution

and secretion at steady state conditions. Domain-specific cell surface biotinylation of MDCK-

Shh cells grown in Transwell chambers revealed Shh either highly polarized at the apical

domain or unpolarized, depending on its expression levels (Figure 1A). For instance, MDCK-

Shh clone-1 distributed Shh mainly apically whereas the MDCK-Shh clone-2 displaying higher

expression levels displayed Shh similarly at apical and basolateral cell surfaces. E-cadherin and

Na-K-ATPase, two basolateral proteins, demonstrated that both MDCK clones are polarized

(Figure 1A). Treatment of MDCK-Shh clone 1 with Na^+-butyrate treatment, known to enhance

the expression of transfected plasmids (Burgos et al., 2004; Gonzalez et al., 1987), led to a loss

of the predominant apical Shh location, while maintaining the polarity of endogenous E-cadherin

Na^+-K^+-ATPase (Figure 1A). These results indicate a predominant apical sorting of Shh through

a process that can be saturated at high expression levels. We then focused on MDCK-Shh

clone-1, named afterwards as MDCK-Shh, which mainly distributed Shh apically. A quantitative

assessment showed ~80% Shh distribution at the apical cell surface and more than 90% apical

secretion of Shh in this clone (Figure 1B). The predominant apical pattern of Shh can be also

observed by indirect immunofluorescence (Figure 1C).

Shh is basolaterally sorted and then transcytosed to the apical cell surface

Shh is known to associate with lipid rafts (Long et al., 2015; Parchure et al., 2018;

Petrov et al., 2017), similar to GPI-APs, which in general are mainly addressed to the apical

domain from the TGN (Hua et al., 2006; Paladino et al., 2006). Therefore, to test whether newly

synthesized Shh is directly sorted to the apical cell surface or follows an indirect basolateral-to-

apical transcytotic route we pulse-labeled MDCK-Shh cells with [^{35}S]-methionine/cysteine and

then chased the arrival of newly synthesized [^{35}S]-Shh to the apical and basolateral cell

surfaces using an established biotinylation targeting assay (Le Bivic et al., 1989; Zurzolo et al.,

1992). This approach showed that [^{35}S]-Shh first became enriched at the basolateral surface

and then progressively increased at the apical cell surface (Figure 2A). E-cadherin used as

internal control displayed a basolateral distribution at all chased times. In contrast with Shh,

GFP-NO-GPI distributed mostly to the apical cell surface at all chased times (Figure 2B),

indicating that our targeting assay is able to detect the TGN-to-apical cell surface pathway

previously described for raft-associated GPI-APs in MDCK cells (Imjeti et al., 2011). We also

found [^{35}S]-Shh in the apical media after 120 minutes of chase, while basolateral secretion was

barely detectable (Figure 2C). These results suggest that newly synthesized Shh is primarily

sorted to the basolateral domain from the TGN and then is transported by transcytosis to the

apical cell surface.

To directly test basolateral-to-apical transcytosis, we first performed an established

assay based on the reducible reagent sulfo-NHS-SS-biotin (Burgos et al., 2004; Zurzolo et al.,

1992). After 120 min of transport from the biotinylated basolateral cell surface, approximately

25% of Shh became protected from reduction by MesNa added to the basolateral chamber

(Figure 2D). MesNa added to the apical side also decreased the biotinylation levels of

basolaterally biotinylated Shh, but not E-cadherin (Figure 2D). Furthermore, we also found in

the apical media Shh previously biotinylated from the basolateral cell surface (Figure 2E).

These results indicate that Shh already present at the basolateral cell surface reaches the

apical cell surface and is apically secreted by transcytosis.

We further corroborate these findings and analyze the route of Shh to the apical domain

we tagged the basolateral pool of Shh with anti-Shh antibodies at 4°C and then shifted the cells

to 37°C for different times. The basolateral pattern of antibody-tagged Shh (0 min) changed

within 60 min to a dot-like pattern typical of the apical recycling endosomes (ARE) and then

acquired an apical pattern (Figure 3A). Colocalization of Shh with Rab11 indicated that the

subapical dot-like compartment actually corresponds to Rab11-ARE (Figure 3B). Therefore,

basolaterally sorted Shh follows the well-known transcytotic route widely characterized for the

polyimmunoglobulin (PIgR), which includes the Rab11- ARE (Apodaca et al., 1994; Brown et

al., 2000; Wang et al., 2000).

Shh can recycle from the apical cell surface back to the basolateral cell surface

Our results so far indicate that newly synthesized Shh follows a transcytotic pathway

from the basolateral to the apical membrane. This is congruent with one of the models proposed

for Hh protein trafficking in *Drosophila* (Gallet et al., 2003; Gallet et al., 2006). However, the

opposite apical-to-basolateral Hh transcytosis has also been proposed in *Drosophila* (Callejo et

al., 2011; Guerrero and Kornberg, 2014). Therefore, we used the same type of antibody-tagging

experiment and looked for apical-to-basolateral transcytosis. We find that apically tagged Shh

can also redistribute to the basolateral cell surface (Figure 3C). Therefore, as suggested in

Drosophila wing disk epithelia (Callejo et al., 2011; Guerrero and Kornberg, 2014), once arriving

to the apical surface Shh can recycle back to the basolateral cell surface by transcytosis.

Role of lipid modifications in Shh polarized sorting

Hhs proteins are synthesized as a precursor protein that is autocleaved and processed

by cholesteroylation at the C-terminus and palmitoylation at the N-terminus (Chen et al., 2011;

Pepinsky et al., 1998; Porter et al., 1996) (Figure 4A). These lipid modifications determine Hhs

membrane association, nano-scale organization, mode of secretion and range of action

(Parchure et al., 2018; Petrov et al., 2017). Therefore, we assessed the polarity of Shh mutants

lacking both palmitoylation and cholesterylation (ShhNC24S), only cholesterylation (ShhN) or

palmitoylation (ShhNpC24S). ShhNC24S that lacks both lipid modifications and does not

associate with membrane (Feng et al., 2004) was secreted without polarity (Figure 4B).

Therefore, the peptidic structure of Shh either lacks specific sorting signals for polarity or

requires membrane association to expose sorting competent signals, perhaps through nano-

scale organization (Vyas et al., 2008; Vyas et al., 2014). We also found the C-terminal truncated

mutant ShhN lacking cholesterylation associated to the basolateral cell surface and secreted

unpolarized (Figure 4C). Instead, ShhNpC24S lacking the cysteine for palmitoylation and

conserving the cholesterol modification preferentially distributed to the basolateral membrane

and was mostly secreted apically (Figure 4D). Pulse-chase domain-specific targeting assays

showed that newly synthesized ShhNpC24S arrived first at the basolateral cell surface and then

at the apical cell surface and medium (Figure 4E), similar to the pathway followed by the fully

lapidated protein (See Figure 2A). The low detection levels at the apical cell surface suggest a

faster release of this mutant. The transcytosis assay corroborated an apical secretion of

basolaterally biotinylated-ShhNpC24S (Figure 4F). These results suggest that ShhNpC24S,

similar to wild-type Shh, is initially sorted to the basolateral cell surface and then becomes

transcytosed to the apical domain for secretion. All these observations reveal that both palmitoyl

and cholesteroyl modifications contribute to Shh sorting to the basolateral cell surface. The

cholesterol adduct would be further required for subsequent transcytosis.

Disp-1 distribution and role in newly synthesized Shh secretion

Disp is required for the release of Hh proteins from expressing cells (Burke et al., 1999;

Caspary et al., 2002; Petrov et al., 2017; Stewart et al., 2018). In polarized epithelial cells, Disp

has been described to promote Hh basolateral secretion (Callejo et al., 2011; Etheridge et al.,

2010; Stewart et al., 2018). We used plasmid microinjection (Cancino et al., 2007; Kreitzer et

al., 2003) to express Shh alone or together with HA-tagged Disp-1 (Stewart et al., 2018).

Microinjection avoids the high levels of overexpression achieved by transient cotransfection

experiments, which can alter polarized sorting (Burgos et al., 2004). After 4 h post-

microinjection we clearly found Shh distributed apically, contrasting with the co-expressed

basolateral marker LDLR-GFP (Cancino et al., 2007; Gravotta et al., 2007; Guo et al., 2013)

(Figure 5A). Using these conditions, we co-expressed Shh together with Disp-1 and incubated

the intact cells with anti-Shh added to the apical side to detect Shh at the cell surface. Then we

permeabilized the cells for Disp-1 immunodetection and found it localized just below the Shh

labeled at the apical cell surface (Figure 5B). In the next experiments, we fixed and

permeabilized the cells to immunolocalize Shh and Disp-1 at the same time. Confocal images

were then merged and shown as either maximal projections or just a basolateral section at the

level of the nucleus (Figure 5C). Shh expressed alone can be seen in its typical apical location

Shh (Figure 5C, upper panel), while when co-expressed with Disp-1 also became visible at the

basolateral border of neighboring non-microinjected cells (Figure 5C, middle panel). This

indicates that Disp-1 enhanced the basolateral secretion of Shh, as previously described

(Etheridge et al., 2010). Shh and Disp-1 mostly segregated from each other, with Disp-1

occupying a subapical location overlaid by Shh. Both proteins also colocalized in few spots

(Figure 5C, middle panel).

Then, we co-expressed Shh together with the cleavage deficient recombinant mutant

Disp-1-R279A/E280A (Disp-1-CS) incompetent for Shh secretion (Stewart et al., 2018). Disp-1-

CS displayed an overall similar distribution as Disp-1. However, Disp-1-CS coexpressing cells

showed Shh predominantly at the subapical compartment, likely corresponding to the ARE

(Figure 5C, bottom panel), and lost the Shh apical location in expressing cells and its

basolateral detection in neighboring cells (Figure 5C, bottom panel). These results suggest that

Disp-1-CS not only inhibited Shh secretion from the basolateral surface where Shh first arrived,

but also arrested Shh at the ARE, while *en route* to the apical cell surface from the basolateral

plasma membrane.

Using less amounts of microinjected plasmids to achieve lower levels of expression we

detected colocalization of Shh and Disp-1 in the Rab11-ARE compartment, which included

tubular structures (Figure 6). High resolution images (3D surface rendering) revealed

colocalization of Shh, Disp-1 and Rab11 in the tubular structures of the ARE compartment

(Figure 6; middle panel, arrowheads). Even though most of Disp-1 is seen at subapical planes,

overexposure of Z confocal planes corresponding to the basolateral volume of the cell clearly

showed Disp-1 near the lateral membrane, where few vesicles containing Shh or Rab11 were

also detectable (Figure 6, bottom panel).

To further analyze the role of Disp-1 upon newly synthesized Shh biosynthetic

trafficking we performed live-cell imaging from the TGN to cell surface transport of Shh coupled

to GFP (Shh-GFP) (Chamberlain et al., 2008). We first corroborated in plasmid-microinjected

MDCK cells for Shh-GFP to acquire an apical distribution (Figure 7A) using the route of Rab-11-

ARE (Figure 7B). Then, we followed Shh-GFP trafficking from the TGN to the cell surface in a

synchronized manner with previously established methods (Cancino et al., 2007; Guo et al.,

2013; Kreitzer et al., 2003). After plasmid microinjection, we incubated the cells for 45 minutes

at 37ºC to allow for detectable expression levels and then incubated the cells at 20ºC for 2h to

arrest the biosynthetic trafficking at the TGN. At this initial time, Shh-GFP expressed alone or in

combination with Disp-1 or Disp-1-CS mutant displayed the perinuclear pattern characteristic of

the TGN (Figure 7C, 0 min) (Cancino et al., 2007). Shifting the cells to 37ºC allowed the TGN

exit in the presence of cycloheximide to avoid further protein synthesis. Shh-GFP first became

detectable at the cell border, reflecting its basolateral arrival, and then progressively acquired

an ARE location and the typical apical pattern (Figure 7C). A halo of receiving cells surrounding

the expressing cells displayed Shh in their border, revealing the Shh basolateral secretion from

producing cells (see also Fig 5C middle panel). This can be seen more clearly in the

overexposed image corresponding to 120 min post-TGN trafficking (Figure 7C, middle panel).

Cells co-expressing Disp-1 showed an enhanced Shh-GFP basolateral secretion, as judged by

the higher intensity and longer distance of the fluorescent borders of neighboring non-

expressing cells (Figure 7C, middle panel). These results indicate that post-TGN basolateral

secretion of Shh is sensitive to Disp-1 expression levels and provide direct evidence of Shh

sorting from the TGN to the basolateral cell surface. In striking contrast, cells co-expressing the

Disp-1-CS mutant showed a decreased basolateral distribution in neighboring cells (Figure 7C,

lower panel), indicating lower basolateral secretion levels. Furthermore, Shh-GFP accumulated

in a punctate compartment suggesting an arrest at the ARE (Figure 7C, lower panel). Taken

together, all these experiments indicate that Disp1 promotes Shh secretion first from the

basolateral cell surface where it arrives from the TGN. The effects of coexpressing Disp-1-CS

corroborate that Shh secretion requires Disp-1 processing by cleavage (Stewart et al., 2018)

and suggest that Disp-1 is also required for trafficking along the basolateral-to-apical

transcytosis route at the level of the ARE, where Shh accumulates if Disp-1 function is

disturbed.

DISCUSSION

In diverse epithelial systems, Hhs are found apically and basolaterally distributed and

secreted with variable polarity (Chamberlain et al., 2008; Mao et al., 2018; Therond, 2012). An

open question is whether Hh subcellular distribution and secretion reflects default-unsorted or

specific-sorting trafficking events. The steady-state conditions of previous studies cannot

answer this question and have even led to controversial models (Ayers et al., 2010; Callejo et

al., 2011; Gallet, 2011; Guerrero and Kornberg, 2014; Therond, 2012). We addressed this

problem using MDCK cells as model system and combined pulse chase and live cell imaging to

follow the cell surface arrival of newly synthesized Shh, while domain specific cell surface

tagging directly assessed transcytosis from each polarized domain. Our results provide new

information regarding the intracellular routes taken by newly synthesized Shh, which can help to

interpret and understand previous observations in different epithelial systems.

We find a predominant (~80%) apical cell surface distribution and secretion of Shh in

permanently transfected MDCK cells under steady state conditions. However, pulse-chase

domain-specific targeting assays reveal that newly synthesized Shh arrives first to the

basolateral cell surface and then to the apical plasma membrane. Live cell imaging after

releasing a 20C° TGN arrest demonstrates Shh basolateral sorting directly from the TGN.

Basolateral tagging with biotinylation or antibody binding directly demonstrate transcytosis of

basolateral Shh to the apical domain. The transcytosis pathway of Shh includes the Rab11-ARE

compartment, similar to the most characterized transcytosis route of pIgR (Apodaca et al., 1994;

Rojas and Apodaca, 2002). A reverse transcytosis of Shh from the apical to the basolateral

domain can also occur, as shown by apical tagging with antibody. All these observations

suggest a model in which newly synthesized Shh is first addressed to the basolateral domain

from the TGN and then becomes transported through the Rab11-ARE transcytotic route to the

opposite apical domain, from which most of its secretion takes place while a small proportion

returns to the basolateral cell surface. Our results imply specific sorting events operating first at

the TGN and then at the basolateral plasma membrane.

MDCK cells possess apical and basolateral exocytic pathways of roughly equivalent

transporting capacity and therefore newly synthesized proteins lacking sorting information would

follow these pathways by default resulting in their unpolarized distribution or secretion

(Gonzalez et al., 1987; Gottlieb et al., 1986; Scheiffele et al., 1995). In contrast, intestinal

epithelial cells and hepatocytes have a prominent basolateral exocytic route into which both

basolateral and apical proteins can enter and once reaching the cell surface the apical proteins

become transcytosed to the opposite domain (Bartles et al., 1987; Hauri and Matter, 1991;

Hubbard, 1989; Rindler and Traber, 1988). Proteins lacking information for TGN-mediated

sorting that are unpolarized in MDCK cells can be basolaterally addressed in intestinal cells

(Rindler and Traber, 1988). Our results in MDCK cells clearly indicate that Shh possess

basolateral sorting information decoded first at the TGN and predicts its primary basolateral cell

surface arrival in different epithelial cells.

We initially considered that Shh sorting behavior might share similarities with GPI-

proteins due to their common association with lipid rafts at the luminal membrane leaflet (Mayor

and Riezman, 2004). GPI-proteins are most commonly addressed to the apical cell surface from

the TGN in MDCK cells (Imjeti et al., 2011; Lebreton et al., 2019; Lisanti et al., 1988; Paladino

et al., 2006). However, our pulse chase experiments revealed basolateral sorting of Shh under

the same conditions that corroborated apical biosynthetic trafficking of a previously

characterized recombinant GPI-protein (Imjeti et al., 2011). Lipid-raft association is certainly not

enough for apical sorting (Lebreton et al., 2019; Paladino et al., 2004). A proposed additional

requirement for at least apical GPI-proteins is clustering at the TGN promoted by cholesterol

and luminal calcium (Lebreton et al., 2021; Paladino et al., 2004), or glycosylation in some

epithelial cells (Imjeti et al., 2011). GPI-proteins lacking this clustering property, such as the

prion protein, are instead addressed to the basolateral domain (Lebreton et al., 2021; Sarnataro

et al., 2002). Prion also becomes transcytosed from the basolateral to the apical domain

(Arkhipenko et al., 2016), similar to what we found here for Shh. The mechanism through which

lipid-raft associated proteins are basolaterally sorted, endocytosed and addressed to the

opposite domain by transcytosis remains unknown.

Epithelial cells have transcytosis pathways from the apical or basolateral domains to the

opposite cell surface (Mostov et al., 1995; Tuma and Hubbard, 2003). Transmembrane proteins

can follow these pathways guided by specific sorting signals or associated with default

movement of membranes. For instance, the cytosolic tail of pIgR contains sorting signals for

basolateral delivery, clathrin-mediated endocytosis and transcytosis (Luton et al., 2009; Mostov

et al., 1995), whereas the basolateral recycling transferrin receptor reaches the apical domain

when cells do not express the AP1B-adaptor responsible for its basolateral sorting (Perez Bay

et al., 2013), suggesting a default post-endocytic trafficking. Both pathways include transit

through the Rab11-ARE compartment (Luton et al., 2009; Mostov et al., 1995; Perez Bay et al.,

2013). We show that Shh still associated with the basolateral plasma membrane moves towards

the apical cell surface using the Rab11-ARE transcytotic route. The mechanism might involve

yet unidentified transmembrane escorting proteins linking Shh to the cytosolic sorting machinery

or a default pathway from the basolateral cell surface.

Cholesterol modification is a crucial determinant in Hh protein membrane association,

mode of secretion and morphogenetic extension range (Callejo et al., 2011; Gallet et al., 2003;

Lewis et al., 2001; Porter et al., 1996). We find that Shh mutant lacking palmitoylation and

cholesterylation is secreted unpolarized. Therefore, the Shh peptidic structure either lacks

specific sorting signals for polarity or requires membrane association to expose sorting

competent proteinaceous signals. Our results also show that mutants lacking either

palmitoylation (ShhNpC24S) or cholesterylation (ShhN) predominantly distribute to the

basolateral cell surface, suggesting that both lipid modifications contribute to the basolateral

sorting of Shh. We could not attribute a particular role of palmitoylation in the basolateral-to-

apical trafficking because Shh defective in cholesterol modification is inefficiently palmitoylated

(Chen et al., 2004; Feng et al., 2004; Kohtz et al., 2001; Sbrogna et al., 2003). On the other

hand, our pulse chase and basolateral tagging experiments indicate that the cholesteryl group

can drive the entire polarized trafficking pathway of Shh, including the primary basolateral

sorting and subsequent transcytosis to the apical domain. It is possible that cholesterol

represents a sorting signal decoded at different steps of the Shh itinerary to the apical cell

surface, perhaps involving the trafficking role proposed for Disp (Callejo et al., 2011; D'Angelo

et al., 2015; Gore et al., 2021; Stewart et al., 2018), as a cholesterol binding protein (Cannac et

al., 2020; Chen et al., 2020).

Disp-1 is a multipass transmembrane protein with a luminal/extracellular sterol-sensing

domain that binds Shh in a cholesterol-dependent manner and is required for its release in a

diffusible form from the membrane (Burke et al., 1999; Cannac et al., 2020; Chen et al., 2020;

Creanga et al., 2012; Tukachinsky et al., 2012). In vertebrates, Disp-1 transfers Shh from the

cell surface to secreted SCUBE2 requiring the sodium gradient across the plasma membrane

(Burke et al., 1999; Cannac et al., 2020; Chen et al., 2020; Petrov et al., 2020). Stewart et al.

(Stewart et al., 2018) recently described that Disp-1 acquires functional competence after its

cleavage by the proprotein convertase furin at the cell surface (Stewart et al., 2018). Because

furin recycles between the TGN and the basolateral cell surface (Simmen et al., 1999), Disp-1

should reach first the basolateral cell surface to become competent in Hhs secretion in epithelial

cells (Stewart et al., 2018). In *Drosophila*, exogenously expressed Disp has been reported at

the plasma membrane of the apical, basal and lateral regions (D'Angelo et al., 2015), or mainly

at the basolateral plasma membrane (Callejo et al., 2011; Stewart et al., 2018). In MDCK cells,

transfection experiments showed Disp-1 distributed to the basolateral region seemingly due to

sorting information contained in its intracellular domain (Etheridge et al., 2010). We find with

RT-PCR that MDCK cells express Disp-1 (not shown) and our microinjection experiments

avoiding a saturating overexpression revealed recently synthesized Disp-1 predominantly

distributed at the subapical region, just underneath Shh, though also spreading throughout the

cell. These observations most likely reflect a dynamic Disp-1 trafficking and variations

depending on its expression levels and experimental conditions.

Our experiments of plasmid microinjection combined with live cell imaging show that

slightly overexpressed Disp-1 enhances basolateral secretion of newly expressed Shh just after

releasing a TGN block. Etheridge et al. (Etheridge et al., 2010) described a decreased

basolateral secretion of Shh under conditions of Disp-1 silencing or Disp-1 mutant expression in

MDCK cells. Accordingly, we find that coexpression of the cleavage defective Disp-CS mutant

(Stewart et al., 2018) decreases the basolateral secretion of Shh. A trade-off between apical

and basolateral Hh secretion has been previously proposed in the wing disk epithelia (Ayers et

al., 2010; Gore et al., 2021). Our results showing that basolateral secretion of Shh directly from

the TGN requires and can be enhanced by increasing the coexpressed level of Disp-1

suggesting that conditions that increase or decrease this process would determine how much

Hhs reaches the apical cell surface, where Disp is also expected to mediate Hhs secretion,

(D'Angelo et al., 2015).

Disp has been proposed to play a role in Hh subcellular distribution and trafficking

through endocytic pathways. This notion derives from studies in Drosophila epithelia models

where Hh secretion and signaling are sensitive to alterations in small GTPases such as Rab4,

Rab5, Rab8 and Rab11, which regulate endosomal compartments (Callejo et al., 2011;

D'Angelo et al., 2015; Gore et al., 2021; Pizette et al., 2021; Stewart et al., 2018). The

expression of Disp-1-CS previously suggested a requirement of Disp-1 cleavage in endosomal

membrane trafficking (Stewart et al., 2018). More recent evidence in *Drosophila* highlights a role

of Disp and Rab11 in Hh delivery from a tubular endosomal compartment named Hherisomes,

which modifies Hh signaling potential (Pizette et al., 2021). It is unknown whether this

compartment is functionally related to the Rab11-ARE compartment of MDCK cells (Perez Bay

et al., 2016). Our microinjection experiments show colocalization of newly synthesized Disp-1

and Shh together with Rab11 in a tubular subapical endosome corresponding to the ARE.

Furthermore, we show that Disp-1-CS not only decreases basolateral secretion from the TGN

but also accumulates Shh at the ARE, thus decreasing its apical cell surface distribution.

Therefore, our results add more evidence to support a role of Disp-1 in the vesicular trafficking

of Hhs, revealing a requirement for Shh trafficking along the basolateral-to-apical transcytosis

pathway at the level of the Rab11-ARE compartment.

The complex trafficking behavior of Shh in MDCK cells is compatible with the apical

and/or basolateral distribution and secretion of Hh seen in different epithelial cells, predicting

also a reciprocal basolateral versus apical secretion variations. In MDCK cells, the predominant

Shh apical location and secretion at steady state would result from low levels of Shh basolateral

secretion. Parietal cells in the stomach distribute and secrete Shh mostly apically, also showing

detectable levels at the basolateral side (Zavros et al., 2008). Basolateral secretion of Shh can

be inferred from stromal mesenchymal cell responses in intestinal cells (Buller et al., 2012;

Shyer et al., 2015) and kidney epithelium (Ding et al., 2012; Edeling et al., 2016; Zhou et al.,

2014), without discarding an apical secretion with other functions. Apical and basolateral

secretion of Shh sustaining distinct roles and signaling have been described in mature airway

epithelia (Mao et al., 2018). Here, immunostaining localizes Shh both at apical and lateral

regions and Shh basolateral release stimulates the stromal mesenchymal cells through primary

cilia, while its apical secretion induces non-canonical signaling via motile cilia (Mao et al., 2018).

In our proposed model the efficiency of Shh endocytic removal for transcytosis and/or secretion

from the basolateral domain would determine the relative distribution and secretion from each

domain.

Our results might also help to reconcile debated models in *Drosophila* epithelial

systems (Guerrero and Kornberg, 2014; Matusek et al., 2020). The relative magnitude of apical

and basolateral exocytic pathways operating in the long studied wing disc epithelia is unknown

(Guerrero and Kornberg, 2014; Matusek et al., 2020). Extensive studies in the wing disc

epithelia proposes a primary addressing of Hh towards the apical plasma membrane, even

though it is detected both in apical and basolateral cell surfaces (Callejo et al., 2011; D'Angelo

et al., 2015; Gradilla and Guerrero, 2013; Therond, 2012). A long controversy stands on

whether secretion for long-range Hh signaling occurs from the apical domain after endocytic

recycling (D'Angelo et al., 2015), or from the basolateral cell surface following apical-to-

basolateral transcytosis (Callejo et al., 2011). However, previous studies suggested a primary

Hh basolateral addressing followed by transcytosis to the apical domain instead, which might

explain the enhanced basolateral Hh distribution seen when Disp is silenced in other embryo

epithelial cells (Gallet et al., 2003). Basolateral Hh accumulation and secretion has also been

found under conditions that silenced the small GTPase Rab8, known to be involved in endocytic

and basolateral trafficking (Gore et al., 2021). These controversial views reflect the difficulties of

attributing a trafficking pathway based on the interpretation of locations and cell target

responses under steady-state conditions. As we show in MDCK cells, a predominant steady-

state apical cell surface distribution can result from a primary basolateral sorting from the TGN

followed by transcytosis. Our results also demonstrate that transcytosis of Shh still bound to the

membrane occurs not only from the basolateral but also from the apical domain. Although we

did not measure apical recycling after apical endocytosis, as described in the wing disk epithelia

(D'Angelo et al., 2015), this pathway might indeed operate together with transcytosis. Therefore,

our results are compatible with the observations in *Drosophila* epithelial systems and perhaps

can be used to reinterpret conflicting Hh trafficking models.

Hh proteins have been found secreted in different ways, including multimers (Gallet et

al., 2006; Zeng et al., 2001), lipoprotein particles (Panakova et al., 2005), extracellular vesicles

(Gradilla et al., 2014; Matusek et al., 2014; Parchure et al., 2015; Tanaka et al., 2005; Vyas et

al., 2014) and associated with cytonemes that are long filopodia (Bischoff et al., 2013; Chen et

al., 2017; Gonzalez-Mendez et al., 2017). How a primary basolateral sorting followed by

transcytosis relates with particular forms of Shh secretion remains a challenge for future

studies. Hh proteins form nano-scaled oligomers via electrostatic interactions and this process

is required for endocytosis followed by targeting to multivesicular bodies and secretion in

exosomes (Parchure et al., 2015; Vyas et al., 2008; Vyas et al., 2014). The role of this Shh

oligomerization in polarized sorting can now be evaluated in future experiments using the

proposed model in MDCK cells.

An indirect trafficking towards the apical cell surface includes several intermediate

stages where regulation might occur impacting upon Shh polarized secretion. This would allow

for reciprocal basolateral versus apical secretion variations in distinct epithelial contexts.

Conditions that increase basolateral secretion would decrease apical secretion. Variations in

Disp-1 expression will first impact the Shh basolateral secretion and then its transcytosis-

dependent apical secretion. This, together with the regulation of endocytic trafficking through

Rab GTPases, can contribute to determine variations in the basolateral versus apical secretion

and the signaling extension of this powerful regulatory molecule in morphogenesis and cancer.

MATERIALS AND METHODS
Experimental procedures
514
515 Key resources table

Reagent type (species) or resource	Designation	Source or reference	Identifiers	Additional information
Cell line (cannis familiaris)	MDCK (Madin-Darby canine kidney) II cells	ATCC provided by Enrique Rodriguez-Boulan (Weill Cornell Medical College, New York, NY)		
Culture media	DMEM	Thermo Fisher	12100046	
Culture media	DMEM methionine and cystine free	Thermo Fisher	21013024	for metabolic label
Culture media	DEMEM-HEPES	Thermo Fisher	12430054	for microinjection and live cell imaging
Transfected/ microinjected construct (M. musculus)	pCDNA3.1-Shh	This paper		Subcloned from pBluescript-mShh (P. A. Beachy) (Development 1994)
Transfected construct (M. musculus)	pCDNA3.1-ShhNC24S	This paper		Subcloned from pRK5-mShhNC24S (P. A. Beachy) (J Cell Sci 1999)
Transfected construct (M. musculus)	pCDNA3.1-ShhN	This paper		PCR cloning from pCDNA3.1-Shh
Transfected construct (M. musculus)	pCDNA3.1-ShhNpC24S	This paper		Subcloned from pSK-C24S (P. A. Beachy)
Microinjected construct (M. musculus)	pCDNA3-Shh-GFP	McMahon Lab (Chamberlain 2008)		
Cell line (cannis familiaris)	MDCK-Rab11-mCherry	This paper		Stably expressing human Rab11a-mCherry
Cell line (cannis familiaris)	MDCK-GFP-NO-GPI	Imjeti 2011		Stably expressing GFP-NO-GPI
Microinjected construct (M. musculus)	pCDNA3 V5-Dispatched-HA	Ogden Lab (Stewart 2018)		pCDNA3 from Invitrogen
Microinjected construct (M. musculus)	pCDNA3 V5-Dispatched-CS-HA	Ogden Lab (Stewart 2018)		pCDNA3 from Invitrogen
Microinjected construct (M. musculus)	LDLR-GFP-Y18A	Matter 1992		(pCB6, Hunziker 1991) LDLR-GFP-Y18A
Antibody	Anti-Shh	DSHB (Developmental Studies Hybridoma Bank)	5E1	IF (1:50) IP (1:50)
Antibody	Anti-Shh	Santa Cruz Biotechnology, Inc. (California, USA)	sc-9024 (H-160)	WB (1:1000)
Antibody	Anti-E-cadherin	Cell signaling	24E10 mAb #3195	IF (1:100)
Antibody	Anti-E-cadherin	BD transduction		IP (2mg/sample)

		laboratories		
Antibody	Anti-GFP (rabbit)	This paper		IP (1:250)
Antibody	Anti-Rab11a (rabbit)	Thermo Fisher Scientific	715300	IF (1:100)
Antibody	Anti-Fibronectin	Sigma	F6140	WB (1:2000)
Antibody	Anti-Na$^+$/K$^+$-ATPase	Santa Cruz Biotechnology, Inc. (California, USA)	sc-21713	WB (1:2000)
Antibody	Anti-HA (chicken)	Millipore	AB3254	IF (1:100)
Antibody	AlexaFluor 488	Life technologies, Carlsbad, CA	A11029 (Mouse) A11034 (Rabbit) A-11039 (Chicken)	1:500
Antibody	AlexaFluor 555	Life technologies	A21424 (Mouse) A21429 (Rabbit) A32932 (Chicken)	1:500
Antibody	AlexaFluor 647	Life technologies	A21235 (Mouse)	1:500
Fluorescent dye	Hoechst	Thermo Fisher Scientific	62249	1:1000
Software, algorithm	Adobe Photoshop CS4	Adobe, San Jose, CA		for making figures
Software, algorithm	LAS X	Leica, Wetzlar, Germany		image analysis
Software, algorithm	Prism	GraphPad, La Jolla, CA		for statistical, analysis and graphs
Software, algorithm	SigmaPlot10 software			for quantification of golgi retention
Software, algorithm	Huygens Professional software			for deconvolution images
Software, algorithm	ImageJ (NIH)		2.0.0-rc-69/1.52o	image analysis
Software, algorithm	OptiQuant software	PerkinElmer	3.0	
Commercial assay or kit	Lipofectamine 2000	Thermo Fisher Scientific	11668027	
Commercial assay or kit	Pierce™ ECL Western Blotting Substrate	Pierce™, Waltham, MA, USA	32106	
Chemical compound, drug	Sulfo-NHS-LC-biotin	Pierce™ /Thermo Fisher Scientific	21335	
Chemical compound, drug	EZ-Link™ NHS-SS-Biotin	Pierce/Thermo Fisher Scientific	21441	
Chemical compound, drug	NeutrAvidin-agarose	Pierce/Thermo Fisher Scientific	29201	
Chemical compound, drug	MesNa	Sigma	PHR1570	
Chemical compound, drug	^{35}S-methionine/cysteine	Perkin Elmer NEN Life Science Products (Boston, USA)	NEG709A005MC	
Chemical compound, drug	Fluoromount-G	Thermo Fisher Scientific	00495802	
Polarized culture	24mm Transwells	Costar	3412	for biochemistry

chamber				
Polarized culture chamber	12mm Transwells	Costar	3401	for immunofluorescence

516
Cloning and Subcloning

cDNA coding for Shh (from pBlueScript-Shh), ShhNpC24S (from pSK-C24S) and
ShhNC24S (from pRK5-ShhNC24S) (kindly provided by Philip A. Beachy, Stanford University,
CA, USA) were subcloned by double digestion with EcoRI and XbaI (for Shh), NotI (for
ShhNpC24S) or NotI and KpnI (for ShhNC24S) restriction enzymes and ligated into pcDNA3.1(-
) with T4 DNA ligase (Thermo Fisher cat #15224017) according to the manufacturer's
instructions. ShhN dsDNA coding for residues 1-198 of murine Shh was obtained by PCR using
Pfu polymerase (Promega cat # M7741) and the following primers: Fw:
GTCTCGAGCCAACTCCGATGTGTTCCG and Rev: CGGAATTCTTAGCCGCCGGATTTGGCC
that include XhoI and EcoRI sites for subcloning into pcDNA3.1(+). All constructs were
confirmed by sequencing.

528
Cell Lines

MDCK cells were grown in DMEM (Cat# 12100046, Thermo Fisher) supplemented with
7.5% fetal bovine serum (FBS, Cat# SH30071.03 HyClone), 50 U/mL antibiotics (Penicillin-
Streptomycin mix (Cat# 15070063, Thermo Fisher), and 5 µg/ml Plasmocin (Cat# ant-mpp,
InvivoGen), in a 37ºC, 5% CO_2 humidified incubator, routinely analyzed by RT-PCR for
mycoplasma contamination. Passages were made before reaching confluence every 2-3 days
by two washes with PBS (15 min each) and incubation in trypsin 0.05% for 5-10 min (Trypsin
Cat#15400054, Thermo Fisher). MDCK cells stable transfected clones (MDCK-Shh, -ShhN, -
ShhNC24S, -ShhNpC24S) were generated by transfection with Lipofectamine 2000 reagent and
selection by G418 resistance.

539
Transfection

MDCK (100.000 cells) grown in 24-multiwell plates were transfected with 1µg of plasmid
DNA and Lipofectamine 2000 reagent, trypsinized the next day and plated at 1/10 and 1/100
dilution into 100mm dishes. Selection media containing G418 (0.8mg/ml) was added the next
day, changed every 5 days until visible colonies were isolated using cloning cylinders and
expanded in 6-multiwell plates. Screening was made by immunofluorescence (see below) and
immunoblot of lysates and media (See "Protein Analysis").

547
Cell lysates and protein analysis

To obtain lysates cells were washed twice in PBS containing 1 mM MgCl2, 0.1mM
CaCl2 (PBS-CM) and lysed in 500 µl RIPA buffer (50 mM Tris-HCl (pH 8.0), 150 mM NaCl,
0.5% Na-deoxycholate, 0.1% SDS, 1% Triton X-100) supplemented with protease inhibitors
(PMSF 2mM, Leupeptin 2µg/ml, Pepstatin 1µg/ml). Recovered lysates were cleared by

centrifugation at 14.000rpm at 4ºC. Protein concentration was determined on the cleared cell
lysates with BCA method and 500 µg total protein was used for immunoprecipitation or
NeutrAvidin-precipitation (see below), 10% volume of each lysate was used as loading control.
556

<u>Steady-state Shh Secretion</u>
Analysis of secreted proteins was performed by collection of conditioned media in low
serum media (0.3% FBS), precipitation in 10% trichloroacetic acid (TCA) and incubation for
30min on ice. Precipitated proteins were centrifuged at 4ºC and 14.000 rpm for 10 min, washed
once with 1ml of 100% acetone and re-centrifuged. Pellets were solubilized in 30µl of NaOH
0.1M and 10µl of 3X Laemmli sample buffer. Lysates and/or secreted proteins were resolved in
12%SDS-PAGE, transferred to PVDF membranes and immunoblotted with the indicated
antibodies.
565

566   <u>Polarized monolayers of MDCK cells grown in Transwell inserts</u>
567   MDCK cells (100,000-400,000 cells) were plated on 12 mm or 24-mm Transwell inserts,
568   0.4-mm pore size (Costar), and grown with daily media changes until the
569   transepithelial resistance reached 400 ohm/cm^2 (4–5 days) measured with an EVOM
570   electrometer (World Precision Instruments, Sarasota, FL).
571

<u>Immunofluorescence</u>
Cells were fixed in freshly prepared 4% PFA in PBS-CM for 15 min at room temperature
or overnight at 4°C. To analyze subcellular distribution of interest proteins, cells were
permeabilized with 0.2% saponin in PBS-CM. All subsequent steps contained 0.2% saponin.
Primary and secondary antibodies were incubated for 30 min at 37°C or overnight at 4°C.
Coverslips were mounted in Fluoromount.
578

<u>Protein immunoprecipitation</u>
The key resources table indicates the amount of antibodies (GFP, E-cadherin, 5E1)
used per sample. Antibodies were pre-incubated with 30µl of protein-A sepharose for 2h at 4°C,
then beads were washed 3 times to remove excess antibodies. Total cell lysate (approx. 500
µg) or collected medium was then added to beads and incubated for 2h at 4ºC, washed 3 times
in RIPA buffer and eluted in 30µl of 2X Laemmli Buffer.
585

<u>Domain-Selective Surface Biotinylation</u>
Steady-state apical and basolateral distribution of Shh and endogenous proteins were
analyzed by domain selective biotinylation of MDCK cells grown in Transwell inserts, as
described (Bravo-Zehnder et al., 2000; Oyanadel et al., 2018). Cells were rinsed twice with ice-
cold PBS-CM followed by two successive 20 min incubations at 4ºC with 0.5 mg/ml Sulfo-NHS-

LC-Biotin in PBS-CM, added to the apical (0.6 ml) or basolateral (1 ml) sides. Cells were then

incubated with 50mM NH_4Cl in PBS-CM for 10 min, rinsed twice with PBS-CM and lysed in

RIPA buffer, supplemented with proteases inhibitors for 30 min at 4ºC. NeutrAvidin-Sepharose

was used to retrieve biotinylated proteins by centrifugation (18,000 x g for 10 min), which then

were analyzed by immunoblot against Shh and endogenous proteins, as described (Bravo-

Zehnder et al., 2000; Oyanadel et al., 2018; Sargiacomo et al., 1989).

598   Pulse-chase domain-selective targeting of newly synthesized proteins

599   Biotinylation targeting assay was used to assess biosynthetic delivery of proteins to

apical and basolateral PM domains. Stable transfected cells were pulse-labeled in methionine-

free and cysteine-free DMEM supplemented with 400 µCi/ml [^{35}S]-methionine/cysteine for 30

min in 120µl of medium applied to the basolateral side of the inverted filter, as described

(Marzolo et al., 1997). Chase was made for different times in culture media containing 300 µM

cycloheximide (CHX) and 3- to 6-fold excess cold methionine (0.6mM) and cysteine and

(1.2mM). Cells were then subjected to domain-selective surface biotinylation, chilled in ice-cold

PBS-CM and lysed with RIPA buffer supplemented with protease inhibitors at 4ºC for 30 min.

Lysates were cleared by centrifugation (18,000 x g for 10 min) before immunoprecipitation with

protein-A-Sepharose preincubated antibodies (5E1-anti-Shh or E-cadherin). The mixture of

biotinylated (surface) plus non-biotinylated (intracellular) immunoprecipitated proteins was

incubated at 95º for 5 min in 30 µl of 10% SDS RIPA to dissociate immune complexes. The

eluted proteins were diluted to 1 ml with RIPA and subjected to NeutrAvidin-Sepharose

precipitation to isolate surface-located Shh. In addition, secreted Shh was immunoprecipitated

from the apical and basolateral media with protein-A-Sepharose bound 5E1, as described for

other proteins (Gonzalez et al., 1993; Gonzalez et al., 1987; Marzolo et al., 1997). The immune

complexes of surface-located and secreted Shh were resolved by SDS-PAGE. [^{35}S]-

methionine/cysteine labeled proteins were visualized in fluorograms made with Amplify^{TM}

fluorography reagent (Amersham), developed in scanner Cyclone Plus (Perkin Elmer) with

OptiQuant software (3.0, Perkin Elmer).

620   Transcytosis assays

621   Trafficking of Shh from one cell surface domain to the opposite (transcytosis) was

622   analyzed by domain-specific biotinylation and cell surface-tagging immunofluorescence assays.

623   *1. Biotinylation transcytosis assay* was performed with reducible biotin reagent Sulfo-

NHS-SS-biotin as described (Burgos et al., 2004). In brief, six Transwell filter cups with

polarized live MDCK monolayer cells were subjected to basolateral-specific biotinylation with

Sulfo-NHS-SS-biotin in *PBS-CM.* One filtercup was used for total basolateral biotinylated Shh

("non-reduced, time 0"), and other two filtercups were subjected to either apical or basolateral

reduction with 22 mM MesNa reagent (0 min) made in buffer 50 mM Tris-HCl, pH 8.6, 100 mM

NaCl. The remaining three filtercups were incubated for 120 min in 37ºC in growth media to
allow trafficking and then MesNa was added to the apical or basolateral side. Monolayers were
alkylated with 20mM iodoacetamide in PBS-CM containing 1%BSA, washed twice in PBS-CM,
lysed in Ripa Buffer supplemented with protease inhibitors and subjected to NeutrAvidin-
Sepharose precipitation and immunoblot. To analyze apical secretion of Shh present in the
basolateral membrane, cells were biotinylated from the basolateral side and then incubated in
DMEM, 0.5 % fetal bovine serum at 37ºC for 2 hours. Biotinylated Shh proteins were retrieved
from apical and basolateral media with NeutrAvidin-Sepharose and immunoblotted.

2.   Cell surface-tagging immunofluorescence transcytosis assay. Live MDCK cells
grown in Transwell inserts were apically or basolaterally incubated with 5E1 antibody diluted in
DMEM (supplemented with 10mM HEPES) for 30 min at 4°C. After 3 washes in PBS-CM, cells
were shifted to 37°C in growth media to allow trafficking for different times, fixed at different
times with PFA-PBS-CM and permeabilized. Shh bound to the primary antibody was detected
by incubation of fluorophore-coupled secondary antibodies for 30 min at 37°C.

3.   Secretion of basolaterally-sorted ShhNpC24S. MDCK cells grown in Transwell
inserts were rinsed twice with ice-cold PBS-CM followed by two successive 20 min incubations
at 4ºC with 0.5 mg/ml Sulfo-NHS-LC-Biotin in PBS-CM, added to the basolateral sides. Cells
were then incubated with 50mM NH_4Cl in PBS-CM for 5 min, rinsed twice with PBS-CM and
shifted to 37°C in growth media for 2h. Then apical and basolateral media was collected and
precipitated with Neutravidin (basolaterally-biotinylated secreted proteins). Samples were
analyzed by western blot.

Microinjection

MDCK cells were plated either in 12mm diameter glass coverslips (5000 cells) the
absence of Plasmocin. After achieving full confluence (4-5 cells) microinjections were performed
in the cell nucleus (plasmids) using back-loaded glass capillaries, an Eppendorf NI2
micromanipulator and an Eppendorf Transjector 5246 system. The expression plasmids DNAs
were used at a concentration of 20 to 100ng/µl in buffer HEPES 10 mM pH7.4, KCl 140 mM.
During the microinjection procedure cells were maintained in DMEM-HEPES.

Live cell imaging

The microinjection approach to express and accumulate exogenous proteins at the
TGN and then assess subsequent trafficking was performed as described (Cancino et al., 2007;
Kreitzer et al., 2000). Briefly, microinjected cells were incubated for 45 min at 37°C in DMEM-
HEPES plus 7.5% FSB to allow protein expression without ER leakage, plasmid concentration
was adjusted (20 to 100ng/µl) to obtain enough protein expression to be detected by
fluorescence microscopy and then the exocytic trafficking was arrested at the TGN by

incubating the cells at 20°C for 2 h in DMEM-HEPES with CHX. After the 20°C block the cells

were mounted on a Leica SP8 microscope in DMEM-HEPES plus CXM and incubated at 37°C.

Images acquisition and processing

Images were collected at 1024 × 1024 resolution in Z stacks of 300-nm steps using a

Leica SP8 confocal microscope and a 63× oil immersion 1.4 ma lens, and processed with

ImageJ free software, NIH. High Resolution images were acquired at Nyquist rate parameters

(Nyquist Calculator, SVI, NL) and the resulting oversampled images were deconvoluted with

Huygens Essential Software (SVI, NL), which was also used for 3D surface rendering

processing.

Statistical Analysis

Significance was determined with Prism software using Student's t-test. For all

statistical analyses **$p < 0.05$ and ****$p < 0.0001$. Error bars indicate standard error of mean

(SEM).

ACKNOWLEDGEMENTS
We are grateful to Dr. Stacey Ogden (St. Jude Children's Research Hospital, USA) for
kindly providing Dispatched plasmids constructs, Dr. Andrew P. McMahon (University
of Southern California, USA) and Dr. Philip A. Beachy (Stanford University School of
Medicine, USA) for kindly providing Shh plasmids constructs.
687

688 **ADDITIONAL INFORMATION**

689 **Funding**

Funder	Grant reference number	Author
Fondo Nacional de Desarrollo Científico y Tecnológico (FONDECYT)	1181907	Alfonso González
Programa de Apoyo a Centros con Financiamiento Basal. Agencia Nacional de Investigación y Desarrollo (ANID)	#AFB170004; #AFB170005; #ACE210009	Alfonso González
Postdoctoral Project (FONDECYT)	3150603	Lisette Sandoval

AUTHOR CONTRIBUTIONS
Lisette Sandoval, Conceptualization, Data curation, Formal analysis, Validation,
Investigation, Visualization, Methodology, Writing - original draft, Writing - review and
editing; Mariana Labarca, Conceptualization, Formal analysis, Validation, Investigation,
Visualization, Methodology, Writing - review and editing; Claudio Retamal, Data
curation, Formal analysis, Methodology; Juan Larraín, Conceptualization, Resources;
Alfonso González, Conceptualization, Supervision, Funding acquisition, Writing
698

REFERENCES

Apodaca, G., Katz, L.A., and Mostov, K.E. (1994). Receptor-mediated transcytosis of IgA in MDCK cells is via apical recycling endosomes. J Cell Biol *125*, 67-86.

Arkhipenko, A., Syan, S., Victoria, G.S., Lebreton, S., and Zurzolo, C. (2016). PrPC Undergoes Basal to Apical Transcytosis in Polarized Epithelial MDCK Cells. PLoS One *11*, e0157991.

Ayers, K.L., Gallet, A., Staccini-Lavenant, L., and Therond, P.P. (2010). The long-range activity of Hedgehog is regulated in the apical extracellular space by the glypican Dally and the hydrolase Notum. Developmental cell *18*, 605-620.

Bartles, J.R., Feracci, H.M., Stieger, B., and Hubbard, A.L. (1987). Biogenesis of the rat hepatocyte plasma membrane in vivo: comparison of the pathways taken by apical and basolateral proteins using subcellular fractionation. J Cell Biol *105*, 1241-1251.

Beachy, P.A., Karhadkar, S.S., and Berman, D.M. (2004). Tissue repair and stem cell renewal in carcinogenesis. Nature *432*, 324-331.

Bischoff, M., Gradilla, A.C., Seijo, I., Andres, G., Rodriguez-Navas, C., Gonzalez-Mendez, L., and Guerrero, I. (2013). Cytonemes are required for the establishment of a normal Hedgehog morphogen gradient in Drosophila epithelia. Nat Cell Biol *15*, 1269-1281.

Bonifacino, J.S. (2014). Adaptor proteins involved in polarized sorting. J Cell Biol *204*, 7-17.

Bravo-Zehnder, M., Orio, P., Norambuena, A., Wallner, M., Meera, P., Toro, L., Latorre, R., and Gonzalez, A. (2000). Apical sorting of a voltage- and Ca2+-activated K+ channel alpha -subunit in Madin-Darby canine kidney cells is independent of N-glycosylation. Proc Natl Acad Sci U S A *97*, 13114-13119.

Briscoe, J., and Therond, P.P. (2013). The mechanisms of Hedgehog signalling and its roles in development and disease. Nature reviews Molecular cell biology *14*, 416-429.

Brown, P.S., Wang, E., Aroeti, B., Chapin, S.J., Mostov, K.E., and Dunn, K.W. (2000). Definition of distinct compartments in polarized Madin-Darby canine kidney (MDCK) cells for membrane-volume sorting, polarized sorting and apical recycling. Traffic *1*, 124-140.

Buller, N.V., Rosekrans, S.L., Westerlund, J., and van den Brink, G.R. (2012). Hedgehog signaling and maintenance of homeostasis in the intestinal epithelium. Physiology (Bethesda) *27*, 148-155.

Burgos, P.V., Klattenhoff, C., de la Fuente, E., Rigotti, A., and Gonzalez, A. (2004). Cholesterol depletion induces PKA-mediated basolateral-to-apical transcytosis of the scavenger receptor class B type I in MDCK cells. Proc Natl Acad Sci U S A *101*, 3845-3850.

Burke, R., Nellen, D., Bellotto, M., Hafen, E., Senti, K.A., Dickson, B.J., and Basler, K. (1999). Dispatched, a novel sterol-sensing domain protein dedicated to the release of cholesterol-modified hedgehog from signaling cells. Cell *99*, 803-815.

Callejo, A., Bilioni, A., Mollica, E., Gorfinkiel, N., Andres, G., Ibanez, C., Torroja, C., Doglio, L., Sierra, J., and Guerrero, I. (2011). Dispatched mediates Hedgehog basolateral release to form the long-range morphogenetic gradient in the Drosophila wing disk epithelium. Proc Natl Acad Sci U S A *108*, 12591-12598.

Cancino, J., Torrealba, C., Soza, A., Yuseff, M.I., Gravotta, D., Henklein, P., Rodriguez-Boulan, E., and Gonzalez, A. (2007). Antibody to AP1B adaptor blocks biosynthetic and recycling routes of basolateral proteins at recycling endosomes. Mol Biol Cell *18*, 4872-4884.

Cannac, F., Qi, C., Falschlunger, J., Hausmann, G., Basler, K., and Korkhov, V.M. (2020). Cryo-EM structure of the Hedgehog release protein Dispatched. Sci Adv *6*, eaay7928.

Caspary, T., Garcia-Garcia, M.J., Huangfu, D., Eggenschwiler, J.T., Wyler, M.R., Rakeman, A.S., Alcorn, H.L., and Anderson, K.V. (2002). Mouse Dispatched homolog1 is required for long-range, but not juxtacrine, Hh signaling. Curr Biol *12*, 1628-1632.

Chamberlain, C.E., Jeong, J., Guo, C., Allen, B.L., and McMahon, A.P. (2008). Notochord-derived Shh
concentrates in close association with the apically positioned basal body in neural target cells and forms a
dynamic gradient during neural patterning. Development *135*, 1097-1106.

Chen, H., Liu, Y., and Li, X. (2020). Structure of human Dispatched-1 provides insights into Hedgehog
ligand biogenesis. Life Sci Alliance *3*.

Chen, M.H., Li, Y.J., Kawakami, T., Xu, S.M., and Chuang, P.T. (2004). Palmitoylation is required for the
production of a soluble multimeric Hedgehog protein complex and long-range signaling in vertebrates.
Genes Dev *18*, 641-659.

Chen, W., Huang, H., Hatori, R., and Kornberg, T.B. (2017). Essential basal cytonemes take up Hedgehog
in the Drosophila wing imaginal disc. Development *144*, 3134-3144.

Chen, X., Tukachinsky, H., Huang, C.H., Jao, C., Chu, Y.R., Tang, H.Y., Mueller, B., Schulman, S.,
Rapoport, T.A., and Salic, A. (2011). Processing and turnover of the Hedgehog protein in the endoplasmic
reticulum. J Cell Biol *192*, 825-838.

Creanga, A., Glenn, T.D., Mann, R.K., Saunders, A.M., Talbot, W.S., and Beachy, P.A. (2012). Scube/You
activity mediates release of dually lipid-modified Hedgehog signal in soluble form. Genes Dev *26*, 1312-
1325.

D'Angelo, G., Matusek, T., Pizette, S., and Therond, P.P. (2015). Endocytosis of Hedgehog through
dispatched regulates long-range signaling. Developmental cell *32*, 290-303.

Danesin, C., Agius, E., Escalas, N., Ai, X., Emerson, C., Cochard, P., and Soula, C. (2006). Ventral neural
progenitors switch toward an oligodendroglial fate in response to increased Sonic hedgehog (Shh) activity:
involvement of Sulfatase 1 in modulating Shh signaling in the ventral spinal cord. J Neurosci *26*, 5037-
5048.

Ding, H., Zhou, D., Hao, S., Zhou, L., He, W., Nie, J., Hou, F.F., and Liu, Y. (2012). Sonic hedgehog
signaling mediates epithelial-mesenchymal communication and promotes renal fibrosis. J Am Soc Nephrol
23, 801-813.

Edeling, M., Ragi, G., Huang, S., Pavenstadt, H., and Susztak, K. (2016). Developmental signalling
pathways in renal fibrosis: the roles of Notch, Wnt and Hedgehog. Nat Rev Nephrol *12*, 426-439.

Etheridge, L.A., Crawford, T.Q., Zhang, S., and Roelink, H. (2010). Evidence for a role of vertebrate Disp1
in long-range Shh signaling. Development *137*, 133-140.

Feng, J., White, B., Tyurina, O.V., Guner, B., Larson, T., Lee, H.Y., Karlstrom, R.O., and Kohtz, J.D.
(2004). Synergistic and antagonistic roles of the Sonic hedgehog N- and C-terminal lipids. Development
131, 4357-4370.

Gallet, A. (2011). Hedgehog morphogen: from secretion to reception. Trends Cell Biol *21*, 238-246.

Gallet, A., Rodriguez, R., Ruel, L., and Therond, P.P. (2003). Cholesterol modification of hedgehog is
required for trafficking and movement, revealing an asymmetric cellular response to hedgehog.
Developmental cell *4*, 191-204.

Gallet, A., Ruel, L., Staccini-Lavenant, L., and Therond, P.P. (2006). Cholesterol modification is necessary
for controlled planar long-range activity of Hedgehog in Drosophila epithelia. Development *133*, 407-418.

Gan, Y., McGraw, T.E., and Rodriguez-Boulan, E. (2002). The epithelial-specific adaptor AP1B mediates
post-endocytic recycling to the basolateral membrane. Nat Cell Biol *4*, 605-609.

Gonzalez, A., Nicovani, S., and Juica, F. (1993). Apical secretion of hepatitis B surface antigen from
transfected Madin- Darby canine kidney cells. J Biol Chem *268*, 6662-6667.

Gonzalez, A., Rizzolo, L., Rindler, M., Adesnik, M., Sabatini, D.D., and Gottlieb, T. (1987). Nonpolarized
secretion of truncated forms of the influenza hemagglutinin and the vesicular stomatitis virus G protein
from MDCK cells. Proc Natl Acad Sci U S A *84*, 3738-3742.

Gonzalez, A., and Rodriguez-Boulan, E. (2009). Clathrin and AP1B: key roles in basolateral trafficking
through trans-endosomal routes. FEBS Lett *583*, 3784-3795.

Gonzalez-Mendez, L., Seijo-Barandiaran, I., and Guerrero, I. (2017). Cytoneme-mediated cell-cell contacts
for Hedgehog reception. Elife *6*.

Gore, T., Matusek, T., D'Angelo, G., Giordano, C., Tognacci, T., Lavenant-Staccini, L., Rabouille, C., and
Therond, P.P. (2021). The GTPase Rab8 differentially controls the long- and short-range activity of the
Hedgehog morphogen gradient by regulating Hedgehog apico-basal distribution. Development *148*.

Gottlieb, T.A., Beaudry, G., Rizzolo, L., Colman, A., Rindler, M., Adesnik, M., and Sabatini, D.D. (1986).
Secretion of endogenous and exogenous proteins from polarized MDCK cell monolayers. Proc Natl Acad
Sci U S A *83*, 2100-2104.

Gradilla, A.C., Gonzalez, E., Seijo, I., Andres, G., Bischoff, M., Gonzalez-Mendez, L., Sanchez, V., Callejo,
A., Ibanez, C., Guerra, M.*, et al.* (2014). Exosomes as Hedgehog carriers in cytoneme-mediated transport
and secretion. Nature communications *5*, 5649.

Gradilla, A.C., and Guerrero, I. (2013). Hedgehog on the move: a precise spatial control of Hedgehog
dispersion shapes the gradient. Curr Opin Genet Dev *23*, 363-373.

Gravotta, D., Deora, A., Perret, E., Oyanadel, C., Soza, A., Schreiner, R., Gonzalez, A., and Rodriguez-
Boulan, E. (2007). AP1B sorts basolateral proteins in recycling and biosynthetic routes of MDCK cells.
Proc Natl Acad Sci U S A *104*, 1564-1569.

Guerrero, I., and Kornberg, T.B. (2014). Hedgehog and its circuitous journey from producing to target cells.
Seminars in cell & developmental biology *33*, 52-62.

Guo, X., Mattera, R., Ren, X., Chen, Y., Retamal, C., Gonzalez, A., and Bonifacino, J.S. (2013). The
adaptor protein-1 mu1B subunit expands the repertoire of basolateral sorting signal recognition in epithelial
cells. Developmental cell *27*, 353-366.

Hall, E.T., Cleverdon, E.R., and Ogden, S.K. (2019). Dispatching Sonic Hedgehog: Molecular Mechanisms
Controlling Deployment. Trends Cell Biol *29*, 385-395.

Hauri, H.P., and Matter, K. (1991). Protein traffic in intestinal epithelial cells. Semin Cell Biol *2*, 355-364.

Hua, W., Sheff, D., Toomre, D., and Mellman, I. (2006). Vectorial insertion of apical and basolateral
membrane proteins in polarized epithelial cells revealed by quantitative 3D live cell imaging. J Cell Biol
172, 1035-1044.

Hubbard, A.L. (1989). Endocytosis. Curr Opin Cell Biol *1*, 675-683.

Imjeti, N.S., Lebreton, S., Paladino, S., de la Fuente, E., Gonzalez, A., and Zurzolo, C. (2011). N-
Glycosylation instead of cholesterol mediates oligomerization and apical sorting of GPI-APs in FRT cells.
Mol Biol Cell *22*, 4621-4634.

Kohtz, J.D., Lee, H.Y., Gaiano, N., Segal, J., Ng, E., Larson, T., Baker, D.P., Garber, E.A., Williams, K.P.,
and Fishell, G. (2001). N-terminal fatty-acylation of sonic hedgehog enhances the induction of rodent
ventral forebrain neurons. Development *128*, 2351-2363.

Kreitzer, G., Marmorstein, A., Okamoto, P., Vallee, R., and Rodriguez-Boulan, E. (2000). Kinesin and
dynamin are required for post-Golgi transport of a plasma- membrane protein. Nat Cell Biol *2*, 125-127.

Kreitzer, G., Schmoranzer, J., Low, S.H., Li, X., Gan, Y., Weimbs, T., Simon, S.M., and Rodriguez-Boulan,
E. (2003). Three-dimensional analysis of post-Golgi carrier exocytosis in epithelial cells. Nat Cell Biol *5*,
126-136.

Le Bivic, A., Real, F.X., and Rodriguez-Boulan, E. (1989). Vectorial targeting of apical and basolateral
plasma membrane proteins in a human adenocarcinoma epithelial cell line. Proc Natl Acad Sci U S A *86*,
9313-9317.

Lebreton, S., Paladino, S., Liu, D., Nitti, M., von Blume, J., Pinton, P., and Zurzolo, C. (2021). Calcium
levels in the Golgi complex regulate clustering and apical sorting of GPI-APs in polarized epithelial cells.
Proc Natl Acad Sci U S A *118*.

Lebreton, S., Paladino, S., and Zurzolo, C. (2019). Clustering in the Golgi apparatus governs sorting and
function of GPI-APs in polarized epithelial cells. FEBS Lett *593*, 2351-2365.

Lehmann, G.L., Hanke-Gogokhia, C., Hu, Y., Bareja, R., Salfati, Z., Ginsberg, M., Nolan, D.J., Mendez-
Huergo, S.P., Dalotto-Moreno, T., Wojcinski, A.*, et al.* (2020). Single-cell profiling reveals an endothelium-
mediated immunomodulatory pathway in the eye choroid. J Exp Med *217*.

Lewis, P.M., Dunn, M.P., McMahon, J.A., Logan, M., Martin, J.F., St-Jacques, B., and McMahon, A.P.
(2001). Cholesterol modification of sonic hedgehog is required for long-range signaling activity and
effective modulation of signaling by Ptc1. Cell *105*, 599-612.

Lisanti, M.P., Sargiacomo, M., Graeve, L., Saltiel, A.R., and Rodriguez-Boulan, E. (1988). Polarized apical
distribution of glycosyl-phosphatidylinositol-anchored proteins in a renal epithelial cell line. Proc Natl Acad
Sci U S A *85*, 9557-9561.

Long, J., Tokhunts, R., Old, W.M., Houel, S., Rodgriguez-Blanco, J., Singh, S., Schilling, N., Capobianco,
A.J., Ahn, N.G., and Robbins, D.J. (2015). Identification of a family of fatty-acid-speciated sonic hedgehog
proteins, whose members display differential biological properties. Cell Rep *10*, 1280-1287.

Luton, F., Hexham, M.J., Zhang, M., and Mostov, K.E. (2009). Identification of a cytoplasmic signal for
apical transcytosis. Traffic *10*, 1128-1142.

Machado, M.V., and Diehl, A.M. (2017). Hedgehog signalling in liver pathophysiology. J Hepatol.

Mao, S., Shah, A.S., Moninger, T.O., Ostedgaard, L.S., Lu, L., Tang, X.X., Thornell, I.M., Reznikov, L.R.,
Ernst, S.E., Karp, P.H.*, et al.* (2018). Motile cilia of human airway epithelia contain hedgehog signaling
components that mediate noncanonical hedgehog signaling. Proc Natl Acad Sci U S A *115*, 1370-1375.

Marzolo, M.P., Bull, P., and Gonzalez, A. (1997). Apical sorting of hepatitis B surface antigen (HBsAg) is
independent of N-glycosylation and glycosylphosphatidylinositol-anchored protein segregation. Proc Natl
Acad Sci U S A *94*, 1834-1839.

Matusek, T., Marcetteau, J., and Therond, P.P. (2020). Functions of Wnt and Hedgehog-containing
extracellular vesicles in development and disease. J Cell Sci *133*.

Matusek, T., Wendler, F., Poles, S., Pizette, S., D'Angelo, G., Furthauer, M., and Therond, P.P. (2014).
The ESCRT machinery regulates the secretion and long-range activity of Hedgehog. Nature *516*, 99-103.

Mayor, S., and Riezman, H. (2004). Sorting GPI-anchored proteins. Nature reviews Molecular cell biology
5, 110-120.

Mellman, I., and Nelson, W.J. (2008). Coordinated protein sorting, targeting and distribution in polarized
cells. Nature reviews Molecular cell biology *9*, 833-845.

Mostov, K.E., Altschuler, Y., Chapin, S.J., Enrich, C., Low, S.H., Luton, F., Richman-Eisenstat, J., Singer,
K.L., Tang, K., and Weimbs, T. (1995). Regulation of protein traffic in polarized epithelial cells: the
polymeric immunoglobulin receptor model. Cold Spring Harb Symp Quant Biol *60*, 775-781.

Oyanadel, C., Holmes, C., Pardo, E., Retamal, C., Shaughnessy, R., Smith, P., Cortes, P., Bravo-Zehnder,
M., Metz, C., Feuerhake, T.*, et al.* (2018). Galectin-8 induces partial epithelial-mesenchymal transition with
invasive tumorigenic capabilities involving a FAK/EGFR/proteasome pathway in Madin-Darby canine
kidney cells. Mol Biol Cell *29*, 557-574.

Paladino, S., Lebreton, S., and Zurzolo, C. (2015). Trafficking and Membrane Organization of GPI-
Anchored Proteins in Health and Diseases. Curr Top Membr *75*, 269-303.

Paladino, S., Pocard, T., Catino, M.A., and Zurzolo, C. (2006). GPI-anchored proteins are directly targeted
to the apical surface in fully polarized MDCK cells. J Cell Biol *172*, 1023-1034.

Paladino, S., Sarnataro, D., Pillich, R., Tivodar, S., Nitsch, L., and Zurzolo, C. (2004). Protein
oligomerization modulates raft partitioning and apical sorting of GPI-anchored proteins. J Cell Biol *167*,
699-709.

Panakova, D., Sprong, H., Marois, E., Thiele, C., and Eaton, S. (2005). Lipoprotein particles are required
for Hedgehog and Wingless signalling. Nature *435*, 58-65.

Parchure, A., Vyas, N., Ferguson, C., Parton, R.G., and Mayor, S. (2015). Oligomerization and
endocytosis of Hedgehog is necessary for its efficient exovesicular secretion. Mol Biol Cell *26*, 4700-4717.

Parchure, A., Vyas, N., and Mayor, S. (2018). Wnt and Hedgehog: Secretion of Lipid-Modified
Morphogens. Trends Cell Biol *28*, 157-170.

Peng, T., Frank, D.B., Kadzik, R.S., Morley, M.P., Rathi, K.S., Wang, T., Zhou, S., Cheng, L., Lu, M.M.,
and Morrisey, E.E. (2015). Hedgehog actively maintains adult lung quiescence and regulates repair and
regeneration. Nature *526*, 578-582.

Pepinsky, R.B., Zeng, C., Wen, D., Rayhorn, P., Baker, D.P., Williams, K.P., Bixler, S.A., Ambrose, C.M.,
Garber, E.A., Miatkowski, K.*, et al.* (1998). Identification of a palmitic acid-modified form of human Sonic
hedgehog. J Biol Chem *273*, 14037-14045.

Perez Bay, A.E., Schreiner, R., Benedicto, I., Paz Marzolo, M., Banfelder, J., Weinstein, A.M., and
Rodriguez-Boulan, E.J. (2016). The fast-recycling receptor Megalin defines the apical recycling pathway of
epithelial cells. Nature communications *7*, 11550.

Perez Bay, A.E., Schreiner, R., Mazzoni, F., Carvajal-Gonzalez, J.M., Gravotta, D., Perret, E., Lehmann
Mantaras, G., Zhu, Y.S., and Rodriguez-Boulan, E.J. (2013). The kinesin KIF16B mediates apical
transcytosis of transferrin receptor in AP-1B-deficient epithelia. EMBO J *32*, 2125-2139.

Petrov, K., Wierbowski, B.M., Liu, J., and Salic, A. (2020). Distinct Cation Gradients Power Cholesterol
Transport at Different Key Points in the Hedgehog Signaling Pathway. Developmental cell *55*, 314-327
e317.

Petrov, K., Wierbowski, B.M., and Salic, A. (2017). Sending and Receiving Hedgehog Signals. Annu Rev
Cell Dev Biol *33*, 145-168.

Pizette, S., Matusek, T., Herpers, B., Therond, P.P., and Rabouille, C. (2021). Hherisomes, Hedgehog
specialized recycling endosomes, are required for high level Hedgehog signaling and tissue growth. J Cell
Sci *134*.

Porter, J.A., Young, K.E., and Beachy, P.A. (1996). Cholesterol modification of hedgehog signaling
proteins in animal development. Science *274*, 255-259.

Rindler, M.J., and Traber, M.G. (1988). A specific sorting signal is not required for the polarized secretion
of newly synthesized proteins from cultured intestinal epithelial cells. J Cell Biol *107*, 471-479.

Rodriguez-Boulan, E., and Macara, I.G. (2014). Organization and execution of the epithelial polarity
programme. Nature reviews Molecular cell biology *15*, 225-242.

Rojas, R., and Apodaca, G. (2002). Immunoglobulin transport across polarized epithelial cells. Nature
reviews Molecular cell biology *3*, 944-955.

Sargiacomo, M., Lisanti, M., Graeve, L., Le Bivic, A., and Rodriguez-Boulan, E. (1989). Integral and
peripheral protein composition of the apical and basolateral membrane domains in MDCK cells. J Membr
Biol *107*, 277-286.

Sarnataro, D., Paladino, S., Campana, V., Grassi, J., Nitsch, L., and Zurzolo, C. (2002). PrPC is sorted to
the basolateral membrane of epithelial cells independently of its association with rafts. Traffic *3*, 810-821.

Sbrogna, J.L., Barresi, M.J., and Karlstrom, R.O. (2003). Multiple roles for Hedgehog signaling in zebrafish
pituitary development. Developmental biology *254*, 19-35.

Scheiffele, P., Peranen, J., and Simons, K. (1995). N-glycans as apical sorting signals in epithelial cells.
Nature *378*, 96-98.

Sezgin, E., Levental, I., Mayor, S., and Eggeling, C. (2017). The mystery of membrane organization:
composition, regulation and roles of lipid rafts. Nature reviews Molecular cell biology *18*, 361-374.

Shyer, A.E., Huycke, T.R., Lee, C., Mahadevan, L., and Tabin, C.J. (2015). Bending gradients: how the
intestinal stem cell gets its home. Cell *161*, 569-580.

Simmen, T., Nobile, M., Bonifacino, J.S., and Hunziker, W. (1999). Basolateral sorting of furin in MDCK
cells requires a phenylalanine- isoleucine motif together with an acidic amino acid cluster. Mol Cell Biol *19*,
3136-3144.

Simon, E., Aguirre-Tamaral, A., Aguilar, G., and Guerrero, I. (2016). Perspectives on Intra- and
Intercellular Trafficking of Hedgehog for Tissue Patterning. J Dev Biol *4*.

Simons, K., and Ikonen, E. (1997). Functional rafts in cell membranes. Nature *387*, 569-572.

Stewart, D.P., Marada, S., Bodeen, W.J., Truong, A., Sakurada, S.M., Pandit, T., Pruett-Miller, S.M., and
Ogden, S.K. (2018). Cleavage activates dispatched for Sonic Hedgehog ligand release. Elife *7*.

Tanaka, Y., Okada, Y., and Hirokawa, N. (2005). FGF-induced vesicular release of Sonic hedgehog and
retinoic acid in leftward nodal flow is critical for left-right determination. Nature *435*, 172-177.

Therond, P.P. (2012). Release and transportation of Hedgehog molecules. Curr Opin Cell Biol *24*, 173-
180.

Tukachinsky, H., Kuzmickas, R.P., Jao, C.Y., Liu, J., and Salic, A. (2012). Dispatched and scube mediate
the efficient secretion of the cholesterol-modified hedgehog ligand. Cell Rep *2*, 308-320.

Tuma, P.L., and Hubbard, A.L. (2003). Transcytosis: crossing cellular barriers. Physiol Rev *83*, 871-932.

Vyas, N., Goswami, D., Manonmani, A., Sharma, P., Ranganath, H.A., VijayRaghavan, K., Shashidhara,
L.S., Sowdhamini, R., and Mayor, S. (2008). Nanoscale organization of hedgehog is essential for long-
range signaling. Cell *133*, 1214-1227.

Vyas, N., Walvekar, A., Tate, D., Lakshmanan, V., Bansal, D., Lo Cicero, A., Raposo, G., Palakodeti, D.,
and Dhawan, J. (2014). Vertebrate Hedgehog is secreted on two types of extracellular vesicles with
different signaling properties. Sci Rep *4*, 7357.

Walton, K.D., and Gumucio, D.L. (2021). Hedgehog Signaling in Intestinal Development and Homeostasis.
Annual review of physiology *83*, 359-380.

Wang, E., Brown, P.S., Aroeti, B., Chapin, S.J., Mostov, K.E., and Dunn, K.W. (2000). Apical and
basolateral endocytic pathways of MDCK cells meet in acidic common endosomes distinct from a nearly-
neutral apical recycling endosome. Traffic *1*, 480-493.

Wang, Q., Asarnow, D.E., Ding, K., Mann, R.K., Hatakeyama, J., Zhang, Y., Ma, Y., Cheng, Y., and
Beachy, P.A. (2021). Dispatched uses Na(+) flux to power release of lipid-modified Hedgehog. Nature.

Zavros, Y., Orr, M.A., Xiao, C., and Malinowska, D.H. (2008). Sonic hedgehog is associated with H+-K+-
ATPase-containing membranes in gastric parietal cells and secreted with histamine stimulation. Am J
Physiol Gastrointest Liver Physiol *295*, G99-G111.

Zeng, X., Goetz, J.A., Suber, L.M., Scott, W.J., Jr., Schreiner, C.M., and Robbins, D.J. (2001). A freely
diffusible form of Sonic hedgehog mediates long-range signalling. Nature *411*, 716-720.

Zhou, D., Li, Y., Zhou, L., Tan, R.J., Xiao, L., Liang, M., Hou, F.F., and Liu, Y. (2014). Sonic hedgehog is a
novel tubule-derived growth factor for interstitial fibroblasts after kidney injury. J Am Soc Nephrol *25*, 2187-
2200.

Zurzolo, C., Le Bivic, A., Quaroni, A., Nitsch, L., and Rodriguez-Boulan, E. (1992). Modulation of
transcytotic and direct targeting pathways in a polarized thyroid cell line. EMBO J *11*, 2337-2344.

Zurzolo, C., Lisanti, M.P., Caras, I.W., Nitsch, L., and Rodriguez-Boulan, E. (1993).
Glycosylphosphatidylinositol-anchored proteins are preferentially targeted to the basolateral surface in
Fischer rat thyroid epithelial cells. J Cell Biol *121*, 1031-1039.

<

Shh / E-cadherin

Figure 1. Shh expressed in MDCK cells is apically distributed and secreted. (A) Domain-selective biotinylation of permanently transfected MDCK-Shh grown in Transwell filters. MDCK-Shh Clone 1 shows Shh mainly at the apical cell surface or non-polarized when its expression is increased by 2mM Na+-butyrate treatment for 24h. MDCK-Shh Clone 2 expressing higher Shh levels displays Shh equally distributed among the apical and basolateral cell surface. The basolateral distribution of E-cadherin and Na+/K+-ATPase corroborates the polarized status of the cells in all conditions. (B) Graphs show Shh relative apical versus basolateral cell surface distribution and secretion in MDCK-Shh clone 1. Domain-specific cell surface biotinylation of six independent experiments (n=6) and immunoblot of TCA precipitated proteins from low-serum media conditioned for 6 h of three independent experiments (n=3) reveal aprox. 80% apical cell surface distribution and more than 90% secretion of Shh, respectively. Data expressed as a percentage of total surface or secreted Shh and analysed by an unpaired t-test. In all figures error bars represent mean ± SEM. ****, P≤0.0001. (C) MDCK-Shh clone 1 cells immunofluorescence staining for Shh (green) and E-cadherin (magenta). Images displayed as maximum projection of confocal z-stack and orthogonal views show an apical pattern of Shh. Scale bar, 10 μm.

Figure 2. Newly synthesized Shh first arrives at the basolateral cell surface and then reaches the apical side by transcytosis for secretion. Polarized monolayers grown in Transwell filters of MDCK-Shh (A, C) or GFP-NO-GPI (B) cells were pulse-labeled with ^{35}S-methionine/cysteine and chased at 37^{0}C for the indicated times by subsequent cell-surface-specific biotinylation at 4^{0}C. Cell surface targeting of newly synthesized proteins (Shh, E-cadherin or GFP-NO-GPI) and loading control (10% of total protein in lysates) was analyzed by SDS-PAGE and Cyclone radioactivity scanner (see details in Materials and Methods). (C) Immunoprecipitation of ^{35}S-Shh shows its secretion to either apical or basolateral chambers. (D) MDCK-Shh were incubated with reducible-Biotin by the basolateral domain at 4^{0}C, then allowed to resume trafficking by incubation at 37^{0}C for 120 min. Biotin was then reduced by either apical or basolateral sides to detect internalization or transcytosis of the pool of Shh initially biotinylated by the basolateral domain. E-cadherin was used as a control of a basolateral resident protein. Data expressed as a percentage of total surface Shh of three independent experiments (n=3) and analyzed by an unpaired t-test. In graph error bars represent mean ± SEM. *, P≤0.05. (E) MDCK-Shh cells were biotinylated from the basolateral side at 4^{0}C and then incubated at 37^{0}C for 2h. Apical and basolateral conditioned media were collected, precipitated with Neutravidin and analyzed by immunoblot for secreted Shh. Data expressed as a percentage of total secreted Shh of four independent experiments (n=4) and analyzed by an unpaired t-test. In graph error bars represent mean ± SEM. ***, P≤0.0001.

Figure 3. Basolateral Shh moves to the apical side by transcytosis through Rab-11-ARE and apical Shh can undergo reverse transcytosis. MDCK-Shh cells grown in Transwell filters were incubated with anti-Shh antibodies at 4^0C from the basolateral (A, B) or apical (C) sides, shifted 37^0C for the indicated times and Shh distribution was followed by immunofluorescence and orthogonal views of confocal images. Shh moved from the basolateral to the apical domain involving a subapical punctate compartment typical of the ARE, as corroborated in by its colocalization with endogenous Rab11 (red) after 60 min of trafficking (B). (C) Apically immunolabeled Shh moving to the basolateral domain is detectable after 60 min. Scale bar, 10 μm.

Figure 4. Sorting of Shh lipidation mutants. (A) Schematic illustration of Shh processing and lipidation. (B-F) MDCK cells stably expressing Shh lipidation mutants lacking both lipids (B, ShhNC24S), containing only palmitoyl (C, ShhN) or cholesteryl (D-F, ShhNpC24S) moieties were analyzed by domain-specific cell-surface biotinylation and secretion (presence in the media after 2h accumulation, TCA precipitation and immunoblot). Basolateral E-cadherin and Na+/K+-ATPase transmembrane proteins and secreted Fibronectin (FN) were used to control polarity. (E) MDCK-ShhNpC24S cells were pulsed labeled with ^{35}S-methionine/cysteine and chased to detect protein arrival to each cell surface along with its polarized secretion, as in Figure 2A. Samples were analyzed by SDS-PAGE and Cyclone radioactivity scanner. (F) MDCK-ShhNpC24S cells were biotinylated from the basolateral side at 4°C and then incubated at 37°C for 2h. To assess for ShhNpC24S secretion, apical and basolateral conditioned media were collected, precipitated with Neutravidin and analyzed by immunoblot. Data expressed as a percentage of total secreted Shh of three independent experiments (n=3) and analysed by an unpaired t-test. In graph error bars represent mean ± SEM. ****, P≤0.0001.

Figure 5. Shh is apically and basolateraly distributed under Dispt-1 expression while it is retained at ARE in cells expressing cleavage deficient Disp-1CS. MDCK cells were plasmid microinjected and protein expression was allowed for 4h. (A) Immunofluorescence of co-microinjected Shh (green) and basolateral marker LDLR-GFP-Y18A (magenta). (B) Co-microinjected cells with Shh and Disp-1 were fixed and immune-stained first by the apical side with anti-Shh (green) antibodies and then permeabilized and stained for Disp-1 with anti-HA (magenta) antibodies. (C) Cells were microinjected with Shh alone (upper panels) or together with Disp-1 (middle panels) or cleavage deficient Disp-1-CS (lower panels). Then cells were analyzed by indirect immunofluorescence for Shh (green) and Disp-1 (HA, magenta). Arrowheads indicate non-microinjected cells with basolateral Shh localization due to secretion from expressing cells. Zoomed image to the right demonstrates Shh apically and basolaterally detected in the presence of Disp-1 (BL view) or retained in the ARE compartment in the presence of Disp-1-CS (maximal projection, also see Fig. 7). In all figures scale bar, 10 μm.

Figure 6. Colocalization of transiently expressed Shh and Disp-1 in Rab11-ARE. Polarized MDCK cells were microinjected with Shh-GFP (green) along with Disp-1 and after 4 hrs expression cells were fixed and immunostained for anti-HA (magenta) and anti-Rab11 (yellow). A microinjected cell is shown in the upper panel (center) and a non-microinjected cell is visible at the down-right corner. Maximal projection of 10 planes of 0.1 microns sections over thshows the 3D surface rendering of the zoomed region, in which colocalization of Shh (green) Disp-1 (magenta) e nucleus shows the subapical Rab11-ARE displaying a tubular pattern. Center panel and Rab11 (yellow), is shown by white arrowheads. The lower panel illustrates a 3D surface rendering of medial and basal planes (40 planes of 0.1 microns sections), in which basal distribution of Disp-1 and few vesicles containing Shh or Rab11 are shown. Scale bar, 10 µm.

A. Shh-GFP Hoechst

B. Shh-GFP / Rab11-CFP

0 min   90 min

C.

Time (37ºC)

| | 0 min | 30 min | 60 min | 90 min | 120 min | 120 min (overexposed) |

Shh-GFP

Shh-GFP + Disp-1

Shh-GFP + Disp-1-CS

Figure 7. Basolateral secretion of Shh from the TGN is enhanced by Disp-1 and decreased by Disp-1-CS. (A) Shh-GFP distribution after 4 h of expression in plasmid microinjected polarized MDCK cells show a similar polarity as Shh at steady state (See Figure 1). (B) Shh-GFP expressed by microinjection in MDCK cells permanently transfected with Rab11-CFP plasmid. After 45 min of Shh-GFP expression, the cells were incubated for 2 h at 20ºC to arrest the protein at the TGN (time = 0) and then shifted to 37ºC to reestablish the exit from TGN for 90 min. Arrows point to Rab11-ARE and arrowheads show basolateral Shh-GFP. (C) Live cell imaging analysis of Shh-GFP trafficking from the TGN to the cell surface for the indicated times. Shh-GFP alone (upper panel), co-microinjected with Disp-1 (middle panel) or Disp-1-CS (lower panel). Shh-GFP detected in the border of neighboring non-expressing cells reflects its basolateral secretion, which is enhanced by co-expressed Disp-1 and decreased by Disp-1-CS, more clearly seen in the overexposed image at time 120 min.