IFN-γ stimulated murine and human neurons mount anti-parasitic

defenses against the intracellular parasite *Toxoplasma gondii*

Sambamurthy Chandrasekaran[1], Joshua A. Kochanowsky[2], Emily F. Merritt[2], Joseph S.

Lagas[3], Ayesha Swannigan[4], Anita A. Koshy* [1,2,5]

[1]BIO5 Institute, University of Arizona, Tucson, Arizona, 85719, United States of

America.

[2]Department of Immunobiology, University of Arizona, Tucson, Arizona, 85719, United

States of America.

[3]Undergraduate Biology Research Program, University of Arizona, Tucson, Arizona,

85719, United States of America.

[4]Undergraduate Research Opportunities Consortium, University of Arizona, Tucson,

Arizona, 85719, United States of America.

[5]Department of Neurology, University of Arizona, Tucson, Arizona, 85719, United States

of America.

Running title: Neurons can clear intracellular parasites

22 Correspondence: Anita A. Koshy, akoshy@arizona.edu

Summary

Dogma holds that *Toxoplasma gondii* persists in neurons because neurons cannot clear intracellular parasites, even with IFN-γ stimulation. As several recent studies questioned this idea, we used primary murine neuronal cultures from wild-type and transgenic mice in combination with IFN-γ stimulation and parental and transgenic parasites to reassess IFN-γ dependent neuronal clearance of intracellular parasites. We found that neurons respond to IFN-γ and that a subset of neurons clear intracellular parasites via immunity regulated GTPases. Whole neuron reconstructions from mice infected with parasites that trigger neuron GFP expression only after full invasion revealed that ~40% of these *T. gondii*-invaded neurons no longer harbor parasites. Finally, IFN-γ stimulated human stem cell derived neurons showed a ~ 50% decrease in parasite infection rate when compared to unstimulated cultures. This work highlights the capability of human and murine neurons to mount cytokine-dependent anti-*T. gondii* defense mechanisms *in vitro* and *in vivo*.

Keywords: Murine neurons, human neurons, *T. gondii*, Immunity related GTPases,

Introduction

A select number of highly divergent intracellular microbes (e.g. measles virus, polio virus, *Toxoplasma gondii*) cause infections of the central nervous system (CNS). Though these microbes infect many cell types, in the CNS, neurons are often preferentially infected. One commonly cited reason for this neuron predominance is that neurons lack the ability to mount traditional cell-intrinsic immune responses (Joly et al., 1991; Oldstone et al., 1986; Rall et al., 1995). For example, neurons have low baseline levels of MHC I and STAT1 (Joly et al., 1991; Neumann et al., 1995; Rose et al., 2007; Wong et al., 1984). However, considerable evidence shows that neurons can respond to type I and type II interferons (Delhaye et al., 2006; Neumann et al., 1997, 1995; Rose et al., 2007) and clear certain viral pathogens (e.g. Sindbis virus (Binder and Griffin, 2001; Orvedahl et al., 2010) and vesicular stomatitis virus (Detje et al., 2009)). Limited work has been done on the capabilities of neurons to clear non-viral pathogens.

Toxoplasma gondii is a protozoan parasite that naturally infects most warm-blooded animals, including humans and mice. In most immune competent hosts, *T. gondii* establishes a persistent or latent infection by switching from its fast, growing lytic form (the tachyzoite) to its slow growing, encysting form (the bradyzoite). In humans and mice, the CNS is a major organ of persistence and neurons are the principal cell in which *T. gondii* cysts are found (D. J. Ferguson and Hutchison, 1987; D. J. P. Ferguson and Hutchison, 1987; Melzer et al., 2010; Cabral et al., 2016). IFN-γ is essential for control of *T. gondii* both systemically and in the CNS (Suzuki et al., 1989, 1988), in part through the activation of the immunity regulated GTPase system (IRGs) which is critical

for parasite control in hematopoetic and non-hematopoetic cells (Collazo et al., 2002;

Halonen et al., 2001; Taylor et al., 2000). Based upon these findings and *in vitro* studies

showing that *T. gondii* readily invades murine astrocytes and neurons, but only IFN-γ-

stimulated astrocytes— not IFN-γ-stimulated neurons— clear intracellular parasites

(Jones et al., 1986; Halonen et al., 1998, 2001; Schluter et al., 2001), our model of CNS

toxoplasmosis was that during natural infection parasites enter the CNS, invade both

astrocytes and neurons, after which astrocytes kill the intracellular parasites, leaving the

immunologically incompetent neuron as the host cell for the persistent, encysted form of

the parasite.

Several recent findings have called this model into question. Pan-cellular ectopic

expression of an MHC I allele (H-2 Ld) associated with low levels of CNS persistence

(Blanchard et al., 2008; Brown et al., 1995) leads to a lower CNS parasite burden than

when mice lack expression of this MHC I allele in neurons only (Salvioni et al., 2019).

Moreover, the use of a Cre-based system that permanently marks CNS cells that have

been injected with *T. gondii* proteins (Koshy et al., 2012, 2010), revealed that parasites

extensively interact with neurons and that the majority (> 90%) of these *T. gondii*-

injected neurons do not actively harbor cysts (Cabral et al., 2016; Koshy et al., 2012).

Together these *in vivo* studies question our prior model by raising the possibility that

neurons clear intracellular parasites.

Given these conflicting *in vitro* and *in vivo* findings, here we used primary murine

neuronal cultures from wild-type and genetically modified mice in combination with

cytokine stimulation and parental and transgenic parasites, including a new engineered

T. gondii-Cre line, to reassess the ability of neurons to clear intracellular parasites in the

setting of IFN-γ stimulation. These data reveal that neurons respond to IFN-γ, including

up-regulating the IRGs, and that a subset of neurons (~20%) clear intracellular parasites

via the IRGs. In addition, in Cre reporter mice infected with *T. gondii*-Cre parasites that

mark CNS cells only after fully invasion, whole neuron reconstructions showed that

~40% of these *T. gondii*-invaded neurons no longer harbor parasites. Finally, IFN-γ

stimulation of human stem cell derived neurons (huSC-neurons) led to an ~50%

decrease in parasite infection rate when compared to unstimulated, infected cultures.

Collectively, these data highly suggest that IFN-γ stimulation leads to parasite

resistance in murine and human neurons and that a subset of murine neurons clear

intracellular parasites both *in vitro* and *in vivo*, likely via the IRGs.

Results

IFN-γ stimulated primary pure murine cortical neurons show classical IFN-γ

responses.

As *T. gondii* primarily infects and encysts in the cortex (Berenreiterová et al., 2011;

Boillat et al., 2020; Mendez et al., 2021), we sought to determine the response of

cortical neurons to IFN-γ stimulation. To accomplish this goal, we exposed pure primary

murine cortical neuronal cultures to 100 U/ml of IFN-γ or vehicle control for 4 and 24hrs,

followed by harvesting of total RNA. We chose these time points because prior work

showed that hippocampal neurons have a delayed IFN-γ response (Rose et al., 2007).

After harvesting the RNA, we used quantitative real time-PCR (qRT-PCR) to quantify

the transcripts levels of traditional IFN-γ-response genes (*STAT1*, *IRF1*, *MHC-I*) as well

as the effector components of the IRG system (*Irga6*, *Irgb6*, and *Gbp2*) (Howard et al.,

2011; Khaminets et al., 2010). We found that *STAT1*, the classical transcription factor

that drives the expression of many IFN-γ response genes, and *IRF1* were highly

upregulated (4hrs:3-5 $\log_2$ fold; 24hrs: 5-7 $\log_2$ fold) compared to unstimulated neurons

(**Fig 1A**), while MHC-I showed a more modest level of upregulation (~ 2 $\log_2$ fold). In

addition, consistent with finding in non-neuronal murine cells, compared to unstimulated

neurons, IFN-γ stimulated neurons also significantly up-regulated *Irga6*, *Irgb6*, and

Gbp2 (4hrs:4-7 $\log_2$ folds; 24hrs: 7-9 $\log_2$ folds) (**Fig 1A**) (Boehm et al., 1998; Degrandi

et al., 2013; Lafuse et al., 1995). To determine how these increased transcript levels

translated to protein levels, we isolated total protein lysates from unstimulated and IFN-

γ stimulated cultures. For STAT1, we blotted both for total STAT1 and for

phosphorylated STAT1, the active form that enters the nucleus and binds DNA.

Compared to unstimulated cultures, IFN-γ stimulation cultures showed a >10-fold

increase in protein levels for total STAT1 and p-STAT1 at 24hrs post-stimulation and an

>35-fold increase at 48hrs post-stimulation (**Fig 1B**). The undetectable level of total

STAT1 in unstimulated neurons is consistent with previously published data suggesting

that cultured neurons have low or no constitutive amounts of STAT1 (O'Donnell et al.,

2015). Similarly, at 24 hours post-stimulation, the IRG complex effector proteins Irga6

and Irgb6 increased ~ 7-fold and 10-fold respectively over unstimulated cultures, a level

that was maintained at 48 hours post stimulation.

To confirm that the detected changes were primarily driven by neurons and not by glial

cells that commonly cause low levels of contamination in "pure" neuronal cultures, we

stained the cultures to determine what percentage of cells were neurons, astrocytes, or

microglia. These analyses showed that our cultures were consistently 95% neurons and

5% astrocytes; no microglia were observed (**Fig S1**). To determine the level of nuclear

translocation pSTAT1 at the single neuron level, we stained unstimulated and

stimulated (100 U/ml for 24hrs) primary neuron cultures with DAPI and antibodies

against Tuj1, a neuronal marker, and p-STAT1. We then quantified the pSTAT1 signal

intensity in Tuj1$^+$ nuclei. In unstimulated cultures, we observed almost no pSTAT1

signal in Tuj1$^+$ nuclei (MFI 2236 ± 353.24, mean ± SEM) (**Fig 1C, D**). Conversely, IFN-γ

stimulated neurons showed robust pSTAT1 staining in Tuj1$^+$ nuclei (MFI 44778 ±

25785, mean ± SEM) (**Fig 1C, D**).

Together these data show that IFN-γ stimulated primary murine cortical neuron cultures

upregulate IFN-γ response genes and proteins in a delayed manner consistent with

what has been observed in hippocampal neurons (Rose et al., 2007). The upregulated

genes and proteins include *STAT1* and the IRG genes known to be required for IFN-γ-

dependent killing of intracellular parasites in murine non-neuronal cells (Khaminets et

al., 2010). Collectively, these data suggest that IFN-γ stimulated murine cortical neurons

upregulate the appropriate machinery to clear intracellular parasites via the IRG system.

IFN-γ pre-stimulation leads to a decrease in the percentage of *T. gondii*-infected

neurons.

To address neuronal capability for clearing intracellular parasites, we infected IFN-γ

stimulated or unstimulated neurons with the two canonical parasite strains (type II and

type III) that have moderate to mild acute virulence in mice because both are IRG-

sensitive (Khaminets et al., 2010). We then monitored infection rates of neurons at 3,

12, and 24 hours post infection (hpi). At 3hpi, stimulated and unstimulated neurons

showed a similar rate of neuron infection, regardless of infecting strain (**Fig 2A, B**). By

12 and 24 hpi, regardless of infecting strain, IFN-γ stimulated cultures showed an ~25%

decrease in neuron infection rate compared to unstimulated cultures (**Fig 2A**, **B**).

GCre-expressing parasites show that neurons clear parasites in response to IFN-

γ.

While the prior data suggested that IFN-γ stimulated neurons might clear intracellular

parasites, they could also be explained by decreased rates of late invasion in the IFN-γ

stimulated neurons, especially as clearance assays in neurons are limited by the

inability to synchronize infection or vigorously wash off uninvaded parasites as either

procedure causes widespread neuronal death. To address the possibility of an invasion

defect versus true clearance of intracellular parasites, we required a way to specifically

track neurons that were infected and subsequently cleared the intracellular parasite. As

noted above, we had previously developed a Cre-based system that leads to GFP

expression only in host cells injected with *T. gondii* proteins. In this system, Cre is fused

to a rhoptry protein (or ROP), which are parasite proteins that are injected into host cells

prior to invasion, which means that parasite-triggered host cell expression does not

require parasite invasion (i.e., aborted invasion) (**Fig 2C**). Thus, the RCre (ROP::Cre)-

expressing parasites do not help us distinguish between aborted invasion versus

clearance of intracellular parasites. Therefore, we fused Cre to a dense granule protein

(GRA) that is released into host cells only *after* invasion (Bougdour et al., 2013; Braun

et al., 2013; Franco et al., 2016; Gay et al., 2016). Thus, GCre (GRA::Cre)-expressing

parasites would not cause Cre-mediated recombination in the setting of aborted

invasion (**Fig 2C**) and would identify only host cells that were or had previously been

infected.

We engineered type II (Prugniaud) parasites to express an HA-tagged GCre (II-GCre).

Using immunofluorescent assays and plaque assays, we determined that GCre was

expressed, did not localize to the rhoptries, and that the expression of GCre did not

affect overall parasite viability (**Fig S2**). To test the capability of II-GCre parasites to

trigger Cre-mediated recombination, we infected fibroblast Cre reporter cells that

express GFP only after Cre-mediated recombination (Koshy et al., 2010) with II-RCre

parasites, II-GCre parasites, or parental type II parasites (no Cre expression). At 24

hours post-infection, we observed that both II-RCre parasites and II-GCre parasites

caused host cell expression of GFP, while the parental strain did not (**Fig. S3A**).

Compared to II-RCre parasites, II-GCre parasites showed a decreased efficiency of

causing Cre-mediated recombination (>90% vs. 50%, **Fig. S3B**), which was expected

because host cell-exported GRA proteins show decreased exportation when fused to

ordered proteins (Bracha et al., 2018; Curt-Varesano et al., 2016; Franco et al., 2016).

Having confirmed that Il-GCre parasites trigger Cre-mediated recombination, we next

assessed the capability of Il-GCre parasites to identify only infected cells. To address

this concern, at 24 hours post-infection, we quantified the number of GFP$^+$ Cre reporter

fibroblasts that harbored parasites. With Il-RCre parasites, ~ 30% of GFP$^+$ cells were

actively infected, while with Il-GCre parasites, ~ 60% percent of GFP$^+$ cells were

infected (**Fig. S3B**). While the Il-GCre parasites doubled the rate of infected GFP$^+$ cells,

~40% were uninfected. As this fibroblast Cre reporter cell line continues to divide after

infection and Cre-mediated recombination (Koshy et al., 2012), we hypothesized that

such division accounted for the uninfected GFP$^+$ cells in Il-GCre infected cultures. To

test this possibility, we used cortical neuron cultures as neurons do not divide. In neuron

cultures from Cre reporter mice, infection with Il-RCre parasites resulted in 67% ±

1.82% of GFP$^+$ neurons being actively infected, while infection with Il-GCre parasites

resulted in 98% ± 0.62% of GFP$^+$ neurons being actively infected (**Fig 2 D, E**). Given

that both RCre and GCre infected cultures showed a substantial increase in actively

infected GFP$^+$ cells when using non-dividing cells, these data suggest that in the

fibroblast Cre reporter cell line, post-Cre-mediated recombination cell division accounts

for ~40% of the uninfected GFP$^+$ cells. For the Il-RCre parasites, the remaining ~30% of

uninfected GFP$^+$ cells (and the ~30% of uninfected GFP$^+$ neurons) likely arise from

aborted invasion (**Fig 2C**). For Il-GCre parasites, that ~100% of GFP$^+$ neurons were

infected confirms that Il-GCre parasites trigger Cre-mediated recombination only after

fully invading the host cell.

We next tested how IFN-γ pre-stimulation affected the rate of infected GFP$^+$ neurons by

stimulating Cre reporter neuron cultures with vehicle alone or IFN-γ (100 U/ml) for 24

hours prior to infection with II-GCre parasites. Consistent with previous results (**Fig 2E**),

in the vehicle treated cultures, 97 ± 1.4% of GFP$^+$ neurons harbored a parasite (**Fig. 2F,**

G). In the setting of pre-treatment with IFN-γ, now only 78 ± 1.6% GFP$^+$ neurons

harbored parasites. The data suggest that the decrease in the rate of infection in IFN-γ

stimulated neurons (**2A, B**) is primarily mediated by IFN-γ stimulated neurons clearing

intracellular parasites, rather than IFN-γ stimulation leading to a decrease in parasite

invasion.

In IFN-γ stimulated murine neurons Irga6 loads onto the PVM in a Irgm1/3-

dependent manner

As IFN-γ stimulated neurons up-regulate the IRG effectors (**Fig 1**) and a portion clear

intracellular parasites (**Fig 2**), we next sought to determine if neurons use the IRG

system to mediate IFN-γ-dependent killing of *T. gondii*. As the loading of IRGs on *T.*

gondii PVM is indispensable for parasite clearance in IFN-γ stimulated non-neuronal

cells (Y. O. Zhao et al., 2009) and as Irga6 is one of the effectors that loads onto the PV

(Khaminets et al., 2010), we analyzed the percentage of PVs that were also Irga6$^+$ in

unstimulated and IFN-γ stimulated neurons infected with type III parasites. Consistent

with findings in non-neuronal murine cells, we found that in unstimulated neurons, type

III parasite PVs showed almost no Irga6 loading, while PVs in IFN-γ stimulated neurons

showed an 8-fold increase in Irga6 loading (**Fig 3A, B**). To further confirm these

findings, we also used a type III "IRG-resistant" strain (Cabral et al., 2016). This strain is

engineered to express high levels of ROP18 (III+ROP18). ROP18 is a *T. gondii* kinase

that phosphorylates Irga6 thereby preventing PV loading and effector oligomerization

(Hermanns et al., 2016). The parental type III strain has minimal expression of ROP18

because of an insertion in the promoter region of the *rop18* gene; it is this lack of

ROP18 expression that renders the parental strain susceptible to the IRGs (Saeij et al.,

2006; Taylor et al., 2006). In cultures infected with III+ROP18 parasites, as expected,

we now found almost no Irga6$^+$ PVs even in the setting of IFN-γ stimulation (**Fig 3C, D**).

Finally, to confirm that IRG-loading followed the same principles in neurons as in non-

neuronal cells, we used neurons that lacked the regulatory IRG components (Irgm1 and

Irgm3) (Collazo et al., 2001). Irgm1 and Irgm3 tether the effector components (e.g.

Irga6) to the appropriate organelle until triggered to release the effectors onto the PV

(Hunn et al., 2008) and thus are required for PV loading of Irga6 (Henry et al., 2009). In

Irgm1/3 KO neurons, we found an increase in Irga6 dispersion throughout the cytosol in

the setting of IFN-γ stimulation but no specific loading onto PVs regardless of IFN-γ

stimulation or infecting strain (**Fig 3E-H**).

Together, these data show that neuronal Irga6 loads onto the parasitophorous

vacuole/PVM of intracellular IRG-sensitive parasites in the setting of IFN-γ pre-

stimulation and when neurons have an intact IRG-system. These data strongly suggest

that murine neurons use the IRG system for IFN-γ-dependent clearance/killing of

intracellular parasites.

IFN-γ stimulated murine neurons kill intracellular parasites using the IRG system.

While the preceding data strongly suggests that murine neurons deploy the IRG-system

to kill intracellular parasites in the settling of IFN-γ stimulation, they do not show that the

IRG-system is essential for neuronal killing. To directly test this possibility, we took

several approaches. First, we tested the ability of IFN-γ stimulated, wild type (WT)

neurons to clear III+ROP18 parasites which show no Irga6$^+$ loading even in the setting

of IFN-stimulation (**Fig 3D**). Indeed, in murine neuronal cultures infected with III+ROP18

parasites, we found the same rate of infection over time, regardless of IFN-γ stimulation

state (**Fig 4A**). Second, we infected Irgm1/3 KO neuronal cultures with parental (type III,

IRG-sensitive) or III+ROP18 parasites (IRG-resistant) and with or without IFN-γ pre-

stimulation. We again found the same rate of infection, regardless of what strain we

utilized (IRG-sensitive or resistant) and IFN-γ pre-stimulation state (**Fig 4B**). As a final

method of confirming that the lack of IRGs specifically affected intracellular parasites,

we bred the Irgm1/3 KO mice to the Cre reporter mice to yield mice homozygous for the

Cre reporter construct and that lack both Irgm1 and Irgm3 (Fig S4) and generated

neuronal cultures from these Cre reporter Irgm1/3 KO mice. We then stimulated these

neuronal cultures with vehicle or IFN-γ, followed by infection with II-GCre parasites. In

these cultures, ~98% (-IFN-γ: 98.12 ± 0.67%; + IFN-γ: 97.8% ± 0.53) of GFP$^+$ neurons

were infected, regardless of IFN-γ stimulation status (**Fig 4C, D**).

Collectively, these data definitively show that *in vitro* IFN-γ stimulated neurons kill

intracellular IRG-sensitive parasites via the IRG system.

Neurons clear intracellular parasites *in vivo*

Having shown that IFN-γ stimulated neurons clear intracellular parasites *in vitro*, we

sought to determine if neurons cleared parasites *in vivo*. As the II-GCre parasites trigger

Cre-mediated recombination only after full host cell invasion (**Fig 2C**), we reasoned that

if we found GFP$^+$, parasite$^-$ neurons in Cre reporter mice infected with II-GCre parasites,

these neurons must have cleared the invading parasite. To assess for GFP$^+$ parasite-

neurons *in vivo*, we created whole neuron reconstructions from 200 µm cleared brain

sections stained with Hoechst from 21-day post-infection II-GCre infected mice (Cabral

et al., 2020; Koshy and Cabral, 2014). As the PV excludes GFP expressed by the host

cell, we looked for areas within the GFP$^+$ neurons devoid of GFP (Fig 5A, B) and then

confirmed that these areas contained parasites using Hoechst staining of the parasite

DNA (Fig 5B). Out of 22 reconstructed GFP$^+$ neurons, we found that 9 (~40%) showed

no evidence of persistent parasite infection (Fig 5C, D). These data highly suggest that

in vivo, a percentage of murine neurons clear intracellular parasites.

IFN-γ stimulated human neurons show resistance to *T. gondii* infection

Having shown that murine neurons clear a portion of intracellular parasites *in vitro* and

in vivo, we sought to translate these findings to human neurons. While mice and murine

cells are good models for human infection (both are naturally infected with *T. gondii*,

have the CNS as a major organ of persistence, have neurons as the major host cell for

cysts, and require IFN-γ and CD8 T cells to control toxoplasmosis), differences exist

between the two. In the current context, the most relevant difference is that human cells

lack the expansive range of Irgms that mice have and instead rely on alternative cell-

specific mechanisms for IFN-γ-dependent control of *T. gondii* (Fisch et al., 2019). To

determine how IFN-γ stimulation influenced control of *T. gondii* in human neurons, we

derived human neurons (huSC neurons) from human neuroprogenitor cells

reprogrammed from an embryonic stem cell line. After confirming that the huSC

neurons expressed appropriate cortical neuronal markers (**Fig S5**), we used these

huSC neurons for the *T. gondii* clearance assay. We pre-treated the huSC neurons with

human IFN-γ for 24 hours, followed by infection with type II or III parasites. We then

monitored the rate of neuron infection at 3, 12, and 24 hpi. At 3 hpi, regardless of

infecting strain, we found equivalent neuron infection rates between unstimulated and

IFN-γ stimulated cultures. At 12 and 24 hpi, IFN-γ stimulated cultures showed an ~50%

decrease in the number of infected neurons compared to unstimulated cultures (**Fig 6**).

Discussion

In this study, we sought to address the question: can neurons directly clear intracellular

parasites? Using *in vitro* primary murine neurons from wild-type and transgenic mice in

combination with IFN-γ pre-stimulation and transgenic parasites, this work shows that a

portion of neurons can and do clear intracellular parasite in an IFN-γ-dependent, IRG-

dependent manner. Using our new GCre parasites, which trigger host cell mediated

recombination only after full invasion, we also show that ~ 40% of neurons clear

parasites *in vivo*. Finally, using huSC neurons, we translated our findings to human

neurons, showing that IFN-γ pre-stimulation decreases the infection rate by ~50%.

The data presented are the first to show that IFN-γ pre-stimulation enables human and

murine neurons to partially resist infection by an intracellular eukaryotic pathogen. In

murine neurons, we leveraged *T. gondii* biology to show that this resistance was

secondary to clearance of intracellular parasites (**Fig 2A, B**), not simply an invasion

defect (an important distinction in cultures where procedures such as aggressively

washing off extracellular parasites cannot be done). In addition, our finding that IFN-γ

stimulated murine neurons clear parasites in an IRG-dependent manner explains why a

prior study found that IFN-γ stimulated primary murine neurons failed to clear

intracellular parasites. The prior study, which was done at a time when neither the IRGs

nor the parasite mechanisms to block the IRGs had been fully described, used a type I

strain (RH) that we now know is IRG-resistant (Hermanns et al., 2016; Schluter et al.,

2001). In human neurons, we currently cannot distinguish between an IFN-γ-dependent

invasion defect or clearance of intracellular parasites or both. Though we cannot

distinguish between these possibilities, the IFN-γ-dependent, anti-parasite effect

appears to have a more robust effect on huSC neuron infection rates compared to

murine neurons. What mechanisms underlie this impressive IFN-γ-dependent

resistance to *T. gondii* infection will be the subject of future studies.

Though we have shown that IFN-γ-stimulated neurons can clear parasites in an IRG-

dependent manner, our data also suggests that major cell-specific differences in IRG

efficiency exist. We found that *in vitro*, ~ 20-25% of intracellular parasites will be cleared

by neurons in an IFN-γ, IRG-dependent manner, while other groups have shown that

IFN-γ stimulated murine astrocytes, macrophages, and fibroblasts have higher rates of

Irga6[+] loading (50-75%) and clearance over a much shorter time (1-2 hours post-

infection) (Khaminets et al., 2010; MacMicking, 2012; Martens et al., 2005). While some

of the difference may be secondary to technical differences (e.g. use of antibody vs.

transfection of Irga6-tagged with GFP, different MOIs), part of the difference is likely

secondary to a blunted cell-intrinsic immune response from neurons (as suggested by

the undetectable levels of baseline STAT1). Other, not mutually exclusive possibilities

include that IRG-clearance differs between neuronal subcellular locations (i.e. it might

be expensive to put IRG-machinery along the whole neuron) and/or that full neuron

responses require direct interactions with other cell types such as astrocytes or T cells.

Our *in vivo* data using GCre parasites are consistent with the possibility that other cell

types influence neuronal clearance of intracellular parasites. *In vivo* we found that ~

40% of GCre-triggered GFP⁺ neurons do not harbor parasites (**Fig 5**). This rate of

neuronal clearance is approximately double what we observed *in vitro*. In the *in vivo*

setting, neurons are in constant communication with other cell types (e.g. astrocytes,

microglia, infiltrating T cells), which may potentiate IRG-dependent clearance or initiate

complementary methods for clearing intracellular parasites (e.g. CD40-dependent

xenophagy (Andrade et al., 2006; Subauste, 2009)). Future studies will focus on

defining these *in vivo* vs. *in vitro* differences.

In summary, our findings offer substantial evidence that IFN-γ pre-stimulation enables

murine and human neurons to mount anti-parasitic defenses against *T. gondii*. While

much work is left to be done to understand these anti-parasitic defenses, the work

presented here suggests that *T. gondii's* persistence in neurons is not simply a foregone

conclusion.

Acknowledgements

The authors would like to thank all members of the Koshy Lab for helpful discussions and critical review of the manuscript. We would like thank Greg Taylor (Duke University), for providing us with breeding pairs for *Irgm1/3^{-/-}*, Jonathan Howard (Instituto Gulbenkian de Ciência), for providing us with anti-Irga6 antibodies. Finally, we thank Dr. Jared Churko (University of Arizona) and Dr. Rita Sattler (Barrow Neurological Institute) and their lab members for support with human neuron differentiations. Funding was provided by the National Institutes of Health (NS095994 (AAK); AI147711 (JAK)) and the BIO5 Institute, University of Arizona (A.A.K). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Author Contributions

Conceptualization: Sambamurthy Chandrasekaran, Joshua Kochanowsky, Anita A. Koshy.

Methodology: Sambamurthy Chandrasekaran, Joshua Kochanowsky, Emily F. Merritt, Anita A. Koshy.

Validation: Sambamurthy Chandrasekaran, Joshua Kochanowsky, Emily F. Merritt

Formal analysis: Sambamurthy Chandrasekaran, Joshua Kochanowsky, Emily F. Merritt, Anita A. Koshy.

Investigation: Sambamurthy Chandrasekaran, Joshua Kochanowsky, Emily F. Merritt, Anita A. Koshy.

Data curation: Sambamurthy Chandrasekaran, Joshua Kochanowsky, Emily F. Merritt

410 **Writing – original draft**: Sambamurthy Chandrasekaran

411 **Writing – review & editing**: Anita A. Koshy

412 **Supervision**: Anita A. Koshy

413 **Funding acquisition**: Joshua A. Kochanowsky, Anita A. Koshy

Declaration of Interests

416 The authors have declared that no competing interests exist.

Materials and Methods

Parasite maintenance

The parasite strains used in this study were maintained through serial passage in

human foreskin fibroblasts (HFFs) using DMEM, supplemented with 10% fetal bovine

serum (FBS), 2mM Glutagro and 100 IU/ml penicillin and 100 µg/ml streptomycin.

Except for the type II (Prugniaud) strain that expresses Gra16::Cre (II-GCre), the *T.*

gondii strains used have been previously described (Cabral et al., 2016; Koshy et al.,

2012). To engineer the II-GCre strain, type II parasites were electroporated with a

plasmid encoding Cre recombinase fused to the dense granule protein, Gra16, and a

separate drug-selectable marker (Donald et al., 1996). Single clones were selected by

limiting dilution.

Mice

All procedures and experiments were carried out in accordance with the Public Health

Service Policy on Human Care and Use of Laboratory Animals and approved by the

University of Arizona's Institutional Animal Care and Use Committee (#12-391). All mice

were bred and housed in specific-pathogen-free University of Arizona Animal Care

facilities. Cre reporter mice (Madisen et al., 2010) (#007906) were originally purchased

from Jackson Laboratories. Breeding pairs of *Irgm1/m3$^{-/-}$* (Irgm1/3 KO) mice (Collazo et

al., 2001) were generously provided by Greg Taylor (Duke University, Durham, NC).

***In vivo* infection with GCre parasites**

Cre reporter mice were intraperitonally infected with 10,000 or 20,000 freshly lysed

Gra16-Cre (GCre) tachyzoites. Mice were anesthetized 21 days post infection,

harvested brains were drop fixed in 4% PFA and stored overnight at 4^{0}C before being

transferred and stored in 30% sucrose until they were sectioned. *T. gondii* infected,

sucrose embedded brains were sagittal sectioned to 200 μm on a vibratome and stored

in cryoprotectant media (0.05 M sodium phosphate buffer containing 30% glycerol and

30% ethylene glycol). Sections were cleared using a modified PACT clearing protocol

previously described (Cabral et al., 2020). In brief, sections were incubated overnight in

a hydrogel monomer solution at 4 degrees Celsius and deoxygenated the next morning

by bubbling nitrogen gas into sample vials. Samples were incubated at 42 degrees

Celsius to initiate crosslinking of proteins, washed, and incubated in 8% sodium dodecyl

sulfate (SDS) at 45 degrees Celsius to remove lipids. After multiple wash steps to

remove SDS from the sections, nuclei were stained using hoechst. Samples were then

washed and submerged in sorbitol refractive index matching solution (sRIMS)

consisting of 70% Sorbitol and 0.01% NaN$_3$ dissolved in 0.02M PB overnight before

being mounted and imaged on a Zeiss NLO 880 confocal microscope (Imaging Core –

Marley, University of Arizona). Brain sections were mounted for imaging on spacer

slides in fresh sRIMS (Koshy and Cabral, 2014).

Images were obtained at 40x magnification to ensure visualization of parasite nuclei.

We created z-stack tile scans of each neuron to ensure capturing as much of the

neuron's axon and processes as were available in each section. Stitched images were

converted to Imaris files and imported into Bitplane Imaris software, where neuronal

projections were rendered using the filaments tool.

Primary murine neuron culturing

The primary neurons were cultured by methods described previously with minor

modifications (Parker et al., 2018). The culturing plates were prepared by coating

overnight with 0.001% poly-L-lysine (Sigma, Cat # P4707) solution for plastic surfaces

and 100 µg/ml poly-L-lysine hydrobromide (Sigma, Cat # P6282) for glass surfaces.

Neurons were seeded in plating media at appropriate densities: 500,000 in 6-well plates

for RNA and protein extraction, 100,000 in 24 well plates with coverslips for imaging and

20,000 in 96 well plates for counting. Four hours after plating, full volume media

exchange to neurobasal media (Thermo Fisher, Cat # 21103049) was performed. On

day in vitro (DIV) 4, neurons received a half volume media change of neurobasal media

with 5 µM cytosine arabinoside to stop glial proliferation. One third media exchanges

with neurobasal media occurred every 3-4 days thereafter. All the experiments were

performed on 12 DIV neurons.

IFN-γ stimulation and *T. gondii* infection

The primary neurons were pre-stimulated with 100 U/ml of mouse recombinant IFN-γ for

4h & 24h (RNA extraction), 24h & 48h (protein extraction), or 24h (*T. gondii* infections).

Freshly syringe-lysed *T. gondii* parasites resuspended in neurobasal media were used

to infect the primary neurons at MOI=4 (For 3 hpi), MOI=2 (For 12 hpi), MOI=1 (For 24

hpi) and MOI=0.2 (for 48 and 72 hpi) time points.

Quantitative real time PCR

For quantification of the genes, RNA was extracted from 4h and 24h stimulated primary

neurons using TRIzol reagent (Life technologies) and following the manufacturer's

protocol. 500 ng of total RNA was converted into first strand cDNA using a High-

Capacity cDNA Reverse Transcription Kit (Applied Biosystems^{TM}; Cat No: 4368814) and

following the manufacturer's instructions. Using the primer listed in Table 1, IFN-γ

response and IRG pathway genes were amplified using SYBR green fluorescence

detection with an Epppendorf Mastercycler ep realplex 2.2 system. GAPDH was used

as a housekeeping gene to normalize DNA levels. Results were calculated using the $2^{-\Delta\Delta CT}$

method (Livak and Schmittgen, 2001).

Protein extraction and Western blotting

Primary neurons were either unstimulated or stimulated with IFN-γ for 24h and 48h,

followed by total protein extraction as previously described (Cabral et al., 2017). Equal

amounts of protein were subjected to SDS-PAGE, transferred to PVDF membrane and

western blotting was done by standard methods. The blots were imaged using the

Odyssey Infrared Imaging Systems (LI-COR Biosciences).

Neuronal clearance assay

Primary neurons (wildtype or Irgm 1/3 KO) plated were plated on a poly-L-lysine coated

96-well plate (20,000/well) and were either unstimulated or pre-stimulated with IFN-γ for

24h prior to infection with *T. gondii* parasites. The cells were labeled with anti-NeuN

(neuronal nuclei), DAPI (all nuclei) and parasites were mcherry positive, wells were

imaged using an Operatta CLS high content analysis microscope (Functional Genomics

Core, University of Arizona). Generated images were then analyzed using Image J

software. The results were represented as percent decrease in the infected neurons in

stimulated compared to the unstimulated group.

For assays involving GCre parasites, neurons from either wildtype (Cre reporter) or Cre

reporter Irgm 1/3 KO mice were plated on poly-L-lysine coated 6 well plate

(500,000/well). At the appropriate DIV, the neurons were either unstimulated or pre-

stimulated with IFN-γ for 24h followed by infection with II-GCre parasites. At 72 hrs

post-infection, the plates were processed as described above. These fixed and stained

plates were then analyzed using an epifluorescent microscope (EVOS). The person

analyzing the images was blinded to the IFN-γ stimulation and/or infecting parasite

strain.

Immunofluorescence assay

521 Cells were grown on poly-L-lysine-coated glass coverslips (described above) and were

522 processed by methods as previously described (Parker et al., 2018).

Antibodies

The following primary antibodies were used in the study: mouse anti-tubulin beta III

isoform (Tuj1),clone TU20 (MAB1637, Millipore, 1:1000); rabbit anti- β3-Tubulin, D71G9

(similar to Tuj1) (5568S, CST, 1:1000); mouse anti-NeuN clone A60 (MAB377, Millipore,

1:1000); rabbit anti-Glial Fibrillary Acidic protein (GFAP) (Z0334, DAKO, 1:500); rabbit

anti S100 (Z0311, DAKO, 1:500); rabbit anti-ALDH1L1 (Ab87117, Abcam, 1:500);

chicken anti-Iba1 (Ab 139590, Abcam, 1:500), rabbit anti-STAT1 (Ab47425, Abcam;

1:500); mouse anti-pSTAT1 pY701 clone14/p-STAT1 (612132, BD Biosciences, 1:250);

rabbit anti-pSTAT1 Tyr701, Clone 58D6 (9167, CST, 1:200); mouse anti-SAG1 DG52

(gift John Boothroyd, 1:10,000); mouse anti-SRS-9 (gift John Boothroyd, 1:10,000);

mouse anti-Irga6 (1:1500), mouse anti-Irgb6 (1:250) (gift Jonathan Howard); rabbit ant-

HA C29F4 (3724S, CST, 1:500); mouse anti-ROP2/3/4 (1:1000, gift John Boothroyd);

DAPI (D3571, Thermo Fisher, 1:1000); Hoechst 33342 Trihydrochloride, Trihydrate

(H3570, Thermo Fisher, 1:1000). The following species-appropriate secondary

antibodies were used: Alexa Fluor 405 goat anti-rabbit IgG, Alexa Fluor 488 goat anti-

mouse IgG, Alexa Fluor 568 goat anti-rabbit IgG, Alexa Fluor 647 goat anti-mouse IgG,

Alexa Fluor 647 goat anti-chicken IgG, Alexa Fluor 647 goat anti-rabbit IgG, donkey

anti-mouse IgG, DyLight 680 conjugate (1:10000), and donkey anti-rabbit IgG, DyLight

800 conjugate (1:10000). Unless otherwise noted, secondary antibodies were obtained

from Life Technologies and used at a concentration of 1:500.

Human stem cell derived neurons

H7 human embryonic Neural stem cells (NSCs) derived from the NIH-approved H7

embyronic stem cells (WiCell WA07) were purchased from the University of Arizona

iPSC core (https://stemcells.arizona.edu/). The NSCs were expanded and differentiated

into cortical layer neurons using a previously described protocol (Yan et al., 2013) with

minor modifications. Briefly, NSCs were expanded on Matrigel® Matrix (Corning®,

#354277) coated plates using NSC expansion medium (NEM) (Thermofisher, Cat #

A1647801). The media was changed every other day until NSCs reached confluence.

The passaged NSCs were used at P2 for differentiation into cortical neurons by plated

them on poly-L-ornithine (20ug/ml) (Sigma, Cat # P4957) and laminin (5ug/ml) (Thermo

Fisher, Cat # 23017015) coated plates. For 14 days, the cells were differentiated into

cortical neurons using neural differentiation medium (NDM) consisting of Neurobasal

medium, 2mM L-Glutamine (Thermo Fisher, Cat # 25030024), 1% B-27 (Thermo

Fisher, Cat # 17504044), 200µM L- Ascorbic acid (Sigma, Cat # A92902), 0.5mM c-

AMP (Stem Cell Technologies Cat # 73886), 20ng/ml BDNF (Stem Cell Technologies

Cat # 78005), 20ng/ml GDNF (Stem Cell Technologies Cat # 78058), 20ng/ml NT-3

(Stem Cell Technologies Cat # 78074), and Penicillin/Streptomycin (Thermo Fisher, Cat

15140122) cocktail. The culture medium was exchanged with fresh NDM every 2-3

days.

Human neuronal clearance assay

The clearance assay in human neurons was performed as described for primary pure

murine neurons except that human IFN-γ (R&D Systems, Cat # 285-MI-100) was used

for pre-stimulation.

571   **Statistical Analyses**

572   Graphs were generated and statistical analyses were performed using Graphpad Prism

573   9.1.2 software. The specific test used (e.g., ANOVA vs. t-test) is noted in each figure.

Fig 1. IFN-γ stimulated primary murine neurons show intact IFN-γ signaling pathway and express genes involved in *T. gondii* clearance.

Primary neurons were cultured for 12 days *in vitro* (DIV), after which they were stimulated with vehicle or IFN-γ (100 U/ml). At the listed times, RNA or protein was isolated or immunofluorescent assays were performed. **A.** Quantification of specified genes using quantitative PCR. Expression is shown as Log_2 fold change compared to unstimulated cultures. **B.** Representative images of western blots for specified proteins from unstimulated and IFN-γ stimulated neuron cultures. **C.** Densitometric quantification of western blots from (**B**). Densitometry of given gene is normalized to densitometry of β-tubulin and then shown as fold change compared to unstimulated cultures. For (**A, C**) Bars, mean ± SEM. N = 3 independent experiments. **D.** *Images*: Representative images of unstimulated or IFN-γ stimulated neurons stained as indicated (anti-Tuj1 antibodies stains neurons). Scale bar = 5μm. *Graph*: Quantification of the mean fluorescent intensity (MFI) of pSTAT1 nuclear signal in unstimulated or IFN-γ stimulated Tuj1$^+$ cells (neurons). N = 48-51 nuclei evaluated/condition/experiment, 2 independent experiments. Bars, mean ± SD. ****$p \leq 0.0001$, unpaired t-test with Welch's correction.

Fig 2. IFN-γ stimulated neurons clear intracellular parasites.

Primary neurons were cultured and stimulated as in Fig 1. **A.** Graph of the percentage

of infected neurons at listed time points for type II parasites, normalized to the

unstimulated culture. **B.** As in (**A**) except for type III parasites. (**A, B**) Primary neurons

were stimulated with IFN-γ (100U/ml) or vehicle for 24hrs, after which the cultures were

infected with type II or type III parasites. At the listed time points, cultures were fixed,

stained with anti-NeuN antibodies (stains neuron nuclei) and DAPI, and analyzed on an

Operatta CLS high content analysis microscope. Bars mean ± SEM. N = 25-30 FOV

(~750-1000 neurons/experiment), 3 independent experiments. ****$p<0.0001$. ns = not

significant, 2-way ANOVA with Sidak's multiple comparisons. **C.** Schematic of host cells

labeled by RCre versus GCre parasites. *Left*, In the RCre system the Cre fusion protein

is secreted prior to invasion of neurons, which leads to GFP-expressing neurons that

can arise from: i) injection, invasion, and chronic infection, ii) injection and invasion of

host cell followed by host cell killing or clearance of the parasite, iii) injection of the

protein <u>*without*</u> invasion (aborted invasion). *Right*, In the GCre system the Cre fusion

protein is secreted post invasion, which leads to GFP-expressing neurons that can arise

from: i) invasion and persistent infection or ii) invasion followed by killing or clearance of

the parasite, but not from aborted invasion. **D.** Representative images of GFP+ neurons

infected with RCre or GCre parasites. *Merge image*: Green = GFP-expression in

neurons, white = parasites stained with anti-*T. gondii* antibodies (a cocktail of anti-SAG1

and anti-SRS9 antibodies to capture both tachyzoites and bradyzoites.) Scale bar = 10

µm **E.** Graph of the percentage of actively infected GFP+ neurons at 72 hpi

(unstimulated cultures). **F.** Representative images of unstimulated or IFN-γ (100 U/ml)

stimulated primary neurons infected with GCre parasites for 72hrs. *Merge image*, as in

(**D**). Scale bar = 10 µm **G.** Graph of the percentage of actively infected GFP$^+$ neurons at

72 hpi. (**E, G**) Bars mean ± SEM. N ≥ 200 GFP$^+$ neurons/well, 3 wells/experiment, 3-4

independent experiments. **p ≤ 0.005, ***p ≤ 0.0005, t-test with Welch's correction.

Tuj1 Irga6 T. gondii

Fig 3. IFN-γ stimulated wild-type murine show increased loading of Irga6 onto the PVM of type III parasites.

Primary neurons were cultured as in Fig 1, followed by 24 hours of IFN-γ (100U/ml) or vehicle stimulation, after which cultures were infected with listed *T. gondii* strains. At 12 hours post infection, cultures were fixed and stained with anti-Irga6 and anti-Tuj1 antibodies. The stained cultures were analyzed by confocal microscope. **A.** Representative images of stained cultures from wild-type (WT) mice infected with type III parasites and pre-stimulated with vehicle or IFN-γ. Blue = anti-Tuj1 antibodies, Green = anti-Irga6 antibodies, Red = mCherry expressing parasites. **B.** Quantification of the percentage of Irga6+ parasitophorous vacuoles (PVs) in the setting of vehicle or IFN-γ pre-stimulation. **C.** Representative images as in (**A**) except infected with III+ROP18 parasites. **D.** As in (**B**). **E.** Representative images as in (**A**) except using cultures from Irgm1/3 KO mice. Scale bar = 10 μm full image, 5 μm inset. **F.** As in (**B**). **G.** Representative images as in (**C**) except using IRG KO neurons. **H.** As in (**B**). (**B**, **D**, **F**, **H**) Bars- mean ± SEM, N = 100-200 PVs/experiment, 3 independent experiments. ns- not significant, **p ≤ 0.01, t-test with Welch's correction.

Fig 4. An intact IRG system is required for IFN-γ-dependent murine neurons

clearance of intracellular parasites.

Primary neurons from wild-type (Cre reporter), Irgm1/3 KO, or Cre reporter Irgm1/3 KO

mice were cultured as in Fig 1 followed by stimulated with IFN-γ (100U/ml) or vehicle for

24hrs, after which the cultures were infected with listed *T. gondii* strain. **A.** Graph of the

percentage of infected WT neurons at listed time points for III+ROP18 parasites,

normalized to the unstimulated culture. **B.** As in (**A**) except using Irgm1/3 KO neurons

and either type III parasites (*left graph*) or III+ROP18 parasites (*right graph*). (**A, B**) At

the listed time points, cultures were fixed, stained, and analyzed as in **Fig 2A, B**. Bars

mean ± SEM. N = 25-30 FOV analyzed/experiment (~750-1000 neurons

analyzed/experiment), 3 independent experiments. *p ≤ 0.05, ns = not significant, 2-way

ANOVA with Sidak's multiple comparisons. **C.** Representative images of GFP⁺ neurons

from Cre Reporter Irgm1/3 KO mice infected with II-GCre parasites. *Merge image*:

Green = GFP-expression in neurons, White = parasites stained with anti-*T. gondii*

antibodies (a cocktail of anti-SAG1 and anti-SRS9 antibodies to capture both

tachyzoites and bradyzoites.) Scale bars = 10 µm **D.** Graph of the percentage of

infected GFP⁺ neurons. Bars, mean ± SEM. Eat dot represents the mean value of 1

experiment. N = N ≥ 200 GFP⁺ neurons analyzed/experiment, 4 independent

experiments. ns = not significant, Welch's t test. (**C, D**) Cultures were infected, fixed,

stained, and analyzed as in **Fig 2 F, G**.

Fig 5. Forty percent of GFP⁺ neurons in mice infected with GCre parasites do not harbor parasites.

Cre reporter mice were infected with GCre parasites. At 21 dpi, brains were harvested and sagittally sectioned into 200 μm thick sections. Thick sections were cleared and imaged at 40x on a confocal microscope. Neurons in resulting images were then rendered using Imaris software. **A.** A representative rendering of a GFP⁺ neuron in which no parasites were identified. **B.** As in (**A**) except now with a GFP⁺ neuron in which parasites were identified. White arrow shows parasites within neuron soma. **C.** Single plane of soma from (**B**) *top image*: GFP channel, *middle image*: blue channel (Hoechst), *bottom image*: merge. Note the GFP displacement, suggesting parasite presence within the neuron, which is then confirmed by visualization of parasite nuclei stained with hoecsht (blue). **D.** Graph of rendered GFP⁺ neurons containing parasites (infected) or not containing parasites (uninfected). N = 22 neurons from 4 mice.

Fig 6. IFN-γ stimulation of human neurons leads to a decrease in the percentage infected with *T. gondii*.

Human neurons were differentiated for 14 DIV from neural stem cells, after which they were stimulated with vehicle or IFN-γ (100 U/ml). At listed time points, cultures were

fixed, stained with anti-Tuj1 antibodies and DAPI, and analyzed on an Operatta CLS

high content analysis microscope. **A. C.** Representative images of human neurons

infected with either type II or III parasites. Green = anti-Tuj1, Red = mCherry expressing

parasites, Blue = DAPI. Scale bars = 20 µm **B.** Graph of the percentage of infected

neurons at listed time points for type II parasites, normalized to the unstimulated culture.

D. As in (**B**) except for type III parasites. Bars mean ± SD. N = 7-20 FOV (~1000-1500

neurons/experiment), 2 independent differentiations. ****$p<0.0001$, **$p<0.005$ ns = not

significant, 2-way ANOVA with Sidak's multiple comparisons.

Fig S1. *In vitro* neuronal cultures are pure with little glial contamination

739 Quantification of the *in vitro* cultures after 12 DIV stained with anti-Tuj1 antibodies,

740 (neuronal marker), an anti-astrocyte cocktail (anti-GFAP, anti-S100B, anti-ALDH1L1),

741 and Iba1 (microglia marker). The numbers are presented as percentage of total cells

742 counted (n= 100/experiment). Bars, mean ± SEM, N= 3 experiments.

A.

B.

Fig S2. GCre is expressed and viable in *T. gondii*.

A. Immunofluorescence for HA-tagged RCre and GCre fusion protein. HFFs were

infected with identified strains (MOI=1) for 24 hours. Images depict listed *T. gondii*-Cre

strain stained with anti-HA (green), anti-ROP2/3/4 (red), and DAPI (blue).

B. Quantification of plaque assay at 10 dpi after infection with 200 parasites of the

identified strains. Each dot = 1 experiment, N = 3 experiments. Bars, ± mean SEM.

Welch's t test. ns = not significant.

DAPI GFP Toxo

DAPI GFP Toxo

Fig S3. GCre is capable of Cre-mediated recombination in Cre-reporter fibroblasts and Cre-reporter neurons.

A. Immunofluorescence for Cre-mediated recombination in Cre reporter fibroblasts infected with either RCre or GCre parasites. Cre reporter fibroblasts were infected with identified strains (MOI 1) for 24 hours and then fixed and stained. Images depict GFP (green), parasites (anti-SAG1; red), and nuclei (DAPI, blue).

B. Quantification of Cre-mediated recombination efficiency. Cre reporter fibroblasts were infected as in (**A**) and efficiency scores were quantified by dividing the number of GFP$^+$ cells by the number of infected cells multiplied by 100.

C. The percentage of GFP$^+$ cells that actively harbored a parasite was quantified by dividing the number of infected GFP+ cells by the total number of GFP$^+$ cells multiplied by 100.

D. Immunofluorescence for Cre-mediated recombination in Cre reporter neurons infected with either RCre or GCre parasites. Cre reporter neurons were infected with identified strains (MOI=0.1) for 72 hours and then fixed and stained. Images depict GFP$^+$ neurons (green), parasites (anti-SRS9 and anti-SAG1; red), and nuclei (DAPI, blue).

B and **C** Values, mean ± SEM. Each dot represents the mean value of 1 experiment. N = 3 experiments. One-way ANOVA with Dunnett's multiple comparison test. ***$p<0.0005$ and ****$p<0.0001$. ND = Not Detected.

Fig S4. Generation of Cre reporter Irgm1/m3 knockout mice

A. Schematic representation of breeding Cre reporter to Irgm1/m3 knockout mice.

B. Agarose gel representing the genotyping of the transgenic lines.

Tuj1 Map2 DAPI   Tuj1 Ctip2 DAPI   Tuj1 Brn2 DAPI   Tuj1 Tbr1 DAPI

Fig S5. Characterization of human neurons for cortical layer markers

Human stem cell derived neurons were grown and differentiated for 14 days, after which

cultures were fixed and stained with DAPI and anti-Tuj1 antibodies (pan neuronal

marker) and one of the following: anti-Map2 antibodies (mature neuronal marker), anti-

Ctip2 antibodies (cortical layer V, VI), anti-Brn2 antibodies (cortical layer II-V), or anti-

Tbr1 antibodies (cortical layer I, V, VI). Scale bar = 25µm

References

1. Andrade, R. M., Wessendarp, M., Gubbels, M.-J., Striepen, B. & Subauste, C. S. CD40 induces macrophage anti-Toxoplasma gondii activity by triggering autophagy-dependent fusion of pathogen-containing vacuoles and lysosomes. *J Clin Invest* **116**, 2366–2377 (2006).

2. Berenreiterová, M., Flegr, J., Kuběna, A. A. & Němec, P. The Distribution of Toxoplasma gondii Cysts in the Brain of a Mouse with Latent Toxoplasmosis: Implications for the Behavioral Manipulation Hypothesis. *PLOS ONE* **6**, e28925 (2011).

3. Binder, G. K. & Griffin, D. E. Interferon-γ-Mediated Site-Specific Clearance of Alphavirus from CNS Neurons. *Science* **293**, 303–306 (2001).

4. Blanchard, N. *et al.* Immunodominant, protective response to the parasite Toxoplasma gondii requires antigen processing in the endoplasmic reticulum. *Nat Immunol* **9**, 937–944 (2008).

5. Boehm, U. *et al.* Two Families of GTPases Dominate the Complex Cellular Response to IFN-γ. *The Journal of Immunology* **161**, 6715–6723 (1998).

6. Boillat, M. *et al.* Neuroinflammation-Associated Aspecific Manipulation of Mouse Predator Fear by Toxoplasma gondii. *Cell Reports* **30**, 320-334.e6 (2020).

7. Bougdour, A. *et al.* Host Cell Subversion by Toxoplasma GRA16, an Exported Dense Granule Protein that Targets the Host Cell Nucleus and Alters Gene Expression. *Cell Host & Microbe* **13**, 489–500 (2013).

8. Bracha, S. *et al.* Engineering Brain Parasites for Intracellular Delivery of Therapeutic Proteins. *bioRxiv* 481192 (2018) doi:10.1101/481192.

9. Braun, L. *et al.* A Toxoplasma dense granule protein, GRA24, modulates the early immune response to infection by promoting a direct and sustained host p38 MAPK activation. *Journal of Experimental Medicine* **210**, 2071–2086 (2013).

10. Brown, C. R. *et al.* Definitive identification of a gene that confers resistance against Toxoplasma cyst burden and encephalitis. *Immunology* **85**, 419–428 (1995).

11. Cabral, C. M., McGovern, K. E., MacDonald, W. R., Franco, J. & Koshy, A. A. Dissecting Amyloid Beta Deposition Using Distinct Strains of the Neurotropic Parasite Toxoplasma gondii as a Novel Tool. *ASN Neuro* **9**, 1759091417724915 (2017).

12. Cabral, C. M., Merritt, E. F. & Koshy, A. A. Three-Dimensional Reconstruction of Toxoplasma-Neuron Interactions In Situ. *Methods Mol Biol* **2071**, 283–295 (2020).

13. Cabral, C. M. *et al.* Neurons are the Primary Target Cell for the Brain-Tropic Intracellular Parasite Toxoplasma gondii. *PLoS Pathog.* **12**, e1005447 (2016).

14. Collazo, C. M. *et al.* The function of gamma interferon-inducible GTP-binding protein IGTP in host resistance to Toxoplasma gondii is Stat1 dependent and requires expression in both hematopoietic and nonhematopoietic cellular compartments. *Infect Immun* **70**, 6933–6939 (2002).

15. Collazo, C. M. *et al.* Inactivation of Lrg-47 and Irg-47 Reveals a Family of Interferon γ–Inducible Genes with Essential, Pathogen-Specific Roles in Resistance to Infection. *J Exp Med* **194**, 181–188 (2001).

16. Curt-Varesano, A., Braun, L., Ranquet, C., Hakimi, M.-A. & Bougdour, A. The aspartyl protease TgASP5 mediates the export of the Toxoplasma GRA16 and GRA24 effectors into host cells. *Cellular Microbiology* **18**, 151–167 (2016).

17. Degrandi, D. *et al.* Murine Guanylate Binding Protein 2 (mGBP2) controls Toxoplasma gondii replication. *PNAS* **110**, 294–299 (2013).

18. Delhaye, S. *et al.* Neurons produce type I interferon during viral encephalitis. *PNAS* **103**, 7835–7840 (2006).

19. Detje, C. N. *et al.* Local type I IFN receptor signaling protects against virus spread within the central nervous system. *J Immunol* **182**, 2297–2304 (2009).

20. Donald, R. G. K., Carter, D., Ullman, B. & Roos, D. S. Insertional Tagging, Cloning, and Expression of the Hypoxanthine-Xanthine-Guanine Phosphoribosyltransferase Gene. *Journal of Biological Chemistry* **271**, 14010–14019 (1996).

21. Ferguson, D. J. & Hutchison, W. M. The host-parasite relationship of Toxoplasma gondii in the brains of chronically infected mice. *Virchows Arch A Pathol Anat Histopathol* **411**, 39–43 (1987).

22. Ferguson, D. J. P. & Hutchison, W. M. An ultrastructural study of the early development and tissue cyst formation of Toxoplasma gondii in the brains of mice. *Parasitol Res* **73**, 483–491 (1987).

23. Fisch, D., Clough, B. & Frickel, E.-M. Human immunity to Toxoplasma gondii. *PLOS Pathogens* **15**, e1008097 (2019).

890   24. Franco, M. *et al.* A Novel Secreted Protein, MYR1, Is Central to Toxoplasma's
891   Manipulation of Host Cells. *mBio* **7**, e02231-02215 (2016).
892

893   25. Gay, G. *et al.* Toxoplasma gondii TgIST co-opts host chromatin repressors
894   dampening STAT1-dependent gene regulation and IFN-γ–mediated host
895   defenses. *Journal of Experimental Medicine* **213**, 1779–1798 (2016).
896

897   26. Halonen, S. K., Chiu, F. & Weiss, L. M. Effect of cytokines on growth of
898   Toxoplasma gondii in murine astrocytes. *Infect. Immun.* **66**, 4989–4993 (1998).
899

900   27. Halonen, S. K., Taylor, G. A. & Weiss, L. M. Gamma Interferon-Induced Inhibition
901   ofToxoplasma gondii in Astrocytes Is Mediated by IGTP. *Infection and Immunity*
902   **69**, 5573–5576 (2001).
903

904   28. Henry, S. C. *et al.* Balance of Irgm protein activities determines IFN-γ-induced
905   host defense. *Journal of Leukocyte Biology* **85**, 877–885 (2009).
906

907   29. Hermanns, T., Müller, U. B., Könen-Waisman, S., Howard, J. C. & Steinfeldt, T.
908   The Toxoplasma gondii rhoptry protein ROP18 is an Irga6-specific kinase and
909   regulated by the dense granule protein GRA7. *Cellular Microbiology* **18**, 244–259
910   (2016).
911

912   30. Howard, J. C., Hunn, J. P. & Steinfeldt, T. The IRG protein-based resistance
913   mechanism in mice and its relation to virulence in Toxoplasma gondii. *Curr. Opin.*
914   *Microbiol.* **14**, 414–421 (2011).
915

916   31. Hunn, J. P. *et al.* Regulatory interactions between IRG resistance GTPases in
917   the cellular response to Toxoplasma gondii. *EMBO J* **27**, 2495–2509 (2008).
918

919   32. Joly, E., Mucke, L. & Oldstone, M. B. A. Viral persistence in neurons explained
920   by lack of major histocompatibility class I expression. *Science* **253**, 1283–1286
921   (1991).
922

923   33. Jones, T. C., Bienz, K. A. & Erb, P. In vitro cultivation of Toxoplasma gondii cysts
924   in astrocytes in the presence of gamma interferon. *Infect Immun* **51**, 147–156
925   (1986).
926

927   34. Khaminets, A. *et al.* Coordinated loading of IRG resistance GTPases on to the
928   Toxoplasma gondii parasitophorous vacuole. *Cell. Microbiol.* **12**, 939–961 (2010).
929

930   35. Koshy, A. A. & Cabral, C. M. 3-D imaging and analysis of neurons infected in
931   vivo with Toxoplasma gondii. *J Vis Exp* (2014) doi:10.3791/52237.
932

933   36. Koshy, A. A. *et al.* Toxoplasma co-opts host cells it does not invade. *PLoS*
934   *Pathog.* **8**, e1002825 (2012).
935

37. Koshy, A. A. *et al.* Toxoplasma secreting Cre recombinase for analysis of host-parasite interactions. *Nat Methods* **7**, 307–309 (2010).

38. Lafuse, W. P., Brown, D., Castle, L. & Zwilling, B. S. Cloning and characterization of a novel cDNA that is IFN-gamma-induced in mouse peritoneal macrophages and encodes a putative GTP-binding protein. *J Leukoc Biol* **57**, 477–483 (1995).

39. Livak, K. J. & Schmittgen, T. D. Analysis of Relative Gene Expression Data Using Real-Time Quantitative PCR and the 2−ΔΔCT Method. *Methods* **25**, 402–408 (2001).

40. MacMicking, J. D. Interferon-inducible effector mechanisms in cell-autonomous immunity. *Nature Reviews Immunology* **12**, 367–382 (2012).

41. Madisen, L. *et al.* A robust and high-throughput Cre reporting and characterization system for the whole mouse brain. *Nat. Neurosci.* **13**, 133–140 (2010).

42. Martens, S. *et al.* Disruption of Toxoplasma gondii Parasitophorous Vacuoles by the Mouse p47-Resistance GTPases. *PLOS Pathogens* **1**, e24 (2005).

43. Melzer, T. C., Cranston, H. J., Weiss, L. M. & Halonen, S. K. Host Cell Preference of Toxoplasma gondii Cysts in Murine Brain: A Confocal Study. *J Neuroparasitology* **1**, (2010).

44. Mendez, O. A. *et al.* Semi-automated quantification and neuroanatomical mapping of heterogeneous cell populations. *Journal of Neuroscience Methods* **305**, 98–104 (2018).

45. Neumann, H., Cavalié, A., Jenne, D. E. & Wekerle, H. Induction of MHC class I genes in neurons. *Science* **269**, 549–552 (1995).

46. Neumann, H., Schmidt, H., Cavalié, A., Jenne, D. & Wekerle, H. Major Histocompatibility Complex (MHC) Class I Gene Expression in Single Neurons of the Central Nervous System: Differential Regulation by Interferon (IFN)-γ and Tumor Necrosis Factor (TNF)-α. *Journal of Experimental Medicine* **185**, 305–316 (1997).

47. O'Donnell, L. A. *et al.* Interferon gamma induces protective non-canonical signaling pathways in primary neurons. *J. Neurochem.* **135**, 309–322 (2015).

48. Oldstone, M. B., Blount, P., Southern, P. J. & Lampert, P. W. Cytoimmunotherapy for persistent virus infection reveals a unique clearance pattern from the central nervous system. *Nature* **321**, 239–243 (1986).

49. Orvedahl, A. *et al.* Autophagy protects against Sindbis virus infection of the central nervous system. *Cell Host Microbe* **7**, 115–127 (2010).

50. Parker, S. S. *et al.* High Fidelity Cryopreservation and Recovery of Primary Rodent Cortical Neurons. *eNeuro* **5**, (2018).

51. Rall, G. F., Mucke, L. & Oldstone, M. B. Consequences of cytotoxic T lymphocyte interaction with major histocompatibility complex class I-expressing neurons in vivo. *Journal of Experimental Medicine* **182**, 1201–1212 (1995).

52. Rose, R. W., Vorobyeva, A. G., Skipworth, J. D., Nicolas, E. & Rall, G. F. Altered levels of STAT1 and STAT3 influence the neuronal response to interferon gamma. *Journal of Neuroimmunology* **192**, 145–156 (2007).

53. Saeij, J. P. J. *et al.* Polymorphic Secreted Kinases Are Key Virulence Factors in Toxoplasmosis. *Science* **314**, 1780–1783 (2006).

54. Salvioni, A. *et al.* Robust Control of a Brain-Persisting Parasite through MHC I Presentation by Infected Neurons. *Cell Rep* **27**, 3254-3268.e8 (2019).

55. Schluter, D., Deckert, M., Hof, H. & Frei, K. Toxoplasma gondii infection of neurons induces neuronal cytokine and chemokine production, but gamma interferon- and tumor necrosis factor-stimulated neurons fail to inhibit the invasion and growth of T. gondii. *Infect Immun* **69**, 7889–7893 (2001).

56. Subauste, C. S. CD40, autophagy and Toxoplasma gondii. *Mem Inst Oswaldo Cruz* **104**, 267–272 (2009).

57. Suzuki, Y., Conley, F. K. & Remington, J. S. Importance of endogenous IFN-gamma for prevention of toxoplasmic encephalitis in mice. *J Immunol* **143**, 2045–2050 (1989).

58. Suzuki, Y., Orellana, M. A., Schreiber, R. D. & Remington, J. S. Interferon-gamma: the major mediator of resistance against Toxoplasma gondii. *Science* **240**, 516–518 (1988).

59. Taylor, G. A. *et al.* Pathogen-specific loss of host resistance in mice lacking the IFN-gamma-inducible gene IGTP. *Proc Natl Acad Sci U S A* **97**, 751–755 (2000).

60. Taylor, S. *et al.* A Secreted Serine-Threonine Kinase Determines Virulence in the Eukaryotic Pathogen Toxoplasma gondii. *Science* **314**, 1776–1780 (2006).

61. Wong, G. H. W., Bartlett, P. F., Clark-Lewis, I., Battye, F. & Schrader, J. W. Inducible expression of H–2 and Ia antigens on brain cells. *Nature* **310**, 688–691 (1984).

1027   62. Yan, Y. *et al.* Efficient and rapid derivation of primitive neural stem cells and
1028   generation of brain subtype neurons from human pluripotent stem cells. *Stem*
1029   *Cells Transl Med* **2**, 862–870 (2013).
1030
1031   63. Zhao, Y. O., Khaminets, A., Hunn, J. P. & Howard, J. C. Disruption of the
1032   Toxoplasma gondii parasitophorous vacuole by IFNgamma-inducible immunity-
1033   related GTPases (IRG proteins) triggers necrotic cell death. *PLoS Pathog.* **5**,
1034   e1000288 (2009).
1035

1036