Centrosomal Enrichment and Proteasomal Degradation of SYS-1/β-catenin Requires the Microtubule
Motor Dynein

Joshua W. Thompson, Maria F. Valdes Michel, Bryan T. Phillips

Abstract

The *C. elegans* Wnt/β-catenin Asymmetry (WβA) pathway utilizes asymmetric regulation of SYS-
1/β-catenin and POP-1/TCF coactivators. This differentially regulates gene expression during cell fate
decisions, specifically by asymmetric localization of determinants in mother cells to produce daughters
biased towards their appropriate cell fate at birth. Despite the induction of asymmetry, β-catenin
localizes symmetrically to mitotic centrosomes in both mammals and *C. elegans*. Due to the mitosis-
specific mobility of centrosomal SYS-1 and 'traffic jam' like enrichment of SYS-1 at kinetochore
microtubules when SYS-1 centrosomal loading is disrupted, we investigated active trafficking in SYS-1
centrosomal localization. Here, we demonstrate that trafficking by microtubule motor dynein is required
to maintain SYS-1 centrosomal enrichment, by dynein RNAi-mediated decreases in SYS-1 centrosomal
enrichment and by temperature-sensitive allele of the dynein heavy chain. Conversely, we observe that
depletion of microtubules by Nocodazole treatment or RNAi of putative dynein-proteasome adapter
ECPS-1 exhibits increased centrosomal enrichment of SYS-1. Moreover, disruptions to SYS-1 or negative
regulator microtubule trafficking are sufficient to significantly exacerbate SYS-1 dependent cell fate
misspecifications. We propose retrograde microtubule-mediated trafficking enables SYS-1 and negative
regulators to enrich at centrosomes, enhancing their interaction and perhaps implicating the
centrosome as a mitotic sink for proteins targeted for degradation.

Keywords

Development, Wnt, β-catenin, microtubule, ECM29, dynein, proteasome, trafficking, elegans,
asymmetry, cell cycle, robustness

Introduction

During metazoan development, cells must robustly respond to and integrate multiple signaling pathways to appropriately specify and populate tissues. This is of further importance in the developing embryo, where cell division rapidly alternates between S-phase and mitosis while simultaneously specifying tissues and entire germ layers (Pintard and Bowerman 2019). As such, the cell's response to these signals and the subsequent establishment of asymmetry often begins before asymmetric daughter cells are even 'born' via an asymmetric cell division (ACD) (Sunchu and Cabernard 2020; Juanes, 2020; Fuchs and Chen, 2013; Gómez-López, Petrisch et al 2014). Among the signaling pathways utilized in development is Wnt/β-catenin signaling, wherein the binding of Wnt ligands disrupts β-catenin degradation via dismantling an intracellular destruction complex. This effectively stabilizes β-catenin, enabling it to activate target genes with co-activator TCF/LEF (Nusse and Clevers 2017). Wnt/β-catenin signaling is specialized in its ability to regulate both polarity and proliferation – enabling developing organisms to orient and expand tissues simultaneously for appropriate size and morphology (Loh et al 2016). For this reason, Wnt/β-catenin signaling is readily adaptable throughout development to the adult organism.

Wnt/β-catenin is particularly well adapted to the organization of a patterned stem cell niche for proliferative tissue maintenance (Pinto, Clevers et al 2003; Reilein, Kalderon et al 2017). However, this function in adult tissue homeostasis also associates deleterious dysregulation of Wnt/β-catenin signaling with disease states. Throughout transduction of the Wnt signal are proteins vulnerable to oncogenic mutation (Yang, yen 2011; Zhunussova, Djansugurova et al 2019; Liu, Thibodeau et al 2000). However, these often affect their tumorigenicity via dysregulation of the pathway effector β-catenin, which can promote proliferation via over-and mis-expression of targets like Cyclin D1 and cMyc (Niehrs, Acebron 2012; He, Kinzler et al 1998; Shtutman, Ben-Ze'ev et al 1999; Tetsu and McCormick, 1999). Moreover, β-

catenin is a bifunctional protein in mammalian systems. That is, it is distributed between the previously

described signaling role while also maintaining cell-cell adhesion at adherens junctions (Niehrs, Acebron

2012; Reya, Clevers 2005), where it facilitates the interaction between membrane protein E-cadherin

and the cytoskeleton. The combination of these roles for β-catenin makes it both an attractive target for

studying tumorigenesis, both due to hyperproliferative phenotypes and ease of metastasis. However,

the dual functions can also lead to confounding pleiotropic phenotypes after manipulation of β-catenin

activity (Reya, Clevers 2005; Tejpar, Alman et al 1999; Geyer, Reis-Filho et al 2011).

C. elegans, unlike the canonically described Wnt signaling pathway, has distributed the adhesive

and signaling roles across its four β-catenins (Sawa, 2012). Here we focus specifically on the *C. elegans*

β-catenin SYS-1 for its role as a transcriptional activator in a modified Wnt/β-cat signaling pathway, the

Wnt/β-catenin Asymmetry pathway(WβA) (Phillips, Kimble et al 2007). The WβA pathway is a

specialized application of the more basic Wnt signaling paradigm, wherein regulation of the co-activator

SYS-1 (β-catenin) modulates the activity of transcription factor POP-1 (TCF), during asymmetric cell

division (Baldwin and Phillips 2018, Lam and Phillips 2017). That is, dividing mother cells polarize

components of the Wnt signaling pathway to a cortical region which will appropriately benefit the

daughter cell inheriting that cortex – distributing positive regulators of a Wnt response to the signaled

pole of the cell, and negative regulators to the unsignaled pole (Baldwin and Phillips 2014; Baldwin,

Phillips et al 2016; Sugioka, Sawa et al 2011). Daughter cells in such a division therefore inherit Wnt

signaling components that bias them toward their appropriate cell fate. Contrary to its known role of

asymmetrically activating Wnt target genes the key transcriptional effector of this pathway, SYS-1 (and

indeed, mammalian β-catenin (Mbom, Barth et al 2014; Vora, Phillips et al 2020)), unexpectedly

localizes symmetrically to mother cell mitotic centrosomes (Phillips, Kimble et al 2007; Huang, Lin et al

2007; Vora and Phillips 2015).

Previous work has demonstrated that mother cell centrosomal localization of SYS-1 limits SYS-1

levels in daughter cells in a proteasome dependent manner (Vora and Phillips 2015). Differential

localization of the proteasome during the cell cycle (Amsterdam, Baumeister et al 1993), as well as

interactor-specific localization of the proteasome for intracellular structuring (Baldin, Coux et al 2008),

suggest that proteasome regulation, localization, and activity are interdependent. Proteasomal subunits

have, for example, been shown to co-sediment with centrosomal proteins and perinuclear inclusions

indicative of centrosomal localization (Wigley, Thomas et al 1999; Wojcik, DeMartino 2003). Similarly,

centrosomal localization is required for mitotic clearance of SYS-1, such that uncoupling SYS-1 from

centrosomes results in daughter cells exhibiting approximately 40% more SYS-1 protein. Moreover, SYS-

1 turnover by FRAP mobile fraction at centrosomes is dependent on RPT-4 proteasomal activity (Vora

and Phillips 2015). Perhaps, then, the centrosome is a region where SYS-1 – proteasome interactions are

concentrated to regulate its retention and effect on cell fate.

The directionality and temporal specificity of this mitotic enrichment and degradation implied

that a more active localization mechanism might be at work, compared to a more passive captured

diffusion mechanism. This is most directly demonstrated in that the cell maintains a proteasome-

dependent, 100% mobile population of centrosomal SYS-1, which is recycled within 120 seconds (Vora

and Phillips 2015). Cells appear to maintain this robust and dynamic localization of SYS-1 specifically on

the mitotic centrosome, with the interphase centrosome undetectable via SYS-1 fluorescence (Vora and

Phillips 2015). However, the mechanism of such temporally specific SYS-1 centrosomal localization is

unknown. Several observations indicate that cells utilize the microtubule cytoskeletal network for rapid

and directional regulation of specific cargo to precise locales, including: that dynein conditionally traffics

transcription factors due to neuronal injury or signaling status, (Ben-Yaakov, Fainzilber et al 2012;

Mikenberg, Kaltschmidt et al 2007) and that dynein more generally can enrich endocytic recycling

components around the Microtubule Organizing Center (MTOC) (Horgan, Mccaffrey et al 2011) or be

sufficient to prevent pathogenic accumulation of RNA binding proteins (Deshiumaru, Tsuboi et al 2021).

Therefore, we hypothesized that the role of the mitotic centrosome as an MTOC could facilitate the

rapid transport of SYS-1 to the pericentriolar material specifically in dividing cells.

The rapid and cell cycle phase-specific enrichment of SYS-1 at the centrosome suggests that it

may also depend on active transport. Further, accumulation at a highly active MTOC like the mitotic

centrosome is most likely accomplished by the microtubule minus end-directed dynein motor complex

(Vora and Phillips 2015; Priyanga, Bhaka-Guha et al 2021). The dynein motor is a multi-subunit complex

composed of a pair of ATP-ase, microtubule-binding heavy chains responsible for the force-generating

steps of the motor, and a variable complement of smaller light, intermediate, and accessory chains that

alter the cargo binding or processivity of the motor (Reck-Peterson, Carter et al 2018; Jha and Surrey

2015). The dynein motor is multifunctional - it provides pulling forces, positions organelles, and can

mediate rapid and conditional movement of proteins and RNAs (Carminati and Stearns 1997; Woźniak,

Allan et al 2009; Harada, Hirokawa et al 1998; Hanz, Fainzilber et al 2003; Bullock and Ish-Horowicz

2001). For example, injured rat axons have demonstrated a dynein intermediate chain-dependent

nuclear localization and coimmunoprecipitation of the transcription factor STAT3, in order to

appropriately adopt an injury-dependent transcriptional response (Ben-Yaakov, Fainzilber et al 2012).

Dynein specifically is the motor complex most often directed towards the MTOC and is known to be

active during mitosis (Jha and Surrey 2015). Therefore, we investigated dynein-dependent trafficking as

an ideal mechanism for SYS-1 localization to the MTOC mitotic centrosomes.

Here we demonstrate that microtubules and minus-end microtubule-mediated trafficking are

required for the centrosomal recruitment of both SYS-1 and the negative regulators responsible for its

regulation there. We evaluate the extent to which SYS-1 utilizes dynein-mediated trafficking to

accumulate at centrosomes (Fig. 1A). Using both long- and short-term disruptions to dynein and

microtubule-mediated trafficking to evaluate SYS-1 localization and SYS-1-dependent cellular activity up

to and including cell fate changes, we demonstrate that SYS-1 and its negative regulators both require

microtubule-mediated trafficking for appropriate regulation of SYS-1 (Fig. 1B-D). Together, these data

characterize a system for enhancing the robustness of the WβA signaling pathway by maintaining

consistent interaction of SYS-1 with negative regulators at the mitotic centrosome.

Results

SYS-1 Overexpression and Centrosomal Uncoupling Results in SYS-1 Accumulation at Regions of

Dynein Enrichment

To test the role of microtubule-mediated trafficking in SYS-1 centrosomal localization, we

investigated what trafficking mechanisms are necessary for SYS-1 regulation. We compared

fluorescently tagged SYS-1 protein localization to that of MT trafficking components. Of most interest

was the cytoplasmic dynein motor complex, since its components have been shown to localize to mitotic

centrosomes, kinetochore microtubules, and cortically across species (Schmidt, Van den Heuvel et al

2017; Tame, Medema et al 2014). This corresponded to GFP::SYS-1 , which localizes to these same

cortical and microtubule dense regions. Moreover, this localization was more common upon

centrosomal uncoupling by RNAi depletion of the spindle assembly and scaffolding protein RSA-2(Fig

2A). While GFP::SYS-1 has been observed at mitotic centrosomes throughout the organism,

accumulation at these cortical and kinetochore 'secondary' sites is uncommon (accumulation at either

or both of these sites occurring in 34.5% of embryos) in untreated organisms but the frequency of this

accumulation approximately doubles, to 69.1%, with *rsa-2(*RNAi) (Fig2A). This finding is consistent with

'traffic jams' observed in microtubule mediated trafficking, often aggravated by decreasing microtubule-

off rate (Leduc, Howard et al 2012).

Untreated, GFP::SYS-1 expression, driven by the strong germline/embryonic PIE-1 promoter, is

likely expressed at higher-than-wildtype levels. This increase, in addition to potentially rate limiting

protein interactions required for RSA-2 mediated centrosomal capture, may bottleneck SYS-1 at the

point of centrosomal capture. The centrosomal uncoupling seen in *rsa-2*(RNAi) has also been shown to

increase SYS-1 protein in resulting daughter cells, in addition to the previously described loss of

centrosomal enrichment (Vora and Phillips 2015). This suggests that increases in intracellular SYS-1,

along with a lower SYS-1/dynein 'off' rate from microtubules by the centrosomal 'sink', are sufficient to

overwhelm the centrosomal degradation mechanisms of SYS-1 and crowd centrosome-bound motor

complexes in microtubule motor 'traffic jams'. That is, by decreasing SYS-1 microtubule on-rate due to

disruptions in motor complex formation, or decreasing its microtubule off-rate by disrupting

centrosomal capture by RSA-2, we can observe GFP::SYS-accumulation throughout the process of

trafficking - at regions of dynein loading and unloading, respectively, as well as regions of high

microtubule occupancyas the off-rate 'traffic jam' worsens (Leduc, Howard et al 2012). The

enhancement of SYS-1 kinetochore and cortex localization seen in *rsa-2(RNAi),* with the coinciding

centrosomal uncoupling, is therefore consistent with *rsa-2(RNAi)* reducing the SYS-1 microtubule-off

rate and exhibiting more frequent, or worse, traffic jams.

Knockdown of Dynein Complex Subunits Reduces SYS-1 Centrosomal Localization

Given the observation that SYS-1 localizes to these sites important for trafficking via the

retrograde dynein motor, we investigated a functional role for the dynein complex in SYS-1 localization.

We began by performing an RNAi screen against various components of the dynein motor complex and

associated proteins, eventually expanding outward to additional motors and microtubule-affecting

proteins (Fig S1). Specifically, by RNAi depletion of these microtubule trafficking proteins, we assayed

the centrosomal steady state of GFP::SYS-1. We expected depletion of transport proteins that play a role

in SYS-1 centrosomal transport to be defective in the maintenance of SYS-1 centrosomal enrichment

when SYS-1 centrosomal clearance was otherwise unaffected. In order to evaluate the centrosome-

specific enrichment of SYS-1, we considered two ways of measuring fluorescently-tagged SYS-1 steady

state levels: 1) the centrosomal enrichment index (CEI)(similar to that described in Gao, Smith et al

2017), the centrosomal fluorescence intensity adjusted by subtraction of the background cytoplasmic

fluorescent intensity of the surrounding cell, to control for backgrounds that may pleiotropically

influence SYS-1 expression, and 2) the SYS-1 CEI evaluated as a ratio to the same enrichment factor of

representative PCM factor γ-tubulin (Tubulin Enrichment Index, or TEI) , to avoid false positives via

defects in PCM maturation (CEI/TEI) (Fig 2B-B''). We then compared these centrosomal enrichment

metrics to those seen in established *rsa-2(*RNAi) centrosomal uncoupling.

As expected, depletion of several dynein subunit knockdowns presented pleiotropic and

deleterious effects, including oogenesis and early patterning defects which precluded our SYS-1

localization assay. We therefore treated organisms with partial RNAi knockdowns by commencing RNAi

feeding 12-24 hours before imaging as necessary, based on the phenotypic severity. While no

treatment disrupted SYS-1 centrosomal steady state sufficiently to lose any apparent SYS-1 centrosomal

accumulation as seen in *rsa-2*(RNAi), we did observe several statistically significant decreases in

centrosomal SYS-1 enrichment (Fig S1,2C – C'). Furthermore, while the total population of some

treatments displayed only an intermediate mean loss of SYS-1 enrichment, the distribution of SYS-1 CEI

values observed in dynein subunit knockdowns extends from wild-type to an *rsa-2(RNAi)*-like disruption

(e.g. Fig. S2). These decreases, and notably lower minimum, in SYS-1 CEI or CEI/TEI are masked

somewhat in the population average by this broad range in phenotype and/or RNAi penetrance (Fig 2C-

C', Fig S2). Most notably, *dlc-1(RNAi)* and *dylt-1(RNAi)* each exhibit a significant γ-tubulin-independent

reduction in centrosomal SYS-1 localization. This centrosomal uncoupling was specifically more visible in

a transgene driven by the endogenous SYS-1 promoter, as compared to SYS-1 driven by a stronger and

likely overexpressing germline/early embryo promoter (Fid 2C-C').

These data suggest that dynein does have a role in the localization of SYS-1 to centrosomes, but

redundancy between components of the dynein motor or between the dynein motor and other

processes may limit the effect any one subunit has on SYS-1 localization. Alternately, while mutants are

often severely deleterious, variability of RNAi efficacy may have resulted in incomplete depletion of SYS-

1 relevant dynein subunits. Similarly, the pleiotropic effects of dynein depletion may also have

superseded effects on SYS-1 localization, leading to the evaluation of only partial knockdowns. In order

to address these issues, we investigated the role of the dynein motor in SYS-1 localization at a more

precise temporal resolution via conditional dynein disruption.

Temporally Restricted Dynein Loss of Function Reveals Dynein Requirement for Centrosomal Steady

State SYS-1 Accumulation.

While our screen showed that dynein knockdown results in significant reduction of centrosomal

SYS-1, with rare individuals exhibiting *rsa-2(RNAi)*-like loss of centrosomal SYS-1 accumulation, we were

unable to completely recreate a population exhibiting consistently severe disruption to SYS-1 CEI. To

distinguish the variety in effect on SYS-1 localization effects between functional redundancies limiting

the effects of dynein depletion and the confounding deleterious effects described above, we utilized a

temperature sensitive allele of the cytoplasmic dynein heavy chain, *dhc-1(or195ts),* to conditionally

inactivate this cytoplasmic dynein heavy chain. This allele, identified in the Bowerman lab, introduces a

lesion to the microtubule binding stalk of DHC-1. This lesion renders the motor slightly hypomorphic,

even at the permissive 15°C (with only 91% of embryos hatching), but still sufficiently functional to form

bipolar spindles and generate pulling force throughout the cell, with homozygotes remaining viable and

fertile at permissive temperatures (Hamill, Bowerman et al 2002). However, after upshifting to 26°C,

dhc-1(ts) mutants undergo complete mitotic spindle collapse as little as 30 seconds, suggesting that

DHC-1 motor function has been nearly completely ablated (O'rourke, Bowerman et al. 2007). Therefore,

we expected that replicating this treatment in a strain containing both *dhc-1(ts)* and fluorescently

tagged SYS-1 would allow us to examine a more penetrant dynein knockdown while overcoming both

potential redundancy between dynein accessory chains in the trafficking of SYS-1 and pleiotropic effects

precluding the observation of mitotic centrosomal SYS-1 (Fig. 3A). As described above, we compared the

CEI of the endogenous promoter-driven VENUS::SYS-1 driven by the endogenous SYS-1 promoter to that

of mCherry::γ-Tubulin (TEI) to control for changes in centrosome size and maturation that may

otherwise confound changes specific to SYS-1 centrosomal steady state. Consistent with the published

hypomorphic nature of the allele at the permissive temperature (O'rourke, Bowerman et al 2007),

imaging *dhc-1(ts)* embryos, shows notable decrease in the SYS-1 CEI of embryos kept at the permissive

temperature (Fig 3 B-D). Following 30-45 seconds of temperature upshift, this hypomorphic phenotype

is enhanced to severely reduce SYS-1 centrosomal enrichment (Fig 3B-C). While there was a decrease in

centrosome size, presumably due to roles for dynein in centrosome maturation (Priyanga, Bhakta-Guha

et al 2021), the CEI/TEI metric accounted for this. We observed a CEI/TEI decrease corresponding to

decreased centrosomal SYS-1 accumulation, proportionally greater than the reduction in γ-tubulin (Fig

3D'). Combined with the individual dynein subunit knockdown, the more penetrant phenotype of *dhc-*

1(ts) mutants indicates that SYS-1 requires the dynein microtubule motor complex to localize

appropriately to centrosomes. Additionally, it seems likely that redundancy of peripheral motor

components or incomplete knockdown is responsible for the variability within individual subunit

depletions seen above (fig 2C-C').

MT depolymerization experiments reveal that SYS-1 centrosomal dynamics are affected both

positively and negatively by Microtubule-Mediated trafficking

To determine the extent to which microtubule-based trafficking in general is responsible for SYS-

1 centrosomal localization, we treated gravid adults and their embryos with >100 µM Nocodazole, a

microtubule-destabilizing agent. This dose, while sufficient for a rapid acute response in adult

organisms, caused no effect on fertilized embryos due to their eggshell and prevented the production of

additional embryos via germline arrest (data not shown). We therefore turned to a lower dose, longer-

term Nocodazole treatment: 12 hours at 25µg/mL (83 µM). This maternal Nocodazole dosing allowed us

to obtain embryos with distinctly limited microtubule networks (Fig 4A). Surprisingly, centrosomal and

cytoplasmic SYS-1 both increased after Nocodazole treatment (Fig 4B, C). This SYS-1 CEI increase is

similar to the phenotype observed in RPT-4 depletion, which decreases proteasome function (Vora and

Phillips, 2015), and is consistent with a model wherein negative regulators also require trafficking. In this

case, we expect nocodazole induced changes in SYS-1 localization to diverge from those seen in dynein

knockdowns by the extent of centrosomal enrichment, but converge on excessive inheritance of

GFP::SYS-1 (Fig 1D).

The dichotomy of centrosomal SYS-1 depletion or enrichment across dynein or microtubule

disruption also provided us an opportunity to evaluate the most basal status of SYS-1 at centrosomes.

Moreover, while both temperature upshift for *dhc-1(ts)* and Nocodazole treatments significantly affect

SYS-1 centrosomal localization, neither treatment appears to completely disrupt microtubule mediated

trafficking. Nocodazole-treated embryos still exhibit some kinetochore microtubules (Fig. 4A) but can

tolerate this treatment for some time. Conversely, *dhc-1(ts)* embryos demonstrate such severe loss of

dynein function at the restrictive temperature that short-term temperature upshifts are required to

establish spindles and maintain viability, suggesting some portion of centrosomal trafficking during

mitosis still occurs. For this reason, we suspected the combination of these treatments could most

severely limit dynein trafficking. A nocodazole-treated, *dhc-1(ts)*-containing organism at the restrictive

temperature will likely exhibit the most severely affected dynein trafficking and thus most severely

disrupted regulation of SYS-1 on the mitotic centrosome.

In fact, studying the combination of these challenges to microtubule-mediated trafficking

allowed us to examine several facets of SYS-1 localization. Due to the difference in response to

nocodazole based on the promoter driving fluorescently-tagged SYS-1 (Fig 4B, D), we were able to utilize

their apparent difference in SYS-1 expression to slightly sensitize organisms to SYS-1/negative regulator

defects. In contrast to the GFP::SYS-1 driven by a promoter for ubiquitous early embryonic PIE-1,

organisms with YFP::SYS-1 driven by the SYS-1 promoter exhibit only a slight, non-significant,

nocodazole-mediated increase in CEI/TEI (Fig. 4D, 1^{st} and 2^{nd} columns). This indicates that the mild

overexpression seen in PIE-1 promoter-driven SYS-1 may be sufficient to stress SYS-1 clearance

mechanisms, allowing it to more readily overwhelm them in nocodazole treatment. Meanwhile,

endogenously driven SYS-1 may be maintained at levels more manageable by the centrosomal clearance

mechanism, resulting in a lesser or variable CEI/TEI increase. Importantly, endogenously driven SYS-1 is

sufficient to overwhelm centrosomal clearance mechanisms when the hypomorphic *dhc-1(ts)* at

permissive temperature is treated with nocodazole, resulting in the same significant increase seen with

GFP::SYS-1 (Fig. 4D 3^{rd} and 4^{th} columns, Fig. 4C). However, this increase is no longer observed when the

dhc-1(ts) organisms are upshifted to the restrictive temperature, indicating that the increasing SYS-1

centrosomal enrichment seen in Nocodazole treated embryos still requires some amount of dynein

trafficking. These observations describe a centrosomal relationship wherein treatments limiting the

coverage of microtubule-mediated dynein trafficking (i.e. nocodazole treatment) increase centrosomal

SYS-1 due to relatively inefficient or rare trafficking events for SYS-1 negative regulators, while transient

severe disruptions to dynein trafficking processivity (dynein subunit RNAi depletion, *dhc-1(ts)*

temperature upshift, upshift + nocodazole) reveal that SYS-1 centrosomal enrichment requires continual

replenishment via a functional microtubule motor.

If microtubule-mediated trafficking has a potential role for proteasome or other SYS-1 negative

regulator centrosomal enrichment, concurrent with SYS-1 localization, we expect approximately similar

localization changes as we would for SYS-1. Therefore, we decided to test the idea that proteasomal

trafficking to, and enrichment at, the centrosome was also perturbed in nocodazole-affected

microtubule networks. We first attempted to localize the proteasome. While the centrosome has been

demonstrated to exhibit proteolytic activity (Vora and Phillips 2015; Vora and Phillips 2016; Hames, Fry

et al 2005; Peel, O'Connell et al 2012; Wigley, Thomas et al 1999; Fabunmi, DeMartino et al 2000),

existing immunohistochemical reagents for mammalian proteasome subunits did not display any distinct

localization pattern. An mCherry tag on proteasome subunit RPT-1 did demonstrate a kinetochore

microtubule enrichment like that of both DHC-1 and SYS-1 but appeared absent from centrosomes. This

may be due to a relatively low RPT-1 copy number per proteasome, or correspondingly low proteasome

number per centrosome, limiting its relative enrichment there. Alternatively, fluorescently tagged

proteins have shown evidence of degradation of the tag at a site of active proteolysis, particularly if such

a tag limits proteasome activity (Huang, Lobel et al 2014). To circumvent this issue, we assayed for

centrosomal proteasome activity indirectly by functionally testing an additional target of the

centrosomal proteasome, the centriole duplicating kinase ZYG-1 (O'Connell, White et al 2001; Peel,

O'Connell et al 2012). In addition, we assayed the centrosomal role of ECPS-1, homolog of mammalian

ECM29 and an adaptor protein proposed to enhance proteolytic activity of proteasomes via its function

as an adapter between the dynein motor complex and the proteasome (Gorbea, Rechsteiner et al 2010).

Knockdown of ECM29 homolog ECPS-1 (formerly d2045.2) resulted in embryos that displayed an

average 33.4% increase in centrosomal GFP::SYS-1 centrosomal enrichment, similar to both Nocodazole-

treated and *rpt-4(*RNAi) embryos (Fig. 5A, 4 A-C). Given the role of ECPS-1/ECM29 in proteasomal

trafficking on MTs rather than direct protein degradation, these data suggest that both SYS-1 and its

negative regulator, the proteasome, are trafficked to the centrosome during mitosis. Therefore, the

overall increase in SYS-1 CEI seen in *ecps-1*(RNAi) embryos is likely an adjustment of the centrosomal

steady state, limiting the trafficking of SYS-1 negative regulators without affecting that of SYS-1. While

fluorescently-tagged SYS-1 exhibits widespread localization and expression, allowing for its capture and

transport on even a limited microtubule network, the several megadalton proteasome may be

dependent on extensive active trafficking. This necessity may be fostered due to both the size and

stoichiometry required of the proteasome complex. Indeed, electron micrographs of ECM29, and its

interactions with the proteasome, demonstrate that the interaction between ECM29 and the

proteasome is at the level of the core 20S proteasome (Leggett, Finley et al 2002). Because of this,

longer term microtubule disruption like that of our Nocodazole treatment may prevent the

enhancement of centrosomal SYS-1-proteasome interactions while not necessarily preventing SYS-1

degradation. That is, the SYS-1 – proteasome interaction is less efficient when not centrosomally

concentrated, and prevention of this co-localization should result in increased SYS-1 levels at the

centrosome and throughout the cell. To test the activity of the centrosomal proteasome after ECPS-1

RNAi treatment, we assayed an additional centrosomal proteasome target – a regulatory kinase of

centrosomal duplication called ZYG-1 (Peel, O'Connell et al 2012). Utilizing a fluorescently tagged ZYG-1

fragment demonstrated to mirror the localization of the endogenous protein (Peters, O'Connell et al

2010), we demonstrated 20% and 27% increased accumulation of ZYG-1 in both proteasome knockdown

rpt-4(RNAi) and *ecps-1(*RNAi) treatments, respectively, supporting the hypothesis that these shared

phenotypes are both proteasome-mediated clearance defects (Fig 5C).

325   Since SYS-1 centrosomal recruitment and regulation of subsequent inheritance depend on rapid

degradation via the centrosomal proteasome (Vora and Phillips 2015), we assayed the accumulating

centrosomal SYS-1 in *ecps-1(RNAi)* for changes to its steady state dynamics by FRAP. This allowed us to

distinguish between increased centrosomal accumulation due to an increased recruitable pool of

cytoplasmic SYS-1 when ECPS-1 is depleted as compared to a change in centrosomal steady state due to

specific changes to the proteasomal throughput in this locale. Comparison of GFP::SYS-1 recovery after

photobleach demonstrates that depletion of ECPS exhibits an intermediate phenotype between

proteasome depletion via *rpt-4(RNAi)* and controls (5G-G"). While the initial rate of *ecps-1(RNAi)*

GFP::SYS-1 recovery was similar to wild type organisms, the late recovery phase of *ecps-1(RNAi)*

organisms showed an early SYS-1 recovery plateau, evidence of a significantly reduced mobile fraction

intermediate to wild type and that of proteasome depleted *rpt-4(RNAI)* organisms (Fig 5E,G). This

suggests that SYS-1 recovers at an approximately wild type rate in *ecps-1(RNAi)* organisms, but remains

approximately as immobile as proteasome-depleted organisms. Therefore, ECPS-1 depleted organisms

appear to degrade SYS-1 incompletely, as observed in the RPT-4 proteasome depletion, but with

sufficient protein turnover to enable wild type recruitment of newly trafficked SYS-1.

This is in contrast to the dynamics observed in other centrosomal proteins. GFP::ZYG-1c exhibits

slightly different centrosomal localization dynamics during mitosis from SYS-1, peaking in centrosomal

representation mid-anaphase (Peel, O'Connell et al 2012). This suggested that GFP::ZYG-1c is

proteasomally degraded, but is most dynamic during early mitosis, maintaining a consistent centrosomal

enrichment in anaphase even in *rpt-4(RNAi)* organisms (Peel, O'Connell et al 2012). SYS-1, conversely,

exhibits strong centrosomal SYS-1 localization through anaphase and only begins to show decreasing

centrosomal levels during telophase (Vora and Phillips 2015). This apparent difference in dynamics is

made evident by comparing the recovery after photobleach of GFP::SYS-1 and GFP::ZYG-1c (Fig. 5E-F),

wherein both SYS-1 and ZYG-1c become less mobile upon *rpt-4(RNAi)* proteasome depletion but only

SYS-1 increases in immobile fraction upon ECPS-1 depletion. This suggests that proteasome depletion

by *rpt-4(RNAi)* leads to decreased proteolytic activity that overwhelms and stabilizes the centrosomal

proteome, including SYS-1, by preventing turnover. Meanwhile, ECPS-1 depleted organisms may be

unaffected in protein clearance per proteasome, but the delayed or partial localization of the

centrosomal proteasome could result in hierarchical stabilization of a spatially or temporally inaccessible

fraction consisting of highly expressed and continually trafficked proteins, like SYS-1.

Dynein Trafficking is Required for Restriction of Wnt-Signaled Cell Fates

Having established a two-fold role for SYS-1 regulation in microtubule-mediated trafficking, we

investigated the role microtubule trafficking played in regulating SYS-1 dependent cell fate decisions.

Since the SYS-1 centrosomal localization pattern has been seen across all examined SYS-1-expressing

tissues (Phillips, Kimble et al 2007; Huang, Lin et al 2007; Baldwin and Phillips 2018; Lam and Phillips

2017), we investigated cell fate changes in embryonic and larval tissues across one embryonic and two

larval SYS-1 dependent cell fate decisions. These included the transcriptional reporter of the WβA target

gene, *end-1,* in the establishment of SYS-1 dependent endoderm, and an anatomical marker for larval

specification of the stem cell-like cells of the hypodermal seam and the germline stem cell niche. First, to

directly assay the effect of centrosomal dysregulation of transcriptional activator SYS-1, we investigated

the nuclear enrichment of GFP::SYS-1 after the Wnt-regulated EMS division (Bei, Mello et al 2002). In

this division, SYS-1 is enriched in the posterior E nucleus, but not that of the anterior MS (Fig. 6A-B).

Analyzing nuclear enrichment of GFP-tagged SYS-1, we were able to observe increased nuclear GFP

signal in both the E and MS daughters in analyzed trafficking mutants (Fig 6C). Interestingly, daughters

affected by both *ecps-1*(RNAi) and *dlc-1*(RNAi) exhibit significant enhancement of nuclear enrichment in

each of the two daughter cells (fig 6C), suggesting that these treatments act negatively on SYS-1, despite

opposing effect on SYS-1 centrosomal enrichment. Following this loss of negative centrosomal

regulation, these data suggest that daughter cell SYS-1 inheritance is increasing and thus affecting the

bias of this asymmetric division, increasing the likelihood of unsignaled daughter cells adopting the

signaled cell fate.

Having established an increase in nuclear GFP::SYS-1 in dynein subunit depletion, even in cells

not signaled to maintain activating SYS-1, we turned to the effect of this SYS-1 expression on target gene

activation. Specifically, we used the endogenous promoter of WβA target gene *end-1* to drive GFP-

tagged Histone 2B in order to evaluate the activity of SYS-1 immediate downstream targets as the

signaled E cell established the endoderm tissue expansion from 2 to 4 to 8 cells (E+1, 2, or 3 divisions,

respectively). Normally, this strain exhibits additional expression of the *end-1* promoter in <10% of

embryos, a frequency which decreases further when SYS-1 is depleted via RNAi (Fig. 6D).. In our

embryos depleted of dynein subunits and RSA-2 by RNAi, however, we observed a different pattern. In

these embryos, misregulation via dynein depletion appeared to instead increasingly overwhelm SYS-1

negative regulation in successive divisions, such that each generation after this SYS-1 dependent

specification increasingly expresses this endodermal reporter ectopically outside of the EMS lineage (Fig

6D). For instance, in *dhc-1(RNAi)* we observe only 3% of embryos mis-expressing one cell division after

the specification of the E cell type (E+1), which increases to 25% by the next division (E+2), and over 83%

three divisions after specification (E+3).

We also investigated larval ACDs to determine the effect this extra transcriptionally active SYS-1

has on organismal development. We examined the seam cells (SCs) and distal tip cells (DTCs) due to

their sensitivity to SYS-1 dependent gene activation (Baldwin, Phillips et al 2016; Baldwin and Phillips

2018; Lam and Phillips 2017; Mila, Putzke et al 2015). In wild-type animals, DTC lineages undergo

asymmetric cell division during L1 to give rise to two DTCs, while *sys-1* mutants and organisms

overexpressing SYS-1 show loss and gain of DTCs, respectively, due to symmetric cell division (fig 7A)

(Miskowski, Kimble et al 2001; Siegfried, Kimble et al 2004; Kidd, Kimble et al 2005; Chesney, Kimble et

al 2009). To further sensitize these cells for changes in SYS-1 dependent cell fate changes, we examined

strains containing additional, transgenic SYS-1 driven by the heat shock promoter. In organisms that are

not subjected to heat shock, dynein depletion only rarely induced development of more than the

endogenous 2 distal tip or 16 seam cells (Fig 7B, 8B). In heat shocked organisms, excess SYS-1 protein

can overwhelm canonical SYS-1 regulation to develop as many as 6 DTCs (fig 7D). While heat shock-

induced overexpression is sufficient to cause a notable shift towards the signaled cell fate, dynein

subunit depletion was able to significantly enhance this (Fig 7C). In the DTC, depletion of DHC-1 or ECPS-

1 was sufficient to induce rare but wild-type-unobserved additional distal tip cells (0.3% of organisms for

these 2 treatments, N= 394, 363, respectively)). Adding heat shock-overexpressed SYS-1 enhanced the

penetrance of ectopic DTCs, with DHC-1 and ECPS-1 depletion resulting in populations where 11.9% and

6.8% of organisms, respectively, develop more than 2 DTCs. Depletion of RSA-2, uncoupling SYS-1 from

centrosomes, and depletion of a dynein light chain, DYLT-1, further enhanced these misspecifications,

increasing the proportion of the population with ectopic DTCs from 26.6% in wild type to 33.6% or 51%

after RNAi depletion, respectively (Fig. 7).

The SC lineage also undergoes several WBA-dependent ACDs throughout larval development (fig

8A). Similarly, in the SC tissue, depletion of dynein light chains alone is sufficient to increase the

frequency of ectopic SCs from 3.3% of larvae with ectopics SCs in control RNAi conditions to 11.7% or

10% of organisms as seen in DYLT-1 and DLC-1 depletions, respectively (Fig 8B). These misspecifications

are further exacerbated by SYS-1 overexpression, wherein more than 60% of the *dylt-1(RNAi)* organisms

have greater than wild type seam cell numbers, developing as many as 21 SCs, (Fig. 8C). These

observations collectively support SYS-1 centrosomal enrichment by dynein as a mechanism to reinforce

SYS-1 function in specifying SYS-1-dependent cell fate decisions. Therefore, compromising this system by

dynein depletion is rarely sufficient to induce misspecifications in otherwise wildtype animals, but

notably enhances the effects of ectopic SYS-1 accumulation, indicating the robustness of WβA-regulated

cell fate decisions after ACD requires SYS-1 negative regulation via dynein-mediated MT transport.

<u>Discussion</u>

Precise control of cell fate determinants is critical for normal development and homeostasis,

particularly in the rapidly dividing cells of the early embryo. This regulation includes the distribution and

inheritance of these determinants by daughter cells. Here we demonstrate that centrosomal negative

regulation of *C. elegans* β-catenin SYS-1 is enhanced during asymmetric division by trafficking to

centrosomes by the microtubule motor dynein. Depletion of individual subunits of the dynein motor

and associated proteins is sufficient to reduce the centrosomal SYS-1 steady state, and temporally

specific ablation of dynein function by *dhc-1(ts)* was sufficient to nearly completely and

disproportionately remove observable SYS-1 from centrosomes. However, some methods of disrupting

dynein/microtubule trafficking result in an unexpected increase in centrosomal accumulation of SYS-1

protein. These increases, caused by microtubule disruption via Nocodazole and depletion of SYS-1

negative regulator adapter ECPS-1, were accompanied by SYS-1 CEI steady state adjustments that

phenocopy centrosomal enrichment seen in disruption of proteasome activity, and further maintain a

SYS-1 immobile fraction like that seen in proteasome disruption experiments. We therefore propose a

model wherein both SYS-1 and its negative regulators, including the proteasome, are dependent on

microtubule-mediated trafficking to centrosomes by the dynein complex in order to appropriately

reinforce SYS-1 asymmetric cell divisions (Fig 9). In this model, SYS-1 requires continual trafficking by

microtubule motors to maintain its observable steady state, suggesting highly efficient interactions with

the dynein molecule while exhibiting vulnerability to subtle disruptions to dynein trafficking. Conversely,

the enzymatic substrate processing of the proteasome requires fewer proteasome complexes to be

trafficked, making it less vulnerable to short term disruptions to trafficking but requiring a robust

microtubule network to secure these relatively fewer events.

While this system of centrosomal localization and clearance of SYS-1 does not appear necessary

for most WβA differentiation events, disruptions to this system increasingly limit its ability to tolerate

perturbations. This suggests that SYS-1 centrosomal localization thereby generates robustness in SYS-1

dependent cell fate decisions (Fig 6-8). This set of parallel and synergistic trafficking mechanisms

accounts for the similar excess SYS-1 found in affected daughter cells of mothers with both increased

and decreased centrosomal recruitment of SYS-1. That is, disruption of either SYS-1 localization

(inefficient trafficking to the site of regulation) or disruption of proteasomal localization (SYS-1 is

trafficked and accumulates, but centrosomal degradation is ineffective) both result in increased SYS-1

protein in daughter cells, consistent with excessive inheritance.

Why does no single dynein disruption treatment fully recapitulate the *rsa-2(RNAi)* phenotype?

While we do see individuals within several dynein knockdown treatments with nearly *rsa-2(RNAi)*-like

SYS-1 centrosomal uncoupling, these populations also include organisms with roughly wild type

centrosomal SYS-1, unlike those in *rsa-2*(RNAi). However, given the severely deleterious nature of

dynein subunit loss-of-functions (Sönnichsen, Neumann et al 2005) and the variability of feeding RNAi

knockdowns as compared to null mutant strains (Tavernarakis, Driscoll et al 2000), it seems likely that

our significant decreases in SYS-1 centrosomal recruitment represent either an intermediate knockdown

phenotype – the ideal hypomorphy between the extremes of catastrophic defects in spindle assembly

and organisms insufficiently affected to demonstrate a phenotype – or indicate that there is partial

redundancy between the SYS-1 relevant functions of the dynein motor. Our *dhc-1(ts)* experiments

support either case by clearly demonstrating a thorough dynein knockdown able to completely

uncouple centrosomal SYS-1. Alternatively, there could be partial redundancy among auxiliary dynein

components for SYS-1 binding/processivity, a possibility suggested by our *dhc-1(ts)* data which showed

the most severe reductions in centrosomal SYS-1.

Why, then, do we observe an increase in centrosomal SYS-1 steady state upon either long term

lower dose nocodazole-mediated knockdowns of trafficking or by RNAi depletion of trafficking subunit

ECPS-1? In response to these data (fig 4), we propose the additional SYS-1 negative regulator trafficking

component to this model. Disrupting the trafficking of centrosomal SYS-1 negative regulators should

result in a previously unseen increase in centrosomal SYS-1 steady state while maintaining a similar

effect on downstream gene expression and cell fate decisions, as seen above (Fig. 4-8). However, the

dichotomy between most dynein depletions and that of ECPS-1 or nocodazole treatment suggests a

differential efficiency in trafficking or centrosomal capture rate between SYS-1 and its negative

regulators, such as the proteasome. Most disruptions to the dynein motor that reduce SYS-1

centrosomal accumulation, mediated by maternally fed RNAi, are consistent but incomplete due to their

requirement for oogenesis and early embryonic viability. This, therefore, should produce embryos

capable of forming wild type dynein complexes, but at a far reduced rate. These dynein-limited embryos

should therefore decrease the volume of SYS-1 capable to be trafficked while not preventing the

centrosomal localization of an occasional proteasome complex sufficient to enzymatically process the

decreased SYS-1 flux. Similarly, depletion of dynein function by *dhc-1(ts)* effectively depletes all dynein

function but can only do so for brief portions of mitosis without inducing mitotic spindle collapse.

While these depletions limit the volume of trafficking by limiting the number of properly

constructed dynein motors, they don't limit the ability of an individual, functional dynein motor to

traffic. The experimental conditions that increase the steady state of centrosomal SYS-1, Nocodazole

and *ecps-1(RNAi)*, instead limit the scope of dynein cargo trafficking. ECPS-1, putative adapter between

dynein and the proteasome, increases the steady state of both SYS-1 and additional centrosomal

proteasome target ZYG-1. This supports the conserved role of ECPS-1, and results in embryos that are

not inhibited for overall dynein function but limited in their ability to traffic proteasome. Nocodazole

treatment similarly does not directly affect the processivity of dynein complexes that form on the

comparatively limited microtubule network. However, it does severely affect the reach of the

microtubule network itself (Fig 4A). Our data therefore suggests that the proteasome requires less

continual reinforcement by microtubule-mediated trafficking but does require more extensive

microtubule formation than is allowed in our Nocodazole-treated embryos.

A role for trafficking in both SYS-1 and its regulatory components, then, suggests a differentially

maintained steady state interaction for the proteasome and centrosomal SYS-1. While SYS-1 is rapidly

and continuously trafficked, it is similarly rapidly processed by centrosomal proteasome. That is, the

apparently rapid processing of centrosomal SYS-1 also requires rapid dynein-mediated flux to maintain

its centrosomal steady state. Any disruption to dynein processivity will therefore allow SYS-1 enrichment

to be overcome by its centrosomal processing, reducing the SYS-1 CEI by slowing the transfer of SYS-1

from cytoplasm to centrosome for degradation. By comparison, the proteasome complex endures at

the centrosome and continually processes SYS-1. However, the mass of the multi-megadalton complex,

or requisite assembling all 30 members of the 26S proteasome in stoichiometrically relevant numbers

(Papaevgeniou and Chondrogianni, 2014), suggests a requirement for continuous and highly processive

dynein trafficking to ensure its establishment at the centrosome. Our proposed model therefore

predicts that short term or partial disruptions to the dynein motor (Fig 9, red inset) allow for the

eventual establishment of a centrosomal proteolysis pathway, which may affect both SYS-1 and

proteasome localization, but nevertheless result in decreasing steady state levels of centrosomal SYS-1

(and presumably other proteasome targets that require active trafficking) as occasional proteasome

centrosomal localization overcomes the impaired localization of SYS-1. Conversely, long term systemic

disruptions to microtubule-mediated motors like that seen in Nocodazole treatment may still capture

SYS-1 in their comparatively limited microtubule network, while being disproportionately deprived of

the larger proteasome complex (Fig 9, blue inset).

Our model that both SYS-1 and its regulators are trafficked was supported but the identification

of a trafficking component that affects one but not both arms of our trafficking model. Most of our

perturbations, generally affecting more core components of the dynein machinery, may exert their

effects by detaching SYS-1 from the motor or by sabotaging all function of the motor complex (Fig S3A).

However, the specificity of our two transport mechanisms was exhibited by one dynein-protein

interaction, via depletion of ECPS-1/the putative ECM29 subunit of the motor. Unlike the other subunits

presenting strong SYS-1 phenotypes, ECM29 has not yet been implicated in dynein function beyond its

adaptor role but has notable effects on proteasome localization and response to stress (Gorbea,

Rechsteiner et al 2010; Haratake, Chiba et al 2016). Because *ecps-1(RNAi)* phenocopies the centrosomal

GFP::SYS-1 accumulation seen in proteasome knockdown via *rpt-4(RNAi)* as well as microtubule network

disruption via Nocodazole treatment, we propose that these observations support a role for

proteasomal localization as well as function for proper centrosomal clearance of SYS-1 (Fig 9).

While this model and our experiments to test it have been largely centered on SYS-1/β-catenin,

many developmental and temporally regulated proteins are known to pass through the centrosome

(Vora and Phillips 2016; Brown, Welch et al 1994; Johnston, Kopito et al 1998; Huang and Raff 1999;

Reck-Peterson, Carter et al 2018), as the centrosomal proteasome is far from SYS-1-specific. It is not

unreasonable, then, to presume that other proteins may benefit from dynein trafficking to something as

short-lived and yet influential to daughter cell inheritance as the mitotic centrosome. Particularly early

in eukaryotic development, subtle changes in the inheritance of cell fate determinants like SYS-1 can

have a profound effect on the ability of an organism to respond to environmental insults. For example,

the asymmetric nuclear localization of SYS-1 indicative of differential cell fate has been shown to arise

early, less than 70 seconds after the establishing division (Baldwin and Phillips 2014; Vora and Phillips

2015). We therefore propose that an active dynein mediated trafficking, at least for cell fate

determinants like SYS-1, could provide a universal buffering against excessive inheritance and

consequentially rare but deleterious cell fate changes.

Materials and Methods

Strains

Strains were maintained on Op50-inoculated NGM plates at 15° (BTP51, BTP220) or 25° C

(TX964, BTP51, LP563, TX691) using typical *C. elegans* methods. These strains contained the following

transgenes and alleles: **N2** (wild type); **TX964** (unc-119(ed3) III; him-3(e1147) IV; teis98 [Ppie-

1::GFP::SYS-1]); **BTP51** (unc-119(ed3) III; tjIs71 [pie-1 promoter::mCherry::H2B, pie-1 promoter::2x

mCherry::tbg-1, unc-119(+)]; qIs95 III) ; **BTP220** (unc-119(ed3) III; tjIs71 [pie-1 promoter::mCherry::H2B,

pie-1 promoter::2x mCherry::tbg-1, unc-119(+)]; qIs95 III; dhc-1(or195) I.) ; **TX691** (unc-119(ed3) III;

teIs46 [pRL1417; end-1p::GFP::H2B + unc-119(+)]) ; **OC341 (**unc-119(ed3) III; bsIs8[pMS5.1:unc -119(+)

pie -1- gfp-zyg-1C-terminus]); **BTP93** (*wIs78[SCMp::GFP + ajm-1p::GFP + F58E10* (cosmid) + *unc-119(+)]*

IV; uiwEx22[pHS::sys-1]); **BTP97** (*qIs56[lag-2p::GFP + unc-119(+)] V;uiwIs3[pHS::sys-1]*); **AZ235** (*unc-*

119(ed3) III; ruIs48[Ppie-1::gfp::tbb-1];

RNAi

To perform RNAi knockdown of target genes, we used HT115 bacteria containing the pL4440

plasmid with a T7-flanked target gene insert. Most of these were obtained from a library supplied by

Ahringer (Kamath and Ahringer 2003). Those which were otherwise obtained (dlc-1) used target gene-

flanking primers with *C. elegans* cDNA to generate a new insert. These bacteria were used to seed IPTG-

containing, inducing plates as described previously. Worms were generally plated to RNAi after Sodium

Hypochlorite Synchronization as L1's. More deleterious knockdowns (*dlc-1, dhc-1)* were washed from

OP50 plates at 24 hours before imaging (L3/L4) or 12 hours before imaging (late L4/early adult)

(Timmons 2006; Bekas and Phillips 2020). DLC-1 WAS DERIVED USING RT PRIMERS DIGESTED WITH Xho I

and HIND III SEQUENCING 7/31

Compound Microscopy, CEI, and Image Processing

To obtain samples, mothers were anesthetized with 250 µM levamisole and dissected *via* a

clean surgical blade. Embryos were then transferred to a 2% agarose pad and imaged on a Zeiss Axio

Imager.D2. All exposure times were kept constant. Images were exported as uncompressed TIF's, and

fluorescence intensity was determined by measuring the mean fluorescence of centrosomal puncta in

ImageJ. The mean intensity of the remaining embryonic cytoplasm was subtracted from the centrosomal

mean to produce the CEI measurement, and identical measurements on 2x mCherry::tbg-1 were used

for TEI. END-1 family nuclei were assayed via the TX691 strain. GFP-tagged Histone intensity was

measured relative to the cytoplasmic autofluorescence.

Temperature Sensitive Allele Imaging

BTP51 and BTP220 were maintained together at 15°C and were imaged as above. Ambient

temperature in the microscope room was decreased to ~18°C for 'permissive' imaging and upshifted to

'restrictive' 26°C by moving slides to a Pecon Tempcontroller 2000-2 heated stage. Embryos were

imaged at the two temperatures sequentially, with approx. 45 seconds between upshift and image, or

were imaged as populations at either the permissive or restrictive temperature.

Nocodazole Treatment

In order to treat embryos before formation of the drug-impermeable eggshell, we exposed late

L4's to a low dose (approx. 83 μM, 25 μg/mL) nocodazole dissolved in 2% DMSO for 16 hours. Organisms

were then harvested by one minute centrifugation at 1000xG and imaged as described above.

Confocal Microscopy and Fluorescence Recovery After Photobleach

Confocal microscopy was performed via a Leica SP8 HyD detector system. The objective used

was a 63x HC PL APO CS2 objective with N.A. = 1.4, utilizing type F immersion oil. Each image analyzed

consisted of the sum of 35 z-images across the embryo, each slice approximately 0.35 μM thick. FRAP

was performed on the same system. Each embryo was imaged twice before bleaching, beginning as

soon as possible after NEBD. The ROI, an approximately 5μm square, was then bleached at an

intermediate focal plane with 100% laser power for 100 iterations. Images were taken as above, once

every 7 seconds. Fluorescence intensity was measured on summed slice Z-projections in FIJI via the

mean fluorescence intensity of ROIs drawn around relevant locales – either centrosome or the entire

dividing cell. Recovery was assayed by subtracting the ROI post-bleach intensity from each

measurement and evaluating it as a percentage of the pre-bleached centrosomal intensity.

Heat Shock-Induced SYS-1 Overexpression

BTP93 and BTP97 worms were exposed to brief 33℃ heat shocks at relevant developmental

timepoints to drive SYS-1 overexpression concurrently with signaled cell fate specification. For DTCs,

BTP97 was subjected to 1-hour heat shock, 30-minutes recovery, and 30-minutes additional heat shock

beginning 11.5 hours after plating. For SCs, BTP93 was subjected to 2-hour heat shock, 30-minute

recovery, and additional 1-hour heat shock beginning 26 hours after plating. Both tissue types were

assayed 50 hours after plating.

Acknowledgements

We thank Kimberley Bekas and Amy Clemons for helpful comments on the manuscript. Strains

were provided by the Caenorhabditis Genetics Center, which is funded by the National Institutes of

Health (NIH) Office of Research Infrastructure Programs [P40 OD01440] and Dr. Kevin F. O'Connell at the

NIDDK (OC341). This work was supported by NIH award NIH GM114007 (B.T.P.).

605
606

References

Amsterdam, A., F. Pitzer and W. Baumeister, 1993 Changes in intracellular localization of proteasomes in

immortalized ovarian granulosa cells during mitosis associated with a role in cell cycle control.

Proc Natl Acad Sci U S A 90: 99-103.

Baldin, V., M. Militello, Y. Thomas, C. Doucet, W. Fic *et al.*, 2008 A novel role for PA28gamma-

proteasome in nuclear speckle organization and SR protein trafficking. Mol Biol Cell 19: 1706-

1716.

Baldwin, A. T., A. M. Clemons and B. T. Phillips, 2016 Unique and redundant beta-catenin regulatory

roles of two Dishevelled paralogs during C. elegans asymmetric cell division. J Cell Sci 129: 983-

993.

Baldwin, A. T., and B. T. Phillips, 2014 The tumor suppressor APC differentially regulates multiple beta-

catenins through the function of axin and CKIalpha during C. elegans asymmetric stem cell

divisions. J Cell Sci 127: 2771-2781.

Baldwin, A. T., and B. T. Phillips, 2018 Chapter 3 - Cell Polarity and Asymmetric Cell Division by the Wnt

Morphogen, pp. 61-102 in *Cell Polarity in Development and Disease*, edited by P. Michael Conn.

Academic Press, Boston.

Bei, Y., J. Hogan, L. A. Berkowitz, M. Soto, C. E. Rocheleau *et al.*, 2002 SRC-1 and Wnt signaling act

together to specify endoderm and to control cleavage orientation in early C. elegans embryos.

Dev Cell 3: 113-125.

Bekas, K. N., and B. T. Phillips, 2020 Generating reliable hypomorphic phenocopies by RNAi using long

dsRNA as diluent. MicroPubl Biol 2020.

Ben-Yaakov, K., S. Y. Dagan, Y. Segal-Ruder, O. Shalem, D. Vuppalanchi *et al.*, 2012 Axonal transcription

factors signal retrogradely in lesioned peripheral nerve. EMBO J 31: 1350-1363.

Bullock, S. L., and D. Ish-Horowicz, 2001 Conserved signals and machinery for RNA transport in

Drosophila oogenesis and embryogenesis. Nature 414: 611-616.

Carminati, J. L., and T. Stearns, 1997 Microtubules orient the mitotic spindle in yeast through dynein-

dependent interactions with the cell cortex. J Cell Biol 138: 629-641.

Chesney, M. A., N. Lam, D. E. Morgan, B. T. Phillips and J. Kimble, 2009 C. elegans HLH-2/E/Daughterless

controls key regulatory cells during gonadogenesis. Dev Biol 331: 14-25.

del Castillo, U., M. Winding, W. Lu and V. I. Gelfand, 2015 Interplay between kinesin-1 and cortical

dynein during axonal outgrowth and microtubule organization in Drosophila neurons. Elife 4:

e10140.

Deshimaru, M., M. Kinoshita-Kawada, K. Kubota, T. Watanabe, Y. Tanaka *et al.*, 2021 DCTN1 Binds to

TDP-43 and Regulates TDP-43 Aggregation. Int J Mol Sci 22.

Fabunmi, R., W. Wigley, P. Thomas and G. DeMartino, 2000 Activity and Regulation of the Centrosome-

associated Proteasome. Journal of Biological Chemistry 275: 409-413.

Fan, J., Q. Zhang, H. Tochio, M. Li and M. Zhang, 2001 Structural basis of diverse sequence-dependent

target recognition by the 8 kDa dynein light chain. J Mol Biol 306: 97-108.

Fuchs, E., and T. Chen, 2013 A matter of life and death: self-renewal in stem cells. EMBO Rep 14: 39-48.

Gao, F. J., L. Shi, T. Hines, S. Hebbar, K. L. Neufeld and D. S. Smith (2017). "Insulin signaling regulates a

functional interaction between adenomatous polyposis coli and cytoplasmic dynein." Mol Biol

Cell 28(5): 587-599.

Geyer, F. C., M. Lacroix-Triki, K. Savage, M. Arnedos, M. B. Lambros *et al.*, 2011 beta-Catenin pathway

activation in breast cancer is associated with triple-negative phenotype but not with CTNNB1

mutation. Mod Pathol 24: 209-231.

Gómez-López, S., R. G. Lerner and C. Petritsch, 2014 Asymmetric cell division of stem and progenitor

cells during homeostasis and cancer. Cell Mol Life Sci 71: 575-597.

Gorbea, C., G. Pratt, V. Ustrell, R. Bell, S. Sahasrabudhe *et al.*, 2010 A protein interaction network for

Ecm29 links the 26 S proteasome to molecular motors and endosomal components. J Biol Chem

285: 31616-31633.

Hames, R. S., R. E. Crookes, K. R. Straatman, A. Merdes, M. J. Hayes *et al.*, 2005 Dynamic recruitment of

Nek2 kinase to the centrosome involves microtubules, PCM-1, and localized proteasomal

degradation. Mol Biol Cell 16: 1711-1724.

Hamill, D. R., A. F. Severson, J. C. Carter and B. Bowerman, 2002 Centrosome maturation and mitotic

spindle assembly in C. elegans require SPD-5, a protein with multiple coiled-coil domains. Dev

Cell 3: 673-684.

Hanz, S., E. Perlson, D. Willis, J. Q. Zheng, R. Massarwa *et al.*, 2003 Axoplasmic importins enable

retrograde injury signaling in lesioned nerve. Neuron 40: 1095-1104.

Harada, A., Y. Takei, Y. Kanai, Y. Tanaka, S. Nonaka *et al.*, 1998 Golgi vesiculation and lysosome

dispersion in cells lacking cytoplasmic dynein. J Cell Biol 141: 51-59.

Haratake, K., A. Sato, F. Tsuruta and T. Chiba, 2016 KIAA0368-deficiency affects disassembly of 26S

proteasome under oxidative stress condition. J Biochem 159: 609-618.

He, T. C., A. B. Sparks, C. Rago, H. Hermeking, L. Zawel *et al.*, 1998 Identification of c-MYC as a target of

the APC pathway. Science 281: 1509-1512.

Horgan, C. P., S. R. Hanscom, R. S. Jolly, C. E. Futter and M. W. McCaffrey, 2010 Rab11-FIP3 links the

Rab11 GTPase and cytoplasmic dynein to mediate transport to the endosomal-recycling

compartment. J Cell Sci 123: 181-191.

Huang, L., D. Pike, D. E. Sleat, V. Nanda and P. Lobel, 2014 Potential pitfalls and solutions for use of

fluorescent fusion proteins to study the lysosome. PLoS One 9: e88893.

Huang, S., P. Shetty, S. M. Robertson and R. Lin, 2007 Binary cell fate specification during C. elegans

embryogenesis driven by reiterated reciprocal asymmetry of TCF POP-1 and its coactivator beta-

catenin SYS-1. Development 134: 2685-2695.

Hutagalung, A. H., and P. J. Novick, 2011 Role of Rab GTPases in membrane traffic and cell physiology.

Physiol Rev 91: 119-149.

Jha, R., and T. Surrey, 2015 Regulation of processive motion and microtubule localization of cytoplasmic

dynein. Biochem Soc Trans 43: 48-57.

Juanes, M. A., 2020 Cytoskeletal Control and Wnt Signaling-APC's Dual Contributions in Stem Cell

Division and Colorectal Cancer. Cancers (Basel) 12.

Kamath, R. S., A. G. Fraser, Y. Dong, G. Poulin, R. Durbin *et al.*, 2003 Systematic functional analysis of the

Caenorhabditis elegans genome using RNAi. Nature 421: 231-237.

Kawaguchi, Y., J. J. Kovacs, A. McLaurin, J. M. Vance, A. Ito *et al.*, 2003 The deacetylase HDAC6 regulates

aggresome formation and cell viability in response to misfolded protein stress. Cell 115: 727-

738.

Kidd, A. R., 3rd, J. A. Miskowski, K. R. Siegfried, H. Sawa and J. Kimble, 2005 A beta-catenin identified by

functional rather than sequence criteria and its role in Wnt/MAPK signaling. Cell 121: 761-772.

Lam, A. K., and B. T. Phillips, 2017 Wnt Signaling Polarizes C. elegans Asymmetric Cell Divisions During

Development. Results Probl Cell Differ 61: 83-114.

Leduc, C., K. Padberg-Gehle, V. Varga, D. Helbing, S. Diez *et al.*, 2012 [How to avoid the formation of

molecular motors traffic jam during intracellular transport]. Med Sci (Paris) 28: 1064-1066.

Leggett, D. S., J. Hanna, A. Borodovsky, B. Crosas, M. Schmidt *et al.*, 2002 Multiple associated proteins

regulate proteasome structure and function. Mol Cell 10: 495-507.

Liu, W., X. Dong, M. Mai, R. S. Seelan, K. Taniguchi *et al.*, 2000 Mutations in AXIN2 cause colorectal

cancer with defective mismatch repair by activating beta-catenin/TCF signalling. Nat Genet 26:

146-147.

Mbom, B. C., K. A. Siemers, M. A. Ostrowski, W. J. Nelson and A. I. Barth, 2014 Nek2 phosphorylates and

stabilizes beta-catenin at mitotic centrosomes downstream of Plk1. Mol Biol Cell 25: 977-991.

Mikenberg, I., D. Widera, A. Kaus, B. Kaltschmidt and C. Kaltschmidt, 2007 Transcription factor NF-

kappaB is transported to the nucleus via cytoplasmic dynein/dynactin motor complex in

hippocampal neurons. PLoS One 2: e589.

Mila, D., A. Calderon, A. T. Baldwin, K. M. Moore, M. Watson *et al.*, 2015 Asymmetric Wnt Pathway

Signaling Facilitates Stem Cell-Like Divisions via the Nonreceptor Tyrosine Kinase FRK-1 in

Caenorhabditis elegans. Genetics 201: 1047-1060.

Miskowski, J., Y. Li and J. Kimble, 2001 The sys-1 gene and sexual dimorphism during gonadogenesis in

Caenorhabditis elegans. Dev Biol 230: 61-73.

Niehrs, C., and S. P. Acebron, 2012 Mitotic and mitogenic Wnt signalling. EMBO J 31: 2705-2713.

Niehrs, C., and S. P. Acebron, 2012 Mitotic and mitogenic Wnt signalling. EMBO J 31: 2705-2713.

Nusse, R., and H. Clevers, 2017 Wnt/beta-Catenin Signaling, Disease, and Emerging Therapeutic

Modalities. Cell 169: 985-999.

O'Connell, K. F., C. Caron, K. R. Kopish, D. D. Hurd, K. J. Kemphues *et al.*, 2001 The C. elegans zyg-1 gene

encodes a regulator of centrosome duplication with distinct maternal and paternal roles in the

embryo. Cell 105: 547-558.

O'Rourke, S. M., M. D. Dorfman, J. C. Carter and B. Bowerman, 2007 Dynein modifiers in C. elegans: light

chains suppress conditional heavy chain mutants. PLoS Genet 3: e128.

Peel, N., M. Dougherty, J. Goeres, Y. Liu and K. F. O'Connell, 2012 The C. elegans F-box proteins LIN-23

and SEL-10 antagonize centrosome duplication by regulating ZYG-1 levels. J Cell Sci 125: 3535-

3544.

Peters, N., D. E. Perez, M. H. Song, Y. Liu, T. Muller-Reichert *et al.*, 2010 Control of mitotic and meiotic

centriole duplication by the Plk4-related kinase ZYG-1. J Cell Sci 123: 795-805.

Phillips, B. T., A. R. Kidd, 3rd, R. King, J. Hardin and J. Kimble, 2007 Reciprocal asymmetry of SYS-1/beta-

catenin and POP-1/TCF controls asymmetric divisions in Caenorhabditis elegans. Proc Natl Acad

Sci U S A 104: 3231-3236.

Pintard, L., and B. Bowerman, 2019 Mitotic Cell Division in Caenorhabditis elegans. Genetics 211: 35-73.

Pinto, D., A. Gregorieff, H. Begthel and H. Clevers, 2003 Canonical Wnt signals are essential for

homeostasis of the intestinal epithelium. Genes Dev 17: 1709-1713.

Priyanga, J., G. Guha and D. Bhakta-Guha, 2021 Microtubule motors in centrosome homeostasis: A

target for cancer therapy? Biochim Biophys Acta Rev Cancer 1875: 188524.

Reck-Peterson, S. L., W. B. Redwine, R. D. Vale and A. P. Carter, 2018 The cytoplasmic dynein transport

machinery and its many cargoes. Nat Rev Mol Cell Biol 19: 382-398.

Reilein, A., D. Melamed, K. S. Park, A. Berg, E. Cimetta *et al.*, 2017 Alternative direct stem cell derivatives

defined by stem cell location and graded Wnt signalling. Nat Cell Biol 19: 433-444.

Reya, T., and H. Clevers, 2005 Wnt signalling in stem cells and cancer. Nature 434: 843-850.

Sawa, H., 2012 Control of cell polarity and asymmetric division in C. elegans. Curr Top Dev Biol 101: 55-

76.

Schmidt, R., L. E. Fielmich, I. Grigoriev, E. A. Katrukha, A. Akhmanova *et al.*, 2017 Two populations of

cytoplasmic dynein contribute to spindle positioning in C. elegans embryos. J Cell Biol 216: 2777-

2793.

745 Shtutman, M., J. Zhurinsky, I. Simcha, C. Albanese, M. D'Amico *et al.*, 1999 The cyclin D1 gene is a target

746   of the beta-catenin/LEF-1 pathway. Proc Natl Acad Sci U S A 96: 5522-5527.

747 Siegfried, K. R., A. R. Kidd, M. A. Chesney and J. Kimble, 2004 The sys-1 and sys-3

748   Genes Cooperate With Wnt Signaling to Establish the Proximal-Distal Axis of the

749   Caenorhabditis elegans Gonad. Genetics 166: 171-186.

750 Sonnichsen, B., L. B. Koski, A. Walsh, P. Marschall, B. Neumann *et al.*, 2005 Full-genome RNAi profiling of

751   early embryogenesis in Caenorhabditis elegans. Nature 434: 462-469.

752 Sugioka, K., K. Mizumoto and H. Sawa, 2011 Wnt regulates spindle asymmetry to generate asymmetric

753   nuclear beta-catenin in C. elegans. Cell 146: 942-954.

754 Sunchu, B., and C. Cabernard, 2020 Principles and mechanisms of asymmetric cell division. Development

755   147.

756 Tame, M. A., J. A. Raaijmakers, B. van den Broek, A. Lindqvist, K. Jalink *et al.*, 2014 Astral microtubules

757   control redistribution of dynein at the cell cortex to facilitate spindle positioning. Cell Cycle 13:

758   1162-1170.

759 Tavernarakis, N., S. L. Wang, M. Dorovkov, A. Ryazanov and M. Driscoll, 2000 Heritable and inducible

760   genetic interference by double-stranded RNA encoded by transgenes. Nat Genet 24: 180-183.

761 Tejpar, S., F. Nollet, C. Li, J. S. Wunder, G. Michils *et al.*, 1999 Predominance of beta-catenin mutations

762   and beta-catenin dysregulation in sporadic aggressive fibromatosis (desmoid tumor). Oncogene

763   18: 6615-6620.

764 Tetsu, O., and F. McCormick, 1999 Beta-catenin regulates expression of cyclin D1 in colon carcinoma

765   cells. Nature 398: 422-426.

766 Timmons, L., 2006 Delivery methods for RNA interference in C. elegans. Methods Mol Biol 351: 119-125.

Valenzuela, J. I., M. Jaureguiberry-Bravo, D. A. Salas, O. A. Ramírez, V. H. Cornejo *et al.*, 2014 Transport

along the dendritic endoplasmic reticulum mediates the trafficking of GABAB receptors. J Cell Sci

127: 3382-3395.

Vora, S., and B. T. Phillips, 2015 Centrosome-Associated Degradation Limits beta-Catenin Inheritance by

Daughter Cells after Asymmetric Division. Curr Biol 25: 1005-1016.

Vora, S. M., J. S. Fassler and B. T. Phillips, 2020 Centrosomes are required for proper beta-catenin

processing and Wnt response. Mol Biol Cell 31: 1951-1961.

Vora, S. M., and B. T. Phillips, 2016 The benefits of local depletion: The centrosome as a scaffold for

ubiquitin-proteasome-mediated degradation. Cell Cycle 15: 2124-2134.

Wigley, W. C., R. P. Fabunmi, M. G. Lee, C. R. Marino, S. Muallem *et al.*, 1999 Dynamic association of

proteasomal machinery with the centrosome. J Cell Biol 145: 481-490.

Wójcik, C., and G. N. DeMartino, 2003 Intracellular localization of proteasomes. The International

Journal of Biochemistry & Cell Biology 35: 579-589.

Wozniak, M. J., B. Bola, K. Brownhill, Y. C. Yang, V. Levakova *et al.*, 2009 Role of kinesin-1 and

cytoplasmic dynein in endoplasmic reticulum movement in VERO cells. J Cell Sci 122: 1979-1989.

Wühr, M., E. S. Tan, S. K. Parker, H. W. Detrich, 3rd and T. J. Mitchison, 2010 A model for cleavage plane

determination in early amphibian and fish embryos. Curr Biol 20: 2040-2045.

Yang, L., X. Wu, Y. Wang, K. Zhang, J. Wu *et al.*, 2011 FZD7 has a critical role in cell proliferation in triple

negative breast cancer. Oncogene 30: 4437-4446.

Zhunussova, G., G. Afonin, S. Abdikerim, A. Jumanov, A. Perfilyeva *et al.*, 2019 Mutation Spectrum of

Cancer-Associated Genes in Patients With Early Onset of Colorectal Cancer. Front Oncol 9: 673.

 FIGURE LEGENDS

Figure 1:

Graphical hypothesis and expected results. 1a) Two potential mechanisms by which SYS-1 (Green) is

recruited to centrosomes. We will distinguish approximately the proportion of SYS-1 actively localized by

motor trafficking (indicated by yellow, blue, and purple complex along black microtubules), or passively

via diffusion-capture (indicated by dashed blue lines). 1B-D) Expected outcomes for centrosomal

recruitment of SYS-1 (top), daughter cell nuclear SYS-1 enrichment (middle), and cell fate (bottom) in

wild type (b) organisms with disrupted SYS-1 centrosomal enrichment (c) and organisms which have

disrupted centrosomal negative regulation of SYS-1 (d). Increasing SYS-1 nuclear enrichment in daughter

cells is indicated by darker shade of green color, while darker orange and blue shade indicates the

robustness of corresponding signaled and unsignaled cell fates, respectively

Figure 2:

Rationale and methodology for microtubule motor RNAi Screen. A) GFP::SYS-1 can be occasionally

observed at regions corresponding to DHC-1 mitotic accumulation. This is particularly enhanced in

centrosomally uncoupled *rsa-2(RNAi)* organisms. The frequency of GFP::SYS-1 enrichment at these

locales in lacZ(RNAi) and rsa-2(RNAi) treatment is distinguished by color. A') GFP::SYS-1 at kinetochore

microtubule (white arrowhead) and cortex(black arrowhead) of the dividing *C. elegans* early embryo.

Inset in black shows cortical localization with increased contrast.. B) Demonstration of CEI and TEI

calculation. CEI was calculated in FIJI by subtracting the mean pixel intensity of a cytoplasmic ROI

encompassing the proximal half of the dividing cell, approximated by the light blue square, from a

circular centrosome ROI, approximated by the white square. B') Expanded view of ROI's as defined in B.

B'') Graphical representation of the raw centrosomal and CEI adjusted fluorescence intensity for the

control (Left) and Affected (right) embryos in B. Control (*lacZ(RNAi)* treated) exhibits both a high raw

centrosomal GFP::SYS-1 and CEI. Affected (*rsa-2(RNAi)* treated) exhibits an intermediate raw GFP::SYS-1,

but severely decreased CEI, better reflecting its visual failure to increase centrosomal GFP::SYS-1 notably

above the bulk of cytoplasmic representation. C) Example YFP::SYS-1 CEI (C) and CEI/TEI (C') of controls

and indicated dynein light chain depletions. Width of the violin plot corresponds to a histogram of

observation density. The median is indicated by the central white dot, with the interquartile range

indicated by the thick black line, and 1.5x additional Interquartile range indicated by the thin black line.

N= lacZ(RNAi), 72; *dlc-1(RNAi)*, 74; *dylt-1(RNAi)*, 52; *rsa-2(RNAi)*, 22 * - $p<0.05$, ** - $p<0.01$, *** - $p<0.001$

by Chi-square test (A) or Student's t-test (C-C').

Figure 3:

tsAllele mediated loss of dynein function has a disproportionate effect on SYS-1 localization. . A)

Diagram of the Dynein motor, and hypothetical model of its role in SYS-1 trafficking. Relative location of

dhc-1(ts) lesion indicated. B) Representative centrosome images of YFP::SYS-1 and g-Tubulin::mCherry in

wildtype and ts allele-containing organisms C) embryos at comparable cell cycle phase before (top) and

after (bottom) temperature upshift. D) SYS-1 CEI and D') SYS-1 CEI per TEI in organisms containing *dhc-*

1(ts) , to control for differences in centrosome assembly, standard error depicted. N= 44, 31, 38, 25 in D

and D' columns, left to right. * - $p<0.05$, ** - $p<0.01$, *** - $p<0.001$ by Student's t-test.

Figure 4:

Nocodazole mediated disruption of microtubules enhances centrosomal GFP::SYS-1 enrichment A)

Representative images of GFP::TBB-1 in DMSO and Nocodazole treated embryos. White arrowheads

indicate astral microtubules (left) and the corresponding lack or dysregulated region in nocodazole

treatment (right) B) Quantification of, and C) Representative images of similarly treated GFP::SYS-1

embryos. For B, N=36 DMSO and 46 nocodazole treated embryos. * - $p<0.05$, ** - $p<0.01$, *** - $p<0.001$

by Student's t-test. D) CEI/TEI as evaluated with combinatorial Wildtype (WT) or *dhc-1(ts)* allele for

Dynein Heavy Chain, and treatment with 2% DMSO vehicle media or 2% DMSO + nocodazole media.

Organisms at the permissive and restrictive temperature are differentiated by blue and red coloring,

respectively. Error bars are SEM. E) Representative yfp::SYS-1 and mCh::g-Tubulin for each treatment in

D, respectively.

Figure 5:

Microtubule-mediated processes can also stabilize centrosomal SYS-1. A) Treatment with *ecps-1(RNAi)*

enhances centrosomal SYS-1 accumulation, N=48 *lacZ(RNAi), N =12 rsa-2(RNAi), N= 124 ecps-1(RNAi).* B)

lacZ(RNAi) and *ecps-1(RNAi)* representative centrosomes in grayscale and intensity dependent LUT. C)

Putative dynein and proteasome adaptor ECPS-1 also appears to also enhance centrosomal ZYG-1

accumulation, by GFP::ZYG-1c CEI (N=31 untreated, N=19 *ecps-1(RNAi),* N= 20 *rpt-4(RNAi),* 14 *rsa-*

2(RNAi)) and D) Representative GFP::ZYG-1c centrosomal images in grayscale and intensity dependent

LUT. E) E) Quantified recovery of GFP::SYS-1. Values indicated are the ratio of bleached centrosomes to

their pre-bleach value, normalized such that the post-bleach value, t = 0, is 0 GFP::SYS-1 intensity. Error

bars are SEM. (F-F'') representative FRAP montage of 2[nd] division P1 cells in indicated embryos. Two

pre-bleach, immediately consecutive timepoints are followed by the 1st post-bleach image at time '0'

and images every 7 seconds thereafter indicated top left. * - $p<0.05$, ** - $p<0.01$, *** - $p<0.001$ by

Student's t-test. One-sided Student's T, asterisk indicates p-value from *lacZ(RNAi)* in E-F.

Figure 6:

Effect of Dynein Depletion on nuclear SYS-1 Localization and Gene Expression in EMS Descendant Cells.

A) The EMS precursor (embryo ventral cell) divides, receiving Wnt and Src specification and orientation

cues, respectively (red arrow). These daughters give rise to posterior, signaled E and anterior, unsignaled

MS daughter cells. Wildtype specification results in depicted approximate GFP::SYS-1 localization

pattern specifically enriched in the signaled daughter nucleus. B) Representative images of wildtype and

ecps-1(RNAi) treated embryos. Orange box encloses the unsignaled MS nucleus, Blue box encloses the

signaled E nucleus. C) Quantification of GFP::SYS-1 nuclear enrichment in indicated dynein RNAi

knockdowns. C') Mean difference in nuclear GFP::SYS-1 enrichment from quantification in C. Error bars

are SEM. D) Embryos ectopically expressing pEND-1::GFP::H2B in cells not descended from the E lineage

in the first (orange) second (blue) and third (green) lineage after the founding of the lineage (the 2, 4,

and 8 cell E descendants, respectively). D') Representative embryos exhibiting WT (all lacZ(RNAi)-

treated, left) and ectopic (*ecps-1(RNAi)*- and 2x*dhc-1(RNAi)*- treated for 1, 2, and 3 generations after E

founding, respectively, right) pEND-1::GFP::H2B

Figure 7:

SYS-1 dysregulation via dynein knockdown can cause rare cell fate conversions. A) Diagram of *C. elegans*

distal tip cell signaled lineage specification in the somatic gonadal precursor. B) Representative image of

WT (top) and ectopically induced (bottom) distal tip cells (DTCs). To distinguish DTC's from pLAG-2

driven vulval expression, DTCs are marked with white arrowheads. C) Number and relative frequency of

DTCs via anatomical marker in untreated organisms and in backgrounds with RNAi depletion of various

dynein subunits. D) Number and relative frequency of DTCs marked by anatomical marker when SYS-1 is

overexpressed via a heat shock-driven transgene in both untreated organisms and in backgrounds with

RNAi depletion of various dynein subunits. * = p<0.05, ** = p<0.01, *** = p<0.001 by one-sided Mann-

Whitney test

Figure 8:

SYS-1 dysregulation via dynein knockdown can cause rare cell fate conversions. A) Diagram of *C. elegans*

seam cell (SC) signaled lineage. B) Representative image of WT (top) and ectopically induced (bottom)

SCs. Likely ectopic SC exhibit spatially distinct 'doublets', highlighted here in brackets. C) Number and

relative frequency of SCs via anatomical marker when in untreated organisms and in backgrounds with

RNAi depletion of various dynein subunits. D) Number and relative frequency of SCs via anatomical

marker when SYS-1 is overexpressed via a heat shock-driven transgene in both untreated organisms and

in backgrounds with RNAi depletion of various dynein subunits. * = $p<0.05$, ** = $p<0.01$, *** = $p<0.001$

by Mann-Whitney test

Figure 9:

Model, demonstrating the various processes by which SYS-1 and its regulators appear to be localized to

mitotic centrosomes. Both SYS-1 and some component of its centrosomal proteasome-dependent

negative regulators, here represented by purple circle, demonstrate some dependence on microtubule

mediated trafficking. While various dynein subunits are requisite components of a functional dynein

complex, the negative effect of their depletion on SYS-1 centrosomal steady state indicates a

preferential or more frequent trafficking of SYS-1. Relatively rare proteasome trafficking events

mediated by ECPS-1 meanwhile, are sufficient to maintain appropriate SYS-1 centrosomal enrichment.

This less efficient proteasomal trafficking, however, renders it vulnerable to Nocodazole-mediated

decrease in overall microtubule trafficking/network formation efficiency. Red inset depicts decreased

processivity or cargo binding capacity induced by dynein light chain RNAi depletions. Blue inset depicts

depletion of adapter ECPS-1 and resulting depletion of proteasome-trafficking specific dynein

complexes, while not otherwise affecting the microtubule motor.

A

2% DMSO   25μg/ml Nocodazole

GFP::Tubulin

B

GFP::SYS-1 CEI

**

DMSO   Nocodazole

C

GFP::SYS-1

1% DMSO

+ Nocodazole

D

CEI/TEI

**   **

*   ns

Permissive   Permissive   Permissive   Permissive   Restrictive   Restrictive   Restrictive   Restrictive
DMSO   nocodazole   DMSO   nocodazole   DMSO   nocodazole   DMSO   nocodazole
WT   WT   dhc-1(ts)   dhc-1(ts)   WT   WT   dhc-1(ts)   dhc-1(ts)

E

YFP::SYS-1

mCh::γ-Tubulin

A

C

E

F

G – GFP::SYS-1 recovery in *lacZ(RNAi)*

G' – GFP::SYS-1 recovery in *rpt-4(RNAi)*

G'' – GFP::SYS-1 recovery in *ecps-1(RNAi)*

A

A'

lacZ(RNAi) – 16 seam cells

dylt-1(RNAi) – 18 seam cells

B

# of SCs	15	16	17	18	19	20	21	
P_{hs}-*sys-1 (no hs); lacZ(RNAi)*	1.7%	95%	3.3%					N=60
P_{hs}-*sys-1 (no hs); dlc-1(RNAi)* *		90%	8.3%			1.7%		N= 60
P_{hs}-*sys-1 (no hs); dylt-1(RNAi)* *		88.3%	10%			1.7%		N=60

Percentage of population

■	100%
■	80-99%
■	50-80%
■	30-50%
■	10-30%
■	5-10%
■	0.1-5%
□	0%

C

# of SCs	15	16	17	18	19	20	21	
P_{hs}-*sys-1 (w/hs); lacZ(RNAi)*	1.7%	80%	13.3%	5%				N=60
P_{hs}-*sys-1 (w/hs); dlc-1(RNAi)* **	1.7%	58.3%	33.3%	5%	1.7%			N=60
P_{hs}-*sys-1 (w/hs); dylt-1(RNAi)* ***	1.7%	38.2%	26.7%	11.7%	11.7%	5%	5%	N= 60

Proteasomal Degradation