Learning enhances behaviorally relevant representations in apical dendrites

Sam E. Benezra[1,3,4], Kripa B. Patel[2,3,4], Citlali Pérez Campos[2,3,4], Elizabeth M.C. Hillman[1,2,3,4], and Randy M. Bruno[1,3,4,5,6*]

[1] Department of Neuroscience, Columbia University, New York, NY 10027
[2] Departments of Biomedical Engineering and Radiology, Columbia University, New York, NY 10027
[3] Kavli Institute for Brain Science, Columbia University, New York, NY 10027
[4] Zuckerman Mind Brain Behavior Institute, Columbia University, New York, NY 10027
[5] Department of Physiology, Anatomy & Genetics, University of Oxford, Oxford, OX1 3PT

[6] Lead contact

* Correspondence: randy.bruno@dpag.ox.ac.uk (R.M.B.)

SUMMARY

Learning alters cortical representations and improves perception. Apical tuft dendrites in Layer 1, which are unique in their connectivity and biophysical properties, may be a key site of learning-induced plasticity. We used both two-photon and SCAPE microscopy to longitudinally track tuft-wide calcium spikes in apical dendrites of Layer 5 pyramidal neurons in barrel cortex as mice learned a tactile behavior. Mice were trained to discriminate two orthogonal directions of whisker stimulation. Reinforcement learning, but not repeated stimulus exposure, enhanced tuft selectivity for both directions equally, even though only one was associated with reward. Selective tufts emerged from initially unresponsive or low-selectivity populations. Animal movement and choice did not account for changes in stimulus selectivity. Enhanced selectivity persisted even after rewards were removed and animals ceased performing the task. We conclude that learning produces long-lasting realignment of apical dendrite tuft responses to behaviorally relevant dimensions of a task.

INTRODUCTION

Learning and memory depend on the ability of biological networks to alter their activity based on past experience. For example, as animals learn the behavioral relevance of stimuli in a sensory discrimination task, neural representations of those stimuli are enhanced[1-7], potentially improving the salience of information relayed to downstream areas. Studies in primary somatosensory (S1)[8] and visual cortex[2] have revealed that top-down signals from distant cortical regions can modify sensory representations during learning, although the cellular and circuit mechanisms underlying this plasticity remain unclear.

Cortical layer 1, comprised mainly of apical tuft dendrites of layer 5 (L5) and layer 2/3 pyramidal

neurons, may be a key site driving the enhancement of sensory representations during learning.

Apical tufts are anatomically well positioned for learning, receiving top-down signals from

numerous cortical and thalamic areas[9-11]. While L5 distal tufts are electrically remote and far from

the soma, they are in close proximity to the highly electrogenic calcium spike initiation zone at

the main bifurcation of the apical dendrite, and form a separate biophysical and processing

compartment from the proximal dendrites[12-16]. Top-down signals arriving at the tuft can trigger

tuft-wide dendritic calcium spikes in L5 neurons[17], which can modulate synaptic plasticity across

the entire dendritic tree[18] and potently drive somatic burst firing[15,19-23]. Consistent with this

observation, L5 apical dendrite activity is highly correlated with somatic activity[24,25]. Therefore,

by strongly influencing somatic activity, L5 apical dendritic calcium spikes can play an important

role in modulating cortical output. Several neuromodulators can augment the excitability of the

apical tuft and increase the likelihood of eliciting calcium spikes[26,27], which could be a substrate

for control of plasticity by behavioral state. Consistent with these ideas, we recently

demonstrated that during behavioral training with positive reinforcements, apical tufts in sensory

cortex acquire associations that extend beyond their normal sensory modality[28]. In mouse models

of dementia and Alzheimer's disease[29,30], tuft dendrites exhibit degeneration which may

contribute to the cognitive and memory deficits.

L5 pyramidal neurons are the major source of output from cortex, targeting numerous subcortical

structures that affect behavior. The activity of apical dendrites is known to correlate with stimulus

intensity, and manipulating L5 apical dendrites and their inputs impacts performance of sensory

tasks[17,31-33]. Apical dendritic calcium spikes of pyramidal cells could be a crucial cellular

mechanism in learning-related plasticity and behavioral modification[18,34,35]. However, sensory

representations of apical tufts, as well as possible changes across learning, have received little

attention.

To address this question, we used two-photon microscopy and a new high-speed volumetric

imaging technique called Swept Confocally-Aligned Planar Excitation (SCAPE)[36,37] to

longitudinally track the activity of GCaMP6f-expressing L5 apical tufts in barrel cortex during a

sensory discrimination task. We found that apical tufts underwent extensive dynamic changes in

selectivity for task-relevant stimuli as performance improved, even though only one of the stimuli

was unrewarded. These changes in responses persisted even after animals disengaged from the

task, demonstrating that learning induced long-lasting changes in tuft sensory representations.

Animals that were exposed to the same stimulation protocol without any reinforcement did not

develop enhanced representations. Our results show for the first time that reinforcement learning

expands apical tuft sensory representations along behaviorally relevant dimensions.

RESULTS

Direction discrimination behavior

We devised an awake head-fixed mouse conditioning paradigm that enables controlled

investigation of reinforcement effects across learning (Fig.1A,B). In addition to discriminating

tactile objects, rodents are known to sense wind direction using their whiskers[38,39] and can be

trained to discriminate different directions of whisker deflections[40,41]. With this in mind, we

directed brief (100-ms) air puffs at the whiskers in either of two directions: rostrocaudal

(backward) or ventrodorsal (upward). One of the directions was paired with a water reward

delivered 500 ms after the air puff and thus constituted a conditioned stimulus (CS+). No reward

was given for the other direction (CS-).

Licking and whisking were monitored throughout the session (Fig.1C,D). Stimuli elicited a brief

passive whisker deflection followed by active whisking over the subsequent ~1.5 seconds

(analyzed below, Fig.6). Any anticipatory licks prior to reward delivery were counted as a

response. Typically, on the first session, mice exhibited few anticipatory licks to either stimulus

(Fig.1C, top, grey shading). By session 2 or 3, mice had learned an association between whisker

deflection and reward, but could not discriminate the CS+ and CS- (middle). Within a week (by

sessions 7-9), every mouse we tested learned to reliably lick to the CS+ while withholding licks

to the CS-, performing substantially above chance after a single week of training (Fig.1C, bottom;

Fig.1E,F). Thus, mice rapidly learned to discriminate the direction of whisker stimuli in our

behavioral task.

Overall stimulus-evoked activity is unbiased and stable across conditioning

To investigate the effects of reinforcement learning on apical tuft activity, we imaged apical tufts

(433 x 433 μm field of view) across conditioning days as well as on an unrewarded pre-

conditioning day to measure naïve stimulus responses and an unrewarded post-conditioning day

to detect any long-lasting changes in responses (Fig.1B). Mice remained water-restricted on the

post-conditioning day and continued licking for reward toward the beginning of the session (see

below). We virally delivered the gene for Cre-dependent GCaMP6f[42] in the barrel cortex of

Rbp4-Cre mice, which labels a heterogeneous population of pyramidal neurons comprising

approximately 50% of layer 5[28,43,44]. By targeting our injections to layer 5B, we predominantly

labeled thick-tufted pyramidal neurons (see Methods). Using intrinsic signal imaging, we mapped

the location of the C2, D2, and gamma whisker barrel columns and identified an overlapping

region in layer 1 with sufficient GCaMP6f expression (Fig.2A). The air puff nozzles were aimed

toward the whiskers corresponding to this region. Dendritic activity was longitudinally recorded

from the same field-of-view (horizontal location and depth) in layer 1 across all sessions

(Supplementary Movie 1).

To extract calcium signals from individual cells, we segmented tufts using CaImAn, a sparse non-

negative matrix factorization method that clusters pixels according to their temporal correlation[45]

(see Methods), and analyzed regions of interest exhibiting apical tuft structure (Fig.2B; 65 ± 15

tufts per mouse; mean $\pm$ SD). Individual segmented tufts were substantial in their spatial extent

(>100 µm), reflecting tuft-wide voltage-gated calcium spikes rather than branch-specific N-

methyl-D-aspartate (NMDA) receptor-mediated spikes. All calcium analyses hereafter refer to

tuft-wide calcium spikes. Average responses to an event include failures. In many tufts, the CS+

and CS- reliably evoked an influx of calcium that robustly activated the tuft (examples in Fig.2C).

Successful calcium events across tufts averaged 28% $\Delta F/F$, consistent with previous studies of

layer 5 apical dendrites[17,31]. Interestingly, during intermediate but not early learning, the average

population response to the CS+ exhibited a two-peak structure (Supp Fig.1, session 4) similar to

tuft reward-related signals we observed previously in barrel cortex[28]. By the last-rewarded and

post sessions, the second CS+ peak was no longer visible, which could be an endpoint of mice

learning that the conditioned stimulus predicts the upcoming reward.

Reward can alter somatic receptive fields in the auditory, visual, and somatosensory cortex of

both rodents and non-human primates such that rewarded stimulus representations become more

robust after learning[4,5,28,46], although cortical sensory responses can remain unchanged during

learning[47]. We investigated whether calcium responses to the CS+ increased in the tuft population

as animals learned its association with reward (Fig.2). Average responses of tufts to the CS+ and

CS- were similar during the pre-conditioning session (Fig.2D; $p = 0.20$, signed rank test, n = 440

pre tufts and 418 post tufts), indicating that there was no inherent bias in the population toward a

particular stimulus in naïve animals. Surprisingly, even after learning, responses to the CS+ and

CS- were similar on the last- and post-conditioning sessions ($p = 0.62$, 0.64, respectively, signed

rank test, Fig.2D,E), revealing that no bias develops for the CS+ among dendritic tufts. Only a

minority of tufts exhibited statistically significant (see Methods) average responses to air puff

stimuli (CS+ responsive: $26 \pm 8\%$; CS- responsive: $25 \pm 8\%$; mean $\pm$ SD across all sessions).

When we excluded responses that were not statistically significant, we again found no difference

between the average response amplitudes to the CS+ and CS- on the pre, last-rewarded, and post

sessions ($p = 0.65, 0.31$, and 0.69, respectively, rank sum test; data not shown). Similarly, the

probability of transients in response to CS+ versus CS- (see Methods) did not differ during pre-

conditioning or post-conditioning sessions ($p = 0.66$ and $p = 0.44$, respectively, data not shown).

Therefore, reinforcement learning in our paradigm does not bias tuft representations toward the

rewarded stimulus.

While a bias for the CS+ did not develop after learning, we wondered whether overall tuft

responses to both conditioned stimuli increased as animals learned the task. Linear regression

analysis revealed that conditioning session number was a poor predictor of both CS+ and CS-

amplitudes (All tufts R^2, CS+: 0.0064, CS-: 0.0035, Fig.2E; Significantly responding tufts R^2,

CS+: 0.014, CS-: 0.014, data not shown). We did find a small but significant decrease in

amplitude from pre to last for CS+ ($p < 0.01$) and CS- ($p < 10^{-7}$), but this was not permanent:

amplitudes did not significantly differ between the pre and post sessions (Fig.2D; $p = 0.53, 0.33$,

CS+ and CS- respectively, Wilcoxon rank sum test). Taken together, these findings demonstrate

that reinforcement learning does not robustly bias the magnitudes of tuft calcium responses to

either stimulus at the population level.

Development of tuft selectivity with task learning

While learning produced no bias in overall tuft activity, learning might enhance selectivity for

conditioned stimuli. Barrel cortex neurons are tuned to the angle of whisker deflection[48-50],

indicating that the sets of synaptic connections activated by the CS+ and CS- may be overlapping

but should not be identical. Therefore, the possibility exists that responses to the CS+ and CS-

can change independently of each other. To examine this, we compared the amplitude of the

average response to CS+ and CS- trials for all segmented tufts on the pre, last-rewarded, and post

sessions (Fig.3A; n = 7 mice; 465 pre, 442 last-rewarded, and 430 post tufts). In agreement with

our previous analysis, we found no significant bias in response amplitude toward CS+ or CS-

during any of the three sessions (Fig.3A; Pre: p = 0.20; last-rewarded: p = 0.43; Post: p = 0.64,

sign-rank test). Under naïve conditions during the pre session, most tufts that responded to air

puff stimuli did not strongly prefer the CS+ or CS- (Fig.3A, left). Surprisingly, on the last-

rewarded session and the unrewarded post-conditioning session, we observed a prominent shift in

the response distribution, where many tufts exhibited more selective responses to one stimulus or

the other (Fig.3A, middle and right).

Plasticity can occur after repeated exposure to stimuli even in the absence of reinforcements[51-55].

To test whether enhanced selectivity depended on reinforcement, we imaged a separate group of

similarly water-restricted mice that were repeatedly exposed to the same stimuli for the same

number of days but without any reward. These mice only received water in their home cage

following each imaging session, but never during stimulus presentation. Repeated exposure mice

exhibited a stable distribution of response selectivity over time (Fig.3B; a separate cohort of 7

mice; 317, 313, and 321 tufts on Day 1, Day 8, and Day 9, respectively). These results suggest

that reinforcement learning, and not simply repeated stimulus exposure, drives apical tufts to

become more selective for either the CS+ or CS-.

To directly quantify the response selectivity of tufts, we computed a selectivity index (SI; see

Methods) ranging from -1 (exclusively CS- responsive) to 1 (exclusively CS+ responsive) for

each tuft. Initially in both the conditioned and repeated exposure mice, the SI distribution was

centered around zero, indicating that most tufts in naïve animals did not strongly prefer either

stimulus (Fig.3C,D, left panels). Consistent with our other analyses (Fig.2D), the mean SI

remained close to zero for each of the three sessions (Fig.3C and Supp.Fig.2D; -0.049, -0.001,

and 0.003 for pre-conditioning, last rewarded, and post-conditioning days, respectively; one-way

ANOVA p = 0.37), confirming that learning produced no overall bias toward one particular

stimulus among the population. During learning, the SI distribution of conditioned but not

repeated exposure mice shifted markedly, whereby a much greater proportion of neurons were

highly selective for either the CS+ or CS- (Fig.3C,D, middle and right panels, |SI| pre versus last-

rewarded: $p < 10^{-6}$, |SI| pre versus post: $p < 10^{-5}$; Wilcoxon rank sum test). These effects can even

be observed within individual mice (Supp.Fig.2). Notably, different tufts within the same animal

exhibited opposite changes in selectivity (Supp.Fig.2A,B). Learning significantly increased tuft

selectivity in individual conditioned mice, but not repeated exposure mice (Supp.Fig.2C). The

degree of enhancement in tuft selectivity was closely correlated with conditioned animals' ability

to discriminate stimuli across sessions (Fig.3E; Pearson's $R = 0.60$, $p < 10^{-5}$).

Whereas selectivity magnitude (|SI|) only considers the amplitude of tuft responses to CS+ and

CS-, their discriminability also depends on their variability. For example, a large difference in

CS+ and CS- responses would not be discriminable if the variability of those responses were very

high; a small difference might be discriminable if the variability were low. We therefore

additionally calculated a d-prime metric of neural discriminability that normalizes differences in

response magnitudes to each stimulus by their variability (see Methods). Similar to selectivity

magnitude, we found that neural discriminability was correlated with behavioral performance

(Fig.3F). In conditioned animals, neural discriminability of CS+ and CS- responses of tufts

increased significantly across learning (Fig.3G, blue; first-rewarded versus last-rewarded: $p < 10^{-3}$,

pre versus post: $p < 10^{-4}$; Wilcoxon rank sum test). By contrast, neural discriminability of tuft

responses in the repeated exposure mice decreased slightly with progressive exposure to the

stimuli (Fig.3G, gray; Day 1 versus Final: $p < 0.01$). Finally, we asked whether the ability to

decode stimulus identity on a trial-by-trial basis increased after learning. To test this, we trained a

support vector machine (SVM) to decode stimulus identity from tuft population activity (see

Methods). We found that decoder performance increased significantly when comparing Pre and

First sessions to Post and Last sessions (Supp.Fig3A; sign-rank test, $p = 0.002$), whereas decoder

performance did not improve over time in the repeated exposure mice (Supp.Fig.3B; sign-rank

test, $p = 0.22$). Taken together, these results show that enhanced stimulus representations can

emerge in apical tufts, but require reinforcement.

The above analyses rely on the accurate measurement of calcium spikes from individual tufts.

While two-photon microscopy acquires images with high resolution and speed, the imaging field

is restricted to a single focal plane. This method can only measure calcium signals from a thin

cross-section of the three-dimensionally complex apical structures. Indeed, many of the spatial

components extracted from our two-photon data were comprised of dendritic branches that cross

the imaging plane at different locations (Supp.Fig.4A), which makes it difficult to determine

whether the segmentation software accurately extracted signals from one tuft or erroneously

merged multiple tufts. For the same reasons, a single apical tuft could be falsely classified as two

different tufts. Such errors could mislead our interpretation of selectivity in the population,

especially given that a single apical tuft can exhibit non-homogenous branch-specific

events[15,56,57].

To confirm that our interpretation was not due to segmentation errors, we repeated the

conditioning experiment using a new, high-speed volumetric imaging approach called

SCAPE[36,37], which allowed us to monitor calcium across entire apical tufts (Supplementary

Movie 2). These three-dimensional datasets ($300 \times 1050 \times 234$ μm field of view) encompassed

large portions of the apical tree which included branches converging on their bifurcation points in

layer 2, enabling us to identify whole apical trees unambiguously (Fig.4A,B; Supp.Fig.4B).

CaImAn effectively demixed overlapping trees in these three-dimensional volumes. Using

SCAPE microscopy, we imaged tuft activity of two additional mice conditioned with the same

behavioral paradigm (Fig.4C). Comparison of tuft responses to the CS+ and CS- on the pre, last-

rewarded, and post sessions (Fig.4D; 241 pre, 215 last-rewarded, 150 post tufts in 2 mice)

revealed again that task learning induced significant increases in tuft selectivity (Fig.4E; pre

versus last-rewarded: $p < 10^{-5}$, pre versus post: $p < 10^{-4}$, Wilcoxon rank sum test of |SI|). On

average, the SI magnitudes were similar between tufts imaged using 2-photon microscopy and

SCAPE (mean ± s.e.m. |SI| for 2-photon versus SCAPE; pre: 0.41±0.01 versus 0.40±0.02; last-

rewarded: 0.54±0.02 versus 0.54±0.02; post: 0.51±0.02 versus 0.53±0.03). These data

demonstrate that the effects in our two-photon dataset are not caused by errors in segmentation,

but rather reflect changes at the level of individual dendritic tufts. Our results, based on two

different imaging approaches, clearly demonstrate that reinforcement increases stimulus

selectivity at the level of the entire apical tuft.

Selective tufts emerge from both initially unresponsive and responsive populations

The striking effect of reinforcement learning on tuft response selectivity could develop in several

ways. For example, initially unresponsive tufts could develop a robust response to either stimulus

after learning (e.g., Fig.5A, top). Conceivably, tufts that were initially unselective in naïve

animals could also maintain their response to one stimulus while losing their response to the other

(e.g., Fig.5A, middle). Either or both scenarios could lead to the increase in neurons that are

selective for stimulus direction. To investigate which changes in individual tufts underlie

population-wide improvements in stimulus selectivity, we longitudinally tracked the same set of

tufts across all sessions and compared their selectivity in pre- and post-conditioning sessions for

both conditioned and repeated exposure mice.

First, we categorized tufts that were unresponsive to either stimulus on the first imaging session,

which accounted for the large majority of tufts (Fig.5E; conditioned: 458/603; repeated exposure:

334/457), and compared their response to the CS+ and CS- on the last session to determine if they

became selective (Fig.5B, see Methods). Stimulus-unresponsive tufts, while on average less

active than responsive ones (median calcium events per minute: 2.65 versus 3.66 for stimulus-

unresponsive and responsive tufts, respectively; $p < 10^{-40}$, Wilcoxon rank sum test;

Supplementary Fig.4), were not silent, with many undergoing tuft-wide calcium influx several

times per minute. Silent tufts that are never active during the session may not have been detected

in our imaging, but we were able to detect tufts that discharged as few as 3 voltage-gated calcium

spikes over a 30-minute behavioral session. Interestingly, in both the conditioned and repeated

exposure mice, approximately 40% of initially unresponsive tufts developed a response to at least

one stimulus by the last session, becoming either selective or unselective (Fig.5B). However, in

conditioned animals, the proportion of initially unresponsive tufts that became selective was

significantly larger than in repeated exposure mice (Fig.5B; $p = 0.04$, 2-sample t-test comparing

mice). Furthermore, while the proportion of selective and unselective tufts in this category was

similar for conditioned animals, unselective tufts were more common in repeated exposure mice

(Fig.5B; $p = 0.03$, paired t-test).

Next, we analyzed tufts that were initially responsive and either selective (Fig.5C; conditioned:

56/603, RE: 43/457) or unselective (Fig.5D; conditioned: 89/603, repeated exposure: 80/457). In

these smaller categories, we found no significant differences in the outcome of selectivity

between the two groups of animals. Together, these results indicate that, while both stimulus

exposure and reinforcement can alter tuft tuning, the presence of reward increases the likelihood

that initially unresponsive tufts develop selectivity for either the CS+ or CS- (summarized in

Fig.5E).

While a greater proportion of tufts from the conditioned animals were selective during the final

session (20.2% versus 10.3% of tufts from conditioned and repeated exposure mice, respectively),

we wondered whether conditioning also impacted the degree of selectivity. Note that some tufts

had very small yet statistically different CS+ and CS- response amplitudes and were thus

classified as selective despite a small SI. First, we compared the SI of initially unresponsive tufts

on the final imaging session (Fig. 5F). Supporting our results in Fig. 5B, the SI distribution was

shifted toward the tails in conditioned, but not repeated exposure mice, indicating that reward

enhances selectivity for either the CS+ or CS- in this subset (|SI| conditioned versus repeated

exposure: $p < 10^{-5}$, Wilcoxon rank sum test, n = 199 and 110 tufts, respectively).

Next, we compared the |SI| of all tufts that were categorized as selective during the last imaging

session in conditioned and repeated exposure mice (Fig. 5G). Interestingly, we found that even

among selective tufts, the |SI| distribution in conditioned mice was significantly greater than in

repeated exposure mice (p = 0.006, Wilcoxon rank sum test, n = 122 and 47 tufts, respectively),

indicating that while selective tufts are present after both conditioning and repeated stimulus

exposure, the magnitude of selectivity is stronger after conditioning.

We then quantified the change in |SI| of all tufts that were responsive in both the first and last

sessions by computing the difference between the two sessions (Fig. 5H). Tufts in conditioned

mice exhibited a greater increase in |SI| across sessions compared to repeated exposure mice (p =

0.01, Wilcoxon rank sum test, n = 48 and 42 tufts, respectively), demonstrating that the

magnitude of selectivity in initially responsive tufts increases after reinforcement learning.

Finally, we found that the degree of selectivity of tufts that eventually became unresponsive on

the last session was overall similar between the two groups (Fig.5I, |SI| conditioned versus

repeated exposure: p = 0.06, Wilcoxon rank sum test, n = 97 and 81 tufts, respectively). However,

tufts that became unresponsive were more likely to be initially highly selective in the conditioned

group than in the repeated exposure group (19 tufts with initial |SI| > 0.75 / 97 tufts ending as

unresponsive in the conditioned group versus 3/81 in the repeated exposure group; p = 0.0013, Z

approximation to binomial). Therefore, learning can involve a loss of responsivity in a small

subset of well-tuned tufts.

In summary, our longitudinal analyses revealed that reinforcement learning biases initially

unresponsive tufts toward becoming selective and enhances the selectivity of tufts that are

initially responsive.

Neither movement nor behavioral choice account for enhanced selectivity

Several plausible factors could underlie the changes in selectivity we observed across learning.

For instance, movements like whisking are correlated with layer 5 somatic action potentials[58-60]

and might have impacted calcium activity in the apical tuft. To investigate whether whisking

could account for the changes in tuft selectivity, we imaged the whiskers with a high-speed

camera and computed whisking amplitude (see Methods) while mice underwent conditioning and

two-photon imaging (Fig.6A). First, we considered whether animals changed their whisker

movements in response to conditioned stimuli over the course of learning. We computed the peak

of the mean stimulus-aligned whisking amplitude for the CS+ and CS- (Fig.1C, left; Fig.6B) for

each session in five mice. Although conditioning alters licking behavior (Fig.1C,E), the

magnitudes of whisker movements following both stimuli were stable across sessions (Fig.6B;

CS+: p = 0.44; CS-: p = 0.45; linear regression). We also computed the standard deviation (SD)

of stimulus-evoked whisker amplitude across trials for all sessions (Fig. 6C). While the whisking

amplitude became slightly more reliable (decreased SD) across sessions ($p < 10^{-4}$), the change in

reliability across sessions was similar for CS+ and CS- (p = 0.53). Therefore, whisking is similar

on both trial types throughout learning.

We next examined whether whisking was correlated with tuft calcium activity by comparing

stimulus-triggered averages and intertrial interval (ITI) whisk-triggered averages of all tufts

during post-conditioning. Whisking amplitude was similar between spontaneous ITI whisking

bouts and evoked whisking responses to stimuli (n = 115 and 617 events, respectively; p = 0.53,

Wilcoxon rank sum test). In contrast to air puff stimuli, ITI whisking bouts were not associated

with a robust calcium response (Fig.6D).

To quantify the relationship of whisking and sensory stimuli to tuft calcium spikes, we performed

a linear regression analysis (see Methods) on 322 tufts using calcium influx as the response

variable and either stimulus or whisking amplitude as a single predictor variable (Fig.6E). Air

puff stimuli more reliably predicted calcium influx than whisking amplitude for each of virtually

all tufts ($p < 10^{-12}$, sign rank test). These results are consistent with other studies that found either

only weak or no correlation between whisking and L5 tuft calcium spikes in S1[28,31,32].

Furthermore, we found no relationship between the whisking response and the median SI

magnitude on a given session (Fig.6F, whisking to CS+ p = 0.22, CS- p = 0.78). Therefore,

changes in whisker movement cannot account for the changes in selectivity during learning that

we observed.

Finally, the possibility remains that other task-related signals relaying information about reward

expectation and behavioral choice could impact apical tuft activity and drive increases in

selectivity. To test this, we compared tuft responses to the CS- in false alarm trials (FA; mouse

incorrectly licked for reward) and correct rejection trials (CR; mouse correctly withheld licks) to

determine if their activity was modulated by behavioral choice. Notice that these two trial types

have the same sensory input but involve different choices. (The corresponding analysis for CS+

trials is not technically possible for lack of sufficient Miss trials after the first conditioning day,

an issue also observed in[1]. A future experiment in which the stimulus strengths are substantially

reduced would drastically increase the error rates, enabling a comparison between Hit and Miss

trials.) Tufts were classified as behaviorally modulated if the FA response was significantly

different from the CR response, and were not behaviorally modulated if CR and FA responses

were statistically indistinguishable (e.g. Fig.7A). Behaviorally modulated tufts accounted for only

~10% of the total tuft population in both early and late learning (50/395 in early; 35/406 in late

learning).

To test whether these behaviorally modulated tufts contributed to increased selectivity during

learning, we excluded them and compared selectivity of the remaining behaviorally-insensitive

tufts. We found that selectivity increased significantly from early to late learning (Fig.7B,C;

median |SI| of 345 tufts early versus 371 tufts late learning: 0.38 versus 0.47, p = 0.02, Wilcoxon

rank sum test), similar to our previous analysis of the entire population. Licking, like whisking,

was a relatively poor predictor of tuft calcium influx (Supp.Fig.6A,B). Because some

behaviorally modulated tufts may not have been statistically detectable, we used multivariate

linear regression to disentangle stimulus responses from licking and whisking, which may have

been confounded with choice. Median coefficients for licking and whisking were on average 3.3

times smaller than median stimulus coefficients for the first rewarded, last rewarded, and post

sessions (all $p < 10^{-6}$, Wilcoxon rank sum test). Even after we factored out possible effects of

movements, CS+ and CS- coefficients were enhanced by learning but not repeated exposure

(Supp.Fig.6C,D), consistent with our other analyses. Together, these results demonstrate that

enhanced selectivity during learning cannot be explained by non-sensory signals related to the

animals' behavior.

Enhanced selectivity in barrel cortex is long-lasting when mice exclusively use whiskers

Mice could conceivably exploit other sensory cues to learn and perform the task, such as auditory

cues from the air nozzles or non-whisker tactile cues from air current eddies contacting the fur or

skin. To determine which mice exclusively used their whiskers to distinguish the CS+ and CS-,

we trimmed all whiskers after the post-conditioning session and assessed performance in five

mice (Figure 8). Performance in each of the five mice decreased after whisker trimming,

indicating that each used some whisker information. Three mice performed the task exclusively

with their whiskers, falling to chance levels after the whisker trim ("whiskers only"). Two other

mice still performed the task above chance after the whisker trim, indicating that they were not

exclusively using their whiskers and exploited information from multiple sensory streams

("whiskers + other senses").

We examined whether these two different behavioral strategies impacted tuft selectivity. Both the

"whiskers only" and "whiskers + other senses" groups exhibited enhanced tuft selectivity in the

last-rewarded session relative to pre-conditioning. This effect was more pronounced in the

"whiskers only" mice (Fig.8A,B, left and middle; whiskers only: median |SI| of 180 pre tufts

versus 169 last-rewarded tufts: 0.36 versus 0.59, $p < 10^{-3}$; "whiskers + other senses": median |SI|

of 144 pre tufts versus 155 last-rewarded tufts: 0.39 versus 0.50, p = 0.01). Surprisingly,

enhanced selectivity persisted during the post-conditioning session for the "whiskers only" group

but not the "whiskers + other senses" group (Fig.8A,B right panels; whiskers only: median |SI| of

pre versus 167 tufts post: 0.36 versus 0.58; $p < 10^{-3}$; whiskers + other senses: median |SI| of 155

pre versus post tufts: 0.39 versus 0.42; p = 0.45). Therefore, tuft selectivity in barrel cortex is

enhanced regardless of behavioral strategy, but outlasts conditioning only when mice rely solely

on their whiskers to perform the task.

We further examined this persistence of enhanced tuft selectivity as experienced mice stopped

performing the task. While the entire post-conditioning session was unrewarded, mice initially

expected rewards and licked for many CS+ trials in the first half of the session. By the second

half of the session, the probability of a lick occurring during the CS+ extinguished, approaching

zero (Fig.8C). We compared the selectivity of tufts during the first and second halves of the post-

conditioning sessions of mice that exclusively used their whiskers and found no difference in the

two distributions (Fig.8D, p = 0.94, Wilcoxon rank sum test of |SI|), demonstrating that selectivity

of the population remained stable throughout the session. Taken together, these results

demonstrate that enhanced stimulus selectivity of apical tuft dendrites after reinforcement

learning is long lasting, persisting even after mice cease performing the task and expecting

reward.

DISCUSSION

Our study is the first to investigate how learning a discrimination task alters apical tuft activity.

Using both novel volumetric whole-tuft imaging and conventional planar microscopy, we

discovered that L5 apical tufts acquire enhanced representations of multiple stimuli during

learning. Rather than simply retuning tufts toward the rewarded stimulus, learning enhanced

selectivity for both stimuli, suggesting that tufts are aligning themselves to the behaviorally

relevant stimulus dimensions. These enhanced sensory representations persist even after mice

cease performing the task. In contrast, representations are slightly degraded by mere repeated

exposure to stimuli outside of a task. Consistent with previous studies[28,31], we found that

movement in and of itself has little direct impact on tuft spikes, indicating that increased

selectivity of apicals reflects alterations in sensory coding as animals learn. This sensitization of

tufts to behaviorally relevant sensory dimensions may be a general feature of all sensory cortical

areas.

Tuft spikes enhance plasticity of synaptic inputs that occur over behavioral (seconds-long)

timescales[18,34]. These new behaviorally relevant tuft representations may therefore prime

subsequent plasticity of synapses across the entire pyramidal neuron. Additionally, tuft events

potently modulate somatic burst firing and enhance how somata respond to their basal inputs[15,61].

As learning and plasticity increase apical selectivity for a behaviorally relevant axis, tuft events

will unavoidably amplify somatic burst output along the same axis. This could enable action

potential output of L5 cells in primary sensory cortex to directly drive behavioral responses via

projections to movement related areas, such as the corticostriatal, corticopontine, and

corticotrigeminal pathways. Thus, tuft spikes have the potential to modify somatic output, both in

the present and in the future.

An open question is whether enhanced stimulus representations in apical tufts are required for

learning this task. One way to address this question would be to silence tuft activity during and

after learning by optogenetically activating NDNF-positive interneurons in layer 1[62]. This

approach is not ideal as NDNF interneurons also inhibit other cells such as Layer 2/3 pyramidal

cells, PV interneurons,[63] and possibly the axons of Layer 5 pyramidal cells, which are known to

densely innervate layer 1. Because this manipulation is not specific to layer 5 apicals, the results

would be difficult to interpret. Focal illumination of inhibitory opsins in tufts has also been used

to assess tuft function[64], but balancing tuft against soma silencing remains challenging and

complicates interpretation. Better tools for selective targeting of apicals would be extremely

useful for addressing such issues.

Enhanced Representation of Behaviorally Relevant Stimuli

Enhancing the representation of relevant stimulus dimensions rather than a singularly important

stimulus, such as a rewarded event, has multiple benefits for behavior. In our paradigm, both the

CS+ and CS- are predictive of whether or not a reward will occur in the future. Explicitly

encoding both stimuli could allow sensory cortical areas to directly elicit actions. In the context of

this task, CS+ preferring tufts in barrel cortex may trigger anticipatory licking while CS-

preferring tufts could suppress licking. L5 cells in sensory cortex via their output to striatum,

pons, brain stem, and spinal cord would thereby be able to directly and rapidly drive action

without further cortical processing, such as by frontal areas including motor cortex[32,65]. Such

rapid sensory-motor transformations by primary sensory areas may be critical for natural time-

constrained behavior.

Furthermore, learning produced a representation in which the degree of selectivity for the two

stimuli was continuous and uniformly distributed. Exclusively CS+ or CS- selective apicals never

dominated the population. Continuous degrees of selectivity across the population, rather than

discrete representations, may allow the system to be more robust to the variability caused by

active movements that alter sensory input. A continuous distribution may also facilitate future

adjustments of neural representations as subjects continue to learn a task or encounter new tasks.

The uniformity we observed may reflect that neurons are high-dimensional, being sensitive to

mixtures of variables[60,66-68], only one of which might be altered here by learning. The uniform

distribution of selectivity corresponds to a full range of pessimism to optimism concerning

stimulus predictions of upcoming rewards. Recent work shows that behavioral performance

benefits from reinforcement learning that incorporates the distribution of reward probabilities

rather than just the average expected reward value[69]. L5 corticostriatal synapses could

theoretically afford a plastic substrate for acquiring the necessary distribution of reward

probabilities.

Surprisingly, past studies in which mice were trained to associate one or more stimuli with a

reward typically show that cortical representations are stronger for the rewarded stimulus [1,3,5]. In

contrast to these studies of layer 2/3 somatic activity, our experiments revealed that the overall

tuft calcium response to the CS+ and CS- at the population level did not change significantly after

animals learned the task (Fig.2). Instead, representations for both stimuli were enhanced by

individual tufts developing selectivity for either the CS+ or the CS- (Fig.3). This divergence in

phenomena may result from several important differences between our work and the

aforementioned studies.

First, enhanced selectivity for both rewarded and unrewarded stimuli could be a phenomenon that

is unique to the apical dendritic tufts. In addition to local inputs, the apical tufts of pyramidal cells

in S1 receive long-range top-down input from several sources, including motor cortex[31,70],

secondary somatosensory cortex[11], and secondary thalamus[9,10,71]. Frontal areas, such as prefrontal

cortex, indeed have enhanced representations of the CS+ and CS- after learning[47]. In contrast,

input to the somata is dominated by the local cortical area and primary thalamus[72,73]. While

somato-dendritic coupling can be strong in L5 neurons[25], it is asymmetric; at least 40% of

somatic transients attenuate in a distance-dependent manner along the apical trunk and distal

tufts[24]. The non-overlapping anatomical inputs and asymmetric coupling together could produce

different learning-related effects on apical tuft and somatic stimulus representations.

Second, learning-related changes may manifest differently in layer 2/3, the usual focus of

previous studies[1,3], and layer 5 pyramidal cells, the tufts of which we studied. With the exception

of a small population of corticostriatal cells, most excitatory cells in layer 2/3 project to other

cortical areas to affect further cortical processing[74,75]. In contrast, many L5 cells project to

subcortical structures including the thalamus, superior colliculus, and brainstem, which may

directly trigger behavioral responses[76-78]. In discrimination paradigms, both stimuli are relevant to

behavior. In our task, the CS+ prompted licking to obtain a reward, and the CS- suppressed

licking that would have no benefit. Thus, an enhanced representation of both stimuli in layer 5

would be advantageous for animals to perform the task efficiently. Recently, it was shown that

apical dendrite activation of subcortical-targeting pyramidal tract L5 cells, but not

intratelencephalic L5 cells that are more like L2/3 cells in their connectivity, determines the

detection of tactile stimuli[32]. The Rbp4-Cre mice we used in this study labels a heterogenous

population of layer 5 pyramidal cells, comprising both pyramidal tract and intratelencephalic

neurons. In the future, it would be interesting to examine whether learning has different effects on

the sensory representations of these two populations. Moreover, direct comparisons of the layers

would be particularly informative.

Finally, it is possible that learning-related changes in sensory representations manifest differently

between a somatosensory modality and a visual modality, the latter being the focus of previous

studies. To our knowledge, we are the first to show changes of sensory representations in

somatosensory cortex within a discrimination paradigm. Mice are known to rely more heavily on

their tactile senses than vision[79]. Their heavy reliance on whisker-mediated touch may make it

advantageous to develop sensory representations of a larger variety of relevant tactile stimuli, in

this case, both the CS+ and CS-.

Candidate Plasticity Mechanisms

Enhanced selectivity could be due to changes in local synaptic connectivity, long-range inputs, or

both. Learning may strengthen and weaken synapses onto barrel cortex neurons from ascending

thalamocortical input or from neighboring cells. Such local plasticity could enhance CS+ or CS-

responsiveness. Alternatively or additionally, other cortical regions encoding task context could

via long-range inputs reconfigure barrel cortex to respond more strongly to these stimuli. The

present results do not completely distinguish between these two scenarios because long-range

inputs may still encode the context while the mouse is in the behavioral apparatus. However, we

found that enhanced representations persist after mice are no longer engaged in the task and

receiving rewards. This result suggests that enhanced representations may be a product of local

plasticity in sensory cortex that alters receptive fields.

Even in the absence of reward, repeated exposure to stimuli can drive plasticity in sensory cortex

and alter response tuning. For instance, repeated exposure to oriented gratings can alter the

orientation tuning of cells in primary visual cortex[51-53], and overstimulation of whiskers induces

plasticity at dendritic spines and alters whisker representations in somatosensory cortex[54,80]. Our

results demonstrate that at the population level enhanced representations developed only when

stimuli were behaviorally relevant. Our longitudinal analysis revealed that while the response

dynamics of some tufts changed after repeated stimuli presentations, overall selectivity of the

population did not increase when rewards were omitted (Figs.3&5). This raises the question:

What are the mechanisms that drive enhanced selectivity under rewarded conditions? In one

possible scenario, reward delivery causes the release of neuromodulators that augment the

activity of apical tufts. Cortical layer 1 is innervated by cholinergic afferents from the nucleus

basalis[81] and adrenergic afferents from the locus coeruleus[82], the main source of acetylcholine and

norepinephrine, respectively. Salient events such as reward and arousal lead to the release of

these neuromodulators in cortex[83,84], which could increase the excitability of apical dendrites by

recruiting disinhibitory circuits or directly influencing dendritic currents[26,27,83,85]. In this model,

the release of reward-driven neuromodulators promotes plasticity and an enhanced representation

of temporally aligned sensory inputs. This phenomenon was demonstrated in auditory cortex,

where tones paired with stimulation of the nucleus basalis shifted the tuning of neurons toward

the frequency of the paired stimulus[86].

Why are representations of the CS- equally enhanced when there is no associated reward? One

explanation is that, as mice learn that the CS- indicates absence of reward, the CS- effectively

signals punishment and acquires negative value. Acetylcholine is released in response to aversive

stimuli, and can activate disinhibitory microcircuits that reduce inhibition onto pyramidal cells

and may be essential for learning[87,88]. Thus, it is possible that both the CS+ and CS-

representations are enhanced by neuromodulatory mechanisms tied to reward and punishment,

respectively. An open question is whether the outcome is due to reinforcement learning or the

behavioral state brought on by the reinforcers rather than their valence. Sensory cortical plasticity

may not be tied to reinforcer valence. Our paradigm creates an environment where mice benefit

from being attentive and engaged in order to maximize reward while minimizing effort. Previous

work has shown that active engagement in a visual discrimination task was associated with

significantly higher selectivity in layer 2/3 cells in visual cortex[1]. Task engagement may lead to a

sustained increase in neuromodulator release throughout the conditioning session, priming the

apical dendrites for plasticity and the development of selective responses for task-relevant stimuli

as they learn.

What determines whether a particular tuft eventually becomes selective for the CS+ or CS-? Our

longitudinal analysis revealed that many tufts that were initially unresponsive to either stimulus

developed a highly selective response to either the CS+ or the CS- (Fig.5). In these tufts, stimulus

preference after learning might be seeded by initially weak, directionally selective inputs on to

the neuron that already exist prior to conditioning and that are potentiated by the learning process.

We also found tufts that initially exhibited robust responses to both stimuli and either lost or

significantly reduced their response to one stimulus after learning. The reduction of an apical

response to a particular stimulus could be driven by local disynaptic inhibition between L5

pyramidal cells mediated by the apical-targeting Martinotti cells[89-91]. Through this mechanism, L5

neurons that are selective for a particular stimulus could inhibit responses to that stimulus in

neighboring L5 apical tufts. Experiments that assess the tuning of excitatory and inhibitory inputs

onto apical dendrites as a function of learning could test such mechanisms.

In addition to demonstrating increased tuft selectivity with learning, we replicated a surprising

phenomenon in a previous instrumental behavior in which a population of apical tufts exhibit

activity around the time of reward[28]. This reward-related activity was observed in four out of the

seven conditioned animals only during CS+ trials and was most prominent during intermediate

conditioning sessions, when most animals were still performing at chance levels, and disappeared

completely by the final conditioning session (Supp.Fig.1). Other than this transient effect,

unconditioned stimuli did not appear to elicit calcium responses, consistent with our previous

findings[28]. The disappearance of this reward-related peak might be attributable to the reward

becoming predictable in later stages of learning. In previous classical conditioning experiments,

dopaminergic cells exhibit responses to rewards early in learning due to the novelty of an

unexpected stimulus. These responses are lost after extended training, as animals learn the

association between the CS and reward[92,93]. While dopaminergic terminals are sparse in primary

sensory areas, they are not entirely absent, nor are dopaminergic receptors. Furthermore, the

excitability of the apical tuft is sensitive to noradrenaline[26]. Interestingly, noradrenergic neurons

in the locus coeruleus exhibit a similar phenomenon to dopaminergic neurons, where responses

shift from temporal alignment with the reward to a predictive conditioned stimulus after

learning[94]. Such mechanisms could explain why reward-related activity is restricted to early-to-

intermediate learning in our paradigm.

Global versus local dendritic spikes

Apical dendrites exhibit not only global spikes that elicit calcium influx across the entire tuft,

which we exclusively analyzed here, but also local events known as NMDA spikes, which

typically engage short (<30-μm) segments of individual dendritic branches[15,31,57]. These local,

NMDA receptor-dependent events can promote prolonged plasticity within individual dendritic

branches in the absence of backpropagating actions potentials, a feature that is unique to the

apical dendrites[16]. In motor cortex, branch-specific NMDA spikes are crucial for establishing the

long-lasting plasticity necessary for learning[56], and depolarization provided by multiple local

NMDA spikes is thought to be essential for the generation of a global calcium spike triggered by

distal synaptic inputs[15]. We focused this study on global tuft-wide calcium events, rather than

local events. Local events are more difficult to unambiguously identify in planar imaging[95], and

their existence *in vivo* is still an open question for L5 apicals in barrel cortex[31,57]. Nonetheless,

they may play important roles in plasticity processes that eventually lead to the emergence of

global tuft spike selectivity for stimuli. Volumetric microscopy studies, the feasibility of which

we showed here, are needed to further investigate the existence of local events in such behaviors

as well as examine possible relationships between local and global tuft events during

reinforcement learning. However, it would be essential to verify that seemingly spatially

overlapping local and global events derive from the same dendritic tree, which requires greater

resolution than was practical for the present study.

To analyze activity of individual tufts, we segmented these structures based on spatiotemporal

covariance[45]. This method does not discount the possibility of errors where one tuft is split

erroneously into two trees, or where two highly correlated tufts are merged. With this in mind, we

used volumetric imaging SCAPE microscopy, which allowed us to visualize the apicals in three

dimensions and unambiguously screen for such artifacts. The results from SCAPE are

quantitatively similar to those from two-photon microscopy, and confirm that our observation of

enhanced selectivity with learning is not an artifact of planar imaging.

Stability of learned tuft representations

In contrast to previous studies of discrimination learning[1-3], we included an unrewarded post-

conditioning session to examine whether learning-related effects persisted through extinction.

Our results show that post-conditioning selectivity of the apical population remains significantly

higher than pre-conditioning, even after animals stop licking in response to the CS+ (Fig.8).

Interestingly, the effects of learning are much more pronounced in animals that relied exclusively

on their whiskers to perform the task. In animals that apparently used other sensory modalities,

we observed a modest increase from the pre to last-rewarded session, which seemed to be largely

absent by the post-conditioning session. Considering that these animals were additionally

exploiting other sensory areas to perform, selectivity may have been more widely distributed and

thus diluted in barrel cortex, diminishing the effect and its stability. How long selectivity persists

in the neuronal population after conditioning and which factors influence stability are interesting

open questions for future study.

Conclusion

In summary, we have shown for the first time that reinforcement learning enhances

representations along behaviorally relevant dimensions in apical tufts. Our results suggest that

dendritic calcium spikes are an important cellular mechanism underlying the changes in sensory

encoding that occur with learning, and provide an avenue for further investigation of cellular and

circuit mechanisms underlying plasticity induced by perceptual experience and reinforcement.

This cellular compartment may be key to understanding pathology in some cognitive, memory,

and learning disorders.

ACKNOWLEDGEMENTS

We thank Venkatakaushik Voleti for help with the design, construction, and alignment of the SCAPE microscope; Dan Kato, Georgia Pierce, and Jung Park for help with pilot experiments; Eftychios Pnevmatikakis and Johannes Friedrich for advice on dendrite segmentation; and Larry Abbott, Stefano Fusi, Ashok Litwin-Kumar, Chris Rodgers, Georgia Pierce, Gordon Petty, and Dan Kato for comments on the manuscript. Funding was provided by a Wellcome Trust Discovery Award, an Academy of Medical Sciences Professorship, NIH/NINDS R01 NS069679, and NIH/NINDS R01 NS094659 (RMB); a Kavli Institute for Brain Science Postdoctoral Fellowship (SEB); NIH/NINDS/NIMH/BRAIN U01 NS094296, UF1 NS108213, U19 NS104649, and RF1 MH114276 (EMCH).

AUTHOR CONTRIBUTIONS

SEB and RMB conceived of the behavioral and two-photon imaging experiments. EMCH and RMB conceived of the SCAPE imaging experiments. SEB built the behavioral apparatus, EMCH, KBP, and CC designed, built, and maintained the SCAPE microscope, and RMB built and maintained the two-photon and intrinsic signal microscopes. SEB performed the experiments and analyzed the data with input from RMB and EMCH. SEB and RMB wrote the manuscript.

DATA AVAILABILITY

Due to the large volume of data (~80TB), data are maintained by the authors and available upon request.

MAIN FIGURES

Fig. 1 | Mice rapidly learn to discriminate stimulus direction in head-fixed paradigm. a, A water droplet is paired with air puffs in one direction (CS+) but not the other (CS-). Licking in anticipation of water is assessed in the response window just after CS+ or CS- and prior to water delivery for the CS+ (grey bar). **b,** Experimental timeline. 2-3 weeks after virus injection, naive tuft responses to stimuli are recorded (pre). The CS+ is then paired with water for 8-9 days (blue). On the last day, stimuli are presented without reward (post). In a separate group of mice, the same stimuli are presented over 9 days in the absence of reward (unrewarded group). **c,** Lick rasters for three different sessions in one example mouse. On session 9, the CS+ but not the CS- reliably elicits licks. **d,** Mean baseline-subtracted whisking amplitude aligned to the CS+ (red) and CS- (navy) across sessions 1, 2, and 9 of an example mouse. **e,** Learning curve demonstrates rapid learning. Mean probability of at least one lick in the response window across sessions. **f,** Behavioral performance of each mouse in the rewarded group (M1 – M7).

Fig. 2 | Overall tuft response to stimuli is unbiased and relatively stable across conditioning. a, Dendritic activity was recorded in layer 1 (i) in the C1/C2 barrel columns (ii). (i) Two-photon image ~60 μm deep relative to pia. Dashed yellow lines denote C1 and C2 boundaries from intrinsic imaging. Single cell reconstruction in left panel from[50]. (ii) Tangential section through layer 4 showing barrels stained with streptavidin-Alexa 647 and GCaMP6f-expressing apical trunks. Red circles indicate location of 2-photon lesions to mark the imaging region for post-hoc analysis. **b,** Overlay of five segmented pseudo-colored tufts from imaging field in A(i). **c,** Time courses of calcium responses of example tufts in **b** to three air puffs (dashes). **d,** Amplitude for CS+ (red) and CS- responses (blue), computed for each segmented tuft in the first 1.5 s post-stimulus (grey points), do not differ within or across sessions. Colored lines indicate median. **e,** Same as in **d**, showing data for all conditioning sessions.

Fig. 3 | Reinforcement learning, but not stimulus exposure, enhances tuft selectivity for CS+

and CS- stimuli. a, Across the indicated sessions, individual tufts (circles) exhibit larger biases

to CS+ or CS- (pooled across all conditioned mice). **b,** Repeated exposure to stimuli does not bias

individual tufts to CS+ or CS-. **c,** Conditioning reshapes distribution of selectivity indices for

tufts from Normal on pre-conditioning session to uniform on post-conditioning session. **d,**

Distribution of tuft selectivity indices remains Normal throughout all repeated exposure sessions.

e, Selectivity (median SI magnitude of tufts for each session) increases with behavioral

performance of 6 animals. **f**, Same as e, but with neural discriminability plotted on the y axis. **g,**

Neural discriminability (mean ± sem) of tufts, pooled across all animals on each session,

increases with conditioning and decreases with repeated exposure.

Fig. 4 | High-speed volumetric imaging of apical tufts confirms the emergence of enhanced

selectivity after learning. a, Top and side view of four example tufts segmented from volumetric

SCAPE imaging. **b,** Time courses of calcium activity from example tufts in **a** during five

presentations of air puff stimuli (dashes). **c,** Performance across all conditioning sessions of two

mice that were imaged with SCAPE. **d,** Across the indicated sessions, individual SCAPE-imaged

tufts (circles) exhibit larger biases to CS+ or CS-. **e,** Conditioning reshapes selectivity distribution

from Normal to uniform.

Fig. 5 | Longitudinal tracking reveals that reward enhances the selectivity of both initially

unresponsive and responsive tufts. a, Three example tufts that were longitudinally tracked

across learning. Top row: An initially unresponsive tuft develops a robust response to the CS+

but not the CS- after learning. Middle row: A responsive but unselective tuft loses its robust CS+

response and becomes selective for the CS-. Bottom row: A CS- selective neuron becomes

unresponsive to both stimuli. **b,** Tufts that were unresponsive during the first session were

longitudinally tracked to the last session. Plotted is the mean proportion of selective and

unselective neurons across all animals in the conditioned (black bars) and repeated exposure

(grey bars) groups. **c,d,** Same analysis as **b** for initially selective (**c**) and unselective (**d**) tufts.

Two-sample t-test was used for comparisons between conditioned and repeated exposure groups.

Paired t-test was used for comparisons within a group. * $p < 0.05$. **e,** Total tuft counts from first to

last session within the 3 response categories for either conditioned (left) or repeated exposure

(right) groups. **f,** SI of responsive tufts on the last session that were initially unresponsive during

the first session. Conditioned tufts have enhanced selectivity compared to repeated exposure. **g,**

Tufts that were selective on the last session are more selective if conditioned (black) rather than

undergoing repeated exposure (grey). **h,** Tufts that responded on both pre and post sessions tend

to have higher selectivity if conditioned rather than undergoing repeated exposure. **i,** SI of

responsive tufts on the first session that later became unresponsive during the last session.

Fig. 6 | Whisking is only weakly correlated with tuft activity and cannot account for changes

in selectivity during learning. a, Whisking amplitude aligned to calcium activity of three

example tufts in one session. Green shading indicates periods of whisking. Red and navy ticks

indicate CS+ or CS- delivery, respectively. **b,** Mean whisking response of five mice to CS+ (red)

and CS- (navy) does not change across sessions during learning (mean ± s.e.m.). **c,** Mean

standard deviation of whisking decreases for both CS+ and CS- across learning, but CS+ and CS-

do not differ. **d,** Event-triggered averages of 322 tufts on the post-conditioning day (grey traces -

individual tufts, black inset - population average) are responsive to stimuli but relatively

unmodulated by whisking. **e,** R^2 values for linear models predicting calcium from stimuli (y axis)

are consistently greater than those predicting calcium from whisking (x axis). Each circle

represents a tuft. (n = 322 tufts) **f,** Magnitude of tuft selectivity does not correlate with mean

whisking amplitude during CS+ (left) and CS- trials (right) on that session.

Fig. 7 | Behavioral responses do not account for enhancement of stimulus selectivity during

learning. a, Mean stimulus responses of four tufts during hit (red), CR (cyan), and FA (black)

trials. Top row: Example tufts whose responses are not behaviorally modulated (CR is similar to

FA). Bottom row: Example tufts with behaviorally modulated responses (CR and FA differ). **b,**

Selectivity index (SI) distribution changes from early (left) and late learning sessions (right) even

when tufts with behaviorally modulated responses (CR≠FA) are excluded. **c,** Median SI

magnitude of tufts in each of six animals (from panel **b**) increases from early to late learning

sessions.

Fig. 8 | Apical tufts in barrel cortex of mice performing the task exclusively with their

whiskers undergo long-lasting changes in selectivity. a, SI histograms of mice performing the

task exclusively with their whiskers exhibit increased selectivity across pre-conditioning, last-

rewarded, and post-conditioning sessions. **b,** Relative to pre-conditioning, mice using their

whiskers and other sensory cues to perform the task have increased selectivity during the last

rewarded session, but not the post-conditioning session. **c,** The probability of anticipatory licks in

response to the CS+ extinguishes across post-conditioning blocks (of 20 trials each). **d,** Tuft

selectively remains uniformly distributed during post-conditioning trial blocks 1-2 (top) while

licking is extinguishing, and blocks 3-4 (bottom) in which licking is extinguished.

METHODS

All experiments complied with the NIH Guide for the Care and Use of Laboratory Animals and were approved by the Institutional Animal Care and Use Committee of Columbia University. Sixteen C57BL/6 mice ranging in age from 77 to 316 days old (mean of 123 days at the time of imaging) were used in these experiments. Six were male, and 10 female. Our results were observed in both male and female individuals, and no sex difference was detected.

Surgery

Animals were administered dexamethasone (1 mg/kg) via intramuscular injection 1-4 hours prior to surgery to reduce edema. Anesthesia was induced with 3% isoflurane in oxygen and maintained at 1%. Mice were head-fixed in a stereotax, and a subcutaneous injection of bupivacaine (0.5%, 0.1 mL) was administered under the scalp. Buprenorphine (0.05 mg/kg) was injected subcutaneously on the back. The scalp was cut, and the skull was covered with a thin layer of Vetbond. A circular craniotomy (3-mm diameter) centered at 1.5 mm posterior and 3.5 mm lateral to bregma was made using a dental drill. The dura was kept moist using artificial cerebrospinal fluid.

For both two-photon and SCAPE microscopy, Rbp4-Cre_KL100 mice were injected with 100 nL of virus (initial titer ~$2x10^{13}$ cfu/mL, diluted 1:4 in artificial cerebrospinal fluid) encoding GCaMP6f in a Cre recombinase-specific manner (AAV1-CAG-flex-GCaMP6f, UPenn Vector Core). The virus was injected in layer 5B of the barrel cortex (1.0 mm deep to the pia) using a pulled pipette (20-30 μm ID) fastened on a Nanoject III, which was mounted on a manipulator angled at ~30° from vertical. The virus was delivered via four injections of 100 nL each, spaced at least 400 μm apart. The depth was chosen to maximize labeling of thick-tufted pyramidal

neurons. In pilot experiments, we found that placing injections 1.0 mm deep resulted primarily in

thick-tufted labeling whereas at more superficial depths (e.g., 0.8 mm deep) we obtained mainly

thin-tufted tufts, consistent with ref [96]. The dura was then removed, and a thin cover glass was

implanted and sealed using superglue. A custom metal head plate was implanted on the skull

using dental cement. Twenty-four hours after surgery, carprofen (5 mg/kg) was administered

subcutaneously. Imaging and behavioral training commenced 3 weeks after surgery.

Behavior

Animals in both rewarded 'conditioning' and unrewarded 'repeated exposure' groups were water

restricted for 2 days prior to starting imaging and habituated to head fixation for ~10 minutes on

each of these 2 days. They were subsequently given ~1 mL of water per day for 9 days either by

pairing water rewards with a specific stimulus (conditioning group), or in their cage following the

imaging session (repeated exposure group). Mice were head restrained in a custom-made

behavioral apparatus by positioning the body in a 3D-printed chamber and fastening the head

plate to metal posts flanking the chamber. Air puff stimuli (10 psi measured before a control

solenoid, 100 ms) were delivered from two nozzles (cut P200 pipette tips) positioned toward the

distal tips of the whiskers, in either the rostrocaudal or ventrodorsal direction. Nozzles were

oriented to prevent air jets from stimulating other parts of the face. One of these directions (CS+)

was paired with a water reward (10 μL), delivered through a lick port 0.5 seconds after the

stimulus onset. The particular direction (rostrocaudal vs ventrodorsal) used as the CS+ was

randomized and counterbalanced across mice. Approximately 180 stimuli were presented over the

course of a 30-minute imaging session (8-12-s intertrial interval). The probability of CS+ or CS-

delivery was 50%. In preliminary experiments, we found that an auditory mask helped prevent

mice from exploiting auditory cues to discriminate the two stimuli: a third air nozzle was

positioned close to the mouse and was active throughout the session.

During the first session (pre-conditioning), stimuli were delivered in the absence of reward to

assess neural and behavioral responses in naïve animals. In the following 7-9 days, the CS+ was

paired with reward. Licks for rewards were detected with a capacitance-based touch sensor

(Sparkfun). A trial response was registered when one or more licks were elicited within a 0.5-

second response window following the stimulus and before reward delivery. To determine

whether behavioral performance was above chance, we computed 95% confidence intervals using

the 'binofit' function in MATLAB. During the final session (post-conditioning), stimuli were

delivered in the absence of reward. Animals in the unrewarded group received the same two

stimuli across 9 days without reward pairing. Behavioral experiments were performed with the

Arduino-based OpenMaze open-source behavioral system, whose designs are fully described at

www.openmaze.org. Whisking was monitored at 125 fps with a camera (Sony PS3eye) and

automatically tracked using published software [97].

Intrinsic signal optical imaging and two-photon imaging

Intrinsic signal optical imaging and two-photon imaging were performed on a Sutter movable

objective microscope. The locations of whisker barrels in S1 were identified using intrinsic signal

optical imaging. Single whiskers in isoflurane-anesthetized mice were stimulated at 5 Hz using a

piezoelectric bimorph while recording the reflectance of 700-nm long-pass incandescent light

with a Rolera CCD camera (QImaging) through a low-magnification objective (Zeiss

5X/0.16NA). Movies were collected using software custom-written in Labview (National

Instruments). Regions of reflectance change were referenced to an image acquired under green

illumination.

Two-photon imaging was conducted on the same microscope under the control of the ScanImage

software package (V. Iyer, Janelia Farms). All calcium imaging data was collected by two-photon

microscopy except for those in figure 4. Scanning during awake conditions was performed at 30

fps using a Chameleon Ultra II laser (Coherent) tuned to 920 nm, precompensated for group

velocity dispersion and focused through a 20x/1.0NA water immersion lens (Zeiss). Aquasonic

clear ultrasound gel was used for the immersion medium. Emitted light was collected with an

HQ535/50 filter (Chroma) and GaAsP photomultiplier tubes (Hamamatsu Photonics). Apical tuft

tufts in Layer 1 were imaged at depths of 40-80 μm from the pial surface (1.5x digital zoom in

ScanImage which yielded a 433 x 433 μm field of view, 512 x 512 pixels).

SCAPE imaging

High-speed volumetric imaging was performed using a custom SCAPE microscope as previously

described, including for dendritic tufts[36,37,98]. Briefly, the cortex was illuminated with an oblique

light sheet through a Olympus XLUMPLFLN 20XW 1.0 NA water immersion objective with a 2-

mm working distance. Fluorescence excited by this sheet (extending in the y-z' direction) was

collected by the same objective lens. A galvanometer mirror in the system was positioned to both

cause the oblique light sheet to scan from side to side across the sample (in the x direction) but

also to de-scan returning fluorescence light. This optical path results in an intermediate, de-

scanned oblique image plane that is stationary yet always co-aligned with the plane in the sample

that is being illuminated by the scanning light sheet. Image rotation optics and a fast sCMOS

camera (Andor Zyla 4.2+) were then focused to capture these y-z' images (750 x 200 pixels) at

>1000 frames per second as the sheet was repeatedly scanned across the cortex in the x direction.

All other system parts, including the objective and sample stage, were stationary during high-

speed 3D image acquisition. Data were reformed into a 3D volume by stacking successive y-z'

planes according to the scanning mirror's x position and de-skewing to correct for the oblique

sheet angle. This rotation of the image volume is responsible for its rectangular appearance

despite the camera's square frames. The resulting volumes were large enough to encompass many

GCaMP6f-labeled tufts in barrel cortex,

In this study, the stationary objective lens in SCAPE was configured on a manual rotation mount

and set to 20º-30º away from the standard upright configuration, so the optical axis was

perpendicular to the cranial window to achieve optimal performance without tilting the head of

the animal. A 488-nm laser (Coherent OBIS) was used for excitation (<10 mW at the sample)

with a 500-nm long-pass filter in the emission path. To achieve optimal spatiotemporal resolution

and volume rate, the sample was imaged with an *x*-direction scanning step of 3 μm over a 300 ×

1050 × 234 μm field of view (*x-y-z*, 3.0 × 1.40 × 1.17 μm per voxel, 100 x 750 x 200 voxels) at

10 volumes per second (VPS). Our imaging involves no special practical considerations or

limitations of field of view or resolution, beyond the usual imaging goal of maximizing

FOV while maintaining sufficient resolution to discern structures of interest (dendrites).

Analysis

Two-photon movies were motion corrected using the NormCorre package [99] in MATLAB. Spatial

and temporal components for individual tufts imaged by two-photon and SCAPE were segmented

using CaImAn v1.8.3, which employs large-scale sparse non-negative matrix factorization [45,100].

CaImAn inherently corrects for background signal. All further analyses used custom-written

routines implemented in MATLAB. Spatial components with tuft structural characteristics were

identified and analyzed, while neuropil components were discarded.

To quantify a tuft's response to stimuli, the mean stimulus-aligned ΔF/F was computed across all

CS+ or CS- trials and corrected by the mean ΔF/F of the second before the trial. Probability of

transients was obtained by taking each trial's ΔF/F in the first 1.5 seconds following either the

CS+ or CS- and fitting these data with a univariate mixture of two Normal distributions: $(1-$

$p)N(\mu_1, \sigma_1) + pN(\mu_2, \sigma_2)$. The smaller Normal reflects the distribution of failures, and the larger

Normal the distribution of transient amplitudes following the stimulus. The parameter p captures

the probability of transients.

From these data, a selectivity index (SI) was defined as $(F_{CS+} - F_{CS-}) / (F_{CS+} + F_{CS-})$, in which F_{CS+}

and F_{CS-} are the mean stimulus-aligned amplitudes ($\Delta F/F$) to the CS+ and CS- within the first 1.5

seconds, respectively. This yielded values that range from -1 (exclusively CS- responsive) to 1

(exclusively CS+ responsive). Neural discriminability was defined as d' = $|F_{CS+} - F_{CS-}| / \sqrt{((\sigma^2_{CS+}}$

$+ \sigma^2_{CS-})/2)$ where σ^2_{CS+} is the variance of the response amplitudes in F_{CS+} and σ^2_{CS-} is the variance

of the response amplitudes in F_{CS-}.

For longitudinal analysis, tufts were categorized as stimulus responsive if they met two criteria:

1) Across all trials, the mean $\Delta F/F$ 1.5 seconds before and 1.5 seconds after the stimulus were

significantly different according to the Wilcoxon rank sum test, for either the CS+ or CS-, and 2)

the average response amplitude for that stimulus was greater than 0.04 $\Delta F/F$. Tufts with a

significant response to only one stimulus were categorized as highly selective and their |SI| was

set to 1. To classify tufts as behaviorally modulated, the mean $\Delta F/F$ of the first 1.5 seconds after

the stimulus was computed for false alarm and correct rejection trials and compared with a rank

sum test. Only sessions with at least 12 false alarm trials were used for this analysis. If the two

distributions were significantly different, the tuft was classified as behaviorally modulated.

Custom MATLAB software was used to compute the median whisker angle, and whisking

amplitude was computed as described previously [101]. The median angle was bandpass filtered

from 4 to 30 Hz and passed through a Hilbert transform to calculate phase. We defined the upper

and lower envelopes of the unfiltered median whisking angle as the points in the whisk cycle

where phase equaled 0 (most protracted) or π (most retracted), respectively. Whisking amplitude

was defined as the difference between these two envelopes. Periods of whisking were defined as

times where whisking amplitude exceeded 20% of maximum for at least 250 ms. Periods of time

where amplitude exceeded this threshold for less than 250 ms were considered ambiguous and

excluded from analysis of whisking versus quiescence. The whisking-triggered average for each

tuft was computed by aligning the calcium signal to the start times of whisking periods during

inter-trial intervals (2-8 seconds after stimulus delivery).

For the linear regression analysis, we excerpted the calcium timeseries 2 seconds before and 6

seconds after each stimulus onset. The whisking amplitude signal was frame aligned to the

calcium signal according to the lag of the calcium-whisking cross-correlation peak for each tuft.

Whisking amplitude was then normalized to the max, yielding values that ranged from 0 to 1. The

stimulus predictor variable was a binary vector with an 800-msec 'on' period (24 frames)

centered at the stimulus time. The timing of the stimulus variable was then aligned to the calcium

signal according to the latency of peak of the mean $\Delta F/F$ of the first 1.5 seconds relative to the

stimulus. The lick predictor variable was a binary vector with 'on' periods denoting lick bouts.

Lick bouts were defined as periods of time where the mouse elicited at least 2 licks, with a

maximum gap of 200 ms, and therefore had variable lengths.

For support vector machine (SVM) analysis, the mean $\Delta F/F$ was computed for a pre-stimulus

epoch (1 second immediately preceding the stimulus, used as a negative control) and a post-

stimulus epoch (0.1 – 1.1 seconds after the stimulus) for each trial. Binary SVMs were trained

separately for each epoch using the MATLAB function fitcsvm. For each iteration, 75% of trials

were randomly chosen to train the SVM, and decoder performance was tested on the remaining

25% of trials. Decoder performance for each session was averaged across 10 iterations.

All statistical tests were two-sided. T-tests were used for Normally distributed data. Otherwise

non-parametric tests were applied.

References
960
961   1   Poort, J. *et al.* Learning Enhances Sensory and Multiple Non-sensory Representations in
962   Primary Visual Cortex. *Neuron* **86**, 1478-1490, doi:10.1016/j.neuron.2015.05.037 (2015).
963   2   Liu, D. *et al.* Orbitofrontal control of visual cortex gain promotes visual associative
964   learning. *Nat Commun* **11**, 2784, doi:10.1038/s41467-020-16609-7 (2020).
965   3   Henschke, J. U. *et al.* Reward Association Enhances Stimulus-Specific Representations
966   in Primary Visual Cortex. *Curr Biol* **30**, 1866-1880 e1865,
967   doi:10.1016/j.cub.2020.03.018 (2020).
968   4   Beitel, R. E., Schreiner, C. E., Cheung, S. W., Wang, X. & Merzenich, M. M. Reward-
969   dependent plasticity in the primary auditory cortex of adult monkeys trained to
970   discriminate temporally modulated signals. *Proc Natl Acad Sci U S A* **100**, 11070-11075,
971   doi:10.1073/pnas.1334187100 (2003).
972   5   Goltstein, P. M., Coffey, E. B., Roelfsema, P. R. & Pennartz, C. M. In vivo two-photon
973   Ca2+ imaging reveals selective reward effects on stimulus-specific assemblies in mouse
974   visual cortex. *J Neurosci* **33**, 11540-11555, doi:10.1523/JNEUROSCI.1341-12.2013
975   (2013).
976   6   David, S. V., Fritz, J. B. & Shamma, S. A. Task reward structure shapes rapid receptive
977   field plasticity in auditory cortex. *Proc Natl Acad Sci U S A* **109**, 2144-2149,
978   doi:10.1073/pnas.1117717109 (2012).
979   7   Fritz, J., Shamma, S., Elhilali, M. & Klein, D. Rapid task-related plasticity of
980   spectrotemporal receptive fields in primary auditory cortex. *Nat Neurosci* **6**, 1216-1223,
981   doi:10.1038/nn1141 (2003).
982   8   Banerjee, A. *et al.* Value-guided remapping of sensory cortex by lateral orbitofrontal
983   cortex. *Nature* **585**, 245-250, doi:10.1038/s41586-020-2704-z (2020).
984   9   Zhang, W. & Bruno, R. M. High-order thalamic inputs to primary somatosensory cortex
985   are stronger and longer lasting than cortical inputs. *eLife* **8**, doi:10.7554/eLife.44158
986   (2019).
987   10   Rubio-Garrido, P., Perez-de-Manzo, F., Porrero, C., Galazo, M. J. & Clasca, F. Thalamic
988   input to distal apical dendrites in neocortical layer 1 is massive and highly convergent.
989   *Cereb Cortex* **19**, 2380-2395, doi:10.1093/cercor/bhn259 (2009).
990   11   Cauller, L. J., Clancy, B. & Connors, B. W. Backward cortical projections to primary
991   somatosensory cortex in rats extend long horizontal axons in layer I. *J Comp Neurol* **390**,
992   297-310 (1998).
993   12   Amitai, Y., Friedman, A., Connors, B. W. & Gutnick, M. J. Regenerative activity in
994   apical dendrites of pyramidal cells in neocortex. *Cereb Cortex* **3**, 26-38 (1993).
995   13   Yuste, R., Gutnick, M. J., Saar, D., Delaney, K. R. & Tank, D. W. Ca2+ accumulations in
996   dendrites of neocortical pyramidal neurons: an apical band and evidence for two
997   functional compartments. *Neuron* **13**, 23-43 (1994).
998   14   Schiller, J., Schiller, Y., Stuart, G. & Sakmann, B. Calcium action potentials restricted to
999   distal apical dendrites of rat neocortical pyramidal neurons. *J Physiol* **505 (Pt 3)**, 605-
1000   616 (1997).
1001   15   Larkum, M. E., Nevian, T., Sandler, M., Polsky, A. & Schiller, J. Synaptic integration in
1002   tuft dendrites of layer 5 pyramidal neurons: a new unifying principle. *Science* **325**, 756-
1003   760, doi:10.1126/science.1171958 (2009).
1004   16   Sandler, M., Shulman, Y. & Schiller, J. A Novel Form of Local Plasticity in Tuft
1005   Dendrites of Neocortical Somatosensory Layer 5 Pyramidal Neurons. *Neuron* **90**, 1028-
1006   1042, doi:10.1016/j.neuron.2016.04.032 (2016).
1007   17   Manita, S. *et al.* A Top-Down Cortical Circuit for Accurate Sensory Perception. *Neuron*
1008   **86**, 1304-1316, doi:10.1016/j.neuron.2015.05.006 (2015).

18   Roelfsema, P. R. & Holtmaat, A. Control of synaptic plasticity in deep cortical networks.
Nat Rev Neurosci **19**, 166-180, doi:10.1038/nrn.2018.6 (2018).
19   Larkum, M. E. & Zhu, J. J. Signaling of layer 1 and whisker-evoked Ca2+ and Na+
action potentials in distal and terminal dendrites of rat neocortical pyramidal neurons in
vitro and in vivo. *J Neurosci* **22**, 6991-7005, doi:20026717 (2002).
20   Larkum, M. E., Senn, W. & Luscher, H. R. Top-down dendritic input increases the gain
of layer 5 pyramidal neurons. *Cereb Cortex* **14**, 1059-1070, doi:10.1093/cercor/bhh065
(2004).
21   Schwindt, P. & Crill, W. Mechanisms underlying burst and regular spiking evoked by
dendritic depolarization in layer 5 cortical pyramidal neurons. *J Neurophysiol* **81**, 1341-
1354, doi:10.1152/jn.1999.81.3.1341 (1999).
22   Larkum, M. E., Zhu, J. J. & Sakmann, B. Dendritic mechanisms underlying the coupling
of the dendritic with the axonal action potential initiation zone of adult rat layer 5
pyramidal neurons. *J Physiol* **533**, 447-466 (2001).
23   Manita, S., Miyakawa, H., Kitamura, K. & Murayama, M. Dendritic Spikes in Sensory
Perception. *Front Cell Neurosci* **11**, 29, doi:10.3389/fncel.2017.00029 (2017).
24   Francioni, V., Padamsey, Z. & Rochefort, N. L. High and asymmetric somato-dendritic
coupling of V1 layer 5 neurons independent of visual stimulation and locomotion. *Elife* **8**,
doi:10.7554/eLife.49145 (2019).
25   Beaulieu-Laroche, L., Toloza, E. H. S., Brown, N. J. & Harnett, M. T. Widespread and
Highly Correlated Somato-dendritic Activity in Cortical Layer 5 Neurons. *Neuron* **103**,
235-241 e234, doi:10.1016/j.neuron.2019.05.014 (2019).
26   Labarrera, C. *et al.* Adrenergic Modulation Regulates the Dendritic Excitability of Layer
5 Pyramidal Neurons In Vivo. *Cell Rep* **23**, 1034-1044, doi:10.1016/j.celrep.2018.03.103
(2018).
27   Brombas, A., Fletcher, L. N. & Williams, S. R. Activity-dependent modulation of layer 1
inhibitory neocortical circuits by acetylcholine. *J Neurosci* **34**, 1932-1941,
doi:10.1523/JNEUROSCI.4470-13.2014 (2014).
28   Lacefield, C. O., Pnevmatikakis, E. A., Paninski, L. & Bruno, R. M. Reinforcement
Learning Recruits Somata and Apical Dendrites across Layers of Primary Sensory
Cortex. *Cell Rep* **26**, 2000-2008 e2002, doi:10.1016/j.celrep.2019.01.093 (2019).
29   Luebke, J. I. *et al.* Dendritic vulnerability in neurodegenerative disease: insights from
analyses of cortical pyramidal neurons in transgenic mouse models. *Brain Struct Funct*
214, 181-199, doi:10.1007/s00429-010-0244-2 (2010).
30   Tsai, J., Grutzendler, J., Duff, K. & Gan, W. B. Fibrillar amyloid deposition leads to local
synaptic abnormalities and breakage of neuronal branches. *Nat Neurosci* **7**, 1181-1183,
doi:10.1038/nn1335 (2004).
31   Xu, N. L. *et al.* Nonlinear dendritic integration of sensory and motor input during an
active sensing task. *Nature* **492**, 247-251, doi:10.1038/nature11601 (2012).
32   Takahashi, N. *et al.* Active dendritic currents gate descending cortical outputs in
perception. *Nat Neurosci*, doi:10.1038/s41593-020-0677-8 (2020).
33   Takahashi, N., Oertner, T. G., Hegemann, P. & Larkum, M. E. Active cortical dendrites
modulate perception. *Science* **354**, 1587-1590, doi:10.1126/science.aah6066 (2016).
34   Bittner, K. C., Milstein, A. D., Grienberger, C., Romani, S. & Magee, J. C. Behavioral
time scale synaptic plasticity underlies CA1 place fields. *Science* **357**, 1033-1036,
doi:10.1126/science.aan3846 (2017).
35   Doron, G. *et al.* Perirhinal input to neocortical layer 1 controls learning. *Science* **370**,
doi:10.1126/science.aaz3136 (2020).
36   Bouchard, M. B. *et al.* Swept confocally-aligned planar excitation (SCAPE) microscopy
for high speed volumetric imaging of behaving organisms. *Nat Photonics* **9**, 113-119,
doi:10.1038/nphoton.2014.323 (2015).

1060	37	Hillman, E. M. *et al.* High-speed 3D imaging of cellular activity in the brain using
1061		axially-extended beams and light sheets. *Curr Opin Neurobiol* **50**, 190-200 (2018).
1062	38	Yu, Y. S., Graff, M. M., Bresee, C. S., Man, Y. B. & Hartmann, M. J. Whiskers aid
1063		anemotaxis in rats. *Sci Adv* **2**, e1600716, doi:10.1126/sciadv.1600716 (2016).
1064	39	Yu, Y. S., Graff, M. M. & Hartmann, M. J. Mechanical responses of rat vibrissae to
1065		airflow. *J Exp Biol* **219**, 937-948, doi:10.1242/jeb.126896 (2016).
1066	40	Nakamura, S., Narumi, T., Tsutsui, K. & Iijima, T. Difference in the functional
1067		significance between the lemniscal and paralemniscal pathways in the perception of
1068		direction of single-whisker stimulation examined by muscimol microinjection. *Neurosci*
1069		*Res* **64**, 323-329, doi:10.1016/j.neures.2009.04.005 (2009).
1070	41	Bernhard, S. M. *et al.* An automated homecage system for multiwhisker detection and
1071		discrimination learning in mice. *PLoS One* **15**, e0232916,
1072		doi:10.1371/journal.pone.0232916 (2020).
1073	42	Chen, T. W. *et al.* Ultrasensitive fluorescent proteins for imaging neuronal activity.
1074		*Nature* **499**, 295-300, doi:10.1038/nature12354 (2013).
1075	43	Kozorovitskiy, Y., Saunders, A., Johnson, C. A., Lowell, B. B. & Sabatini, B. L.
1076		Recurrent network activity drives striatal synaptogenesis. *Nature* **485**, 646-650,
1077		doi:10.1038/nature11052 (2012).
1078	44	Glickfeld, L. L., Andermann, M. L., Bonin, V. & Reid, R. C. Cortico-cortical projections
1079		in mouse visual cortex are functionally target specific. *Nat Neurosci* **16**, 219-226,
1080		doi:10.1038/nn.3300 (2013).
1081	45	Giovannucci, A. *et al.* CaImAn an open source tool for scalable calcium imaging data
1082		analysis. *Elife* **8**, doi:10.7554/eLife.38173 (2019).
1083	46	Pakan, J. M. P., Currie, S. P., Fischer, L. & Rochefort, N. L. The Impact of Visual Cues,
1084		Reward, and Motor Feedback on the Representation of Behaviorally Relevant Spatial
1085		Locations in Primary Visual Cortex. *Cell Rep* **24**, 2521-2528,
1086		doi:10.1016/j.celrep.2018.08.010 (2018).
1087	47	Wang, P. Y. *et al.* Transient and Persistent Representations of Odor Value in Prefrontal
1088		Cortex. *Neuron* **108**, 209-224 e206, doi:10.1016/j.neuron.2020.07.033 (2020).
1089	48	Bruno, R. M., Khatri, V., Land, P. W. & Simons, D. J. Thalamocortical angular tuning
1090		domains within individual barrels of rat somatosensory cortex. *J Neurosci* **23**, 9565-9574
1091		(2003).
1092	49	Bruno, R. M. & Sakmann, B. Cortex is driven by weak but synchronously active
1093		thalamocortical synapses. *Science* **312**, 1622-1627, doi:10.1126/science.1124593 (2006).
1094	50	Ramirez, A. *et al.* Spatiotemporal receptive fields of barrel cortex revealed by reverse
1095		correlation of synaptic input. *Nat Neurosci* **17**, 866-875, doi:10.1038/nn.3720 (2014).
1096	51	Yao, H. & Dan, Y. Stimulus timing-dependent plasticity in cortical processing of
1097		orientation. *Neuron* **32**, 315-323, doi:10.1016/s0896-6273(01)00460-3 (2001).
1098	52	Dragoi, V., Sharma, J., Miller, E. K. & Sur, M. Dynamics of neuronal sensitivity in visual
1099		cortex and local feature discrimination. *Nat Neurosci* **5**, 883-891, doi:10.1038/nn900
1100		(2002).
1101	53	Dragoi, V., Sharma, J. & Sur, M. Adaptation-induced plasticity of orientation tuning in
1102		adult visual cortex. *Neuron* **28**, 287-298, doi:10.1016/s0896-6273(00)00103-3 (2000).
1103	54	Zhang, Y., Cudmore, R. H., Lin, D. T., Linden, D. J. & Huganir, R. L. Visualization of
1104		NMDA receptor-dependent AMPA receptor synaptic plasticity in vivo. *Nat Neurosci* **18**,
1105		402-407, doi:10.1038/nn.3936 (2015).
1106	55	Chu, M. W., Li, W. L. & Komiyama, T. Balancing the Robustness and Efficiency of
1107		Odor Representations during Learning. *Neuron* **92**, 174-186,
1108		doi:10.1016/j.neuron.2016.09.004 (2016).
1109	56	Cichon, J. & Gan, W. B. Branch-specific dendritic Ca(2+) spikes cause persistent
1110		synaptic plasticity. *Nature* **520**, 180-185, doi:10.1038/nature14251 (2015).

57   Palmer, L. M. *et al.* NMDA spikes enhance action potential generation during sensory
input. *Nat Neurosci* **17**, 383-390, doi:10.1038/nn.3646 (2014).
58   Derdikman, D. *et al.* Layer-specific touch-dependent facilitation and depression in the
somatosensory cortex during active whisking. *J Neurosci* **26**, 9538-9547,
doi:10.1523/JNEUROSCI.0918-06.2006 (2006).
59   de Kock, C. P. & Sakmann, B. Spiking in primary somatosensory cortex during natural
whisking in awake head-restrained rats is cell-type specific. *Proc Natl Acad Sci U S A*
106, 16446-16450, doi:10.1073/pnas.0904143106 (2009).
60   Rodgers, C. C. *et al.* Sensorimotor strategies and neuronal representations for shape
discrimination. *Neuron* **109**, 2308-2325 e2310, doi:10.1016/j.neuron.2021.05.019 (2021).
61   Larkum, M. A cellular mechanism for cortical associations: an organizing principle for
the cerebral cortex. *Trends Neurosci* **36**, 141-151, doi:10.1016/j.tins.2012.11.006 (2013).
62   Abs, E. *et al.* Learning-Related Plasticity in Dendrite-Targeting Layer 1 Interneurons.
Neuron **100**, 684-699 e686, doi:10.1016/j.neuron.2018.09.001 (2018).
63   Cohen-Kashi Malina, K. *et al.* NDNF interneurons in layer 1 gain-modulate whole
cortical columns according to an animal's behavioral state. *Neuron* **109**, 2150-2164
e2155, doi:10.1016/j.neuron.2021.05.001 (2021).
64   Ranganathan, G. N. *et al.* Active dendritic integration and mixed neocortical network
representations during an adaptive sensing behavior. *Nat Neurosci* **21**, 1583-1590,
doi:10.1038/s41593-018-0254-6 (2018).
65   Park, J. M. *et al.* Deep and superficial layers of the primary somatosensory cortex are
critical for whisker-based texture discrimination in mice. *bioRxiv*,
2020.2008.2012.245381, doi:10.1101/2020.08.12.245381 (2020).
66   Rigotti, M. *et al.* The importance of mixed selectivity in complex cognitive tasks. *Nature*
497, 585-590, doi:10.1038/nature12160 (2013).
67   Stringer, C., Pachitariu, M., Steinmetz, N., Carandini, M. & Harris, K. D. High-
dimensional geometry of population responses in visual cortex. *Nature* **571**, 361-365,
doi:10.1038/s41586-019-1346-5 (2019).
68   Kim, J., Erskine, A., Cheung, J. A. & Hires, S. A. Behavioral and Neural Bases of Tactile
Shape Discrimination Learning in Head-Fixed Mice. *Neuron* **108**, 953-967 e958,
doi:10.1016/j.neuron.2020.09.012 (2020).
69   Dabney, W. *et al.* A distributional code for value in dopamine-based reinforcement
learning. *Nature* **577**, 671-675, doi:10.1038/s41586-019-1924-6 (2020).
70   Petreanu, L. *et al.* Activity in motor-sensory projections reveals distributed coding in
somatosensation. *Nature* **489**, 299-303, doi:10.1038/nature11321 (2012).
71   Wimmer, V. C., Bruno, R. M., de Kock, C. P., Kuner, T. & Sakmann, B. Dimensions of a
Projection Column and Architecture of VPM and POm Axons in Rat Vibrissal Cortex.
Cereb Cortex **20**, 2265-2276, doi:bhq068 [pii] 10.1093/cercor/bhq068 (2010).
72   Feldmeyer, D. Excitatory neuronal connectivity in the barrel cortex. *Front Neuroanat* **6**,
24, doi:10.3389/fnana.2012.00024 (2012).
73   Constantinople, C. M. & Bruno, R. M. Deep cortical layers are activated directly by
thalamus. *Science* **340**, 1591-1594, doi:10.1126/science.1236425 (2013).
74   Petersen, C. C. & Crochet, S. Synaptic computation and sensory processing in neocortical
layer 2/3. *Neuron* **78**, 28-48, doi:10.1016/j.neuron.2013.03.020 (2013).
75   Yamashita, T. *et al.* Diverse Long-Range Axonal Projections of Excitatory Layer 2/3
Neurons in Mouse Barrel Cortex. *Front Neuroanat* **12**, 33, doi:10.3389/fnana.2018.00033
(2018).
76   Llinas, R., Ribary, U., Contreras, D. & Pedroarena, C. The neuronal basis for
consciousness. *Philos Trans R Soc Lond B Biol Sci* **353**, 1841-1849,
doi:10.1098/rstb.1998.0336 (1998).

77 Krauzlis, R. J., Lovejoy, L. P. & Zenon, A. Superior colliculus and visual spatial
attention. *Annu Rev Neurosci* **36**, 165-182, doi:10.1146/annurev-neuro-062012-170249
(2013).
78 Parvizi, J. & Damasio, A. Consciousness and the brainstem. *Cognition* **79**, 135-160,
doi:10.1016/s0010-0277(00)00127-x (2001).
79 Petty, G. H. & Bruno, R. M. Attentional modulation of secondary somatosensory and
visual thalamus of mice. *eLife* **13** (2024).
80 Feldman, D. E. & Brecht, M. Map plasticity in somatosensory cortex. *Science* **310**, 810-
815, doi:10.1126/science.1115807 (2005).
81 Mechawar, N., Cozzari, C. & Descarries, L. Cholinergic innervation in adult rat cerebral
cortex: a quantitative immunocytochemical description. *J Comp Neurol* **428**, 305-318
(2000).
82 Freedman, R., Foote, S. L. & Bloom, F. E. Histochemical characterization of a
neocortical projection of the nucleus locus coeruleus in the squirrel monkey. *J Comp
Neurol* **164**, 209-231, doi:10.1002/cne.901640205 (1975).
83 Chubykin, A. A., Roach, E. B., Bear, M. F. & Shuler, M. G. A cholinergic mechanism for
reward timing within primary visual cortex. *Neuron* **77**, 723-735,
doi:10.1016/j.neuron.2012.12.039 (2013).
84 Thiele, A. & Bellgrove, M. A. Neuromodulation of Attention. *Neuron* **97**, 769-785,
doi:10.1016/j.neuron.2018.01.008 (2018).
85 Hangya, B., Ranade, S. P., Lorenc, M. & Kepecs, A. Central Cholinergic Neurons Are
Rapidly Recruited by Reinforcement Feedback. *Cell* **162**, 1155-1168,
doi:10.1016/j.cell.2015.07.057 (2015).
86 Froemke, R. C., Merzenich, M. M. & Schreiner, C. E. A synaptic memory trace for
cortical receptive field plasticity. *Nature* **450**, 425-429, doi:10.1038/nature06289 (2007).
87 Letzkus, J. J. *et al.* A disinhibitory microcircuit for associative fear learning in the
auditory cortex. *Nature* **480**, 331-335, doi:10.1038/nature10674 (2011).
88 Gasselin, C., Hohl, B., Vernet, A., Crochet, S. & Petersen, C. C. H. Cell-type-specific
nicotinic input disinhibits mouse barrel cortex during active sensing. *Neuron*,
doi:10.1016/j.neuron.2020.12.018 (2021).
89 Berger, T. K., Silberberg, G., Perin, R. & Markram, H. Brief bursts self-inhibit and
correlate the pyramidal network. *PLoS Biol* **8**, doi:10.1371/journal.pbio.1000473 (2010).
90 Kapfer, C., Glickfeld, L. L., Atallah, B. V. & Scanziani, M. Supralinear increase of
recurrent inhibition during sparse activity in the somatosensory cortex. *Nat Neurosci* **10**,
743-753, doi:10.1038/nn1909 (2007).
91 Naka, A. & Adesnik, H. Inhibitory Circuits in Cortical Layer 5. *Front Neural Circuits* **10**,
35, doi:10.3389/fncir.2016.00035 (2016).
92 Ljungberg, T., Apicella, P. & Schultz, W. Responses of monkey dopamine neurons
during learning of behavioral reactions. *J Neurophysiol* **67**, 145-163,
doi:10.1152/jn.1992.67.1.145 (1992).
93 Pan, W. X., Schmidt, R., Wickens, J. R. & Hyland, B. I. Dopamine cells respond to
predicted events during classical conditioning: evidence for eligibility traces in the
reward-learning network. *J Neurosci* **25**, 6235-6242, doi:10.1523/JNEUROSCI.1478-
05.2005 (2005).
94 Bouret, S. & Sara, S. J. Reward expectation, orientation of attention and locus coeruleus-
medial frontal cortex interplay during learning. *Eur J Neurosci* **20**, 791-802,
doi:10.1111/j.1460-9568.2004.03526.x (2004).
95 Sheffield, M. E. & Dombeck, D. A. Calcium transient prevalence across the dendritic
arbour predicts place field properties. *Nature* **517**, 200-204, doi:10.1038/nature13871
(2015).

96   Oberlaender, M. *et al.* Cell type-specific three-dimensional structure of thalamocortical
circuits in a column of rat vibrissal cortex. *Cereb Cortex* **22**, 2375-2391,
doi:10.1093/cercor/bhr317 (2012).
97   Clack, N. G. *et al.* Automated tracking of whiskers in videos of head fixed rodents. *PLoS
Comput Biol* **8**, e1002591, doi:10.1371/journal.pcbi.1002591 (2012).
98   Voleti, V. *et al.* Real-time volumetric microscopy of in vivo dynamics and large-scale
samples with SCAPE 2.0. *Nat Methods* **16**, 1054-1062, doi:10.1038/s41592-019-0579-4
(2019).
99   Pnevmatikakis, E. A. & Giovannucci, A. NoRMCorre: An online algorithm for piecewise
rigid motion correction of calcium imaging data. *J Neurosci Methods* **291**, 83-94,
doi:10.1016/j.jneumeth.2017.07.031 (2017).
100   Pnevmatikakis, E. A. *et al.* Simultaneous Denoising, Deconvolution, and Demixing of
Calcium Imaging Data. *Neuron* **89**, 285-299, doi:10.1016/j.neuron.2015.11.037 (2016).
101   Petty, G. H., Kinnischtzke, A. K., Hong, Y. K. & Bruno, R. M. Effects of arousal and
movement on secondary somatosensory and visual thalamus. *bioRxiv*,
2020.2003.2004.977348, doi:10.1101/2020.03.04.977348 (2020).
1227
1228

Fig. 1 l Mice rapidly learn to discriminate stimulus direction in head-fixed paradigm.
a, A water droplet is paired with air puffs in one direction (CS+) but not the other (CS-).
Licking in anticipation of water is assessed in the response window just after CS+ or CS-
and prior to water delivery for the CS+ (grey bar). **b,** Experimental timeline. 2-3 weeks after
virus injection, naive tuft responses to stimuli are recorded (pre). The CS+ is then paired
with water for 8-9 days (blue). On the last day, stimuli are presented without reward (post).
In a separate group of mice, the same stimuli are presented over 9 days in the absence of
reward (unrewarded group). **c,** Lick rasters for three different sessions in one example
mouse. On session 9, the CS+ but not the CS- reliably elicits licks. **d,** Mean baseline-sub-
tracted whisking amplitude aligned to the CS+ (red) and CS- (navy) across sessions 1, 2,
and 9 of an example mouse. **e,** Learning curve demonstrates rapid learning. Mean proba-
bility of at least one lick in the response window across sessions. f, Behavioral perfor-
mance of each mouse in the rewarded group (M1 – M7).

Fig. 2 | Overall tuft response to stimuli is unbiased and relatively stable across conditioning. a, Dendritic activity was recorded in layer 1 (i) in the C1/C2 barrel columns (ii). (i) Two-photon image ~60 μm deep relative to pia. Dashed yellow lines denote C1 and C2 boundaries from intrinsic imaging. (ii) Tangential section through layer 4 showing barrels stained with streptavidin-Alexa 647 and GCaMP6f-expressing apical trunks. Red circles indicate location of 2-photon lesions to mark the imaging region for post-hoc analysis. b, Overlay of five segmented pseudo-colored tufts from imaging field in A(i). c, Time courses of calcium responses of example tufts in b to three air puffs (dashes). d, Amplitude for CS+ (red) and CS-responses (blue), computed for each segmented tuft in the first 1.5 s post-stimulus (grey points), do not differ within or across sessions. Colored lines indicate median. e, Same as in d, showing data for all conditioning sessions.

Fig. 3 | Reinforcement learning, but not stimulus exposure, enhances tuft selectivity for CS+ and CS- stimuli. a, Across the indicated sessions, individual tufts (circles) exhibit larger biases to CS+ or CS- (pooled across all conditioned mice). b, Repeated exposure to stimuli does not bias individual tufts to CS+ or CS-. c, Conditioning reshapes distribution of selectivity indices for tufts from Normal on pre-conditioning session to uniform on post-conditioning session. d, Distribution of tuft selectivity indices remains Normal throughout all repeated exposure sessions. e, Selectivity (median SI magnitude of tufts for each session) increases with behavioral performance of 6 animals. f, Same as e, but with neural discriminability plotted on the y axis. g, Neural discriminability (mean ± sem) of tufts, pooled across all animals on each session, increases with conditioning and decreases with repeated exposure.

Fig. 4 | High-speed volumetric imaging of apical tufts confirms the emergence of enhanced selectivity after learning. a, Top and side view of four example tufts segmented from volumetric SCAPE imaging. **b,** Time courses of calcium activity from example tufts in **a** during five presentations of air puff stimuli (dashes). **c,** Performance across all conditioning sessions of two mice that were imaged with SCAPE. **d,** Across the indicated sessions, individual SCAPE-imaged tufts (circles) exhibit larger biases to CS+ or CS-. **e,** Conditioning reshapes selectivity distribution from Normal to uniform.

Fig. 5 | Longitudinal tracking reveals that reward enhances the selectivity of both initially unresponsive and responsive tufts. a, Three example tufts that were longitudinally tracked across learning. Top row: An initially unresponsive tuft develops a robust response to the CS+ but not the CS- after learning. Middle row: A responsive but unselective tuft loses its robust CS+ response and becomes selective for the CS-. Bottom row: A CS- selective neuron becomes unresponsive to both stimuli. **b,** Tufts that were unresponsive during the first session were longitudinally tracked to the last session. Plotted is the mean proportion of selective and unselective neurons across all animals in the conditioned (black bars) and repeated exposure (grey bars) groups. **c,d,** Same analysis as **b** for initially selective (**c**) and unselective (**d**) tufts. Two-sample t-test was used for comparisons between conditioned and repeated exposure groups. Paired t-test was used for comparisons within a group. * $p < 0.05$. **e,** Total tuft counts from first to last session within the 3 response categories for either conditioned (left) or repeated exposure (right) groups. **f,** SI of responsive tufts on the last session that were initially unresponsive during the first session. Conditioned tufts have enhanced selectivity compared to repeated exposure. **g,** Tufts that were selective on the last session are more selective if conditioned (black) rather than undergoing repeated exposure (grey). **h,** Tufts that responded on both pre and post sessions tend to have higher selectivity if conditioned rather than undergoing repeated exposure. **i,** SI of responsive tufts on the first session that later became unresponsive during the last session.

Fig. 6 | Whisking is only weakly correlated with tuft activity and cannot account for changes in selectivity during learning. a, Whisking amplitude aligned to calcium activity of three example tufts in one session. Green shading indicates periods of whisking. Red and navy ticks indicate CS+ or CS- delivery, respectively. **b,** Mean whisking response of four mice to CS+ (red) and CS- (navy) does not change across sessions during learning (mean ± s.e.m.). **c,** Mean standard deviation of whisking decreases for both CS+ and CS- across learning, but CS+ and CS- do not differ. **d,** Event-triggered averages of 322 tufts on the post-conditioning day (grey traces - individual tufts, black inset - population average) are responsive to stimuli but relatively unmodulated by whisking. **e,** R2 values for linear models predicting calcium from stimuli (y axis) are consistently greater than those predicting calcium from whisking (x axis). Each circle represents a tuft. **f,** Magnitude of tuft selectivity does not correlate with mean whisking amplitude during CS+ (left) and CS- trials (right) on that session.

Fig. 7 | Behavioral responses do not account for enhancement of stimulus selectivity during learning. a, Mean stimulus responses of four tufts during hit (red), CR (cyan), and FA (black) trials. Top row: Example tufts whose responses are not behaviorally modulated (CR is similar to FA). Bottom row: Example tufts with behaviorally modulated responses (CR and FA differ). **b,** Selectivity index (SI) distribution changes from early (left) and late learning sessions (right) even when tufts with behaviorally modulated responses (CR≠FA) are excluded. **c,** Median SI magnitude of tufts in each of six animals (from panel **b**) increases from early to late learning sessions.

Fig. 8 | Apical tufts in barrel cortex of mice performing the task exclusively with their whiskers undergo long-lasting changes in selectivity. a, SI histograms of mice performing the task exclusively with their whiskers exhibit increased selectivity across pre-conditioning, last-rewarded, and post-conditioning sessions. **b,** Relative to pre-conditioning, mice using their whiskers and other sensory cues to perform the task have increased selectivity during the last rewarded session, but not the post-conditioning session. **c,** The probability of anticipatory licks in response to the CS+ extinguishes across post-conditioning blocks (of 20 trials each). **d,** Tuft selectively remains uniformly distributed during post-conditioning trial blocks 1-2 (top) while licking is extinguishing, and blocks 3-4 (bottom) in which licking is extinguished.