Learning enhances behaviorally relevant representations in apical dendrites

Sam E. Benezra[1,3,4], Kripa B. Patel[2,3,4], Citlali Pérez Campos[2,3,4], Elizabeth M.C. Hillman[1,2,3,4], and Randy M. Bruno[1,3,4,5*]

[1] Department of Neuroscience, Columbia University, New York, NY 10027
[2] Departments of Biomedical Engineering and Radiology, Columbia University, New York, NY 10027
[3] Kavli Institute for Brain Science, Columbia University, New York, NY 10027
[4] Zuckerman Mind Brain Behavior Institute, Columbia University, New York, NY 10027

[5] Lead contact

* Correspondence: randybruno@columbia.edu (R.M.B.)

SUMMARY

Learning alters cortical representations and improves perception. Apical tuft dendrites in Layer 1, which are unique in their connectivity and biophysical properties, may be a key site of learning-induced plasticity. We used both two-photon and SCAPE microscopy to longitudinally track tuft-wide calcium spikes in apical dendrites of Layer 5 pyramidal neurons as mice learned a tactile behavior. Mice were trained to discriminate two orthogonal directions of whisker stimulation. Reinforcement learning, but not repeated stimulus exposure, enhanced tuft selectivity for both directions equally, even though only one was associated with reward. Selective tufts emerged from initially unresponsive or low-selectivity populations. Animal movement and choice did not account for changes in stimulus selectivity. Enhanced selectivity persisted even after rewards were removed and animals ceased performing the task. We conclude that learning produces long-lasting realignment of apical dendrite tuft responses to behaviorally relevant dimensions of a task.

INTRODUCTION

Learning and memory depend on the ability of biological networks to alter their activity based on past experience. For example, as animals learn the behavioral relevance of stimuli in a sensory discrimination task, neural representations of those stimuli are enhanced (Beitel et al., 2003; David et al., 2012; Fritz et al., 2003; Goltstein et al., 2013; Henschke et al., 2020; Liu et al., 2020; Poort et al., 2015), potentially improving the salience of information relayed to downstream areas. Studies in primary somatosensory (S1) (Banerjee et al., 2020) and visual cortex (Liu et al., 2020) have revealed that top-down signals from distant cortical regions can modify sensory representations during learning, although the cellular and circuit mechanisms underlying this plasticity remain unclear.

Cortical layer 1, comprised mainly of apical tuft dendrites of layer 5 (L5) and layer 2/3 pyramidal neurons, may be a key site driving the enhancement of sensory representations during learning. Apical tufts are anatomically well positioned for learning, receiving top-down signals from numerous cortical and thalamic areas (Cauller et al., 1998; Rubio-Garrido et al., 2009; Zhang and Bruno, 2019). While L5 distal tufts are electrically remote and far from the soma, they are in close proximity to the highly electrogenic calcium spike initiation zone at the main bifurcation of the apical dendrite, and form a separate biophysical and processing compartment from the proximal dendrites (Amitai et al., 1993; Larkum et al., 2009; Sandler et al., 2016; Schiller et al., 1997; Yuste et al., 1994). Top-down signals arriving at the tuft can trigger tuft-wide dendritic calcium spikes in L5 neurons (Manita et al., 2015), which can modulate synaptic plasticity across the entire dendritic tree (Roelfsema and Holtmaat, 2018) and potently drive somatic burst firing (Larkum et al., 2009; Larkum et al., 2004; Larkum and Zhu, 2002; Larkum et al., 2001; Manita et al., 2017; Schwindt and Crill, 1999). Consistent with this observation, L5 apical dendrite activity is highly correlated with somatic activity (Beaulieu-Laroche et al., 2019; Francioni et al., 2019). Therefore, by strongly influencing somatic activity, L5 apical dendritic calcium spikes can play an important role in modulating cortical output. Several neuromodulators can augment the excitability of the apical tuft and increase the likelihood of eliciting calcium spikes (Brombas et al., 2014; Labarrera et al., 2018), which could be a substrate for control of plasticity by behavioral state. Consistent with these ideas, we recently demonstrated that during behavioral training with positive reinforcements, apical tufts in sensory cortex acquire associations that extend beyond their normal sensory modality (Lacefield et al., 2019). In mouse models of dementia and Alzheimer's disease (Luebke et al., 2010; Tsai et al., 2004), tuft dendrites exhibit degeneration which may contribute to the cognitive and memory deficits.

L5 pyramidal neurons are the major source of output from cortex, targeting numerous subcortical
structures that affect behavior. The activity of apical dendrites is known to correlate with stimulus
intensity, and manipulating L5 apical dendrites and their inputs impacts performance of sensory
tasks (Manita et al., 2015; Takahashi et al., 2020; Takahashi et al., 2016; Xu et al., 2012). Apical
dendritic calcium spikes of pyramidal cells could be a crucial cellular mechanism in learning-
related plasticity and behavioral modification (Bittner et al., 2017; Roelfsema and Holtmaat, 2018).
However, sensory representations of apical tufts, as well as possible changes across learning, have
received little attention.
59
To address this question, we used two-photon microscopy and a new high-speed volumetric
imaging technique called Swept Confocally-Aligned Planar Excitation (SCAPE) (Bouchard et al.,
2015; Hillman et al., 2018) to longitudinally track the activity of GCaMP6f-expressing L5 apical
tufts in barrel cortex during a sensory discrimination task. We found that apical tufts underwent
extensive dynamic changes in selectivity for task-relevant stimuli as performance improved, even
though only one of the stimuli was unrewarded. These changes in responses persisted even after
animals disengaged from the task, demonstrating that learning induced long-lasting changes in tuft
sensory representations. Animals that were exposed to the same stimulation protocol without any
reinforcement did not develop enhanced representations. Our results show for the first time that
reinforcement learning expands apical tuft sensory representations along behaviorally relevant
dimensions.
71
72 **RESULTS**
73
Direction discrimination behavior
We devised an awake head-fixed mouse conditioning paradigm that enables controlled
investigation of reinforcement effects across learning (Fig.1A,B). In addition to discriminating
tactile objects, rodents are known to sense wind direction using their whiskers (Yu et al., 2016a;
Yu et al., 2016b) and can be trained to discriminate different directions of whisker deflections
(Bernhard et al., 2020; Nakamura et al., 2009). With this in mind, we directed brief (100-ms) air
puffs at the whiskers in either of two directions: rostrocaudal (backward) or ventrodorsal (upward).
One of the directions was paired with a water reward delivered 500 ms after the air puff and thus
constituted a conditioned stimulus (CS+). No reward was given for the other direction (CS-).
83
Licking and whisking were monitored throughout the session (Fig.1C,D). Stimuli elicited a brief
passive whisker deflection followed by active whisking over the subsequent ~1.5 seconds (analyzed
below, Fig.6). Any anticipatory licks prior to reward delivery were counted as a response.
Typically, on the first session, mice exhibited few anticipatory licks to either stimulus (Fig.1C, top,
grey shading). By session 2 or 3, mice had learned an association between whisker deflection and
reward, but could not discriminate the CS+ and CS- (middle). Within a week (by sessions 7-9),
every mouse we tested learned to reliably lick to the CS+ while withholding licks to the CS-,
performing substantially above chance after a single week of training (Fig.1C, bottom; Fig.1E,F).
Thus, mice rapidly learned to discriminate the direction of whisker stimuli in our behavioral task.
93
Overall stimulus-evoked activity is unbiased and stable across conditioning
To investigate the effects of reinforcement learning on apical tuft activity, we imaged apical tufts
(433 x 433 μm field of view) across conditioning days as well as on an unrewarded pre-conditioning
day to measure naïve stimulus responses and an unrewarded post-conditioning day to detect any
long-lasting changes in responses (Fig.1B). We virally delivered the gene for Cre-dependent
GCaMP6f (Chen et al., 2013) in the barrel cortex of Rbp4-Cre mice, which labels a heterogeneous
population of pyramidal neurons comprising approximately 50% of layer 5 (Glickfeld et al., 2013;
Kozorovitskiy et al., 2012; Lacefield et al., 2019). By targeting our injections to layer 5B, we

predominantly labeled thick-tufted pyramidal neurons (see Methods). Using intrinsic signal
imaging, we mapped the location of the C2, D2, and gamma whisker barrel columns and identified
an overlapping region in layer 1 with sufficient GCaMP6f expression (Fig.2A). The air puff nozzles
were aimed toward the whiskers corresponding to this region. Dendritic activity was longitudinally
recorded from the same field-of-view (horizontal location and depth) in L1 across all sessions
(Supplementary Movie 1).
108
To extract calcium signals from individual cells, we segmented tufts using CaImAn, a sparse non-
negative matrix factorization method that clusters pixels according to their temporal correlation
(Giovannucci et al., 2019) (see Methods), and analyzed regions of interest exhibiting apical tuft
structure (Fig.2B; 65 ± 15 tufts per mouse; mean $\pm$ SD). Individual segmented tufts were substantial
in their spatial extent (>100 μm), reflecting tuft-wide voltage-gated calcium spikes rather than
branch-specific N-methyl-D-aspartate (NMDA) receptor-mediated spikes. All calcium analyses
hereafter refer to tuft-wide calcium spikes. Average responses to an event include failures. In many
tufts, the CS+ and CS- reliably evoked an influx of calcium that robustly activated the tuft (e.g.
Fig.2C). Interestingly, during intermediate but not early learning, the average population response
to the CS+ exhibited a two-peak structure (Supp Fig.1, session 4) similar to tuft reward-related
signals we observed previously in barrel cortex (Lacefield et al., 2019). By the last-rewarded and
post sessions, the second CS+ peak was no longer visible, which could be an endpoint of mice
learning that the conditioned stimulus predicts the upcoming reward.
122
Reward can alter somatic receptive fields in the auditory, visual, and somatosensory cortex of both
rodents and non-human primates such that rewarded stimulus representations become more robust
after learning (Beitel et al., 2003; Goltstein et al., 2013; Lacefield et al., 2019; Pakan et al., 2018),
although cortical sensory responses can remain unchanged during learning (Wang et al., 2020). We
investigated whether calcium responses to the CS+ increased in the tuft population as animals
learned its association with reward. Average responses of tufts to the CS+ and CS- were similar
during the pre-conditioning session (Fig.2D; $p = 0.20$, signed rank test), indicating that there was
no inherent bias in the population toward a particular stimulus in naïve animals. Surprisingly, even
after learning, responses to the CS+ and CS- were similar on the last- and post-conditioning
sessions ($p = 0.62$, 0.64, respectively, signed rank test), revealing that no bias develops for the CS+
among dendritic tufts. Only a minority of tufts exhibited statistically significant average responses
to air puff stimuli (CS+ responsive: $26 \pm 8\%$; CS- responsive: $25 \pm 8\%$; mean $\pm$ SD across all
sessions). When we excluded responses that were not statistically significant (see Methods), we
again found no difference between the average response amplitudes to the CS+ and CS- on the pre,
last-rewarded, and post sessions ($p = 0.65$, 0.31, and 0.69, respectively, rank sum test; data not
shown). Similarly, the probability of transients in response to CS+ versus CS- (see Methods) did
not differ during pre-conditioning ($p = 0.66$) or post-conditioning sessions ($p = 0.44$). Therefore,
reinforcement learning in our paradigm does not bias tuft representations toward the rewarded
stimulus.
142
While a bias for the CS+ did not develop after learning, we wondered whether overall tuft responses
to both conditioned stimuli increased as animals learned the task. Linear regression analysis
revealed that conditioning session number was a poor predictor of both CS+ and CS- amplitudes
(All tufts R^2, CS+: 0.0064, CS-: 0.0035, Fig.2E; Significantly responding tufts R^2, CS+: 0.014, CS-
: 0.014, data not shown). We did find a small but significant decrease in amplitude from pre to last
for CS+ ($p < 0.01$) and CS- ($p < 10^{-7}$), but this was not permanent: amplitudes did not significantly
differ between the pre and post sessions (Fig.2D; $p = 0.53$, 0.33, CS+ and CS- respectively,
Wilcoxon rank sum test). Taken together, these findings demonstrate that reinforcement learning
does not robustly bias the magnitudes of tuft calcium responses to either stimulus at the population
level.

Development of tuft selectivity with task learning

Development of tuft selectivity with task learning
While learning produced no bias in overall tuft activity, learning might enhance selectivity for
conditioned stimuli. Barrel cortex neurons are tuned to the angle of whisker deflection (Bruno et
al., 2003; Bruno and Sakmann, 2006; Ramirez et al., 2014), indicating that the sets of synaptic
connections activated by the CS+ and CS- may be overlapping but should not be identical.
Therefore, the possibility exists that responses to the CS+ and CS- can change independently of
each other. To examine this, we compared the amplitude of the average response to CS+ and CS-
trials for all segmented tufts on the pre, last-rewarded, and post sessions (Fig.3A; n = 7 mice; 465
pre, 442 last-rewarded, and 430 post tufts). In agreement with our previous analysis, we found no
significant bias toward CS+ or CS- during any of the three sessions (Pre: p = 0.20; last-rewarded:
p = 0.43; Post: p = 0.64, sign-rank test). Under naïve conditions during the pre session, most tufts
that responded to air puff stimuli did not strongly prefer the CS+ or CS- (Fig.3A, left). Surprisingly,
on the last-rewarded session and the unrewarded post-conditioning session, we observed a
prominent shift in the response distribution, where many tufts exhibited more selective responses
to one stimulus or the other (Fig.3A, middle and right).

Plasticity can occur after repeated exposure to stimuli even in the absence of reinforcements (Chu
et al., 2016; Dragoi et al., 2002; Dragoi et al., 2000; Yao and Dan, 2001; Zhang et al., 2015). To
test whether enhanced selectivity depended on reinforcement, we imaged a separate group of mice
that were repeatedly exposed to the same stimuli for the same number of days but without any
reward. Repeated exposure mice exhibited a stable distribution of response selectivity over time
(Fig.3B; n = 7 mice; 317, 313, and 321 tufts on Day 1, Day 8, and Day 9, respectively). These
results suggest that reinforcement learning, and not simply repeated stimulus exposure, drives
apical tufts to become more selective for either the CS+ or CS-.

To directly quantify the response selectivity of tufts, we computed a selectivity index (SI; see
Methods) ranging from -1 (exclusively CS- responsive) to 1 (exclusively CS+ responsive) for each
tuft. Initially in both the conditioned and repeated exposure mice, the SI distribution was centered
around zero, indicating that most tufts in naïve animals did not strongly prefer either stimulus
(Fig.3C,D, left panels). Consistent with our other analyses (Fig.2D), the mean SI remained close to
zero for each of the three sessions (-0.049, -0.001, and 0.003 for pre-conditioning, last rewarded,
and post-conditioning days, respectively), confirming that learning produced no overall bias toward
one particular stimulus among the population. During learning, the SI distribution of conditioned
but not repeated exposure mice shifted markedly, whereby a much greater proportion of neurons
were highly selective for either the CS+ or CS- (Fig.3C,D, middle and right panels, |SI| pre versus
last-rewarded: $p < 10^{-6}$, |SI| pre versus post: $p < 10^{-5}$; Wilcoxon rank sum test). These effects can
even be observed within individual mice, with learning significantly increasing tuft selectivity in
individual conditioned mice, but not repeated exposure mice (Supp.Fig.2). The degree of
enhancement in tuft selectivity was closely correlated with conditioned animals' ability to
discriminate stimuli across sessions (Fig.3E; Pearson's R = 0.60, $p < 10^{-5}$).

Whereas selectivity magnitude (|SI|) only considers the amplitude of tuft responses to CS+ and CS-
, their discriminability also depends on their variability. For example, a large difference in CS+ and
CS- responses would not be discriminable if the variability of those responses were very high; a
small difference might be discriminable if the variability were low. We therefore additionally
calculated a d-prime metric of neural discriminability that normalizes differences in response
magnitudes to each stimulus by their variability (see Methods). In conditioned animals, neural
discriminability of CS+ and CS- responses of tufts increased significantly across learning (Fig.3F,
blue; first-rewarded versus last-rewarded: $p < 10^{-3}$, pre versus post: $p < 10^{-4}$; Wilcoxon rank sum
test). By contrast, neural discriminability of tuft responses in the repeated exposure mice decreased

slightly with progressive exposure to the stimuli (Fig.3F, gray; Day 1 versus Final: p < 0.01). Taken
together, these results show that enhanced stimulus representations can emerge in apical tufts, but
require reinforcement.
207
The above analyses rely on the accurate measurement of calcium spikes from individual tufts.
While two-photon microscopy acquires images with high resolution and speed, the imaging field
is restricted to a single focal plane. This method can only measure calcium signals from a thin
cross-section of the three-dimensionally complex apical structures. Indeed, many of the spatial
components extracted from our two-photon data were comprised of dendritic branches that cross
the imaging plane at different locations (Supp.Fig.3A), which makes it difficult to determine
whether the segmentation software accurately extracted signals from one tuft or erroneously
merged multiple tufts. For the same reasons, a single apical tuft could be falsely classified as two
different tufts. Such errors could mislead our interpretation of selectivity in the population,
especially given that a single apical tuft can exhibit non-homogenous branch-specific events
(Cichon and Gan, 2015; Larkum et al., 2009; Palmer et al., 2014).
219
To confirm that our interpretation was not due to segmentation errors, we repeated the conditioning
experiment using a new, high-speed volumetric imaging approach called SCAPE (Bouchard et al.,
2015; Hillman et al., 2018), which allowed us to monitor calcium across entire apical tufts
(Supplementary Movie 2). These three-dimensional datasets ($300 \times 1050 \times 234$ μm field of view)
encompassed large portions of the apical tree which included branches converging on their
bifurcation points in layer 2, enabling us to identify whole apical trees unambiguously (Fig.4A,B;
Supp.Fig.3B). CaImAn effectively demixed overlapping trees in these three-dimensional volumes.
Using SCAPE microscopy, we imaged tuft activity of additional mice conditioned with the same
behavioral paradigm (Fig.4C). Comparison of tuft responses to the CS+ and CS- on the pre, last-
rewarded, and post sessions (Fig.4D; 241 pre, 215 last-rewarded, 150 post tufts in 2 mice) revealed
again that task learning induced significant increases in tuft selectivity (Fig.4E; pre versus last-
rewarded: $p < 10^{-5}$, pre versus post: $p < 10^{-4}$, Wilcoxon rank sum test of |SI|). On average, the SI
magnitudes were similar between tufts imaged using 2-photon microscopy and SCAPE (mean ±
s.e.m. |SI| for 2-photon versus SCAPE; pre: 0.41±0.01 versus 0.40±0.02; last-rewarded: 0.54±0.02
versus 0.54±0.02; post: 0.51±0.02 versus 0.53±0.03). These data demonstrate that the effects in our
two-photon dataset are not caused by errors in segmentation, but rather reflect changes at the level
of individual dendritic tufts. Our results, based on two different imaging approaches, clearly
demonstrate that reinforcement increases stimulus selectivity at the level of the entire apical tuft.
238
Selective tufts emerge from both initially unresponsive and responsive populations
The striking effect of reinforcement learning on tuft response selectivity could develop in several
ways. For example, initially unresponsive tufts could develop a robust response to either stimulus
after learning (e.g., Fig.5A, top). Conceivably, tufts that were initially unselective in naïve animals
could also maintain their response to one stimulus while losing their response to the other (e.g.,
Fig.5A, middle). Either or both scenarios could lead to the increase in neurons that are selective for
stimulus direction. To investigate which changes in individual tufts underlie population-wide
improvements in stimulus selectivity, we longitudinally tracked the same set of tufts across all
sessions and compared their selectivity in pre- and post-conditioning sessions for both conditioned
and repeated exposure mice.
249
First, we categorized tufts that were unresponsive to either stimulus on the first imaging session,
which accounted for the large majority of tufts (conditioned: 458/603; repeated exposure: 334/457),
and compared their response to the CS+ and CS- on the last session to determine if they became
selective (Fig.5B, see Methods). Stimulus-unresponsive tufts, while on average less active than
responsive ones (median calcium events per minute: 2.65 versus 3.66 for stimulus-unresponsive

and responsive tufts, respectively; $p < 10^{-40}$, Wilcoxon rank sum test; Supplementary Fig.4), were
not silent, with many undergoing tuft-wide calcium influx several times per minute. Silent tufts that
are never active during the session may not have been detected in our imaging, but we were able to
detect tufts that discharged as few as 3 voltage-gated calcium spikes over a 30-minute behavioral
session. Interestingly, in both the conditioned and repeated exposure mice, approximately 40% of
initially unresponsive tufts developed a response to at least one stimulus by the last session,
becoming either selective or unselective (Fig.5B). However, in conditioned animals, the proportion
of initially unresponsive tufts that became selective was significantly larger than in repeated
exposure mice ($p < 0.05$, 2-sample t-test comparing subjects). Furthermore, while the proportion
of selective and unselective tufts in this category was similar for conditioned animals, unselective
tufts were more common in repeated exposure mice ($p < 0.05$, paired t-test).
266
Next, we analyzed tufts that were initially responsive and either selective (Fig.5C; conditioned:
56/603, RE: 43/457) or unselective (Fig.5D; conditioned: 89/603, repeated exposure: 80/457). In
these smaller categories, we found no significant differences in the outcome of selectivity between
the two groups of animals. Together, these results indicate that, while both stimulus exposure and
reinforcement can alter tuft tuning, the presence of reward increases the likelihood that initially
unresponsive tufts develop selectivity for either the CS+ or CS- (summarized in Fig.5E).
273
While a greater proportion of tufts from the conditioned animals were selective during the final
session (20.2% versus 10.3% of tufts from conditioned and repeated exposure mice, respectively),
we wondered whether conditioning also impacted the degree of selectivity. First, we compared the
SI of initially unresponsive tufts on the final imaging session (Fig. 5F). Supporting our results in
Fig. 5B, the SI distribution was shifted toward the tails in conditioned, but not repeated exposure
mice, indicating that reward enhances selectivity for either the CS+ or CS- in this subset (|SI|
conditioned versus repeated exposure: $p < 10^{-5}$, Wilcoxon rank sum test).
281
Next, we compared the |SI| of all tufts that were categorized as selective during the last imaging
session in conditioned and repeated exposure mice (Fig. 5G). Interestingly, we found that even
among selective tufts, the |SI| distribution in conditioned mice was significantly greater than in
repeated exposure mice ($p = 0.006$, Wilcoxon rank sum test), indicating that while selective tufts
are present after both conditioning and repeated stimulus exposure, the magnitude of selectivity is
stronger after conditioning.
288
289 We then quantified the change in |SI| of all tufts that were responsive in both the first and last
290 sessions by computing the difference between the two sessions (Fig. 5H). Tufts in conditioned mice
291 exhibited a greater increase in |SI| across sessions compared to repeated exposure mice ($p = 0.01$,
292 Wilcoxon rank sum test), demonstrating that the magnitude of selectivity in initially responsive
293 tufts increases after reinforcement learning.
294
Finally, we found that the SI of tufts that eventually became unresponsive on the last session was
similar between the two groups (Fig.5I, |SI| conditioned versus repeated exposure: $p = 0.06$,
Wilcoxon rank sum test). Therefore, the loss of stimulus responsivity that occurred in many tufts
cannot account for differences in selectivity after conditioning or repeated exposure, which must
instead involve tufts that maintain or acquire selectivity. In summary, our longitudinal analyses
revealed that reinforcement learning biases initially unresponsive tufts toward becoming selective
and enhances the selectivity of tufts that are initially responsive.
302
Neither movement nor behavioral choice account for enhanced selectivity
Several plausible factors could underlie the changes in selectivity we observed across learning. For
instance, movements like whisking are correlated with layer 5 somatic action potentials (de Kock

and Sakmann, 2009; Derdikman et al., 2006; Rodgers et al., 2021) and might have impacted
calcium activity in the apical tuft. To investigate whether whisking could account for the changes
in tuft selectivity, we imaged the whiskers with a high-speed camera and computed whisking
amplitude (see Methods) while mice underwent conditioning and two-photon imaging (Fig.6A).
First, we considered whether animals changed their whisker movements in response to conditioned
stimuli over the course of learning. We computed the peak of the mean stimulus-aligned whisking
amplitude for the CS+ and CS- (Fig.1C, left; Fig.6B) for each session. Although conditioning alters
licking behavior (Fig.1C,E), the magnitudes of whisker movements following both stimuli were
stable across sessions (Fig.6B; CS+: p = 0.44; CS-: p = 0.45; linear regression). We also computed
the standard deviation (SD) of stimulus-evoked whisker amplitude across trials for all mice and
sessions (Fig. 6C). While the whisking amplitude became slightly more reliable (decreased SD)
across sessions ($p < 10^{-4}$), the change in reliability across sessions was similar for CS+ and CS- (p
= 0.53). Therefore, whisking is similar on both trial types throughout learning.

We next examined whether whisking was correlated with tuft calcium activity by comparing
stimulus-triggered averages and intertrial interval (ITI) whisk-triggered averages of all tufts during
post-conditioning. Whisking amplitude was similar between spontaneous ITI whisking bouts and
evoked whisking responses to stimuli (p = 0.53, Wilcoxon rank sum test). In contrast to air puff
stimuli, ITI whisking bouts were not associated with a robust calcium response (Fig.6D).

To quantify the relationship of whisking and sensory stimuli to tuft calcium spikes, we performed
a linear regression analysis (see Methods) on 322 tufts using calcium influx as the response variable
and either stimulus or whisking amplitude as a single predictor variable (Fig.6E). Air puff stimuli
more reliably predicted calcium influx than whisking amplitude for each of virtually all tufts (p <
10^{-12}, sign rank test). These results are consistent with other studies that found either only weak or
no correlation between whisking and L5 tuft calcium spikes in S1 (Lacefield et al., 2019; Takahashi
et al., 2020; Xu et al., 2012). Furthermore, we found no relationship between the whisking response
and the median SI magnitude on a given session (Fig.6F, whisking to CS+ p = 0.22, CS- p = 0.78).
Therefore, changes in whisker movement cannot account for the changes in selectivity during
learning that we observed.

Finally, the possibility remains that other task-related signals relaying information about reward
expectation and behavioral choice could impact apical tuft activity and drive increases in
selectivity. To test this, we compared tuft responses to the CS- in false alarm trials (FA; mouse
incorrectly licked for reward) and correct rejection trials (CR; mouse correctly withheld licks) to
determine if their activity was modulated by behavioral choice. Notice that these two trial types
have the same sensory input but involve different choices. (The corresponding analysis for CS+
trials is not technically possible for lack of sufficient Miss trials after the first conditioning day, an
issue also observed in (Poort et al., 2015).) Tufts were classified as behaviorally modulated if the
FA response was significantly different from the CR response, and were not behaviorally
modulated if CR and FA responses were statistically indistinguishable (e.g. Fig.7A). Behaviorally
modulated tufts accounted for only ~10% of the total tuft population in both early and late learning
(50/395 in early; 35/406 in late learning).

To test whether these behaviorally modulated tufts contributed to increased selectivity during
learning, we excluded them and compared selectivity of the remaining behaviorally-insensitive
tufts. We found that selectivity increased significantly from early to late learning (Fig.7B,C; median
|SI| of 345 tufts early versus 371 tufts late learning: 0.38 versus 0.47, p = 0.02, Wilcoxon rank sum
test), similar to our previous analysis of the entire population. Licking, like whisking, was a
relatively poor predictor of tuft calcium influx (Supp.Fig.5A,B). Because some behaviorally
modulated tufts may not have been statistically detectable, we used multivariate linear regression

to disentangle stimulus responses from licking and whisking, which may have been confounded
with choice. Median coefficients for licking and whisking were on average 3.3 times smaller than
median stimulus coefficients for the first rewarded, last rewarded, and post sessions (all $p < 10^{-6}$,
Wilcoxon rank sum test). Even after we factored out possible effects of movements, CS+ and CS-
coefficients were enhanced by learning but not repeated exposure (Supp.Fig.5C,D), consistent with
our other analyses. Together, these results demonstrate that enhanced selectivity during learning
cannot be explained by non-sensory signals related to the animals' behavior.
364
Enhanced selectivity in barrel cortex is long-lasting when mice exclusively use whiskers
Mice could conceivably exploit other sensory cues to learn and perform the task, such as auditory
cues from the air nozzles or non-whisker tactile cues from air current eddies contacting the fur or
skin. To determine which mice exclusively used their whiskers to distinguish the CS+ and CS-, we
trimmed all whiskers after the post-conditioning session and assessed performance in five mice.
Performance in each of the five mice decreased after whisker trimming, indicating that each used
some whisker information. Three mice performed the task exclusively with their whiskers, falling
to chance levels after the whisker trim ("whiskers only"). Two other mice still performed the task
above chance after the whisker trim, indicating that they were not exclusively using their whiskers
and exploited information from multiple sensory streams ("whiskers + other senses").
375
We examined whether these two different behavioral strategies impacted tuft selectivity. Both the
"whiskers only" and "whiskers + other senses" groups exhibited enhanced tuft selectivity in the
last-rewarded session relative to pre-conditioning. This effect was more pronounced in the
"whiskers only" mice (Fig.8A,B, left and middle; whiskers only: median |SI| of 180 pre tufts versus
169 last-rewarded tufts: 0.36 versus 0.59, $p < 10^{-3}$; "whiskers + other senses": median |SI| of 144
pre tufts versus 155 last-rewarded tufts: 0.39 versus 0.50, $p = 0.01$). Surprisingly, enhanced
selectivity persisted during the post-conditioning session for the "whiskers only" group but not the
"whiskers + other senses" group (Fig.8A,B right panels; whiskers only: median |SI| of pre versus
167 tufts post: 0.36 versus 0.58; $p < 10^{-3}$; whiskers + other senses: median |SI| of 155 pre versus
post tufts: 0.39 versus 0.42; $p = 0.45$). Therefore, tuft selectivity in barrel cortex is enhanced
regardless of behavioral strategy, but outlasts conditioning only when mice rely solely on their
whiskers to perform the task.
388
We further examined this persistence of enhanced tuft selectivity as experienced mice stopped
performing the task. While the entire post-conditioning session was unrewarded, mice initially
expected rewards and licked for many CS+ trials in the first half of the session. By the second half
of the session, the probability of a lick occurring during the CS+ extinguished, approaching zero
(Fig.8C). We compared the selectivity of tufts during the first and second halves of the post-
conditioning sessions of mice that exclusively used their whiskers and found no difference in the
two distributions (Fig.8D, $p = 0.94$, Wilcoxon rank sum test of |SI|), demonstrating that selectivity
of the population remained stable throughout the session. Taken together, these results demonstrate
that enhanced stimulus selectivity of apical tuft dendrites after reinforcement learning is long
lasting, persisting even after mice cease performing the task and expecting reward.
399
400   **DISCUSSION**
401
Our study is the first to investigate how learning a discrimination task alters apical tuft activity.
Using both novel volumetric whole-tuft imaging and conventional planar microscopy, we
discovered that L5 apical tufts acquire enhanced representations of multiple stimuli during learning.
Rather than simply retuning tufts toward the rewarded stimulus, learning enhanced selectivity for
both stimuli, suggesting that tufts are aligning themselves to the behaviorally relevant stimulus
dimensions. These enhanced sensory representations persist even after mice cease performing the

task. In contrast, representations are slightly degraded by mere repeated exposure to stimuli outside
of a task. Consistent with previous studies (Lacefield et al., 2019; Xu et al., 2012), we found that
movement in and of itself has little direct impact on tuft spikes, indicating that increased selectivity
of apicals reflects alterations in sensory coding as animals learn. This sensitization of tufts to
behaviorally relevant sensory dimensions may be a general feature of all sensory cortical areas.

Tuft spikes enhance plasticity of synaptic inputs that occur over behavioral (seconds-long)
timescales (Bittner et al., 2017; Roelfsema and Holtmaat, 2018). These new behaviorally relevant
tuft representations may therefore prime subsequent plasticity of synapses across the entire
pyramidal neuron. Additionally, tuft events potently modulate somatic burst firing and enhance
how somata respond to their basal inputs (Larkum, 2013; Larkum et al., 2009). As learning and
plasticity increase apical selectivity for a behaviorally relevant axis, tuft events will unavoidably
amplify somatic burst output along the same axis. This could enable action potential output of L5
cells in primary sensory cortex to directly drive behavioral responses via projections to movement
related areas, such as the corticostriatal, corticopontine, and corticotrigeminal pathways. Thus, tuft
spikes have the potential to modify somatic output, both in the present and in the future.

Enhanced Representation of Behaviorally Relevant Stimuli

Enhancing the representation of relevant stimulus dimensions rather than a singularly important
stimulus, such as a rewarded event, has multiple benefits for behavior. In our paradigm, both the
CS+ and CS- are predictive of whether or not a reward will occur in the future. Explicitly encoding
both stimuli could allow sensory cortical areas to directly elicit actions. In the context of this task,
CS+ preferring tufts in barrel cortex may trigger anticipatory licking while CS- preferring tufts
could suppress licking. L5 cells in sensory cortex via their output to striatum, pons, brain stem, and
spinal cord would thereby be able to directly and rapidly drive action without further cortical
processing, such as by frontal areas including motor cortex (Park et al., 2020; Takahashi et al.,
2020). Such rapid sensory-motor transformations by primary sensory areas may be critical for
natural time-constrained behavior.

Furthermore, learning produced a representation in which the degree of selectivity for the two
stimuli was continuous and uniformly distributed. Exclusively CS+ or CS- selective apicals never
dominated the population. Continuous degrees of selectivity across the population, rather than
discrete representations, may allow the system to be more robust to the variability caused by active
movements that alter sensory input. A continuous distribution may also facilitate future adjustments
of neural representations as subjects continue to learn a task or encounter new tasks. The uniformity
we observed may reflect that neurons are high-dimensional, being sensitive to mixtures of variables
(Kim et al., 2020; Rigotti et al., 2013; Rodgers et al., 2021; Stringer et al., 2019), only one of which
might be altered here by learning. The uniform distribution of selectivity corresponds to a full range
of pessimism to optimism concerning stimulus predictions of upcoming rewards. Recent work
shows that behavioral performance benefits from reinforcement learning that incorporates the
distribution of reward probabilities rather than just the average expected reward value (Dabney et
al., 2020). L5 corticostriatal synapses could theoretically afford a plastic substrate for acquiring the
necessary distribution of reward probabilities.

Surprisingly, past studies in which mice were trained to associate one or more stimuli with a reward
typically show that cortical representations are stronger for the rewarded stimulus (Goltstein et al.,
2013; Henschke et al., 2020; Poort et al., 2015). In contrast to these studies of layer 2/3 somatic
activity, our experiments revealed that the overall tuft calcium response to the CS+ and CS- at the
population level did not change significantly after animals learned the task (Fig.2). Instead,
representations for both stimuli were enhanced by individual tufts developing selectivity for either

the CS+ or the CS- (Fig.3). This divergence in phenomena may result from several important
differences between our work and the aforementioned studies.
461
First, enhanced selectivity for both rewarded and unrewarded stimuli could be a phenomenon that
is unique to the apical dendritic tufts. In addition to local inputs, the apical tufts of pyramidal cells
in S1 receive long-range top-down input from several sources, including motor cortex (Petreanu et
al., 2012; Xu et al., 2012), secondary somatosensory cortex (Cauller et al., 1998), and secondary
thalamus (Rubio-Garrido et al., 2009; Wimmer et al., 2010; Zhang and Bruno, 2019). Frontal areas,
such as prefrontal cortex, indeed have enhanced representations of the CS+ and CS- after learning
(Wang et al., 2020). In contrast, input to the somata is dominated by the local cortical area and
primary thalamus (Constantinople and Bruno, 2013; Feldmeyer, 2012). While somato-dendritic
coupling can be strong in L5 neurons (Beaulieu-Laroche et al., 2019), it is asymmetric; at least 40%
of somatic transients attenuate in a distance-dependent manner along the apical trunk and distal
tufts (Francioni et al., 2019). The non-overlapping anatomical inputs and asymmetric coupling
together could produce different learning-related effects on apical tuft and somatic stimulus
representations.
475
Second, learning-related changes may manifest differently in layer 2/3, the usual focus of previous
studies (Henschke et al., 2020; Poort et al., 2015), and layer 5 pyramidal cells, the tufts of which
we studied. With the exception of a small population of corticostriatal cells, most excitatory cells
in layer 2/3 project to other cortical areas to affect further cortical processing (Petersen and Crochet,
2013; Yamashita et al., 2018). In contrast, many L5 cells project to subcortical structures including
the thalamus, superior colliculus, and brainstem, which may directly trigger behavioral responses
(Krauzlis et al., 2013; Llinas et al., 1998; Parvizi and Damasio, 2001). In discrimination paradigms,
both stimuli are relevant to behavior. In our task, the CS+ prompted licking to obtain a reward, and
the CS- suppressed licking that would have no benefit. Thus, an enhanced representation of both
stimuli in layer 5 would be advantageous for animals to perform the task efficiently. Recently, it
was shown that apical dendrite activation of subcortical-targeting pyramidal tract L5 cells, but not
intratelencephalic L5 cells that are more like L2/3 cells in their connectivity, determines the
detection of tactile stimuli (Takahashi et al., 2020). The Rbp4-Cre mice we used in this study labels
a heterogenous population of layer 5 pyramidal cells, comprising both pyramidal tract and
intratelencephalic neurons. In the future, it would be interesting to examine whether learning has
different effects on the sensory representations of these two populations. Moreover, direct
comparisons of the layers would be particularly informative.
493
Candidate Plasticity Mechanisms
Enhanced selectivity could be due to changes in local synaptic connectivity, long-range inputs, or
both. Learning may strengthen and weaken synapses onto barrel cortex neurons from ascending
thalamocortical input or from neighboring cells. Such local plasticity could enhance CS+ or CS-
responsiveness. Alternatively or additionally, other cortical regions encoding task context could via
long-range inputs reconfigure barrel cortex to respond more strongly to these stimuli. The present
results do not completely distinguish between these two scenarios because long-range inputs may
still encode the context while the mouse is in the behavioral apparatus. However, we found that
enhanced representations persist after mice are no longer engaged in the task and receiving rewards.
This result suggests that enhanced representations may be a product of local plasticity in sensory
cortex that alters receptive fields.
505
Even in the absence of reward, repeated exposure to stimuli can drive plasticity in sensory cortex
and alter response tuning. For instance, repeated exposure to oriented gratings can alter the
orientation tuning of cells in primary visual cortex (Dragoi et al., 2002; Dragoi et al., 2000; Yao
and Dan, 2001), and overstimulation of whiskers induces plasticity at dendritic spines and alters

whisker representations in somatosensory cortex (Feldman and Brecht, 2005; Zhang et al., 2015).
Our results demonstrate that at the population level enhanced representations developed only when
stimuli were behaviorally relevant. Our longitudinal analysis revealed that while the response
dynamics of some tufts changed after repeated stimuli presentations, overall selectivity of the
population did not increase when rewards were omitted (Figs.3&5). This raises the question: What
are the mechanisms that drive enhanced selectivity under rewarded conditions? In one possible
scenario, reward delivery causes the release of neuromodulators that augment the activity of apical
tufts. Cortical layer 1 is innervated by cholinergic afferents from the nucleus basalis (Mechawar et
al., 2000) and adrenergic afferents from the locus coeruleus (Freedman et al., 1975), the main
source of acetylcholine and norepinephrine, respectively. Salient events such as reward and arousal
lead to the release of these neuromodulators in cortex (Chubykin et al., 2013; Thiele and Bellgrove,
2018), which could increase the excitability of apical dendrites by recruiting disinhibitory circuits
or directly influencing dendritic currents (Brombas et al., 2014; Chubykin et al., 2013; Hangya et
al., 2015; Labarrera et al., 2018). In this model, the release of reward-driven neuromodulators
promotes plasticity and an enhanced representation of temporally aligned sensory inputs. This
phenomenon was demonstrated in auditory cortex, where tones paired with stimulation of the
nucleus basalis shifted the tuning of neurons toward the frequency of the paired stimulus (Froemke
et al., 2007).
528
Why are representations of the CS- equally enhanced when there is no associated reward? One
explanation is that, as mice learn that the CS- indicates absence of reward, the CS- effectively
signals punishment and acquires negative value. Acetylcholine is released in response to aversive
stimuli, and can activate disinhibitory microcircuits that reduce inhibition onto pyramidal cells and
may be essential for learning (Gasselin et al., 2021; Letzkus et al., 2011). Thus, it is possible that
both the CS+ and CS- representations are enhanced by neuromodulatory mechanisms tied to reward
and punishment, respectively. An open question is whether the outcome is due to reinforcement
learning or the behavioral state brought on by the reinforcers rather than their valence. Sensory
cortical plasticity may not be tied to reinforcer valence. Our paradigm creates an environment
where mice benefit from being attentive and engaged in order to maximize reward while
minimizing effort. Previous work has shown that active engagement in a visual discrimination task
was associated with significantly higher selectivity in layer 2/3 cells in visual cortex (Poort et al.,
2015). Task engagement may lead to a sustained increase in neuromodulator release throughout the
conditioning session, priming the apical dendrites for plasticity and the development of selective
responses for task-relevant stimuli as they learn.
544
What determines whether a particular tuft eventually becomes selective for the CS+ or CS-? Our
longitudinal analysis revealed that many tufts that were initially unresponsive to either stimulus
developed a highly selective response to either the CS+ or the CS- (Fig.5). In these tufts, stimulus
preference after learning might be seeded by initially weak, directionally selective inputs on to the
neuron that already exist prior to conditioning and that are potentiated by the learning process. We
also found tufts that initially exhibited robust responses to both stimuli and either lost or
significantly reduced their response to one stimulus after learning. The reduction of an apical
response to a particular stimulus could be driven by local disynaptic inhibition between L5
pyramidal cells mediated by the apical-targeting Martinotti cells (Berger et al., 2010; Kapfer et al.,
2007; Naka and Adesnik, 2016). Through this mechanism, L5 neurons that are selective for a
particular stimulus could inhibit responses to that stimulus in neighboring L5 apical tufts.
Experiments that assess the tuning of excitatory and inhibitory inputs onto apical dendrites as a
function of learning could test such mechanisms.
558
559   In addition to demonstrating increased tuft selectivity with learning, we replicated a surprising
560   phenomenon in a previous instrumental behavior in which a population of apical tufts exhibit

activity around the time of reward (Lacefield et al., 2019). This reward-related activity was observed only during CS+ trials and was most prominent during intermediate conditioning sessions, when most animals were still performing at chance levels, and disappeared completely by the final conditioning session (Supp.Fig.1). Other than this transient effect, unconditioned stimuli did not appear to elicit calcium responses, consistent with our previous findings (Lacefield et al., 2019). The disappearance of this reward-related peak might be attributable to the reward becoming predictable in later stages of learning. In previous classical conditioning experiments, dopaminergic cells exhibit responses to rewards early in learning due to the novelty of an unexpected stimulus. These responses are lost after extended training, as animals learn the association between the CS and reward (Ljungberg et al., 1992; Pan et al., 2005). While dopaminergic terminals are sparse in primary sensory areas, they are not entirely absent, nor are dopaminergic receptors. Furthermore, the excitability of the apical tuft is sensitive to noradrenaline (Labarrera et al., 2018). Interestingly, noradrenergic neurons in the locus coeruleus exhibit a similar phenomenon to dopaminergic neurons, where responses shift from temporal alignment with the reward to a predictive conditioned stimulus after learning (Bouret and Sara, 2004). Such mechanisms could explain why reward-related activity is restricted to early-to-intermediate learning in our paradigm.

Global versus local dendritic spikes

Apical dendrites exhibit not only global spikes that elicit calcium influx across the entire tuft, which we exclusively analyzed here, but also local events known as NMDA spikes, which typically engage short (<30-μm) segments of individual dendritic branches (Larkum et al., 2009; Palmer et al., 2014; Xu et al., 2012). These local, NMDA receptor-dependent events can promote prolonged plasticity within individual dendritic branches in the absence of backpropagating actions potentials, a feature that is unique to the apical dendrites (Sandler et al., 2016). In motor cortex, branch-specific NMDA spikes are crucial for establishing the long-lasting plasticity necessary for learning (Cichon and Gan, 2015), and depolarization provided by multiple local NMDA spikes is thought to be essential for the generation of a global calcium spike triggered by distal synaptic inputs (Larkum et al., 2009). We focused this study on global tuft-wide calcium events, rather than local events. Local events are more difficult to unambiguously identify in planar imaging (Sheffield and Dombeck, 2015), and their existence *in vivo* is still an open question for L5 apicals in barrel cortex (Palmer et al., 2014; Xu et al., 2012). Nonetheless, they may play important roles in plasticity processes that eventually lead to the emergence of global tuft spike selectivity for stimuli. Volumetric microscopy studies, the feasibility of which we showed here, are needed to further investigate the existence of local events in such behaviors as well as examine possible relationships between local and global tuft events during reinforcement learning.

To analyze activity of individual tufts, we segmented these structures based on spatiotemporal covariance (Giovannucci et al., 2019). This method does not discount the possibility of errors where one tuft is split erroneously into two trees, or where two highly correlated tufts are merged. With this in mind, we used volumetric imaging SCAPE microscopy, which allowed us to visualize the apicals in three dimensions and unambiguously screen for such artifacts. The results from SCAPE are quantitatively similar to those from two-photon microscopy, and confirm that our observation of enhanced selectivity with learning is not an artifact of planar imaging.

Stability of learned tuft representations

In contrast to previous studies of discrimination learning (Henschke et al., 2020; Liu et al., 2020; Poort et al., 2015), we included an unrewarded post-conditioning session to examine whether learning-related effects persisted through extinction. Our results show that post-conditioning selectivity of the apical population remains significantly higher than pre-conditioning, even after animals stop licking in response to the CS+ (Fig.8). Interestingly, the effects of learning are much more pronounced in animals that relied exclusively on their whiskers to perform the task. In animals

that apparently used other sensory modalities, we observed a modest increase from the pre to last-
rewarded session, which seemed to be largely absent by the post-conditioning session. Considering
that these animals were additionally exploiting other sensory areas to perform, selectivity may have
been more widely distributed and thus diluted in barrel cortex, diminishing the effect and its
stability. How long selectivity persists in the neuronal population after conditioning and which
factors influence stability are interesting open questions for future study.
618
Conclusion
In summary, we have shown for the first time that reinforcement learning enhances representations
along behaviorally relevant dimensions in apical tufts. Our results suggest that dendritic calcium
spikes are an important cellular mechanism underlying the changes in sensory encoding that occur
with learning, and provide an avenue for further investigation of cellular and circuit mechanisms
underlying plasticity induced by perceptual experience and reinforcement. This cellular
compartment may be key to understanding pathology in some cognitive, memory, and learning
disorders.
627

ACKNOWLEDGEMENTS
We thank Venkatakaushik Voleti for help with the design, construction, and alignment of the SCAPE microscope; Dan Kato, Georgia Pierce, and Jung Park for help with pilot experiments; Eftychios Pnevmatikakis and Johannes Friedrich for advice on dendrite segmentation; and Larry Abbott, Stefano Fusi, Ashok Litwin-Kumar, Chris Rodgers, Georgia Pierce, Gordon Petty, and Dan Kato for comments on the manuscript. Funding was provided by NIH/NINDS R01 NS069679, NIH/NINDS R01 NS094659, a Grossman-Kavli Scholar Award (RMB), a Kavli Institute for Brain Science Postdoctoral Fellowship (SEB), NIH/NINDS/NIMH/BRAIN U01 NS094296, UF1 NS108213, U19 NS104649, and RF1 MH114276 (EMCH).

AUTHOR CONTRIBUTIONS
SEB and RMB conceived of the behavioral and two-photon imaging experiments. EMCH and RMB conceived of the SCAPE imaging experiments. SEB built the behavioral apparatus, EMCH, KBP, and CC designed, built, and maintained the SCAPE microscope, and RMB built and maintained the two-photon and intrinsic signal microscopes. SEB performed the experiments and analyzed the data with input from RMB and EMCH. SEB and RMB wrote the manuscript.

MAIN FIGURES

Figure 1. Mice rapidly learn to discriminate stimulus direction in head-fixed paradigm.
(A) A water droplet is paired with air puffs in one direction (CS+) but not the other (CS-).
Licking in anticipation of water is assessed in the response window just after CS+ or CS- and
prior to water delivery for the CS+ (grey bar).
(B) Experimental timeline. 2-3 weeks after virus injection, naive tuft responses to stimuli are
recorded (pre). The CS+ is then paired with water for 8-9 days (blue). On the last day, stimuli are
presented without reward (post). In a separate group of mice, the same stimuli are presented over
9 days in the absence of reward (unrewarded group).
(C) Lick rasters for three different sessions in one example mouse. On session 9, the CS+ but not
the CS- reliably elicits licks.
(D) Mean baseline-subtracted whisking amplitude aligned to the CS+ (red) and CS- (navy) across
sessions 1, 2, and 9 of an example mouse.
(E) Learning curve demonstrates rapid learning. Mean probability of at least one lick in the
response window across sessions.
(F) Behavioral performance of each mouse in the rewarded group (M1 – M7).

**Figure 2. Overall tuft response to stimuli is unbiased and relatively stable across
conditioning.**
(A) Dendritic activity was recorded in layer 1 (i) in the C1/C2 barrel columns (ii). (i) Two-photon
image ~60 µm deep relative to pia. Dashed yellow lines denote C1 and C2 boundaries from
intrinsic imaging. Reconstruction from (Ramirez et al., 2014).
(ii) Tangential section through layer 4 showing barrels stained with streptavidin-Alexa 647 and
GCaMP6f-expressing apical trunks. Red circles indicate location of 2-photon lesions to mark the
imaging region for post-hoc analysis.
(B) Overlay of five segmented pseudo-colored tufts from imaging field in A(i).
(C) Time courses of calcium responses of example tufts in (B) to three air puffs (dashes).
(D) Amplitude for CS+ (red) and CS- responses (blue), computed for each segmented tuft in the
first 1.5 s post-stimulus (grey points), do not differ within or across sessions. Colored lines
indicate median.
(E) Same as in D, showing data for all conditioning sessions.

**Figure 3. Reinforcement learning, but not stimulus exposure, enhances tuft selectivity for
CS+ and CS- stimuli.**
(A) Across the indicated sessions, individual tufts (circles) exhibit larger biases to CS+ or CS-
(pooled across all conditioned mice).
(B) Repeated exposure to stimuli does not bias individual tufts to CS+ or CS-.
(C) Conditioning reshapes distribution of selectivity indices for tufts from Normal on pre-
conditioning session to uniform on post-conditioning session.
(D) Distribution of tuft selectivity indices remains Normal throughout all repeated exposure
sessions.
(E) Selectivity (median SI magnitude of tufts for each session) increases with behavioral
performance of 6 animals.
(F) Neural discriminability (mean ± sem) of tufts, pooled across all animals on each session,
increases with conditioning and decreases with repeated exposure.

**Figure 4. High-speed volumetric imaging of apical tufts confirms the emergence of
enhanced selectivity after learning.**
(A) Top and side view of four example tufts segmented from volumetric SCAPE imaging.

(B) Time courses of calcium activity from example tufts in (A) during five presentations of air puff stimuli (dashes).

(C) Performance across all conditioning sessions of two mice that were imaged with SCAPE.

(D) Across the indicated sessions, individual SCAPE-imaged tufts (circles) exhibit larger biases to CS+ or CS-.

(E) Conditioning reshapes selectivity distribution from Normal to uniform.

Figure 5. Longitudinal tracking reveals that reward enhances the selectivity of both initially unresponsive and responsive tufts.

(A) Three example tufts that were longitudinally tracked across learning. Top row: An initially unresponsive tuft develops a robust response to the CS+ but not the CS- after learning. Middle row: A responsive but unselective tuft loses its robust CS+ response and becomes selective for the CS-. Bottom row: A CS- selective neuron becomes unresponsive to both stimuli.

(B) Tufts that were unresponsive during the first session were longitudinally tracked to the last session. Plotted is the mean proportion of selective and unselective neurons across all animals in the conditioned (black bars) and repeated exposure (grey bars) groups.

(C,D) Same analysis as C for initially selective (D) and unselective (E) tufts. Two-sample t-test was used for comparisons between conditioned and repeated exposure groups. Paired t-test was used for comparisons within a group. * $p < 0.05$

(E) Total tuft counts from first to last session within the 3 response categories for either conditioned (left) or repeated exposure (right) groups.

(F) SI of responsive tufts on the last session that were initially unresponsive during the first session. Conditioned tufts have enhanced selectivity compared to repeated exposure.

(G) Tufts that were selective on the last session are more selective if conditioned (black) rather than undergoing repeated exposure (grey).

(H) Tufts that responded on both pre and post sessions tend to have higher selectivity if conditioned rather than undergoing repeated exposure.

(I) SI of responsive tufts on the first session that later became unresponsive during the last session.

Figure 6. Whisking is only weakly correlated with tuft activity and cannot account for changes in selectivity during learning.

(A) Whisking amplitude aligned to calcium activity of three example tufts in one session. Green shading indicates periods of whisking. Red and navy ticks indicate CS+ or CS- delivery, respectively.

(B) Mean whisking response of four mice to CS+ (red) and CS- (navy) does not change across sessions during learning (mean ± s.e.m.).

(C) Mean standard deviation of whisking decreases for both CS+ and CS- across learning, but CS+ and CS- do not differ.

(D) Event-triggered averages of 322 tufts on the post-conditioning day (grey traces - individual tufts, black inset - population average) are responsive to stimuli but relatively unmodulated by whisking.

(E) R^2 values for linear models predicting calcium from stimuli (y axis) are consistently greater than those predicting calcium from whisking (x axis). Each circle represents a tuft.

(F) Magnitude of tuft selectivity does not correlate with mean whisking amplitude during CS+ (left) and CS- trials (right) on that session.

Figure 7. Behavioral responses do not account for enhancement of stimulus selectivity during learning.

(A) Mean stimulus responses of four tufts during hit (red), CR (cyan), and FA (black) trials. Top
row: Example tufts whose responses are not behaviorally modulated (CR is similar to FA).
Bottom row: Example tufts with behaviorally modulated responses (CR and FA differ).
(B) Selectivity index (SI) distribution changes from early (left) and late learning sessions (right)
even when tufts with behaviorally modulated responses (CR ≠ FA) are excluded.
(C) Median SI magnitude of tufts in each of six animals (from panel B) increases from early to
late learning sessions.
751
Figure 8. Apical tufts in barrel cortex of mice performing the task exclusively with their
whiskers undergo long-lasting changes in selectivity.
(A) SI histograms of mice performing the task exclusively with their whiskers exhibit increased
selectivity across pre-conditioning, last-rewarded, and post-conditioning sessions.
(B) Relative to pre-conditioning, mice using their whiskers and other sensory cues to perform the
task have increased selectivity during the last rewarded session, but not the post-conditioning
session.
(C) The probability of anticipatory licks in response to the CS+ extinguishes across post-
conditioning blocks (of 20 trials each).
(D) Tuft selectively remains uniformly distributed during post-conditioning trial blocks 1-2 (top)
while licking is extinguishing, and blocks 3-4 (bottom) in which licking is extinguished.

SUPPLEMENTAL FIGURES

Supplementary Figure 1. CS+ trials evoke a second, long-latency peak during early learning, but not late learning.
(A) Left: Population average of stimulus-responsive tufts aligned to CS+ (red) or CS- (blue) trials from an example mouse. Right: Normalized ΔF/F of individual tufts during CS+ trials.
(B) Same as in A, combining data across four mice whose imaging regions were mapped with intrinsic imaging.

Supplementary Figure 2. Selectivity was enhanced in individual animals that received rewards.
Median SI magnitude for each animal across three sessions for conditioned (left) and repeated exposure groups (right). * $p < 0.05$, *** $p < 10^{-3}$

Supplementary Figure 3. Segmented tufts from two-photon and SCAPE microscopy.
(A,B) 12 example tufts extracted from either two-photon (A) or SCAPE microscopy (B). Tufts segmented from SCAPE microscopy are shown as maximum intensity projections from the top and side. Scale bars: 100μm.

Supplementary Figure 4. Calcium event rate of tufts that were either unresponsive or responsive to air puff stimuli.
The number of calcium events per minute was quantified for all tufts during each conditioning session. Data from each group was pooled across all sessions.

Supplementary Figure 5. Licking cannot account for changes in selectivity during learning.
(A) ITI lick-bout-triggered averages of 232 tufts on the 5th conditioning day, when ITI licks were still common (grey traces - individual tufts, black inset - population average), exhibit little or no lick-related calcium influx.
(B) R^2 values for linear models predicting calcium from stimuli (y axis) are consistently greater than those predicting calcium from licking (x axis). Each circle represents one tuft out of 442 tufts on last-rewarded sessions.
(C) Coefficients from a multivariate regression analysis with calcium as the response variable and the CS+, CS-, whisking, and licking as the predictors. CS+ and CS- coefficients are therefore disentangled from correlations with whisking and licking. Conditioning biases individual tufts (circles) to have larger CS+ or CS- coefficients. n = 304, 324, and 322 tufts for First rewarded, Last rewarded, and Post conditioning, respectively.
(D) Similar analysis to C but for repeated exposure group, with calcium as the response variable and the CS+, CS-, and whisking as the predictors. n = 223, 208, and 218 tufts for Day 2, Final - 1, and Final session, respectively.

Supplementary Movie 1. Example two-photon microscopy movie during behavioral session.
Playback speed is in real time. "CS+" and "CS-" denote times of stimulus onset. 433 x 433 μm field of view.

Supplementary Movie 2. Example SCAPE microscopy movie during behavioral session.
Top, maximum intensity projection (MIP) across the dorsal-ventral dimension showing horizontal extent of dendritic activity. *Bottom*, MIP across the medial-lateral dimension showing vertical extent of dendritic activity. Playback speed is in real time. "CS+" and "CS-" denote times of stimulus onset. 300 × 1050 × 234 μm field of view.

METHODS

All experiments complied with the NIH Guide for the Care and Use of Laboratory Animals and were approved by the Institutional Animal Care and Use Committee of Columbia University. Sixteen C57BL/6 mice were used in these experiments.

Surgery

Animals were administered dexamethasone (1 mg/kg) via intramuscular injection 1-4 hours prior to surgery to reduce edema. Anesthesia was induced with 3% isoflurane in oxygen and maintained at 1%. Mice were head-fixed in a stereotax, and a subcutaneous injection of bupivacaine (0.5%, 0.1 mL) was administered under the scalp. Buprenorphine (0.05 mg/kg) was injected subcutaneously on the back. The scalp was cut, and the skull was covered with a thin layer of Vetbond. A circular craniotomy (3-mm diameter) centered at 1.5 mm posterior and 3.5 mm lateral to bregma was made using a dental drill. The dura was kept moist using artificial cerebrospinal fluid.

Rbp4-Cre_KL100 mice were injected with 100 nL of virus (initial titer ~$2x10^{13}$ cfu/mL, diluted 1:4 in artificial cerebrospinal fluid) encoding GCaMP6f in a Cre recombinase-specific manner (AAV1-CAG-flex-GCaMP6f, UPenn Vector Core). The virus was injected in layer 5B of the barrel cortex (1 mm ventral to the pia) using a pulled pipette (20-30 μm ID) fastened on a Nanoject III. This depth was chosen to maximize labeling of thick-tufted pyramidal neurons. In pilot experiments, we found that placing injections 1.0 mm deep resulted primarily in thick-tufted labeling whereas at more superficial depths (e.g., 0.8 mm deep), we obtained mainly thin-tufted tufts, consistent with (Oberlaender et al., 2012). The dura was then removed, and a thin cover glass was implanted and sealed using superglue. A custom metal head plate was implanted on the skull using dental cement. Twenty-four hours after surgery, carprofen (5 mg/kg) was administered subcutaneously. Imaging and behavioral training commenced 3 weeks after surgery.

Behavior

Animals in both rewarded 'conditioning' and unrewarded 'repeated exposure' groups were water restricted for 2 days prior to starting imaging and habituated to head fixation for ~10 minutes on each of these 2 days. They were subsequently given ~1 mL of water per day for 9 days either by pairing water rewards with a specific stimulus (conditioning group), or in their cage following the imaging session (repeated exposure group). Mice were head restrained in a custom-made behavioral apparatus by positioning the body in a 3D-printed chamber and fastening the head plate to metal posts flanking the chamber. Air puff stimuli (10 psi measured before a control solenoid, 100 ms) were delivered from two nozzles (cut P200 pipette tips) positioned toward the distal tips of the whiskers, in either the rostrocaudal or ventrodorsal direction. Nozzles were oriented to prevent air jets from stimulating other parts of the face. One of these directions (CS+) was paired with a water reward (10 μL), delivered through a lick port 0.5 seconds after the stimulus onset. The particular direction (rostrocaudal vs ventrodorsal) used as the CS+ was randomized and counterbalanced across mice. Approximately 180 stimuli were presented over the course of a 30-minute imaging session (8-12-s intertrial interval). The probability of CS+ or CS- delivery was 50%. In preliminary experiments, we found that an auditory mask helped prevent mice from exploiting auditory cues to discriminate the two stimuli: a third air nozzle was positioned close to the mouse and was active throughout the session.

During the first session (pre-conditioning), stimuli were delivered in the absence of reward to assess neural and behavioral responses in naïve animals. In the following 7-9 days, the CS+ was paired with reward. Licks for rewards were detected with a capacitance-based touch sensor (Sparkfun). A trial response was registered when one or more licks were elicited within a 0.5-second response window following the stimulus and before reward delivery. To determine whether behavioral

performance was above chance, we computed 95% confidence intervals using the 'binofit' function
in MATLAB. During the final session (post-conditioning), stimuli were delivered in the absence
of reward. Animals in the unrewarded group received the same two stimuli across 9 days without
reward pairing. Behavioral experiments were performed with the Arduino-based OpenMaze open-
source behavioral system, whose designs are fully described at www.openmaze.org. Whisking was
monitored at 125 fps with a camera (Sony PS3eye) and automatically tracked using published
software (Clack et al., 2012).
875
Intrinsic signal optical imaging and two-photon imaging
Intrinsic signal optical imaging and two-photon imaging were performed on a Sutter movable
objective microscope. The locations of whisker barrels in S1 were identified using intrinsic signal
optical imaging. Single whiskers in isoflurane-anesthetized mice were stimulated at 5 Hz using a
piezoelectric bimorph while recording the reflectance of 700-nm long-pass incandescent light with
a Rolera CCD camera (QImaging) through a low-magnification objective (Zeiss 5X/0.16NA).
Movies were collected using software custom-written in Labview (National Instruments). Regions
of reflectance change were referenced to an image acquired under green illumination.
884
Two-photon imaging was conducted on the same microscope under the control of the ScanImage
software package (V. Iyer, Janelia Farms). Scanning during awake conditions was performed at 30
fps using a Chameleon Ultra II laser (Coherent) tuned to 920 nm, precompensated for group
velocity dispersion and focused through a 20x/1.0NA water immersion lens (Zeiss). Emitted light
was collected with an HQ535/50 filter (Chroma) and GaAsP photomultiplier tubes (Hamamatsu
Photonics). Apical tuft tufts in Layer 1 were imaged at depths of 40-80 μm from the pial surface
(433 x 433 μm field of view, 512 x 512 pixels).
892
SCAPE imaging
High-speed volumetric imaging was performed using a custom SCAPE microscope as previously
described (Bouchard et al., 2015; Hillman et al., 2018; Voleti et al., 2019). Briefly, the cortex was
illuminated with an oblique light sheet through a Olympus XLUMPLFLN 20XW 1.0 NA water
immersion objective with a 2-mm working distance. Fluorescence excited by this sheet (extending
in the y-z' direction) was collected by the same objective lens. A galvanometer mirror in the system
was positioned to both cause the oblique light sheet to scan from side to side across the sample (in
the x direction) but also to de-scan returning fluorescence light. This optical path results in an
intermediate, de-scanned oblique image plane that is stationary yet always co-aligned with the
plane in the sample that is being illuminated by the scanning light sheet. Image rotation optics and
a fast sCMOS camera (Andor Zyla 4.2+) were then focused to capture these y-z' images at >1000
frames per second as the sheet was repeatedly scanned across the cortex in the x direction. All other
system parts, including the objective and sample stage, were stationary during high-speed 3D image
acquisition. Data were reformed into a 3D volume by stacking successive y-z' planes according to
the scanning mirror's x position and de-skewing to correct for the oblique sheet angle.
908
In this study, the stationary objective lens in SCAPE was configured on a manual rotation mount
and set to 20º-30º away from the standard upright configuration, so the optical axis was
perpendicular to the cranial window to achieve optimal performance without tilting the head of the
animal. A 488-nm laser (Coherent OBIS) was used for excitation (<10 mW at the sample) with a
500-nm long-pass filter in the emission path. To achieve optimal spatiotemporal resolution and
volume rate, the sample was imaged with an x-direction scanning step of 3 μm over a 300×1050
$\times 234$ μm field of view (x-y-z, $3.0 \times 1.40 \times 1.17$ μm per pixel) at 10 volumes per second (VPS).
916
917
918

Analysis

Two-photon movies were motion corrected using the NormCorre package (Pnevmatikakis and Giovannucci, 2017) in MATLAB. Spatial and temporal components for individual tufts imaged by two-photon and SCAPE were segmented using CaImAn, which employs large-scale sparse non-negative matrix factorization (Giovannucci et al., 2019; Pnevmatikakis et al., 2016). CaImAn inherently corrects for background signal. All further analyses used custom-written routines implemented in MATLAB. Spatial components with tuft structural characteristics were identified and analyzed, while neuropil components were discarded.

To quantify a tuft's response to stimuli, the mean stimulus-aligned $\Delta F/F$ was computed across all CS+ or CS- trials and corrected by the mean $\Delta F/F$ of the second before the trial. Probability of transients was obtained by taking each trial's $\Delta F/F$ in the first 1.5 seconds following either the CS+ or CS- and fitting these data with a univariate mixture of two Normal distributions: $(1-p)N(\mu_1, \sigma_1) + pN(\mu_2, \sigma_2)$. The smaller Normal reflects the distribution of failures, and the larger Normal the distribution of transient amplitudes following the stimulus. The parameter p captures the probability of transients.

From these data, a selectivity index (SI) was defined as $(F_{CS+} - F_{CS-}) / (F_{CS+} + F_{CS-})$, in which F_{CS+} and F_{CS-} are the mean stimulus-aligned amplitudes ($\Delta F/F$) to the CS+ and CS- within the first 1.5 seconds, respectively. This yielded values that range from -1 (exclusively CS- responsive) to 1 (exclusively CS+ responsive). Neural discriminability was defined as $d' = |F_{CS+} - F_{CS-}| / \sqrt{((\sigma^2_{CS+} + \sigma^2_{CS-})/2)}$ where σ^2_{CS+} is the variance of the response amplitudes in F_{CS+} and σ^2_{CS-} is the variance of the response amplitudes in F_{CS-}.

For longitudinal analysis, tufts were categorized as stimulus responsive if they met two criteria: 1) Across all trials, the mean $\Delta F/F$ 1.5 seconds before and 1.5 seconds after the stimulus were significantly different according to the Wilcoxon rank sum test, for either the CS+ or CS-, and 2) the average response amplitude for that stimulus was greater than 0.04 $\Delta F/F$. Tufts with a significant response to only one stimulus were categorized as highly selective and their |SI| was set to 1. To classify tufts as behaviorally modulated, the mean $\Delta F/F$ of the first 1.5 seconds after the stimulus was computed for false alarm and correct rejection trials and compared with a rank sum test. Only sessions with at least 12 false alarm trials were used for this analysis. If the two distributions were significantly different, the tuft was classified as behaviorally modulated.

Custom MATLAB software was used to compute the median whisker angle, and whisking amplitude was computed as described previously (Petty et al., 2020). The median angle was bandpass filtered from 4 to 30 Hz and passed through a Hilbert transform to calculate phase. We defined the upper and lower envelopes of the unfiltered median whisking angle as the points in the whisk cycle where phase equaled 0 (most protracted) or π (most retracted), respectively. Whisking amplitude was defined as the difference between these two envelopes. Periods of whisking were defined as times where whisking amplitude exceeded 20% of maximum for at least 250 msec. Periods of time where amplitude exceeded this threshold for less than 250 msec were considered ambiguous and excluded from analysis of whisking versus quiescence. The whisking-triggered average for each tuft was computed by aligning the calcium signal to the start times of whisking periods during inter-trial intervals (2-8 seconds after stimulus delivery).

For the linear regression analysis, we excerpted the calcium timeseries 2 seconds before and 6 seconds after each stimulus onset. The whisking amplitude signal was frame aligned to the calcium signal according to the lag of the calcium-whisking cross-correlation peak for each tuft. Whisking amplitude was then normalized to the max, yielding values that ranged from 0 to 1. The stimulus

predictor variable was a binary vector with an 800-msec 'on' period (24 frames) centered at the
stimulus time. The timing of the stimulus variable was then aligned to the calcium signal according
to the latency of peak of the mean $\Delta F/F$ of the first 1.5 seconds relative to the stimulus. The lick
predictor variable was a binary vector with 'on' periods denoting lick bouts. Lick bouts were
defined as periods of time where the mouse elicited at least 2 licks, with a maximum gap of 200
msec, and therefore had variable lengths.
975

References

Amitai, Y., Friedman, A., Connors, B.W., and Gutnick, M.J. (1993). Regenerative activity in apical dendrites of pyramidal cells in neocortex. Cereb Cortex *3*, 26-38.

Banerjee, A., Parente, G., Teutsch, J., Lewis, C., Voigt, F.F., and Helmchen, F. (2020). Value-guided remapping of sensory cortex by lateral orbitofrontal cortex. Nature *585*, 245-250.

Beaulieu-Laroche, L., Toloza, E.H.S., Brown, N.J., and Harnett, M.T. (2019). Widespread and Highly Correlated Somato-dendritic Activity in Cortical Layer 5 Neurons. Neuron *103*, 235-241 e234.

Beitel, R.E., Schreiner, C.E., Cheung, S.W., Wang, X., and Merzenich, M.M. (2003). Reward-dependent plasticity in the primary auditory cortex of adult monkeys trained to discriminate temporally modulated signals. Proc Natl Acad Sci U S A *100*, 11070-11075.

Berger, T.K., Silberberg, G., Perin, R., and Markram, H. (2010). Brief bursts self-inhibit and correlate the pyramidal network. PLoS Biol *8*.

Bernhard, S.M., Lee, J., Zhu, M., Hsu, A., Erskine, A., Hires, S.A., and Barth, A.L. (2020). An automated homecage system for multiwhisker detection and discrimination learning in mice. PLoS One *15*, e0232916.

Bittner, K.C., Milstein, A.D., Grienberger, C., Romani, S., and Magee, J.C. (2017). Behavioral time scale synaptic plasticity underlies CA1 place fields. Science *357*, 1033-1036.

Bouchard, M.B., Voleti, V., Mendes, C.S., Lacefield, C., Grueber, W.B., Mann, R.S., Bruno, R.M., and Hillman, E.M. (2015). Swept confocally-aligned planar excitation (SCAPE) microscopy for high speed volumetric imaging of behaving organisms. Nature photonics *9*, 113-119.

Bouret, S., and Sara, S.J. (2004). Reward expectation, orientation of attention and locus coeruleus-medial frontal cortex interplay during learning. Eur J Neurosci *20*, 791-802.

Brombas, A., Fletcher, L.N., and Williams, S.R. (2014). Activity-dependent modulation of layer 1 inhibitory neocortical circuits by acetylcholine. J Neurosci *34*, 1932-1941.

Bruno, R.M., Khatri, V., Land, P.W., and Simons, D.J. (2003). Thalamocortical angular tuning domains within individual barrels of rat somatosensory cortex. J Neurosci *23*, 9565-9574.

Bruno, R.M., and Sakmann, B. (2006). Cortex is driven by weak but synchronously active thalamocortical synapses. Science *312*, 1622-1627.

Cauller, L.J., Clancy, B., and Connors, B.W. (1998). Backward cortical projections to primary somatosensory cortex in rats extend long horizontal axons in layer I. J Comp Neurol *390*, 297-310.

Chen, T.W., Wardill, T.J., Sun, Y., Pulver, S.R., Renninger, S.L., Baohan, A., Schreiter, E.R., Kerr, R.A., Orger, M.B., Jayaraman, V.*, et al.* (2013). Ultrasensitive fluorescent proteins for imaging neuronal activity. Nature *499*, 295-300.

Chu, M.W., Li, W.L., and Komiyama, T. (2016). Balancing the Robustness and Efficiency of Odor Representations during Learning. Neuron *92*, 174-186.

Chubykin, A.A., Roach, E.B., Bear, M.F., and Shuler, M.G. (2013). A cholinergic mechanism for reward timing within primary visual cortex. Neuron *77*, 723-735.

Cichon, J., and Gan, W.B. (2015). Branch-specific dendritic Ca(2+) spikes cause persistent synaptic plasticity. Nature *520*, 180-185.

Clack, N.G., O'Connor, D.H., Huber, D., Petreanu, L., Hires, A., Peron, S., Svoboda, K., and Myers, E.W. (2012). Automated tracking of whiskers in videos of head fixed rodents. PLoS Comput Biol *8*, e1002591.

Constantinople, C.M., and Bruno, R.M. (2013). Deep cortical layers are activated directly by thalamus. Science *340*, 1591-1594.

Dabney, W., Kurth-Nelson, Z., Uchida, N., Starkweather, C.K., Hassabis, D., Munos, R., and Botvinick, M. (2020). A distributional code for value in dopamine-based reinforcement learning. Nature *577*, 671-675.

David, S.V., Fritz, J.B., and Shamma, S.A. (2012). Task reward structure shapes rapid receptive field plasticity in auditory cortex. Proc Natl Acad Sci U S A *109*, 2144-2149.

de Kock, C.P., and Sakmann, B. (2009). Spiking in primary somatosensory cortex during natural whisking in awake head-restrained rats is cell-type specific. Proc Natl Acad Sci U S A *106*, 16446-16450.

Derdikman, D., Yu, C., Haidarliu, S., Bagdasarian, K., Arieli, A., and Ahissar, E. (2006). Layer-specific touch-dependent facilitation and depression in the somatosensory cortex during active whisking. J Neurosci *26*, 9538-9547.

Dragoi, V., Sharma, J., Miller, E.K., and Sur, M. (2002). Dynamics of neuronal sensitivity in visual cortex and local feature discrimination. Nat Neurosci *5*, 883-891.

Dragoi, V., Sharma, J., and Sur, M. (2000). Adaptation-induced plasticity of orientation tuning in adult visual cortex. Neuron *28*, 287-298.

Feldman, D.E., and Brecht, M. (2005). Map plasticity in somatosensory cortex. Science *310*, 810-815.

Feldmeyer, D. (2012). Excitatory neuronal connectivity in the barrel cortex. Front Neuroanat *6*, 24.

Francioni, V., Padamsey, Z., and Rochefort, N.L. (2019). High and asymmetric somato-dendritic coupling of V1 layer 5 neurons independent of visual stimulation and locomotion. Elife *8*.

Freedman, R., Foote, S.L., and Bloom, F.E. (1975). Histochemical characterization of a neocortical projection of the nucleus locus coeruleus in the squirrel monkey. J Comp Neurol *164*, 209-231.

Fritz, J., Shamma, S., Elhilali, M., and Klein, D. (2003). Rapid task-related plasticity of spectrotemporal receptive fields in primary auditory cortex. Nat Neurosci *6*, 1216-1223.

Froemke, R.C., Merzenich, M.M., and Schreiner, C.E. (2007). A synaptic memory trace for cortical receptive field plasticity. Nature *450*, 425-429.

Gasselin, C., Hohl, B., Vernet, A., Crochet, S., and Petersen, C.C.H. (2021). Cell-type-specific nicotinic input disinhibits mouse barrel cortex during active sensing. Neuron.

Giovannucci, A., Friedrich, J., Gunn, P., Kalfon, J., Brown, B.L., Koay, S.A., Taxidis, J., Najafi, F., Gauthier, J.L., Zhou, P., *et al.* (2019). CaImAn an open source tool for scalable calcium imaging data analysis. Elife *8*.

Glickfeld, L.L., Andermann, M.L., Bonin, V., and Reid, R.C. (2013). Cortico-cortical projections in mouse visual cortex are functionally target specific. Nat Neurosci *16*, 219-226.

Goltstein, P.M., Coffey, E.B., Roelfsema, P.R., and Pennartz, C.M. (2013). In vivo two-photon Ca2+ imaging reveals selective reward effects on stimulus-specific assemblies in mouse visual cortex. J Neurosci *33*, 11540-11555.

Hangya, B., Ranade, S.P., Lorenc, M., and Kepecs, A. (2015). Central Cholinergic Neurons Are Rapidly Recruited by Reinforcement Feedback. Cell *162*, 1155-1168.

Henschke, J.U., Dylda, E., Katsanevaki, D., Dupuy, N., Currie, S.P., Amvrosiadis, T., Pakan, J.M.P., and Rochefort, N.L. (2020). Reward Association Enhances Stimulus-Specific Representations in Primary Visual Cortex. Curr Biol *30*, 1866-1880 e1865.

Hillman, E.M., Voleti, V., Patel, K., Li, W., Yu, H., Campos, C.P., Benezra, S.E., Bruno, R., and Galwaduge, P.T. (2018). High-speed 3D imaging of cellular activity in the brain using axially-extended beams and light sheets. Curr Opin Neurobiol *50*, 190-200.

Kapfer, C., Glickfeld, L.L., Atallah, B.V., and Scanziani, M. (2007). Supralinear increase of recurrent inhibition during sparse activity in the somatosensory cortex. Nat Neurosci *10*, 743-753.

Kim, J., Erskine, A., Cheung, J.A., and Hires, S.A. (2020). Behavioral and Neural Bases of Tactile Shape Discrimination Learning in Head-Fixed Mice. Neuron *108*, 953-967 e958.

Kozorovitskiy, Y., Saunders, A., Johnson, C.A., Lowell, B.B., and Sabatini, B.L. (2012). Recurrent network activity drives striatal synaptogenesis. Nature *485*, 646-650.

Krauzlis, R.J., Lovejoy, L.P., and Zenon, A. (2013). Superior colliculus and visual spatial attention. Annu Rev Neurosci *36*, 165-182.

Labarrera, C., Deitcher, Y., Dudai, A., Weiner, B., Kaduri Amichai, A., Zylbermann, N., and London, M. (2018). Adrenergic Modulation Regulates the Dendritic Excitability of Layer 5 Pyramidal Neurons In Vivo. Cell Rep *23*, 1034-1044.

Lacefield, C.O., Pnevmatikakis, E.A., Paninski, L., and Bruno, R.M. (2019). Reinforcement Learning Recruits Somata and Apical Dendrites across Layers of Primary Sensory Cortex. Cell Rep *26*, 2000-2008 e2002.

Larkum, M. (2013). A cellular mechanism for cortical associations: an organizing principle for the cerebral cortex. Trends Neurosci *36*, 141-151.

Larkum, M.E., Nevian, T., Sandler, M., Polsky, A., and Schiller, J. (2009). Synaptic integration in tuft dendrites of layer 5 pyramidal neurons: a new unifying principle. Science *325*, 756-760.

Larkum, M.E., Senn, W., and Luscher, H.R. (2004). Top-down dendritic input increases the gain of layer 5 pyramidal neurons. Cereb Cortex *14*, 1059-1070.

Larkum, M.E., and Zhu, J.J. (2002). Signaling of layer 1 and whisker-evoked Ca2+ and Na+ action potentials in distal and terminal dendrites of rat neocortical pyramidal neurons in vitro and in vivo. J Neurosci *22*, 6991-7005.

Larkum, M.E., Zhu, J.J., and Sakmann, B. (2001). Dendritic mechanisms underlying the coupling of the dendritic with the axonal action potential initiation zone of adult rat layer 5 pyramidal neurons. J Physiol *533*, 447-466.

Letzkus, J.J., Wolff, S.B., Meyer, E.M., Tovote, P., Courtin, J., Herry, C., and Luthi, A. (2011). A disinhibitory microcircuit for associative fear learning in the auditory cortex. Nature *480*, 331-335.

Liu, D., Deng, J., Zhang, Z., Zhang, Z.Y., Sun, Y.G., Yang, T., and Yao, H. (2020). Orbitofrontal control of visual cortex gain promotes visual associative learning. Nat Commun *11*, 2784.

Ljungberg, T., Apicella, P., and Schultz, W. (1992). Responses of monkey dopamine neurons during learning of behavioral reactions. J Neurophysiol *67*, 145-163.

Llinas, R., Ribary, U., Contreras, D., and Pedroarena, C. (1998). The neuronal basis for consciousness. Philos Trans R Soc Lond B Biol Sci *353*, 1841-1849.

Luebke, J.I., Weaver, C.M., Rocher, A.B., Rodriguez, A., Crimins, J.L., Dickstein, D.L., Wearne, S.L., and Hof, P.R. (2010). Dendritic vulnerability in neurodegenerative disease: insights from analyses of cortical pyramidal neurons in transgenic mouse models. Brain Struct Funct *214*, 181-199.

Manita, S., Miyakawa, H., Kitamura, K., and Murayama, M. (2017). Dendritic Spikes in Sensory Perception. Front Cell Neurosci *11*, 29.

Manita, S., Suzuki, T., Homma, C., Matsumoto, T., Odagawa, M., Yamada, K., Ota, K., Matsubara, C., Inutsuka, A., Sato, M*., et al.* (2015). A Top-Down Cortical Circuit for Accurate Sensory Perception. Neuron *86*, 1304-1316.

Mechawar, N., Cozzari, C., and Descarries, L. (2000). Cholinergic innervation in adult rat cerebral cortex: a quantitative immunocytochemical description. J Comp Neurol *428*, 305-318.

Naka, A., and Adesnik, H. (2016). Inhibitory Circuits in Cortical Layer 5. Front Neural Circuits *10*, 35.

Nakamura, S., Narumi, T., Tsutsui, K., and Iijima, T. (2009). Difference in the functional significance between the lemniscal and paralemniscal pathways in the perception of direction of single-whisker stimulation examined by muscimol microinjection. Neuroscience research *64*, 323-329.

Oberlaender, M., de Kock, C.P., Bruno, R.M., Ramirez, A., Meyer, H.S., Dercksen, V.J., Helmstaedter, M., and Sakmann, B. (2012). Cell type-specific three-dimensional structure of thalamocortical circuits in a column of rat vibrissal cortex. Cereb Cortex *22*, 2375-2391.

Pakan, J.M.P., Currie, S.P., Fischer, L., and Rochefort, N.L. (2018). The Impact of Visual Cues, Reward, and Motor Feedback on the Representation of Behaviorally Relevant Spatial Locations in Primary Visual Cortex. Cell Rep *24*, 2521-2528.

Palmer, L.M., Shai, A.S., Reeve, J.E., Anderson, H.L., Paulsen, O., and Larkum, M.E. (2014).
NMDA spikes enhance action potential generation during sensory input. Nat Neurosci *17*, 383-
390.
Pan, W.X., Schmidt, R., Wickens, J.R., and Hyland, B.I. (2005). Dopamine cells respond to
predicted events during classical conditioning: evidence for eligibility traces in the reward-
learning network. J Neurosci *25*, 6235-6242.
Park, J.M., Hong, Y.K., Rodgers, C.C., Dahan, J.B., Schmidt, E.R., and Bruno, R.M. (2020).
Deep and superficial layers of the primary somatosensory cortex are critical for whisker-based
texture discrimination in mice. bioRxiv, 2020.08.12.245381.
Parvizi, J., and Damasio, A. (2001). Consciousness and the brainstem. Cognition *79*, 135-160.
Petersen, C.C., and Crochet, S. (2013). Synaptic computation and sensory processing in
neocortical layer 2/3. Neuron *78*, 28-48.
Petreanu, L., Gutnisky, D.A., Huber, D., Xu, N.L., O'Connor, D.H., Tian, L., Looger, L., and
Svoboda, K. (2012). Activity in motor-sensory projections reveals distributed coding in
somatosensation. Nature *489*, 299-303.
Petty, G.H., Kinnischtzke, A.K., Hong, Y.K., and Bruno, R.M. (2020). Effects of arousal and
movement on secondary somatosensory and visual thalamus. bioRxiv, 2020.03.04.977348.
Pnevmatikakis, E.A., and Giovannucci, A. (2017). NoRMCorre: An online algorithm for
piecewise rigid motion correction of calcium imaging data. J Neurosci Methods *291*, 83-94.
Pnevmatikakis, E.A., Soudry, D., Gao, Y., Machado, T.A., Merel, J., Pfau, D., Reardon, T., Mu,
Y., Lacefield, C., Yang, W.*, et al.* (2016). Simultaneous Denoising, Deconvolution, and
Demixing of Calcium Imaging Data. Neuron *89*, 285-299.
Poort, J., Khan, A.G., Pachitariu, M., Nemri, A., Orsolic, I., Krupic, J., Bauza, M., Sahani, M.,
Keller, G.B., Mrsic-Flogel, T.D.*, et al.* (2015). Learning Enhances Sensory and Multiple Non-
sensory Representations in Primary Visual Cortex. Neuron *86*, 1478-1490.
Ramirez, A., Pnevmatikakis, E.A., Merel, J., Paninski, L., Miller, K.D., and Bruno, R.M. (2014).
Spatiotemporal receptive fields of barrel cortex revealed by reverse correlation of synaptic input.
Nat Neurosci *17*, 866-875.
Rigotti, M., Barak, O., Warden, M.R., Wang, X.J., Daw, N.D., Miller, E.K., and Fusi, S. (2013).
The importance of mixed selectivity in complex cognitive tasks. Nature *497*, 585-590.
Rodgers, C.C., Nogueira, R., Pil, B.C., Greeman, E.A., Park, J.M., Hong, Y.K., Fusi, S., and
Bruno, R.M. (2021). Sensorimotor strategies and neuronal representations for shape
discrimination. Neuron *109*, 2308-2325 e2310.
Roelfsema, P.R., and Holtmaat, A. (2018). Control of synaptic plasticity in deep cortical
networks. Nat Rev Neurosci *19*, 166-180.
Rubio-Garrido, P., Perez-de-Manzo, F., Porrero, C., Galazo, M.J., and Clasca, F. (2009).
Thalamic input to distal apical dendrites in neocortical layer 1 is massive and highly convergent.
Cereb Cortex *19*, 2380-2395.
Sandler, M., Shulman, Y., and Schiller, J. (2016). A Novel Form of Local Plasticity in Tuft
Dendrites of Neocortical Somatosensory Layer 5 Pyramidal Neurons. Neuron *90*, 1028-1042.
Schiller, J., Schiller, Y., Stuart, G., and Sakmann, B. (1997). Calcium action potentials restricted
to distal apical dendrites of rat neocortical pyramidal neurons. J Physiol *505 (Pt 3)*, 605-616.
Schwindt, P., and Crill, W. (1999). Mechanisms underlying burst and regular spiking evoked by
dendritic depolarization in layer 5 cortical pyramidal neurons. J Neurophysiol *81*, 1341-1354.
Sheffield, M.E., and Dombeck, D.A. (2015). Calcium transient prevalence across the dendritic
arbour predicts place field properties. Nature *517*, 200-204.
Stringer, C., Pachitariu, M., Steinmetz, N., Carandini, M., and Harris, K.D. (2019). High-
dimensional geometry of population responses in visual cortex. Nature *571*, 361-365.
Takahashi, N., Ebner, C., Sigl-Glockner, J., Moberg, S., Nierwetberg, S., and Larkum, M.E.
(2020). Active dendritic currents gate descending cortical outputs in perception. Nat Neurosci.

Takahashi, N., Oertner, T.G., Hegemann, P., and Larkum, M.E. (2016). Active cortical dendrites
modulate perception. Science *354*, 1587-1590.
Thiele, A., and Bellgrove, M.A. (2018). Neuromodulation of Attention. Neuron *97*, 769-785.
Tsai, J., Grutzendler, J., Duff, K., and Gan, W.B. (2004). Fibrillar amyloid deposition leads to
local synaptic abnormalities and breakage of neuronal branches. Nat Neurosci *7*, 1181-1183.
Voleti, V., Patel, K.B., Li, W., Perez Campos, C., Bharadwaj, S., Yu, H., Ford, C., Casper, M.J.,
Yan, R.W., Liang, W., *et al.* (2019). Real-time volumetric microscopy of in vivo dynamics and
large-scale samples with SCAPE 2.0. Nat Methods *16*, 1054-1062.
Wang, P.Y., Boboila, C., Chin, M., Higashi-Howard, A., Shamash, P., Wu, Z., Stein, N.P.,
Abbott, L.F., and Axel, R. (2020). Transient and Persistent Representations of Odor Value in
Prefrontal Cortex. Neuron *108*, 209-224 e206.
Wimmer, V.C., Bruno, R.M., de Kock, C.P., Kuner, T., and Sakmann, B. (2010). Dimensions of a
Projection Column and Architecture of VPM and POm Axons in Rat Vibrissal Cortex. Cereb
Cortex *20*, 2265-2276.
Xu, N.L., Harnett, M.T., Williams, S.R., Huber, D., O'Connor, D.H., Svoboda, K., and Magee,
J.C. (2012). Nonlinear dendritic integration of sensory and motor input during an active sensing
task. Nature *492*, 247-251.
Yamashita, T., Vavladeli, A., Pala, A., Galan, K., Crochet, S., Petersen, S.S.A., and Petersen,
C.C.H. (2018). Diverse Long-Range Axonal Projections of Excitatory Layer 2/3 Neurons in
Mouse Barrel Cortex. Front Neuroanat *12*, 33.
Yao, H., and Dan, Y. (2001). Stimulus timing-dependent plasticity in cortical processing of
orientation. Neuron *32*, 315-323.
Yu, Y.S., Graff, M.M., Bresee, C.S., Man, Y.B., and Hartmann, M.J. (2016a). Whiskers aid
anemotaxis in rats. Sci Adv *2*, e1600716.
Yu, Y.S., Graff, M.M., and Hartmann, M.J. (2016b). Mechanical responses of rat vibrissae to
airflow. J Exp Biol *219*, 937-948.
Yuste, R., Gutnick, M.J., Saar, D., Delaney, K.R., and Tank, D.W. (1994). Ca2+ accumulations
in dendrites of neocortical pyramidal neurons: an apical band and evidence for two functional
compartments. Neuron *13*, 23-43.
Zhang, W., and Bruno, R.M. (2019). High-order thalamic inputs to primary somatosensory cortex
are stronger and longer lasting than cortical inputs. eLife *8*.
Zhang, Y., Cudmore, R.H., Lin, D.T., Linden, D.J., and Huganir, R.L. (2015). Visualization of
NMDA receptor-dependent AMPA receptor synaptic plasticity in vivo. Nat Neurosci *18*, 402-
407.
1212

Figure 1. Mice rapidly learn to discriminate stimulus direction in head-fixed paradigm.

(A) A water droplet is paired with air puffs in one direction (CS+) but not the other (CS-). Licking in anticipation of water is assessed in the response window just after CS+ or CS- and prior to water delivery for the CS+ (grey bar).

(B) Experimental timeline. 2-3 weeks after virus injection, naive tuft responses to stimuli are recorded (pre). The CS+ is then paired with water for 8-9 days (blue). On the last day, stimuli are presented without reward (post). In a separate group of mice, the same stimuli are presented over 9 days in the absence of reward (unrewarded group).

(C) Lick rasters for three different sessions in one example mouse. On session 9, the CS+ but not the CS- reliably elicits licks.

(D) Mean baseline-subtracted whisking amplitude aligned to the CS+ (red) and CS- (navy) across sessions 1, 2, and 9 of an example mouse.

(E) Learning curve demonstrates rapid learning. Mean probability of at least one lick in the response window across sessions.

(F) Behavioral performance of each mouse in the rewarded group (M1 – M7).

Figure 2. Overall tuft response to stimuli is unbiased and relatively stable across conditioning.

(A) Dendritic activity was recorded in layer 1 (i) in the C1/C2 barrel columns (ii). (i) Two-photon image ~60 μm deep relative to pia. Dashed yellow lines denote C1 and C2 boundaries from intrinsic imaging. Reconstruction from (Ramirez et al., 2014).

(ii) Tangential section through layer 4 showing barrels stained with streptavidin-Alexa 647 and GCaMP6f-expressing apical trunks. Red circles indicate location of 2-photon lesions to mark the imaging region for post-hoc analysis.

(B) Overlay of five segmented pseudo-colored tufts from imaging field in A(i).

(C) Time courses of calcium responses of example tufts in (B) to three air puffs (dashes).

(D) Amplitude for CS+ (red) and CS- responses (blue), computed for each segmented tuft in the first 1.5 s post-stimulus (grey points), do not differ within or across sessions. Colored lines indicate median.

(E) Same as in D, showing data for all conditioning sessions.

Figure 3. Reinforcement learning, but not stimulus exposure, enhances tuft selectivity for CS+ and CS- stimuli.

(A) Across the indicated sessions, individual tufts (circles) exhibit larger biases to CS+ or CS- (pooled across all conditioned mice).

(B) Repeated exposure to stimuli does not bias individual tufts to CS+ or CS-.

(C) Conditioning reshapes distribution of selectivity indices for tufts from Normal on pre-conditioning session to uniform on post-conditioning session.

(D) Distribution of tuft selectivity indices remains Normal throughout all repeated exposure sessions.

(E) Selectivity (median SI magnitude of tufts for each session) increases with behavioral performance of 6 animals.

(F) Neural discriminability (mean ± sem) of tufts, pooled across all animals on each session, increases with conditioning and decreases with repeated exposure.

Figure 4. High-speed volumetric imaging of apical tufts confirms the emergence of enhanced selectivity after learning.

(A) Top and side view of four example tufts segmented from volumetric SCAPE imaging.

(B) Time courses of calcium activity from example tufts in (A) during five presentations of air puff stimuli (dashes).

(C) Performance across all conditioning sessions of two mice that were imaged with SCAPE.

(D) Across the indicated sessions, individual SCAPE-imaged tufts (circles) exhibit larger biases to CS+ or CS-.

(E) Conditioning reshapes selectivity distribution from Normal to uniform.

Figure 5. Longitudinal tracking reveals that reward enhances the selectivity of both initially unresponsive and responsive tufts.

(A) Three example tufts that were longitudinally tracked across learning. Top row: An initially unresponsive tuft develops a robust response to the CS+ but not the CS- after learning. Middle row: A responsive but unselective tuft loses its robust CS+ response and becomes selective for the CS-. Bottom row: A CS- selective neuron becomes unresponsive to both stimuli.

(B) Tufts that were unresponsive during the first session were longitudinally tracked to the last session. Plotted is the mean proportion of selective and unselective neurons across all animals in the conditioned (black bars) and repeated exposure (grey bars) groups.

(C,D) Same analysis as C for initially selective (D) and unselective (E) tufts.

Two-sample t-test was used for comparisons between conditioned and repeated exposure groups. Paired t-test was used for comparisons within a group. * p < 0.05

(E) Total tuft counts from first to last session within the 3 response categories for either conditioned (left) or repeated exposure (right) groups.

(F) SI of responsive tufts on the last session that were initially unresponsive during the first session. Conditioned tufts have enhanced selectivity compared to repeated exposure.

(G) Tufts that were selective on the last session are more selective if conditioned (black) rather than undergoing repeated exposure (grey).

(H) Tufts that responded on both pre and post sessions tend to have higher selectivity if conditioned rather than undergoing repeated exposure.

(I) SI of responsive tufts on the first session that later became unresponsive during the last session.

Figure 6. Whisking is only weakly correlated with tuft activity and cannot account for changes in selectivity during learning.
(A) Whisking amplitude aligned to calcium activity of three example tufts in one session. Green shading indicates periods of whisking. Red and navy ticks indicate CS+ or CS- delivery, respectively.
(B) Mean whisking response of four mice to CS+ (red) and CS- (navy) does not change across sessions during learning (mean ± s.e.m.).
(C) Mean standard deviation of whisking decreases for both CS+ and CS- across learning, but CS+ and CS- do not differ.
(D) Event-triggered averages of 322 tufts on the post-conditioning day (grey traces - individual tufts, black inset - population average) are responsive to stimuli but relatively unmodulated by whisking.
(E) R2 values for linear models predicting calcium from stimuli (y axis) are consistently greater than those predicting calcium from whisking (x axis). Each circle represents a tuft.
(F) Magnitude of tuft selectivity does not correlate with mean whisking amplitude during CS+ (left) and CS- trials (right) on that session.

Figure 7. Behavioral responses do not account for enhancement of stimulus selectivity during learning.

(A) Mean stimulus responses of four tufts during hit (red), CR (cyan), and FA (black) trials. Top row: Example tufts whose responses are not behaviorally modulated (CR is similar to FA). Bottom row: Example tufts with behaviorally modulated responses (CR and FA differ).

(B) Selectivity index (SI) distribution changes from early (left) and late learning sessions (right) even when tufts with behaviorally modulated responses (CR≠FA) are excluded.

(C) Median SI magnitude of tufts in each of six animals (from panel B) increases from early to late learning sessions.

Figure 8. Apical tufts in barrel cortex of mice performing the task exclusively with their whiskers undergo long-lasting changes in selectivity.

(A) SI histograms of mice performing the task exclusively with their whiskers exhibit increased selectivity across pre-conditioning, last-rewarded, and post-conditioning sessions.

(B) Relative to pre-conditioning, mice using their whiskers and other sensory cues to perform the task have increased selectivity during the last rewarded session, but not the post-conditioning session.

(C) The probability of anticipatory licks in response to the CS+ extinguishes across post-conditioning blocks (of 20 trials each).

(D) Tuft selectively remains uniformly distributed during post-conditioning trial blocks 1-2 (top) while licking is extinguishing, and blocks 3-4 (bottom) in which licking is extinguished.

Supplementary Figure 1. CS+ trials evoke a second, long-latency peak during early learning, but not late learning.

(A) Left: Population average of stimulus-responsive tufts aligned to CS+ (red) or CS- (blue) trials from an example mouse. Right: Normalized ΔF/F of individual tufts during CS+ trials.

(B) Same as in A, combining data across four mice whose imaging regions were mapped with intrinsic imaging.

Supplementary Figure 2. Selectivity was enhanced in individual animals that received rewards.
Median SI magnitude for each animal across three sessions for conditioned (left) and repeated exposure groups (right). * p < 0.05, *** p < 10^{-3}

Supplementary Figure 3. Segmented tufts from two-photon and SCAPE microscopy.
(A,B) 12 example tufts extracted from either two-photon (A) or SCAPE microscopy (B). Tufts segmented from SCAPE microscopy are shown as maximum intensity projections from the top and side. Scale bars: 100μm.

Supplementary Figure 4. Calcium event rate of tufts that were either unresponsive or responsive to air puff stimuli.
The number of calcium events per minute was quantified for all tufts during each conditioning session. Data from each group was pooled across all sessions.

Supplementary Figure 5. Licking cannot account for changes in selectivity during learning.
(A) ITI lick-bout-triggered averages of 232 tufts on the 5th conditioning day, when ITI licks were still common (grey traces - individual tufts, black inset - population average), exhibit little or no lick-related calcium influx.
(B) R2 values for linear models predicting calcium from stimuli (y axis) are consistently greater than those predicting calcium from licking (x axis). Each circle represents one tuft out of 442 tufts on last-rewarded sessions.
(C) Coefficients from a multivariate regression analysis with calcium as the response variable and the CS+, CS-, whisking, and licking as the predictors. CS+ and CS- coefficients are therefore disentangled from correlations with whisking and licking. Conditioning biases individual tufts (circles) to have larger CS+ or CS- coefficients. n = 304, 324, and 322 tufts for First rewarded, Last rewarded, and Post conditioning, respectively.
(D) Similar analysis to C but for repeated exposure group, with calcium as the response variable and the CS+, CS-, and whisking as the predictors. n = 223, 208, and 218 tufts for Day 2, Final - 1, and Final session, respectively.