Atypical relationships between neurofunctional features of print-sound integration and reading abilities in Chinese children with dyslexia

Zhichao Xia [a, b †*,] Ting Yang [c, d †], Xin Cui [a, e], Fumiko Hoeft [e, f, g, h], Hong Liu [c, d, f],
Xianglin Zhang [a], Xiangping Liu [c, d *], Hua Shu [a *]

[a] State Key Laboratory of Cognitive Neuroscience and Learning &
IDG/McGovern Institute for Brain Research, Beijing Normal University, China

[b] School of Systems Science, Beijing Normal University, China

[c] Faculty of Psychology, Beijing Normal University, China

[d] Beijing Key Laboratory of Applied Experimental Psychology, National
Demonstration Center for Experimental Psychology Education, Faculty of
Psychology, Beijing Normal University, China

[e] Haskins Laboratories, USA

[f] Department of Psychological Sciences and Brain Imaging Research Center,
University of Connecticut, USA

[g] Department of Psychiatry and Weill Institute for Neurosciences, University of
California, San Francisco, USA

[h] Department of Neuropsychiatry, Keio University School of Medicine, Japan

[†] These authors have contributed equally to this work and share first authorship

[*] Correspondence:

Zhichao Xia, State Key Lab of Cognitive Neuroscience and Learning, Beijing
Normal University, China. Email: xiazc.psy@gmail.com;

Hua Shu, State Key Lab of Cognitive Neuroscience and Learning, Beijing
Normal University, China. Email: shuhua@bnu.edu.cn;

Xiangping Liu, Faculty of Psychology, Beijing Normal University, China. Email:
lxp599@163.com

Running Title: Print-sound integration in Chinese dyslexia

Abstract

Conquering grapheme-phoneme correspondence is necessary for developing fluent reading in alphabetic orthographies. In neuroimaging research, this ability is associated with brain activation differences between the audiovisual congruent against incongruent conditions, especially in the left superior temporal cortex. Studies have also shown such a neural audiovisual integration effect is reduced in individuals with dyslexia. However, existing evidence is almost restricted to alphabetic languages. Whether and how multisensory processing of print and sound is impaired in Chinese dyslexia remains underexplored. Of note, semantic information is deeply involved in Chinese character processing. In this study, we applied a functional magnetic resonance imaging audiovisual integration paradigm to investigate the possible dysfunctions in processing character-sound pairs and pinyin-sound pairs in Chinese dyslexic children compared with typically developing readers. Unexpectedly, no region displayed significant group difference in the audiovisual integration effect in either the character or pinyin experiment. However, the results revealed atypical correlations between neurofunctional features accompanying audiovisual integration with reading abilities in Chinese children with dyslexia. Specifically, while the audiovisual integration effect in the left inferior cortex in processing character-sound pairs correlated with silent reading comprehension proficiency in both dyslexia and control group, it was associated with morphological awareness in the control group but with rapid naming in dyslexics. As for pinyin-sound associations processing, while the stronger activation in the congruent than incongruent conditions in the left occipito-temporal cortex and bilateral superior temporal cortices was associated with better oral word reading in the control group, an opposite pattern was found in children with dyslexia. On the one hand, this pattern suggests Chinese dyslexic children have yet to develop an efficient grapho-semantic processing system as typically developing children do. On the other hand, it indicates dysfunctional recruitment of the regions that process pinyin-sound pairs in dyslexia, which may impede character learning.

Keywords

audiovisual integration, character, Chinese, dyslexia, individual differences, pinyin

Highlights

- Neural bases of print-sound integration in Chinese children with and without dyslexia are investigated.
- Dyslexia shows atypical relationships between neurofunctional features of audiovisual integration in the reading network and different aspects of reading.
- Chinese children with dyslexia are likely to use inefficient strategies to process characters and pinyin.

1 Introduction

Reading consists of multiple cognitive processes, and it takes years of formal instruction to achieve a high proficiency level. During this process, establishing a robust correspondence between orthographic code (e.g., grapheme) and phonologic representation (e.g., phoneme) is one initial and fundamental step (Perfetti & Harris, 2013). Behavioral and neuroimaging studies of alphabetic languages reveal that it is critical to conquer the grapheme-phoneme correspondence (GPC) rules to develop fluent reading skills. The failure will impede establishing efficient grapho-semantic mapping and eventually result in reading difficulties such as dyslexia (Blomert, 2011; Di Folco, Guez, Peyre, & Ramus, 2020; Richlan, 2019; Shaywitz, 1998). However, while the manifestation of dyslexia is associated with linguistic features in a given language (Richlan, 2020), whether and to what extent grapho-phonological processing is impaired in Chinese children with dyslexia remains underexplored, especially at the neurofunctional level.

Chinese has a morpheme-based logographic writing system (Perfetti, Cao, & Booth, 2013). Semantics is strongly involved in, even at the most fundamental processing level—character recognition, in addition to grapho-phonological mapping (Bi, Han, Weekes, & Shu, 2007; Guan, Fraundorf, & Perfetti, 2020; Liu et al., 2017; Yang, Shu, McCandliss, & Zevin, 2013; Zhao et al., 2014). At the behavioral level, longitudinal and meta-analytic studies have demonstrated the importance of both phonological-related (e.g., phonological awareness [PA], rapid naming [RAN]) and semantic-related (e.g., morphological awareness [MA]) skills in Chinese reading development and impairments (Pan et al., 2016; Lei et al., 2011; Liu et al., 2017; Ruan, Georgiou, Song, Li, & Shu, 2018). However, most tasks used in these studies require explicit processing of the written scripts. Hence, whether implicit and automatic processing is impaired in dyslexia remains largely unknown. A similar situation exists at the brain level. Previous functional magnetic resonance imaging (fMRI) studies revealed hypoactivation in the left inferior and middle frontal areas during visual rhythming and visual lexical decision in children with dyslexia, suggesting dysfunctions of the neural substrates underlying both print-to-sound and print-to-meaning mappings in tasks requiring explicit processing (Cao et al., 2016; Liu et al., 2012; Siok, Perfetti, Jin, & Tan, 2004). Paralleling the fMRI research, structural and diffusion imaging studies also provided evidence on alterations in these regions and white matter tracts connecting them (Siok, Niu, Jin, Perfetti, & Tan, 2008; Su et al., 2018; Xia, Hoeft, Zhang, & Shu, 2016). While these findings indicate deficits in grapho-semantic and grapho-phonological processing in Chinese children with dyslexia, evidence about the automaticity remains scarce.

In this study, we adopted the fMRI audiovisual paradigm, which is appropriate for investigating automaticity in reading-related processing. This paradigm has been used in shallow orthographies such as Dutch and demonstrated the impaired letter-sound automatized integration as a likely proximal cause of dyslexia that is independent of phonological processing deficits (Blau et al., 2010; Blau, van Atteveldt, Ekkebus, Goebel, & Blomert, 2009; Blau,

van Atteveldt, Formisano, Goebel, & Blomert, 2008). This paradigm's basic logic
is that if a brain area integrates auditory and visual inputs, the neural responses
should differ between congruent versus incongruent conditions. This effect is
usually named "congruency effect" when the activation in the congruent
condition was stronger than the incongruent condition and is named
"incongruency effect" otherwise. Here we used the term "audiovisual integration
effect" given that both directions indicate successful multimodal information
integration. Since it has been repeatedly observed when no task or a passive task
is used, researchers regard the condition differences to reflect implicit processing
(Blau et al., 2010; van Atteveldt, Formisano, Goebel, & Blomert, 2007). To date,
this audiovisual integration effect has been demonstrated in skilled adult
readers and typically developing children (Blau et al., 2010; van Atteveldt &
Ansari, 2014; van Atteveldt, Formisano, Goebel, & Blomert, 2004; van Atteveldt,
Formisano, Blomert, & Goebel, 2007).

Of importance, the direction and strength of this effect can be affected by
several factors, including characteristics of the participants and orthographic
depth (Blau et al., 2010; Blau et al., 2009; Holloway, van Atteveldt, Blomert, &
Ansari, 2015; Kronschnabel, Brem, Maurer, & Brandeis, 2014; Wang, Karipidis,
Pleisch, Fraga-Gonzalez, & Brem, 2020). Researchers found that children with
dyslexia showed atypical patterns in brain areas such as the superior temporal
cortex (STC) (Blau et al., 2010; Blau et al., 2009). It was mainly driven by the
hypoactivation in the congruent condition in dyslexia, indicating a reduced
neural integration, along with higher activation in the incongruent condition,
indicating a lack of suppression. In terms of orthography depth, recently, similar
paradigms have been applied in studies with Chinese adults (Xu, Kolozsvari,
Oostenveld, Leppanen, & Hamalainen, 2019) and typically developing children
(Xia et al., 2020). In particular, Xia et al. (2020) used Chinese characters (deep
orthography) and pinyin (a transparent alphabetic coding system that represents
the pronunciation of characters, which is taught at the earliest stage of Chinese
reading development and used as a scaffold in learning new characters) as
experimental materials and observed a significant audiovisual integration effect
(congruent < incongruent) in the left inferior frontal cortex (IFC) and bilateral
STC for characters. Moreover, this effect in the left IFC in response to character-
sound pairs and that in the left STC in response to pinyin-sound pairs were
associated with children's performance in silent reading comprehension that
relies on grapho-semantic mapping and oral word reading that relies on grapho-
phonological processing, respectively. In the current study, we aimed to examine
the impaired audiovisual integrations of character-sounds and pinyin-sounds
and their associations with different reading abilities in Chinese dyslexia with
the same design.

Notably, while the group comparison approach has been widely used to
identify neural deficits in dyslexia, approaches focusing on individual differences
also provide valuable insights. In this case, brain-behavior correlation is a useful
strategy (Jednorog et al., 2015; Pernet, Andersson, Paulesu, & Demonet, 2009).
Two primary patterns can be identified with it. The first is a universal
correlation between brain and behavioral performance regardless of group,

which indicates the neural system supports the cognitive processing in both
groups. For example, children's PA was correlated with the microstructural
feature of the left arcuate fasciculus, even after controlling reading status
(dyslexic; typical) (Su et al., 2018; Vandermosten et al., 2012). Alternatively,
there could be distinct ways in which reading abilities correlated with brain
measures between children with and without dyslexia, indicating either
dysfunction or compensation (Hoeft et al., 2011; Pernet et al., 2009; Rumsey et
al., 1999; Tschentscher, Ruisinger, Blank, Diaz, & von Kriegstein, 2018). For
example, while the typical readers relied more (higher regional cerebral blood
flow) on the left inferior parietal lobule (IPL), higher activation in this area was
associated with worse reading performance in dyslexia (Rumsey et al., 1999).
However, in previous studies, correlation analyses were commonly conducted
across the dyslexic and control groups. Since the subjects were selected on
purpose, group differences could drive the results (Blau et al., 2010). As
mentioned above, this issue can be addressed by controlling the effect of group in
the statistic model or directly comparing correlations between dyslexia and
typical readers.

To summarize, the main aim of the present study was to examine the
impairment of print-sound integration and related processes in Chinese dyslexia.
We asked two specific questions. First, whether the neurofunctional correlates of
print-sound integration differ between Chinese children with and without
dyslexia. Second, whether the relationships between such neural features and
reading abilities differ between the two groups. For this, we adopted an fMRI
audiovisual paradigm and used characters-sound pairs and pinyin-sound pairs
as materials. Both group comparison and individual differences analytic
approaches were performed. Based on the prior research in typically developing
children (Xia et al., 2020), we predicted that Chinese children with dyslexia
would show a reduced audiovisual integration effect. In addition, dyslexic
children may also display atypical brain-behavior correlations or recruit other
brain regions to integrate cross-modal information.

2 Methods

2.1 Participants and behavioral measures

Initially, one hundred children in grades 3-6 (45 dyslexia) from local elementary
schools attended the experiment. According to inclusion criteria, 23 dyslexic
children (10 girls; age 111-144 months, M [SD] = 122 [10]) were included in the
final analysis. Twenty-one children were excluded due to uncompleted data
collection (n = 9), head motion artifact (n = 9), and poor in-scanner performance
(n = 3). One child with a history of dyslexia diagnosis but performed well in the
character recognition task at the time of data collection was also excluded (this
child had received an intensive behavior intervention program before). The
control group consisted of 22 typically developing children (12 girls; age 118-140

months, *M* [*SD*] = 127 [6]) who were chosen to match the dyslexic group on
grade, age, and sex.

In this study, dyslexia was operationalized by the criteria of having
normal intelligence (⩾ 80 on the abbreviated version of the *Chinese Wechsler*
Intelligence Scale for Children (Wechsler, 1974)) but manifesting reading
difficulty (below -1 *SD* of the norm on a standardized reading screening task
Character Recognition (Xue, Shu, Li, Li, & Tian, 2013)). All the children in the
control group had normal intelligence (≥ 80) and scored above -0.5 *SD* of the
norm on the reading screening task (the aim was to increase the gap in reading
skills between the two groups). In addition, children in both groups were all
right-handed (Oldfield, 1971) native speakers of Chinese, with normal hearing
and normal or corrected-to-normal vision and were free from neurological or
psychiatric disorders. Finally, only the children that completed 4 task fMRI runs,
with an overall accuracy equal to or higher than 75% in the in-scanner passive
task and less than 25% time-points labeled as outliers (i.e., "bad volume") in each
run (data preprocessing section) were included.

Each child received a battery of behavioral tests on reading and cognitive-
linguistic skills individually in a silent room on the same day of the MRI session.
The reading measurements contained: (1) an untimed character naming task
(*Character Recognition*) to estimate the number of characters children had
conquered (Xue et al., 2013); (2) an oral word reading task (*Word List Reading*)
to measure how fast the participant accurately retrieved phonological
representation from visually presented high-frequency two-character words
(Zhang et al., 2012); and (3) a timed comprehension task (*Silent Reading*
Comprehension) to assess the proficiency of meaning access and semantic
judgment (Lei et al., 2011). In addition, PA, RAN, and MA—the three most
critical cognitive-linguistic skills in Chinese reading acquisition—were measured
by *Phoneme Deletion* (Li, Shu, McBride-Chang, Liu, & Peng, 2012), *Digit RAN*
(Liu et al., 2017), and *Morphological Production* (Shu, McBride-Chang, Wu, &
Liu, 2006).

Written informed consents were obtained from all the children and their
guardians after a detailed explanation of the objectives and procedure of the
study. After the experiment, each child received one book and a set of stationery
as compensation. This study was approved by the Institutional Review Board of
the State Key Laboratory of Cognitive Neuroscience and Learning at Beijing
Normal University, and data collection was conducted in 2018.

2.2 Experimental design

This study adopted the audiovisual paradigm that has been repeatedly used in
the previous studies investigating letter-sound integration (**Figure 1**). The
details about the stimuli and procedure can also be found in Xia et al. (2020). In
brief, 56 high-frequency (*M* ± *SD* = 929 ± 1486 per million; Chinese Single-
Character Word Database; https://blclab.org/pyscholinguistic-norms-database/)
pictographic characters that are frequently used as radicals in phonograms were

selected. These characters are visually simple (number of strokes: M = 4.34,
Range = 1-9) and learned early (age of acquire: M = 3 years, *Range* = 2-5 years).
The pinyin spellings of these characters were used as the visual stimuli in the
pinyin experiment. The auditory stimuli (duration: $M \pm SD$ = 476.3 ± 87.5 ms)
were the characters' sounds (i.e., syllables). A native Chinese male recorded the
audio files with a sampling rate of 44.1 kHz and 16-bit quantization, then
normalized to 85 dB and bandpass (100-4000 Hz) filtered with Audacity
(https://www.audacityteam.org/).

The study consisted of 4 task fMRI runs, with the first 2 for the pinyin
experiment and the last 2 for the character experiment. Using such a fixed order
was to prevent priming from characters on visually presented pinyin stimuli.
Two unimodal (auditory [Aud]; visual [Vis]) and two cross-modal conditions
(congruent [avC]; incongruent [avI]) were created for both the character and
pinyin experiments (**Figure 1**). In this study, we focused on the audiovisual
integration effect (i.e., the activation difference between the congruent condition
against the incongruent condition). A block design was used to present stimuli.
There were 8 task blocks (duration = 20.8 s; 2 for each condition) interleaved
with 9 rest blocks (duration = 20.8 s) in a single run. A task block contained 4
mini-blocks. A 1.5 s period was used to collect a whole-brain volume within each
mini-block, and a silent period of 3.7 s was used to present stimuli (see "image
acquisition" part). The stimuli were presented in white at the center of a black
background ("*KaiTi*" font, 96 pt for characters; "*Century Schoolbook*" font, 90 pt
for pinyin). A crosshair was presented at the center of the screen whenever there
was no stimulus. To help children keep their attention on the stimuli while
avoiding explicit congruency judgment, we used a target detection task (Blau et
al., 2010). Specifically, in each task block, two out of 16 experimental stimuli
were randomly replaced with the auditory target (440 Hz pure tone), visual (an
unpronounceable symbol) target or their combination. The participant was asked
to press a button with the right index finger as accurately and quickly as
possible whenever the target appeared.

Insert Figure 1 about here

2.3 Image acquisition

All brain images were collected at Beijing Normal University Imaging Center for
Brain Research using a 3-Tesla Siemens MAGNETOM Trio Tim scanner with a
12-channel head coil. The children first attended a training session to get
familiar with the experimental environment and the scanning noise. During the
formal scan, foam pads were used to hold their heads secure during scanning to
improve image quality. Between sequences, children were told to relax to reduce
the fatigue effect. For each participant, 2 functional runs for the pinyin
experiment, one run for anatomical images, and 2 functional runs for the
character experiment were administered sequentially. The quality of the brain

images was evaluated immediately by a radiologist who was blinded to the
details of this study.

The parameters of the functional images (Gradient Echo Planar Imaging
[EPI]) were: repetition time, 5.2 s; echo time, 32 ms; acquisition time, 1.5 s; flip
angle, 90 degrees; slice thickness, 4.5 mm; interscan gap, 0.675 mm, voxel size,
$3.0 \times 3.0 \times 4.5$ mm^3; 24 slices; 68 volumes). Since this study contained auditory
stimuli, to avoid artifacts induced by the noise during scanning, a sparse
sampling with 1.5 s for image collection and 3.7 s delay for stimuli presentation
was used (Shah et al., 2000). The parameters of the structural images
(Magnetization-Prepared Rapid Acquisition with Gradient Echo [MPRAGE])
were: repetition time, 5.2 s; echo time, 3.39 ms; inversion time, 1.1 s; flip angle, 7
degrees; slice thickness, 1.33 mm; interscan gap, 0 mm; voxel size, $1.3 \times 1.0 \times 1.3$
mm^3; 144 axial slices).

2.4 Data preprocessing

Functional data were analyzed with SPM12 (http://www.fil.ion.ucl). Since
dummy scans were added at the beginning of each run to avoid the T1
equilibration effect, no additional volumes were discarded during the
preprocessing. The sequence was then corrected for head motion. The "bad
volumes" were identified with ART-based outlier detection
(https://www.nitrc.org/projects/artifact_detect). Considering the age range of
participants, a liberal threshold (intensity > 9 *SD*; frame-to-frame head motion >
2 mm) was used. T1 images were segmented and used for transferring the fMRI
data from native space to the standard Montreal Neurological Institute (MNI)
space. The normalized images were smoothed with an 8-mm Full-Width Half-
Maximum Gaussian kernel, and the resulting data were used in the subsequent
model estimation. In the 1st level analysis, 4 experimental conditions, 7
parameters for head motion (3 for translations, 3 for rotations, and 1 for
framewise displacement), and the time point of each "bad volume" were included
in the model. The contrast map referring to the difference between avC against
avI conditions (positive value stands for higher activation in avC than avI;
negative value stands for lower activation in avC than avI) was calculated for
each child and used in the subsequent analyses.

2.5 Statistics

Reading and reading-related cognitive impairments in children with dyslexia
were examined at the behavioral level. We first performed descriptive statistics
and then compared children's performance in multiple behavioral tasks between
groups. Next, we calculated correlations between reading and cognitive-linguistic
skills within each group and compared between groups. For the variable that
was not normally distributed, the Spearman method was used.

As for in-scanner performance, a correct response was defined as the
button press to the target between 200 and 2000 ms. Only the correct trials were
used to calculate the average reaction time (RT). Finally, the effects of group and

run and their interaction were examined on accuracy (ACC) and RT with
analysis of variance (ANOVA).

We conducted brain analyses focusing on the differences between the two
cross-modal conditions (i.e., audiovisual integration effect) for character and
pinyin experiments, respectively, with the same analytic approaches. The
nuisance variables of age, sex, and performance IQ were controlled in all the
analyses. First, a voxel-wise whole-brain 2 (group: control vs. dyslexic) × 2
(condition: avC vs. avI) ANOVA (i.e., the group comparison approach) was
conducted to examine whether the brain regions showing the audiovisual
integration effect in children with dyslexia were the same as the control group.
Significant clusters were identified with the FWE-corrected threshold of p-
cluster < 0.05 (p-voxel < 0.001 for height). These clusters were then used as
regions-of-interest (ROIs), and post hoc t-tests were conducted to interrogate the
effects. Complementary ROI analyses were performed to examine correlations
between the magnitude of the neural integration effect with reading and
reading-related cognitive-linguistic skills in each group, to identify the reading-
related processes most involved in audiovisual integration of character/pinyin for
typical and dyslexic children, respectively. Once a significant correlation was
revealed, we further conducted a correlation coefficients comparison between
groups.

Next, we used two individual differences approaches to examine the
shared and differential brain-behavior correlations in children with and without
dyslexia in the character and pinyin experiments at the whole-brain level. First,
to identify the shared neural basis associated with behavioral performance
between groups, we conducted voxel-wise whole-brain regression analyses on the
contrast maps of avC against avI across all the participants while controlling for
reading status (i.e., dyslexic; control), age, sex, and performance IQ. Children's
performance in *Word List Reading* and *Silent Reading Comprehension* tasks
were used as regressors in separate models to examine the relationships between
neural processing features during print-sound integration with reading abilities
that rely more on grapho-phonological mapping and grapho-semantic mapping,
respectively. Second, we examined whether the relationships between the
neurofunctional features of multisensory processing and reading abilities differ
between the two groups. Same as the previous analysis, we used *Word List*
Reading and *Silent Reading Comprehension* scores as variates of interest in
separate models, along with the factor of group and the interaction. In each
analysis, an F-test was administered with the FWE-corrected threshold of p-
cluster < 0.05 (p-voxel < 0.001 for height), followed by ROI analysis to
interrogate the significant effects.

To visualize results, significant clusters were presented on a FreeSurfer
surface template with BrainNet Viewer (Xia, Wang, & He, 2013). Anatomical
labeling was performed by using the AAL atlas with DPABI
(http://rfmri.org/dpabi).

All the behavioral and ROI analyses were administered with SPSS (v24;
SPSS Inc., Chicago, IL, USA). Effects were considered significant at $p < 0.05$,
and $0.05 < p < 0.1$ was considered indicative of a trend.

3 Results

3.1 Behavior

3.1.1 Description of reading measures and group comparisons

Statistical metrics including *M*, *SD*, *Range*, and result of the Shapiro-Wilk test of
each behavioral measurement were presented in Table 1 (see Supplementary
Figure 1 for plots). The dyslexic group performed worse in all the reading and
reading-related cognitive-linguistic tasks than typically developing children (all
p's < 0.007). No significant differences were found between the two groups on age
or sex (both p's > 0.05). In addition, the IQs of all the children were within the
normal range, while typical readers had higher scores on both verbal and
performance sub-scores.

392   ---------------------------------------

393   Insert Table 1 about here

394   ---------------------------------------

3.1.2 Correlations between reading and reading-related cognitive skills

Same and differential correlations between reading and cognitive-linguistic skills
were revealed between children with and without dyslexia. In typical readers,
character recognition was significantly correlated with MA ($r = 0.589$, $p = 0.004$)
but not PA ($r = 0.117$, $p = 0.606$) or RAN ($r = -0.301$, $p = 0.174$). In dyslexics,
however, this ability was associated with PA ($r = 0.489$, $p = 0.018$), but not RAN
($r = -0.269$, $p = 0.214$) or MA ($r = 0.008$, p $= 0.970$). The group difference on the
correlation coefficients between character recognition and MA was significant (Z
$= 2.09$, $p = 0.037$). A similar pattern was found in reading comprehension
proficiency, where the scores were correlated with MA ($r = 0.456$, $p = 0.033$) in
controls but not dyslexia ($r = -0.012$, $p = 0.957$). On the contrast, oral reading
fluency was significantly correlated with RAN in both groups (controls: $r = -$
0.531, $p = 0.012$; dyslexics: $r = -0.578$, $p = 0.004$).

3.1.3 In-scanner performance

The aim of using a passive target detection task was to ensure the participant
focused their attention on the stimuli delivered via auditory and visual
modalities. The results revealed that children in both groups performed the task
with high ACC (controls: M [SD] = 96.7% [2.9]; dyslexics: M [SD] = 92.4% [5.0]).
In the ANOVA, the main effects of group were significant on both ACC ($p =$
0.001) and RT ($p = 0.012$). The post-hoc analyses showed that the children with

dyslexia had lower accuracy and used more time to complete the tasks than the
normal controls. The main effect of run was significant on RT ($p = 0.012$; faster
as the experiment proceeds) but not ACC ($p = 0.645$). No significant group × run
interaction was observed on either ACC or RT (both p's > 0.05).

419   -------------------------------------

420   Insert Table 2 about here

421   -------------------------------------

3.2 fMRI

We used group comparison (ANOVA), and individual differences (brain-behavior
correlation) approaches to investigate impaired neurofunctional features
accompanying print-sound integration in Chinese children with dyslexia. In this
section, we first present the character experiment results, followed by the pinyin
experiment.

3.2.1 Character: whole-brain ANOVA and ROI analysis

In the voxel-wise whole-brain 2 (group) × 2 (condition) ANOVA, we observed the
left IFC and STC showed a significant main effect of condition that survived the
FWE corrected p-cluster < 0.05 (p-voxel < 0.001 for height; **Table 2**; **Figure 2A**).
The follow-up analysis revealed a similar pattern of lower activation in the
congruent than incongruent conditions in both the control (LIFC: $t = 4.361$, $p <$
0.001; LSTC: $t = 3.646$, $p = 0.002$) and the dyslexic (LIFC: $t = 2.346$, $p = 0.028$;
LSTC: $t = 3.427$, $p = 0.002$; **Figure 2B**). No main effect of group or group ×
condition interaction survived the whole-brain FWE correction.

In the complementary ROI analyses, we observed similar correlations in
the two groups between the audiovisual integration effect and reading
comprehension proficiency in the left IFC (controls: $r = 0.571$, $p = 0.011$;
dyslexics: $r = 0.391$, $p = 0.088$, marginally significant; **Figure 2C**). Interestingly,
distinct contributions of the cross-modal conditions within each group were
observed. To be specific, the correlation was driven by the incongruent condition
in typical readers (avC: $r = 0.222$, $p = 0.360$; avI: $r = -0.526$, $p = 0.021$) but was
related to the congruent condition in dyslexia (avC: $r = 0.400$, $p = 0.080$,
marginally significant; avI: $r = -0.185$, $p = 0.436$). Group difference on the
correlation coefficients between reading comprehension proficient and brain
activation in the incongruent condition was significant ($Z = 2.41$, $p = 0.016$).
Correlations between the audiovisual integration effect and reading-related
cognitive skills further support the idea that different mechanisms underlie the
integration-comprehension relationships in the two groups (**Figure 2C**): the
audiovisual integration effect in this region was correlated with MA in the
controls ($r = 0.551$, $p = 0.014$) but not in the dyslexics ($r = -0.025$, $p = 0.915$).
Group difference on the correlation coefficients was significant ($Z = 2.01$, $p =$
0.044). On the other hand, the effect was correlated with RAN in the dyslexics (r
$= -0.509$, $p = 0.022$) but not in the controls ($r = -0.338$, $p = 0.157$).

3.2.2 Character: whole-brain group × reading interaction

Regarding the individual differences approach, no significant clusters survived
the FWE-corrected threshold of p-cluster < 0.05 (p-voxel < 0.001 for height) in
the analysis investigating relationships between the neural audiovisual
integration effect and reading abilities across or that explored the difference
between groups.

462   ------------------------------------

463   Insert Figure 2 about here

464   ------------------------------------

3.2.3 Pinyin: whole-brain group comparison

The same analytic approach was used in the pinyin experiment. In the voxel-
wise whole-brain 2 (group) × 2 (condition) ANOVA, no regions showed significant
main effect or interaction at the FWE corrected threshold of p-cluster < 0.05 (p-
voxel < 0.001 for height).

3.2.4 Pinyin: whole-brain group × reading interaction and ROI analysis

Then, we investigated the neural deficits with the individual differences
approaches in a whole-brain fashion. While no region displayed significant brain-
behavior correlation similar across groups, clusters located in the left
occipitotemporal cortex (OTC) and bilateral STC showed significant group
differences in the correlation between the audiovisual integration effect and oral
reading fluency (FWE corrected p-cluster < 0.05, p-voxel < 0.001 for height;
Table 2; **Figure 3A**). The subsequent analyses revealed positive brain-reading
correlations in typical readers and negative correlations in children with
dyslexia (**Figure 3B, C and D; Table 3**). Furthermore, the correlations in the
left OTC (avC: $r = 0.685$, $p = 0.001$; avI: $r = 0.118$, $p = 0.631$) and STC (avC: $r =$
0.588, $p = 0.008$; avI: $r = -0.014$, $p = 0.956$) were driven by the congruent
condition in the control group, while fluent oral reading was not correlated with
brain activation of the right STC in either congruent ($r = 0.240$, $p = 0.321$) or
incongruent condition ($r = -0.259$, $p = 0.284$). In contrast, in dyslexia, the
correlations were driven by the avI condition (left OTC: $r = 0.674$; $p = 0.001$; left
STC: $r = 0.445$, $p = 0.049$; right STC: $r = 0.543$, $p = 0.013$) but not the avC
condition (left OTC: $r = 0.026$, $p = 0.913$; left STC: $r = -0.050$, $p = 0.833$; right
STC: $r = -0.181$, $p = 0.444$) in all three regions. Group differences on the
correlations were significant between fluent oral reading and activation in the
congruent condition in the left OTC ($Z = 2.54$, $p = 0.011$) and left STC ($Z = 2.26$,
$p = 0.024$), and between fluent oral reading and activation in the in incongruent
condition in the left OTC ($Z = -2.18$, $p = 0.029$) and right STC ($Z = -2.73$, $p =$
0.006).

Regarding correlations between neural audiovisual integration effect
(avC-avI) with reading-related cognitive-linguistic skills, the left STC was

negatively associated with RAN in controls ($r = -0.479$, $p = 0.038$) and showed a
trend positively correlated with RAN in dyslexics ($r = 0.394$, $p = 0.085$,
marginally significant). Group difference was significant ($Z = -2.93$, $p = 0.003$).

Insert Figure 3 about here

Insert Table 3 about here

----

4 Discussion

This study investigated the neurofunctional features of implicitly integrating
print and sound and their relationships with reading abilities in Chinese
children with and without dyslexia. We adopted an fMRI audiovisual paradigm
with a passive target detection task. Character-sounds and pinyin-sounds, whose
orthographic depths are dramatically different, were used as materials. Of
importance, due to the morpho-syllabic nature of characters, semantic
information can be automatically activated during character recognition, at least
in typical readers. That is, this study enabled us to tap into the three-way
relationship between orthography, phonology, and semantics in normal and
impaired readers. Although no group differences on the audiovisual integration
effect were found, the results revealed strikingly atypical correlations between
neural integration of both character-sounds and pinyin-sounds with different
aspects of reading ability in Chinese children with dyslexia. On the one hand,
these findings indicate that children with dyslexia rely on articulatory
phonological information during the implicit character processing, which
indicates a less developed automatic grapho-semantic mapping and integration.
On the other hand, it suggests a malfunctional grapho-phonological mapping in
dyslexia and implies that these children may have difficulty in developing a
typical pinyin processing strategy and transferring it to learning characters as
their typically developing peers do.

4.1 Left IFC: inefficient grapho-semantic mapping in Chinese dyslexia

First, this study indicates insufficient semantic information access from visual
input in Chinese children with dyslexia. At the behavioral level, we observed
significantly less proficiency in the dyslexic group on silent reading
comprehension. Moreover, while both character recognition accuracy and reading
comprehension proficiency were associated with MA in typical readers, children
with dyslexia showed significant correlations between reading abilities with

other cognitive-linguistic skills instead of MA. Specifically, character recognition
was correlated with PA in dyslexia, in line with a previous study in an
independent and larger sample (Song, Zhang, Shu, Su, & McBride, 2020).
Regarding silent reading comprehension, although no correlation was
significant, there was a trend with RAN, a phonological processing skill that
consists of rapid and accurate phonological representation access, retrieval, and
articulatory operations. This result is in line with recent studies in which
phonological skill contributes to reading more at the early stages of reading
acquisition, and the contribution of morphological processing skill increases as
children grow (Liu et al., 2017). In addition to the direct effect, MA also mediates
the effect of PA on reading (Pan et al., 2016).

Furthermore, we used the fMRI audiovisual paradigm to probe the neural
bases of processing character-sound associations in Chinese children with and
without dyslexia. In terms of brain activation in the left IFC and STC, dyslexic
children seem to integrate information from auditory and visual modalities the
same way as controls, reflected as a strong audiovisual integration effect
(congruent < incongruent) in both groups. Moreover, the effect in the left IFC
was associated with reading comprehension proficiency, regardless of reading
status. Of note, group differences were uncovered in the subsequent ROI
analysis. First, while the integration-comprehension association was driven more
by the incongruent condition in typical readers, the correlation was driven more
by the congruent condition in children with dyslexia, suggesting dyslexic and
typical children may use different strategies in processing characters and
corresponding sounds, and this difference enlarges with reading abilities
increasing within each group. Second, while the audiovisual integration effect in
the left IFC was associated with MA in the control group, it was correlated with
RAN in dyslexia. These findings indicate that articulatory phonological
processing is more likely involved in implicit processing of character-sound pairs
in children with dyslexia. In previous studies investigating neural impairment in
Chinese dyslexia, both hypo- and hyper-activation of the left frontal areas have
been reported (Cao et al., 2016; Liu et al., 2012; Siok et al., 2004). In a meta-
analysis, different parts of the left IFC were distinguished based on
functionality, where individuals with dyslexia displayed reduced activation in
the ventral part associated with semantic processing but increased activation in
the dorsal part that was associated with articulatory processing, presumably
compensating for their less efficient grapho-semantic route (Hancock, Richlan, &
Hoeft, 2017; Richlan, Kronbichler, & Wimmer, 2011). Given that the frontal
region is multifunctional (Fedorenko & Blank, 2020; Hagoort, 2014), dyslexia
may recruit it in reading-related processing in a different way compared with
typical readers.

Of note, our in-scanner task did not require any sound-semantic or print-
semantic processing. However, Chinese has a morpheme-based logographic
writing system, which involves semantic information in character recognition
(Guan et al., 2020; Liu et al., 2017; Yang et al., 2013; Zhao et al., 2014).
Therefore, phonological and semantic information could both be accessed
effortlessly, at least in typically developing children after 4-5 years of formal

instruction. Thus, it is also reasonable to predict that semantic processing skill
plays an equal or even more critical role in reading development than
phonological processing skill, and its impairments will result in reading
difficulties. In line with this hypothesis, previous studies of Chinese
demonstrated morphological awareness uniquely predicted reading outcomes
and dyslexia status (Pan et al., 2016; Ruan et al., 2018; Song et al., 2020). The
current results also showed that while the left IFC was strongly involved in both
groups, it was more associated with articulatory phonological processing in
dyslexia and semantic processing in typical children. The region is close to the
one found to underlie morphological processing and show hypo-activation in
children with dyslexia during tasks requiring explicit semantic processing (Liu
et al., 2013; Zou, Packard, Xia, Liu, & Shu, 2015).

These findings suggest that Chinese children with dyslexia have yet to
develop a well-functioning brain system for automated semantic access and
integration during implicit character/word recognition. In contrast, they are
more likely to reply on an articulatory strategy by recruiting the multifunctional
frontal area.

4.2 Bilateral TPC and left OTC: malfunctioning grapho-phonological
mapping in Chinese dyslexia

In addition to inefficient grapho-semantic mapping, this study also indicates that
children with dyslexia may neither develop a typical grapho-phonological
mapping. At the behavior level, the dyslexic group performed worse in the tasks
measuring oral reading fluency, PA, and RAN, in line with previous studies in
larger samples of Chinese children (e.g., Lei et al., 2011). In the pinyin fMRI
experiment, we observed differential brain-reading correlations in the classic
reading-related areas. These regions have also been associated with oral reading
fluency in terms of cortical thickness (Xia et al., 2018). In particular, the
magnitude of the audiovisual integration effect in response to pinyin-sound pairs
in the left OTC and STC was positively associated with oral word reading
fluency in typical controls: the better the children performed in the oral reading
task, the higher the activation was in the congruent condition than incongruent
condition. In contrast, the direction of the brain-reading correlation was reversed
in dyslexia: children with higher reading fluency showed higher activation in the
incongruent condition than congruent condition. Additionally, the correlations in
the control group were driven more by individual differences in brain responses
to the congruent condition. In contrast, the correlations in the dyslexic group
were driven more by the incongruent condition. These findings suggest that
while the same regions were recruited for implicit audiovisual integration of
pinyin, children with dyslexia may use different strategies.

The left OTC and STC have been regarded as critical nodes in the classic
reading network. Deficits in these areas have also been reported in dyslexia
(Richlan, Kronbichler, & Wimmer, 2009; Richlan et al., 2011). On the one hand,
the left OTC has been considered the interface for initially integrating
orthographic, phonological, and semantic information (Price & Devlin, 2011).
The left OTC contains a specific portion in the fusiform gyrus named the visual

word form area that has been found to respond specifically to word and word-like
stimuli. The left STC, on the other hand, is a central area that presents
phonological information (Boets et al., 2013; Glezer et al., 2016), including lexical
tone—the supramarginal phoneme in tonal languages such as Chinese (Si, Zhou,
& Hong, 2017; L. Zhang et al., 2011). It is functionally and structurally
connected to the left OTC, which can be shaped by learning grapho-phonological
mappings (Stevens, Kravitz, Peng, Tessler, & Martin, 2017; Thiebaut de
Schotten, Cohen, Amemiya, Braga, & Dehaene, 2014). In the current study,
besides the left-hemispheric regions, the right STC also showed significant group
differences in brain-behavior relationships. While this region has been less
frequently reported in previous studies, it is a substrate for supporting lexical
tone processing (Liang & Du, 2018; Si et al., 2017; Zhang et al., 2011). In
addition, the cortical thickness of this area is also associated with oral reading in
typically developing Chinese children (Xia et al., 2018).

The differential relationship between the audiovisual integration effect
during pinyin-sound processing and oral word reading fluency in dyslexia can be
interpreted in at least two ways. First, if pinyin processing skill is a continuum
and dyslexia presents the lower end, the current finding then hints at the
expansion and renormalization hypothesis of brain plasticity associated with
skill learning (Wenger, Brozzoli, Lindenberger, & Lövdén, 2017). That is, the
growth curve of print-sound integration is an inverse U-shape. When the child
starts learning pinyin at the earliest stages, the brain response to incongruent
audiovisual pairs is lower than congruent pairs. With learning, mismatching in
information extracted from different modalities induces higher activation during
integration. Children turn to focus on the overlearned visual scripts by efficiently
suppressing attractive auditory information at the highly familiar stage. In this
case, activation in the incongruent condition will be suppressed and lower than
that in the congruent condition. This interpretation is in line with our findings
that brain activation in the congruent condition was associated with oral reading
in typical readers, whereas incongruent condition was related to oral reading in
dyslexia. The alternative explanation is also associated with development but
assumes that individuals with dyslexia process pinyin differently compared with
typical controls. In general, typical readers shift from assembled to addressed
phonology with reading experience increases (Mei et al., 2014). Pinyin is
assembled in nature. But since there are only approximately 400 syllables, it can
be expected that typical readers in upper elementary grades who are highly
familiar with it could achieve the addressed phonology. Given that children learn
Chinese characters as holistic syllable-level units, children who read pinyin with
the same addressed phonology may benefit more. In this case, the differential
brain-reading correlations probably reflect the inefficient assembled phonology
adopted by dyslexic children in processing pinyin-sound pairs. However,
although this explanation is appealing, conclusions cannot be made with the
current evidence. To date, evidence on the developmental trajectory of pinyin
reading is still lacking. More studies on preliterate and emerging readers with a
longitudinal design are needed.

Nonetheless, these findings indicate impaired automatic grapho-
phonological mapping in dyslexia from the perspective of individual differences.
This anomaly could be underpinned by the altered recruitment of cortical areas
such as the left OTC and bilateral STC. Such deficits may further impede the
development of the ventral pathway for rapid character/word recognition and
result in reading problems. Recruiting preliterate children and conducting
longitudinal neuroimaging research is necessary to further answer this "cause
vs. consequence" question (Nash et al., 2017).

4.3 Limitations and future directions

There are several limitations in this study, and caution should be taken when
interpreting the results. First, we administered the pinyin experiment ahead of
the character experiment to reduce the prime effect of characters on processing
visually presented pinyin stimuli. This may influence brain activation to speech
sounds in character runs because the same auditory stimuli were used. Second,
to have sufficient statistical power, we used the liberal criteria to assess imaging
data quality and exclude participants with poor quality data accordingly. Third,
while the overall pattern indicates that children in both groups maintained their
attention throughout the experiment, the dyslexia group performed worse. To
deal with this issue, we controlled performance IQ in all the analyses. This could
be why no group differences were detected but demonstrated that the main
findings of brain-behavior relationships are robust. At last, we did not have
enough cases for looking into different subtypes of dyslexia. Studies with larger
sample sizes are needed, and a counterbalance design for estimating the order
effect, much stricter criteria for controlling MRI data quality, well-matching on
in-scanner performance between groups, and dividing dyslexia into subtypes can
be applied.

4.4 Conclusion

The present study examined impaired audiovisual integration of character-sound
associations and pinyin-sound associations in Chinese children with dyslexia at
the neurofunctional level. The results revealed that dyslexia manifested deficits
in grapho-semantic mapping, with the left IFC as the most relevant area, and
the grapho-phonological mapping, with the left OTC and bilateral STC as central
regions. Moreover, the current findings also imply that Chinese children with
dyslexia may process pinyin—the alphabetic coding system representing the
pronunciation of characters—in a lagged or deviated way, which impedes the
development of the direct route for rapid character/word recognition and
semantic access.

5 Author Contributions

709 Conceptualization: ZX, HS, XL; Investigation: ZX, TY, XC, HL, XZ; Formal
710 Analysis: ZX; Data Curation: ZX; Writing – Original Draft Preparation: ZX;

711 Writing – Review & Editing: ZX, TY, XC, FH, HL, XZ, HS, XL; Funding
712 Acquisition: ZX, HS, XL; Supervision: ZX, HS, XL

6 Acknowledgment

715 The authors thank all the participating children, their families, and all the
716 examiners.

7 Funding

This work was supported by the Key Program of the National Social Science
Foundation of China (14ZDB157), National Key Basic Research Program of
China (grant number 2014CB846103), National Natural Science Foundation of
China grants (grant number 31271082, 31671126, 31611130107, 61374165,
81801782), Beijing Municipal Science & Technology Commission (grant number
Z151100003915122), Fundamental Research Fund for the Central Universities,
China Postdoctoral Science Foundation (grant number 2019T120062,
2018M641235), and Social Science Fund of Beijing (grant number 17YYA004).

8 Declaration of interest

729 The authors declare that there is no conflict of interest.

9 Data availability statement

732 The data that support the findings of this study are available from the
733 corresponding authors upon reasonable request.

10 References

Bi, Y., Han, Z., Weekes, B., & Shu, H. (2007). The interaction between semantic
and the nonsemantic systems in reading: Evidence from Chinese.
Neuropsychologia, 45(12), 2660-2673.

Blau, V., Reithler, J., van Atteveldt, N., Seitz, J., Gerretsen, P., Goebel, R., et al.
(2010). Deviant processing of letters and speech sounds as proximate
cause of reading failure: a functional magnetic resonance imaging study of
dyslexic children. *Brain, 133*(Pt 3), 868-879.

Blau, V., van Atteveldt, N., Ekkebus, M., Goebel, R., & Blomert, L. (2009). Reduced neural integration of letters and speech sounds links phonological and reading deficits in adult dyslexia. *Curr Biol, 19*(6), 503-508.

Blau, V., van Atteveldt, N., Formisano, E., Goebel, R., & Blomert, L. (2008). Task-irrelevant visual letters interact with the processing of speech sounds in heteromodal and unimodal cortex. *Eur J Neurosci, 28*(3), 500-509.

Blomert, L. (2011). The neural signature of orthographic-phonological binding in successful and failing reading development. *Neuroimage, 57*(3), 695-703.

Boets, B., Op de Beeck, H. P., Vandermosten, M., Scott, S. K., Gillebert, C. R., Mantini, D., et al. (2013). Intact but less accessible phonetic representations in adults with dyslexia. *Science, 342*(6163), 1251-1254.

Cao, F., Yan, X., Wang, Z., Liu, Y., Wang, J., Spray, G. J., et al. (2016). Neural signatures of phonological deficits in Chinese developmental dyslexia. *Neuroimage.*

Di Folco, C., Guez, A., Peyre, H., & Ramus, F. (2020). Epidemiology of developmental dyslexia: A comparison of DSM-5 and ICD-11 criteria. *medRxiv*, 2020.2012.2018.20248189.

Fedorenko, E., & Blank, I. A. (2020). Broca's Area Is Not a Natural Kind. *Trends Cogn Sci, 24*(4), 270-284.

Glezer, L. S., Eden, G., Jiang, X., Luetje, M., Napoliello, E., Kim, J., et al. (2016). Uncovering phonological and orthographic selectivity across the reading network using fMRI-RA. *NeuroImage, 138*, 248-256.

Guan, C. Q., Fraundorf, S. H., & Perfetti, C. A. (2020). Character and child factors contribute to character recognition development among good and poor Chinese readers from grade 1 to 6. *Annals of Dyslexia.*

Hagoort, P. (2014). Nodes and networks in the neural architecture for language: Broca's region and beyond. *Curr Opin Neurobiol, 28*, 136-141.

Hancock, R., Richlan, F., & Hoeft, F. (2017). Possible roles for fronto-striatal circuits in reading disorder. *Neurosci Biobehav Rev, 72*, 243-260.

Hoeft, F., McCandliss, B. D., Black, J. M., Gantman, A., Zakerani, N., Hulme, C., et al. (2011). Neural systems predicting long-term outcome in dyslexia. *Proc Natl Acad Sci U S A, 108*(1), 361-366.

Holloway, I. D., van Atteveldt, N., Blomert, L., & Ansari, D. (2015). Orthographic Dependency in the Neural Correlates of Reading: Evidence from Audiovisual Integration in English Readers. *Cerebral Cortex, 25*(6), 1544-1553.

Jednorog, K., Marchewka, A., Altarelli, I., Monzalvo Lopez, A. K., van Ermingen-
Marbach, M., Grande, M., et al. (2015). How reliable are gray matter
disruptions in specific reading disability across multiple countries and
languages? Insights from a large-scale voxel-based morphometry study.
Hum Brain Mapp, 36(5), 1741-1754.

Jinger, P., Shuang, S., Mengmeng, S., Catherine, M., Hongyun, L., Yuping, Z., et
al. (2016). On the relationship between phonological awareness,
morphological awareness and Chinese literacy skills: evidence from an 8-
year longitudinal study. *Developmental Science, 19*(6), 982-991.

Kronschnabel, J., Brem, S., Maurer, U., & Brandeis, D. (2014). The level of
audiovisual print-speech integration deficits in dyslexia.
Neuropsychologia, 62, 245-261.

Lei, L., Pan, J., Liu, H., McBride-Chang, C., Li, H., Zhang, Y., et al. (2011).
Developmental trajectories of reading development and impairment from
ages 3 to 8 years in Chinese children. *J Child Psychol Psychiatry, 52*(2),
212-220.

Li, H., Shu, H., McBride-Chang, C., Liu, H., & Peng, H. (2012). Chinese
children's character recognition: Visuo-orthographic, phonological
processing and morphological skills. *Journal of Research in Reading,*
35(3), 287-307.

Liang, B., & Du, Y. (2018). The Functional Neuroanatomy of Lexical Tone
Perception: An Activation Likelihood Estimation Meta-Analysis.
Systematic Review. *Frontiers in neuroscience. 12*(495).

Liu, L., Tao, R., Wang, W., You, W., Peng, D., & Booth, J. R. (2013). Chinese
dyslexics show neural differences in morphological processing.
Developmental Cognitive Neuroscience, 6, 40-50.

Liu, L., Wang, W., You, W., Li, Y., Awati, N., Zhao, X., et al. (2012). Similar
alterations in brain function for phonological and semantic processing to
visual characters in Chinese dyslexia. *Neuropsychologia, 50*(9), 2224-2232.

Liu, Y., Georgiou, G. K., Zhang, Y., Li, H., Liu, H., Song, S., et al. (2017).
Contribution of cognitive and linguistic skills to word-reading accuracy
and fluency in Chinese. *International Journal of Educational Research,*
82, 75-90.

Mei, L., Xue, G., Lu, Z.-L., He, Q., Zhang, M., Wei, M., et al. (2014). Artificial
Language Training Reveals the Neural Substrates Underlying Addressed
and Assembled Phonologies. *PLOS ONE, 9*(3), e93548.

Nash, H. M., Gooch, D., Hulme, C., Mahajan, Y., McArthur, G., Steinmetzger, K.,
et al. (2017). Are the literacy difficulties that characterize developmental

dyslexia associated with a failure to integrate letters and speech sounds? *Developmental science, 20*(4).

Oldfield, R. C. (1971). The assessment and analysis of handedness: the Edinburgh inventory. *Neuropsychologia, 9*(1), 97-113.

Perfetti, C., Cao, F., & Booth, J. (2013). Specialization and Universals in the Development of Reading Skill: How Chinese Research Informs a Universal Science of Reading. *Sci Stud Read, 17*(1), 5-21.

Perfetti, C. A., & Harris, L. N. (2013). Universal reading processes are modulated by language and writing system. *Language Learning and Development, 9*(4), 296-316.

Pernet, C., Andersson, J., Paulesu, E., & Demonet, J. F. (2009). When all hypotheses are right: a multifocal account of dyslexia. *Hum Brain Mapp, 30*(7), 2278-2292.

Price, C. J., & Devlin, J. T. (2011). The interactive account of ventral occipitotemporal contributions to reading. *Trends Cogn Sci, 15*(6), 246-253.

Richlan, F. (2019). The Functional Neuroanatomy of Letter-Speech Sound Integration and Its Relation to Brain Abnormalities in Developmental Dyslexia. [Mini Review]. *Frontiers in Human Neuroscience, 13*(21).

Richlan, F. (2020). The Functional Neuroanatomy of Developmental Dyslexia Across Languages and Writing Systems. *Front Psychol, 11*(155), 155.

Richlan, F., Kronbichler, M., & Wimmer, H. (2009). Functional abnormalities in the dyslexic brain: a quantitative meta-analysis of neuroimaging studies. *Hum Brain Mapp, 30*(10), 3299-3308.

Richlan, F., Kronbichler, M., & Wimmer, H. (2011). Meta-analyzing brain dysfunctions in dyslexic children and adults. *Neuroimage, 56*(3), 1735-1742.

Ruan, Y., Georgiou, G. K., Song, S., Li, Y., & Shu, H. (2018). Does writing system influence the associations between phonological awareness, morphological awareness, and reading? A meta-analysis. *Journal of Educational Psychology, 110*(2), 180.

Rumsey, J. M., Horwitz, B., Donohue, B. C., Nace, K. L., Maisog, J. M., & Andreason, P. (1999). A Functional Lesion in Developmental Dyslexia: Left Angular Gyral Blood Flow Predicts Severity. *Brain and Language, 70*(2), 187-204.

Shah, N. J., Steinhoff, S., Mirzazade, S., Zafiris, O., Grosse-Ruyken, M. L., Jancke, L., et al. (2000). The effect of sequence repeat time on auditory

cortex stimulation during phonetic discrimination. *Neuroimage, 12*(1),
100-108.

Shaywitz, S. E. (1998). Dyslexia. *N Engl J Med, 338*(5), 307-312.

Shu, H., McBride-Chang, C., Wu, S., & Liu, H. Y. (2006). Understanding Chinese
developmental dyslexia: Morphological awareness as a core cognitive
construct. *Journal of Educational Psychology, 98*(1), 122-133.

Si, X., Zhou, W., & Hong, B. (2017). Cooperative cortical network for categorical
processing of Chinese lexical tone. *Proc Natl Acad Sci U S A, 114*(46),
12303-12308.

Siok, W. T., Niu, Z., Jin, Z., Perfetti, C. A., & Tan, L. H. (2008). A structural-
functional basis for dyslexia in the cortex of Chinese readers. *Proc Natl
Acad Sci U S A, 105*(14), 5561-5566.

Siok, W. T., Perfetti, C. A., Jin, Z., & Tan, L. H. (2004). Biological abnormality of
impaired reading is constrained by culture. *Nature, 431*(7004), 71-76.

Song, S., Zhang, Y., Shu, H., Su, M., & McBride, C. (2020). Universal and
Specific Predictors of Chinese Children With Dyslexia – Exploring the
Cognitive Deficits and Subtypes. *Frontiers in Psychology, 10*(2904).

Stevens, W. D., Kravitz, D. J., Peng, C. S., Tessler, M. H., & Martin, A. (2017).
Privileged Functional Connectivity Between the Visual Word Form Area
and the Language System. *Journal of Neuroscience*, 0138-0117.

Su, M., Zhao, J., Thiebaut de Schotten, M., Zhou, W., Gong, G., Ramus, F., et al.
(2018). Alterations in white matter pathways underlying phonological and
morphological processing in Chinese developmental dyslexia. *Dev Cogn
Neurosci, 31*, 11-19.

Thiebaut de Schotten, M., Cohen, L., Amemiya, E., Braga, L. W., & Dehaene, S.
(2014). Learning to read improves the structure of the arcuate fasciculus.
Cereb Cortex, 24(4), 989-995.

Tschentscher, N., Ruisinger, A., Blank, H., Diaz, B., & von Kriegstein, K. (2018).
Reduced structural connectivity between left auditory thalamus and the
motion-sensitive planum temporale in developmental dyslexia. *arXiv
preprint arXiv:1811.11658.*

van Atteveldt, N., & Ansari, D. (2014). How symbols transform brain function: A
review in memory of Leo Blomert. *Trends in Neuroscience and Education,
3*(2), 44-49.

van Atteveldt, N., Formisano, E., Goebel, R., & Blomert, L. (2004). Integration of
letters and speech sounds in the human brain. *Neuron, 43*(2), 271-282.

van Atteveldt, N. M., Formisano, E., Blomert, L., & Goebel, R. (2007). The effect of temporal asynchrony on the multisensory integration of letters and speech sounds. *Cereb Cortex, 17*(4), 962-974.

van Atteveldt, N. M., Formisano, E., Goebel, R., & Blomert, L. (2007). Top–down task effects overrule automatic multisensory responses to letter–sound pairs in auditory association cortex. *NeuroImage, 36*(4), 1345-1360.

Vandermosten, M., Boets, B., Poelmans, H., Sunaert, S., Wouters, J., & Ghesquiere, P. (2012). A tractography study in dyslexia: neuroanatomic correlates of orthographic, phonological and speech processing. *Brain, 135*(Pt 3), 935-948.

Wang, F., Karipidis, II, Pleisch, G., Fraga-Gonzalez, G., & Brem, S. (2020). Development of Print-Speech Integration in the Brain of Beginning Readers With Varying Reading Skills. *Front Hum Neurosci, 14*(289), 289.

Wechsler, D. (1974). *WISC-R, Wechsler intelligence scale for children, revised*: Psychological Corporation.

Wenger, E., Brozzoli, C., Lindenberger, U., & Lövdén, M. (2017). Expansion and Renormalization of Human Brain Structure During Skill Acquisition. *Trends in Cognitive Sciences, 21*(12), 930-939.

Xia, M., Wang, J., & He, Y. (2013). BrainNet Viewer: a network visualization tool for human brain connectomics. *PloS one, 8*(7), e68910.

Xia, Z., Hoeft, F., Zhang, L., & Shu, H. (2016). Neuroanatomical anomalies of dyslexia: Disambiguating the effects of disorder, performance, and maturation. *Neuropsychologia, 81*, 68-78.

Xia, Z., Yang, T., Cui, X., Hoeft, F., Liu, H., Shu, H., et al. (2020). Neurofunctional mechanisms underlying audiovisual integration of characters and *pinyin* in Chinese children. *bioRxiv*, 2020.2005.2031.126128.

Xia, Z., Zhang, L., Hoeft, F., Gu, B., Gong, G., & Shu, H. (2018). Neural correlates of oral word reading, silent reading comprehension, and cognitive subcomponents. *International Journal of Behavioral Development, 0*(0), 0165025417727872.

Xu, W., Kolozsvari, O. B., Oostenveld, R., Leppanen, P. H. T., & Hamalainen, J. A. (2019). Audiovisual Processing of Chinese Characters Elicits Suppression and Congruency Effects in MEG. *Front Hum Neurosci, 13*(18), 18.

Xue, J., Shu, H., Li, H., Li, W., & Tian, X. J. J. o. P. R. (2013). The Stability of Literacy-Related Cognitive Contributions to Chinese Character Naming and Reading Fluency. *42*(5), 433-450.

Yang, J., Shu, H., McCandliss, B. D., & Zevin, J. D. (2013). Orthographic
influences on division of labor in learning to read Chinese and English:
Insights from computational modeling. *Biling (Camb Engl), 16*(Spec Iss 2),
354-366.

Zhang, L., Xi, J., Xu, G., Shu, H., Wang, X., & Li, P. (2011). Cortical dynamics of
acoustic and phonological processing in speech perception. *PLoS One, 6*(6),
e20963.

Zhang, Y., Zhang, L., Shu, H., Xi, J., Wu, H., Zhang, Y., et al. (2012).
Universality of categorical perception deficit in developmental dyslexia: an
investigation of Mandarin Chinese tones. *J Child Psychol Psychiatry,
53*(8), 874-882.

Zhao, J., Wang, X., Frost, S. J., Sun, W., Fang, S.-Y., Mencl, W. E., et al. (2014).
Neural division of labor in reading is constrained by culture: A training
study of reading Chinese characters. *Cortex, 53*, 90-106.

Zou, L., Packard, J. L., Xia, Z., Liu, Y., & Shu, H. (2015). Neural Correlates of
Morphological Processing: Evidence from Chinese. *Frontiers in human
neuroscience, 9*.
947

Figure legends

Figure 1 Schematic description of the fMRI experimental procedure (A, B, C for a single run, a block, and a mini-block, respectively) and stimuli examples (D). Abbreviations: Aud = auditory, avC = audiovisual congruent, avI = audiovisual incongruent, Fix = fixation, Vis = visual.

Figure 2 (A) Results of the voxel-wise whole-brain ANOVA on the character conditions. Only clusters showing a significant main effect of condition were identified at the FWE corrected threshold of p-cluster < 0.05 (p-voxel < 0.001 for height). No region showed a significant main effect of group or group × condition interaction. (B) Bar plots present the brain activation level in the audiovisual congruent and incongruent conditions for the control group and dyslexic group. (C) Scatterplots display correlations between the neural audiovisual integration effect (avC-avI) in the left IFC with silent reading comprehension proficiency and related cognitive-linguistic skills (morphological awareness, rapid naming). Abbreviations: avC = audiovisual congruent, avI = audiovisual incongruent, IFC = inferior frontal cortex, STC = superior temporal cortex.

Figure 3 (A) Brain map of regions showing significant differences between dyslexic and control groups in correlation between the audiovisual integration effect and oral reading fluency (FWE corrected p-cluster < 0.05, p-voxel < 0.001 for height). Scatterplots presents correlations in each group (blue: control; red: dyslexic) in the left OTC (B), STC (C), and right STC (D). Abbreviations: avC = audiovisual congruent, avI = audiovisual incongruent, OTC = occipito-temporal cortex, STC = superior temporal cortex.

Supplementary Figure 1 The summary of demographic and behavioral measures is shown as boxplots, with the box indicating the interquartile range (IQR). The whiskers show the range of values within 1.5 × IQR and a horizontal line indicating the median. Individual data are shown as dots. The color coding is indicated in the legend below to the plot. Data visualization was done with PlotsOfData (https://huygens.science.uva.nl/PlotsOfData/).

Table 1 Demographic and behavioral profiles

Measures	Typical Readers (n = 22)			Reading Disorder (n = 23)			Comparison	
	Mean	SD	Range	Mean	SD	Range	t/X^2	p-value
Age (month)	127	6	118 ~ 140	122 †	10	111 ~ 144	1.938	0.061
Sex (female/male)	12/10			10/13			0.552	0.556
Verbal IQ (standard score)	109	12	88 ~ 139	97	11	67 ~ 118	3.668	< 0.001
Performance IQ (standard score)	115	13	82 ~ 137	105	12	82 ~ 135	2.512	0.016
Full-Scale IQ (standard score)	113	11	93 ~ 130	101	9	84 ~ 116	3.986	< 0.001
Character Recognition (item)	122	8	107 ~ 140	86 †	11	53 ~ 99	11.830	< 0.001
Standard score	1.38	0.66	0.41 ~ 2.43	-2.18	0.83	-4.520 ~ -1.020	12.701	< 0.001
Word List Reading (word/minute)	94 ‡	16	72 ~ 140	63	11	40 ~ 83	7.612	< 0.001
Standard score	0.95 †	1.11	-0.38 ~ 4.06	-1.584	0.67	-2.930 ~ -0.130	7.278	< 0.001
Reading Fluency (character/minute)	361 ‡	104	220 ~ 565	163	56	70 ~ 304	7.890	< 0.001
Standard score	0.98 ‡	1.04	-0.38 ~ 3.12	-1.083	0.57	-2.250 ~ 0.430	7.998	< 0.001
Phoneme Deletion (item)	21 †	4	7 ~ 25	17 †	6	1 ~ 25	2.848	0.007
Rapid Naming (second)	17	2	11 ~ 20	22	4	15 ~ 30	-5.481	< 0.001
Morphological Production (item)	24 ‡	3	20 ~ 29	22	3	13 ~ 28	2.927	0.005

Note. † Shapiro-Wilk p < 0.05, ‡ 0.05 < Shapiro-Wilk p < 0.01

Table 2 Significant clusters in the voxel-wise whole-brain analyses

Material and contrast	Label	Brain area	$P_{\text{FWE-corrected}}$	Size	Peak F	X	Y	Z
Character								
Main effect of condition	LSTC	Left middle and superior temporal	0.007	391	26.86	-60	-32	6
	LIFC	Left inferior frontal gyrus,	0.024	281	23.90	-46	16	14
Pinyin								
Group difference in brain-reading	LOTC	Inferior and middle occipital gyri,	0.004	358	29.28	-44	-72	-2
(Reading: *Oral reading fluency*)	LSTC	Left middle and superior temporal	0.039	197	23.02	-52	-44	10
	RSTC	Right superior temporal gyrus,	0.047	185	38.68	62	-8	4

Note. The significant clusters were identified with the FWE-corrected threshold of *p*-cluster < 0.05 (*p*-voxel < 0.001 for height). Brain area labeling is based on the AAL atlas. Cluster size refers to the number of voxels.

Table 3 Brain-behavior correlations in the significant clusters in the voxel-wise whole-brain regression analyses.

Behavior measure:	Group	Integration Effect (avC-avI)		avC		avI	
Oral reading fluency		*r*-value	*p*-value	*r*-value	*p*-value	*r*-value	*p*-value
L OTC	Control	0.525 *	0.021	0.685 **	0.001	0.118	0.631
	Dyslexic	-0.804 ***	< 0.001	0.026	0.913	0.674 **	0.001
L STC	Control	0.693 **	0.001	0.588 **	0.008	-0.014	0.956
	Dyslexic	-0.668 **	0.001	-0.050	0.833	0.445 *	0.049
R STC	Control	0.655 **	0.002	0.240	0.321	-0.259	0.284
	Dyslexic	-0.706 ***	< 0.001	-0.181	0.444	0.543 *	0.013

Note. Age, sex, and performance IQ were controlled statistically. Abbreviations: avC = audiovisual congruent, avI = audiovisual incongruent. *** $p < 0.001$, ** $p < 0.01$, * $p < 0.05$

A

Left IFC   Left STC

F-value

Character: main effect of condition

B

LSTC   LIFC

Control   Dyslexic

C

Left IFC-Reading Comprehension

Left IFC-Cognitive Skills

Control

avC-avI (residuals) vs Silent reading comprehension (residuals)

avC-avI: *r* = 0.571, *p* = 0.011 *
- avC: *r* = 0.222, *p* = 0.360
- **avI: *r* = -0.526, *p* = 0.021 ***

avC-avI (residuals) vs Morphological awareness (residuals)

avC-avI: *r* = 0.551, *p* = 0.014 *
- **avC: *r* = 0.828, *p* < 0.001 *****
- avI: *r* = 0.102, *p* = 0.679

Dyslexic

avC-avI (residuals) vs Silent reading comprehension (residuals)

avC-avI: *r* = 0.391, *p* = 0.088 †
- avC: *r* = 0.400, *p* = 0.080 †
- avI: *r* = -0.185, *p* = 0.436

avC-avI (residuals) vs Rapid naming (residuals)

avC-avI: *r* = -0.509, *p* = 0.022 *
- avC: *r* = -0.400, *p* = 0.081 †
- avI: *r* = 0.340, *p* = 0.143

A

Left STC

Left OTC

Right STG

F-value
+6
0

Pinyin: group difference on brain-behavior correlation

B **Left OTC-Oral Reading**

■ Control
■ Dyslexic

avC-avI (residuals)

Control:
r = 0.525, *p* = 0.021 *

Dyslexic:
r = -0.804, *p* < 0.001 ***

Oral reading fluency (residuals)

C **Left STC-Oral Reading**

avC-avI (residuals)

Control:
r = 0.693, *p* = 0.001 ***

Dyslexic:
r = -0.668, *p* = 0.001 ***

Oral reading fluency (residuals)

D **Right STC-Oral Reading**

avC-avI (residuals)

Control:
r = 0.655, *p* = 0.002 **

Dyslexic:
r = -0.707, *p* < 0.001 ***

Oral reading fluency (residuals)