Atypical relationships between neurofunctional features of print-sound integration and reading abilities in Chinese children with dyslexia

Zhichao Xia [a, b †*], Ting Yang [c, d †], Xin Cui [a, e], Fumiko Hoeft [e, f, g, h], Hong Liu [c, d, f], Xianglin Zhang [a], Xiangping Liu [c, d *], Hua Shu [a *]

[a] State Key Laboratory of Cognitive Neuroscience and Learning & IDG/McGovern Institute for Brain Research, Beijing Normal University, China

[b] School of Systems Science, Beijing Normal University, China

[c] Faculty of Psychology, Beijing Normal University, China

[d] Beijing Key Laboratory of Applied Experimental Psychology, National Demonstration Center for Experimental Psychology Education, Faculty of Psychology, Beijing Normal University, China

[e] Haskins Laboratories, USA

[f] Department of Psychological Sciences and Brain Imaging Research Center, University of Connecticut, USA

[g] Department of Psychiatry and Weill Institute for Neurosciences, University of California, San Francisco, USA

[h] Department of Neuropsychiatry, Keio University School of Medicine, Japan

[†] These authors have contributed equally to this work and share first authorship

[*] Correspondence:

Zhichao Xia, State Key Lab of Cognitive Neuroscience and Learning, Beijing Normal University, China. Email: xiazc.psy@gmail.com;

Hua Shu, State Key Lab of Cognitive Neuroscience and Learning, Beijing Normal University, China. Email: shuhua@bnu.edu.cn;

Xiangping Liu, Faculty of Psychology, Beijing Normal University, China. Email: lxp599@163.com

Running Title: Print-sound integration in Chinese dyslexia

Abstract

Conquering print-sound mappings (e.g., grapheme-phoneme correspondence rules) is vital for developing fluent reading skills. In neuroimaging research, this ability can be indexed by activation differences between audiovisual congruent against incongruent conditions in brain areas such as the left superior temporal cortex. In line with it, individuals with dyslexia have difficulty in tasks requiring print-sound processing, accompanied by a reduced neural integration. However, existing evidence is almost restricted to alphabetic languages. Whether and how multisensory processing of print and sound is impaired in Chinese dyslexia remains underexplored. In this study, we applied a passive audiovisual integration paradigm with functional magnetic resonance imaging to investigate the possible dysfunctions in processing character-sound (opaque; semantics can be automatically accessed) and pinyin-sound associations (transparent; no particular meaning can be confirmed) in Chinese dyslexic children. Unexpectedly, the dyslexic group did not show reduced neural integration compared with typically developing readers in either character or pinyin experiment. However, the results revealed atypical correlations between neural integration and different reading abilities in dyslexia. Specifically, while the neural integration in the left inferior frontal cortex in processing character-sound pairs correlated with silent reading comprehension in both children with and without dyslexia, it was associated with morphological awareness (semantic-related) in controls but with rapid naming (phonological-related) in dyslexics. This result indicates Chinese dyslexic children may not use the same grapho-semantic processing strategy as their typical peers do. As for pinyin-sound processing, while a stronger neural integration in the direction of "congruent > incongruent" in the left occipito-temporal cortex and bilateral superior temporal cortices was associated with better oral reading fluency in the control group, an opposite pattern was found in dyslexia. This finding may reflect dyslexia's dysfunctional recruitment of the regions in grapho-phonological processing, which further impedes character learning.

Keywords

audiovisual integration, character, Chinese, dyslexia, individual differences, pinyin

Highlights

- Neurofunctional correlates of print-sound integration in Chinese children with and without dyslexia are investigated.
- Dyslexic children show atypical relationships between neural audiovisual integration and reading abilities.
- Chinese children with dyslexia are likely to use inefficient strategies to process characters and pinyin.

1 Introduction

Reading consists of multiple cognitive processes, and it takes years of formal instruction to achieve a high proficiency. In this process, establishing robust links between orthographic and phonological representations (e.g., grapheme-phoneme correspondence [GPC] rules) is one initial and fundamental step (Perfetti & Harris, 2013). Behavioral and neuroimaging studies of alphabetic languages reveal that it is critical to conquer the GPC rules to develop fluent reading skills. The failure will impede building efficient grapho-semantic mapping and eventually result in reading difficulties (Blomert, 2011; Di Folco, Guez, Peyre, & Ramus, 2020; Richlan, 2019; Shaywitz, 1998). Nowadays, most researchers agree that the manifestation of dyslexia is associated with linguistic features in a given language (Richlan, 2020). However, while existing evidence is almost restricted to alphabetic orthographies, the question of whether and to what extent print-sound integration is impaired in Chinese children with dyslexia remains underexplored, especially at the neurofunctional level.

Chinese has a morpheme-based logographic writing system (Perfetti, Cao, & Booth, 2013). In addition to phonological information, semantics is strongly involved in even the most fundamental processing—character recognition (Bi, Han, Weekes, & Shu, 2007; Guan, Fraundorf, & Perfetti, 2020; Liu et al., 2017; Yang, Shu, McCandliss, & Zevin, 2013; Zhao et al., 2014). At the behavioral level, longitudinal and meta-analytic studies have demonstrated the importance of both phonological-related (e.g., phonological awareness [PA], rapid naming [RAN]) and semantic-related (e.g., morphological awareness [MA]) skills in Chinese reading development (Pan et al., 2016; Lei et al., 2011; Liu et al., 2017; Ruan, Georgiou, Song, Li, & Shu, 2018). However, it should also be noted that the tasks used in these studies required explicit processing of the written scripts. A similar situation exists at the brain level. Previous functional magnetic resonance imaging (fMRI) studies revealed hypoactivation in the left inferior and middle frontal areas during visual rhythming and lexical decision in children with dyslexia, suggesting dysfunctions of the neural substrates underlying both print-to-sound and print-to-meaning mappings in tasks requiring explicit processing (Cao et al., 2016; Liu et al., 2012; Siok, Perfetti, Jin, & Tan, 2004). Paralleling the fMRI research, structural and diffusion imaging studies also provided evidence on alterations in morphometry of these regions and white matter tracts connecting them (Siok, Niu, Jin, Perfetti, & Tan, 2008; Su et al., 2018; Xia, Hoeft, Zhang, & Shu, 2016). Hence, while these findings indicate deficits in grapho-semantic and grapho-phonological processing in Chinese children with dyslexia, the question of whether implicit and automatic processing is impaired remains largely unknown.

In this study, we adopted a passive fMRI audiovisual paradigm (i.e., without explicit phonological or semantic judgment), which is appropriate for investigating automaticity in reading-related processing. This paradigm has been used in shallow orthographies such as Dutch and demonstrated the impaired letter-sound automatized integration as a likely proximal cause of dyslexia that is independent of phonological processing deficits (Blau et al., 2010;

Blau, van Atteveldt, Ekkebus, Goebel, & Blomert, 2009; Blau, van Atteveldt,
Formisano, Goebel, & Blomert, 2008). This paradigm's basic logic is that if a
brain area integrates auditory and visual inputs or is involved in the subsequent
higher-level cognitive processes, its activation should differ between the
congruent and incongruent conditions. This effect is usually named "congruency
effect" when the activation in the congruent condition was stronger than the
incongruent condition and is named "incongruency effect" otherwise. Here we
used the term "audiovisual integration effect" (or "neural integration" for short),
given that both directions indicate successful multimodal information
integration. Since this effect can be observed even when no task or a passive task
is used, researchers regard it to reflect implicit processing (Blau et al., 2010; van
Atteveldt, Formisano, Goebel, & Blomert, 2007). To date, the neural integration
has been demonstrated in skilled adult readers and typically developing children
(Blau et al., 2010; van Atteveldt & Ansari, 2014; van Atteveldt, Formisano,
Goebel, & Blomert, 2004; van Atteveldt, Formisano, Blomert, & Goebel, 2007).

131   Of importance, direction and strength of the neural integration are
affected by several factors, such as characteristics of participants and
orthographic depth of languages (Blau et al., 2010; Blau et al., 2009; Holloway,
van Atteveldt, Blomert, & Ansari, 2015; Kronschnabel, Brem, Maurer, &
Brandeis, 2014; Wang, Karipidis, Pleisch, Fraga-Gonzalez, & Brem, 2020). For
example, individuals with dyslexia showed an atypical pattern in brain areas
such as the superior temporal cortex (STC) (Blau et al., 2010; Blau et al., 2009).
This anomaly was driven by hypo-activation in the congruent condition along
with hyper-activation in the incongruent condition in dyslexia, indicating
reduced neural integration and lack of suppression, respectively. In terms of
orthographic depth, investigations were administered with Chinese adults (Xu,
Kolozsvari, Oostenveld, Leppanen, & Hamalainen, 2019) and typically
developing children recently (Xia et al., 2020). In particular, Xia et al. (2020)
used Chinese characters and pinyin (a transparent alphabetic coding system
that represents the pronunciation of characters, which is taught at the earliest
stage of Chinese reading development and used as a scaffold in learning new
characters) as experimental materials and observed a significant audiovisual
integration effect in the direction of "congruent < incongruent" in the left inferior
frontal cortex (IFC) and bilateral STC in processing character-sound
associations. Moreover, neural integration in the left IFC in response to
character-sound pairs and that in the left STC in response to pinyin-sound pairs
were associated with children's performance in silent reading comprehension
that relies on grapho-semantic mapping and oral word reading fluency that
relies on grapho-phonological processing, respectively. This pattern is likely to be
driven by stimuli's linguistic properties, including orthographic transparency
and involvement of semantics. Using the same experimental design, the current
fMRI study aimed to examine whether the neural audiovisual integrations of
character-sounds and pinyin-sounds are impaired in Chinese dyslexia and how
they associate with different levels of reading abilities.

160   Notably, while group comparison has been widely used to identify neural
deficits in dyslexia, approaches focusing on individual differences also provide

invaluable insights. In this case, the brain-behavior correlation is a useful
strategy (Jednorog et al., 2015; Pernet, Andersson, Paulesu, & Demonet, 2009),
with which two primary patterns can be identified. The first is a universal brain-
behavior correlation regardless of reading status (dyslexia vs. control), indicating
that the same neural system supports the cognitive processing in both groups.
For example, children's PA is correlated with the microstructural feature of the
left arcuate fasciculus, even after controlling group effect (Su et al., 2018;
Vandermosten et al., 2012). Alternatively, there could be distinct ways in which
reading abilities correlated with brain measures between children with and
without dyslexia, indicating dysfunction or compensation (Hoeft et al., 2011;
Pernet et al., 2009; Rumsey et al., 1999; Tschentscher, Ruisinger, Blank, Diaz, &
von Kriegstein, 2018). For example, while typical readers rely more (higher
regional cerebral blood flow) on the left inferior parietal lobule (IPL), higher
activation in this area is associated with worse reading performance in dyslexia
(Rumsey et al., 1999). However, previous studies commonly conducted
correlation analyses while pooling individuals from different groups. Since the
participants were selected on purpose, between-group differences could drive the
results (Blau et al., 2010). As mentioned above, this issue can be addressed by
controlling the effect of group in the statistic model or directly comparing
correlations between dyslexia and typical readers.

To summarize, the main aim of this study was to investigate the possible
impairments in Chinese children with dyslexia in implicit processing of print-
sound associations and related information (e.g., semantics). We asked two
specific questions. First, whether the neurofunctional correlates of print-sound
integration differ between the dyslexics and controls. Second, whether the
relationships between neural integration and reading abilities differ between
groups. We adopted a passive fMRI audiovisual paradigm and used characters
and pinyin—scripts with contrasting linguistic features—as experimental
materials. Both group comparison and individual differences analytic approaches
were performed. Based on the prior research in typically developing children (Xia
et al., 2020), we predicted a reduced neural integration in Chinese children with
dyslexia. In addition, dyslexic children might display atypical brain-behavior
correlations or recruit other brain regions to integrate cross-modal information.

196 2 Methods

197 2.1 Participants and behavioral measures

In this study, dyslexia was operationalized by the criteria of having normal
intelligence (≥ 80 on the abbreviated version of the *Chinese Wechsler*
Intelligence Scale for Children; Wechsler, 1974) but manifesting reading
difficulty (below -1 *SD* of the norm on a standardized reading screening task
Character Recognition; Xue, Shu, Li, Li, & Tian, 2013). On the other hand, each
child in the control group should have normal intelligence (≥ 80) and a score
above -0.5 *SD* of the norm on the reading screening task (the aim was to increase

the gap in reading skills between groups). In addition, children in both groups
should be right-handed (Oldfield, 1971) native speakers of Chinese, with normal
hearing and normal or corrected-to-normal vision, and were free from
neurological or psychiatric disorders. Finally, only the children that completed
all the task fMRI runs, with an overall accuracy equal to or higher than 75% in
the in-scanner passive task and less than 25% time-points labeled as outliers
(i.e., "bad volume") in each run (data preprocessing section) were included.
Initially, one hundred children in grades 3-6 (including 45 dyslexia) were
recruited from local elementary schools. According to the inclusion criteria
described above, 23 dyslexic children (10 girls; age 111-144 months, M [SD] =
122 [10]) were included in the final analysis. Twenty-one dyslexic children were
excluded due to uncompleted MRI data collection (n = 9), severe head motion
artifacts (n = 9), or poor in-scanner performance (n = 3). One child with a history
of dyslexia diagnosis but performed normally in the character recognition task at
the time of data collection was also excluded (this child had received an intensive
behavior intervention program). The controls were chosen to match the dyslexic
group on grade, age, and sex. The final control group consisted of 22 typically
developing children with qualified neuroimaging data (12 girls; age 118-140
months, M [SD] = 127 [6]).

Each child received a battery of behavioral tests on reading and cognitive-
linguistic skills individually in a silent room on the same day of the MRI session.
The reading measurements contained: (1) an untimed character naming task
(*Character Recognition*) to estimate the number of characters children had
conquered (Xue et al., 2013); (2) an oral word reading task (*Word List Reading*)
to measure how fast the participant accurately retrieved phonological
representations from visually presented high-frequency two-character words
(Zhang et al., 2012); and (3) a timed comprehension task (*Silent Reading
Comprehension*) to assess the proficiency of meaning access and semantic
judgment (Lei et al., 2011). In addition, PA, RAN, and MA—the three most
critical cognitive-linguistic skills in Chinese reading acquisition—were measured
by *Phoneme Deletion* (Li, Shu, McBride-Chang, Liu, & Peng, 2012), *Digit RAN*
(Liu et al., 2017), and *Morphological Production* (Shu, McBride-Chang, Wu, &
Liu, 2006).

Written informed consents were obtained from all the children and their
guardians after a detailed explanation of the objectives and procedure of the
study. After the experiment, each child received a book and a set of stationery as
compensation. This study was approved by the Institutional Review Board of the
State Key Laboratory of Cognitive Neuroscience and Learning at Beijing Normal
University. The data collection was conducted in 2018.

2.2 Experimental design

This study adopted a passive audiovisual paradigm that has been widely used in
the previous fMRI studies investigating the neural basis of letter-sound
integration (**Figure 1**). The details about the stimuli and procedure can be found
in Xia et al. (2020). In brief, 56 pictographic characters with high-frequency ($M \pm$

SD = 929 ± 1486 per million; Chinese Single-Character Word Database;
https://blclab.org/pyscholinguistic-norms-database/; Liu et al., 2007) and are
frequently used as radicals in phonograms were selected. These characters are
visually simple (number of strokes: *M* = 4.34, *Range* = 1-9), learned early (age of
acquisition: *M* = 3 years, *Range* = 2-5 years), and with high rating scores (7 as
the highest value) on concreteness (5.76 ± 1.19) and imageability (5.96 ± 1.00).
The pinyin spellings of these characters were used as the visual stimuli in the
pinyin experiment. The auditory stimuli (duration: *M* ± *SD* = 476.3 ± 87.5 ms)
were the characters' sounds (i.e., syllables). A native Chinese male recorded the
audio files with a sampling rate of 44.1 kHz and 16-bit quantization. The sound
files were then normalized to 85 dB and bandpass (100-4000 Hz) filtered with
Audacity (https://www.audacityteam.org/).

The study consisted of 4 task fMRI runs, with the first and second runs for
the pinyin experiment and the third and fourth runs for the character
experiment. We used the fixed order to prevent priming from characters on
visually presented pinyin stimuli. Two unimodal (auditory [Aud]; visual [Vis])
and two cross-modal conditions (congruent [avC]; incongruent [avI]) were created
for each experiment (**Figure 1**). In this study, we focused on activation
differences between the congruent against incongruent conditions—the neural
integration. A block design was used to deliver stimuli. There were 8 task blocks
(duration = 20.8 s; 2 blocks for each condition) interleaved with 9 rest blocks
(duration = 20.8 s) in a single run. A task block contained 4 mini-blocks. A 1.5 s
period was used to collect a whole-brain volume within each mini-block, and a
silent period of 3.7 s was used to present stimuli (see "image acquisition" part).
The stimuli were presented in white at the center of a black background ("*KaiTi*"
font, 96 pt for characters; "*Century Schoolbook*" font, 90 pt for pinyin). A
crosshair was presented at the center of the screen whenever there was no
stimulus. To help children keep their attention on the stimuli while avoiding
explicit congruency judgment, we used a target detection task (Blau et al., 2010).
Specifically, in each task block, two out of 16 experimental stimuli were
randomly replaced with the auditory target (440 Hz pure tone), visual (an
unpronounceable symbol) target or their combination. The participant was asked
to press a button with the right index finger as accurately and quickly as
possible whenever the target appeared.

283   -------------------------------------

284   Insert Figure 1 about here

285   -------------------------------------

2.3 Image acquisition

All brain images were collected at Beijing Normal University Imaging Center for
Brain Research using a 3-Tesla Siemens MAGNETOM Trio Tim scanner with a
12-channel head coil. The children first attended a training session to get
familiar with the experimental environment and the scanning noise. During the
formal scan, foam pads were used to hold their heads secure during scanning to

improve image quality. In addition, children could take a break between
sequences to reduce the possible fatigue effect. For each participant, two
functional runs for the pinyin experiment, one anatomical run for structural
images, and two functional runs for the character experiment were administered
sequentially. The quality of the brain images was evaluated immediately by a
radiologist who was blinded to the details of this study.

The parameters of the functional images (Gradient Echo Planar Imaging
[EPI]) were as follows, repetition time, 5.2 s; echo time, 32 ms; acquisition time,
1.5 s; flip angle, 90 degrees; slice thickness, 4.5 mm; interscan gap, 0.675 mm,
voxel size, $3.0 \times 3.0 \times 4.5$ mm^3; 24 slices; 68 volumes). Since this study contained
auditory stimuli, to avoid artifacts induced by the noise during scanning, we
used a sparse sampling design with 1.5 s for image collection and 3.7 s delay for
stimuli presentation (Shah et al., 2000). The parameters of the structural images
(Magnetization-Prepared Rapid Acquisition with Gradient Echo [MPRAGE])
were as follows, repetition time, 5.2 s; echo time, 3.39 ms; inversion time, 1.1 s;
flip angle, 7 degrees; slice thickness, 1.33 mm; interscan gap, 0 mm; voxel size,
$1.3 \times 1.0 \times 1.3$ mm^3; 144 axial slices).

2.4 Data preprocessing

Functional data were analyzed with SPM12 (http://www.fil.ion.ucl). Three
dummy scans were added at the beginning of each run to avoid the T1
equilibration effect. No additional volumes were discarded during the
preprocessing. The sequence was corrected for head motion. The "bad volumes"
were identified with ART-based outlier detection
(https://www.nitrc.org/projects/artifact_detect). Considering the age range of
participants, a liberal threshold (intensity > 9 SD; frame-to-frame head motion >
2 mm) was used. T1 images were segmented and used for transferring the fMRI
data from native space to the standard Montreal Neurological Institute (MNI)
space. The normalized images were smoothed with an 8-mm Full-Width Half-
Maximum Gaussian kernel, and the resulting data were used in the subsequent
model estimation. In the 1st level analysis, 4 experimental conditions, 7 head
motion parameters (3 for translations, 3 for rotations, and 1 for framewise
displacement), and the time point of each "bad volume" were included in the
model. The contrast map in which the value of each voxel refers to "avC-avI" (i.e.,
positive value stands for higher activation in avC than avI; negative value
stands for lower activation in avC than avI) was calculated for each child and
used in the subsequent analyses.

2.5 Statistics

Deficits in reading and reading-related cognitive skills in children with dyslexia
were examined at the behavioral level. We first performed descriptive statistics
of multiple behavioral tasks in each group and then conducted group
comparisons. Next, we calculated correlations between reading and cognitive-
linguistic skills within each group and compared them between groups. The
Spearman method was used for the variable that was not normally distributed.
As for in-scanner performance, a correct response was defined as the button

press to the target with a reaction time (RT) ranged from 200 to 2000 ms. Only correct trials were used to calculate the average RT. Finally, the effects of group and run and their interaction were examined on accuracy (ACC) and RT with analysis of variance (ANOVA).

We conducted brain analyses focusing on differences between the two cross-modal conditions for the character and pinyin experiments, respectively, with the same analytic approaches. The nuisance variables of age, sex, and performance IQ were controlled in all the analyses. First, a voxel-wise whole-brain 2 (group: control vs. dyslexic) × 2 (condition: avC vs. avI) ANOVA (i.e., the group comparison approach) was conducted to examine whether the brain regions showing neural integration in children with dyslexia were the same as those in the control group. Significant clusters were identified with the FWE-corrected threshold of p-cluster < 0.05 (p-voxel < 0.001 for height). These clusters were used as regions-of-interest (ROIs), and post hoc t-tests were conducted to interrogate the effects. Complementary ROI analyses were performed to examine correlations between the neural integration with reading and reading-related cognitive-linguistic skills in each group, to identify the reading-related processes involved in audiovisual integration of character/pinyin for typical and dyslexic children, respectively. Once a significant correlation was revealed, we conducted a correlation coefficient comparison between groups.

Next, we used two individual differences approaches to examine the shared and different brain-behavior correlations in children with and without dyslexia in the character and pinyin experiments at the whole-brain level. First, to identify the shared neural basis associated with behavioral performance between groups, we conducted voxel-wise whole-brain regression analyses on the contrast maps of avC against avI across all the participants while controlling for the effects of group, age, sex, and performance IQ. Children's performance in *Word List Reading* and *Silent Reading Comprehension* tasks were used as regressors in separate models to examine the relationships between neural processing features during print-sound integration with reading abilities that rely more on grapho-phonological mapping and grapho-semantic mapping, respectively. Second, we examined whether the associations between neural integration and reading abilities differ between groups. Same as the previous analysis, we used *Word List Reading* and *Silent Reading Comprehension* as variates of interest in separate models, along with the factor of group and the interaction. In each analysis, an F-test was administered with the FWE-corrected threshold of p-cluster < 0.05 (p-voxel < 0.001 for height), followed by ROI analysis to interrogate the significant effects.

To visualize the results, significant clusters were presented on a FreeSurfer surface template with BrainNet Viewer (Xia, Wang, & He, 2013). Anatomical labeling was performed using the AAL atlas with DPABI (http://rfmri.org/dpabi).

All the behavioral and ROI analyses were administered with SPSS (v24; SPSS Inc., Chicago, IL, USA). Effects were considered significant at $p < 0.05$, and $0.05 < p < 0.1$ was considered indicative of a trend.

3 Results

3.1 Behavior

3.1.1 Reading measures and group comparisons

Statistical metrics including *M*, *SD*, *Range*, and result of the Shapiro-Wilk test of each behavioral measurement are presented in Table 1 (see Supplementary Figure 1 for plots). The dyslexic group performed worse in all the reading and reading-related cognitive tasks than the typically developing children (all *p*'s < 0.007). No significant between-group differences were found on age or sex (both *p*'s > 0.05). In addition, the IQs of all the children were within the normal range, while the typical readers had higher scores on both verbal and performance subscales.

Insert Table 1 about here

3.1.2 Correlations between reading and reading-related cognitive skills

Both same and different correlations between reading and cognitive-linguistic skills were observed between children with and without dyslexia. In typical readers, character recognition was significantly correlated with MA ($r = 0.589$, $p = 0.004$) but not PA ($r = 0.117$, $p = 0.606$) or RAN ($r = -0.301$, $p = 0.174$). In dyslexics, however, this ability was associated with PA ($r = 0.489$, $p = 0.018$), but not RAN ($r = -0.269$, $p = 0.214$) or MA ($r = 0.008$, $p = 0.970$). The group difference on the correlation coefficients between character recognition and MA was significant ($Z = 2.09$, $p = 0.037$). A similar pattern was found in silent reading comprehension proficiency, where the scores were correlated with MA ($r = 0.456$, $p = 0.033$) in controls but not dyslexia ($r = -0.012$, $p = 0.957$). On the contrast, oral reading fluency was significantly correlated with RAN in both groups (controls: $r = -0.531$, $p = 0.012$; dyslexics: $r = -0.578$, $p = 0.004$).

3.1.3 In-scanner performance

The aim of using a passive target detection task was to ensure that the participant focused their attention on the stimuli delivered via auditory and visual modalities without performing explicit congruency judgment. The results revealed that children in both groups performed the task with high ACC (controls: *M* [SD] = 96.7% [2.9]; dyslexics: *M* [*SD*] = 92.4% [5.0]). In the ANOVA, the main effects of group were significant on both ACC ($p = 0.001$) and RT ($p = 0.012$). The post-hoc analyses showed that the children with dyslexia had lower accuracy and used more time to complete the tasks than the normal controls. The main effect of run was significant on RT ($p = 0.012$; faster as the experiment

419 proceeds) but not on ACC ($p = 0.645$). No significant group × run interaction was
420 observed on either ACC or RT (both p's > 0.05).

421   -------------------------------------

422   Insert Table 2 about here

423   -------------------------------------

3.2 fMRI

We used group comparison (ANOVA), and individual differences (brain-behavior
correlation) approaches to investigate the impaired neurofunctional features
accompanying print-sound integration in Chinese children with dyslexia. In this
section, we first present the results of the character experiment, followed by the
pinyin experiment.

3.2.1 Character: whole-brain ANOVA and ROI analysis

In the voxel-wise whole-brain ANOVA, the left IFC and STC showed a
significant main effect of condition that survived the FWE corrected p-cluster <
0.05 (p-voxel < 0.001 for height; **Table 2**; **Figure 2A**). The follow-up analysis
revealed less activation in the congruent than incongruent conditions in both the
controls (LIFC: $t = 4.361$, $p < 0.001$; LSTC: $t = 3.646$, $p = 0.002$) and the dyslexics
(LIFC: $t = 2.346$, $p = 0.028$; LSTC: $t = 3.427$, $p = 0.002$; **Figure 2B**). No main
effect of group or group × condition interaction survived the whole-brain FWE
correction.

In the complementary ROI analysis, we observed similar correlations
between the neural integration and reading comprehension proficiency in the left
IFC in both groups (controls: $r = 0.571$, $p = 0.011$; dyslexics: $r = 0.391$, $p = 0.088$,
marginally significant; **Figure 2C**). However, the relative contribution of each
cross-modal condition was different. To be specific, the correlation was driven by
the incongruent condition in typical readers (avC: $r = 0.222$, $p = 0.360$; avI: $r = -$
0.526, $p = 0.021$) but was more related to the congruent condition in dyslexia
(avC: $r = 0.400$, $p = 0.080$, marginally significant; avI: $r = -0.185$, $p = 0.436$).
Between-group difference on the correlation coefficients between reading
comprehension proficiency and brain activation in the incongruent condition was
significant ($Z = 2.41$, $p = 0.016$). The distinct patterns of the correlations between
the neural integration and reading-related cognitive skills further support the
idea that different mechanisms underlie the integration-comprehension
relationships in the two groups (**Figure 2C**): the neural integration in the left
IFC was correlated with MA in the controls ($r = 0.551$, $p = 0.014$) but not in the
dyslexics ($r = -0.025$, $p = 0.915$). Group difference on the correlation coefficients
was significant ($Z = 2.01$, $p = 0.044$). On the other hand, the effect was correlated
with RAN in the dyslexics ($r = -0.509$, $p = 0.022$) but not in the controls ($r = -$
0.338, $p = 0.157$).

3.2.2 Character: whole-brain group × behavior interaction

Regarding the individual differences approaches, no cluster survived the FWE-
corrected threshold of p-cluster < 0.05 (p-voxel < 0.001 for height) in the analysis
investigating the same relationships between the neural integration and reading
abilities across groups or that explored the correlation differences between
groups.

464   ---------------------------------------

465   Insert Figure 2 about here

466   ---------------------------------------

3.2.3 Pinyin: whole-brain group comparison

The same analytic approaches were used in the pinyin experiment. In the voxel-
wise whole-brain ANOVA, no regions showed significant main effect of group or
condition or their interaction at the FWE corrected threshold of p-cluster < 0.05
(p-voxel < 0.001 for height).

3.2.4 Pinyin: whole-brain group × behavior interaction and ROI analysis

We investigated the neural deficits with the individual differences approaches in
a whole-brain fashion. While no region displayed the same brain-behavior
correlation across groups, clusters located in the left occipitotemporal cortex
(OTC) and bilateral STC showed significant between-group differences in the
correlation between the neural integration and oral reading fluency (FWE
corrected p-cluster < 0.05, p-voxel < 0.001 for height; **Table 2**; **Figure 3A**). The
subsequent analyses revealed positive brain-reading correlations in typical
readers and negative correlations in children with dyslexia (**Figure 3B, C and**
D; Table 3). Furthermore, the correlations in the left OTC (avC: $r = 0.685$, $p = $
0.001; avI: $r = 0.118$, $p = 0.631$) and STC (avC: $r = 0.588$, $p = 0.008$; avI: $r = -$
0.014, $p = 0.956$) were driven by the congruent condition in the control group,
while oral word reading fluency was not correlated with brain activation of the
right STC in either the congruent condition ($r = 0.240$, $p = 0.321$) or incongruent
condition ($r = -0.259$, $p = 0.284$). In contrast, in dyslexia, the correlations were
driven by the incongruent condition (left OTC: $r = 0.674$; $p = 0.001$; left STC: $r = $
0.445, $p = 0.049$; right STC: $r = 0.543$, $p = 0.013$) but not the congruent condition
(left OTC: $r = 0.026$, $p = 0.913$; left STC: $r = -0.050$, $p = 0.833$; right STC: $r = -$
0.181, $p = 0.444$) in all three brain regions. Between-group differences on the
correlations of the oral reading fluency and activation in the congruent condition
in the left OTC ($Z = 2.54$, $p = 0.011$) and left STC ($Z = 2.26$, $p = 0.024$), and on the
correlations of the oral reading fluency and activation in the in incongruent
condition in the left OTC ($Z = -2.18$, $p = 0.029$) and right STC ($Z = -2.73$, $p = $
0.006) were significant.

Regarding correlations between the neural integration and reading-
related cognitive-linguistic skills, the left STC was negatively associated with
RAN in controls ($r = -0.479$, $p = 0.038$) and showed a trend positively correlated

with RAN in dyslexics ($r = 0.394$, $p = 0.085$, marginally significant). The
between-group difference was significant ($Z = -2.93$, $p = 0.003$).

501   -------------------------------------

502   Insert Figure 3 about here

503   -------------------------------------

504   -------------------------------------

505   Insert Table 3 about here

506   -------------------------------------

4 Discussion

This study investigated the neurofunctional features of implicit print-sound
integration and their relationships with reading abilities in Chinese children
with and without dyslexia. We adopted an fMRI audiovisual paradigm with a
passive target detection task, where characters and pinyin—scripts with
dramatically different orthographic depths—were used as experimental
materials. Of importance, due to the morpho-syllabic nature of characters,
semantic information can be automatically activated during character
recognition, at least in typical readers. That is, this study enabled us to tap into
the three-way relationship between orthography, phonology, and semantics in
normal and impaired readers without demanding explicit phonological or
semantic processing. Although no between-group differences on the audiovisual
integration effect at the brain level were found, the results revealed strikingly
atypical correlations between the neural integration of both character-sounds
and pinyin-sounds with reading abilities in Chinese children with dyslexia. On
the one hand, these anomalies indicate that children with dyslexia rely more on
articulatory phonological information during implicit character processing,
reflecting a less developed automatic grapho-semantic mapping and integration.
On the other hand, it also suggests a malfunctional grapho-phonological
mapping in dyslexia and implies that these children may have difficulty in
developing the same pinyin processing strategy and transferring it to learning
characters as their typically developing peers do.

**4.1 Left IFC and inefficient grapho-semantic mapping in Chinese
dyslexia**

First, this study revealed inefficient semantic information access from visual
input in Chinese children with dyslexia. We observed slower silent reading
comprehension in the dyslexia group. Moreover, while both character recognition
accuracy and reading comprehension proficiency were associated with MA in
typical readers, children with dyslexia showed significant correlations between

reading abilities with other cognitive-linguistic skills instead of MA. Specifically,
character recognition was correlated with PA, in line with the previous study
with a large independent sample (Song, Zhang, Shu, Su, & McBride, 2020).
Regarding silent reading comprehension, although no correlation was
significant, there was a trend with RAN, a phonological processing skill that
consists of rapid and accurate phonological representation access, retrieval, and
articulatory operations. This result is in line with recent studies in which
phonological skill contributes to reading more at the early stages of reading
acquisition, and the contribution of morphological processing skill increases as
children grow (Liu et al., 2017). In addition to the direct effect, MA also mediates
the effect of PA on reading (Pan et al., 2016).

Furthermore, we used a passive fMRI audiovisual paradigm to probe the
neural bases of processing character-sound associations in Chinese children with
and without dyslexia. In terms of brain activation in the left IFC and STC,
dyslexic children seem to integrate information from auditory and visual
modalities the same way as controls, reflected as a strong audiovisual
integration effect in the direction of "congruent < incongruent". Moreover, the
neural integration in the left IFC was associated with silent reading
comprehension proficiency, regardless of reading status. However, between-
group differences were uncovered in the subsequent ROI analysis. First, while
the integration-comprehension association was driven more by the incongruent
condition in typical readers, it was driven more by the congruent condition in
children with dyslexia, suggesting dyslexic and typical children may use
different strategies in processing characters and corresponding sounds, and this
difference enlarge with reading abilities increasing within each group. Second,
while the audiovisual integration effect in the left IFC was associated with MA
in the control group, it was correlated with RAN in dyslexia. These findings
indicate that articulatory phonological processing is more likely involved in
implicit processing of character-sound pairs in children with dyslexia. In the
previous studies investigating neural impairment in Chinese dyslexia, both
hypo- and hyper-activation of the left frontal areas were reported (Cao et al.,
2016; Liu et al., 2012; Siok et al., 2004). In a meta-analysis, different parts of the
left IFC were distinguished based on functionality, where individuals with
dyslexia displayed reduced activation in the ventral part associated with
semantic processing but increased activation in the dorsal part that was
associated with articulatory processing, presumably compensating for their less
efficient grapho-semantic route (Hancock, Richlan, & Hoeft, 2017; Richlan,
Kronbichler, & Wimmer, 2011). Given that the frontal region is multifunctional
(Fedorenko & Blank, 2020; Hagoort, 2014), dyslexia may recruit it in reading-
related processing in a different way compared with typical readers.

Of note, the in-scanner task used in this study did not require any sound-
semantic or print-semantic processing. Nevertheless, since Chinese has a
morpheme-based logographic writing system that involves semantic information
even at the character processing level (Guan et al., 2020; Liu et al., 2017; Yang et
al., 2013; Zhao et al., 2014), both phonological and semantic information could be
accessed effortlessly, at least in typical readers who have received 4-5 years of

formal instruction. Thus, it is reasonable to predict that semantic processing skill plays an equal or even more critical role in reading development than phonological processing skill, and its impairments will result in reading difficulties. In line with this hypothesis, previous studies of Chinese demonstrated morphological awareness uniquely predicted reading outcomes and dyslexia status (Pan et al., 2016; Ruan et al., 2018; Song et al., 2020). The current results also showed that while the left IFC was strongly involved in both groups, it was more associated with articulatory phonological processing in dyslexia and semantic-related morphological processing in typical children. The region is close to the one found to underlie morphological processing and show hypo-activation in children with dyslexia during tasks requiring explicit semantic processing (Liu et al., 2013; Zou, Packard, Xia, Liu, & Shu, 2015).

In short, the findings of the character experiment suggest that Chinese children with dyslexia have yet to develop the same brain system for automated semantic access and integration during implicit character/word recognition as that in typically developing children. In contrast, these children are more likely to rely on an articulatory strategy by recruiting the multifunctional frontal area, which may underpin their slow reading comprehension.

4.2 Bilateral TPC, left OTC and malfunctioning grapho-phonological mapping in Chinese dyslexia

In addition to the inefficient grapho-semantic mapping, this study also indicates that children with dyslexia may not develop a typical grapho-phonological mapping. At the behavior level, the dyslexic group performed worse in the tasks measuring oral reading fluency, PA, and RAN, in line with previous studies (e.g., Lei et al., 2011). In the pinyin experiment, we observed differential brain-behavior correlations in the classic reading-related areas, including the left OTC and bilateral STC. The morphometric measurements in these regions have also been associated with oral reading fluency in Chinese school-age children (Xia et al., 2018). In particular, the neural integration in the direction of "congruent > incongruent" in response to pinyin-sound pairs was positively associated with oral reading fluency in typical controls: the better the children performed in the oral word reading task, the higher the activation was in the congruent condition than incongruent condition. In contrast, the correlation was negative in dyslexia: children with higher reading fluency showed higher activation in incongruent than congruent conditions. Additionally, the correlations in the control group were driven more by individual differences in brain responses to the congruent stimuli. In contrast, the correlations in the dyslexic group were driven more by the incongruent condition. These findings suggest that while the same brain regions were recruited for both groups' implicit audiovisual integration of pinyin, children with dyslexia may use them differently.

The OTC and STC in the left hemisphere have been regarded as critical nodes in the classic reading network. Deficits in these areas have also been repeatedly reported in dyslexia (Richlan, Kronbichler, & Wimmer, 2009; Richlan et al., 2011). On the one hand, the left OTC has been considered the interface for initially integrating orthographic, phonological, and semantic information (Price

& Devlin, 2011). In addition, the left OTC contains a specific portion in the
fusiform gyrus named Visual Word Form Area that has been found to respond
specifically to word and word-like stimuli. The left STC, on the other hand, is a
central area that represents phonological information (Boets et al., 2013; Glezer
et al., 2016), including lexical tone—the supramarginal phoneme in tonal
languages such as Chinese (Si, Zhou, & Hong, 2017; L. Zhang et al., 2011). The
left STC is functionally and structurally connected to the left OTC, which can be
shaped by learning grapho-phonological mappings (Stevens, Kravitz, Peng,
Tessler, & Martin, 2017; Thiebaut de Schotten, Cohen, Amemiya, Braga, &
Dehaene, 2014). In the current study, besides the left-hemispheric regions, the
right STC also showed significant group differences in brain-behavior
relationships. Although this region has been less frequently reported in previous
studies in alphabetic languages, it subserves lexical tone processing in Chinese
(Liang & Du, 2018; Si et al., 2017; Zhang et al., 2011). In addition, the cortical
thickness of this area is also associated with oral reading fluency in typically
developing Chinese children (Xia et al., 2018).

The differential relationship between the audiovisual integration effect
during pinyin-sound processing and oral word reading fluency in dyslexia can be
interpreted in at least two ways. First, suppose pinyin processing skill is a
continuum, and dyslexia represents the lower end. In that case, the current
finding then hints at the expansion and renormalization hypothesis of brain
plasticity associated with skill learning (Wenger, Brozzoli, Lindenberger, &
Lövdén, 2017). That is, the growth curve of print-sound integration is an inverse
U-shape. When the child starts learning pinyin, the brain response to
incongruent audiovisual pairs is lower than congruent pairs. With learning,
mismatched information extracted from visual and auditory modalities induces
higher activation during integration. Then, children turn to focus on overlearned
visual scripts by efficiently suppressing attractive auditory information at the
highly familiar stage. In this case, activation in the incongruent condition will be
suppressed and weaker than the congruent condition. This interpretation is in
line with our findings that brain activation in the congruent condition was
associated with oral reading in typical readers, whereas incongruent condition
was related to oral reading in dyslexia. The alternative explanation is also
associated with development but assumes that individuals with dyslexia process
pinyin differently from typical controls. In general, typical readers shift from
assembled to addressed phonology with reading experience increases (Mei et al.,
2014). Pinyin is assembled in nature. But since there are only approximately 400
syllables, it can be expected that typical readers in upper elementary grades who
are highly familiar with it could achieve the addressed phonology. Given that
children learn Chinese characters as holistic syllable-level units, children who
read pinyin with the same addressed phonology may benefit more. In this case,
the differential brain-behavior correlations probably reflect the assembled
phonology adopted by dyslexic children in processing pinyin-sound pairs.
Although this explanation is appealing, conclusions cannot be made without
further examination. To date, research on the developmental trajectory of pinyin
reading is still lacking. More studies on preliterate and emerging readers with a
longitudinal design are needed.

Nonetheless, these findings indicate impaired automatic grapho-
phonological mapping in dyslexia from the perspective of individual differences.
This anomaly could be underpinned by the altered recruitment of cortical areas
such as the left OTC and bilateral STC. As alphabetic languages, learning to
read in Chinese requires establishing links between visual forms and linguistic
representations (Perfetti & Harris, 2013). Chinese children rely on phonological
mediation in reading comprehension at the earliest stages and later gradually
shift to rapid grapho-semantic processing with a large amount of practice (Zhou
et al., 2017). In this case, deficits in grapho-phonological mapping and
corresponding neural basis can impede the development of the ventral pathway
for rapid character/word recognition and result in reading problems. Recruiting
preliterate children and conducting longitudinal neuroimaging research is
necessary to further examine the causal relationship (Nash et al., 2017).

4.3 Limitations and future directions

This study has several limitations, and caution should be taken when
interpreting the results. First, since we adopted a passive audiovisual
integration paradigm here, we could not directly measure the involvement of
semantic processing in print-sound processing. Second, we administered the
pinyin experiment ahead of the character experiment to reduce the possible
prime effect of characters on processing visually presented pinyin stimuli. This
may influence brain activation to speech sounds in the character experiment
because the same auditory stimuli were used. Third, to have sufficient statistical
power, we used the liberal criteria to assess imaging data quality and exclude
participants with poor quality data accordingly. Forth, while the overall pattern
indicates that children in both groups maintained their attention throughout the
experiment, the dyslexia group performed significantly worse. We controlled
performance IQ in all the analyses to deal with this issue. The results
demonstrated that the main findings of brain-behavior relationships are robust.
At last, we did not have enough cases for looking into different subtypes of
dyslexia. In the future, studies using multiple experimental designs related to
print-sound integration should be conducted with a larger sample size, where a
counterbalance design for estimating the order effect, much stricter criteria for
controlling MRI data quality, strategies for well-matching on in-scanner
performance between groups, and dividing dyslexia into subtypes can be applied.

4.4 Conclusion

The present study explored the impaired audiovisual integration of character-
sound associations and pinyin-sound associations in Chinese children with
dyslexia at the neurofunctional level. The results revealed that dyslexia
manifested an atypical relationship between silent reading comprehension and
the neural integration of character-sounds in the left IFC and between oral
reading fluency and the neural integration of pinyin-sounds in the left OTC and
bilateral STC, providing possible neural substrates underpinning inefficient
grapho-semantic mapping and grapho-phonological mapping, respectively.
Importantly, the current findings also imply that Chinese children with dyslexia
may process pinyin—the alphabetic coding system representing the

pronunciation of characters—in a lagged or deviated way, which can further
impede the development of the direct route for rapid character/word recognition
and semantic access.

5 Author Contributions

725 Conceptualization: ZX, HS, XL; Investigation: ZX, TY, XC, HL, XZ; Formal
726 Analysis: ZX; Data Curation: ZX; Writing – Original Draft Preparation: ZX;
727 Writing – Review & Editing: ZX, TY, XC, FH, HL, XZ, HS, XL; Funding
728 Acquisition: ZX, HS, XL; Supervision: ZX, HS, XL

6 Acknowledgment

731 The authors thank all the participating children, their families, and examiners.

7 Funding

This work was supported by the Key Program of the National Social Science
Foundation of China (14ZDB157), National Key Basic Research Program of
China (grant number 2014CB846103), National Natural Science Foundation of
China grants (grant number 31271082, 31671126, 31611130107, 61374165,
81801782), Beijing Municipal Science & Technology Commission (grant number
Z151100003915122), Fundamental Research Fund for the Central Universities,
China Postdoctoral Science Foundation (grant number 2019T120062,
2018M641235), and Social Science Fund of Beijing (grant number 17YYA004).

8 Declaration of interest

744 The authors declare that there is no conflict of interest.

9 Data availability statement

747 The data that support the findings of this study are available from the
748 corresponding authors upon reasonable request.

10 References

Bi, Y., Han, Z., Weekes, B., & Shu, H. (2007). The interaction between semantic and the nonsemantic systems in reading: Evidence from Chinese. *Neuropsychologia, 45*(12), 2660-2673.

Blau, V., Reithler, J., van Atteveldt, N., Seitz, J., Gerretsen, P., Goebel, R., et al. (2010). Deviant processing of letters and speech sounds as proximate cause of reading failure: a functional magnetic resonance imaging study of dyslexic children. *Brain, 133*(Pt 3), 868-879.

Blau, V., van Atteveldt, N., Ekkebus, M., Goebel, R., & Blomert, L. (2009). Reduced neural integration of letters and speech sounds links phonological and reading deficits in adult dyslexia. *Curr Biol, 19*(6), 503-508.

Blau, V., van Atteveldt, N., Formisano, E., Goebel, R., & Blomert, L. (2008). Task-irrelevant visual letters interact with the processing of speech sounds in heteromodal and unimodal cortex. *Eur J Neurosci, 28*(3), 500-509.

Blomert, L. (2011). The neural signature of orthographic-phonological binding in successful and failing reading development. *Neuroimage, 57*(3), 695-703.

Boets, B., Op de Beeck, H. P., Vandermosten, M., Scott, S. K., Gillebert, C. R., Mantini, D., et al. (2013). Intact but less accessible phonetic representations in adults with dyslexia. *Science, 342*(6163), 1251-1254.

Cao, F., Yan, X., Wang, Z., Liu, Y., Wang, J., Spray, G. J., et al. (2017). Neural signatures of phonological deficits in Chinese developmental dyslexia. *Neuroimage, 146*, 301-311.

Di Folco, C., Guez, A., Peyre, H., & Ramus, F. (2020). Epidemiology of developmental dyslexia: A comparison of DSM-5 and ICD-11 criteria. *medRxiv*, 2020.2012.2018.20248189.

Fedorenko, E., & Blank, I. A. (2020). Broca's Area Is Not a Natural Kind. *Trends Cogn Sci, 24*(4), 270-284.

Glezer, L. S., Eden, G., Jiang, X., Luetje, M., Napoliello, E., Kim, J., et al. (2016). Uncovering phonological and orthographic selectivity across the reading network using fMRI-RA. *NeuroImage, 138*, 248-256.

Guan, C. Q., Fraundorf, S. H., & Perfetti, C. A. (2020). Character and child factors contribute to character recognition development among good and poor Chinese readers from grade 1 to 6. *Annals of Dyslexia, 70*(2), 220-242.

Hagoort, P. (2014). Nodes and networks in the neural architecture for language: Broca's region and beyond. *Curr Opin Neurobiol, 28*, 136-141.

Hancock, R., Richlan, F., & Hoeft, F. (2017). Possible roles for fronto-striatal circuits in reading disorder. *Neurosci Biobehav Rev, 72*, 243-260.

Hoeft, F., McCandliss, B. D., Black, J. M., Gantman, A., Zakerani, N., Hulme, C., et al. (2011). Neural systems predicting long-term outcome in dyslexia. *Proc Natl Acad Sci U S A, 108*(1), 361-366.

Holloway, I. D., van Atteveldt, N., Blomert, L., & Ansari, D. (2015). Orthographic Dependency in the Neural Correlates of Reading: Evidence from Audiovisual Integration in English Readers. *Cerebral Cortex, 25*(6), 1544-1553.

Jednorog, K., Marchewka, A., Altarelli, I., Monzalvo Lopez, A. K., van Ermingen-Marbach, M., Grande, M., et al. (2015). How reliable are gray matter disruptions in specific reading disability across multiple countries and languages? Insights from a large-scale voxel-based morphometry study. *Hum Brain Mapp, 36*(5), 1741-1754.

Jinger, P., Shuang, S., Mengmeng, S., Catherine, M., Hongyun, L., Yuping, Z., et al. (2016). On the relationship between phonological awareness, morphological awareness and Chinese literacy skills: evidence from an 8-year longitudinal study. *Developmental Science, 19*(6), 982-991.

Kronschnabel, J., Brem, S., Maurer, U., & Brandeis, D. (2014). The level of audiovisual print-speech integration deficits in dyslexia. *Neuropsychologia, 62*, 245-261.

Lei, L., Pan, J., Liu, H., McBride-Chang, C., Li, H., Zhang, Y., et al. (2011). Developmental trajectories of reading development and impairment from ages 3 to 8 years in Chinese children. *J Child Psychol Psychiatry, 52*(2), 212-220.

Li, H., Shu, H., McBride-Chang, C., Liu, H., & Peng, H. (2012). Chinese children's character recognition: Visuo-orthographic, phonological processing and morphological skills. *Journal of Research in Reading, 35*(3), 287-307.

Liang, B., & Du, Y. (2018). The Functional Neuroanatomy of Lexical Tone Perception: An Activation Likelihood Estimation Meta-Analysis. Systematic Review. *Frontiers in neuroscience. 12*(495).

Liu, L., Tao, R., Wang, W., You, W., Peng, D., & Booth, J. R. (2013). Chinese dyslexics show neural differences in morphological processing. *Developmental Cognitive Neuroscience, 6*, 40-50.

Liu, L., Wang, W., You, W., Li, Y., Awati, N., Zhao, X., et al. (2012). Similar alterations in brain function for phonological and semantic processing to visual characters in Chinese dyslexia. *Neuropsychologia, 50*(9), 2224-2232.

Liu, Y., Georgiou, G. K., Zhang, Y., Li, H., Liu, H., Song, S., et al. (2017). Contribution of cognitive and linguistic skills to word-reading accuracy

and fluency in Chinese. *International Journal of Educational Research,*
82, 75-90.

Liu, Y., Shu, H., & Li, P. (2007). Word naming and psycholinguistic norms:
Chinese. Behav Res Methods, 39(2), 192-198.

Mei, L., Xue, G., Lu, Z.-L., He, Q., Zhang, M., Wei, M., et al. (2014). Artificial
Language Training Reveals the Neural Substrates Underlying Addressed
and Assembled Phonologies. *PLOS ONE, 9*(3), e93548.

Nash, H. M., Gooch, D., Hulme, C., Mahajan, Y., McArthur, G., Steinmetzger, K.,
et al. (2017). Are the literacy difficulties that characterize developmental
dyslexia associated with a failure to integrate letters and speech sounds?
Developmental science, 20(4).

Oldfield, R. C. (1971). The assessment and analysis of handedness: the
Edinburgh inventory. *Neuropsychologia, 9*(1), 97-113.

Perfetti, C., Cao, F., & Booth, J. (2013). Specialization and Universals in the
Development of Reading Skill: How Chinese Research Informs a Universal
Science of Reading. *Sci Stud Read, 17*(1), 5-21.

Perfetti, C. A., & Harris, L. N. (2013). Universal reading processes are
modulated by language and writing system. *Language Learning and*
Development, 9(4), 296-316.

Pernet, C., Andersson, J., Paulesu, E., & Demonet, J. F. (2009). When all
hypotheses are right: a multifocal account of dyslexia. *Hum Brain Mapp,*
30(7), 2278-2292.

Price, C. J., & Devlin, J. T. (2011). The interactive account of ventral
occipitotemporal contributions to reading. *Trends Cogn Sci, 15*(6), 246-
253.

Richlan, F. (2019). The Functional Neuroanatomy of Letter-Speech Sound
Integration and Its Relation to Brain Abnormalities in Developmental
Dyslexia. *Frontiers in Human Neuroscience, 13*(21).

Richlan, F. (2020). The Functional Neuroanatomy of Developmental Dyslexia
Across Languages and Writing Systems. *Front Psychol, 11*(155), 155.

Richlan, F., Kronbichler, M., & Wimmer, H. (2009). Functional abnormalities in
the dyslexic brain: a quantitative meta-analysis of neuroimaging studies.
Hum Brain Mapp, 30(10), 3299-3308.

Richlan, F., Kronbichler, M., & Wimmer, H. (2011). Meta-analyzing brain
dysfunctions in dyslexic children and adults. *Neuroimage, 56*(3), 1735-
1742.

Ruan, Y., Georgiou, G. K., Song, S., Li, Y., & Shu, H. (2018). Does writing system
influence the associations between phonological awareness, morphological
awareness, and reading? A meta-analysis. *Journal of Educational*
Psychology, 110(2), 180.

Rumsey, J. M., Horwitz, B., Donohue, B. C., Nace, K. L., Maisog, J. M., &
Andreason, P. (1999). A Functional Lesion in Developmental Dyslexia:
Left Angular Gyral Blood Flow Predicts Severity. *Brain and Language,*
70(2), 187-204.

Shah, N. J., Steinhoff, S., Mirzazade, S., Zafiris, O., Grosse-Ruyken, M. L.,
Jancke, L., et al. (2000). The effect of sequence repeat time on auditory
cortex stimulation during phonetic discrimination. *Neuroimage, 12*(1),
100-108.

Shaywitz, S. E. (1998). Dyslexia. *N Engl J Med, 338*(5), 307-312.

Shu, H., McBride-Chang, C., Wu, S., & Liu, H. Y. (2006). Understanding Chinese
developmental dyslexia: Morphological awareness as a core cognitive
construct. *Journal of Educational Psychology, 98*(1), 122-133.

Si, X., Zhou, W., & Hong, B. (2017). Cooperative cortical network for categorical
processing of Chinese lexical tone. *Proc Natl Acad Sci U S A, 114*(46),
12303-12308.

Siok, W. T., Niu, Z., Jin, Z., Perfetti, C. A., & Tan, L. H. (2008). A structural-
functional basis for dyslexia in the cortex of Chinese readers. *Proc Natl*
Acad Sci U S A, 105(14), 5561-5566.

Siok, W. T., Perfetti, C. A., Jin, Z., & Tan, L. H. (2004). Biological abnormality of
impaired reading is constrained by culture. *Nature, 431*(7004), 71-76.

Song, S., Zhang, Y., Shu, H., Su, M., & McBride, C. (2020). Universal and
Specific Predictors of Chinese Children With Dyslexia – Exploring the
Cognitive Deficits and Subtypes. *Frontiers in Psychology, 10*(2904).

Stevens, W. D., Kravitz, D. J., Peng, C. S., Tessler, M. H., & Martin, A. (2017).
Privileged Functional Connectivity Between the Visual Word Form Area
and the Language System. *Journal of Neuroscience, 37*(21), 5288-5297.

Su, M., Zhao, J., Thiebaut de Schotten, M., Zhou, W., Gong, G., Ramus, F., et al.
(2018). Alterations in white matter pathways underlying phonological and
morphological processing in Chinese developmental dyslexia. *Dev Cogn*
Neurosci, 31, 11-19.

Thiebaut de Schotten, M., Cohen, L., Amemiya, E., Braga, L. W., & Dehaene, S.
(2014). Learning to read improves the structure of the arcuate fasciculus.
Cereb Cortex, 24(4), 989-995.

Tschentscher, N., Ruisinger, A., Blank, H., Diaz, B., & von Kriegstein, K. (2018). Reduced structural connectivity between left auditory thalamus and the motion-sensitive planum temporale in developmental dyslexia. *arXiv preprint arXiv:1811.11658*.

van Atteveldt, N., & Ansari, D. (2014). How symbols transform brain function: A review in memory of Leo Blomert. *Trends in Neuroscience and Education, 3*(2), 44-49.

van Atteveldt, N., Formisano, E., Goebel, R., & Blomert, L. (2004). Integration of letters and speech sounds in the human brain. *Neuron, 43*(2), 271-282.

van Atteveldt, N. M., Formisano, E., Blomert, L., & Goebel, R. (2007). The effect of temporal asynchrony on the multisensory integration of letters and speech sounds. *Cereb Cortex, 17*(4), 962-974.

van Atteveldt, N. M., Formisano, E., Goebel, R., & Blomert, L. (2007). Top–down task effects overrule automatic multisensory responses to letter–sound pairs in auditory association cortex. *NeuroImage, 36*(4), 1345-1360.

Vandermosten, M., Boets, B., Poelmans, H., Sunaert, S., Wouters, J., & Ghesquiere, P. (2012). A tractography study in dyslexia: neuroanatomic correlates of orthographic, phonological and speech processing. *Brain, 135*(Pt 3), 935-948.

Wang, F., Karipidis, II, Pleisch, G., Fraga-Gonzalez, G., & Brem, S. (2020). Development of Print-Speech Integration in the Brain of Beginning Readers With Varying Reading Skills. *Front Hum Neurosci, 14*(289), 289.

Wechsler, D. (1974). *WISC-R, Wechsler intelligence scale for children, revised*: Psychological Corporation.

Wenger, E., Brozzoli, C., Lindenberger, U., & Lövdén, M. (2017). Expansion and Renormalization of Human Brain Structure During Skill Acquisition. *Trends in Cognitive Sciences, 21*(12), 930-939.

Xia, M., Wang, J., & He, Y. (2013). BrainNet Viewer: a network visualization tool for human brain connectomics. *PloS one, 8*(7), e68910.

Xia, Z., Hoeft, F., Zhang, L., & Shu, H. (2016). Neuroanatomical anomalies of dyslexia: Disambiguating the effects of disorder, performance, and maturation. *Neuropsychologia, 81*, 68-78.

Xia, Z., Yang, T., Cui, X., Hoeft, F., Liu, H., Shu, H., et al. (2020). Neurofunctional mechanisms underlying audiovisual integration of characters and *pinyin* in Chinese children. *bioRxiv*, 2020.2005.2031.126128.

Xia, Z., Zhang, L., Hoeft, F., Gu, B., Gong, G., & Shu, H. (2018). Neural correlates of oral word reading, silent reading comprehension, and

cognitive subcomponents. *International Journal of Behavioral*
Development, 42(3), 342-356.

Xu, W., Kolozsvari, O. B., Oostenveld, R., Leppanen, P. H. T., & Hamalainen, J.
A. (2019). Audiovisual Processing of Chinese Characters Elicits
Suppression and Congruency Effects in MEG. *Front Hum Neurosci,*
13(18), 18.

Xue, J., Shu, H., Li, H., Li, W., & Tian, X. J. J. o. P. R. (2013). The Stability of
Literacy-Related Cognitive Contributions to Chinese Character Naming
and Reading Fluency. *42*(5), 433-450.

Yang, J., Shu, H., McCandliss, B. D., & Zevin, J. D. (2013). Orthographic
influences on division of labor in learning to read Chinese and English:
Insights from computational modeling. *Biling (Camb Engl), 16*(Spec Iss 2),
354-366.

Zhang, L., Xi, J., Xu, G., Shu, H., Wang, X., & Li, P. (2011). Cortical dynamics of
acoustic and phonological processing in speech perception. *PLoS One, 6*(6),
e20963.

Zhang, Y., Zhang, L., Shu, H., Xi, J., Wu, H., Zhang, Y., et al. (2012).
Universality of categorical perception deficit in developmental dyslexia: an
investigation of Mandarin Chinese tones. *J Child Psychol Psychiatry,*
53(8), 874-882.

Zhao, J., Wang, X., Frost, S. J., Sun, W., Fang, S.-Y., Mencl, W. E., et al. (2014).
Neural division of labor in reading is constrained by culture: A training
study of reading Chinese characters. *Cortex, 53*, 90-106.

Zhou, W., Shu, H., Miller, K., & Yan, M. (2018). Reliance on orthography and
phonology in reading of Chinese: A developmental study. Journal of
Research in Reading, 41(2), 370-391.

Zou, L., Packard, J. L., Xia, Z., Liu, Y., & Shu, H. (2015). Neural Correlates of
Morphological Processing: Evidence from Chinese. *Frontiers in human*
neuroscience, 9.
967

Figure legends

Figure 1 Schematic description of the fMRI experimental procedure (A, B, C for a run, a block, and a mini-block, respectively) and stimuli examples (D). Abbreviations: Aud = auditory, avC = audiovisual congruent, avI = audiovisual incongruent, Fix = fixation, Vis = visual.

Figure 2 (A) Results of the voxel-wise whole-brain ANOVA on the character conditions. Only clusters showing a significant main effect of condition were identified at the FWE corrected threshold of p-cluster < 0.05 (p-voxel < 0.001 for height). No region showed significant main effect of group or group × condition interaction. (B) The bar plots present brain activation in the audiovisual congruent and incongruent conditions for the control and dyslexic groups. (C) The scatterplots display correlations between the neural audiovisual integration effect (avC-avI) in the left IFC with silent reading comprehension proficiency and related cognitive-linguistic skills (morphological awareness, rapid naming). Abbreviations: avC = audiovisual congruent, avI = audiovisual incongruent, IFC = inferior frontal cortex, STC = superior temporal cortex.

Figure 3 (A) Brain map of regions showing significant between-group differences in the correlation between the audiovisual integration effect and oral reading fluency (FWE corrected p-cluster < 0.05, p-voxel < 0.001 for height). The scatterplots presents correlations in each group (blue: control; red: dyslexic) in the left OTC (B), STC (C), and right STC (D). Abbreviations: avC = audiovisual congruent, avI = audiovisual incongruent, OTC = occipito-temporal cortex, STC = superior temporal cortex.

Supplementary Figure 1 The summary of demographic and behavioral measures is shown as boxplots, with the box indicating the IQR. The whiskers show the range of values within 1.5 × IQR and a horizontal line indicating the median. Individual data are shown as dots. The color coding is indicated in the legend below to the plot. Data visualization was performed with PlotsOfData (https://huygens.science.uva.nl/PlotsOfData/). Abbreviations: IQR = interquartile range.

Table 1 Demographic and behavioral profiles

Measures	Typical Readers (n = 22)			Reading Disorder (n = 23)			Comparison	
	Mean	SD	Range	Mean	SD	Range	t/X^2	p-value
Age (month)	127	6	118 ~ 140	122 †	10	111 ~ 144	1.938	0.061
Sex (female/male)	12/10			10/13			0.552	0.556
Verbal IQ (standard score)	109	12	88 ~ 139	97	11	67 ~ 118	3.668	< 0.001
Performance IQ (standard score)	115	13	82 ~ 137	105	12	82 ~ 135	2.512	0.016
Full-Scale IQ (standard score)	113	11	93 ~ 130	101	9	84 ~ 116	3.986	< 0.001
Character Recognition (item)	122	8	107 ~ 140	86 †	11	53 ~ 99	11.830	< 0.001
Standard score	1.38	0.66	0.41 ~ 2.43	-2.18	0.83	-4.520 ~ -1.020	12.701	< 0.001
Word List Reading (word/minute)	94 ‡	16	72 ~ 140	63	11	40 ~ 83	7.612	< 0.001
Standard score	0.95 †	1.11	-0.38 ~ 4.06	-1.584	0.67	-2.930 ~ -0.130	7.278	< 0.001
Silent Reading Comprehension (character/minute)	361 ‡	104	220 ~ 565	163	56	70 ~ 304	7.890	< 0.001
Standard score	0.98 ‡	1.04	-0.38 ~ 3.12	-1.083	0.57	-2.250 ~ 0.430	7.998	< 0.001
Phoneme Deletion (item)	21 †	4	7 ~ 25	17 †	6	1 ~ 25	2.848	0.007
Rapid Naming (second)	17	2	11 ~ 20	22	4	15 ~ 30	-5.481	< 0.001
Morphological Production (item)	24 ‡	3	20 ~ 29	22	3	13 ~ 28	2.927	0.005

Note. † Shapiro-Wilk p < 0.05, ‡ 0.05 < Shapiro-Wilk p < 0.01

Table 2 Significant clusters in the voxel-wise whole-brain analyses

Experiment and contrast	Label	Brain area	$P_{\text{FWE-corrected}}$	Size	Peak F	X	Y	Z
Character								
Main effect of condition	LSTC	Left middle and superior temporal	0.007	391	26.86	-60	-32	6
	LIFC	Left inferior frontal gyrus,	0.024	281	23.90	-46	16	14
Pinyin								
Group difference in correlation	LOTC	Inferior and middle occipital gyri,	0.004	358	29.28	-44	-72	-2
Brain: avC-avI	LSTC	Left middle and superior temporal	0.039	197	23.02	-52	-44	10
Behavior: *Oral reading fluency*	RSTC	Right superior temporal gyrus,	0.047	185	38.68	62	-8	4

Note. The significant clusters were identified with the FWE-corrected threshold of *p*-cluster < 0.05 (*p*-voxel < 0.001 for height). Brain area labeling is based on the AAL atlas. Cluster size refers to the number of voxels.

Table 3 Brain-behavior correlations in the significant clusters in the voxel-wise whole-brain regression analyses.

Behavior:	Group	Integration Effect (avC-avI)		avC		avI	
		r-value	*p*-value	*r*-value	*p*-value	*r*-value	*p*-value
Oral reading fluency							
L OTC	Control	0.525 *	0.021	0.685 **	0.001	0.118	0.631
	Dyslexic	-0.804 ***	< 0.001	0.026	0.913	0.674 **	0.001
L STC	Control	0.693 **	0.001	0.588 **	0.008	-0.014	0.956
	Dyslexic	-0.668 **	0.001	-0.050	0.833	0.445 *	0.049
R STC	Control	0.655 **	0.002	0.240	0.321	-0.259	0.284
	Dyslexic	-0.706 ***	< 0.001	-0.181	0.444	0.543 *	0.013

Note. Age, sex, and performance IQ were controlled statistically. Abbreviations: avC = audiovisual congruent, avI = audiovisual incongruent. *** $p < 0.001$, ** $p < 0.01$, * $p < 0.05$

1000

A

Left STC

Left IFC

F-value

+6

Character: main effect of condition

B

LSTC

LIFC

avC   avI

avC   avI

■ **Control**   ■ **Dyslexic**

C

Left IFC-Reading Comprehension

Left IFC-Cognitive Skills

Control

avC-avI (residuals)

Silent reading comprehension (residuals)

avC-avI: *r* = 0.571, *p* = 0.011 *
- avC: *r* = 0.222, *p* = 0.360
- **avI:** *r* = -0.526, *p* = 0.021 *

avC-avI (residuals)

Morphological awareness (residuals)

avC-avI: *r* = 0.551, *p* = 0.014 *
- **avC:** *r* = 0.828, *p* < 0.001 ***
- avI: *r* = 0.102, *p* = 0.679

Dyslexic

avC-avI (residuals)

Silent reading comprehension (residuals)

avC-avI: *r* = 0.391, *p* = 0.088 †
- avC: *r* = 0.400, *p* = 0.080 †
- avI: *r* = -0.185, *p* = 0.436

avC-avI (residuals)

Rapid naming (residuals)

avC-avI: *r* = -0.509, *p* = 0.022 *
- avC: *r* = -0.400, *p* = 0.081 †
- avI: *r* = 0.340, *p* = 0.143

A

Left STC

Left OTC

Right STG

Pinyin: group difference on brain-behavior correlation

B **Left OTC-Oral Reading**

■ **Control**
■ **Dyslexic**

avC-avI (residuals)

Oral reading fluency (residuals)

Control:
r = 0.525, *p* = 0.021 *

Dyslexic:
r = -0.804, *p* < 0.001 ***

C **Left STC-Oral Reading**

avC-avI (residuals)

Oral reading fluency (residuals)

Control:
r = 0.693, *p* = 0.001 ***

Dyslexic:
r = -0.668, *p* = 0.001 ***

D **Right STC-Oral Reading**

avC-avI (residuals)

Oral reading fluency (residuals)

Control:
r = 0.655, *p* = 0.002 **

Dyslexic:
r = -0.707, *p* < 0.001 ***