BeeDNA: microfluidic environmental DNA metabarcoding as a tool for connecting plant and pollinator communities

Lynsey R. Harper[1,2], Matthew L. Niemiller[3], Joseph B. Benito[3], Lauren E. Paddock[4,5], E. Knittle[4], Brenda Molano-Flores[1], and Mark A. Davis[1*]

[1] Illinois Natural History Survey, Prairie Research Institute, University of Illinois at Urbana-Champaign, Champaign IL 61820, USA

[2] NatureMetrics Ltd, 1 Occam Court, Surrey Research Park, Guildford, GU2 7HJ, UK

[3] Department of Biological Sciences, College of Science, The University of Alabama in Huntsville, Huntsville AL 35889, USA

[4] Department of Animal Biology, University of Illinois Urbana-Champaign, Urbana IL 61801, USA

[5] Duchossois Family Institute, Knapp Center for Biomedical Discovery Biological Sciences Division, The University of Chicago, Chicago IL 60637, USA

*Corresponding author:

Email: davis63@illinois.edu

Running title: eDNA survey of pollinator communities

24 **Abstract**

25

26 Pollinators are imperiled by global declines that can reduce plant reproduction, erode essential

27 ecosystem services and resilience, and drive economic losses. Monitoring pollinator biodiversity

28 trends is key for adaptive conservation and management, but conventional surveys are often

29 costly, time consuming, and require taxonomic expertise. Environmental DNA (eDNA)

30 metabarcoding surveys are booming due to their rapidity, non-invasiveness, and cost efficiency.

31 Microfluidic technology allows multiple primer sets from different markers to be used in eDNA

32 metabarcoding for more comprehensive species inventories whilst minimizing biases associated

33 with individual primer sets. We evaluated microfluidic eDNA metabarcoding for pollinator

34 community monitoring by introducing a bumblebee colony to a greenhouse flower assemblage

35 and sampling natural flower plots. We collected nectar draws, flower swabs, or whole flower

36 heads from four flowering species, including two occurring in both the greenhouse and field.

37 Samples were processed using two eDNA isolation protocols before amplification with 15

38 primer sets for two markers (COI and 16S). Microfluidic eDNA metabarcoding detected the

39 target bumblebee and greenhouse insects as well as common regional arthropods. Pollinator

40 detection was maximized using whole flower heads preserved in ATL buffer and extracted with

41 a modified Qiagen® DNeasy protocol for amplification with COI primers. eDNA surveillance

42 could enhance pollinator assessment by detecting protected and endangered species and being

43 more applicable to remote, inaccessible locations, whilst reducing survey time, effort, and

44 expense. Microfluidic eDNA metabarcoding requires optimization to address remaining efficacy

45 concerns but this approach shows potential in revealing complex networks underpinning critical

46 ecosystem functions and services, enabling more accurate assessments of ecosystem resilience.

Keywords: arthropods, community ecology, ecosystem assessment, eDNA, mutualistic

interactions, pollinators

1. Introduction

Biodiversity is being lost at an alarming rate across habitats, ecosystems, and geopolitical

boundaries (Barnosky et al., 2011; Wake & Vredenburg, 2008). Current extinction rates exceed

those of the last five mass extinction events, with the sixth mass extinction event being driven by

ever-growing anthropogenic activities (Ehlers & Krafft, 2006). The Anthropocene has been

particularly detrimental to pollinators (Cameron et al., 2011; Hallmann et al., 2017; Lever et al.,

2014; Macgregor et al., 2019; Potts et al., 2010), leading to numerous pollinator species being

listed or petitioned for listing in the United States Federal Register (81 FR 67786, 82 FR 10285)

as threatened or endangered. Pollinators are of vital importance to North American ecosystems

and economies (Losey & Vaughan, 2006; Southwick & Southwick, 1992). The loss of pollinator

biodiversity is of substantial concern as it reduces plant reproductive success (Klein et al., 2007;

Ollerton et al., 2011; Thomann et al., 2013), erodes ecosystem services provided by pollination

(Ollerton et al., 2011; Vanbergen & the Insect Pollinators Initiative, 2013; Winfree et al., 2011),

and ultimately drives staggering economic losses (Gallai et al., 2009; Kevan & Phillips, 2001;

Klein et al., 2007). Therefore, rapid, efficient, and accurate assessment of pollinator communities

is a conservation imperative to inform adaptive management strategies to mitigate or prevent

further loss of pollinator biodiversity (Carvalheiro et al., 2013; Didham et al., 2020; Montgomery

et al., 2020; Thomas et al., 2019).

Such assessments are both technically difficult and financially costly (Bartomeus, 2013;

Didham et al., 2020; Plein et al., 2017). Conventional monitoring is often restricted in terms of

comparing multiple communities, requires high time investment to document presence or

absence of pollinator species, and is reliant on dwindling taxonomic expertise to identify

observed pollinators (Sheffield et al., 2009; Tur et al., 2013; Weiner et al., 2014). Moreover,

monitoring the overwhelming majority of pollinator species is unrealistic (Didham et al., 2020;

Young et al., 2017). These challenges are likely exacerbated for threatened and endangered

species, whose populations tend to be numerically small and patchily distributed (Matthies et al.,

2004). Yet, pollinator assessment is critical to monitor status and trends of rare, threatened and

endangered pollinators, and evaluate the impacts of biodiversity loss on ecosystem services

(Carvalheiro et al., 2013; Montgomery et al., 2020). Approaches that can estimate pollinator

biodiversity in both a time-efficient and cost-effective manner will drastically enhance the

development of effective interventions to mitigate or prevent further losses (Didham et al.,

2020).

Environmental DNA (eDNA) analysis has emerged as a viable candidate for rapid, non-

invasive biodiversity assessment (Deiner et al., 2017; Taberlet et al., 2012). Organisms deposit

genetic material into their environment (e.g., via secretions and excretions) that can be harnessed

to identify single species or entire communities in freshwater (Biggs et al., 2015; Broadhurst et

al., 2021), marine (K. J. Harper et al., 2020; Valsecchi et al., 2021) and terrestrial (Clare et al.,

2021; Katz et al., 2020) contexts. Single-species eDNA assays have been increasingly leveraged

as a reliable means of surveying rare or invasive species (Thalinger et al., 2021). However, the

costs associated with single-species assay development, validation and application, coupled with

increasing numbers of imperiled species requiring monitoring (especially pollinators),

necessitates methods that can simultaneously screen for multiple taxa, increasing efficiencies and

reducing financial burdens (L. R. Harper et al., 2018; Moss et al., 2022; Wilcox et al., 2020).

eDNA metabarcoding may provide a solution for more comprehensive and less costly

assessment of invertebrate communities across larger spatial and temporal scales (L. R. Harper et

al., 2020; Mächler et al., 2019; Roger et al., 2021).

eDNA metabarcoding is based on PCR amplification of short DNA fragments using

primers with conserved binding sites across multiple taxa, flanking a region of highly variable

DNA sequence that enables taxa to be distinguished from each other, followed by high-

throughput sequencing (Deiner et al., 2017; Taberlet et al., 2012). The mitochondrial cytochrome

c oxidase subunit I (COI) locus is the standard genetic marker for DNA barcoding of metazoans

due to high variability among species (enabling high taxonomic resolution) and extensive

reference database representation (Elbrecht et al., 2018). However, this variability has proven

problematic for metabarcoding due to a paucity of conserved primer binding sites (Clarke et al.,

2017; Leese et al., 2020). Degenerate primers can counter this issue but have inherent risks of

primer mismatch, primer slippage, non-target amplification (e.g., bacteria, algae and fungi), and

binding of non-target regions. All of these issues can introduce bias, complicating bioinformatic

processing and taxonomic assignment (Clarke et al., 2017; Elbrecht et al., 2016, 2018; Leese et

al., 2020).

Using different markers (e.g., 16S ribosomal RNA) or COI in conjunction with other

markers has been investigated, but reference databases for other markers are comparatively

sparse, limiting taxonomic resolution, and use of multiple markers increases PCR and

sequencing costs (Clarke et al., 2017; Elbrecht et al., 2016; Elbrecht et al., 2017). Furthermore,

good *in silico* performance of individual primer sets may not translate to *in situ* success, with

performance also varying depending on the desired level of taxonomic resolution (Corse et al.,

2019). Using multiple primer sets targeting the same marker has been proposed to combat this

discrepancy in performance (Corse et al., 2019; Polanco F. et al., 2021), but does not avoid

increased PCR and sequencing costs. Microfluidic technology may offer a solution by enabling

PCR amplification with multiple primer sets targeting different taxa, different loci, or different

regions of the same locus simultaneously, with all amplicons then being sequenced in single run

(Brown et al., 2016). Hauck et al. (2019) demonstrated the potential of microfluidic eDNA

metabarcoding to elucidate broader communities, provide high taxonomic resolution, and

scrutinize performance of individual primer sets.

To our knowledge, only one study has evaluated eDNA metabarcoding of flowers as an

approach to document pollinator communities (Thomsen & Sigsgaard, 2019). The authors

successfully detected a wide range of terrestrial arthropods, including at least 135 species, but

also highlighted issues of primer choice (despite pairing a COI primer set with a 16S primer set)

and reference database representation. Our study builds upon this work using a combinatorial

experimental design under both controlled laboratory and field conditions that tests the influence

of flower species, sample type, and eDNA isolation protocol on pollinator communities detected

by microfluidic eDNA metabarcoding. Based on our results, we propose a workflow that should

enhance eDNA detection of pollinator communities and facilitate further research in this field.

2. Materials and methods

2.1 Greenhouse experiment

A mixed-species flowering plant assemblage was established in the University of Illinois Urbana-Champaign (UIUC) Plant Care Facility (Table S1; Figure 1a), hereafter greenhouse. Plantings were staggered so that a constant supply of pollen and nectar were available. A subset of flowering plants was moved to a secure room where a Natupol colony (Koppert Biological Systems) of common eastern bumblebees (*Bombus impatiens*) was released and allowed to acclimate for 10 days (Figure 1b). Temperature was set at 21-24°C during the day and 12-16°C during the night, humidity was ambient, and light was set to 14-hr day length.

We selected four focal flower species with different flower morphologies to separately introduce to the secure room over two months: *Penstemon* sp., *Monarda* sp., *Solanum* sp., and *Cynoglossum amabile* (Figure 1c-e). Prior to the introduction of a flower species, flowers were sampled using up to three methods to provide a detection baseline for other insect species present in the greenhouse (baseline controls): 1) whole flower heads were harvested (*Penstemon*, *Monarda*, and *Solanum*), 2) the flower was swabbed with a flocked cotton swab (*Penstemon* and *Monarda* only), and 3) nectar was extracted from the flower via microcapillary tube (*Cynoglossum amabile* only). All three sampling approaches could not be applied to all four flower species because *Solanum* does not produce nectar and attempted nectar draws from *Penstemon* and *Monarda* were unsuccessful. Samples were preserved in Qiagen® ATL buffer or cetyl trimethyl ammonium bromide (CTAB; Teknova) in sterile 2 mL microcentrifuge tubes (MidWest Scientific). Volumes of 300, 600, 900 or 1200 µL were used depending on sample size/type (see Table S2).

Upon introducing a flower species, we observed the flowers for bumblebee visits. After a

visit by a bumblebee and any handling was observed, the flower was swabbed (*Penstemon* and

Monarda only). A total of 50 flowers (one flower per plant) were sampled in one day, with 25

each preserved in ATL buffer and CTAB in sterile 2 mL microcentrifuge tubes. Flowers were

left unobserved in the greenhouse for 24 hrs, then an approximately equal number of flowers

from each of three benches were swabbed unsystematically to assess the ability to detect

pollinator visits in the absence of direct observation. Again, 50 flowers (one flower per plant)

were sampled in one day and divided equally between the two preservation buffers. This two-day

process was repeated for harvesting of whole flower heads (*Penstemon*, *Monarda*, and *Solanum*),

then nectar extraction from flowers via microcapillary tube (*Cynoglossum amabile* only). On

each day that a flower species was sampled, sterile 2 mL microcentrifuge tubes containing 1200

μL of ATL buffer and CTAB were taken into the secure room but no sample was added. These

tubes were transported, stored, and processed alongside samples (field blanks).

2.2 Field experiment

Two of the four focal flower species (*Penstemon* and *Monarda*) in the greenhouse also occurred

on the UIUC campus enabling a pilot field component to the greenhouse experiment using a

similar sampling approach. We concentrated field sampling effort on *Penstemon* first, followed

by *Monarda* due to differences in timing of peak flowering. Flowers were observed between 9am

and 12pm on days when the weather was suitable (sunny with temperatures >13°C). When a bee

species visited a flower, the bee was identified (https://beespotter.org/) and a photo was taken.

After the bee left the flower, the flower was swabbed. A total of 50 flowers (one flower per

plant) were sampled in one day, with 25 each preserved in ATL buffer and CTAB in sterile 2 mL

microcentrifuge tubes. On the next suitable day, 50 unsystematic swabs (i.e., no observation of

pollinator visits) were acquired (one flower per plant) and divided equally between the two

preservation buffers. This process was then repeated for harvesting of whole flower heads. Field

blanks were included on each day that sampling occurred.

2.3 DNA extraction, quantification and plating

DNA extraction was performed at the Collaborative Ecological Genetics Laboratory, Illinois

Natural History Survey, UIUC, where dedicated labs and established procedures for working

with environmental samples of low DNA concentration are in place. Procedures include regular

bleaching (50% v/v solution) of work surfaces, ultraviolet (UV) irradiation of reagents,

consumables, and workspaces, isolated PCR hoods, and physical separation of pre-PCR (i.e., a

dedicated eDNA "clean room" where no high-copy DNA has been introduced) and post-PCR

processes.

A modified Qiagen® DNeasy® Blood & Tissue (QBT) protocol (Thomsen & Sigsgaard,

2019) was used for extraction of samples preserved in ATL buffer (see Supporting Information).

Lysis was conducted in the tubes containing samples by adding 60, 100, 200, or 300 μL of

proteinase K respectively depending on the volume of preservative used (Table S2). DNA was

eluted in 2×60 μL of AE buffer, with a 15 min incubation step at 37°C each time before

centrifugation. A modified Phenol-Chloroform-Isoamyl (PCI) extraction and ethanol

precipitation method (Renshaw et al., 2015) was employed for samples preserved in CTAB (see

Supporting Information). DNA was rehydrated with 100 μL of 1x TE buffer (ThermoFisher

Scientific). Buffer-only extraction blanks were included for each batch of extractions ($n = 23$).

DNA extracts were stored at -20°C until quantification and plating.

Samples extracted with both protocols were quantified using a Qubit™ 3.0 Fluorometer

(Life Technologies, Grand Island, NY) with the Qubit™ dsDNA HS Assay Kit (ThermoFisher

Scientific). Three readings were taken for each sample with the average used as the final

concentration (ng/µL). We aliquoted 30 µL of each sample into a 96-well 0.2 mL optical reaction

plate (ThermoFisher Scientific). A total of 42 baseline controls/eDNA samples, two field blanks,

two extraction blanks, and two PCR positive controls (one of ten bee species known to occur in

Illinois, listed in Figure 2 excluding *B. griseocollis*) were included on each plate. Plates were

stored at -20°C until submission (typically less than 24 hrs) for microfluidic eDNA

metabarcoding (see section 2.5).

2.4 Primer selection and validation

We conducted a comprehensive literature review to identify suitable primer sets for arthropod

eDNA metabarcoding. We refined this list of primers to those with Fluidigm compatible

annealing temperatures (~55°C) that amplified a 150-400 bp fragment (lower bound to meet

post-amplification removal of primer-adapted dimers, upper bound to limit amplicon length for

paired-end sequencing) for microfluidic eDNA metabarcoding.

For *in silico* primer validation, we constructed a reference sequence database for

invertebrate species that occur in Illinois using a list of binomial names based on observations by

greenhouse staff, light-trapping data from the Illinois Bat Conservation Program (IBCP),

arthropods and molluscs catalogued in the Illinois Natural History Survey (INHS) Insect

Collection Database (http://inhsinsectcollection.speciesfile.org/InsectCollection.aspx), cicadas

recorded by Catherine Dana (INHS), and threatened and endangered priorities provided by

Angella Moorehouse (Illinois Nature Preserves Commission). With this list, all available

sequences for markers amplified by the selected primer sets were downloaded from GenBank

using Reprophylo v1.3 (Szitenberg et al., 2015) in July 2019. Reference sequences in GenBank

format were converted to ecoPCR (Ficetola et al., 2010) database format using obiconvert

(Boyer et al., 2016). ecoPCR parameters were set to allow a 50–500 bp fragment and up to 3

mismatches between each primer and each sequence in the reference database.

For *in vitro* primer validation, DNA extracts from 11 bee species known to occur in

Illinois were amplified with the selected primer sets. (Figure 2). PCR was performed in triplicate

using reaction volumes of 25 μL, including 12.5 μL of sterile molecular grade water

(MidwestScientific), 11.1 μL of GoTaq® Colorless Master Mix (ThermoFisher Scientific), 0.2

μL of forward primer and 0.2 μL of reverse primer (10 μM, Integrated DNA Technologies), and

1 μL of template DNA. Thermocycling conditions were 5 min at 94°C, followed by 40 cycles of

45 sec at 94°C, 1 min at 55°C and 1 min 30 sec at 72°C, then 5 min at 72°C. PCR amplification

was confirmed using agarose gel electrophoresis (2 g agarose (Midwest Scientific), 100 mL 1x

TAE buffer (Midwest Scientific)) at 120 V for 90 min, and gels were visualized with a LabNet

Systems EnduroGel Gel Documentation System. Amplification was classed as successful (3 of 3

PCR replicates), moderately successful (1 or 2 of 3 PCR replicates), or unsuccessful (0 of 3 PCR

replicates).

2.5 Microfluidic eDNA metabarcoding

Microfluidic eDNA metabarcoding, including PCR amplification, Illumina sequencing, and

demultiplexing using QIIME 2™ (Bolyen et al., 2019), was conducted by the Roy J. Carver

Biotechnology Center Functional Genomic Unit at the UIUC. The Fluidigm 48.48 Access

Array™ (Fluidigm Corporation, 2016) was used for PCR amplification of each eDNA sample (2

ng) once with our validated primer panel (see Supporting Information). Products were quantified

on a Qubit™ Fluorometer and stored at -20°C. All samples were run on a Fragment Analyzer

(Advanced Analytics) and amplicon regions and expected sizes confirmed. Samples were then

pooled in equal amounts according to product concentration. The pooled products were size

selected on a 2% agarose E-gel (Life Technologies) and extracted from the isolated gel slice with

a QIAquick Gel Extraction Kit (Qiagen). Purified products were run on an Agilent Bioanalyzer

to confirm appropriate amplicon profile and determination of average size. The products were

quantified using a Qubit™ Fluorometer, pooled evenly, and the final libraries were quantified by

qPCR on a BioRad CFX Connect Real-Time System (Bio-Rad Laboratories, Inc.). The libraries

were denatured, spiked with 20% PhiX Control v3, and loaded at 8 pM to an Illumina MiSeq for

cluster formation and sequencing with a MiSeq Reagent Kit v3 (600-cycle) (Illumina, Inc.). In

total, 40 baseline controls, 336 eDNA samples, 18 field blanks, 18 extraction blanks, two PCR

negative controls, and 18 PCR positive controls were sequenced across two runs (288

samples/controls and 144 samples/controls, respectively). Sequences were sorted and

demultiplexed by primer set and sample, then quality and yield assessed using the FastX Tool

Kit (http://hannonlab.cshl.edu/fastx_toolkit/index.html).

2.6 Bioinformatic processing

The demultiplexed FASTQ files for each primer set were processed using metaBEAT v0.97.11

(https://github.com/HullUnibioinformatics/metaBEAT) and the Anacapa Toolkit v1 (Curd et al.,

2019), which has been deposited on GitHub (https://github.com/limey-bean/Anacapa) and

permanently archived (http://doi.org/10.5281/zenodo.3064152). Each pipeline incorporates a

different suite of open-source software for bioinformatic processing (see Supporting

Information), which allowed us to assess the robustness of eDNA detections. metaBEAT

produced comparable or better detection than Anacapa for our focal pollinator species and

species known to occur in the greenhouse (see Supporting Information and Figure S1), thus we

used these results for downstream analyses.

2.7 Data analysis

All data manipulation and downstream analyses were conducted in R v3.6.3 (R Core Team,

2020). Datasets for each primer set were combined into a single master dataset, retaining primer

information, and detections in eDNA samples and process controls were assessed (Figures S2,

S3). Taxon richness within samples produced by each primer pair and taxon overlap between

different primer pairs was assessed (Figures S4, S5) before removing primer information and

merging reads for each taxon. A highly conservative false positive sequence threshold was

calculated based on the maximum frequency of arthropod DNA (2.05%) found in the negative

process controls (i.e., field blanks, extraction blanks, and PCR negative controls) and applied to

the metabarcoding data. PCR positive controls were not used for threshold determination as

DNA was isolated from bumblebee specimens sourced from pinned museum collections

resulting in high contamination frequency, and the species that specimens belonged to could

naturally occur in our study system (see Discussion). The data were filtered to remove non-

arthropod taxonomic assignments before taxonomic assignments were filtered on a sample-by-

sample basis to retain the lowest taxonomic ranks where assignments to multiple taxonomic

ranks were present (e.g., species, genus and family). The filtered data were then combined with

greenhouse and field metadata.

We examined taxa detected in the baseline controls before flower species were

introduced to the secure room containing common eastern bumblebees (Figure S6), in eDNA

samples after flower species were introduced to the secure room (Figure S7), and in eDNA

samples taken from flowers around the UIUC campus (Figure S8). We then investigated

biological and technical grouping variables that potentially influence pollinator eDNA detection:

flower species, sample type, and eDNA isolation protocol. The read count data were converted to

binary data as potential bias introduced by PCR amplification may prevent reliable abundance or

biomass estimation from sequence reads produced by DNA or eDNA metabarcoding (Elbrecht et

al., 2017).

First, alpha diversity (i.e., taxon richness) of eDNA samples according to each grouping

variable was compared. The data were non-normally distributed and the assumptions of a one-

way Analysis of Variance (ANOVA) were violated (Table S5, Figures S9, S10, S11), which

Generalised Linear Models (GLMs) with different error families and link-functions did not

resolve. Therefore, we compared taxon richness according to each grouping variable using the

non-parametric Kruskal-Wallis test from the package stats v3.6.3, and performed multiple

pairwise comparisons using the non-parametric Dunn's test (p-values adjusted with the

Benjamini-Hochberg method where applicable) from the package FSA v0.9.1 (Ogle et al., 2020).

The package betapart v1.5.4 (Baselga & Orme, 2012) was used to estimate total beta

diversity, partitioned by nestedness-resultant (i.e., community dissimilarity due to taxon subsets)

and turnover (i.e., community dissimilarity due to taxon replacement), across all samples with

the *beta.multi* function. Greenhouse samples 80PEWF7-P2-G3, 80MOSW09-P3-C3,

80PEWF23-P1-G4 and 80PEWF73-P9-C5 and UIUC campus samples 80PESW105-P5-G2 and

80PEWF37-P2-A3 skewed visualisation of beta diversity components, thus these samples were

removed from the dataset prior to performing these analyses. The three components of beta

diversity (Jaccard dissimilarity) were estimated for eDNA samples according to each grouping

variable using the *beta.pair* function. For each component of beta diversity, we compared the

variance in each group of samples by calculating homogeneity of multivariate dispersions

(MVDISP) using the *betadisper* function in the package vegan v2.5-7 (Oksanen et al., 2019).

Differences in MVDISP between groups of eDNA samples were statistically tested using an

ANOVA. Community dissimilarity for each component of beta diversity was visualized using

Non-metric Multidimensional Scaling (NMDS) with the *metaMDS* function and tested

statistically using permutational multivariate analysis of variance (PERMANOVA) with the

function *adonis* in the package vegan. Pre-defined cut-off values were used for effect size, where

PERMANOVA results were interpreted as moderate and strong effects if $R^2 > 0.09$ and $R^2 >$

0.25 respectively. These values are broadly equivalent to correlation coefficients of $r = 0.3$ and

0.5 which represent moderate and strong effects accordingly (Nakagawa & Cuthill, 2007). All

plots were produced using the packages ggplot2 v3.3.5 (Wickham, 2016) and ggpubr v0.4.0

(Kassambara, 2020).

3. Results

3.1 Primer selection and validation

Our literature review revealed almost 120 individual primers that yielded over 60 primer

combinations across two mitochondrial and two nuclear genes that were suitable for invertebrate

metabarcoding, of which 22 were tested *in silico* (Table S3). A total of 15 primer sets designed

to amplify regions of the mitochondrial COI and 16S genes were deemed suitable for

microfluidic eDNA metabarcoding based on Fluidigm 48.48 Access Array™ parameters and

taxonomic coverage (Table S3). These primer sets have been used in studies investigating

diversity of terrestrial arthropods (Clarke et al., 2014), freshwater macroinvertebrates (Elbrecht

& Leese, 2017; Vamos et al., 2017), freshwater molluscs (Klymus et al., 2017), and marine

metazoans (Günther et al., 2018; Leray et al., 2013; Meusnier et al., 2008) as well as diet of

various predators (Corse et al., 2019; Zeale et al., 2011). Several of the selected primer sets used

here were recently vetted by Elbrecht et al. (2019) for terrestrial arthropod biodiversity

assessment using DNA metabarcoding, where eight published primer sets outperformed 28

others that were tested, including: BF3/BR2 (Elbrecht et al., 2019), BF1/BR2 (Elbrecht & Leese,

2017), fwhF2/fwhR2n (Vamos et al., 2017), and mlCOIintF/jgHCO2198 (Geller et al., 2013;

Leray et al., 2013). Our *in vitro* testing confirmed that the 15 selected primer sets successfully

amplified pollinators, specifically 11 bee species that occur in Illinois (Figure 2). Given these

results, we chose to retain the full primer panel for microfluidic eDNA metabarcoding.

3.2 Microfluidic eDNA metabarcoding

Each sequencing run generated 20,771,510 and 19,116,230 sequence reads, respectively. On

average, each primer set produced 1,384,767 and 1,274,415 sequence reads for samples/controls

on each sequencing run (average of 4,808 and 8,850 sequence reads per sample/control with each

primer set). UniMinibarF1/UniMinibarR2 produced the fewest sequence reads (6,824 and 4,556)

whereas mlCOIintF/jgHCO2198 (3,844,444 and 3,516,470) and Ins16S-1F/Ins16S-1Rshort

(4,743,478 and 5,848,630) produced the highest reads for COI and 16S, respectively. The

majority of taxonomically assigned reads belonged to Arthropoda for most primer sets,

excluding BF2/BR2 (70.79% and 80.44%), BF3/BR2 (66.19% and 81.39%) and

MOL16S_F/MOL16S_R (26.52% and 25.63%). The primer sets that proportionately had the

most reads assigned to Apidae were nsCOIFo/mlCOIintK (2.01% and 14.11%), fwhF2/fwhR2n

(2.91% and 7.94%), BF1/BR2 (1.23% and 3.72%), LepLCO/MLepF1rev (2.60% and 2.15%),

and mlCOIintF/jgHCO2198 (1.98% and 3.60%). Raw read counts, bioinformatics parameters,

taxonomically assigned read counts, and unassigned read counts are summarised for each primer

set in Table S4.

Microfluidic eDNA metabarcoding correctly identified seven out of 10 bee species used

as PCR positive controls to species level, with three species identified to genus or family level.

However, the PCR positive controls contained a number of contaminants (Figure S3) due to the

manner in which bee specimens were collected, stored, and preserved by the INHS Insect

Collection. In contrast, the negative process controls (field, extraction, and PCR) exhibited very

little contamination (Figure S3). Before false positive threshold application and filtering of

taxonomic assignments, we detected 358 taxa in the 40 baseline controls and 336 eDNA

samples. A total of 279 taxa were either consistently detected below the sequence threshold

(2.05%) calculated using our negative process controls or non-invertebrate and of coarse

taxonomic resolution. The final dataset after threshold application contained 79 taxa for

downstream analyses.

3.3 Greenhouse experiment

The common eastern bumblebee was not detected in any baseline controls collected from flower

species before they were introduced to the secure room (Figure S6), but was successfully

detected in eDNA samples collected from flower species after their introduction. *Monarda* whole

flowers ($n = 1$) and swabs ($n = 3$) preserved in CTAB yielded positive detections for the target

species. All sample types from *Penstemon*, *Solanum*, and *Cyngolossum amabile* failed to detect

the target species (Figure S7). However, *Solanum* whole flowers preserved in ATL buffer ($n = 1$)

produced weak eDNA signals for common eastern bumblebee, but these were removed by our

false positive sequence threshold.

In addition to the target species, pest control and resident arthropods were detected in

samples collected from the greenhouse. Specifically, the silverleaf whitefly (*Bemisia tabaci*),

dark-winged fungus gnat (*Bradysia impatiens*), western flower thrip (*Frankliniella occidentalis*),

insidious flower bug (*Orius* spp.), and aphids (*Aphis* spp.) were detected (Figures S6 and S7).

However, a number of beneficial insect species as well as species known to occur in the

greenhouse were not detected, including *Amblyseius californicus*, *Brasysia coprophilia*, camel

crickets (*Ceuthophilus* spp.), *Encarsia formosa*, convergent lady beetle (*Hippodamia*

convergens), bold jumping spider (*Phidippus audax*), shore fly (*Scatella stagnalis*), spider mites

(Tetranychidae spp.), banded-wing whitefly (*Trialeurodes abutiloneus*), and greenhouse whitefly

(*T. vaporariorum*). Of these species, only shore fly was detected in *Cynoglossum amabile* nectar

preserved in ATL buffer ($n = 1$) and *Penstemon* whole flowers preserved in CTAB ($n = 2$) before

false positive threshold application. Nonetheless, several species not previously known to occur

in the greenhouse were detected, for example, cigarette beetles (*Lasioderma serricorne*).

3.4 Field experiment

We observed flower visitation from seven bee species in the field: common eastern bumblebee,

black-and-gold bumblebee (*B. auricomus*), two-spotted bumblebee (*B. bimaculatus*), brown-

belted bumblebee (*B. griseocollis*), American bumblebee (*B. pensylvanicus*), European honey

bee (*Apis mellifera*), and eastern carpenter bee (*Xylocopa virginica*). Prior to false positive

sequence threshold application, the brown-belted bumblebee and half-black bumblebee (*B.*

vagans) were detected at low frequencies from *Penstemon* whole flowers preserved in CTAB,

but not *Penstemon* swabs or any *Monarda* samples. None of the observed bee species were

detected after false positive sequence threshold application (Figure S8). This included the eastern

carpenter bee despite widespread observations and extensive evidence of nectar robbing by this

species (Figure S12). Arthropods that were not directly observed but were recovered via

microfluidic eDNA metabarcoding included the eastern calligrapher (*Toxomerus geminatus*),

margined calligrapher (*T. marginatus*), cigarette beetle (*Melanophthalma inermis*), banded

garden spider (*Argiope trifasciata*), sheet weavers (*Islandiana flaveola* and *Tennesseellum*

formica), house fly (*Musca domestica*), egg parasitoid wasps (*Telenomus podisi*), Asian tiger

mosquito (*Aedes albopictus*), and firebrat (*Thermobia domestica*). Most taxa were detected from

Penstemon whole flowers or swabs (Figure S8).

3.5 Influence of biological and technical factors on eDNA detection

Overall, flower species did not influence alpha diversity of eDNA samples collected from the

greenhouse (H = 3.070, *P* = 0.381; Figure 3ai) or UIUC campus (H = 0.009, *P* = 0.925; Figure

3bi). Taxon richness for *Penstemon* tended to be higher than other greenhouse flower species

(*Cynoglossum amabile*: Z = -1.457, *P* = 0.871; *Monarda* Z = -0.166, *P* = 0.868; *Solanum*: Z =

1.116, *P* = 0.529). Taxon richness for *Penstemon* was comparable to *Monarda* around the UIUC

campus (Z = 0.094, unadjusted *P* = 0.925), but maximum taxon richness for *Penstemon* was

higher. Beta diversity among greenhouse (Figure 3aii-iv) and field (Figure 3bii-iv) eDNA

samples was largely driven by turnover as opposed to nestedness-resultant (Table 1). MVDISP

did not significantly differ between flower species for any beta diversity component except nestedness-resultant in the UIUC campus samples (Table 1). Flower species had a moderate positive effect on turnover (Figure 3aii) and total beta diversity (Figure 3aiv) of greenhouse arthropod communities, and a weak positive effect on turnover (Figure 3bii) and total beta diversity (Figure 3biv) of UIUC campus arthropod communities (Table 1).

Sample type did not influence alpha diversity of eDNA samples collected from the greenhouse (H = 3.923 P = 0.141; Figure 4ai), but did influence alpha diversity of UIUC campus eDNA samples (H = 6.459, P = 0.011; Figure 4bi). Taxon richness from nectar draws (*Cynoglossum amabile* only) was generally lower than swabs (Z = -0.692, P = 0.489) and whole flowers (Z = -1.494, P = 0.406) collected during the greenhouse experiment. In both the greenhouse and field experiments, swabs tended to produce lower taxon richness than whole flowers (greenhouse: Z = -1.473, P = 0.211; UIUC campus: Z = -2.541, unadjusted P = 0.011; Figures 4ai, bi). MVDISP did not differ between sample types for any beta diversity component except turnover in the UIUC campus samples (Table 2). Sample type had a weak positive effect on nestedness-resultant (Figure 4biii) of UIUC campus arthropod communities, but not any other component of beta diversity for greenhouse or UIUC campus samples (Table 2).

eDNA isolation protocol did not influence alpha diversity of eDNA samples collected from the greenhouse (H = 0.012, P = 0.915; Figure 5ai) or the UIUC campus (H = 3.569, P = 0.059; Figure 5bi). There was very little difference in taxon richness between protocols for the greenhouse experiment (Z = 0.107, unadjusted P = 0.915; Figure 5ai). In the field experiment, taxon richness was typically greater in samples preserved in ATL buffer for QBT extraction (Z = 1.889, unadjusted P = 0.059) as opposed to CTAB preservation and PCI extraction (Figure 5bi). MVDISP did not differ between eDNA isolation protocols for any beta diversity component

(Table 3). eDNA isolation protocol did not influence any beta diversity component (Table 3;

Figures 5aii-iv, bii-iv).

4. Discussion

The past decade has yielded remarkable technological advances to address the critical need to

execute rapid, large-scale surveys of pollinators using metabarcoding (Elbrecht et al., 2019;

Roger et al., 2021; Thomsen & Sigsgaard, 2019). More recently, microfluidic eDNA

metabarcoding has enabled researchers to "cast a broader net" and compensate for taxonomic

biases or blind-spots inherent to individual primer sets (Hauck et al., 2019). We applied

microfluidic eDNA metabarcoding to flower samples and found this approach to be a viable

means of documenting arthropod communities, including pollinators, confirming the work of

Thomsen & Sigsgaard (2019). Moreover, we tested biological and technical sources of

variability in arthropod eDNA detection to develop an optimized workflow for eDNA-based

pollinator biodiversity assessment. Below we contextualize our results and provide a framework

for subsequent field research aimed at further validation of eDNA metabarcoding for assessment

of all pollinator groups (e.g., insects, bats, and birds).

4.1 Greenhouse and field experiments

Both experiments provided important insights for field survey design. The most notable result of

our greenhouse experiment was that microfluidic eDNA metabarcoding was effective in

detecting the target common eastern bumblebee from one of our focal flower species (*Monarda*).

We also detected insects that had been released for pest control and residents known to occur in

the greenhouse facilities, further underscoring the utility of the approach to document arthropod

communities. However, detection was imperfect as only *Monarda* swabs and whole flowers

preserved in CTAB for PCR extraction yielded positive bumblebee detections, and several

beneficial insect species as well as species known to occur in the greenhouse were not detected.

We drew nectar from, swabbed, and harvested multiple flowers from each focal species

immediately after visits from bumblebees, but *Penstemon*, *Solanum*, and *Cynoglossum amabile*

did not yield any detections. The positive *Monarda* detections were flowers that had sat in the

greenhouse for prolonged periods of time, suggesting that eDNA may accumulate to high enough

levels for detection only after repeated visits by pollinator species, similar to other terrestrial

species and aquatic or soil eDNA (Kucherenko et al., 2018; Williams et al., 2018).

Expanding upon the greenhouse experiment, eDNA metabarcoding detected pollinators

and other plant-associated arthropods in natural flower plots around the UIUC campus, though

again with interesting caveats. Numerous arthropod taxa were detected from our focal flower

species (*Monarda* and *Penstemon*), including Arachnida, Coleoptera, Diptera, Hymenoptera, and

Lepidoptera. Yet, not all observed bumblebee species were detected, suggesting that inherent

characteristics of some pollinator species and/or their interactions with various plant

morphologies may make them difficult to detect with this approach. Conventional surveys (e.g.,

visual, acoustic, and trapping) of arthropod communities that incorporate expert taxonomic

identification would be needed for comprehensive estimation of non-detection rates and to

pinpoint taxa which are frequently missed by eDNA metabarcoding.

Nevertheless, our controlled greenhouse experiment and pilot field experiment suggest

that the development of an intensive sampling regime using flower swabbing and whole flower

harvest across multiple flower species differing in their morphology, flowering phenology, etc. at

fine spatiotemporal scales would yield a robust approach to rapidly assess pollinator

communities in diverse ecosystems. We expand upon the measures needed to advance eDNA-

based pollinator biodiversity assessment below.

4.2 Influence of biological and technical factors on eDNA detection

Thomsen and Sigsgaard (2019) provided the first evidence that eDNA metabarcoding could be

used to monitor pollinator communities and other plant-associated arthropods by sequencing

wild flowers. Our study advanced this monitoring approach by evaluating the influence of

methodological choices throughout the eDNA metabarcoding workflow on the pollinator

communities detected. Specifically, we tested four flower species with distinct morphological

characteristics, three sample types, and two eDNA isolation protocols. While not extensively

compared here, we also used 15 primer sets for two markers and processed our data using two

bioinformatics pipelines. Our results show that eDNA detection of pollinator communities (and

other plant-associated arthropods) is likely driven by synergistic or antagonistic effects of

biological and technical factors. While differences in alpha diversity were negligible, community

composition was influenced by flower species and sample type in both the greenhouse and field

experiments. Taxa detected from one flower species were different taxa to those detected from

other flower species, but taxa detected from one sample type were subsets of the communities

detected with other sample types. These differences may arise from several sources, but suggest

the diversity of arthropod communities revealed is likely dependent upon the diversity of the

plant community sampled.

Flower morphology likely affects the ability of flower species to capture and retain

eDNA from pollinators. Three of our focal flower species have distinct morphologies (Figures

1c-e). *Penstemon* has deep, conical flowers that require insects to crawl into the flower to seek

pollen and nectar. *Monarda* has multiple small flowers arranged in a head-like cluster providing

pollen and nectar. *Solanum* flowers are considerably smaller, with petals that do not obscure the

stamen and no nectar. We hypothesize that the small size, low surface area, and exposed stamens

of *Solanum* flowers coupled with the buzz pollination strategy required for this plant resulted in

limited contact and thus less DNA deposition from the target common eastern bumblebee in the

greenhouse, ultimately lowering detection probability. Although common eastern bumblebee

DNA was detected in a *Solanum* sample, the signal was weak and subsequently removed by our

false positive sequence threshold. In contrast, *Monarda* inflorescences are large, complex, have

more overall nectar, and thus have a morphology that promotes longer handling and

consequently increased deposition of DNA and probability of capturing and retaining pollinator

eDNA. *Penstemon* has similar potential, as inflorescences are deep, ostensibly resulting in more

pollinator surface area contact, and flower surfaces less exposed to the elements. Environmental

exposure (e.g., temperature, rainfall, UV, and microbial activity) was previously found to lower

detection of prey items from faeces with metabarcoding due to faster DNA degradation

(McInnes et al., 2016; Oehm et al., 2011), and could similarly affect eDNA on flowers.

Morphological differences may also lead to differential handling times for insect species.

Though we did not quantify handling time in this study, our observations indicate that

bumblebees spent considerably longer on *Monarda* inflorescences, spent less time on *Penstemon*

flowers, and spent the least amount of time on *Solanum* flowers. Our observations echo

numerous studies that have identified differential handling times by bumblebees across a variety

of flowers (Franco et al., 2011; Ivey, Martinez, & Wyatt, 2003), which is associated with

morphological variation among bumblebee species (Harder, 1983). Increased contact and

handling time with flowers may confer higher detection probabilities, ultimately influencing the

ability of eDNA metabarcoding to recover pollinator communities. Curiously, some species that

were frequently observed left no trace of eDNA on flowers, such as eastern carpenter bees. In

Penstemon flower beds on the UIUC campus, we observed numerous visits by carpenter bees

and collected multiple samples immediately after visitation. Moreover, nearly every flower we

inspected showed evidence of nectar robbing (Figure S12), yet no reads were assigned to this

species before or after false positive threshold application. Eastern carpenter bee DNA was used

as a PCR positive control and successfully detected, thus lack of detection was not due to PCR,

sequencing, or bioinformatics failure.

Sample type also influenced pollinator community composition. Sampling the entire

flower head, irrespective of flower species, produced more species detections than swabbing or

nectar draws. This method has already been established as an effective means of documenting

pollinator communities (Thomsen & Sigsgaard, 2019), and likely results in improved chance of

eDNA capture as well as increased DNA yield. Nevertheless, swabbing was able to detect

pollinators, including the target common eastern bumblebee, and may be a more appropriate

means of eDNA sampling in remote locations where it is unreasonable to transport the requisite

sampling materials for whole flower harvesting, or in situations where a flower species is rare

and/or legally protected. Nectar draws may be a less practical method to detect eDNA from

pollinators for a number of reasons. First, some flowering plants do not produce nectar (e.g.,

Solanum), limiting the plant species that could be sampled with this approach. Second, nectar

robbing may render nectar-producing flowers unable to be sampled due to nectar depletion.

Finally, volumes of nectar among flower species can be variable and, in many instances,

incredibly low, which may reduce detection probability. The effect of interplay between nectar

depletion by a pollinator and nectar regeneration by a plant on eDNA detection is also unknown.

Further exploration with high volume nectar-producing plants pollinated by insects as well as

bird- and bat-pollinated plants is critical to close this knowledge gap.

eDNA isolation protocols are known to substantially influence eDNA detection of target

taxa (e.g., Djurhuus et al., 2017; Hinlo et al., 2017; Renshaw et al., 2015; Spens et al., 2016), but

comparisons have largely focused on aquatic eDNA and may not be transferable to eDNA on

flowers (but see Lear et al., 2018). In our study, eDNA isolation protocol did not influence

pollinator eDNA detection. Overall, we found ATL preservation with a modified QBT extraction

(Thomsen & Sigsgaard, 2019) generally recovered greater taxon richness than CTAB

preservation and PCI extraction (Renshaw et al., 2015), but the latter was more effective in

detecting our target bumblebee species in the greenhouse experiment. Community dissimilarity

stemmed from communities produced by CTAB preservation and PCI extraction being

comprised of different taxa to the communities produced by ATL preservation and QBT

extraction. These results reaffirm findings from aquatic eDNA research (Djurhuus et al., 2017;

Hinlo et al., 2017; Lear et al., 2018), but further investigations using samples from natural flower

plots where species inventories from conventional surveys are available would be worthwhile.

We used microfluidic eDNA metabarcoding to counteract some of the biases introduced

in standard eDNA metabarcoding by primer choice. When performance of individual primer sets

was examined, there was substantial variation in the total number of taxa within (Figure S4) and

across (Figure S5) samples as well as the number of unique taxa detected by each primer set and

distinct taxa shared by different combinations of primer sets (Figure S5). This reinforces the

importance of carefully evaluating primer sets *in silico* and *in vitro*, and using primers targeting

different markers and different regions (Corse et al., 2019; Polanco F. et al., 2021). Where

studies are limited to selection of one primer set or marker for budgetary reasons, COI primers

maintain their position in providing greater taxonomic coverage and resolution for arthropods,

specifically mlCOIintF/jgHCO2198 (Geller et al., 2013; Leray et al., 2013), ZBJArtF1c-

ZBJArtR2c (Zeale et al., 2011), BF3-BR2 and BF2-BR2 (Elbrecht et al., 2019; Elbrecht &

Leese, 2017). However, metabarcoding is an evolving field with new primers targeting

established regions as well as new markers targeting novel regions constantly emerging, and the

best primers and markers will likely vary with target taxonomic group. Therefore, we advocate

for continued assessment of individual primer sets and primer set combinations to maximize the

biodiversity recovered whilst minimising effort and costs associated with eDNA metabarcoding.

Similarly, different bioinformatics pipelines can produce variable results due to the

different softwares, parameterizations and algorithms implemented. We assessed two

bioinformatics pipelines using different tools, namely clustering (metaBEAT) versus ASV

parsing (Anacapa) and a custom reference database (metaBEAT) versus a CRUX database

(Anacapa) (Curd et al., 2019; Hänfling et al., 2016). metaBEAT detected more arthropod species

known to occur in the greenhouse facilities (Figure S1), thus data produced by this pipeline were

used for downstream analyses. As bioinformatics pipelines continue to be developed and refined

for processing metabarcoding data and public reference databases are bolstered by additional

sequences, continued evaluation of bioinformatics pipelines is necessary to assess potential

biases and taxon recovery (Mathon et al., 2021; O'Rourke et al., 2020). As such, we recommend

that future studies assess different bioinformatic pipelines and their effects on species detection

against known species inventories to help minimise the occurrence of false negatives and false

positives in eDNA metabarcoding datasets.

4.3 Using hindsight to inform foresight

In our study, samples were PCR amplified and sequenced once. This was likely a main factor

contributing to low eDNA detection of plant-associated arthropods present in the greenhouse

facilities and observed in natural flower plots as there were no additional PCR replicates to offset

PCR stochasticity and enable an additive strategy to bolster species inventories (Alberdi et al.,

2018). Typically, metabarcoding studies have used three PCR replicates that are either pooled for

sequencing or sequenced independently (Alberdi et al., 2018; Beentjes et al., 2019; Weigand &

Macher, 2018), but eight or more PCR replicates may be required when detection probabilities

are likely to be low (Ficetola et al., 2015). Eight PCR replicates were used by Hauck et al. (2019)

for microfluidic eDNA metabarcoding, but some standard eDNA metabarcoding studies have

used as many as 12 (Valentini et al., 2016) or 15 (Sato et al., 2017) PCR replicates. Greater PCR

replication can improve detection rates for rare and/or low-density species, and enable

identification of potential false positives if taxa occur in only one PCR replicate. However,

removal of taxa occurring in a single replicate may increase the possibility of false negatives for

rare and/or low-density taxa (Weigand & Macher, 2018). Conversely, some studies have

suggested there may be little impact of including taxa that occur in only one PCR replicate on

inferences of biodiversity patterns and community dissimilarly (Beentjes et al., 2019). Therefore,

PCR replication may be of greater importance in alpha diversity estimation than beta diversity

estimation. This is a key area of investigation for future pollinator and plant-associated arthropod

eDNA research.

Similarly, more biological replication may be necessary to enhance pollinator eDNA

detection rates, as has been observed for eDNA detection of terrestrial vertebrates from water

(Broadhurst et al., 2021; T.-H. Macher et al., 2021). Here, only one flower was sampled per

plant, but plant-associated arthropods may not have interacted with this particular flower. Since

this study was conceived, alternative strategies for eDNA detection of terrestrial insects have

been successfully trialled, such as eDNA aggregation via water (Valentin et al., 2020), airborne

eDNA (Roger et al., 2022), and eDNA deposited on leaves (Krehenwinkel et al., 2022). These

approaches may circumvent the need for biological replication and invasive sampling of flowers

but require extensive validation against the current sampling strategies of swabbing flowers

and/or whole flower harvest for pollinator eDNA used here and by Thomsen and Sigsgaard

(2019).

Another important consideration is the use of microfluidic technology. High sequencing

depth is key alongside PCR replication to improve detection of rare and/or low-density taxa in

complex environmental samples (Alberdi et al., 2018; Doble et al., 2019; Hajibabaei et al.,

2019). However, sequencing depth is inherently lower with microfluidic eDNA metabarcoding

as it is split across the number of primer sets being used. Although we sequenced eDNA samples

across two MiSeq runs, sequencing depth may have been insufficient for our primer panel

consisting of 15 primer sets. There is an important trade-off to consider, as using a larger primer

panel with the Fluidigm 48.48 Access Array™ allows the performance of more individual primer

sets to be evaluated. Prior to false positive threshold application, we found that the primer set

mlCOIintF/jgHCO2198 (Geller et al., 2013; Leray et al., 2013) detected the most taxa, followed

by BF2/BR2 and BF3/BR2 (Elbrecht et al., 2019; Elbrecht & Leese, 2017), but each of the 15

primer sets detected taxa not identified by other primer sets, and different combinations of

primer sets detected taxa not identified by other combinations (Figure S5). More sequencing

depth is possible with the use of more powerful sequencers, such as the Illumina NovaSeq, with

cost of these sequencing platforms being the main barrier to uptake at present. However, as the

cost of Illumina sequencing continues to fall, greater sequencing depth can be achieved at

equivalent or lower price points (even compared to one year ago). A magnitudinal increase in

reads returned should also increase the likelihood of detecting DNA traces present at lower

concentrations because they were left behind by species that are rarer, smaller, present at lower

density or exhibited less flower handling time. This would greatly improve the efficacy of

pollinator eDNA metabarcoding. However, where rare pollinators are the focus of monitoring

efforts, it may be more appropriate to use targeted eDNA analysis (e.g. qPCR) for increased

sensitivity (L. R. Harper et al., 2018; Moss et al., 2022).

PCR reaction volumes are also much lower using the Fluidigm 48.48 Access Array™ (5

μL) than they would be in standard PCR (typically 25 μL or 50 μL). Consequently, the amount

of template DNA added to Fluidigm 48.48 Access Array™ reactions is a fraction of that used in

standard PCR reactions, which is problematic as higher volumes of template DNA reduce

uncertainty in detection rates (Mächler et al., 2015). In our study, 2 ng of template DNA was

loaded to the Fluidigm 48.48 Access Array™, providing 0.13 ng DNA for amplification with

each primer set in each reaction chamber. Although the initial volume of template DNA could be

increased, the risk of inhibition or overwhelming PCR reactions also increases (Lance & Guan,

2020). Furthermore, the Fluidigm 48.48 Access Array™ must be run at a fixed annealing

temperature which may be suboptimal for certain primer sets, thus impairing species detection

(Clarke et al., 2017; Elbrecht et al., 2019). Therefore, metabarcoding projects may wish to

exploit microfluidic technology in the first instance to determine which primer sets maximise

community diversity, provide the most informative data, and should be used for processing of

future samples (Brown et al., 2016), whilst acknowledging the caveats associated with this

technology.

Contamination of eDNA samples can arise from many sources, including field sampling,

DNA extraction, PCR, or sequencing errors (Ficetola et al., 2015; Ficetola et al., 2016;

Sepulveda et al., 2020). We identified low levels of contamination in field and extraction blanks

as well as PCR negative controls, yet considerably higher levels of contamination in PCR

positive controls. The specimens used for our PCR positive controls were collected using pan

traps, bulk preserved in ethanol, and sorted by hand without sterile gloves in unsterile conditions.

The samples were then pinned and stored using standard entomological practices in open

drawers at the INHS for several years. Given the above, we speculate that the frequency and

composition of contaminants (including other pollinators, mites, and decomposers) likely

occurred from contact with: 1) other species in the field prior to sampling (pre-catch), 2) other

species during sampling and preservation (catch), and/or 3) common arthropod species living in

museum collections (post-catch). This contamination led to the implementation of an ostensibly

high and exclusionary false positive sequence threshold based on our negative process controls

which likely removed true detections. Therefore, careful selection and procurement of specimens

and their DNA for PCR positive controls will reduce threshold stringency, further enhancing the

performance of pollinator eDNA metabarcoding.

5. Conclusions

Our study adds to a growing body of evidence that eDNA metabarcoding is a viable means of surveying pollinator communities. Methodological choices throughout the entire eDNA metabarcoding workflow, from what to sample, how to sample, eDNA isolation protocol, primer choice, and bioinformatics pipeline, can impact the community diversity elucidated using this approach. Based on our results, we have synthesised a recommended workflow for pollinator biodiversity assessment using eDNA metabarcoding (Figure 6). First, given the observed differences in detection across flower species, we recommend that replicate samples from a diverse flower species assemblage from a given site be established. By maximizing flower diversity (and considering spatial and temporal variation in pollinator communities), one should maximize the probability of detecting a comprehensive pollinator (and other plant-associated arthropod) community. Second, to maximize detection of diverse arthropod/pollinator communities, we recommend that whole flower heads are sampled. This increases the surface area available for eDNA to be captured from and may require less time to sample than swabbing or nectar draws. However we note that in certain contexts (i.e., in remote locations and/or in dealing with threatened/endangered plants that cannot be destructively sampled) swabbing may be a viable alternative. Third, ATL preservation followed by a modified QBT extraction generally revealed more diverse pollinator communities. Moreover, this approach tends to recover clean, long-stranded DNA. Fourth, the COI locus performed better than the 16S locus in recovering pollinator communities and minimizing the number of unassigned reads due to greater reference database coverage. Additionally, use of multiple primer sets helped minimize taxonomic biases of individual primer sets. On this basis, we recommend using a suite of COI

primers in future metabarcoding studies of insect pollinators, but this is unlikely to be optimal for

other pollinator groups (e.g., bats and birds). As such, *in silico* and *in vitro* primer validation

should be undertaken for available primer sets for common metabarcoding markers (e.g., COI,

12S rRNA, 16S rRNA, 18S rRNA) depending on the pollinator community one seeks to

document. Finally, metaBEAT seemingly recovered more diversity and detected more species

known to be present in greenhouse facilities than Anacapa, but results may have been influenced

by the use of different reference databases and tools for bioinformatic processing. Therefore, we

advise future studies should undertake comparisons with available pipelines and compare data to

known species inventories to select the most appropriate pipeline for their dataset.

Acknowledgements

We are grateful for the Strategic Environmental Research and Development Program for funding

this research (grant number RC19-1102). We thank Patricia Dickerson for technical assistance.

We are deeply grateful to Rosalie Metallo, Montgomery Flack, and Clinton Shipley at the UIUC

Plant Care facility for technical, logistical, and moral support of our research. We appreciate Dr.

Mark R. Band, Dr. Alvaro G. Hernandez, Dr. Christy L. Wright and the staff at the Roy J. Carver

Biotechnology Center Functional Genomics Unit for technical support and conducting

microfluidic metabarcoding. We thank Emily Curd for assistance with implementing Anacapa.

We recognize Danielle Ruffatto, Tiffany Jolley, and Joe Spencer for imaging and design support.

We thank Catherine Dana, Tara Hohoff, and Tommy McElrath for providing samples, as well as

encouragement and creative criticism to improve this research. We thank Angella Moorehouse at

the Illinois Nature Preserves Commission for supplying species lists used to develop the

reference DNA sequence database.

Authors contributions

M.A.D., B.M-F., and M.L.N. conceived and designed the study. B.M-F., L.E.P., and E.K.

established the greenhouse flower assemblage and M.A.D., L.E.P., E.K., and B.M-F. collected

samples. M.A.D., L.E.P., and E.K. performed laboratory work before samples were sent to the

sequencing facility. M.L.N., J.B., and L.R.H. performed bioinformatics processing, and L.R.H.

analysed the data. L.R.H. and M.A.D. wrote the manuscript, which all authors revised.

Data availability statement

Raw sequence reads, bioinformatics scripts, R scripts, and corresponding

data have been permanently archived at Zenodo (https://doi.org/10.5281/zenodo.6756972).

References
773

Alberdi, A., Aizpurua, O., Gilbert, M. T. P., & Bohmann, K. (2018). Scrutinizing key steps for
reliable metabarcoding of environmental samples. *Methods in Ecology and Evolution*, *9*, 134–
147. https://doi.org/10.1111/2041-210X.12849
Barnosky, A. D., Matzke, N., Tomiya, S., Wogan, G. O. U., Swartz, B., Quental, T. B., …
Ferrer, E. A. (2011) Has the Earth's sixth mass extinction already arrived? *Nature*, *471*, 51–
57. https://doi.org/10.1038/nature09678

Bartomeus, I. (2013). Understanding linkage rules in plant-pollinator networks by using hierarchical models that incorporate pollinator detectability and plant traits. *PLoS ONE*, *8*, e69200. https://doi.org/10.1371/journal.pone.0069200

Baselga, A., & Orme, C. D. L. (2012). betapart: an R package for the study of beta diversity : Betapart package. *Methods in Ecology and Evolution*, *3*, 808–812. https://doi.org/10.1111/j.2041-210X.2012.00224.x

Beentjes, K. K., Arjen G C, Schilthuizen, M., Hoogeveen, M., & van der Hoorn, B. B. (2019). The effects of spatial and temporal replicate sampling on eDNA metabarcoding. *PeerJ*, *7*, e7335. https://doi.org/10.7717/peerj.7335

Biggs, J., Ewald, N., Valentini, A., Gaboriaud, C., Dejean, T., Griffiths, R. A., … Dunn, F. (2015). Using eDNA to develop a national citizen science-based monitoring programme for the great crested newt (*Triturus cristatus*). *Biological Conservation*, *183*, 19–28. https://doi.org/10.1016/j.biocon.2014.11.029

Bolger, A. M., Lohse, M., & Usadel, B. (2014). Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics*, *30*, 2114–2120. https://doi.org/10.1093/bioinformatics/btu170

Bolyen, E., Rideout, J. R., Dillon, M. R., Bokulich, N. A., Abnet, C. C., Al-Ghalith, G. A., … Caporaso, J. G. (2019). Reproducible, interactive, scalable and extensible microbiome data science using QIIME 2. *Nature Biotechnology*, *37*, 852–857. https://doi.org/10.1038/s41587-019-0209-9

Boyer, F., Mercier, C., Bonin, A., Le Bras, Y., Taberlet, P., & Coissac, E. (2016). obitools: a unix-inspired software package for DNA metabarcoding. *Molecular Ecology Resources*, *16*, 176–182. https://doi.org/10.1111/1755-0998.12428

Broadhurst, H. A., Gregory, L. M., Bleakley, E. K., Perkins, J. C., Lavin, J. V., Bolton, P., … McDevitt, A. D. (2021). Mapping differences in mammalian distributions and diversity using environmental DNA from rivers. *The Science of the Total Environment*, *801*, 149724. https://doi.org/10.1016/j.scitotenv.2021.149724

Brown, S. P., Ferrer, A., Dalling, J. W., & Heath, K. D. (2016). Don't put all your eggs in one basket: a cost-effective and powerful method to optimize primer choice for rRNA environmental community analyses using the Fluidigm Access Array. *Molecular Ecology Resources*, *16*, 946–956. https://doi.org/10.1111/1755-0998.12507

Cameron, S. A., Lozier, J. D., Strange, J. P., Koch, J. B., Cordes, N., Solter, L. F., & Griswold, T. L. (2011). Patterns of widespread decline in North American bumble bees. *Proceedings of the National Academy of Sciences of the United States of America*, *108*, 662–667. https://doi.org/10.1073/pnas.1014743108

Carvalheiro, L. G., Kunin, W. E., Keil, P., Aguirre-Gutiérrez, J., Ellis, W. N., Fox, R., … Biesmeijer, J. C. (2013). Species richness declines and biotic homogenisation have slowed down for NW-European pollinators and plants. *Ecology Letters*, *16*, 870–878. https://doi.org/10.1111/ele.12121

Clare, E. L., Economou, C. K., Bennett, F. J., Dyer, C. E., Adams, K., McRobie, B., … Littlefair, J. E. (2021). Measuring biodiversity from DNA in the air. *bioRxiv*. https://doi.org/10.1101/2021.07.15.452392

Clarke, L. J., Beard, J. M., Swadling, K. M., & Deagle, B. E. (2017). Effect of marker choice and thermal cycling protocol on zooplankton DNA metabarcoding studies. *Ecology and Evolution*, *7*, 873–883. https://doi.org/10.1002/ece3.2667

Clarke, L. J., Czechowski, P., Soubrier, J., Stevens, M. I., & Cooper, A. (2014). Modular tagging of amplicons using a single PCR for high-throughput sequencing. *Molecular Ecology Resources*, *14*, 117–121. https://doi.org/10.1111/1755-0998.12162

Corse, E., Tougard, C., Archambaud-Suard, G., Agnèse, J.-F., Messu Mandeng, F. D., Bilong Bilong, C. F., … Dubut, V. (2019). One-locus-several-primers: A strategy to improve the taxonomic and haplotypic coverage in diet metabarcoding studies. *Ecology and Evolution*, *9*, 4603–4620. https://doi.org/10.1002/ece3.5063

Curd, E. E., Gold, Z., Kandlikar, G. S., Gomer, J., Ogden, M., O'Connell, T., … Meyer, R. S. (2019). Anacapa Toolkit : An environmental DNA toolkit for processing multilocus metabarcode datasets. *Methods in Ecology and Evolution*, *10*, 1469–1475. https://doi.org/10.1111/2041-210X.13214

Deiner, K., Bik, H. M., Mächler, E., Seymour, M., Lacoursière-Roussel, A., Altermatt, F., … Bernatchez, L. (2017). Environmental DNA metabarcoding: Transforming how we survey animal and plant communities. *Molecular Ecology*, *26*, 5872–5895. https://doi.org/10.1111/mec.14350

Didham, R. K., Basset, Y., Matilda Collins, C., Leather, S. R., Littlewood, N. A., Menz, M. H. M., … Hassall, C. (2020). Interpreting insect declines: seven challenges and a way forward. *Insect Conservation and Diversity*, *13*, 103–114. https://doi.org/10.1111/icad.12408

Djurhuus, A., Port, J., Closek, C., Yamahara, K., Romero-Maraccini, O., Walz, K., … Chavez, F. (2017). Evaluation of filtration and DNA extraction methods for environmental DNA biodiversity assessments across multiple trophic levels. *Frontiers in Marine Science*, *4*, 314. https://doi.org/10.3389/fmars.2017.00314

Doble, C. J., Hipperson, H., Salzburger, W., Horsburgh, G. J., Mwita, C., Murrell, D. J., & Day, J. J. (2019). Testing the performance of environmental DNA metabarcoding for surveying highly diverse tropical fish communities: A case study from Lake Tanganyika. *Environmental DNA*, *12*, 1819. https://doi.org/10.1002/edn3.43

Ehlers, E., & Krafft, T. (2006). *Earth System Science in the Anthropocene: Emerging Issues and Problems*. Springer Nature.

Elbrecht, V., Braukmann, T. W. A., Ivanova, N. V., Prosser, S. W. J., Hajibabaei, M., Wright, M., … Steinke, D. (2019). Validation of COI metabarcoding primers for terrestrial arthropods. *PeerJ*, *7*, e7745. https://doi.org/10.7717/peerj.7745

Elbrecht, V., Hebert, P. D. N., & Steinke, D. (2018). Slippage of degenerate primers can cause variation in amplicon length. *Scientific Reports*, *8*, 10999. https://doi.org/10.1038/s41598-018-29364-z

Elbrecht, V., & Leese, F. (2017). Validation and Development of COI Metabarcoding Primers for Freshwater Macroinvertebrate Bioassessment. *Frontiers in Environmental Science*, *5*. https://doi.org/10.3389/fenvs.2017.00011

Elbrecht, V., Taberlet, P., Dejean, T., Valentini, A., Usseglio-Polatera, P., Beisel, J.-N., … Leese, F. (2016). Testing the potential of a ribosomal 16S marker for DNA metabarcoding of insects. *PeerJ*, *4*, e1966. https://doi.org/10.7717/peerj.1966

Elbrecht, V., Vamos, E. E., Meissner, K., Aroviita, J., & Leese, F. (2017). Assessing strengths and weaknesses of DNA metabarcoding-based macroinvertebrate identification for routine stream monitoring. *Methods in Ecology and Evolution*, *8*, 1265–1275. https://doi.org/10.1111/2041-210X.12789

Ficetola, G. F., Coissac, E., Zundel, S., Riaz, T., Shehzad, W., Bessière, J., … Pompanon, F. (2010). An *In silico* approach for the evaluation of DNA barcodes. *BMC Genomics*, *11*, 434. https://doi.org/10.1186/1471-2164-11-434

Ficetola, G. F., Pansu, J., Bonin, A., Coissac, E., Giguet-Covex, C., De Barba, M., … Taberlet, P. (2015). Replication levels, false presences and the estimation of the presence/absence from eDNA metabarcoding data. *Molecular Ecology Resources*, *15*, 543–556. https://doi.org/10.1111/1755-0998.12338

Ficetola, G. F., Taberlet, P., & Coissac, E. (2016). How to limit false positives in environmental DNA and metabarcoding? *Molecular Ecology Resources*, *16*, 604–607. https://doi.org/10.1111/1755-0998.12508

Fluidigm Corporation (2016) Access array system for Illumina sequencing systems user guide (PN 100–3770 J1). San Francisco, CA: Fluidigm Corporation. www.fluidigm.com

Franco, A. M., Goldenberg, R., & Varassin, I. G. (2011). Pollinator guild organization and its consequences for reproduction in three synchronopatric species of Tibouchina (Melastomataceae). *Revista Brasileira de Entomologia*, *55*, 381–388. https://doi.org/10.1590/S0085-56262011005000040

Gallai, N., Salles, J.-M., Settele, J., & Vaissière, B. E. (2009). Economic valuation of the vulnerability of world agriculture confronted with pollinator decline. *Ecological Economics: The Journal of the International Society for Ecological Economics*, *68*, 810–821. https://doi.org/10.1016/j.ecolecon.2008.06.014

Geller, J., Meyer, C., Parker, M., & Hawk, H. (2013). Redesign of PCR primers for mitochondrial cytochrome c oxidase subunit I for marine invertebrates and application in all-taxa biotic surveys. *Molecular Ecology Resources*, *13*, 851–861. https://doi.org/10.1111/1755-0998.12138

Günther, B., Knebelsberger, T., Neumann, H., Laakmann, S., & Martínez Arbizu, P. (2018). Metabarcoding of marine environmental DNA based on mitochondrial and nuclear genes. *Scientific Reports*, *8*, 14822. https://doi.org/10.1038/s41598-018-32917-x

Hajibabaei, M., Porter, T. M., Robinson, C. V., Baird, D. J., Shokralla, S., & Wright, M. T. G. (2019). Watered-down biodiversity? A comparison of metabarcoding results from DNA extracted from matched water and bulk tissue biomonitoring samples. *PLoS One*, *14*, e0225409. https://doi.org/10.1371/journal.pone.0225409

Hallmann, C. A., Sorg, M., Jongejans, E., Siepel, H., Hofland, N., Schwan, H., … de Kroon, H. (2017). More than 75 percent decline over 27 years in total flying insect biomass in protected areas. *PLoS ONE*, *12*, e0185809. https://doi.org/10.1371/journal.pone.0185809

Hänfling, B., Lawson Handley, L., Read, D. S., Hahn, C., Li, J., Nichols, P., … Winfield, I. J. (2016). Environmental DNA metabarcoding of lake fish communities reflects long-term data from established survey methods. *Molecular Ecology*, *25*, 3101–3119. https://doi.org/10.1111/mec.13660

Harder, L. D. (1983). Flower handling efficiency of bumble bees: morphological aspects of probing time. *Oecologia*, *57*(1-2), 274–280. https://doi.org/10.1007/BF00379591

Harper, K. J., Goodwin, K. D., Harper, L. R., LaCasella, E. L., Frey, A., & Dutton, P. H. (2020). Finding Crush: Environmental DNA Analysis as a Tool for Tracking the Green Sea Turtle *Chelonia mydas* in a Marine Estuary. *Frontiers in Marine Science*, *6*, 810. https://doi.org/10.3389/fmars.2019.00810

Harper, L. R., Lawson Handley, L., Hahn, C., Boonham, N., Rees, H. C., Gough, K. C., … Hänfling, B. (2018). Needle in a haystack? A comparison of eDNA metabarcoding and targeted qPCR for detection of the great crested newt (*Triturus cristatus*). *Ecology and Evolution*, *8*, 6330–6441. https://doi.org/10.1002/ece3.4013

Harper, L. R., Lawson Handley, L., Sayer, C. D., Read, D. S., Benucci, M., Blackman, R. C., … Hänfling, B. (2020). Assessing the impact of the threatened crucian carp (*Carassius carassius*) on pond invertebrate diversity: A comparison of conventional and molecular tools. *Molecular Ecology*, *30*, 3252–3269. https://doi.org/10.1111/mec.15670

Hauck, L. L., Weitemier, K. A., Penaluna, B. E., Garcia, T. S., & Cronn, R. (2019). Casting a broader net: Using microfluidic metagenomics to capture aquatic biodiversity data from diverse taxonomic targets. *Environmental DNA*, *1*, 2008. https://doi.org/10.1002/edn3.26

Hinlo, R., Gleeson, D., Lintermans, M., & Furlan, E. (2017). Methods to maximise recovery of environmental DNA from water samples. *PLoS ONE*, *12*, e0179251. https://doi.org/10.1371/journal.pone.0179251

Ivey, C. T., Martinez, P., & Wyatt, R. (2003). Variation in pollinator effectiveness in swamp milkweed, Asclepias incarnata (Apocynaceae). *American Journal of Botany*, *90*, 214–225. https://doi.org/10.3732/ajb.90.2.214

Kassambara, A. (2020). ggpubr: 'ggplot2' Based Publication Ready Plots. R package version 0.3.0. https://CRAN.R-project.org/package=ggpubr

Katz, A. D., Harper, L. R., Sternhagen, E. C., Pearce, S. E., Melder, C. A., Sperry, J. H., & Davis, M. A. (2020). Environmental DNA is effective in detecting the federally threatened Louisiana Pinesnake (*Pituophis ruthveni*). *Environmental DNA*, 3, 409–425. https://doi.org/10.1002/edn3.126

Kevan, P. G., & Phillips, T. P. (2001). The economic impacts of pollinator declines: an approach to assessing the consequences. *Conservation Ecology*, *5*, 368–383. https://doi.org/10.1017/S1068280500007371

Klein, A.-M., Vaissière, B. E., Cane, J. H., Steffan-Dewenter, I., Cunningham, S. A., Kremen, C., & Tscharntke, T. (2007). Importance of pollinators in changing landscapes for world crops. *Proceedings of The Royal Society B*, *274*, 303–313. https://doi.org/10.1098/rspb.2006.3721

Klymus, K. E., Marshall, N. T., & Stepien, C. A. (2017). Environmental DNA (eDNA) metabarcoding assays to detect invasive invertebrate species in the Great Lakes. *PloS One*, *12*(5), e0177643. https://doi.org/10.1371/journal.pone.0177643

Krehenwinkel, H., Weber, S., Broekmann, R., Melcher, A., Hans, J., Wolf, R., Hochkirch, A., Kennedy, S.R., Koschorreck, J., Kuenzel, S., Müller, C., Retzlaff, R., Teubner, D., Schanzer, S., Klein, R., Paulus, M., Udelhoven, T. & Veith, M. (2022) Environmental DNA from archived leaves reveals widespread temporal turnover and biotic homogenization in forest arthropod communities. *bioRxiv*, 2022.04.27.489699. https://doi.org/10.1101/2022.04.27.489699

Kucherenko, A., Herman, J. E., Iii, E. M. E., & Urakawa, H. (2018). Terrestrial Snake Environmental DNA Accumulation and Degradation Dynamics and its Environmental Application. *Herpetologica*, *74*, 38–49. https://doi.org/10.1655/Herpetologica-D-16-00088

Lance, R. F., & Guan, X. (2020). Variation in inhibitor effects on qPCR assays and implications for eDNA surveys. *Canadian Journal of Fisheries and Aquatic Sciences*, *77*, 23–33. https://doi.org/10.1139/cjfas-2018-0263

Lear, G., Dickie, I., Banks, J., Boyer, S., Buckley, H., Buckley, T., … Holdaway, R. (2018). Methods for the extraction, storage, amplification and sequencing of DNA from environmental samples. *New Zealand Journal of Ecology*, *42*, 10. https://doi.org/10.20417/nzjecol.42.9

Leese, F., Sander, M., Buchner, D., Elbrecht, V., Haase, P., & Zizka, V. M. A. (2020). Improved freshwater macroinvertebrate detection from environmental DNA through minimized nontarget amplification. *Environmental DNA*, *3*, 261–276. https://doi.org/10.1002/edn3.177

Leray, M., Yang, J. Y., Meyer, C. P., Mills, S. C., Agudelo, N., Ranwez, V., … Machida, R. J. (2013). A new versatile primer set targeting a short fragment of the mitochondrial COI region for metabarcoding metazoan diversity: application for characterizing coral reef fish gut contents. *Frontiers in Zoology*, *10*, 34. https://doi.org/10.1186/1742-9994-10-34

Lever, J. J., van Nes, E. H., Scheffer, M., & Bascompte, J. (2014). The sudden collapse of pollinator communities. *Ecology Letters*, *17*, 350–359. https://doi.org/10.1111/ele.12236

Losey, J. E., & Vaughan, M. (2006). The Economic Value of Ecological Services Provided by Insects. *Bioscience*, *56*, 311–323. https://doi.org/10.1641/0006-3568(2006)56[311:TEVOES]2.0.CO;2

Macgregor, C. J., Williams, J. H., Bell, J. R., & Thomas, C. D. (2019). Moth biomass increases and decreases over 50 years in Britain. *Nature Ecology & Evolution*, *3*, 1645–1649. https://doi.org/10.1038/s41559-019-1028-6

Macher, T.-H., Schütz, R., Arle, J., Beermann, A. J., Koschorreck, J., & Leese, F. (2021). Beyond fish eDNA metabarcoding: Field replicates disproportionately improve the detection of stream associated vertebrate species. *Metabarcoding and Metagenomics*, *5*, e66557. https://doi.org/10.3897/mbmg.5.66557

Mächler, E., Deiner, K., Spahn, F., & Altermatt, F. (2015). Fishing in the water: effect of sampled water volume on environmental DNA-based detection of macroinvertebrates. *Environmental Science & Technology*, *50*, 305–312. https://doi.org/10.1021/acs.est.5b04188

Mächler, E., Little, C. J., Wüthrich, R., Alther, R., Fronhofer, E. A., Gounand, I., … Altermatt, F. (2019). Assessing different components of diversity across a river network using eDNA. *Environmental DNA*, *1*, 290–301. https://doi.org/10.1002/edn3.33

Mathon, L., Valentini, A., Guérin, P.-E., Normandeau, E., Noel, C., Lionnet, C., … Manel, S. (2021). Benchmarking bioinformatic tools for fast and accurate eDNA metabarcoding species identification. *Molecular Ecology Resources*, 21, 2565–2579. https://doi.org/10.1111/1755-0998.13430

Matthies, D., Bräuer, I., Maibom, W., & Tscharntke, T. (2004). Population size and the risk of local extinction: empirical evidence from rare plants. *Oikos*, *105*, 481–488. https://doi.org/10.1111/j.0030-1299.2004.12800.x

McInnes, J. C., Alderman, R., Deagle, B. E., Lea, M.-A., Raymond, B., & Jarman, S. N. (2016). Optimised scat collection protocols for dietary DNA metabarcoding in vertebrates. *Methods in Ecology and Evolution*, *8*, 192–202. https://doi.org/10.1111/2041-210X.12677

Meusnier, I., Singer, G. A. C., Landry, J.-F., Hickey, D. A., Hebert, P. D. N., & Hajibabaei, M. (2008). A universal DNA mini-barcode for biodiversity analysis. *BMC Genomics*, *9*, 214. https://doi.org/10.1186/1471-2164-9-214

Montgomery, G. A., Dunn, R. R., Fox, R., Jongejans, E., Leather, S. R., Saunders, M. E., … Wagner, D. L. (2020). Is the insect apocalypse upon us? How to find out. *Biological Conservation*, *241*, 108327. https://doi.org/10.1016/j.biocon.2019.108327

Moss, W.E., Harper, L.R., Davis, M.A., Goldberg, C.S., Smith, M.M. & Johnson, P.T.J. (2022) Navigating the trade-offs between environmental DNA and conventional field surveys for improved amphibian monitoring. *Ecosphere*, **13**, e3941. https://doi.org/10.1002/ecs2.3941

Nakagawa, S., & Cuthill, I. C. (2007). Effect size, confidence interval and statistical significance: a practical guide for biologists. *Biological Reviews*, *82*, 591–605. https://doi.org/10.1111/j.1469-185X.2007.00027.x

Oehm, J., Juen, A., Nagiller, K., Neuhauser, S., & Traugott, M. (2011). Molecular scatology: how to improve prey DNA detection success in avian faeces? *Molecular Ecology Resources*, *11*, 620–628. https://doi.org/10.1111/j.1755-0998.2011.03001.x

Ogle, D. H., Wheeler, P., & Dinno, A. (2020). FSA: Fisheries Stock Analysis. R package version 0.8.30. https://github.com/droglenc/FSA

Oksanen, J., Blanchet, F. G., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., … Wagner, H. (2019). vegan: Community Ecology Package. R package version 2.5-6. https://CRAN.R-project.org/package=vegan

Ollerton, J., Winfree, R., & Tarrant, S. (2011). How many flowering plants are pollinated by animals? *Oikos*, 120, 321–326. https://doi.org/10.1111/j.1600-0706.2010.18644.x

O'Rourke, D. R., Bokulich, N. A., Jusino, M. A., MacManes, M. D., & Foster, J. T. (2020). A total crapshoot? Evaluating bioinformatic decisions in animal diet metabarcoding analyses. *Ecology and Evolution*, *10*, 9721–9739. https://doi.org/10.1002/ece3.6594

Plein, M., Morris, W. K., Moir, M. L., & Vesk, P. A. (2017). Identifying species at coextinction risk when detection is imperfect: Model evaluation and case study. *PLoS ONE*, *12*, e0183351. https://doi.org/10.1371/journal.pone.0183351

Polanco F., A., Richards, E., Flück, B., Valentini, A., Altermatt, F., Brosse, S., … Pellissier, L. (2021). Comparing the performance of 12S mitochondrial primers for fish environmental DNA across ecosystems. *Environmental DNA*. https://doi.org/10.1002/edn3.232

Potts, S. G., Biesmeijer, J. C., Kremen, C., Neumann, P., Schweiger, O., & Kunin, W. E. (2010). Global pollinator declines: trends, impacts and drivers. *Trends in Ecology & Evolution*, *25*, 345–353. https://doi.org/10.1016/j.tree.2010.01.007

Renshaw, M. A., Olds, B. P., Jerde, C. L., McVeigh, M. M., & Lodge, D. M. (2015). The room temperature preservation of filtered environmental DNA samples and assimilation into a phenol-chloroform-isoamyl alcohol DNA extraction. *Molecular Ecology Resources*, *15*, 168–176. https://doi.org/10.1111/1755-0998.12281

Roger, F., Ghanavi, H.R., Danielsson, N., Wahlberg, N., Löndahl, J., Pettersson, L.B., Andersson, G.K.S., Boke Olén, N. & Clough, Y. (2022) Airborne environmental DNA metabarcoding for the monitoring of terrestrial insects - A proof of concept from the field. *Environmental DNA*. https://doi.org/10.1002/edn3.290

Sato, H., Sogo, Y., Doi, H., & Yamanaka, H. (2017). Usefulness and limitations of sample pooling for environmental DNA metabarcoding of freshwater fish communities. *Scientific Reports*, *7*, 14860. https://doi.org/10.1038/s41598-017-14978-6

Sepulveda, A. J., Hutchins, P. R., Forstchen, M., Mckeefry, M. N., & Swigris, A. M. (2020). The Elephant in the Lab (and Field): Contamination in Aquatic Environmental DNA Studies. *Frontiers in Ecology and Evolution*, *8*, 440. https://doi.org/10.3389/fevo.2020.609973

Sheffield, C. S., Hebert, P. D. N., Kevan, P. G., & Packer, L. (2009). DNA barcoding a regional bee (Hymenoptera: Apoidea) fauna and its potential for ecological studies. *Molecular Ecology Resources*, *9*, 196–207. https://doi.org/10.1111/j.1755-0998.2009.02645.x

Southwick, E. E., & Southwick, L. (1992). Estimating the Economic Value of Honey Bees (Hymenoptera: Apidae) as Agricultural Pollinators in the United States. *Journal of Economic Entomology*, *85*, 621–633. https://doi.org/10.1093/jee/85.3.621

Spens, J., Evans, A. R., Halfmaerten, D., Knudsen, S. W., Sengupta, M. E., Mak, S. S. T., … Hellström, M. (2016). Comparison of capture and storage methods for aqueous macrobial eDNA using an optimized extraction protocol: advantage of enclosed filter. *Methods in Ecology and Evolution*, *8*, 635–645. https://doi.org/10.1111/2041-210X.12683

Szitenberg, A., John, M., Blaxter, M. L., & Lunt, D. H. (2015). ReproPhylo: An Environment for Reproducible Phylogenomics. *PLoS Computational Biology*, *11*, e1004447. https://doi.org/10.1371/journal.pcbi.1004447

Taberlet, P., Coissac, E., Pompanon, F., Brochmann, C., & Willerslev, E. (2012). Towards next-generation biodiversity assessment using DNA metabarcoding. *Molecular Ecology*, *21*, 2045–2050. https://doi.org/10.1111/j.1365-294X.2012.05470.x

Thalinger, B., Deiner, K., Harper, L. R., Rees, H. C., Blackman, R. C., Sint, D., … Bruce, K. (2021). A validation scale to determine the readiness of environmental DNA assays for routine species monitoring. *Environmental DNA*, *3*, 823–836. https://doi.org/10.1002/edn3.189

Thomann, M., Imbert, E., Devaux, C., & Cheptou, P.-O. (2013). Flowering plants under global pollinator decline. *Trends in Plant Science*, *18*, 353–359. https://doi.org/10.1016/j.tplants.2013.04.002

Thomas, C. D., Jones, T. H., & Hartley, S. E. (2019). "Insectageddon": A call for more robust data and rigorous analyses. *Global Change Biology*, *25*, 1891–1892. https://doi.org/10.1111/gcb.14608

Thomsen, P. F., & Sigsgaard, E. E. (2019). Environmental DNA metabarcoding of wild flowers reveals diverse communities of terrestrial arthropods. *Ecology and Evolution*, *9*, 1665-1679. https://doi.org/10.1002/ece3.4809

Tur, C., Castro-Urgal, R., & Traveset, A. (2013). Linking plant specialization to dependence in interactions for seed set in pollination networks. *PLoS ONE*, *8*, e78294. https://doi.org/10.1371/journal.pone.0078294

Valentini, A., Taberlet, P., Miaud, C., Civade, R., Herder, J., Thomsen, P. F., … Dejean, T. (2016). Next-generation monitoring of aquatic biodiversity using environmental DNA metabarcoding. *Molecular Ecology*, *25*, 929–942. https://doi.org/10.1111/mec.13428

Valentin, R. E., Fonseca, D. M., Gable, S., Kyle, K. E., Hamilton, G. C., Nielsen, A. L., & Lockwood, J. L. (2020). Moving eDNA surveys onto land: Strategies for active eDNA aggregation to detect invasive forest insects. *Molecular Ecology Resources*, *20*, 746–755. https://doi.org/10.1111/1755-0998.13151

Valsecchi, E., Arcangeli, A., Lombardi, R., Boyse, E., Carr, I. M., Galli, P., & Goodman, S. J. (2021). Ferries and Environmental DNA: Underway Sampling From Commercial Vessels Provides New Opportunities for Systematic Genetic Surveys of Marine Biodiversity. *Frontiers in Marine Science*, *8*, 1136. https://doi.org/10.3389/fmars.2021.704786

Vamos, E. E., Elbrecht, V., & Leese, F. (2017). Short COI markers for freshwater macroinvertebrate metabarcoding. *Metabarcoding and Metagenomics*, *1*, e14625. https://doi.org/10.3897/mbmg.1.14625

Vanbergen, A. J., & the Insect Pollinators Initiative. (2013). Threats to an ecosystem service: pressures on pollinators. *Frontiers in Ecology and the Environment*, *11*, 251–259. https://doi.org/10.1890/120126

Wake, D. B., & Vredenburg, V. T. (2008). Are we in the midst of the sixth mass extinction? A
view from the world of amphibians. *Proceedings of the National Academy of Sciences of the*
United States of America, 105, 11466–11473. https://doi.org/10.1073/pnas.0801921105

Weigand, A. M., & Macher, J.-N. (2018). A DNA metabarcoding protocol for hyporheic
freshwater meiofauna: Evaluating highly degenerate COI primers and replication strategy.
Metabarcoding and Metagenomics, *2*, e26869. https://doi.org/10.3897/mbmg.2.26869

Weiner, C. N., Werner, M., Linsenmair, K. E., & Blüthgen, N. (2014). Land-use impacts on
plant-pollinator networks: interaction strength and specialization predict pollinator declines.
Ecology, *95*, 466–474. https://doi.org/10.1890/13-0436.1

Wickham, H. (2016). *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag New York.
https://ggplot2.tidyverse.org

Wilcox, T. M., McKelvey, K. S., Young, M. K., Engkjer, C., Lance, R. F., Lahr, A., …
Schwartz, M. K. (2020). Parallel, targeted analysis of environmental samples via high-
throughput quantitative PCR. *Environmental DNA*, *2*, 544–553.
https://doi.org/10.1002/edn3.80

Williams, K. E., Huyvaert, K. P., Vercauteren, K. C., Davis, A. J., & Piaggio, A. J. (2018).
Detection and persistence of environmental DNA from an invasive, terrestrial mammal.
Ecology and Evolution, *8*, 688–695. https://doi.org/10.1002/ece3.3698

Winfree, R., Gross, B. J., & Kremen, C. (2011). Valuing pollination services to agriculture.
Ecological Economics: The Journal of the International Society for Ecological Economics,
71, 80–88. https://doi.org/10.1016/j.ecolecon.2011.08.001

Young, B. E., Auer, S., Ormes, M., Rapacciuolo, G., Schweitzer, D., & Sears, N. (2017). Are
pollinating hawk moths declining in the Northeastern United States? An analysis of collection
records. *PLoS ONE*, *12*, e0185683. https://doi.org/10.1371/journal.pone.0185683

Zeale, M. R. K., Butlin, R. K., Barker, G. L. A., Lees, D. C., & Jones, G. (2011). Taxon-specific
PCR for DNA barcoding arthropod prey in bat faeces. *Molecular Ecology Resources*, *11*,
236–244. https://doi.org/10.1111/j.1755-0998.2010.02920.x

1120

Table 1. Summary of analyses statistically comparing homogeneity of multivariate dispersions (MVDISP) between the communities of different flower species (ANOVA) and variation in community composition of eDNA samples from different flower species (PERMANOVA). Relative contributions of taxon turnover and nestedness-resultant to total beta diversity (Jaccard dissimilarity) for each flower species are given in brackets. P-values in bold indicate statistical significance at 0.05.

	Homogeneity of multivariate dispersions (ANOVA)			Community similarity (PERMANOVA)			
	Mean distance to centroid ± SE	df	F	P	df	F	R^2	P
Greenhouse							
Turnover		3	0.886	0.454	3	2.897	0.139	**0.007**
Cynoglossum amabile (100.00%)	0.333 ± 0.333							
Monarda (96.70%)	0.531 ± 0.056							
Penstemon (97.65%)	0.333 ± 0.232							
Solanum (95.60%)	0.370 ± 0.134							
Nestedness-resultant		3	0.815	0.491	3	-1.854	-0.115	0.992
Cynoglossum amabile (0.00%)	0.000 ± 0.000							
Monarda (3.30%)	0.111 ± 0.032							
Penstemon (2.35%)	0.153 ± 0.039							
Solanum (4.40%)	0.157 ± 0.026							
Total beta diversity		3	0.772	0.515	3	2.284	0.113	**0.005**
Cynoglossum amabile (100.00%)	0.333 ± 0.333							
Monarda (100.00%)	0.595 ± 0.012							
Penstemon (100.00%)	0.472 ± 0.197							
Solanum (100.00%)	0.472 ± 0.060							
UIUC campus							
Turnover		1	2.570	0.111	1	7.999	0.044	**0.001**
Monarda (98.40%)	0.586 ± 0.012							
Penstemon (99.55%)	0.418 ± 0.163							
Nestedness		1	4.262	**0.041**	1	-7.080	-0.043	1.000
Monarda (1.60%)	0.127 ± 0.013							
Penstemon (0.45%)	0.207 ± 0.021							
Total beta diversity		1	1.406	0.237	1	5.721	0.032	**0.001**
Monarda (100.00%)	0.612 ± 0.005							
Penstemon (100.00%)	0.498 ± 0.138							

Table 2. Summary of analyses statistically comparing homogeneity of multivariate dispersions (MVDISP) between the communities produced by different sample types (ANOVA) and variation in community composition of eDNA samples sourced from different plant material (PERMANOVA). Relative contributions of taxon turnover and nestedness-resultant to total beta diversity (Jaccard dissimilarity) for each sample type are given in brackets. P-values in bold indicate statistical significance at 0.05.

	Homogeneity of multivariate dispersions (ANOVA)				Community similarity (PERMANOVA)			
	Mean distance to centroid ± SE	df	F	*P*	df	F	R^2	*P*
Greenhouse								
Turnover		2	1.115	0.335	2	2.166	0.073	0.061
Nectar (100.00%)	0.333 ± 0.333							
Swab (97.88%)	0.582 ± 0.038							
Whole flower (98.51%)	0.397 ± 0.198							
Nestedness		2	1.816	0.172	2	-0.402	-0.015	0.816
Nectar (0.00%)	0.000 ± 0.000							
Swab (2.12%)	0.129 ± 0.012							
Whole flower (1.49%)	0.168 ± 0.029							
Total beta diversity		2	0.825	0.444	2	1.868	0.064	0.053
Nectar (100.00%)	0.333 ± 0.333							
Swab (100.00%)	0.591 ± 0.039							
Whole flower (100.00%)	0.508 ± 0.115							
UIUC campus								
Turnover		1	4.730	**0.031**	1	-0.950	-0.006	0.999
Swab (99.27%)	0.333 ± 0.225							
Whole flower (99.44%)	0.469 ± 0.118							
Nestedness		1	0.307	0.580	1	13.989	0.075	**0.002**
Swab (0.73%)	0.160 ± 0.046							
Whole flower (0.56%)	0.176 ± 0.022							
Total beta diversity		1	1.478	0.226	1	0.774	0.005	0.654
Swab (100.00%)	0.478 ± 0.198							
Whole flower (100.00%)	0.544 ± 0.061							

Table 3. Summary of analyses statistically comparing homogeneity of multivariate dispersions
(MVDISP) between the communities produced by different eDNA isolation protocols (ANOVA)
and variation in community composition of eDNA samples preserved and extracted using
different methods (PERMANOVA). Relative contributions of taxon turnover and nestedness-
resultant to total beta diversity (Jaccard dissimilarity) for each eDNA isolation protocol are given
in brackets. P-values in bold indicate statistical significance at 0.05.

	Homogeneity of multivariate dispersions (ANOVA)				Community similarity (PERMANOVA)			
	Mean distance to centroid ± SE	df	F	*P*	df	F	R^2	*P*
Greenhouse								
Turnover		1	0.673	0.416	1	1.446	0.025	0.226
ATL/Qiagen (98.47%)	0.429 ± 0.171							
CTAB/PCI (97.90%)	0.509 ± 0.093							
Nestedness		1	0.035	0.852	1	-1.257	-0.023	0.859
ATL/Qiagen (1.53%)	0.138 ± 0.024							
CTAB/PCI (2.10%)	0.146 ± 0.028							
Total beta diversity		1	0.479	0.492	1	1.150	0.020	0.303
ATL/Qiagen (100.00%)	0.524 ± 0.091							
CTAB/PCI (100.00%)	0.572 ± 0.043							
UIUC campus								
Turnover		1	3.602	0.059	1	0.799	0.005	0.582
ATL/Qiagen (99.48%)	0.463 ± 0.140							
CTAB/PCI (99.12%)	0.346 ± 0.190							
Nestedness		1	0.019	0.890	1	4.938	0.028	0.065
ATL/Qiagen (0.52%)	0.187 ± 0.022							
CTAB/PCI (0.88%)	0.191 ± 0.032							
Total beta diversity		1	2.019	0.157	1	1.255	0.007	0.208
ATL/Qiagen (100.00%)	0.545 ± 0.075							
CTAB/PCI (100.00%)	0.470 ± 0.183							

Figure 1. Images depicting greenhouse experiment and flower morphology, size, and characteristics of our focal flower species: *Penstemon* sp., *Monarda* sp., and *Solanum* sp.. We established a mixed-species colony of flowering plants to ensure a constant supply of pollen and nectar **(a)**. A subset of plants was then moved to an isolated greenhouse room where a colony of common eastern bumblebees was established **(b)**. The large, conical flowers of *Penstemon* **(c)** require pollinators to crawl in to access nectar/pollen. *Monarda* **(d)** inflorescences are composed of crowded, head-like clusters of flowers providing nectar and pollen. *Solanum* **(e)** inflorescences are considerably smaller and star-shaped with exposed stamens but no nectar. Photos by Mark Davis and Tiffany Jolley.

Figure 2. Tile plot depicting the results of *in vitro* primer validation. The subset of primers with approximately appropriate annealing temperature and fragment length were tested with 11 locally occurring bee species using PCR and verified via gel electrophoresis. Blue cells represent comparisons in which all three PCR replicates were positive, yellow cells represent comparisons in which one or two PCR replicates were positive, and grey cells represent comparisons in which all three PCR replicates failed.

Figure 3. Summaries of alpha and beta diversity comparisons made between flower species sampled during **(a)** greenhouse and **(b)** field experiments: **(i)** boxplot showing the number of taxa detected in samples from each flower species, followed by non-metric multidimensional scaling (NMDS) plots of invertebrate communities for **(ii)** turnover, **(iii)** nestedness-resultant, and **(iv)** total beta diversity. Boxes show 25th, 50th, and 75th percentiles, and whiskers show 5th and 95th percentiles.

1174
1175
Figure 4. Summaries of alpha and beta diversity comparisons made between sample types
collected during **(a)** greenhouse and **(b)** field experiments: **(i)** boxplot showing the number of
taxa detected in samples from each sample type, followed by non-metric multidimensional
scaling (NMDS) plots of invertebrate communities for **(ii)** turnover, **(iii)** nestedness-resultant,
and **(iv)** total beta diversity. Boxes show 25th, 50th, and 75th percentiles, and whiskers show 5th
and 95th percentiles.
1182

Figure 5. Summaries of alpha and beta diversity comparisons made between methods used for
preservation and extraction of samples collected during **(a)** greenhouse and **(b)** field
experiments: **(i)** boxplot showing the number of taxa detected in samples from each sample type,
followed by non-metric multidimensional scaling (NMDS) plots of invertebrate communities for
(ii) turnover, **(iii)** nestedness-resultant, and **(iv)** total beta diversity. Boxes show 25th, 50th, and
75th percentiles, and whiskers show 5th and 95th percentiles.

1191

	FLOWER MORPHOLOGY	SAMPLING METHOD	PRESERVATION/ EXTRACTION	PRIMER SET	BIOINFORMATICS PIPELINE
OPTIMAL WORKFLOW	Replicate flowers across diverse flower species assemblage	Whole flower head harvesting	ATL preservation + modified Qiagen DNeasy Blood and Tissue extraction	Suite of multiple COI primer sets	metaBEAT* *Available pipelines should be evaluated for each new study
OTHER APPROACHES EXAMINED	*Monarda* *Penstemon* *Solanum* (whole flower head harvest only) *Cynglossum amabile* (nectar only)	Whole flower head harvesting Swabbing Nectar extraction	CTAB preservation + Phenol-Chloroform-Isoamyl extraction ATL preservation + modified Qiagen DNeasy Blood and Tissue extraction	COI 16S	Anacapa metaBEAT

1192
1193
1194 **Figure 6.** A recommended workflow for pollinator eDNA metabarcoding. This workflow was
1195 developed based on the results of this study and includes technical recommendations that should
1196 yield the most robust estimate of pollinator communities.
1197