A single-cell massively parallel reporter assay detects cell type specific cis-regulatory activity

Siqi Zhao[1,2], Clarice KY Hong[1,2], Connie A Myers[3], David M Granas[1,2], Michael A White[1,2], Joseph C Corbo[3], Barak A Cohen[1,2]

[1]Edison Family Center for Systems Biology and Genome Sciences, [2]Department of Genetics, Washington University School of Medicine, USA. [3]Department of Pathology, Washington University School of Medicine, USA.

Massively parallel reporter gene assays are key tools in regulatory genomics, but cannot be used to identify cell-type specific regulatory elements without performing assays serially across different cell types. To address this problem, we developed a single-cell massively parallel reporter assay (scMPRA) to measure the activity of libraries of *cis*-regulatory sequences (CRSs) across multiple cell-types simultaneously. We assayed a library of core promoters in a mixture of HEK293 and K562 cells and showed that scMPRA is a reproducible, highly parallel, single-cell reporter gene assay that detects cell-type specific *cis*-regulatory activity. We then measured a library of promoter variants across multiple cell types in *ex vivo* mouse retinas and showed that subtle genetic variants can produce cell-type specific effects on *cis*-regulatory activity. We anticipate that scMPRA will be widely applicable for studying the role of CRSs across diverse cell types.

Introduction

The majority of heritable variation for human diseases maps to the non-coding portions of the genome[1–6]. This observation has led to the hypothesis that genetic variation in the *cis*-regulatory sequences (CRSs) that control gene expression underlies a large fraction of disease burden[7–10]. Because many CRSs function only in specific cell types[11], there is intense interest in high-throughput assays that can measure the effects of cell-type-specific CRSs and their genetic variants.

Massively Parallel Reporter Assays (MPRAs) are one family of techniques that allow investigators to assay libraries of CRSs and their non-coding variants *en masse*[12–18]. In an MPRA experiment, every CRS drives a reporter gene carrying a unique DNA barcode in its 3' UTR, which allows investigators to quantify the activity of each CRS by the ratio of its barcode abundances in the output RNA and input DNA. This approach allows investigators to identify new CRSs, assay the effects of non-coding variants, and discover general rules governing the functions of CRSs[12,19–23]. One limitation of MPRAs is that they are generally performed in monocultures, or as bulk assays across the cell types of a tissue. Performing cell-type specific MPRAs in tissues will require methods to simultaneously readout reporter gene activities and cell type information in heterogeneous pools of cells.

To address this problem, we developed scMPRA, a procedure that combines single-cell RNA sequencing with MPRA. scMPRA simultaneously measures the activities of reporter genes

in single cells and the identities of those cells using their single-cell transcriptomes. The key
component of scMPRA is a two-level barcoding scheme that allows us to measure the copy
number of all reporter genes present in a single cell from mRNA alone. A specific barcode
marks each CRS of interest (CRS barcode, "cBC") and a second random barcode (rBC) acts as
a proxy for DNA copy number of reporter genes in single cells (**Fig. 1a**). The critical aspect of
the rBC is that it is complex enough to ensure that the probability of the same cBC-rBC
appearing in the same cell more than once is vanishingly small. In this regime, the number of
different cBC-rBC pairs in a single cell becomes an effective proxy for the copy number of a
CRS in that cell. Even if a cell carries reporter genes for multiple different CRSs, and each of
those reporter genes is at a different copy number, we can still normalize each reporter gene in
each individual cell to its plasmid copy number. With this barcoding scheme, we can measure
the activity of many CRSs with different input abundances at single-cell resolution, which allows
us to measure the activity of CRSs simultaneously across different populations of cells.

54   Results

scMPRA enables single-cell measurement of CRS activity
As a proof of principle, we first used scMPRA to test whether different classes of core promoters
show different activities in different cell types. Core promoters are the non-coding sequences
that surround transcription start sites, where general cofactors interact with RNA polymerase
II[24,25]. Core promoters are divided into different classes by the functions of their host genes
(housekeeping vs developmental), as well as by the sequence motifs they contain (TATA-box,
downstream promoter element (DPE), and CpG islands). We selected 676 core promoters that
we previously tested[24,25] and cloned them into a double-barcoded MPRA library. Given the
complexity of the library (>1x10^7 unique cBC-rBC pairs), we calculated that the probability of
plasmids with the same cBC-rBC pair occurring in the same cell is less than 0.01 with our
transfection protocols. Given this low likelihood, the number of rBC per cBC in a cell represents
the copy number of a CRS in that cell. Knowing the copy number of CRSs in single cells allows
us to normalize reporter gene expression from each CRS to its copy number in individual cells.
68   We performed a cell mixing experiment to test whether scMPRA could measure cell
69   type specific expression of reporter genes. We transfected K562 (chronic myelogenous
70   leukemia) and HEK293 (human embryonic kidney) cells, and performed scMPRA on a 1:1
71   mixture of those cell lines (**Fig. 1b**). The mRNA from single cells was captured, converted to
72   cDNA, and sequenced. The resulting cBC-rBC abundances and transcriptome of each single
73   cell are linked by their shared 10x cell barcode.

We recovered a total of 3112 cells (97%) that could be unambiguously assigned to one
of the two cell types (**Fig. 2a, Supplementary Fig. 1a,b**) and computed the mean expression of
each core promoter in the library in each cell type (**Methods**). The measurements were
reproducible in both cell types (K562: Pearson's R = 0.89, Spearman's ρ = 0.57, HEK293:
Pearson' R = 0.96, Spearman's ρ = 0.92) (**Figs. 2b,c, Supplementary Table 1**), and we
obtained measurements for 99.5% of core promoters in K562 cells and 100% in HEK293 cells,
highlighting the efficiency of scMPRA. The median number of cells in which each core promoter
was measured was 76 for K562 cells and 287 for HEK293 cells (**Figs. 2d,e**). We also tabulated
the number of cBC-rBC pairs in each individual cell and found that the median per cell was 164
in K562 cells and 341 in HEK293 cells (**Supplementary Figs. 1c,d**). On average we detected
10 rBCs per promoter in individual HEK293 cells and 2 rBCs per promoter in K562 cells
(**Supplementary Fig. 1e,f**). To validate the scMPRA measurements, we conducted bulk MPRA
of the core promoter library in the two cell types separately. Bulk MPRA measurements are not
corrected for PCR amplification biases with UMIs, and we found that the bulk measurements
correlate well with aggregated single-cell measurements without UMI correction (**Figs. 2f,g**).
That correlation drops with the UMI-corrected single-cell measurements (**Supplementary Figs.**
1g,h), which suggests that bulk measurements may suffer from over counting because of
uneven amplification during PCR.

101
Fig. 2. scMPRA detects cell type specific CRS activity. (**a**) UMAP of the transcriptome from the mixed-cell scMPRA experiment.
3312 out of 3417 cells are assigned to either K562 or HEK293 cells. (**b,c**) Reproducibility of replicate measurements of the mean
expression from each core promoter in both K562 and HEK293 cells (**d,e**) Histogram of the number of cells in which each core
promoter was measured for HEK293 and K562 cells. (**f,g**) Correlations between scMPRA and bulk MPRA using mRNA
abundances (cBC counts per cell) to make the two methods comparable.(**h**) Boxplot of the activities of core promoters from different
categories in K562 (orange) and HEK293 (blue) cells. The promoter categories are taken from Haberle et al. Because the average
expression of all promoters were different between K562 and HEK293, we plotted each category according to its deviation from the
average expression (z-score) of all promoters in each cell type (**i**) Volcano plot for differential expression (DE) of core promoters in
K562 and HEK293 cells. Significant DE reporters (red dots) have adjusted p-value <0.01 and log-2 fold change greater than 0.3. (**j**)
Venn diagram of the functional characterization (housekeeping vs developmental) of down-regulated core promoters in K562 cells.
Housekeeping promoters are enriched (p-value = 1.08×10^{-11} from hypergeometric test). (**k**) Pie chart of the sequence features (CpG,
DPE, TATA) of down-regulated core promoters. CpG promoters are enriched (p=2.18×10^{-6}, from hypergeometric test).

scMPRA detects cell type specific CRS activity

We asked whether the data allowed us to detect core promoters with differential activity
between K562 and HEK293 cells. While different classes of core promoters generally had
similar activities in both cell lines (**Fig. 2h**), our differential analysis using DEseq2[26] identified a
small number of promoters (11 out of 669) that are upregulated in K562 cells, and 59 promoters
that are downregulated in K562 cells (adjusted p-value < 0.01, log2 fold change > 0.3, **Fig. 2i,**
Supplementary Table 2). Among the down-regulated promoters, 48 out of 59 core promoters
belong to housekeeping genes (p=1.08×10^{-11}, **Fig. 2j**), and 46 out of 59 core promoters are
CpG-island-containing core promoters (p=2.18×10^{-6}, **Fig. 2k**). This result is not due to
differences in the quality of measurements between housekeeping and developmental

promoters (**Supplementary Figs. 1i,j**). These results demonstrate the ability of scMPRA to
detect CRSs with cell-type specific activities.
127
scMPRA detects cell sub-state specific CRS activity
Single-cell studies have revealed heterogeneity in cell states even within isogenic cell
types[27–30]. Therefore, we asked if scMPRA can identify CRSs with cell-state specific activity. We
repeated scMPRA on K562 cells alone and obtained a total of 5141 cells from two biological
replicates. Measurements of each library member were again highly correlated between
replicates and agree well with independent bulk measurement (**Supplementary Figs. 2a,b**).
Because the phases of the cell cycle represent distinct cell-states, we asked whether
scMPRA could identify reporter genes with differential activity through the cell cycle. We
assigned cell cycle phases to each cell using their single cell transcriptome data (**Fig. 3a**) and
calculated the mean expression of each reporter gene in different cell cycle phases. We found
that most core promoters in our library are upregulated in the G1 phase of the cell cycle, and
that some housekeeping promoters are highly expressed through all cell cycle phases (**Fig. 3b,**
Supplementary Table 3). We also identified core promoters with different expression dynamics
through the cell cycle. For example, we found that the core promoter of *UBA52* remains highly
expressed in the S phase, whereas the core promoter of *CXCL10* is lowly expressed throughout
(**Supplementary Fig. 2c**). This analysis illustrates the ability of scMPRA to identify CRSs
whose expression naturally fluctuates with cellular dynamics.
We then asked whether scMPRA could detect reporter genes with activities that were
specific to other cell-states in K562 cells, after normalizing for cell cycle effects. We focused on
two specific sub-states that have been reported and experimentally validated for high
proliferation rates in K562 cells[31,32]. The first is the CD34$^+$/CD38$^-$ sub-state that has been
identified as a leukemia stem cell subpopulation, and the second is the CD24$^+$ sub-state that is
linked to selective activation of proliferation genes by bromodomain transcription factors[28,29]. To
identify these sub-states in our single-cell transcriptome data, we first regressed out the cell
cycle effects and confirmed that the single cell transcriptome data no longer clustered by cell
cycle phase (**Supplementary Fig. 2d**). We then identified clusters within K562 cells that have
the CD34$^+$/CD38$^-$ expression signature, or the CD24$^+$ signature (**Fig. 3c**). Although the
CD34$^+$/CD38$^-$ cells represent only 9.3% of the cells, scMPRA revealed two distinct classes of
core promoters that are upregulated and downregulated in these cells relative to the CD24+ and
"differentiated" clusters (**Fig. 3d**). Conversely, the expression patterns of promoters are similar
between the CD24$^+$ and "differentiated" clusters (**Fig. 3d, Supplementary Table 4**). Motif
analysis of the up/down regulated classes of promoters in CD34$^+$/CD38$^-$ cells showed that
different core promoter motifs are enriched in each class, with the TATA box and Motif 5 being
enriched in the upregulated class and MTE and TCT motifs being enriched in downregulated
class (**Fig. 3e**). This result suggests that differences in core promoter usage might be driving the
differences between CD34$^+$/CD38$^-$ and the other clusters. Because the TATA box is mostly
found in developmental core promoters, the CD34$^+$/CD38$^-$ subpopulation likely reflects the more
"stem-like" cellular environment in these cells. Our analysis highlights the ability of scMPRA to
identify CRSs with differential activity in rare cell populations.
167
168

Fig. 3. scMPRA detects sub-state-specific CRS activity. (**a**) PCA plot of K562 cells classified by their cell cycle scores. (**b**) Heatmap of core promoter activities in different cell cycle phases (Color bar indicates housekeeping (blue) vs developmental (red) promoters). Core promoter activities have been normalized within each cell cycle phase to highlight the differences between housekeeping and developmental promoters. (**c**) UMAP embedding of K562 cells with high proliferation sub-states (CD34+/CD38- and CD24+). (**d**) Hierarchical clustering showing two clusters ("up" and "down") based on expression patterns in the three substates. The promoter activities are plotted as their z-score from the average across cell states to highlight the difference between cell states. (**e**) Proportion of promoters in the up and down clusters that contain the indicated core promoter motif. * represents significant enrichment in one cluster over the other ($p < 0.05$, Fisher's exact test).

scMPRA is reproducible and accurate in murine retinas

To demonstrate that scMPRA is applicable in a complex tissue with multiple cell types, we performed experiments in explanted murine retinas. Intact retina from newborn mice can be cultured and transfected *ex vivo*. This system has been useful for bulk MPRA experiments[13,19,33], but the results from those experiments report the aggregate expression of library members across the cell types of the retina. Performing scMPRA in *ex vivo* retina provided a chance to assay an MPRA library in a living tissue with multiple cell types in their proper three-dimensional organization.

For this analysis we designed a library consisting of two independently synthesized wild-
type copies, and 113 variants, of the full-length *Gnb3* promoter (115 library members,
Supplementary Table 5). We chose the *Gnb3* promoter because it has high activity in
photoreceptors and bipolar cells, but lower expression in other interneurons (i.e. amacrine cells)
and Müller glia cells. The library contains mutations in the known transcription factor binding
sites (TFBSs) in the *Gnb3* promoter as well as mutations that scan across two phylogenetically
conserved regions of the promoter (details in Fig. 6 below). We constructed this library of *Gnb3*
promoter variants using the double barcoding strategy described above, with one key
modification that we now describe.
198
In the *Gnb3* promoter library we addressed the inability of scMPRA to measure silent
library members. In the first iteration of scMPRA, when a library member produces no mRNA
barcodes its corresponding plasmid cannot be detected, and thus, a cell containing a silent
plasmid is indistinguishable from a cell without a plasmid. This was not a limitation in our
experiments with K562 and HEK293 cells because we detected expressed mRNAs from every
member of the library. However, to avoid this potential problem in our retina experiments, we
included an additional cassette on the *Gnb3* promoter library that allows us to detect the
presence of plasmids carrying silent promoter variants. We included a cassette in which the U6
promoter drives the expression of a second copy of the cBC coupled to the 10x Capture
Sequence (**Fig. 4a**). The U6 promoter drives strong RNA Polymerase III-dependent
transcription, and is independent of the activity of the *Gnb3* promoter. The Capture Sequence is
a specific sequence that is typically used to identify gRNAs in Perturb-seq experiments[34], but we
use it here to isolate U6 expressed cBCs (**Fig. 4b**). When a cell contains a U6 cBC without the
corresponding *Gnb3* promoter cBC, it indicates the presence of a silent library member.
213
We introduced the *Gnb3* promoter variant library into newborn mouse retinas and
assessed the cell types into which the library entered by scRNA-seq (**Fig. 4c**). We obtained a
total of 22,161 cells from two replicate experiments with a mean of 22,528 reads per cell and
1,642 genes per cell. The scRNA-seq data showed that we recovered rod photoreceptors
(87.3%), bipolar cells (3.5%), interneurons (i.e. amacrine cells) (5.2%), and Müller glia cells
(3.9%) (**Fig. 4d, Supplementary Figure 3 a,b**).
220
We then computed the expression of each *Gnb3* promoter variant in each cell type by
sequencing the *Gnb3*-expressed barcodes and the U6 barcodes from single cells. Cells with
U6-expressed cBC counts, but no *Gnb3*-expressed cBC counts, represented cells in which that
promoter variant was silent. On average, *Gnb3* promoter variants were silent in 22% of cells, but
this number varied widely (**Fig. 4e**) and was linearly related to the strength of the promoter, with
stronger promoters

Fig. 4. scMPRA design and workflow in mouse retina. (**a**) Schematic of *Gnb3* promoter library constructs. In addition to the cBC and rBC barcodes, the *Gnb3* promoter library contains an additional cassette in which the constitutive U6 promoter expresses a second copy of the cBC with a capture sequence for isolating these transcripts on gel beads. (**b**) Two different types of transcripts produced from the *Gnb3* promoter library to measure promoter expression and detect unexpressed promoters respectively. The two types of transcripts originating from the same cell share the same 10x cell barcodes (**c**) Experimental workflow for scMPRA in *ex vivo* mouse retinas. (**d**) UMAP of all cells measured in scMPRA with four major cell types identified. (**e**) For each *Gnb3* variant in the library, we determined the proportion of cells that contain barcoded poly(A) transcripts out of all the cells that contained the variant. (**f**) Reproducibility of promoter activities between replicates in each of the four major cell types.

expressing in a larger fraction of cells (**Supplementary Fig. 3c**). Using both the *Gnb3*-driven and U6-driven counts allowed us to compute the average expression of a promoter variant across all the cells of a given cell type, while still accounting for cells in which that promoter variant is silent (**Methods**).

Replicate measurements of the *Gnb3* promoter variant library were reproducible in all four cell types (**Fig. 4f, Supplementary Table 6**). Reproducibility was highest in rod cells (Spearman's ρ: 0.97, Pearson's R: 0.98) because rod cells are the most abundant cell type in the mouse retina. The reproducibility was slightly lower in the rarer cell types (bipolar cells: Spearman's ρ: 0.88, Pearson's R: 0.92, Müller glia: Spearman's ρ: 0.93, Pearson's R: 0.95, and

interneurons: Spearman's ρ: 0.95, , Pearson's R: 0.98), but remained high enough to assess the
expression of individual library members. We determined how reproducibility scales with the
number of cells in scMPRA by subsampling the expression data. The minimum number of cells
required for reproducible measurements (Spearman's ρ > 0.75) of mean reporter gene levels is
75 cells (**Supplementary Fig. 3d**). Our results show that scMPRA works well for measuring
reporter gene levels across cell types in complex tissues using small numbers of cells.

253
254   **Fig. 5. scMPRA recapitulates *Gnb3* expression patterns.** (**a**) The expression of the wild-type *Gnb3* promoter in scMPRA reflects
255   endogenous expression levels of *Gnb3* in the respective cell types (Error bar denotes 95% C.I from two biological replicates). (**b**)
256   The expression of the entire *Gnb3* library in different cell types also follows endogenous *Gnb3* expression (****: p-value < 0.0001 for
257   Mann-Whitney U test). (**c**) scMPRA recapitulates the effects of a known *Gnb3* variant, where the CRX3[Q50]/CRX5[Q50] variant reduces
258   expression in bipolar cells specifically (*: p-value < 0.05 for Welch's t-test).
259
Two additional observations suggest that scMPRA measurements are accurate in *ex*
vivo retinas. First, the expression of the wild type *Gnb3* reporter, as well as the average
expression of all *Gnb3* promoter variants, correlates with endogenous *Gnb3* expression in the
corresponding scRNA-seq data (**Fig. 5a,b**). Second, our scMPRA data reproduced the known
effect of a cell type specific *Gnb3* promoter variant. Murphy *et al.*[42] showed that altering two of
the K50 homeobox sites in the *Gnb3* promoter to Q50 sites reduces expression in bipolar cells
while leaving expression in rod cells relatively unaffected. We observed the same reduction in
bipolar cells when compared with rod cells for this same mutant (**Fig. 5c**). In addition, scMPRA
also revealed that this mutant shows increased activity in Müller glia and interneurons. Taken
together, these observations demonstrate that scMPRA is reproducible and accurate when
applied to cell types in a complex tissue.
271
272   **scMPRA reveals cell-type specific promoter variants.**
273
The *Gnb3* library was designed to probe components of the promoter including five
binding sites for K50-type homeodomain TFs, an E-box binding site, and two evolutionarily
conserved regions (**Fig. 6a**). In this experiment we define the effect of a mutation as its relative
fold-change to the WT *Gnb3* promoter in each cell type because the *Gnb3* promoter is
expressed at different levels across cell types (**Fig. 5a**). We labeled the homeobox sites as
Cone Rod Homeobox (CRX) sites because *CRX* is a K50-homeodomain protein which plays an
important role in rods and bipolar cells and is required for *Gnb3* expression. Inactivating
mutations in any individual CRX site decreased *Gnb3* reporter expression in bipolar and rod
cells, but deletion of either CRX1 or CRX5 also resulted in increased expression in interneurons
(**Fig. 6b**). The CRX2 disruption had the largest effect on expression, and mutating the CRX2

site in combination with any other CRX site also caused large reductions in expression in rods
and bipolar cells. Single and double swaps of K50 CRX binding sites with Q50 binding sites
tended to yield cell-type specific effects, primarily because interneurons displayed larger
responses to the Q50 swaps compared with rod and bipolar cells (**Fig. 6c**). Increasing the
affinity of CRX sites tended to have mild effects on expression in rods and bipolar cells, but
increased expression significantly in interneurons (**Fig. 6d**). The results from modifying CRX
sites demonstrated that perturbations to single binding sites can produce cell-type specific
effects.

292
Fig. 6. Mutations in the *Gnb3* promoter display cell-type specific effects. (**a**) Schematic of the *Gnb3* promoter showing the
location of the five CRX binding sites and the E-box. (**b**) Effects of individual and pairwise deletions of CRX binding sites. (**c**) Effects
of individual and pairwise mutations of CRX K50 binding sites to Q50 binding sites. (**d**) Effects of changing CRX binding site
affinities. (**e**) Effects of saturation mutagenesis of the E-box. (**f**) Effects of shuffle mutants in conserved regions of the *Gnb3*
promoter. Each region was split into 5bp windows and the nucleotides in each window were shuffled. Labels above the heatmap
indicate locations where the mutations impact CRX or E-box binding sites. All plots show log2 fold changes of the mutant relative to
WT *Gnb3* expression in that cell type. Stars above the plot indicate a significant cell-type specific effect by one-way ANOVA.

We next examined the effects of single nucleotide changes in the E-box binding site
(**Fig. 6e**). Helix-Loop-Helix (bHLH) transcription factors, which bind E-box motifs, are critical for
the development of multiple retinal cell types[35]. Several single-nucleotide substitutions in the E-
box resulted in strong effects on expression, although only one substitution produced significant
cell-type specific effects. While the E-box is critical for strong expression of the *Gnb3* promoter,
subtle changes to its sequence do not generally result in cell-type specific changes to its
activity.

To examine the effects of more severe sequence changes, and to assess the effects of
perturbations outside the known TFBSs, we tiled mutations through the two evolutionarily
conserved regions shuffling 5 bp at a time (**Fig. 6f**). Mutations in all six TFBSs resulted in cell-
type specific changes in expression, but several mutations in the *Gnb3* promoter outside of the
known TFBS also resulted in cell-type specific changes in expression. Thus, other information in
the *Gnb3* promoter provides important cell-type context for the functioning of the CRX and E-
box motifs.

Our analysis of the *Gnb3* promoter shows that single-binding site and single-nucleotide
variants can result in cell-type specific changes to *cis*-regulation, and that scMPRA is a powerful
tool for identifying these changes across cell-types in mammalian tissues. The *cis*-regulatory
logic of the *Gnb3* promoter keeps it expressed at high levels in rods and bipolar cells in the early
postnatal period, and at much lower levels in interneurons, which we speculate is why most cell-
type specific perturbations result in effects of different sizes in interneurons when compared with
rods and bipolar cells.

Conclusions

We have presented a single-cell MPRA method to measure the cell-type and cell-state
specific effects of CRSs. We demonstrated that scMPRA detects cell-type specific reporter gene
activity in a mixed population of cells as well as in living retinas, and cell-state specific activity in
isogenic K562 cells. The assay is reproducible and reports accurate mean levels of reporter
gene activity in as few as 75 cells in a complex tissue. New methods that increase the number
of single cells measured per experiment[36] will increase the size of libraries that can be assayed
by scMPRA. The dynamic range was relatively small in this study (8-fold between the strongest
and weakest *Gnb3* variants), which may reflect the activity of these specific sequences, but may
also arise from the low efficiency of mRNA capture in single cells. scMPRA will therefore benefit
from continuing improvements of methods to capture and recover mRNA from single cells.

With the burgeoning of Adeno-associated viral delivery systems[37–41], we anticipate that
scMPRA will be widely used to study *cis*-regulatory effects in a variety of complex tissues. Given
the hypothesis that non-coding variants with cell-type specific effects underlie a large fraction of
human disease, an important application of scMPRA will be to identify genetic variants with cell-
type specific *cis*-regulatory effects.

342
343

Methods

Cell Culture

346 K562 cells were cultured in Iscove's Modified Dulbecco's Medium (IMDM) + 10% Fetal Bovine
347 Serum (FBS) + 1% non-essential amino acids + 1% pen/strep at 37□ with 5% CO_2. HEK293
348 cells were cultured in Eagle's Minimum Essential Medium (EMEM) + 10% Fetal Bovine Serum
349 (FBS) + 1% pen/strep at 37□ with 5% CO_2.

Core Promoter Library Cloning

We developed a two-level barcoding strategy to enable single-cell normalization of plasmid copy
number. We applied this strategy to a library of core promoters we previously tested by bulk
MPRA[24]. That core promoter library contains 676 core promoters, each with a length of 133bp.
The library cloning was done in three steps: First, we synthesized a library of 676 core
promoters each barcoded with 10 different cBCs and cloned this library into a backbone[24]. In a
second step, a dsRed fluorescent reporter cassette was cloned between each core promoter
and its associated cBCs as described[24]. Thirdly, we modified this library for scMPRA by adding
random barcodes downstream of the cBCs, but upstream of the polyA site.

To add the random barcodes (rBCs) we synthesized a single-stranded 90 bp DNA
oligonucleotide (oligo) containing a 25 bp random sequence (the rBC), a restriction site, and 30
bp homology to the library vector on each side of the rBC region. We used NEBuilder® HiFi
DNA Assembly Master Mix (E2621) to clone this oligo into the core promoter library. 4µg of the
plasmid library was split into four reactions and digested with 2µl of SalI for 1.5 hours at 37□.
The digested product was run at 100V for 2 hours on a 0.7% agarose gel. The correct size band
was cut and purified with the Monarch Gel Extraction Kit (New England BioLabs T1020L). The
insert single-stranded DNA was diluted to 1 µM with H2O. Three assembly reactions were
pooled together, each reaction containing 100 ng of digested library backbone, 1µM of insert
DNA, 1µl of NEBuffer 2, 10 µl of 2X Hifi assembly mix, and H2O up to 20 µl. The reaction was
incubated at 50°C for 1 hour. The assembled product was purified with the Monarch PCR&DNA
Cleanup kit (New England BioLabs T1030L) and eluted in 12 µl of H2O.

The assembled plasmid was transformed using Gene Pulser Xcell Electroporation Systems by
electroporation (BIO-RAD 1652661) into 50 µl of ElectroMax DH10B electrocompetent cells
(Invitrogen 18290015) with 1 µl of assembled product at 2 kV, 2000 Ω, 25 nF, with 1 mm gap.
950 µl of SOC medium (Invitrogen 15544034) was added to the cuvette and then transferred to
a 15 ml Falcon tube. Two transformations were performed, and each tube was incubated at 37
□ for 1 hour on a rotator with 300 rpm. The culture was then added to pre-warmed 150 µl
LB/Amp medium and grown overnight at 37□. 1 µl of the culture was also diluted 1:100 and 50

µl of the diluted cultured was plated on a LB agar plate to estimate the transformation efficiency.
For the core promoter library, we prepared DNA from more than 4x10[8] colonies. Shallow
sequencing of this library (below) showed that the majority of library members encoded unique
cBC-rBC combinations.

Gnb3 Promoter Variant Library Design and Cloning

The *Gnb3* library was designed to probe components of the promoter including five binding sites
for K50-type homeodomain TFs, an E-box binding site, and two evolutionarily conserved
regions. We labeled the K50 homeobox sites as Cone Rod Homeobox (CRX) sites because
CRX is a K50-homeodomain protein required for *Gnb3* expression and a key-lineage
determining factor in retina, even though other K50-type homeobox proteins are also expressed
in retinas. To test whether the disruption of CRX sites in the *Gnb3* promoter has cell-type
specific effect, we made the following three types of mutations: (1) All individual and pairwise
deletions of the CRX binding sites by mutating [42] the CRX sites to 5'-CTACTCCC-3'. (2) All
individual and pairwise mutations of CRX binding sites from K50 homebox to Q50 homeobox
motifs: 5'-CTAATTAC-3'. (3) All individual mutations of CRX binding sites to high (5'-CTAATCCC-
3'), medium (5'-CTAAGCCC-3') and low affinity (5'-CTTATCCC-3') K50 homeobox sites [22]. Our
unpublished data suggested that the E-box is important for the Gnb3 promoter activity and E-
box motif is bound by many neuronal specific TFs [35], hence we mutated each base pair in the E-
box to every other base pair and made pairwise mutations of the two core base pairs in the E-
box motif. Lastly, we took an unbiased approach to screen for potential cell-type-specific
mutations by shuffling mutations across the two conserved regions in the *Gnb3* promoter. Each
conserved region was tiled into 5bp windows and the nucleotides within each window were
shuffled. All library sequences and the corresponding cBCs can be found in Supplementary
Table 5.
405
The library of *Gnb3* promoter variants was constructed in four steps. In the first step, we cloned
the *Gnb3* promoter variant library into the core promoter library vector backbone. We ordered
double-stranded DNA fragments from Integrated DNA Technologies (Coralville, Iowa) encoding
the varying part of the (520 bp) *Gnb3* promoter and 113 promoter variants. The wild type *Gnb3*
promoter sequence was included twice, each time fused to a different cBC. The DNA fragments
were manually pooled and cloned together as a library. In the second step, we cloned the
remaining *Gnb3* promoter (300bp) and an mEmerald reporter cassette between the *Gnb3*
promoter variants and the first cBC copy using HiFi assembly. In the third step, we used NEB
HiFi DNA Assembly Master Mix (New England Biolabs E2621) to insert the U6 promoter
between the two copies of the cBCs where it drives expression of the downstream copy of the
cBC. In the fourth step, we introduced high-complexity rBCs between the first cBC and the U6
promoter. We synthesized a DNA oligo containing a 25 bp random sequence (the rBC), a
restriction site, and 30 bp homology to the library vector on each side of the rBC barcode region.
We then used HiFi Assembly to clone the rBC oligos into the *Gnb3* promoter variant library. In
this final library each plasmid contains a *Gnb3* promoter variant driving mEmerald with a unique
cBC-rBC combination in its 3' UTR, which is followed by a polyA signal and the U6 promoter
driving a second copy of the cBC, a capture sequence, and a termination signal. A total of eight
HiFi Assembly reactions were pooled together to increase the library complexity. This library

was transformed and amplified in *E. coli* as described above, and DNA was prepared from
2x10^9 colonies.

Estimating Library Complexity

To estimate the complexity of the core promoter library, we sequenced the DNA library using a
nested PCR-based Illumina library preparation protocol. Briefly, we first used Q5 polymerase
(New England BioLabs M0515) to amplify the region containing the two barcodes with SCARED
17 and SCARED P18 (primer sequences can be found in Supplementary Table 8). The total
reaction volume was 50µl using 50ng of plasmid library with 2.5 µl of 10uM primer each. After
25 cycles of amplification the product was purified with the Monarch PCR&DNA Cleanup kit
(New England BioLabs T1030), and eluted with 20 µl of ddH2O. For the second round of PCR
we used the primers SCARED P19 and SCARED P20 in a 25 µl reaction with 0.25 µl product
from the previous step. After 10 rounds of amplification the product was purified using the
Monarch PCR&DNA Cleanup kit. For the last PCR we added the P5 and P7 Illumina adapters
with SCARED P5, SCARED P7 with 10 cycles of amplification in a 25 µl reaction with 2 µl of
purified product. This final product was sequenced on an Illumina MiSeq, and we obtained a
total of 1,693,933 reads. After filtering out reads without a cBC or rBC of the correct length, we
obtained a total of 1,359,176 reads (80% of the total reads) and 99.5% represented unique
cBC-rBC pairs. For the *Gnb3* library, we performed shallow sequencing, and obtained a total of
1,939,479 reads. After filtering out reads without correct cBC or rBC, we obtained a total of
1,838,415 reads (94.7% of the total reads). Among the 1,838,415 correct reads, 99.5%
represented unique cBC-rBC pairs.

Estimating the Probability of Identical cBC-rBC Pairs in the Same Cell

448   We estimated the probability that more than one copy of a plasmid carrying the same cBC-rBC
449   pair would be transfected into the same cell. We call this probability the collision rate. If the
450   library is transfected into *n* cells, and a specific cBC-rBC pair is present at *m* copies in the
451   library, then the expected number of collisions per experiment is given by:

$$n^{-m} \sum_{k=0}^{n} \binom{n}{k} \sum_{q=0}^{(n-k)} \binom{n-k}{q} \binom{m}{q} q! \left\{ \begin{matrix} m-q \\ n-k-q \end{matrix} \right\}_{n \geq 2} (n-k-q)!(m-q)$$

453   where k denotes the number of cells that received no plasmid, q denotes the number of cells
454   transfected with exactly one plasmid, parentheses denote the binomial coefficient, and brackets
455   denote the partition function. The above expression was simplified by substituting with the
456   bivariate generating function, and the expected number of collisions per experiment is:

$$m(1 - (\frac{n-1}{n})^{m-1})$$

460   The expected number of collisions per cell (λ) is given by,

$$\lambda = \frac{m(1 - (\frac{n - 1}{n})^{m-1})}{n}$$

463 And, assuming collisions are a Poisson process, the probability of at least one collision in a cell
464 is:

$$P(Collision) = 1 - e^{-\lambda}$$

Using this framework, we can estimate the probability of a collision in our experiment. We
assume one million cells (*n*) are transfected using 10 ug of plasmid DNA, and that the effective
number of plasmids that enter the nucleus is 10% of that input amount (1 ug)[43]1 ug of plasmid
DNA is 2.3×10^{11}. Thus, the value of *m* in the nucleus is 2.3×10^{11} divided by the number of
unique members of the library. This allows us to calculate P(Collision) for a library of any given
size. This framework shows that we require a library with 1.6×10^6 unique members to achieve
P(Collision)=0.01. To be 99% sure that a library has at least 1.6×10^6 unique members requires
preparing that library 4.5 times as many independent colonies (7.2×10^6), assuming a Poisson
distributed library. The core promoter library was prepared from 4×10^8 colonies, 55 times more
than required for P(Collision)=0.01, and the *Gnb3* variant library was prepared from 2×10^9
colonies, 277 times more than required for P(Collision)=0.01.

Cell Line Transfections

K562 cells were transfected with the core promoter library using electroporation with the Neon
transfection system (Invitrogen MPK5000). One million cells were transfected with 2 µg of
plasmid DNA (mixed-cell experiment) or 10 µg of plasmid DNA (K562 sub-state experiment),
with 3 pulses of 1450 V for 10 ms.

HEK293 cells were transfected with the core promoter library using the Lipofectamine3000
protocol. 4 µl of p3000 reagent, 4µl of Lipofectamine, and OptiMEM were mixed with 2 µg of
plasmid DNA to a volume of 250 µl. The lipofectamine reagents and plasmid were mixed and
incubated at room temp for 15 minutes and then added dropwise to 1 million cells. We
harvested K562 and HEK293 cells 24 hours after transfections for scMPRA.

Ex vivo culturing and transfection of mouse retinas

Retinas from newborn mice were dissected, cultured, and electroporated as described in White
et al.[44] For each replicate experiment, three retinas were electroporated with 0.5 µg/µL of the
Gnb3 promoter variant library, and co-electroporated with 0.5 µg/µL of a plasmid in which the
Rhodopsin promoter drives the dsRED fluorophore. Electroporated retinas were harvested and
dissociated as in Murphy et al.[42] with modifications as outlined below.

Briefly, three retinas/replicate were washed 3x in cold Hanks' Balanced Salt Solution (HBSS)
(Thermo Fisher) and were then incubated in 400 ul of HBSS containing 0.65 mg papain
(Worthington Biochem) for 10 min at 37˚C. 600ul of Dulbecco's Modified Eagle Medium (DMEM)
(Thermo Fisher) containing 10% fetal calf serum (FCS) (Gibco) was added and the tissue was

gently triturated with P1000 to achieve single cells suspension. 100 units DNase1 (Roche) was
added, the cell suspension and incubated an additional 5 min at 37˚C. Cells were centrifuged at
400g for 4 min then resuspended in 600 ml of sorting buffer (2.5 mM EDTA, 25 mM HEPES, 1%
BSA in HBSS) and passed through a 35um filter and used directly for Fluorescence Activated
Cell Sorting (FACS).

Because the majority of cells in murine retinas are rod photoreceptors, we attempted to enrich
other cell types using FACS. The co-electroporated Rhodopsin-DsRed construct marks rod cells
specifically. Therefore, we used FACS to generate a 1:1 mixture of dsRED$^+$ to dsRED$^-$ cells
from dissociated retinas. This procedure should yield a mix of cells in which rod cells comprise
50% of the total cells. In practice, rod cells still comprised 87% of the cells that were analyzed
by scMPRA.

Bulk MPRA from Cell Lines

For both K562 cells and HEK293 cells, we harvested cells 24 hours after transfection, extracted
total mRNA, and performed reverse transcription using the Superscript IV Reverse
Transcriptase kit (Invitrogen 18090010). A sequencing library was then constructed using a
nested PCR strategy. Briefly, we first used Q5 (New England BioLabs M0515) polymerase to
amplify the region containing the cBC-rBC barcodes with SCARED P17 and SCARED P18. The
total reaction volume was 50µl with 50ng of backbone and 2.5 µl of 10uM primer each. After 25
cycles of amplification the product was purified with the Monarch PCR&DNA Cleanup kit (New
England BioLabs T1030L) and eluted with 20 µl of ddH2O. Sequencing adapters were then
added with 2 rounds of PCR, 10 cycles each. 0.25µl of PCR product was used with SCARED
P19 and P20, then purified using the Monarch PCR&DNA Cleanup kit (New England BioLabs
T1030L). Finally, we used 2 µl of this product with primers SCARED P5 and SCARED P7, and
then sequenced the resulting product on an Illumina Mi-seq instrument. The activity of library
members was computed as described previously[21].
524
525

Single-cell RNA-seq for scMPRA

527   To perform scMPRA we targeted 2000 cells from the HEK/K562 mixed pool per replicate for
528   each mixed cell experiment, 2500 cells per replicate for the K562-only experiment, and 2500
529   cells (after sorting) per replicate for the retina experiment. The cells were prepared according to
530   the 10x Chromium Single Cell 3' Feature Barcode Library Kit (PN-1000079) protocol.
531
532   Our goal was two-fold: to quantify the cBC-rBC pairs from each single cell and to sequence the
533   cellular mRNAs from those same single cells. We captured all polyadenylated RNAs (barcoded
534   reporter RNAs and cellular mRNAs) from single cells following the manufacturer's protocol up to
535   the cDNA amplification step.
536
For the cellular mRNAs (transcriptome), we followed the 10x protocol, using 1/4 of the cDNA
library to generate dual-indexed transcriptomes. To quantify the cBC-rBC pairs, we performed

separate PCRs using primers specifically targeting the reporter gene to improve barcode
recovery efficiencies. Because the 10x protocol only uses 1/4 of the generated cDNA, we
separately amplified the barcodes from another 1/4 of the pellet cleanup. We first used Q5 (New
England BioLabs M0515) polymerase to amplify the region containing the cBC-rBC pairs with
SCARED P17 and SCARED P18 with 10 cycles. The sample was divided equally into eight
PCR reactions, each with 50 µl of total volume to reduce possible jackpotting. The product was
then purified with the Monarch PCR&DNA Cleanup kit (New England BioLabs T1030L), and
eluted with 20 µl of ddH2O. We then added sequencing adapters using an additional two rounds
of PCR. The first adapter PCR was performed with SCARED P21 and SCARED PP2 with a total
of 10 ng of product from the barcode PCR. Again, we pooled eight PCR reactions, each with 50
µl of total volume and 10 PCR cycles. The PCR product was purified using the Monarch
PCR&DNA Cleanup kit (New England BioLabs T1030L). For the last PCR, to add the P5 and P7
Illumina adapters, we used the primers SCARED P45, SCARED PP3 with 10ng of product and
pooled eight PCR reactions, each with 50 µl of total volume and 10 PCR cycles.
553
554   For the U6 promoter library construction, we followed Step 4 of the 10x feature barcoding library
555   preparation protocol (Chromium Next GEM Single Cell 3' Reagent Kits v3.1 (Dual Index)
556   CG000316 Rev C).
557
558   The transcriptome and barcode libraries were combined and sequenced on the Illumina
559   NextSeq 500 with 28x105 cycles. Read1 was limited to 28bp to avoid sequencing the constant
560   poly(A) sequence.

scRNA-seq data processing

The single-cell RNAseq data was processed using Cellranger 6.0.1
(https://github.com/10xGenomics/cellranger) and Scanpy 1.8.1[45]
(https://github.com/theislab/scanpy) following the standard pipeline. Briefly, different sequencing
runs from the same biological replicate were pooled together and processed with CellRanger
6.1.1; the final output expression matrix was then imported into Scanpy for further processing.
We first removed cells with less than 1000 genes, genes that were present in less than three
cells, and cells with high counts of mitochondrial genes. Next, we normalized the UMI counts to
the total cell UMI counts. The normalized expression matrix was used for clustering and
visualization with Scanpy.

scMPRA Data Processing

For each promoter library, paired-end reads generated from barcoded reporter RNAs were
processed with custom scripts that can be found on Github (https://github.com/barakcohenlab
/scMPRA). In each paired-end read, Read1 contains a 10x cell barcode and a UMI, while Read2
contains the cBC and rBC sequences. We define a "quad" as a 10x Cell Barcode, UMI, cBC,
and rBC originating from the same individual paired-end read. To tabulate the cBC-rBCs we first
matched the constant sequences flanking both barcodes, filtering out reads where either
barcode was not the correct length. We performed this filtering using a stand-alone program
(https://github.com/szhao045/scMPRA_parsingtools). Second, we filtered out incorrect 10x Cell

Barcodes based on the CellRanger output barcode list using error-correction with a maximum
hamming distance of one. Third, to mitigate the effect of template-switching during the PCR
steps, we plotted the rank read depth for each unique quad and identified an "elbow point" at a
minimum depth of 1 read for the mixed-cell and the retina experiment, and 10 reads for the
K562 alone experiment. We kept all reads above the minimum depth, and also kept a low-depth
unique quad if it contained a cBC-rBC matching a high-depth pair with a hamming distance of at
most one. Lastly, for the mixed-cell experiment and the K562 cell alone experiment, we
removed any cell with less than 100 scMPRA-associated UMIs, since the scMPRA reads from
those cells were poorly sampled. For the last step, because the retina experiment contains
additional information from the U6 promoter, we did not threshold based on cells. Since U6
promoter data provides information on whether a given cBC in a given cell is sampled well, we
removed all unique barcode pairs containing only 1 UMI for a cBC.
592

Calculating the Single-Cell Activities of Promoters

595 Once the high-confidence quads were identified, we computed A, the activity of a promoter in an
596 individual cell using,

$$A = \frac{\sum_{i=1}^{n} UMI\ count\ for\ cBC_i}{\sum_{i=1}^{n} rBC\ count\ for\ cBC_i}$$

598 where n is the number of unique cBCs that mark a single promoter in the library, and the UMI
599 and rBC counts are summed over all quads with a given 10x cell barcode. We then compute C,
600 the cell-type specific activity of a promoter as,

$$C = \frac{\sum_{j=1}^{m} A_j}{m}$$

602 where m is the number of cells in a given cell type, and all 10x cell barcodes assigned to a given
603 cell type are identified from their matched scRNA-seq profiles. For scMPRA data from the retina
604 we modified the equation for cell-type specific activity as follows,

$$C = \frac{\sum_{j=1}^{P} A_j}{P + U}$$

606 where P is the number of cells of a given cell type in which *Gnb3*-driven cBCs were detected
607 and U is the number of cells of that cell type for which a U6 promoter cBC was detected without
608 detecting any corresponding *Gnb3*-driven cBC. This modification has the effect of adding
609 activities of zero for all cells with U6-driven cBCs that did not express a *Gnb3*-driven cBC.

Cell cycle analysis

612 Cell cycle analysis for the scRNA-seq experiment was done with Scanpy 1.8.1 with cell cycle
613 genes[46]. The expression profile of each cell was projected onto a PCA plot based on the list of
614 cell cycle genes using Scanpy.

Motif analysis

The core promoters were first clustered according to their expression levels in the different cell
sub-state populations by hierarchical clustering. We categorized our data into up/down
regulated clusters at the first branching point, aiming to preserve the large structure. We then
identified core promoter motifs in each promoter according to the parameters in Zabidi et al[47].
using MAST v4.10.0[48] and plotted the proportion of promoters containing each motif in each
promoter class.

Statistical Analyses

All statistical analyses were done using Python 3.9.6, Numpy 1.12.1[49], Scipy 1.6.3 and R 4.0.2.

Ethics

This study was performed in strict accordance with the recommendations in the Guide for the
Care and Use of Laboratory Animals of the National Institutes of Health. All of the animals were
handled according to protocol # A-3381-01 approved by the Institutional Animal Care and Use
Committee of Washington University in St. Louis. Euthanasia of mice was performed according
to the recommendations of the American Veterinary Medical Association Guidelines on
Euthanasia. Appropriate measures are taken to minimize pain and discomfort to the animals
during experimental procedures.

Data and Code Availability

Next-generation sequencing data that support the findings of the study are available in the Gene
Expression Omnibus using accession code GSE188639.

The code that supports the findings of this study is available on Github Repository
(https://github.com/barakcohenlab/scMPRA).

Acknowledgements

We thank the members of the Cohen laboratory for their critical feedback on the manuscript. We
thank Jess Hoistington-Lopez and MariaLynn Crosby for assistance with high-throughput
sequencing. This work is supported by grants to B.A.C from the National Institutes of Health,
R01 GM140711 and R01 GM092910, and by a grant to J.C.C from the National Institutes of
Health R01 EY030075.

Author Contributions

S.Z. and B.A.C. conceived and designed the project. All experiments and analyses were performed by S.Z. with technical contributions from C.K.Y. H. and D.M.G., except for the electroporation and culturing of mouse retinas, which was performed by C.A.M. J.C.C and M.A.W provided critical input into the design of the *Gnb3* promoter variant library. S.Z., C.K.Y.H., and B.A.C. wrote the manuscript with input and feedback from all authors.

Competing Interests

S.Z. and B.A.C. are inventors on a pending patent filed by Washington University in St. Louis which may encompass the methods, reagents, and data disclosed in this manuscript. B.A.C is on the scientific advisory board of Patch Biosciences.

Supplemental Tables

Supplemental Table 1. Mixed Cell Experiment Expression
Supplemental Table 2. Differential expression of the core promoter library between K562 and HEK293 cells
Supplemental Table 3. K562 Cell Cell Cycle Expression
Supplemental Table 4. K562 Cell Cell Sub-state Expression
Supplemental Table 5. Gnb3 Promoter Variant Library
Supplemental Table 6. Gnb3 Library Expression in Retina
Supplemental Table 7. Sequences of Oligonucleotides Used in this Study

Supplementary Figures

Supplementary Figure 1. scMPRA measures cell-type specific CRS activity. (**a**) UMAP of the single-cell transcriptome from the mixed-cell experiment. 105 out of 3417 cells (3%) are labeled by both K562 and HEK293 cell genes. (**b**) UMAP of the mixed-cell

676 experiment with cells marked by other representative markers for K562 and HEK293 cell expression. (**c,d**) Histogram of the number
677 of plasmids (unique cBC-rBC pairs) transfected into K562 cells and HEK293 cells. (**e,f**) Histogram of the mean number of rBC per
678 cBC (CRS) per cell for K562 cells and HEK293 cells. (g,h) Correlation of bulk MPRA versus scMPRA where only the scMPRA data
679 has been UMI normalized (**i,j**) Scatterplot of scMPRA reproducibility for housekeeping and developmental promoters in K562 cells
680 and HEK293 cells.

681
682

683
Supplementary Figure 2 scMPRA measures CRS activity in K562 cell substates. (**a**) Reproducibility for mean expression of
core promoters in K562 cells. (**b**) Correlation of bulk and scMPRA (non-UMI corrected) in K562 cells (**c**) Different dynamics of
expression. For *UBA52*, the promoter is most highly expressed in S phase; whereas for *CSF1*, the promoter is most highly
expressed in G1 phase. For *CXCL10,* the promoter is expressed evenly through cell cycle (Stars indicate significance from Wilcoxon
rank sum test, *: p < 0.05.) (**d**) Cells no longer cluster together based on cell cycle genes after the effects of the cell cycle are
removed.

709
710 **Supplementary Figure 3 (a)** Expression of marker genes by scRNA-seq used to identify cell types in the retina **(b)** Percentage of
711 the total cells recovered represented by each retinal cell type **(c)** Plot showing the relationship between the mean activity of a *Gnb3*
712 promoter variant in a given cell type (x-axis) and the proportion of cells in which that promoter variant is silent (y-axis). Individual
713 cells in which a given *Gnb3* variant is silent are identified as cells with U6-expressed cBC, but no *Gnb3*-expressed cBC. **(d)** The
714 correlation between biological replicates is plotted as a function of the number of cells used in the analysis.

References

732 1. Schaub, M. A., Boyle, A. P., Kundaje, A., Batzoglou, S. & Snyder, M. Linking disease

733   associations with regulatory information in the human genome. *Genome Res.* **22**, 1748–

734   1759 (2012).

735 2. Maurano, M. T. *et al.* Systematic localization of common disease-associated variation in

736   regulatory DNA. *Science* **337**, 1190–1195 (2012).

737 3. Hindorff, L. A. *et al.* Potential etiologic and functional implications of genome-wide

738   association loci for human diseases and traits. *Proc. Natl. Acad. Sci. U. S. A.* **106**, 9362–

739   9367 (2009).

740 4. Yang, J. *et al.* Common SNPs explain a large proportion of the heritability for human height.

741   *Nat. Genet.* **42**, 565–569 (2010).

742 5. Vattikuti, S., Guo, J. & Chow, C. C. Heritability and genetic correlations explained by

743   common SNPs for metabolic syndrome traits. *PLoS Genet.* **8**, e1002637 (2012).

744 6. Shi, H., Kichaev, G. & Pasaniuc, B. Contrasting the Genetic Architecture of 30 Complex

745   Traits from Summary Association Data. *Am. J. Hum. Genet.* **99**, 139–153 (2016).

746 7. Aygün, N. *et al.* Brain-trait-associated variants impact cell-type-specific gene regulation

747   during neurogenesis. *Am. J. Hum. Genet.* **108**, 1647–1668 (2021).

748 8. Nott, A. *et al.* Brain cell type–specific enhancer–promoter interactome maps and disease-

749   risk association. *Science* (2019).

750 9. Spielmann, M. & Mundlos, S. Looking beyond the genes: the role of non-coding variants in

751   human disease. *Hum. Mol. Genet.* **25**, R157–R165 (2016).

752 10. Zhang, F. & Lupski, J. R. Non-coding genetic variants in human disease. *Hum. Mol. Genet.*

753   **24**, R102–10 (2015).

754 11. Ong, C.-T. & Corces, V. G. Enhancer function: new insights into the regulation of tissue-

specific gene expression. *Nat. Rev. Genet.* **12**, 283–293 (2011).

12. Arnold, C. D. *et al.* Genome-wide quantitative enhancer activity maps identified by STARR-

seq. *Science* **339**, 1074–1077 (2013).

13. Kwasnieski, J. C., Mogno, I., Myers, C. A., Corbo, J. C. & Cohen, B. A. Complex effects of

nucleotide variants in a mammalian cis-regulatory element. *Proc. Natl. Acad. Sci. U. S. A.*

109, 19498–19503 (2012).

14. Ireland, W. T. *et al.* Deciphering the regulatory genome of Escherichia coli, one hundred

promoters at a time. *Elife* **9**, (2020).

15. Patwardhan, R. P. *et al.* Massively parallel functional dissection of mammalian enhancers

in vivo. *Nat. Biotechnol.* **30**, 265–270 (2012).

16. Sharon, E. *et al.* Inferring gene regulatory logic from high-throughput measurements of

thousands of systematically designed promoters. *Nat. Biotechnol.* **30**, 521–530 (2012).

17. Kinney, J. B., Murugan, A., Callan, C. G., Jr & Cox, E. C. Using deep sequencing to

characterize the biophysical mechanism of a transcriptional regulatory sequence. *Proc.*

Natl. Acad. Sci. U. S. A. **107**, 9158–9163 (2010).

18. Melnikov, A. *et al.* Systematic dissection and optimization of inducible enhancers in human

cells using a massively parallel reporter assay. *Nat. Biotechnol.* **30**, 271–277 (2012).

19. White, M. A. *et al.* A Simple Grammar Defines Activating and Repressing cis-Regulatory

Elements in Photoreceptors. *Cell Rep.* **17**, 1247–1254 (2016).

20. Kwasnieski, J. C., Fiore, C., Chaudhari, H. G. & Cohen, B. A. High-throughput functional

testing of ENCODE segmentation predictions. *Genome Res.* **24**, 1595–1602 (2014).

21. Chaudhari, H. G. & Cohen, B. A. Local sequence features that influence AP-1 cis-

regulatory activity. *Genome Res.* **28**, 171–181 (2018).

22. Hughes, A. E. O., Myers, C. A. & Corbo, J. C. A massively parallel reporter assay reveals

context-dependent activity of homeodomain binding sites in vivo. *Genome Res.* **28**, 1520–

1531 (2018).

23. Tewhey, R. *et al.* Direct Identification of Hundreds of Expression-Modulating Variants using

a Multiplexed Reporter Assay. *Cell* **165**, 1519–1529 (2016).

24. Hong, C. K. Y. & Cohen, B. A. Genomic environments scale the activities of diverse core

promoters. *bioRxiv* 2021.03.08.434469 (2021) doi:10.1101/2021.03.08.434469.

25. Haberle, V. *et al.* Transcriptional cofactors display specificity for distinct types of core

promoters. *Nature* **570**, 122–126 (2019).

26. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for

RNA-seq data with DESeq2. *Genome Biol.* **15**, 550 (2014).

27. Shaffer, S. M. *et al.* Rare cell variability and drug-induced reprogramming as a mode of

cancer drug resistance. *Nature* **546**, 431–435 (2017).

28. Moudgil, A. *et al.* Self-Reporting Transposons Enable Simultaneous Readout of Gene

Expression and Transcription Factor Binding in Single Cells. *Cell* **182**, 992–1008.e21

(2020).

29. Litzenburger, U. M. *et al.* Single-cell epigenomic variability reveals functional cancer

heterogeneity. *Genome Biol.* **18**, 15 (2017).

30. Min, M. & Spencer, S. L. Spontaneously slow-cycling subpopulations of human cells

originate from activation of stress-response pathways. *PLoS Biol.* **17**, e3000178 (2019).

31. Bonnet, D. & Dick, J. E. Human acute myeloid leukemia is organized as a hierarchy that

originates from a primitive hematopoietic cell. *Nat. Med.* **3**, 730–737 (1997).

32. Ishikawa, F. *et al.* Chemotherapy-resistant human AML stem cells home to and engraft

within the bone-marrow endosteal region. *Nat. Biotechnol.* **25**, 1315–1321 (2007).

33. Friedman, R. Z. *et al.* Information content differentiates enhancers from silencers in mouse

photoreceptors. *Elife* **10**, (2021).

34. Dixit, A. *et al.* Perturb-Seq: Dissecting Molecular Circuits with Scalable Single-Cell RNA

Profiling of Pooled Genetic Screens. *Cell* **167**, 1853–1866.e17 (2016).

35. Reese, B. E. Development of the retina and optic pathway. *Vision Res.* **51**, 613–632

(2011).

36. Cao, J. *et al.* Comprehensive single-cell transcriptional profiling of a multicellular organism.

Science **357**, 661–667 (2017).

37. Chan, Y. K. *et al.* Engineering adeno-associated viral vectors to evade innate immune and

inflammatory responses. *Sci. Transl. Med.* **13**, (2021).

38. Byrne, L. C. *et al.* In vivo-directed evolution of adeno-associated virus in the primate retina.

JCI Insight **5**, (2020).

39. Wang, D., Tai, P. W. L. & Gao, G. Adeno-associated virus vector as a platform for gene

therapy delivery. *Nat. Rev. Drug Discov.* **18**, 358–378 (2019).

40. Bryant, D. H. *et al.* Deep diversification of an AAV capsid protein by machine learning. *Nat.*

Biotechnol. **39**, 691–696 (2021).

41. Shen, S. Q. *et al.* Massively parallel cis-regulatory analysis in the mammalian central

nervous system. *Genome Res.* **26**, 238–255 (2016).

42. Murphy, D. P., Hughes, A. E., Lawrence, K. A., Myers, C. A. & Corbo, J. C. Cis-regulatory

basis of sister cell type divergence in the vertebrate retina. *Elife* **8**, (2019).

43. Cohen, R. N., van der Aa, M. A. E. M., Macaraeg, N., Lee, A. P. & Szoka, F. C., Jr.

Quantification of plasmid DNA copies in the nucleus after lipoplex and polyplex transfection.

J. Control. Release **135**, 166–174 (2009).

44. White, M. A., Myers, C. A., Corbo, J. C. & Cohen, B. A. Massively parallel in vivo enhancer

assay reveals that highly local features determine the cis-regulatory function of ChIP-seq

peaks. *Proc. Natl. Acad. Sci. U. S. A.* **110**, 11952–11957 (2013).

45. Wolf, F. A., Angerer, P. & Theis, F. J. SCANPY: large-scale single-cell gene expression

data analysis. *Genome Biol.* **19**, 15 (2018).

46. Tirosh, I. *et al.* Dissecting the multicellular ecosystem of metastatic melanoma by single-cell

RNA-seq. *Science* **352**, 189–196 (2016).

47. Zabidi, M. A. *et al.* Enhancer–core-promoter specificity separates developmental and

833  housekeeping gene regulation. *Nature* **518**, 556–559 (2014).

834 48. Bailey, T. L. & Gribskov, M. Combining evidence using p-values: application to sequence

835  homology searches. *Bioinformatics* **14**, 48–54 (1998).

836 49. Harris, C. R. *et al.* Array programming with NumPy. *Nature* **585**, 357–362 (2020).

837