# Genomic landscape of drug response reveals novel mediators of anthelmintic resistance

**Authors:** Stephen R. Doyle[1,†,*], Roz Laing[2,†,*], David Bartley[3], Alison Morrison[3], Nancy Holroyd[1], Kirsty Maitland[2], Alistair Antonopoulos[2], Umer Chaudhry[4§], Ilona Flis[2], Sue Howell[5], Jennifer McIntyre[2], John S. Gilleard[6], Andy Tait[2], Barbara Mable[2], Ray Kaplan[5+], Neil Sargison[4], Collette Britton[2], Matthew Berriman[1], Eileen Devaney[2,‡], James A. Cotton[1,‡]

† These authors contributed equally to this work

‡ These authors contributed equally to this work

§ Current address: School of Veterinary Medicine, University of Surrey; Surrey, United Kingdom

+ Current address: Department of Pathobiology, School of Veterinary Medicine, St. George's University; St. George's, Grenada

*Corresponding authors: email: stephen.doyle@sanger.ac.uk; email: Rosalind.Laing@glasgow.ac.uk

**Affiliations:**
1. Wellcome Sanger Institute; Hinxton, United Kingdom
2. Institute of Biodiversity Animal Health and Comparative Medicine, College of Medical, Veterinary and Life Sciences, University of Glasgow; Glasgow, United Kingdom
3. Moredun Research Institute; Penicuik, United Kingdom
4. Royal (Dick) School of Veterinary Studies, University of Edinburgh; Edinburgh, United Kingdom
5. Department of Infectious Diseases, College of Veterinary Medicine, University of Georgia; Athens, United States
6. Department of Comparative Biology and Experimental Medicine, Host-Parasite Interactions Program, Faculty of Veterinary Medicine, University of Calgary; Calgary, Canada

**Abstract:**

Understanding the genetic basis of anthelmintic drug resistance in parasitic nematodes

is key to improving the efficacy and sustainability of parasite control. Here, we use a

genetic cross in a natural host-parasite system to simultaneously map resistance loci

for the three major classes of anthelmintics. This approach identifies novel alleles for

resistance to benzimidazoles and levamisole and implicates the transcription factor,

*cky-1*, in ivermectin resistance. This gene is within a locus under selection in ivermectin

resistant populations worldwide; functional validation using knockout experiments

supports a role for *cky-1* overexpression in ivermectin resistance. Our work

demonstrates the feasibility of high-resolution forward genetics in a parasitic

nematode, and identifies variants for the development of molecular diagnostics to

combat drug resistance in the field.

**One-Sentence Summary:**

Genetic mapping of known and novel anthelmintic resistance-associated alleles in a

multi-drug resistant parasitic nematode

**Main Text:** Over a billion people and countless livestock and companion animals

require at least annual treatment with anthelmintic drugs to control parasitic worm

(helminth) infections. The rapid and widespread evolution of resistance to these drugs

is a significant health concern in livestock (*1*) and places an economic burden on food

production. Resistance is present on every continent where anthelmintics are used; in

many places, individual drug classes are now ineffective, and some farms have

resistance to every major class of drug (*2*), threatening the economic viability of livestock

farming. In Europe, gastrointestinal helminths of livestock are responsible for annual

production losses of €686 million, of which €38 million is associated with anthelmintic

resistance (*3*). Drug resistance is also now a major concern in the treatment of

helminths infecting dogs (*4, 5*), with multiple drug resistance to all major anthelmintic

classes in the dog hookworm now common in the USA (*6*). The same classes of drugs

to which veterinary parasites have rapidly evolved resistance are also used to control

related human-infective helminths, which are targeted at scale by some of the largest

preventative chemotherapy programmes in the world. Although less established in

human-infective helminths, the emergence of widespread anthelmintic resistance –

echoing the current global emergency around antimicrobial resistance – will have

serious socio-economic and welfare impacts on people infected with parasitic worms

and derail hard-won progress towards the proposed eradication and elimination of

helminths over the next decade (*7, 8*).

Despite extensive efforts, the causal mutations and mechanisms of resistance in parasitic helminths remain largely unresolved. Many candidate "resistance genes" have been proposed for most drug classes; these candidates are primarily homologues of genes that confer resistance in the free-living model nematode *Caenorhabditis elegans,* and are subsequently assayed for differences in genetic variation and/or gene expression in parasite isolates that vary in their response to treatment (*9–11*). A successful example of this approach is the identification of variants of β-tubulin that inhibit tubulin-depolymerisation by benzimidazole-class anthelmintics (*12, 13*). These variants, particularly at amino acid positions 167, 198 and 200 of β-tubulin isotype 1 (*14–16*), have subsequently been shown to be associated with resistance in many parasitic species for which benzimidazoles have been extensively used, and a number of these parasite-specific mutations have been functionally validated in *C. elegans* (*17, 18*). However, these three variants are unlikely to explain all phenotypic variation associated with resistance (*19, 20*), and it is unknown to what degree other variants contribute to benzimidazole resistance in parasitic species. For other drug classes, few candidate genes have been functionally validated or shown to be important in natural parasite populations. For example, concurrent mutation of three glutamate-gated chloride channels (*glc-1, avr-14, avr-15*) enables resistance to high levels of ivermectin by *C. elegans (21)*, yet no strong evidence of selection on these channels in any parasitic species has been demonstrated to date. On the one hand, the many genes proposed may reflect that resistance is a complex, quantitative trait where similar resistance phenotypes can be derived from variation in multiple loci. Alternatively, resistance may be species and/or population-specific, and evolve independently under

subtly different selection pressures (*22*). However, some candidates are likely to have

been falsely associated with resistance, as most studies present relatively weak

genetic evidence from the analysis of single or few candidate loci in small numbers of

helminth populations that often differ in both drug susceptibility and geographic origin.

Many helminth species are exceptionally genetically diverse (*23–26*), and consequently,

candidate gene approaches have limited power to disentangle causal variation from

linked but unrelated background genetic variation, a situation that is exacerbated by

the experimental intractability and inadequate genomic resources available for many

parasitic helminths (*9*).

Here we describe a genome-wide forward genetics approach using the parasitic

nematode *Haemonchus contortus* as a model to identify genetic variation associated

with resistance to three of the most important broad-spectrum anthelmintic drugs

globally: ivermectin, levamisole, and benzimidazole. *H. contortus* is an economically

important gastrointestinal parasite of livestock and one of only a few genetically

tractable parasites used for drug discovery (*27, 28*), vaccine development (*29, 30*), and

anthelmintic resistance research (*22*). Our approach has exploited a genetic cross

between the susceptible MHco3(ISE) and multi-drug resistant MHco18(UGA) strains of

*H. contortus* (**fig. S1 A**), allowing us to investigate resistance in a natural host-parasite

system while controlling for confounding genetic diversity that differentiates parasite

strains (see **Supplementary materials** regarding the establishment and validation of

the cross). Using an eXtreme Quantitative Trait Locus (X-QTL) (*31, 32*) analysis

framework, whereby pools of F3-generation progeny from F2 adults treated *in vivo*

were sampled pre- and post-treatment for each drug (**fig. S1 B**; n = 3 parasite

populations per drug class maintained in independent sheep; **fig. S2**) and analysed by

whole-genome sequencing (**table S1**), we aimed to identify drug-specific quantitative

trait loci (QTLs) associated with resistance throughout the genome. These QTLs and

specific variants were independently validated using genome-wide variation from

populations of *H. contortus* obtained from ten US farms of known resistance

phenotype (see **Supplementary materials** for a description of the US farms and

quantitative phenotyping**; table S2, fig. S3, fig. S4**), and from more than 350 individual

parasites sampled throughout the world where *H. contortus* is endemic (*25*, *33*).

***A genetic cross between genetically-distinct susceptible and multi-drug resistant***

***strains reveals drug-specific QTL after selection***

A key feature and thus advantage of using a genetic cross to map anthelmintic

resistance loci is that the high degree of within-strain diversity and genome-wide

genetic divergence is controlled by admixture in the F1 generation of the cross. The

susceptible and resistant parental *H. contortus* strains of the cross are highly

genetically differentiated throughout their genomes (**Fig. 1A;** mean $F_{ST}$ = 0.089 ± 0.066

SD; n = 16,794,366 single nucleotide variant sites), typical of two parasite strains

sampled from different continents (*25, 34*). In subsequent generations, both

susceptible and resistant alleles segregate at moderate frequencies in the absence of

selection, and genetic recombination breaks down the linked genetic variation that

defines and differentiates the parental strains. This was evident by a significantly lower

genome-wide genetic differentiation in the F3-generation control population (genome-

wide mean $F_{ST}$ = 0.012 ± 0.004) and absence of discrete peaks of high genetic

differentiation (**Fig. 1B: Control**). In contrast, after each drug treatment, discrete QTLs

that differ between each drug class were revealed: after benzimidazole treatment, we

identified a major QTL on chromosome 1 (**Fig. 1B: Benzimidazole**); after levamisole,

two QTLs on chromosome 4 and 5 (**Fig. 1B: Levamisole**); and after ivermectin, a major

QTL on chromosome 5 and minor QTLs on chromosomes 2 and 5 (**Fig. 1B:**

**Ivermectin**).

**Fig. 1. A genetic cross followed by drug selection reveals discrete QTLs associated with**

**each anthelmintic drug class.**

(**A**) Genome-wide comparison of susceptible MHco3(ISE) and multidrug-resistant MHco18(UGA) parental strains revealed broad-scale genetic differentiation on all chromosomes. (**B**) Comparison of genome-wide differentiation between F3 generation pooled infective-stage larvae (L$_3$, n = 200) sampled pre- and post-treatment revealed distinct genomic regions or QTLs associated with benzimidazole, levamisole, and ivermectin drug treatment. An untreated control where sampling was time-matched to the treated groups is shown for comparison. In all plots, each point represents the mean genetic differentiation ($F_{ST}$) from three biological replicates in five kb sliding windows, and the dashed line represents the genome-wide mean $F_{ST}$ + 3 SD for each comparison (See **fig. S2** for genome-wide replicate data). Individual chromosomes are indicated by alternating dark and light blue shading.

### *Variation at β-tubulin isotype 1 and a novel β-tubulin isotype 2 variant is associated with high levels of benzimidazole resistance*

The β-tubulin isotype 1 (HCON_00005260) gene and, in particular, nonsynonymous changes at coding positions 167, 198, and 200 have been widely associated with benzimidazole resistance in *H. contortus* (*13*, *15*, *16*, *35*) and other nematodes frequently exposed to benzimidazole treatment (*17*, *36*). After benzimidazole selection, a single broad QTL was found on chromosome 1 (**Fig. 2A;** see **Supplementary materials** for further discussion of the genetic structure of the QTL) containing the β-tubulin isotype 1 locus. Within this gene, we identified a significant increase in the frequency of a Phe200Tyr variant (a phenylalanine [reference susceptible variant] to tyrosine [resistant variant] substitution at position 200) from pre- to post-treatment and relative to untreated controls (**Fig. 2B;** *P* = 1.7e-26, genome-wide Cochran–Mantel–

Haenszel (CMH) test between replicates). We also identified a small increase in

frequency of the Phe167Tyr variant (mean $freq_{\text{pre-treatment}}$ = 0.14 to $freq_{\text{post-treatment}}$ = 0.20),

however, no variation was found at the Glu198 position. Considering the previous

association between these variants and benzimidazole resistance, we conclude that

the Phe200Tyr variant is the primary driver of phenotypic resistance in the X-QTL

population.

*Haemonchus contortus* has multiple β-tubulin genes (*37*), and deletion of the β-tubulin

isotype 2 gene (HCON_00043670) on chromosome 2 has been associated with

increased levels of resistance beyond that of mutations in the isotype 1 gene alone

(*14*). Here, we found no evidence of deletions in isotype 2. However, a minor but not

significant increase in genetic differentiation between pre- and post-treatment

populations was found at this locus, and a Glu198Val variant at a homologous site to a

known resistance variant in isotype 1 was present at a low frequency in the genetic

cross ($freq_{\text{pre-treatment}}$ = 0.260 to $freq_{\text{post-treatment}}$ = 0.323; not significant genome-wide

CMH). However, on the US farms, the Glu198Val variant did vary in frequency between

farms and was significantly correlated (r = 0.978, $P$ = 1.02e-6; Pearson's correlation)

with $EC_{50}$ values for benzimidazole resistance (**Fig. 2C**). The variance observed in $EC_{50}$

among resistant farm populations was not caused by variation in the frequency of the

Phe200Tyr mutation of the isotype 1 gene, as this variant was already at high

frequency in all populations, except for the farm that was susceptible to

benzimidazoles (Farm 1; **fig. S5**). These data suggest that once the isotype 1

Phe200Tyr variant has reached near fixation in the population, the Glu198Val variant of

isotype 2 mediates higher levels of benzimidazole resistance than conferred by the

Phe200Tyr variant alone. As such, this novel allele present in β-tubulin isotype 2 should

be considered, in addition to the well-characterised isotype 1 variants, as a genetic

marker for benzimidazole resistance.

In addition to the association with benzimidazole resistance, it has been suggested

that the β-tubulin isotype 1 Phe200Tyr variant in *H. contortus* (*38–40*) and also at an

equivalent variant site in a β-tubulin gene in the human-infective filarial nematode

*Onchocerca volvulus* (*41*) is associated with ivermectin resistance. Here we found no

evidence of selection on either the Phe167Tyr or Phe200Tyr variants (or any variant

found in the region) in X-QTL analyses of ivermectin treatment (**fig. S6A**), nor any

correlation with ivermectin $EC_{50}$ on the US farms (**fig. S6B**). These data reaffirm that

mutations in β-tubulin isotype 1 are specific to benzimidazole resistance.

**Fig. 2. Characterisation of QTL associated with benzimidazole resistance.**

**(A)** Chromosome-wide genetic differentiation between pre- and post-benzimidazole treatment

on chromosome 1. Each point represents the mean $F_{ST}$ in a five kb window; points are coloured

based on the concordance of individual replicates indicated by none (blue), 1 of 3 (yellow), 2 of

3 (orange), or all 3 (red) above the genome-wide threshold. The genome-wide threshold is

defined as the mean + 3 SD of the chromosome-wide $F_{ST}$ indicated by the horizontal dashed

line, whereas the vertical dashed line highlights the position of the β-tubulin isotype 1

(HCON_00005260) gene. **(B)** Allele frequency change at Phe167Tyr and Phe200Tyr variant

positions of β-tubulin isotype 1 pre- and post-treatment, including untreated time-matched

control. Coloured lines show biological replicates. *P*-values are calculated using pairwise t-

tests of allele frequency by sampling time point (i.e., pre- and post-treatment). **(C)** Correlation

between benzimidazole $EC_{50}$ concentration (µM) observed at particular farms and Glu198Val

variant frequency of β-tubulin isotype 2 (HCON_00043670) on US farms. Pearson's correlation

(r) and associated *P*-value together with the trendline and standard error of the linear

regression are shown.

***Levamisole selection implicates acetylcholine receptors, including a novel acr-8***

***variant, with resistance***

The anthelmintic activity of levamisole is due to its antagonistic effect on nematode

nicotinic acetylcholine receptors (*42*), and resistance in *C. elegans* is typically

associated with variation in subunits of these receptors or other accessory proteins

that contribute to acetylcholine-mediated signalling (*43*). Here we identified two major

QTLs on chromosomes 4 and 5 that contain a tandem duplication of the acetylcholine

receptor subunit β-type *lev-1* (HCON_00107690 & HCON_00107700) and acetylcholine

receptor subunit *acr-8* (HCON_00151270), respectively (**Fig. 3A**).

The *H. contortus acr-8* gene (**Fig. 3B**) has long been implicated in levamisole

resistance; a truncated isoform of *acr-8* containing the two first exons and a part of

intron 2 (previously called *Hco-acr-8b*) (*44*), and subsequently, a 63 bp indel between

exons 2 and 3 have been associated with resistance based on their presence in several

resistant isolates (*45*). However, the functional consequence of these variants in

mediating levamisole resistance *in vivo* is not yet clear. Here, we identified two larger

deletion variants spanning 31,527,022 to 31,527,119 (97 bp) or 31,527,121 (99 bp) on

chromosome 5 that increased in frequency from 73.47% in the pre-treatment

population to 86.58% after levamisole treatment (**Fig. 3C;** paired t-test across

replicates, $P = 0.1$). However, the *acr-8* indel was present in the levamisole susceptible

parental MHco3(ISE) strain (59.05%) and was present at only a slightly higher

frequency in the resistant MHco18(UGA) strain (63.55%). Thus, these data argue that

the *acr-8* indel is a poor marker of levamisole resistance.

We did, however, identify a nonsynonymous variant (Ser168Thr) in *acr-8* that was

strongly correlated with resistance across multiple datasets. In the X-QTL analyses,

Ser168Thr increased to a high frequency after drug selection in the F2 generation (**Fig.**

**3D**; position 31,521,884; genome-wide CMH: $P = 1.6e\text{-}15$; allele frequency change

pre- vs post-treatment: $P = 1.0e\text{-}4$; in time-matched no-treatment control: $P = 0.4$). It

was also found at a high frequency in the USA field population with the highest

levamisole drug resistance phenotype (Farm 7; $freq_{Ser168Thr} = 0.64$). This association was

supported in global diversity data of *H. contortus* (*25*), where we found the Ser168Thr

variant fixed in parasites from the Kokstad (KOK; South Africa) population ($freq_{Ser168Thr} =$

1.0; n = 4), the only population with confirmed levamisole resistance in that study,

whereas the variant was absent in all other populations analysed. The identification of

Ser168Thr prompted us to look beyond *H. contortus*; a reanalysis of levamisole

resistance in resequencing data from the closely related clade V parasitic nematode

*Teladorsagia circumcincta (46)* revealed a homologous non-synonymous variant at high

frequency in resistant parasites (Ser140Thr in Cont419:G75849C ; $freq_{Ser140Thr} = 0.972$),

which was absent in the susceptible population to which it was compared. Although a

serine to threonine substitution is a relatively conservative change, we found the serine

residue to be highly conserved among clade V nematodes (**fig. S7**), particularly among

the parasite species, whereas in the free-living *Caenorhabditis spp.*, threonine is

encoded at this position.

In *C. elegans, acr-8* is genetically and functionally distinct relative to *acr-8* of parasitic

nematodes and is not a component of the native levamisole receptor (*47*); the *C.*

*elegans* functional homolog *lev-8*, which can be transgenically substituted by *H.*

*contortus acr-8* to produce a functional receptor (*48*), does encode a serine at this

homologous position. The *H. contortus* ACR-8 Ser168Thr variant lies immediately

downstream of the cys-loop domain within the ligand-binding pocket and is

immediately adjacent to a highly conserved tryptophan residue essential for ligand

binding (*49*, *50*) (**Fig. 3E**). Importantly, key residues downstream of the conserved

tryptophan have previously been shown to influence levamisole sensitivity of closely

related receptor subunits (*51*). Thus, we hypothesise that the Ser168Thr variant

facilitates a change in the molecular interactions within the binding pocket of ACR-8,

resulting in a decreased sensitivity to levamisole.

The identification of *lev-1* genes within the chromosome 4 QTL is compelling, with

three intronic variants of *lev-1* (top variant position 14,995,062 in HCON_00107700; *P* =

1.7e-20; CMH test) among of the top ten most differentiated SNPs on this

chromosome. However, it remains unclear what effect the overall observed variation in

the *lev-1* genes has on levamisole resistance. Although multiple non-synonymous

variants were also identified (seven and three variants for HCON_00107690 and

HCON_00107700, respectively), none were predicted to cause high-effect changes in

the protein sequence and exhibited only relatively minor shifts in allele frequency upon

levamisole treatment. In *C. elegans*, several dominant resistant variants of *lev-1* have

been described (not found in the data described here); however, *lev-1* can be lost

without affecting the function of the receptor (*43*). Examination of variation in *lev-1*

expression in addition to genetic variation may be required to elucidate the role of *H.*

*contortus lev-1* subunits in levamisole resistance. Close to the *lev-1* genes and toward

the centre of the QTL, four of the top ten variants in chromosome 4 were found in

HCON_00107560 (top non-synonymous variant: Arg934His at position 14,781,344; *P* =

1.0e-21; CMH test), an ortholog of *C. elegans kdin-1*. Highly conserved with

mammalian orthologs (*52*), *kdin-1* has been shown to co-localise with acetylcholine

receptors at rat neuromuscular junctions during development (*53*) where, via a PDZ

domain, it participates in the coordination of signalling components including ion

channels and neurotransmitters. The precise role of HCON_00107560 or *kdin-1* in *H.*

*contortus* or *C. elegans*, respectively, remains unknown; however, its putative

association with levamisole response here warrants further investigation.

Signals of selection on two components of the pentameric acetylcholine receptor

prompted us to look for selection on the remaining subunits. Although the expression

of *unc-63* (HCON_00024380) and *unc-29.3* (HCON_00003520) mRNAs were

significantly reduced in the larvae of resistant MHco18(UGA) strain (*54*), we found no

evidence of selection on the region of the genome containing these genes.

**Fig. 3. Characterisation of QTL associated with levamisole resistance.**

(**A**) QTL between pre-treatment and levamisole-treated parasites on chromosome 4 (top) and

chromosome 5 (bottom). Each point represents the mean $F_{ST}$ in a five kb window; points are

coloured based on the concordance of individual replicates indicated by none (blue), 1 of 3

(yellow), 2 of 3 (orange), or all 3 (red) above the genome-wide threshold (horizontal dashed line;

mean + 3 SD of the chromosome-wide $F_{ST}$). (**B**) Gene model for *acr-8* (top; HCON_00151270)

and a cuticle collagen (bottom; HCON_00151260), highlighting the position of the overlapping

*acr-8*/levamisole-associated indel and the Ser168Thr variant of *acr-8*. (**C**) Visualisation of

sequencing reads supporting the *acr-8* intronic indel. Mapped reads are coloured to reflect the

degree to which they have been clipped to allow correct mapping in the presence of the

deletion, i.e. reads that have not been clipped are blue, whereas reads that are moderate to

highly clipped are coloured red to yellow, respectively. (**D**) Comparison of Ser168Thr variant

frequency between pre- and post-levamisole treatment (red) and time-matched untreated

controls (green). (**E**) Structure of the pentameric cys-loop acetylcholine receptor of *Torpedo*

*marmorata* (Protein Data Base ID: 4AQ9), one of the few species from which the receptor's

structure has been resolved (*55*). The Trp[Ser/Thr]Tyr motif is highly conserved among the

clade V nematodes (**fig. S7**) and the distantly related alpha subunit of *T. marmorata*; Thr174,

the homologous position of the *H. contortus* Hc_Ser168Thr variant of *acr-8,* lies within the

acetylcholine binding pocket at the interface of the alpha and gamma subunits and adjacent to

Trp173 (*H. contortus* Hc_Trp167), a residue essential for ligand binding.

***A resolved ivermectin QTL implicates* cky-1 *as a novel mediator of resistance***

Ivermectin is a critically important broad-spectrum drug used to control several

human- and veterinary-infective helminths worldwide and is also widely used as an

acaricide targeting ticks and mites. We recently identified a ~5 Mb QTL associated with

ivermectin resistance from 37 to 42 Mb on chromosome 5 from the analysis of a

backcross experiment (*34, 56*), and subsequently, we identified evidence of selection in

the same chromosomal region in ivermectin-resistant field populations from Africa and

Australia (*25*). Although the introgression region from the backcross was broad (*57*), the

genetic architecture of the QTL was consistent with a single dominant variant driving

resistance, and we were able to demonstrate that most candidate genes previously

proposed to be associated with resistance were not under direct ivermectin selection.

However, we were unable to confidently identify any novel candidate driving mutation

among the ~360 genes lying within the region (*34*).

Here, we confirm the QTL within the previously implicated chromosome 5 region at

~37.5 Mb (*34*, *56*) but with significantly increased resolution (**Fig. 4A**). We have

narrowed the genetic association to approximately 300 kb wide (region: ~37,250,000-

37,550,000), based on the region of highest differentiation between independently

replicated pre- and post-treatment X-QTL samples (**Fig. 4B**). This region was also

highly differentiated between pre-treatment larvae and adult male worms that survived

ivermectin treatment *in vivo* **(fig. S8 A,B),** and between larvae that survived treatment

with an $EC_{75}$ dose of ivermectin and those sensitive to an $EC_{50}$ dose *in vitro* **(fig. S8**

**C,D)**. Together, these results confirm that this locus is under direct selection and

mediates resistance in both the parasitic stages *in vivo* and in the free-living stages *in*

*vitro* (see **Supplementary materials** for further discussion). Finally, this was the only

region in the genome where increased levels of ivermectin resistance (i.e., $EC_{50}$) was

associated with a loss of genetic diversity in moderately or highly resistant field

populations relative to susceptible populations (**Fig 4C**), consistent with a selective

sweep in response to ivermectin-mediated selection.

The main chromosome 5 QTL contained 25 genes and included an expansion of

protein kinases (8/21 genes present in the genome with the InterPro identifier

IPR015897), some of which had the highest statistical association with resistance; for

example, HCON_00155240 (intronic position 37,336,132, $P$ = 3.3e-13; position

37,235,944, $P$ = 1.2e-12) and HCON_00155270 (intronic position 37,343,439, $P$ = 1.0e-

10). These protein kinases are, however, novel leads with no previous association to

drug resistance, and a lack of functional orthologs and observed gene expansion made

it difficult to further infer and test a role for these genes in ivermectin resistance.

Towards the middle of the QTL, we identified *cky-1* (HCON_00155390; positions

37,487,982 - 37,497,398) as a new mediator of resistance, based on several lines of

evidence. In the X-QTL data, *cky-1* contained eight moderately to highly differentiated

significant non-synonymous variants (top variant: position 37,497,061 [Ser583Pro], $P$ =

9.6e-09; CMH test). In a complementary study, we showed *cky-1* was the only gene in

the region significantly upregulated in both males and females of the resistant

MHco18(UGA) isolate relative to MHco3(ISE) and was one of the most upregulated

genes genome-wide (*58*). In this study, RT-qPCR of *cky-1* from the parental isolates of

the cross and two unrelated ivermectin-resistant *H. contortus* strains revealed

significant overexpression in ivermectin-resistant relative to sensitive strains (**Fig. 4E**),

an observation that was replicated between sensitive and ivermectin-resistant strains

of the related parasite, *T. circumcincta* (**Fig. 4E**). To explore *cky-1* further, we assayed

*C. elegans* developmental and pumping behavioural phenotypes, both known to be

perturbed by ivermectin exposure (*59*), to test the role of differential expression of *cky-*

*1* on the resistant phenotype in the presence of ivermectin. While complete knockout

of *cky-1* was non-viable, both a balanced deletion (VC2274) (**Fig. 4E**) and RNAi

knockdown (**Fig. 4F**) of *cky-1* increased the sensitivity of *C. elegans* to ivermectin

relative to the ivermectin-susceptible N2 strain. The level of *cky-1* expression is,

therefore, associated with the ivermectin resistance phenotype in nematodes.

Two additional, less prominent QTLs on chromosome 5 at ~46 Mb and on

chromosome 2 at ~3 Mb were also identified after ivermectin treatment (**Fig. 4A;** see

**Supplementary materials** for a description of the two QTLs). The second

chromosome 5 QTL was identified as a candidate region associated with resistance in

the backcross (*34*); however, we did not have the statistical power to differentiate it

from the main QTL in that experiment. Here, the QTL appeared to segregate

independently of the prominent 37.5 Mb peak, providing more robust evidence of a

second resistance-conferring variant on chromosome 5. Although the main

chromosome 5 QTL at 37.5 Mb was present in all selection experiments with

ivermectin, the secondary QTLs were variable between replicates and experiments. To

further refine the association, we exposed the F5 generation of the cross to a half

standard dose of ivermectin, followed by a double dose thereafter. The rationale was

first to identify low-effect variants (responding to the half dose treatment), then select a

subset of variants that conferred resistance at high doses (see **Supplementary**

**materials** for additional background). In these experiments, we consistently detected

the main chromosome 5 QTL but not the less prominent chromosome 2 QTL.

Additionally, we detected the presence of at least three new minor QTLs (**fig. S10**,

supplementary text). Of practical significance, the identification of novel replicate-

specific variants in addition to the main chromosome 5 QTL highlights the

consequence of under-dosing in selecting novel variants, and emphasises the

importance of correct dosing in the field.

**Fig. 4. Characterisation of QTL associated with ivermectin resistance.**

(**A**) QTL between pre- and post-ivermectin treatment on chromosome 5. Each point represents the mean $F_{ST}$ in a five kb window; points are coloured based on the concordance of individual replicates indicated by none (blue), 1 of 3 (yellow), 2 of 3 (orange), or all 3 (red) above the genome-wide threshold (horizontal dashed line; mean + 3 SD of the chromosome-wide $F_{ST}$). A

magnified aspect of the main chromosome 5 QTL, highlighting (**B**) genetic differentiation ($F_{ST}$) in

the X-QTL cross, and (**C**) nucleotide diversity (Pi) on US farms, where each farm is coloured by

the degree of ivermectin resistance ($EC_{50}$) measured by larval development assays. In A, B and

C, the position of *cky-1* is indicated by the vertical dashed line. (**D**) RT-qPCR analysis of *cky-1*

expression in both *H. contortus* and *T. circumcincta* strains that differ in their ivermectin

resistance phenotype. Data represents $\log_2$-transformed expression normalised to actin or

GAPDH control genes for *H. contortus* and *T. circumcincta,* respectively. Downregulation of

*cky-1* expression in *C. elegans* by either (**E**) a balanced deletion or (**F**) RNAi-knockdown

increases ivermectin sensitivity relative to the control N2 strain, based on developmental

assays measuring the percentage of progeny surviving to adulthood relative to DMSO controls.

(In **E** and **F,** each point represents an independent treatment condition, which is normalised to

a DMSO control without ivermectin. A Kruskal-Wallis test was used to determine whether

treatment condition differed from untreated control, where ns = not significant, * $p < 0.05$, ** $p$

$< 0.01$, and **** $p < 0.0001$.

**Discussion**

Anthelmintics are currently the most important tool for controlling parasitic worm

infections in humans and animals worldwide, and this is likely to remain true for the

foreseeable future. However, this paradigm of control is threatened by the emergence

and spread of anthelmintic-resistant parasites. Despite the large health and economic

impacts resulting from increasing levels of anthelmintic resistance, multiple

complicating factors have hindered the ability to determine the genetic loci responsible

for resistance. Here we demonstrate an efficient approach to map multiple drug

resistance-conferring loci for three of the most important anthelmintic drugs in the

globally distributed and genetically tractable parasitic nematode, *H. contortus*. We

have identified novel variants and loci likely involved in resistance to each of these drug

classes; these include the β-tubulin isotype 2 Glu198Val variant correlated with

benzimidazole resistance in field populations, the *acr-8* Ser168Thr variant associated

with levamisole resistance in both the cross and field populations of *H. contortus*, and

*cky-1* as a novel candidate gene that mediates ivermectin response. Our approach was

validated by identifying QTLs and variants previously associated with drug resistance,

for example, the β-tubulin isotype 1 Phe200Tyr variant associated with benzimidazole

resistance and the *acr-8* indel variant associated with levamisole resistance. However,

for the latter, we provide evidence against the indel being a reliable marker of

resistance. Finally, we note an absence of many previously proposed ivermectin-

associated candidate genes in the QTL described, highlighting both the limitation of

candidate gene approaches and the power of genome-wide forward-genetic strategies

to robustly identify regions of the genome containing known and novel mediators of

resistance (*9*).

We have refined a previously identified QTL for ivermectin resistance on chromosome

5 (*34*) to ~300 kb, and together with functional genetic evidence from expression and

knockout experiments, we have explicitly tested the role of our proposed candidate in

the main ivermectin QTL on chromosome 5, the NPAS4 ortholog *cky-1*. This gene

encodes an activity-dependent basic Helix-Loop-Helix (bHLH)-PAS family transcription

factor shown in mammals to regulate the excitation/inhibition balance upon neuronal

activation to limit excitotoxicity (*60*) and during the development of inhibitory synapses

to control the expression of activity-dependent genes (*61*). It is yet to be determined if

this is a conserved molecular function in nematodes; however, it is tempting to

speculate that the hyperexcitability as a result of induced activation of ion channels by

ivermectin at the neuromuscular junction is, at least in part, controlled by a "retuning"

of the excitation/inhibition balance to limit toxicity. The role of *cky-1* in ivermectin

resistance is supported by: (i) genetic differentiation between susceptible and resistant

strains around this locus relative to genome-wide variation that is replicated in

geographically and genetically diverse strains here and elsewhere (*25*, *34*, *62*), (ii) the

presence of non-synonymous variants that are highly differentiated before and after

treatment, (iii) increased gene expression of *cky-1* in resistant strains relative to

susceptible strains (supported by genome-wide RNA-seq (*58*)) and (iv) knockdown of

the *C. elegans* ortholog leading to hypersensitivity to ivermectin. We acknowledge that

overexpression of *cky-1* in *C. elegans* does not recapitulate the high levels of

ivermectin resistance seen in *H. contortus* or, for example, by concurrent mutation of

glutamate-gated chloride channels in *C. elegans* (*21*); while this may argue against *cky-*

*1* as a universal mediator of resistance, it likely reflects the challenge of using a

heterologous expression system in which there is an assumption that the biology (and,

therefore, response to treatment) is concordant between the free-living and parasitic

species, and/or may reflect the multigenic nature of ivermectin resistance in different

species (*63–65*). Given the lack of an obvious causal non-synonymous variant, we

hypothesise that a non-coding variant that influences the expression of *cky-1* is under

selection in resistant strains of *H. contortus*; however, such variants are difficult to

validate without genotype and transcriptional phenotype data from a large number of

individual worms.

It is broadly accepted that the mode of action of ivermectin is on ligand-gated ion

channels, and ivermectin resistance has been associated with variants in glutamate-

gated channels (*66*). Concurrent mutation of a number of these channels (*glc-1, avr-14*

and *avr-15*) confers high-level resistance in *C. elegans* (*21*) and selection on at least

one of these channels (*glc-1*) in wild strains (*67*) has been demonstrated. We find no

evidence to suggest that genetic variation in these channels confers ivermectin

resistance in *H. contortus*. Transcriptional changes in these channels in resistant,

relative to drug-susceptible, parasite strains have been demonstrated previously; for

example, the glutamate-gated chloride channel subunits (*glc-3, glc-5*), as well as p-

glycoprotein ABC transporters (*pgp-1, pgp-2, pgp-9*) (*54*) in the MHco18(UGA) strain.

Similarly, a *pgp-9* copy number variant was associated with ivermectin resistance in a

genetic cross and bulk segregant experiment in the related nematode *T. circumcincta*

*(46)*, while transgenic overexpression of the equine parasitic nematode *Parascaris*

*univalens pgp-9* modulated ivermectin sensitivity in *C. elegans* (*68*). However, none of

these genes were identified in regions of differentiation after treatment in this study,

suggesting these genes are not the direct target of selection. However, we cannot

exclude that variation in expression of these genes may be a downstream response to

selection on a transcriptional regulator such as *cky-1*.

The use of genetic crosses, in which the genetics of the parasites can be controlled, is

the ideal way to generate populations of individuals in which the relationship between

genotype and phenotype can be assayed. Our approach here relies on selecting

populations of parasites using drug treatment, however, advances are still required to

improve phenotyping of resistance in individual parasites. The ability to do so would

improve our understanding of the molecular basis of drug resistance phenotypes and

enable more sophisticated genetic approaches to unravel the role of the minor

signatures of selection we observe in this experiment. Recent advances in single larvae

whole-genome sequencing (*69*) and low-input RNA sequencing (*70*), even at single-cell

resolution (*71*), now provide the tools to allow a more precise understanding of

molecular and cellular phenotypes for drug response and may help to fully understand

the role of *cky-1*. The identification of *cky-1* as a putative candidate offers new

plausible hypotheses relevant to a resistant phenotype, whereby *cky-1* may act: (i)

during development to establish a neuronal architecture that is more tolerant to

hyperexcitability such as that caused by ivermectin, and/or (ii) in response to

ivermectin exposure by initiating transcription of downstream genes to modulate the

excessive excitation/inhibition imbalance, thereby mitigating the lethal effect. These

hypotheses will require further validation, aided in the first instance by identifying the

downstream targets of *cky-1*. However, it is clear that the molecular mechanisms by

which parasites develop ivermectin resistance are more complex than previously

appreciated. Broader, systems biology approaches are likely needed to understand the

relationship between direct evidence of selection in the genome and the downstream

transcriptional responses that enable parasite survival when challenged with

ivermectin. By defining the genomic landscape of anthelmintic resistance even in a

single resistant strain, our results refocus effort away from candidate genes with limited

support and redefine our understanding of the evolution of anthelmintic resistance in

helminths of veterinary and medical importance.

**References and Notes:**

1.   R. M. Kaplan, Drug resistance in nematodes of veterinary importance: a status report.
*Trends Parasitol.* **20**, 477–481 (2004).

2.   R. M. Kaplan, A. N. Vidyashankar, An inconvenient truth: global worming and anthelmintic
resistance. *Vet. Parasitol.* **186**, 70–78 (2012).

3.   J. Charlier, L. Rinaldi, V. Musella, H. W. Ploeger, C. Chartier, H. R. Vineer, B. Hinney, G.
von Samson-Himmelstjerna, B. Băcescu, M. Mickiewicz, T. L. Mateus, M. Martinez-
Valladares, S. Quealy, H. Azaizeh, B. Sekovska, H. Akkari, S. Petkevicius, L. Hektoen, J.
Höglund, E. R. Morgan, D. J. Bartley, E. Claerebout, Initial assessment of the economic
burden of major parasitic helminth infections to the ruminant livestock industry in Europe.
*Prev. Vet. Med.* **182**, 105103 (2020).

4.   P. D. Jimenez Castro, S. B. Howell, J. J. Schaefer, R. W. Avramenko, J. S. Gilleard, R. M.
Kaplan, Multiple drug resistance in the canine hookworm *Ancylostoma caninum*: an
emerging threat? *Parasit. Vectors*. **12**, 576 (2019).

5.   C. Bourguinat, A. C. Y. Lee, R. Lizundia, B. L. Blagburn, J. L. Liotta, M. S. Kraus, K. Keller,
C. Epe, L. Letourneau, C. L. Kleinman, T. Paterson, E. C. Gomez, J. A. Montoya-Alonso, H.
Smith, A. Bhan, A. S. Peregrine, J. Carmichael, J. Drake, R. Schenker, R. Kaminsky, D. D.
Bowman, T. G. Geary, R. K. Prichard, Macrocyclic lactone resistance in *Dirofilaria immitis*:
Failure of heartworm preventives and investigation of genetic markers for resistance. *Vet.*
*Parasitol.* **210**, 167–178 (2015).

6.   P. D. Jimenez Castro, A. Venkatesan, E. Redman, R. Chen, A. Malatesta, H. Huff, D. A.
Zuluaga Salazar, R. Avramenko, J. S. Gilleard, R. M. Kaplan, Multiple drug resistance in
hookworms infecting greyhound dogs in the USA. *Int. J. Parasitol. Drugs Drug Resist.* **17**,
107–117 (2021).

7.   World Health Organization, "Ending the neglect to attain the Sustainable Development
Goals: A road map for neglected tropical diseases 2021–2030" (World Health
Organization, Geneva, 2020), (available at
https://www.who.int/neglected_diseases/resources/who-ucn-ntd-2020.01/en/).

8.   A. Montresor, D. Mupfasoni, A. Mikhailov, P. Mwinzi, A. Lucianez, M. Jamsheed, E.
Gasimov, S. Warusavithana, A. Yajima, Z. Bisoffi, D. Buonfrate, P. Steinmann, J. Utzinger,
B. Levecke, J. Vlaminck, P. Cools, J. Vercruysse, G. Cringoli, L. Rinaldi, B. Blouin, T. W.
Gyorkos, The global progress of soil-transmitted helminthiases control in 2020 and World
Health Organization targets for 2030. *PLoS Negl. Trop. Dis.* **14**, e0008505 (2020).

9.   S. R. Doyle, J. A. Cotton, Genome-wide approaches to investigate anthelmintic resistance.
*Trends Parasitol.* **35**, 289–301 (2019).

10. A. C. Kotze, P. W. Hunt, P. Skuce, G. von Samson-Himmelstjerna, R. J. Martin, H. Sager,
J. Krücken, J. Hodgkinson, A. Lespine, A. R. Jex, J. S. Gilleard, R. N. Beech, A. J.
Wolstenholme, J. Demeler, A. P. Robertson, C. L. Charvet, C. Neveu, R. Kaminsky, L.
Rufener, M. Alberich, C. Menez, R. K. Prichard, Recent advances in candidate-gene and
whole-genome approaches to the discovery of anthelmintic resistance markers and the
description of drug/receptor interactions. *Int. J. Parasitol. Drugs Drug Resist.* **4**, 164–184
(2014).

11. S. R. Hahnel, C. M. Dilks, I. Heisler, E. C. Andersen, D. Kulke, *Caenorhabditis elegans* in
anthelmintic research - Old model, new perspectives. *Int. J. Parasitol. Drugs Drug Resist.*
**14**, 237–248 (2020).

12. M. Driscoll, E. Dean, E. Reilly, E. Bergholz, M. Chalfie, Genetic and molecular analysis of a
*Caenorhabditis elegans* beta-tubulin that conveys benzimidazole sensitivity. *J. Cell Biol.*
**109**, 2993–3003 (1989).

13. M. H. Roos, J. H. Boersema, F. H. Borgsteede, J. Cornelissen, M. Taylor, E. J. Ruitenberg,
Molecular analysis of selection for benzimidazole resistance in the sheep parasite
*Haemonchus contortus*. *Mol. Biochem. Parasitol.* **43**, 77–88 (1990).

14. M. S. Kwa, F. N. Kooyman, J. H. Boersema, M. H. Roos, Effect of selection for
benzimidazole resistance in *Haemonchus contortus* on beta-tubulin isotype 1 and isotype
2 genes. *Biochem. Biophys. Res. Commun.* **191**, 413–419 (1993).

15. A. Silvestre, J. Cabaret, Mutation in position 167 of isotype 1 beta-tubulin gene of
Trichostrongylid nematodes: role in benzimidazole resistance? *Mol. Biochem. Parasitol.*
**120**, 297–300 (2002).

16. M. Ghisi, R. Kaminsky, P. Mäser, Phenotyping and genotyping of *Haemonchus contortus*
isolates reveals a new putative candidate mutation for benzimidazole resistance in
nematodes. *Vet. Parasitol.* **144**, 313–320 (2007).

17. C. M. Dilks, S. R. Hahnel, Q. Sheng, L. Long, P. T. McGrath, E. C. Andersen, Quantitative
benzimidazole resistance and fitness effects of parasitic nematode beta-tubulin alleles. *Int.*
*J. Parasitol. Drugs Drug Resist.* **14**, 28–36 (2020).

18. M. S. G. Kwa, J. G. Veenstra, M. Van Dijk, M. H. Roos, β-Tubulin genes from the parasitic
nematode *Haemonchus contortus* modulate drug resistance in *Caenorhabditis elegans*.
*Journal of Molecular Biology*. **246**, 500–510 (1995).

19. S. R. Hahnel, S. Zdraljevic, B. C. Rodriguez, Y. Zhao, P. T. McGrath, E. C. Andersen,
Extreme allelic heterogeneity at a *Caenorhabditis elegans* beta-tubulin locus explains
natural resistance to benzimidazoles. *PLoS Pathog.* **14**, e1007226 (2018).

20. M. Zamanian, D. E. Cook, S. Zdraljevic, S. C. Brady, D. Lee, J. Lee, E. C. Andersen,

Discovery of genomic intervals that underlie nematode responses to benzimidazoles. *PLoS Negl. Trop. Dis.* **12**, e0006368 (2018).

21. J. A. Dent, M. M. Smith, D. K. Vassilatis, L. Avery, The genetics of ivermectin resistance in *Caenorhabditis elegans*. *Proc. Natl. Acad. Sci. U. S. A.* **97**, 2674–2679 (2000).

22. J. S. Gilleard, *Haemonchus contortus* as a paradigm and model to study anthelmintic drug resistance. *Parasitology*. **140**, 1506–1522 (2013).

23. S. R. Doyle, C. Bourguinat, H. C. Nana-Djeunga, J. A. Kengne-Ouafo, S. D. S. Pion, J. Bopda, J. Kamgno, S. Wanji, H. Che, A. C. Kuesel, M. Walker, M.-G. Basáñez, D. A. Boakye, M. Y. Osei-Atweneboana, M. Boussinesq, R. K. Prichard, W. N. Grant, Genome-wide analysis of ivermectin response by *Onchocerca volvulus* reveals that genetic drift and soft selective sweeps contribute to loss of drug sensitivity. *PLoS Negl. Trop. Dis.* **11**, e0005816 (2017).

24. J. S. Gilleard, Understanding anthelmintic resistance: the need for genomics and genetics. *Int. J. Parasitol.* **36**, 1227–1239 (2006).

25. G. Sallé, S. R. Doyle, J. Cortet, J. Cabaret, M. Berriman, N. Holroyd, J. A. Cotton, The global diversity of *Haemonchus contortus* is shaped by human intervention and climate. *Nat. Commun.* **10**, 4811 (2019).

26. D. J. Berger, T. Crellen, P. H. L. Lamberton, F. Allan, A. Tracey, J. D. Noonan, N. B. Kabatereine, E. M. Tukahebwa, M. Adriko, N. Holroyd, J. P. Webster, M. Berriman, J. A. Cotton, Whole-genome sequencing of *Schistosoma mansoni* reveals extensive diversity with limited selection despite mass drug administration. *Nature Communications*. **12** (2021), doi:10.1038/s41467-021-24958-0.

27. R. Kaminsky, P. Ducray, M. Jung, R. Clover, L. Rufener, J. Bouvier, S. S. Weber, A. Wenger, S. Wieland-Berghausen, T. Goebel, N. Gauvry, F. Pautrat, T. Skripsky, O. Froelich, C. Komoin-Oka, B. Westlund, A. Sluder, P. Mäser, A new class of anthelmintics effective against drug-resistant nematodes. *Nature*. **452**, 176–180 (2008).

28. S. Preston, A. Jabbar, C. Nowell, A. Joachim, B. Ruttkowski, J. Baell, T. Cardno, P. K. Korhonen, D. Piedrafita, B. R. E. Ansell, A. R. Jex, A. Hofmann, R. B. Gasser, Low cost whole-organism screening of compounds for anthelmintic activity. *Int. J. Parasitol.* **45**, 333–343 (2015).

29. C. C. Bassetto, A. F. T. Amarante, Vaccination of sheep and cattle against haemonchosis. *J. Helminthol.* **89**, 517–525 (2015).

30. D. P. Knox, D. L. Redmond, G. F. Newlands, P. J. Skuce, D. Pettit, W. D. Smith, The nature and prospects for gut membrane proteins as vaccine candidates for *Haemonchus contortus* and other ruminant trichostrongyloids. *Int. J. Parasitol.* **33**, 1129–1137 (2003).

31. A. Burga, E. Ben-David, T. L. Vergara, J. Boocock, L. Kruglyak, Fast genetic mapping of
complex traits in *C. elegans* using millions of individuals in bulk. *Nature Communications*.
**10**, 2680 (2019).

32. F. D. Chevalier, C. L. L. Valentim, P. T. LoVerde, T. J. C. Anderson, Efficient linkage
mapping using exome capture and extreme QTL in schistosome parasites. *BMC
Genomics*. **15**, 617 (2014).

33. S. R. Doyle, A. Tracey, R. Laing, N. Holroyd, D. Bartley, W. Bazant, H. Beasley, R. Beech,
C. Britton, K. Brooks, U. Chaudhry, K. Maitland, A. Martinelli, J. D. Noonan, M. Paulini, M.
A. Quail, E. Redman, F. H. Rodgers, G. Sallé, M. Z. Shabbir, G. Sankaranarayanan, J. Wit,
K. L. Howe, N. Sargison, E. Devaney, M. Berriman, J. S. Gilleard, J. A. Cotton, Genomic
and transcriptomic variation defines the chromosome-scale assembly of *Haemonchus
contortus*, a model gastrointestinal worm. *Commun Biol*. **3**, 656 (2020).

34. S. R. Doyle, C. J. R. Illingworth, R. Laing, D. J. Bartley, E. Redman, A. Martinelli, N.
Holroyd, A. A. Morrison, A. Rezansoff, A. Tracey, E. Devaney, M. Berriman, N. Sargison, J.
A. Cotton, J. S. Gilleard, Population genomic and evolutionary modelling analyses reveal a
single major QTL for ivermectin drug resistance in the pathogenic nematode, *Haemonchus
contortus*. *BMC Genomics*. **20**, 218 (2019).

35. M. S. G. Kwa, J. G. Veenstra, M. H. Roos, Benzimidazole resistance in *Haemonchus
contortus* is correlated with a conserved mutation at amino acid 200 in β-tubulin isotype 1.
*Molecular and Biochemical Parasitology*. **63**, 299–303 (1994).

36. R. W. Avramenko, E. M. Redman, L. Melville, Y. Bartley, J. Wit, C. Queiroz, D. J. Bartley, J.
S. Gilleard, Deep amplicon sequencing as a powerful new tool to screen for sequence
polymorphisms associated with anthelmintic resistance in parasitic nematode populations.
*Int. J. Parasitol.* **49**, 13–26 (2019).

37. G. I. Saunders, J. D. Wasmuth, R. Beech, R. Laing, M. Hunt, H. Naghra, J. A. Cotton, M.
Berriman, C. Britton, J. S. Gilleard, Characterization and comparative analysis of the
complete *Haemonchus contortus* β-tubulin gene family and implications for benzimidazole
resistance in strongylid nematodes. *Int. J. Parasitol.* **43**, 465–475 (2013).

38. M. de Lourdes Mottier, R. K. Prichard, Genetic analysis of a relationship between
macrocyclic lactone and benzimidazole anthelmintic selection on *Haemonchus contortus*.
*Pharmacogenet. Genomics*. **18**, 129–140 (2008).

39. J. K. L. Eng, W. J. Blackhall, M. Y. Osei-Atweneboana, C. Bourguinat, D. Galazzo, R. N.
Beech, T. R. Unnasch, K. Awadzi, G. W. Lubega, R. K. Prichard, Ivermectin selection on
beta-tubulin: evidence in *Onchocerca volvulus* and *Haemonchus contortus*. *Mol. Biochem.
Parasitol.* **150**, 229–235 (2006).

40. J. M. L. dos Santos, J. M. L. dos Santos, J. F. Vasconcelos, G. A. Frota, W. L. C. Ribeiro,

W. P. P. André, L. da Silva Vieira, M. Teixeira, C. M. L. Bevilaqua, J. P. Monteiro,
*Haemonchus contortus* β-tubulin isotype 1 gene F200Y and F167Y SNPs are both
selected by ivermectin and oxfendazole treatments with differing impacts on anthelmintic
resistance. *Veterinary Parasitology*. **248**, 90–95 (2017).

41.  M. Y. Osei-Atweneboana, D. A. Boakye, K. Awadzi, J. O. Gyapong, R. K. Prichard,
Genotypic analysis of β-tubulin in *Onchocerca volvulus* from communities and individuals
showing poor parasitological response to ivermectin treatment. *Int. J. Parasitol. Drugs*
*Drug Resist.* **2**, 20–28 (2012).

42.  R. J. Martin, A. P. Robertson, S. K. Buxton, R. N. Beech, C. L. Charvet, C. Neveu,
Levamisole receptors: a second awakening. *Trends Parasitol.* **28**, 289–296 (2012).

43.  J. T. Fleming, M. D. Squire, T. M. Barnes, C. Tornoe, K. Matsuda, J. Ahn, A. Fire, J. E.
Sulston, E. A. Barnard, D. B. Sattelle, J. A. Lewis, *Caenorhabditis elegans* levamisole
resistance genes lev-1, unc-29, and unc-38 encode functional nicotinic acetylcholine
receptor subunits. *J. Neurosci.* **17**, 5843–5857 (1997).

44.  A. Fauvin, C. Charvet, M. Issouf, J. Cortet, J. Cabaret, C. Neveu, cDNA-AFLP analysis in
levamisole-resistant *Haemonchus contortus* reveals alternative splicing in a nicotinic
acetylcholine receptor subunit. *Molecular and Biochemical Parasitology*. **170**, 105–107
(2010).

45.  V. Barrère, R. N. Beech, C. L. Charvet, R. K. Prichard, Novel assay for the detection and
monitoring of levamisole resistance in *Haemonchus contortus*. *Int. J. Parasitol.* **44**, 235–
241 (2014).

46.  Y.-J. Choi, S. A. Bisset, S. R. Doyle, K. Hallsworth-Pepin, J. Martin, W. N. Grant, M.
Mitreva, Genomic introgression mapping of field-derived multiple-anthelmintic resistance
in *Teladorsagia circumcincta*. *PLoS Genet.* **13**, e1006857 (2017).

47.  G. Hernando, I. Bergé, D. Rayes, C. Bouzat, Contribution of subunits to *Caenorhabditis*
*elegans* levamisole-sensitive nicotinic receptor function. *Mol. Pharmacol.* **82**, 550–560
(2012).

48.  A. Blanchard, F. Guégnard, C. L. Charvet, A. Crisford, E. Courtot, C. Sauvé, A. Harmache,
T. Duguet, V. O'Connor, P. Castagnone-Sereno, B. Reaves, A. J. Wolstenholme, R. N.
Beech, L. Holden-Dye, C. Neveu, Deciphering the molecular determinants of cholinergic
anthelmintic sensitivity in nematodes: When novel functional validation approaches
highlight major differences between the model *Caenorhabditis elegans* and parasitic
species. *PLoS Pathog.* **14**, e1006996 (2018).

49.  D. K. Williams, C. Stokes, N. A. Horenstein, R. L. Papke, Differential regulation of receptor
activation and agonist selectivity by highly conserved tryptophans in the nicotinic
acetylcholine receptor binding site. *J. Pharmacol. Exp. Ther.* **330**, 40–53 (2009).

50.   T. Lynagh, S. A. Pless, Principles of agonist recognition in Cys-loop receptors. *Front.*
*Physiol.* **5**, 160 (2014).

51.   D. Rayes, M. J. De Rosa, M. Bartos, C. Bouzat, Molecular basis of the differential
sensitivity of nematode and mammalian muscle to the anthelmintic agent levamisole. *J.*
*Biol. Chem.* **279**, 36372–36381 (2004).

52.   H. Kong, J. Boulter, J. L. Weber, C. Lai, M. V. Chao, An evolutionarily conserved
transmembrane protein that is a novel downstream target of neurotrophin and ephrin
receptors. *J. Neurosci.* **21**, 176–185 (2001).

53.   S. Luo, Y. Chen, K.-O. Lai, J. C. Arévalo, S. C. Froehner, M. E. Adams, M. V. Chao, N. Y.
Ip, α-Syntrophin regulates ARMS localization at the neuromuscular junction and enhances
EphA4 signaling in an ARMS-dependent manner. *Journal of Cell Biology*. **169** (2005), pp.
813–824.

54.   S. M. Williamson, B. Storey, S. Howell, K. M. Harper, R. M. Kaplan, A. J. Wolstenholme,
Candidate anthelmintic resistance-associated gene expression and sequence
polymorphisms in a triple-resistant field isolate of *Haemonchus contortus*. *Mol. Biochem.*
*Parasitol.* **180**, 99–105 (2011).

55.   N. Unwin, Y. Fujiyoshi, Gating movement of acetylcholine receptor caught by plunge-
freezing. *J. Mol. Biol.* **422**, 617–634 (2012).

56.   E. Redman, N. Sargison, F. Whitelaw, F. Jackson, A. Morrison, D. J. Bartley, J. S. Gilleard,
Introgression of ivermectin resistance genes into a susceptible *Haemonchus contortus*
strain by multiple backcrossing. *PLoS Pathog.* **8**, e1002534 (2012).

57.   S. R. Doyle, R. Laing, D. J. Bartley, C. Britton, U. Chaudhry, J. S. Gilleard, N. Holroyd, B.
K. Mable, K. Maitland, A. A. Morrison, A. Tait, A. Tracey, M. Berriman, E. Devaney, J. A.
Cotton, N. D. Sargison, A genome resequencing-based genetic map reveals the
recombination landscape of an outbred parasitic nematode in the presence of polyploidy
and polyandry. *Genome Biol. Evol.* **10**, 396–409 (2018).

58.   R. Laing, S. R. Doyle, J. McIntyre, K. Maitland, A. Morrison, D. J. Bartley, R. Kaplan, U.
Chaudhry, N. Sargison, A. Tait, J. A. Cotton, C. Britton, E. Devaney, Transcriptomic
analyses implicate neuronal plasticity and chloride homeostasis in ivermectin resistance
and recovery in a parasitic nematode. *bioRxiv* (2021), p. 2021.11.12.468372.

59.   J. A. Dent, M. W. Davis, L. Avery, avr-15 encodes a chloride channel subunit that mediates
inhibitory glutamatergic neurotransmission and ivermectin sensitivity in *Caenorhabditis*
*elegans*. *EMBO J.* **16**, 5867–5879 (1997).

60.   I. Spiegel, A. R. Mardinly, H. W. Gabel, J. E. Bazinet, C. H. Couch, C. P. Tzeng, D. A.
Harmin, M. E. Greenberg, Npas4 regulates excitatory-inhibitory balance within neural

circuits through cell-type-specific gene programs. *Cell*. **157**, 1216–1229 (2014).

61. Y. Lin, B. L. Bloodgood, J. L. Hauser, A. D. Lapan, A. C. Koon, T.-K. Kim, L. S. Hu, A. N.
Malik, M. E. Greenberg, Activity-dependent regulation of inhibitory synapse development
by Npas4. *Nature*. **455**, 1198–1204 (2008).

62. A. M. Rezansoff, R. Laing, J. S. Gilleard, Evidence from two independent backcross
experiments supports genetic linkage of microsatellite Hcms8a20, but not other candidate
loci, to a major ivermectin resistance locus in *Haemonchus contortus*. *Int. J. Parasitol.* **46**,
653–661 (2016).

63. K. S. Evans, J. Wit, L. Stevens, S. R. Hahnel, B. Rodriguez, G. Park, M. Zamanian, S. C.
Brady, E. Chao, K. Introcaso, R. E. Tanny, E. C. Andersen, Two novel loci underlie natural
differences in *Caenorhabditis elegans* abamectin responses. *PLoS Pathog.* **17**, e1009297
(2021).

64. J. Wit, C. M. Dilks, E. C. Andersen, Complementary approaches with free-living and
parasitic nematodes to understanding anthelmintic resistance. *Trends Parasitol.* **37**, 240–
250 (2021).

65. A. Streit, Opinion: What do rescue experiments with heterologous proteins tell us and what
not? *Parasitol. Res.* (2021), doi:10.1007/s00436-021-07247-z.

66. R. Laing, V. Gillan, E. Devaney, Ivermectin – Old Drug, New Tricks? *Trends in Parasitology*.
**33** (2017), pp. 463–472.

67. R. Ghosh, E. C. Andersen, J. A. Shapiro, J. P. Gerke, L. Kruglyak, Natural variation in a
chloride channel subunit confers avermectin resistance in *C. elegans*. *Science*. **335**, 574–
578 (2012).

68. A. P. Gerhard, J. Krücken, C. Neveu, C. L. Charvet, A. Harmache, G. von Samson-
Himmelstjerna, Pharyngeal pumping and tissue-specific transgenic P-Glycoprotein
expression influence macrocyclic lactone susceptibility in *Caenorhabditis elegans*.
*Pharmaceuticals*. **14** (2021), doi:10.3390/ph14020153.

69. S. R. Doyle, G. Sankaranarayanan, F. Allan, D. Berger, P. D. Jimenez Castro, J. B. Collins,
T. Crellen, M. A. Duque-Correa, P. Ellis, T. G. Jaleta, R. Laing, K. Maitland, C. McCarthy, T.
Moundai, B. Softley, E. Thiele, P. T. Ouakou, J. V. Tushabe, J. P. Webster, A. J. Weiss, J.
Lok, E. Devaney, R. M. Kaplan, J. A. Cotton, M. Berriman, N. Holroyd, Evaluation of DNA
extraction methods on individual helminth egg and larval stages for whole-genome
sequencing. *Front. Genet.* **10**, 826 (2019).

70. S. Sun, C. Roedelsperger, R. J. Sommer, Single worm transcriptomics identifies a
developmental core network of oscillating genes with deep conservation across
nematodes. *Genome Res.* (2021), doi:10.1101/gr.275303.121.

71.  C. L. Diaz Soria, J. Lee, T. Chong, A. Coghlan, A. Tracey, M. D. Young, T. Andrews, C.
Hall, B. L. Ng, K. Rawlinson, S. R. Doyle, S. Leonard, Z. Lu, H. M. Bennett, G. Rinaldi, P. A.
Newmark, M. Berriman, Single-cell atlas of the first intra-mammalian developmental stage
of the human parasite *Schistosoma mansoni*. *Nat. Commun.* **11**, 6411 (2020).

72.  E. Redman, E. Packard, V. Grillo, J. Smith, F. Jackson, J. S. Gilleard, Microsatellite
analysis reveals marked genetic differentiation between *Haemonchus contortus* laboratory
isolates and provides a rapid system of genetic fingerprinting. *Int. J. Parasitol.* **38**, 111–122
(2008).

73.  J. H. Gill, J. M. Redwin, J. A. Van Wyk, E. Lacey, Avermectin inhibition of larval
development in *Haemonchus contortus* — Effects of ivermectin resistance. *International
Journal for Parasitology*. **25**, 463–470 (1995).

74.  S. B. Howell, J. M. Burke, J. E. Miller, T. H. Terrill, E. Valencia, M. J. Williams, L. H.
Williamson, A. M. Zajac, R. M. Kaplan, Prevalence of anthelmintic resistance on sheep and
goat farms in the southeastern United States. *J. Am. Vet. Med. Assoc.* **233**, 1913–1919
(2008).

75.  M. M. George, L. Lopez-Soberal, B. E. Storey, S. B. Howell, R. M. Kaplan, Motility in the
L3 stage is a poor phenotype for detecting and measuring resistance to
avermectin/milbemycin drugs in gastrointestinal nematodes of livestock. *Int. J. Parasitol.
Drugs Drug Resist.* **8**, 22–30 (2018).

76.  I. Kozarewa, Z. Ning, M. A. Quail, M. J. Sanders, M. Berriman, D. J. Turner, Amplification-
free Illumina sequencing-library preparation facilitates improved mapping and assembly of
(G C)-biased genomes. *Nat. Methods*. **6**, 291–295 (2009).

77.  P. Ewels, M. Magnusson, S. Lundin, M. Käller, MultiQC: summarize analysis results for
multiple tools and samples in a single report. *Bioinformatics*. **32**, 3047–3048 (2016).

78.  A. M. Bolger, M. Lohse, B. Usadel, Trimmomatic: a flexible trimmer for Illumina sequence
data. *Bioinformatics*. **30**, 2114–2120 (2014).

79.  H. Li, Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM.
*arXiv [q-bio.GN]* (2013), (available at http://arxiv.org/abs/1303.3997).

80.  A. McKenna, M. Hanna, E. Banks, A. Sivachenko, K. Cibulskis, A. Kernytsky, K. Garimella,
D. Altshuler, S. Gabriel, M. Daly, M. A. DePristo, The Genome Analysis Toolkit: a
MapReduce framework for analyzing next-generation DNA sequencing data. *Genome Res.*
**20**, 1297–1303 (2010).

81.  P. Cingolani, A. Platts, L. L. Wang, M. Coon, T. Nguyen, L. Wang, S. J. Land, X. Lu, D. M.
Ruden, A program for annotating and predicting the effects of single nucleotide
polymorphisms, SnpEff: SNPs in the genome of *Drosophila melanogaster* strain w1118;

826  iso-2; iso-3. *Fly*. **6**, 80–92 (2012).

827 82. L. Ferretti, S. E. Ramos-Onsins, M. Pérez-Enciso, Population genomics from pool
828   sequencing. *Mol. Ecol.* **22**, 5561–5576 (2013).

829 83. R. Kofler, R. V. Pandey, C. Schlötterer, PoPoolation2: identifying differentiation between
830   populations using sequencing of pooled DNA samples (Pool-Seq). *Bioinformatics*. **27**,
831   3435–3436 (2011).

832 84. P. Danecek, A. Auton, G. Abecasis, C. A. Albers, E. Banks, M. A. DePristo, R. E.
833   Handsaker, G. Lunter, G. T. Marth, S. T. Sherry, G. McVean, R. Durbin, 1000 Genomes
834   Project Analysis Group, The variant call format and VCFtools. *Bioinformatics*. **27**, 2156–
835   2158 (2011).

836 85. K. Katoh, D. M. Standley, MAFFT multiple sequence alignment software version 7:
837   improvements in performance and usability. *Mol. Biol. Evol.* **30**, 772–780 (2013).

838 86. J. Yang, R. Yan, A. Roy, D. Xu, J. Poisson, Y. Zhang, The I-TASSER Suite: protein
839   structure and function prediction. *Nat. Methods*. **12**, 7–8 (2015).

840 87. R. H. Baker, S. Buschbaum, J. B. Matthews, I. J. McKendrick, T. Schnieder, C. Strube, A.
841   J. Nisbet, GTP-cyclohydrolase and development in *Teladorsagia circumcincta* and
842   *Dictyocaulus viviparus* (Nematoda: Strongylida). *Exp. Parasitol.* **128**, 309–317 (2011).

843 88. R. T. Bell, B. X. H. Fu, A. Z. Fire, Cas9 Variants Expand the Target Repertoire in
844   *Caenorhabditis elegans*. *Genetics*. **202**, 381–388 (2016).

845 89. C. C. Mello, J. M. Kramer, D. Stinchcomb, V. Ambros, Efficient gene transfer in *C.elegans*:
846   extrachromosomal maintenance and integration of transforming sequences. *The EMBO
847   Journal*. **10**, 3959–3970 (1991).

848 90. L. Avery, H. R. Horvitz, Effects of starvation and neuroactive drugs on feeding in
849   *Caenorhabditis elegans*. *J. Exp. Zool.* **253**, 263–270 (1990).

850 91. S. T. Laing, A. Ivens, V. Butler, S. P. Ravikumar, R. Laing, D. J. Woods, J. S. Gilleard, The
851   transcriptional response of *Caenorhabditis elegans* to ivermectin exposure identifies novel
852   genes involved in the response to reduced food intake. *PLoS One*. **7**, e31367 (2012).

853 92. N. C. Sangster, J. M. Redwin, H. Bjorn, Inheritance of levamisole and benzimidazole
854   resistance in an isolate of *Haemonchus contortus*. *Int. J. Parasitol.* **28**, 503–510 (1998).

855 93. L. F. Le Jambre, J. H. Gill, I. J. Lenane, P. Baker, Inheritance of avermectin resistance in
856   *Haemonchus contortus*. *Int. J. Parasitol.* **30**, 105–111 (2000).

857 94. S. Shompole, D. P. Jasmer, Cathepsin B-like cysteine proteases confer intestinal cysteine
858   protease activity in *Haemonchus contortus*. *J. Biol. Chem.* **276**, 2928–2934 (2001).

859    95. D. P. Jasmer, C. Yao, A. Rehman, S. Johnson, Multiple lethal effects induced by a
860           benzimidazole anthelmintic in the anterior intestine of the nematode *Haemonchus*
861           *contortus*. *Mol. Biochem. Parasitol.* **105**, 81–90 (2000).

862    96. I. J. I. Janssen, J. Krücken, J. Demeler, G. von Samson-Himmelstjerna, *Caenorhabditis*
863           *elegans*: modest increase of susceptibility to ivermectin in individual P-glycoprotein loss-
864           of-function strains. *Exp. Parasitol.* **134**, 171–177 (2013).

865    97. A. P. Page, The sensory amphidial structures of *Caenorhabditis elegans* are involved in
866           macrocyclic lactone uptake and anthelmintic resistance. *Int. J. Parasitol.* **48**, 1035–1042
867           (2018).

**Acknowledgements:** This work was supported by the Biotechnology and Biological
Sciences Research Council (BBSRC) [BB/M003949]; the Scottish Government's Rural
and Environment Science and Analytical Services (RESAS) division; a University of
Glasgow James Herriot Scholarship; the Wellcome Trust [206194 and 216614/Z/19/Z];
and UKRI [MR/T020733/1]. For the purpose of Open Access, the authors have applied
a CC BY public copyright licence to any Author Accepted Manuscript version arising
from this submission. We would like to acknowledge members of the BUG Consortium
and Parasite Genomics group at the Wellcome Sanger Institute for insightful
discussions throughout this project. We also thank Pathogen Informatics and DNA
Pipelines (WSI) for their support and expertise and the Biosciences Division at the
Moredun Research Institute for expert care and animal assistance.

**Funding:** Biotechnology and Biological Sciences Research Council (BBSRC;
BB/M003949) (RL, MB, ED, JAC, NS, BM, DB, CB). Wellcome Trust's core funding of
the Wellcome Sanger Institute (grant WT206194) (MB). University of Glasgow James
Herriot Scholarship (AA). Wellcome Clinical Research Career Development Fellowship
(216614/Z/19/Z) (RL). UKRI Future Leaders Fellowship (MR/T020733/1) (SRD).

**Author contributions:** Conceptualisation: RL, AT, ED, JAC. Investigation: SRD, RL,
DB, AM, KM, AA, CB, UC, IF, JM. Formal analysis: SRD, RL. Software: SRD.
Resources: DB, AM, RK, NS. Supervision: CB, JSG, NS, MB, ED, JAC. Project
administration: NH, ED. Writing - Original Draft: SRD, JAC, RL, ED. Writing - Review &
Editing: All authors. Funding acquisition: RL, AT, MB, ED, JAC, NS, BM, DB, CB, SRD.

**Competing interests:** Authors declare that they have no competing interests.

**Data and materials availability:** Raw sequencing data for this study are outlined in
**table S1** and are archived under the ENA study accession PRJEB4207. The *H.
contortus* genome assembly and manually curated annotation resources are publicly
available at
https://parasite.wormbase.org/Haemonchus_contortus_prjeb506/Info/Index/. The code
used to generate and analyse data and to plot figures can be found at
https://github.com/stephenrdoyle/hcontortus_X-QTL.

**Supplementary Materials**

Materials and Methods

Supplementary Text

Figs. S1 to S10

Tables S1 to S2

References (72 - 97)