1 **TITLE:** *Marchantia polymorpha* gene expression atlas reveals the hierarchy of abiotic
2 stress responses and conservation of diurnal gene expression
3

4 **AUTHORS:** Qiao Wen Tan[1], Peng Ken Lim[1], Zhong Chen[2], Asher Pasha[3], Nicholas
5 Provart[3], Marek Mutwil[1*]
6
7 [1]School of Biological Sciences, Nanyang Technological University, 60 Nanyang Drive,
8 Singapore, 637551, Singapore
9
10 [2]Amoeba Education Hub, 1 West Coast Road, 128020, Singapore
11
12 [3]Department of Cell and Systems Biology/Centre for the Analysis of Genome Evolution
13 and Function, University of Toronto, Toronto, Ontario M5S 3B2, Canada
14
15
*Corresponding author:
Marek Mutwil
School of Biological Sciences,
Nanyang Technological University,60 Nanyang Drive,
637551, Singapore,
Singapore
Email: mutwil@ntu.edu.sg

Abstract
Abiotic stress has a dramatic impact on ecosystems and the yield of crops, and global
warming will likely result in more frequent and intense stresses. However, due to the
complexity of the underlying signalling pathways and the fact that stresses often occur
in combinations in nature, our knowledge of how plants acclimatize to abiotic stress is
still lacking. Here, we used a plant with minimal regulatory network redundancy,
Marchantia polymorpha, to study how seven abiotic stresses alone and in combination
affect the phenotype, gene expression, and activity of biological pathways. Worryingly,
we see no evidence of conservation of gene expression to abiotic stress between model
organisms, suggesting that each species develops unique strategies to cope with
abiotic stress. Interestingly, we observed that certain stresses are able to suppress
others, while specific combinations can elucidate novel transcriptomic responses.
Finally, we provide two online databases to study abiotic stress responses and diurnal
gene expression data in Marchantia.

Introduction
The colonization of land by plants, which occurred around 470 Ma, was essential to
establish habitable environments on land for all kingdoms of life [1]. Bryophytes (which
include mosses, liverworts, and hornworts) represent the earliest diverging group of
non-vascular land plants [2–4]. Morphology of the earliest land plant fossils (which
consists primarily of tissue fragments and spores from the Middle Ordovician around
470 Ma) showed early land plants were liverwort-like [5,6]. As a liverwort, *Marchantia*
polymorpha is a valuable model to better understand the emergence and evolution of
land plants, as it allows us to compare the biology of aquatic algae, non-vascular plants
to vascular, seed, and flowering plants. Studying the Marchantia biology can reveal the
reason behind the successful terrestrialization event, as *Marchantia* contains traits
essential for terrestrialization and increased complexity (e.g., hormones auxin,
jasmonate, salicylic acid, protection mechanisms against desiccation, photooxidative
damage)[7], albeit at a less complex organization than vascular plants [8].
Apart from its interesting evolutionary position among land plants, Marchantia is
a valuable model to study basic plant biology. Largely due to the lack of whole-genome
duplications in the liverwort lineage, *Marchantia* shows a simpler, redundancy-free
regulatory genome [8], that together with the ease of growth and genetic manipulation [9],
make Marchantia an excellent model to study general plant biology. The *Marchantia*
genome contains all necessary components for most land-plant signalling pathways, but
compared to flowering plants, the components are reduced to the minimum number,
making it easier to dissect the pathways [8]. For example, the auxin signalling network in
Marchantia is simple yet functional, with all relevant genes existing as single orthologs
[10]. Similarly, cellulose biosynthesis in Marchantia uses the same but simplified
machinery; while *Arabidopsis thaliana* contains ten cellulose synthases that exist in

multimeric complexes, Marchantia has only two [11]. Furthermore, since the dominant
generation of *Marchantia* is the haploid gametophyte, heterozygosity can be
circumvented to study mutant and transgenic phenotypes directly. Different genetic
combinations can be obtained by crossing, where eggs produced by the archegonia are
fertilized by motile sperm originating from antheridia [12]. Alternatively, clonal lines can be
easily produced asexually by small disc-shaped propagules known as gemmae, which
are found on the dorsal side of the haploid thallus [13]. Further advantages of *Marchantia*
include the option to cryopreserve of gemmae [14] and spermatozoa [15], ~3 months sexual
generation time, 2–3 weeks for asexual reproduction, and a small genome size (225.8
Mb, nine chromosomes)[8,9]. These features make *Marchantia* an attractive model to
dissect the function of genes and biological pathways. Indeed, with >500 publications
available on NCBI at the time of writing and with >50 publications in 2021 alone,
Marchantia is a model organism on the rise.
Extreme environmental conditions can cause a multitude of biotic and abiotic
stresses that can devastate crops and induce the collapse of entire ecosystems [16,17].
Plants have evolved sophisticated mechanisms to perceive and respond to the different
stresses, which induces an acclimation process that allows the plant to survive the
stress [18], but often at the cost of reduced growth [16,17]. Many studies have analysed the
effect of stress on plant growth by identifying differentially expressed genes between
stress-treated and untreated plants (e.g., [19–21]), or by identifying single nucleotide
polymorphisms associated with stress resistance in *Arabidopsis* and maize [22–24]. The
studies performed on non-agricultural species such as *Arabidopsis* can shed light on
fundamental processes of stress acclimation, but it is currently unclear whether the
acclimation mechanisms between different species or even accessions are conserved
and transferable from model to crop.
In addition, plants are often exposed to a combination of stresses in the natural
environment which can require opposing strategies for survival. For example, drought
causes plants to close their stomata to minimize water loss [25,26], while heat causes the
stomata to open to cool down their leaves via transpiration [21,27,28]. The stress signalling
is mediated by a diverse ensemble of stress-specific sensors/receptors, networks of
protein kinases/phosphatases, calcium channels/pumps, and transcription factors that
can be localized to different organelles [29,30]. Stress signalling is further communicated
and attenuated by hormones, other signalling molecules (e.g., reactive oxygen species)
and protein modifications (s-nitrosylation, ubiquitination, myristoylation) [31,32]. Due to this
complexity, it is challenging to elucidate the molecular basis of stress acclimation to one
stress. The problem becomes even more intangible when multiple simultaneous
stresses are considered. Apart from antagonistic reactions to combined stresses, it has
been observed that a plant's response to different stresses is not equal and is
dominated by the stress that is perceived to be more severe by the plant. This
phenomenon has been observed when abiotic and biotic stresses intersect (refer to the

chapter by [33] for examples) and in combinations of heat and drought stress [34].
Therefore, a simpler model plant system, such as *Marchantia,* would be valuable in
studying stress acclimation.
In this study, we set to take advantage of *Marchantia*'s less complex signalling
architecture to understand better how plants respond to environmental cues such as
stress and time of the day to modulate the expression of genes and biological
pathways. To this end, we constructed an abiotic stress gene expression atlas of
Marchantia comprising seven abiotic stresses (darkness, high light, cold, heat, nitrogen
deficiency, salt, mannitol) and their pairwise combinations (e.g., cold + salt). We first
compared the stress acclimation responses to other members of the plant kingdom to
determine whether these responses are conserved across species. We then identified
stress-responsive biological pathways and investigated whether new gene families
evolved specifically to confer adaptations to abiotic stress. Next, we compared the
transcriptomic responses of the different stress combinations (cross-stress) to identify
the differentially regulated pathways and how the different stresses enhance or
suppress one another. We then constructed a diurnal gene expression atlas and
investigated whether diurnal gene expression is conserved between *Marchantia* and
Archaeplastida. Finally, to make the data readily accessible to non-bioinformaticians, we
provide i) an eFP browser for Marchantia that allows the visualization of gene
expression in organs and stress conditions and ii) an updated CoNekT platform,
allowing sophisticated comparative gene expression and co-expression analyses.
124
Material and Methods
Maintenance of *Marchantia polymorpha*
Male *Marchantia polymorpha*, accession Takaragaike-1 (kindly provided by Prof. Chen
Zhong from National Institute of Education, Singapore) was propagated on half-strength
Gamborg B-5 Basal agar (1% sucrose, pH 5.5, 1.4% agar)[35] in deep well plates (SPL
Biosciences, SPL 310101) at $24^{\circ}C$ under continuous LED light at 60 $\mu Em^{-2}s^{-1}$.
131
Experimental setup for stress experiment
In order to determine the ideal stress condition to be used for cross-stress experiments,
a series of severity was used for heat, cold, salt, osmotic, light, dark, and nitrogen
deficiency stresses. For each stress variation, three agar plates containing nine
gemmae each were plated, where two plates were used as material for RNA
sequencing and one plate was kept for observation until 21 days after plating (DAP).
For salt, osmotic, and nitrogen deficiency stress, the gemmae were plated on
half-strength Gamborg B-5 Basal agar (pH 5.5, 1.4% agar, 12.4 mM KNO_3)
supplemented with 20 - 200 mM NaCl (20 mM steps), 50-400 mM mannitol (50 mM
steps) and KNO_3 concentration ranging from 90 to 0% (22.3 - 0 mM, 2.5 mM steps),
respectively. The potassium concentration in nitrogen deficiency agar was maintained

using equimolar concentrations of KCl. Gemmae subjected to heat, cold, light, and dark
stresses were plated on normal half-strength Gamborg B5 agar (pH 5.5, 1.4% agar,
12.4 mM KNO_3, 0.5 mM $(NH_4)_2SO_4$).
Plates were grown at 24^oC under continuous LED light at 60 $\mu Em^{-2}s^{-1}$ from days
0 to 13. For dark stress, plates were moved to the plant growth chamber (HiPoint M-
313) on days 8, 9, 10, 11, 12, 13, and 14 for growth in darkness at 24^oC for 7, 6, 5, 4, 3,
2 and 1 day(s), respectively. On day 14, all plates were transferred to the plant growth
chamber (HiPoint M-313) for 24 hours. Control and plates for salt, osmotic, and nitrogen
deficiency stresses were maintained at 24^oC under continuous LED light at 60 $\mu Em^{-2}s^{-1}$
in the plant growth chamber. Normal growth conditions were used for heat, cold, dark
and light stresses with the following modifications. For heat and cold stress, 27^oC to
36^oC (3^oC steps) and 3^oC to 12^oC (3^oC steps) were used, respectively. For light stress,
plants were subjected to 20% to 100% chamber capacity of light output (115 to 535
$\mu Em^{-2}s^{-1}$), at steps of 20%, approximating to 100 $\mu Em^{-2}s^{-1}$ per step.
On day 15, whole plants were harvested by pooling three plants into 2 mL
Eppendorf tubes, flash-frozen in liquid nitrogen, and stored at -80^oC. This was done in
triplicates, resulting in 3 (replicates) * 3 (pooled plants) used for each RNA-seq sample.
Pictures of the front and back of the observation plates were taken, and the plates were
returned to normal growth conditions. Pictures of the front and back of the observation
plates were taken on day 21 without cover.
Cross stress experiments were carried out similarly with the following parameters
for the various stress combinations - heat (33^oC), cold (3^oC), salt (40 mM NaCl),
osmotic (100 mM mannitol), light (435 $\mu Mm^{-2}s^{-1}$), darkness (3 days) and nitrogen
deficiency (0 mM KNO_3).
167
Diurnal gene expression
For diurnal gene expression, three agar plates containing nine gemmae each were
plated for each sampling time point on half-strength Gamborg B-5 Basal agar (0%
sucrose, pH 5.5, 1.4% agar). The plants were grown at 24^oC under continuous LED
light at 60 $\mu Em^{-2}s^{-1}$. On day 8, plates were transferred to the plant growth chamber and
grown at 24^oC under LED light at 60 $\mu Em^{-2}s^{-1}$ with a 12 h/12 h day/night cycle. On day
15, plants were harvested at Zeitgeber time (ZT) 2, 6, 10, 14, 18, and 22. The plants
were stored in 2 mL Eppendorf tubes (3 plants per tube) and flash-frozen in liquid
nitrogen before storage at -80^oC.
177
Size measurement of *Marchantia polymorpha*
The photos of the observation plates were scaled and measured on Adobe Illustrator.
Length and breadth of the thallus were taken with relation to the central axis (Figure 1B
shows the plants with the central axis in a horizontal position). The thallus's approximate

area was derived by the product of the length and breadth of the thallus. Abnormally
small plants (outliers) were excluded from further analysis.
184

RNA extraction and sequencing

Plants from each Eppendorf tube were ground into fine powder in liquid nitrogen using
mortar and pestle. Total RNA was extracted using the Spectrum™ Plant Total RNA Kit
(Sigma, STRN-250) using Protocol A (750 µL Binding Solution) according to
manufacturer's instruction with on-column DNase digestion using 60 µL of DNase
mixture (15 µL RQ1 RNase-Free DNase (Promega, M6101), 6 µL RQ1 DNase 10X
Reaction Buffer and 19 µL nuclease-free water) per column.

Preliminary quality control of the extracted RNA (triplicates for each condition)
was done using Nanodrop before further quality control checks by Novogene
(Singapore) for sample quantitation, integrity, and purity using Nanodrop agarose gel
electrophoresis and Agilent 2100 Bioanalyzer. Library construction from total RNA
including eukaryotic mRNA enrichment by oligo(dT) beads, library size selection, and
PCR enrichment was performed by Novogene using NEBNext® Ultra™ II Directional
RNA Library Prep Kit for Illumina®. The libraries were then sequenced with Illumina
Novaseq-6000, paired-end sequencing at 150 base pairs and at sequencing depths of
~20 million reads per sample.
201

Expression quantification

203 RNA sequencing data were mapped against the *Marchantia polymorpha* CDS (v5.1
204 revision 1, MarpolBase), quantified, and TPM-normalised (transcript per million) using
205 Kallisto v 0.46.1. Publicly available RNA sequencing data for *Oryza sativa* were
206 downloaded from ENA as fastq files and mapped against the CDS of *Oryza sativa* v7.0
207 obtained from Phytozome v13 using Kallisto.
208

Identification of differentially expressed genes

Non-normalized counts from Kallisto were used for the analyses of differentially
expressed genes (DEGs) using R package DESeq2 [36] where various stress conditions
were compared against their respective controls.

For our *M. polymorpha* dataset, only genes that were found to be differentially
expressed against controls from two different batches were considered for further
analysis. For downstream analysis, only genes with an adjusted p-value < 0.05 and a

$-1 > \log_2$-fold > 1 were considered as differentially expressed. DEGs of *Arabidopsis*
thaliana, *Chlamydomonas reinhardtii*, and *Cyanophora paradoxa* were obtained from
the study of [20].

Identification of transcription factors, kinases, and differentially expressed pathways

The biological function and pathway membership of genes of *Cyanophora paradoxa, Chlamydomonas reinhardtii, Marchantia polymorpha, Oryza sativa,* and *Arabidopsis thaliana* were annotated using Mercator 4 v2.0[37]. The annotation of transcription factors belonging to *C. reinhardtii*, *M. polymorpha*, *O. sativa*, and *A. thaliana* was retrieved from PlantTFDB v5.0[38], while the annotation of kinases was obtained from iTAK v18.12[39]. Transcription factors and kinases of *C. paradoxa* were inferred from orthologs of the four species identified by OrthoFinder (data from [40]).

Significantly differentially expressed pathways were determined through hypergeometric testing, where the observed number of DEGs in a pathway was compared to the permuted number of DEGs. The p-values were adjusted for multiple testing using Benjamini Hochberg correction (p-value < 0.05)[41].

Comparison of species and stresses with differentially expressed biological pathways and orthogroups

To estimate the similarities of species/stresses by the differentially expressed biological pathways, we determined for each species/stress the identity biological pathways showing significant enrichment upregulated (U), downregulated (D), both (UD) or not changed (N). Then, each experiment was represented as a vector of U, D, UD, or N values. The Jaccard distance between each experiment was calculated and used as the linkage for the cluster map. For a fair comparison of the experiments and species, biological pathways that were N (not changed) for both species/stresses were excluded. The modified Jaccard's distance between species/stress experiments X and Y is:

$$JD_{(X,Y)} = 1 - \frac{Gene\ family\ statuses\ in\ X\ \cap\ Gene\ family\ statuses\ in\ Y}{Gene\ family\ statuses\ in\ X\ \cup\ Gene\ family\ statuses\ in\ Y}$$

For the analysis where we used orthogroups as a basis for comparison, we used the same procedure, but with the following changes: for each species, an orthogroup was labeled as "None" if it did not contain any genes, "no change" if it did not contain DEGs (i.e., the expression was not affected by stress), "ambivalent" if it contained up and downregulated genes, and "upregulated" and "downregulated" for orthogroups that only contain up or downregulated genes, respectively. Then, each experiment was represented as a vector, where the items represented orthogroups with 'None', 'no change', 'ambiguous', 'upregulated' or 'downregulated' items. For a fair comparison of the experiments and species, gene families that are not present in either experiment (due to certain orthogroups not being present in all species), and the absence of DEGs from gene families in both experiments, were excluded.

255   The JD values range from 0 (the two species/stresses contain identical
256 differentially expressed orthogroups or biological pathways) to 1 (the two
257 species/stresses are entirely different). The JD values were used as linkages for
258 clustering of the species/stresses.
259

Stress responsiveness of biological pathways and phylostrata
We compared the stress responsiveness of the orthogroups across various species and
stresses by assigning gene families into five categories (as done above) for each stress:
no genes found in family ('None'), no change in gene expression (NC, i.e., no DEGs),
only upregulated DEGs are present (UP), only downregulated DEGs are present
(DOWN), and both up- and down-regulated DEGs are present (AMB). We looked at the
distribution of biological processes of differentially expressed genes that are found in
the orthogroups that were responsive to stress. The counts were logged and row
normalised by the total number of genes present in a Mapman bin.
A phylostratigraphic analysis was performed on the *Marchantia* stress
perturbations and the corresponding stresses in the four other species. Through the
statuses of the orthogroups assigned earlier, we considered an orthogroup to be stress-
responsive if there were at least two of the same status (UP/DOWN/AMB) across
species that are not None or NC. Alternatively, if there was a mixture of statuses, the
orthogroup is assigned MIXED for that stress. The age of the orthogroup was
determined by the presence of the oldest species present in our analysis. Similarly,
species-specific orthogroups were assigned statuses based on their initial assignments.
All other scenarios were considered as non-responsive and assigned the status of 'NR'.
278

Comparison of single and cross stresses with UpSet plot
280 Using the DEGs obtained from differential gene expression analysis, we looked at the
281 top 50 intersections of all perturbations of *Marchantia* in terms of up and downregulated
282 genes through an upset plot, provided by https://upsetplot.readthedocs.io/en/stable/.
283

Determination of rhythmic genes and biological pathways
Rhythmic genes were identified using JTK_Cycle v3[42] with parameters set at six
timepoints with three replicates, temporal spacing of 4 hours between samples, and
target period of 6 hours. Genes are considered expressed if the TPM value is more than
1 in at least one replicate. Genes are only considered rhythmic if it is supported by an
adjusted p-value of < 0.05. The LAG value was corrected by two hours (i.e., LAG 0 to
LAG 2) as our first timepoint is ZT2, meaning two hours after initial exposure to light.

To observe the rhythmicity of genes according to their biological function, we
calculated the percentage of rhythmic genes that peak at the respective LAGs identified
by JTK_Cycle for each Mapman bin. The linkage between the Mapman bins was
determined by scipy hierarchical clustering.
295

Comparison of diurnal gene expression in *Marchantia* to other species
We compared the diurnal expression of 1-to-1 orthologs as done in [43]. JTK data from
this publication included species *Cyanophora paradoxa, Chlamydomonas reinhardtii,*
Porphyridium purpureum, Klebsormidium nitens, Physcomitrella patens, Selaginella
moellendorffii, Picea abies, Oryza sativa, and *Arabidopsis thaliana.* In addition, we also
looked at the number of orthologs that have the same phase in *Marchantia* and another
species after applying a phase shift in the other species.
303

304 **Data availability**
305 The RNA-seq data capturing the expression of controls, single and double stresses are
306 available from https://www.ebi.ac.uk/ena as E-MTAB-11141.
307

Script availability
Python and bash scripts used to generate the figures in the paper are available from:
https://github.com/tqiaowen/marchantia-stress

Results
Subjecting *Marchantia polymorpha* to a combination of abiotic stresses
To capture gene expression changes caused by single stress or a combination of
stresses, we first set to establish the type and magnitude of the different stresses. We
defined two types of stresses: i) the environmental stresses comprised heat, cold,
excess light, and prolonged darkness, while ii) media stresses comprised nitrogen
deficiency, excess salt (representing ionic and osmotic stress), and excess mannitol
(representing osmotic stress). Next, the magnitude of the stresses was modulated to
identify sub-lethal stress conditions or growth decrease by ~50% (inferred from
approximate thallus area). To this end, gemmae grown on sealed, sterile agar plates at
24 hours light were subjected to varying degrees of stress, and their phenotypes were
observed on days 15 and 21. For media stresses, the gemmae were subjected to the
stress from day 0, while for the environmental stresses, the stress was applied at day
14 for 24 hours (Figure 1A). Prolonged darkness was an exception to this design, as
plates were subjected to darkness on days 8, 9, 10, 11, 12, 13, and 14 to expose the
plants to 7, 6, 5, 4, 3, 2, and 1 day(s) of darkness at day 15, respectively (Figure 1A).
Single stresses showed varying degrees of effect on plant growth at day 15 (day
of harvest, Figure S1 shows growth measurements, Supplemental Data 1 shows agar
plates) and day 21 (6 days post-stress for environmental stresses, Figure 1B). One day

of cold stress did not affect the growth at the temperature range tested (3-12°C), and we
selected 3°C for further analysis. The heat stress experiment showed that the plants
abruptly died when the heat treatment temperature was increased from 33°C (no
phenotype) to 36°C (death), and we selected 33°C for further study. For light stress, we
selected 435 $\mu Em^{-2}s^{-1}$) as we observed necrosis at the next higher light intensity (535
$\mu Em^{-2}s^{-1}$, Supplemental Data 1). For osmotic (100 mM selected) and salt stress (40mM
selected), we observed an expected negative growth gradient when the concentration of
the two compounds was increased. For nitrogen deficiency, at 0 mM KNO_3, we
observed a decrease in growth and an accumulation of a red pigment, which likely
represents auronidin, a flavonoid shown to accumulate during phosphate deficiency [44].
Finally, for darkness, we observed that the growth of plants decreased proportionately
with the duration of stress, and we selected plants grown in three days of darkness, as
they showed a growth decrease of 50% (Figure S1). On day 21 (i.e., six days of normal
growth condition), all dark-grown plants showed increased size, indicating that the
seven days of darkness are not lethal.
Next, we set out to determine how *Marchantia* responds to a combination of two
stresses. To this end, we tested all 19 possible pairs of stresses (cold+heat and
dark+light combinations are mutually exclusive and hence, excluded) using the same
experimental outline as for single stresses (Figure 1A). We did not observe any
unexpected additive effects of any two stresses (Figure 1D), as typically, a combination
of stresses resulted in an expected combination of phenotypes (e.g., nitrogen: small,
pigmented, mannitol: small, nitrogen+mannitol: even smaller and pigmented) except for
salt-nitrogen (SN), which is significantly larger than its salt counterpart but not different
from its nitrogen counterpart (Figure 1E). While plants subjected to a combination of
sub-lethal heat (33°C) and high light (435 $\mu Em^{-2}s^{-1}$) died (Figure 1D), this was most
likely due to a greenhouse effect caused by high irradiation and sealed plates, as the
temperature of agar climbed to 38°C (i.e., lethal temperature, Figure 1C). Interestingly,
for high light and cold, we observed yellowing of the thalli, indicating that lowering the
temperature makes the plants more sensitive to high light (Figure 1C).
Similar to the single stress experiments, the RNA of plants subjected to the
combination of two stresses were isolated. The resulting panel of 7 single stresses, 18
combinations of two stresses and 2 untreated controls (controls from single stress and
combined stress batches) were sent for RNA sequencing (Table S1 contains sample
labels). The resulting data was used to estimate gene expression (Table S2, S3
contains Transcript Per Million (TPM) values and raw counts, respectively). Overall, we
observed a good agreement between the sample replicates, as samples showed
expected clustering (Figure S2), and the correlation between expression profiles of
replicates was >0.97 (Table S4).

369
Figure 1. Influence of different abiotic stresses on the growth of *Marchantia*
polymorpha. A) Overview of the experimental setup for stress experiments. Gemmae
were plated on ½ Gamborg B-5 agar with necessary modifications relevant to the stress
tested. Black lines below the timeline indicate the duration where plants were exposed
to stress. All plates were moved to the plant growth chamber on day 14 at the growth
conditions appropriate to the respective stresses. Plants were sampled on day 15,
where three plants were pooled as a replicate sample for RNA-seq. Observation plates
were returned to normal growth conditions on day 15, and photographs of the plates
were taken on days 15 and 21. B) Phenotype of plants on day 21 for heat, cold,
osmotic, salt, light, dark and nitrogen deficiency stresses. Conditions with underline

represent the stress intensities used for combined stress. Controls F and L are
representatives from the batches at the start and end of independent stresses in panel
B. C) Phenotypes of plants on day 21 for combined stresses. Controls O and R are
representatives from the start and end of combined stresses in panel C. D and E)
Thallus size of gemmalings on Day 15 and 21 of growth (1/7 day after physical stress,
15/21 days after chemical stress, error bars are represented by standard deviation and
Student's two-tailed t-test, p-value < 0.05. Asterisk (*) represents significant difference
to control, caret (^) represents significant difference to first single stress control and plus
sign (+) indicates significant difference to second single stress control.

No evidence for conservation of abiotic stress responses in Archaeplastida

Model organisms, such as *Arabidopsis thaliana*, are studied with the aim to apply the
gained knowledge to other plants, such as food crops. The usage of model organisms
assumes that there is a degree of conservation of gene sequences, biological
pathways, and how the different organisms respond to stress. To study the conservation
of stress responses to abiotic stress across multiple members of Archaeplastida, we
compared our stress expression data of *Marchantia polymorpha* (Mpo: heat, cold, high
light, darkness, low nitrogen, salt, mannitol) with publicly available expression data for
basally diverging alga *Cyanophora paradoxa* (Cpa: heat, cold, high light, darkness, low
nitrogen, low sulphur, low phosphate, salt, mannitol), rice *Oryza sativa* (Osa: cold, salt,
drought) and *Arabidopsis thaliana* (Ath: heat, cold, high light, darkness, low nitrogen,
low sulphur, low phosphate, low iron, salt) (Table 1, Table S5)[45,46].
To assess how abiotic stress affects global gene expression, we first identified
the percentage of differentially expressed genes (DEGs). Since our experiment was
done in batches, we used two controls taken at the beginning and the middle of the
experiment and set the requirement for a gene to show conserved differential
expression in both controls to be deemed a DEG (Table S6 and S7). Overall, we
observed a good agreement between the two controls, as both volcano plots (Figure
S3) and set comparisons (Figure S4) indicated a similar set of DEGs. Standard
differential gene analysis was also done for *O. sativa*, and a summary of the DESeq2
output and list of DEGs can be found in Tables S8 and 9, respectively. We observed
that typically, only a small percentage of genes are differentially expressed, and the
percentage of DEGs varies across species and stress, with dark stress having the most
DEGs across all species (Figure 2A). The transcription factors (TFs) also comprised a
small number of DEGs (Figure 2B), with the highest percentage in *A. thaliana* and *O.
sativa* in general (list of TFs Table S10). The percentage of protein kinase DEGs follow
the same trend as TFs, with the highest percentage in Arabidopsis and rice (Figure 2C)
(list of kinases Table S11). In most experiments, there is an approximately even amount
of up and downregulated genes except in *Arabidopsis* salt stress, where there is a
disproportionate amount of downregulated genes and a larger proportion of upregulated
genes in *Cyanophora* light, *Marchantia* mannitol and nitrogen, *Oryza* salt-1, and
Arabidopsis low phosphate and low iron (Figure 2D).

Next, we wanted to understand which biological pathways are differentially
expressed during stress and how these responses are conserved in Archaeplastida. To
this aim, we first identified pathways that were significantly upregulated (more genes
from the pathway are DEGs than expected by chance, adjusted p-value<0.05),
downregulated, or both (a pathway contains more up- and down-regulated genes than
expected by chance). This analysis was performed for each species and stress
combination in our dataset, giving us stress/species (x-axis) and biological process
enrichment (y-axis) matrix. The matrix was then subjected to hierarchical clustering,
which grouped similar species/stress responses (by clustering columns) and similarly
responding biological pathways (by clustering rows) (Figure 2E). Overall, we observed
no apparent conservation between stresses in different organisms as the different
stresses did not cluster well. At the same time, we did not observe any biologically
meaningful clustering of biological pathways. For example, unrelated protein
translocation pathways, chromatin organization and vesicle trafficking showed the
highest similarity across all species/stresses (Figure 2E). To confirm these findings, we
also investigated whether there are similarities in responses of gene families (Table
S12). In this analysis, we assumed that if two species/stresses are similar, they should
show a similar pattern of up- or down-regulated gene families. However, similar to the
results for biological pathways, we observed no clear grouping of species/stresses
(Figure S5), suggesting that gene expression responses are not conserved across
species.

Table 1. Summary of stresses analysed across the five species in this study.

	Cpa	Cre	Mpo	Osa	Ath
Heat	37^{o}C, 12 h (in darkness)	-	33^{o}C, 24 h	-	44^{o}C, 4h
Cold	4^{o}C, 12 h (in darkness)	-	3^{o}C, 24 h	10^{o}C, 10h	10^{o}C, 24 h
Osmotic	100 mM mannitol, 96 h	-	100 mM mannitol, 15 days	10% available water content (AWC), 3 days	-
Salt	75 mM NaCl, 96 h	-	40 mM NaCl, 15 days	300 mM NaCl, 24 h	150 mM NaCl, 13 days
High light	150 µEm^{-2}s^{-1}, 24 h	145 µEm^{-2}s^{-1}, 8 h	435 µEm^{-2}s^{-1}, 24 h	-	300 µEm^{-2}s^{-1}, 4 h
Prolonged darkness	0 µEm^{-2}s^{-1}, 3 days	-	0 µEm^{-2}s^{-1}, 3 days	-	0 µEm^{-2}s^{-1}, 2 days
Nitrogen deficiency	10% recommended KNO$_3$, 96 h	No nitrogen source, 2 days	0 mM KNO$_3$, 15 days	-	No nitrogen source, 2 days
Other nutrient deficiencies	Y	Y	-	-	Y

Figure 2. Comparative analysis of differential gene and pathway changes across species and stresses. A) Percentage of genes that are significantly differentially expressed. B) Percentage of differentially expressed transcription factors. C) Percentage of differentially expressed kinases. D) Percentage of up and downregulation of differentially expressed genes. E) Summary of biological pathways, represented by **M**apman bins, that are significantly enriched for differentially expressed genes. Colours in the heatmap, red, pink, blue, and gray correspond to significant upregulation (U), up and downregulation (UD), downregulation (D), and no significant change (N), respectively. Experiment labels are coloured according to species, with blue, orange, green, purple, and red corresponding to *C. paradoxa*, *C. reindhardii*, *M. polymorpha*, *Oryza sativa,* and *A. thaliana,* respectively. The following abbreviations are used for the

457 description of stresses: cold (C), dark (D), heat (H), light (L), mannitol (M), nitrogen
458 deficiency (N), salt (S), iron deficiency (minFe), sulfur deficiency (minS) and phosphorus
459 deficiency (minP).

460
No evidence for the specialization of newly evolved gene families to stress
acclimation
While gene expression response was not conserved in different species, the enrichment
analysis in the previous figure revealed that specific biological pathways are more
stress-responsive than others. For example, MapMan bin comprising 'Nutrient uptake'
was more significantly differentially expressed across the different species/stresses than
'Vesicle transport' (Figure 2E).
To identify which biological pathways are highly responsive to the seven tested
stresses in *M. polymorpha*, we first identified how many stresses a given gene family
(represented by orthogroups) responds to. We observed that certain orthogroups are
responsive to multiple stresses and have genes that are significantly upregulated (up;
Table S13), downregulated (down), ambivalent (amb; some genes are up-, others are
down-regulated), or not changed (nc) for one or more stresses. We then assigned the
gene families to biological processes and plotted the stress responsiveness
occurrences of the gene families (Figure 3A). Overall, we observed that gene families
involved in DNA damage response, cell cycle, chromatin organization, protein
biosynthesis/translocation are typically responding to one or two abiotic stresses,
suggesting that these pathways are typically not affected by abiotic stress. Conversely,
gene families involved in various uncharacterized enzymatic reactions ('Enzyme
classification'), transcriptional gene regulation ('RNA biosynthesis') and protein
degradation, cell wall, solute transport, and nutrient uptake were frequently differentially
regulated during up to seven abiotic stresses, suggesting that these pathways are
highly responsive to abiotic stress (Figure 3A).
Recently established gene families could be important for adaptation to the local
environment. To test this hypothesis, we investigated whether there is a dependency
between the age of a gene family and its stress responsiveness. First, by using
phylostratigraphy (Table S14[47]), we assigned gene families to eight phylostrata. We
sorted them from oldest to youngest (archaeplastida:Plantae, viridiplantae: green plants,
embrophyta: land plants), and the species-specific phylostrata Mpo (*Marchantia*), Cpa
(*Cyanophora*), Ath (*Arabidopsis*), Osa (*rice*), and Cre (*Chlamydomonas*). We then
plotted the number of gene families, and the number of abiotic stresses they respond to
and color-coded them by their phylostrata (Figure 3B). For all phylostrata, we observed
a clear negative relationship between the number of gene families (y-axis) and the
stress responsiveness (x-axis), indicating that most gene families typically respond to at
most one or two stresses. Interestingly, the plots for the different phylostrata were
largely parallel (e.g., Archaeplastida and Marchantia-specific lines are parallel and

overlapping), suggesting that there is no dependence between the age of gene families
and their age. The only exceptions to this trend were rice- (Osa) and Chlamydomonas-
specific (Cpa) gene families that showed a more negative slope, but this can be
explained by these species only having data for two and three stresses, respectively.
We conclude that there is no dependency between the age of gene families and
their responsiveness to stress. Thus, more recent gene families have likely not evolved
to help plants acclimatize to abiotic stress.

**Figure 3. Analysis of stress responsiveness, biological function, and age of gene
families.** A) The proportion of functionally annotated orthogroups responding to the
seven single (e.g., cold, heat) abiotic stresses in *Marchantia* and the equivalent stresses
in the four other species. The data was row-normalized to show the proportion of the
orthogroups responding to the number of abiotic stresses. B) Number of orthogroups
found in phylostrata at various levels of stress responsiveness.

Combinatorial analysis reveals the hierarchy of stress responses in Marchantia

Plants experience many stresses when growing in nature, and these stresses need to
be perceived and responded to. To better understand how Marchantia responds to
single stress or a combination of two stresses, we first identified differentially expressed
genes in the 18 combinations of two stresses and then compared the number of up- and
down-regulated genes with the single stresses. We observed that the combination of
stresses typically contains a similar or higher number of DEGs when compared to single
stresses (Figure 4A). Next, to see if a severe growth phenotype results in a large
number of differentially expressed genes, we plotted the plant size (x-axis) and the
number of differentially expressed genes (y-axis, Figure 4B). We observed no
significant correlation between these two variables for plants at days 15 and 21 (p-value
> 0.05, Figure 4B). We concluded that there is no correlation between the severity of
growth phenotype and the transcriptomic response to stress, as, e.g., the smallest
plants (salt+mannitol, SM) also had the fewest number of differentially expressed
genes.

The different single and combined stresses are likely to elicit similar and unique
gene expression responses, resulting in the stresses having similar sets of DEGs. To
elucidate these similarities, we used the UpSet plot, which shows the intersection of
multiple sets for up-regulated (Figure 4C) and down-regulated (Figure 4D) genes.
Interestingly, the largest set of up- and down-regulated genes was unique to
heat+darkness combined stress (HD), suggesting that HD elicits the most unique and
dramatic transcriptional response among the tested stresses. Other unique stress
responses comprised upregulated cold+darkness (CD), cold+high light (CL),
cold+nitrogen deficiency (CN) and heat+nitrogen deficiency (HN, Figure 4C); and
downregulated cold+darkness (CD), cold+nitrogen deficiency (CN), cold (C) and
cold+high light (CL). Darkness alone (D) and in combination with other stresses (e.g.,
ND, CD, MD, SD, HD) also contained a high number of up-regulated (connected dots in
column 4,5, Figure 4C) and down-regulated (column 2,4, Figure 4D) genes, suggesting
a conserved, core darkness response. Similarly, we observed core responses to heat
(e.g., 6th column, Figure 4C) and cold (10th column, Figure 4C). Interestingly, we also
observed DEGs across heat and darkness experiments (Figure 4C, D), suggesting that
these two stresses elicit a similar response to a degree.

Figure 4. The number of differentially expressed genes in the tested stress combinations. A) The number of significantly (adjusted p-value < 0.05 against two controls) upregulated (red) and downregulated (blue) differentially expressed genes. B) The number of DEGs (x-axis) versus the size (y-axis) of Marchantia plants at day 15 (blue) and day 21 (orange). The R^2 and p-values are shown in the legend. Upset plot showing top 50 intersections of C) upregulated genes and D) downregulated genes. The bar charts on the top show the number of DEGs in the intersections, while the bar charts on the left show the number of DEGs in a given stress combination. The back dots indicate a given stress combination, while vertically connected dots indicate which

stress combination DEGs comprise a given intersection. For example, the 4th column in
C) indicates that CD, ND, D, MD, SD, and HD stresses share ~200 upregulated DEGs.

To better understand how Marchantia responds to a combination of two different
stresses, we compared the combined response (AB) to the response to individual
stresses (A and B), with three different metrics measuring the shared response,
dominance of a stress and novel responses induced by combined stress. We first
produced Venn diagrams for upregulated (Figure S6) and downregulated (Figure S7)
gene sets. The first metric measures the similarity between A and B (Figure 5A, green
area, Jaccard Index) and ranges from 0 (A, B do not have any DEGs in common) to 1
(A, B elicit identical DEGs). The second metric measures whether one stress
suppresses the other (Figure 5B, the difference between red and blue area) and ranges
from -1 (AB is similar to A but not to B, i.e., A suppresses B) to 1 (AB is similar to B but
not A, i.e., B suppresses A). The third metric measures whether a combination of two
stresses elicits a different, novel response when compared to the two individual
stresses (Figure 5C, purple area specific to AB) and ranges from 0 (all DEGs are found
in A and B, i.e., no novel response) to 1 (all DEGs in AB are unique, i.e., the
combination of AB elicited a unique response).
Stresses showing the highest similarity in terms of DEG responses comprise salt
and mannitol for up- and downregulated genes (Figure 5D-E, dark area for SM), salt
and nitrogen deficiency for upregulated genes (Figure 5D), and heat and darkness for
downregulated genes (dark area for HD, Figure 5E). The suppression analysis showed
that darkness is a strong suppressor for many stresses (red boxes in CD, MD, ND, SD),
except HD, indicating that heat stress and darkness are comparably dominant (Figure
5D-E). To support this observation, heat stress could also suppress other stresses (blue
boxes in HS, HN, HM, Figure 5D-E). Finally, the novelty analysis revealed that the
salt+mannitol combination elicited DEGs that were not found in the individual stresses
(dark blue boxes for SM, Figure 5D-E), suggesting that the two stresses can activate
new responses when combined.
To better understand the hierarchies of the stresses and how these stresses
affect the activity of biological processes, we identified which processes contain
significantly more upregulated or downregulated DEGs than expected by chance for
each stress combination (Figure 5F). The clustering of the rows (biological processes)
and columns (stresses) revealed that darkness-containing stresses form a clear group
of similar responses (six stress combinations comprising D, HD, CD, MD, SD and ND),
confirming the previous observation of the darkness suppressing other stresses.
Interestingly, the darkness caused a dramatic decrease in gene expression of
numerous pathways (Figure 5F, blue squares). The next largest group contained nearly
all heat stress combinations (four stresses H, HD, HS, HN) and a similar but less
dramatic decrease in the activity of many biological processes. Interestingly, despite the
dramatic downregulation of biological processes in most dark and heat stresses, a

subset of stresses (H, HS, HD, MD) was significantly upregulated for uncharacterized
genes (bin not annotated), suggesting that these responses employ poorly understood
genes. The other groups comprised high light (four stresses L, NL, LS, ML), nitrogen
deficiency (three stresses N, MN, and SN), and cold (three stresses C, CM, and CS). In
contrast, salt- and mannitol-containing stresses did not form any groups, suggesting
that despite dramatic phenotypic changes, these stresses are suppressed by other
stresses (Figure 5F, M and S are not grouped with other stresses). Interestingly,
salt+mannitol, cold+high light, and cold+low nitrogen were also not grouped, indicating
that these combinations result in novel transcriptomic responses.
Taken together, we can rank the strength of dominance of abiotic stresses as
darkness (six stress combinations), heat and light (four each), nitrogen deficiency and
cold (three each), salt and mannitol (none).

Figure 5. Comparison of differentially expressed genes and biological pathways in single and combined stresses. Illustration and equation of metrics used to measure A) similarity between two stresses, B) suppression of one stress when two stresses are combined, and C) novel genes that are differentially regulated when stresses are combined. Interaction between independent and cross stresses for D) upregulated genes and E) downregulated genes. In each stress combination, the first ('A') and second ('B') stress is the first and second letter of the combined ('AB') stress, respectively. For the suppression measure, a darker shade of red and blue indicates that more genes from the first ('A') and second ('B') stress are not represented in the combined stress, respectively. F) Heatmap of biological pathways (captured by second-level Mapman bins) that are significantly differentially expressed (hypergeometric test, adjusted p-value < 0.05). For brevity, we only show Mapman bins that are differentially expressed in at least three stress perturbations. Arbitrary clades of stress perturbations are highlighted by the different colours red, grey, green, orange, and blue squares at the tip of the dendrogram, and the leaf labels are also coloured. Abbreviations used to describe the categories of regulation are upregulation ('U', red), up and downregulation ('UD', purple), downregulation ('D', blue), and no change ('N', gray).

Comparative analysis reveals the conservation of diurnal gene expression between Marchantia and other Archaeplastida

Most members of the Archaeplastida kingdom subject more than one-third of their genes to diurnal gene expression, underlining the importance of diurnal regulation[43]. To study diurnal gene expression in Marchantia, the plants were grown in a 12 hour light / 12 dark cycle, and six timepoints were collected. The timepoints are 2, 6, 8 hours after light on and 2, 6, 8 hours after the light was switched off, corresponding to zeitgeber timepoints ZT2, 4, 6, 8, 12, 16, respectively.

The resulting 18 RNA-seq samples (six timepoints in triplicates) revealed the expected circular PCA plot (Figure S8A), and the gene expression from the individual samples showed an expected phasing pattern (Figure 6A). We identified 51.1% genes with significantly rhythmic gene expression (adjusted p-value < 0.05) with Jonckheere-Terpstra-Kendall (JTK_Cycle) algorithm[42], which revealed the time of the day where each rhythmic gene peaked (phase, Table S15). We also observed an expected circular pattern of the PCA plot of the genes (Figure S8B). The distribution of phases showed two peaks at ZT10 (i.e., 10 hours after lights on) and ZT20 (i.e., 4 hours before lights on, Figure 6B), which was reminiscent of phase distribution in Arabidopsis and rice[43]. Our previous study indicated that cell division genes are peaking just before the light / dark change in single cellular algae *Cyanophora paradoxa*, *Porphyridium purpureum*, *Chlamydomonas reinhardtii* and simple multicellular algae *Klebsormidium nitens*[43], showing that the diurnal rhythm regulates cell division in these algae. In contrast, Marchantia showed a more uniform expression of cell division markers (Figure 6C), indicating that, similarly to other land plants, Marchantia's cell division is not under the control of diurnal rhythm.

To elucidate how the different biological pathways are active during a typical day
of Marchantia, we visualized the phase distribution of rhythmic genes and their
pathways and performed a clustering analysis to reveal which pathways show similar
diurnal gene expression (Figure 6D). Not surprisingly, we observed that, e.g.,
photosynthesis genes peak in expression during the day (Figure 6D, photosynthesis is
labeled green during ZT 0-10) and shows a similar pattern with redox homeostasis,
which is important for scavenging harmful reactive oxygen species generated by light
capture [48]. Interestingly, and in line with the phase distribution plot (Figure 6A), we
observed that most biological pathways tend to peak at two times of the day: during the
day at ~ZT10 and the night at ~ZT20.

Our previous study showed that diurnal gene expression is conserved across >1
billion years of plant evolution[43]. To investigate whether Marchantia's diurnal gene
expression is also conserved, we compared the distribution of phases of Marchantia
genes and their orthologs in four algae (*Cyanophora paradoxa*, *Porphyridium*
purpureum, *Chlamydomonas reinhardtii,* and *Klebsormidium nitens*) and five land plants
(*Physcomitrella patens, Selaginella moellendorffii, Picea abies, Oryza sativa,* and
Arabidopsis thaliana). The analysis plots the phases of Marchantia genes (y-axis,
phases are arranged from 0 (top) to 23 (bottom)) versus the phases of the 1-to-1
orthologues from other species (x-axis, phases are arranged from 0 (left) to 23 (right)).
Two species with similar diurnal gene expression would show an accumulation of points
at similar x and y coordinates, indicating that the orthologs of the two species peak at
the same time of the day. Conversely, the points would be randomly distributed in the
plot for two species with different diurnal gene expressions. The analysis revealed that
for most species, with the exception of Selaginella and Picea, we observe groups of
genes at similar x and y coordinates, suggesting that the diurnal gene expression of
Marchantia is conserved (Figure 6E). To further quantify the conservation of diurnal
gene expression, we plotted the distribution of the differences between phase values of
one-to-one orthologues of Marchantia and the nine Archaeplastida. For all species,
except for Selaginella, Picea, and Arabidopsis, we observed a clear peak at around 0
(Figure 6F), confirming that the diurnal gene expression of Marchantia is conserved to
most Archaeplastida. Taken together, the conservation between Marchantia and the
other Archaeplastida makes Marchantia an attractive model to study diurnal gene
expression.

Figure 6. Comparison of diurnal rhythms of Marchantia and other Archaeplastida.
A) Z-score-normalised expression of genes (y-axis) taken at different zeitgeber timepoints (x-axis). Yellow and grey boxes indicate the presence and absence of light at the time of sampling, respectively. The ticks indicate the biological replicates. The data was collected in triplicates. B) Distribution of rhythmic genes (as %, y-axis) across phases (x-axis). C) Z-score-normalised expression of significantly rhythmic (adjusted p-value < 0.05) cell cycle genes (cyclin A, B, D, cyclin-dependent kinase (CDK), DNA primase, and DNA polymerase). The y-axis and x-axis indicate the gene expression and average expression at given ZT, respectively. D) Diurnal phase peak distribution of biological pathways. Rows represent biological pathways, while the columns indicate the phases at which the genes in a pathway peak. The colors of the cells indicate the % of rhythmic genes in a given pathway at each phase. The sum of each row is 100%. E) The heatmap shows the phase distribution of phases of Marchantia genes (y-axis, phases are arranged from 0:top to 23:bottom) versus the phases of the 1-to-1 orthologs from other species (x-axis, phases are arranged from 0:left to 23:right). The intensity of

the cells in the heatmap **is** proportional to the number of genes that peak at a given
phase for the two species. Black lines indicate the light to dark transition. Abbreviations
used to represent species are as follows: *Cyanophora paradoxa* (Cpa)*,*
Chlamydomonas reinhardtii (Cre)*, Porphyridium purpureum* (Ppu)*, Klebsormidium*
nitens (Kni)*, Physcomitrella patens* (Ppa)*, Selaginella moellendorffii* (Smo)*, Picea abies*
(Pab)*, Oryza sativa* (Osa) and *Arabidopsis thaliana* (Ath). F) Distribution of differences
of phases of the 1-to-1 orthologues between Marchantia and the nine members of
Archaeplastida. Value of 0 indicates that the 1-to-1 orthologs peak at the same time of
the day**; the** value of -10 and 10 shows that the orthologs peak 10 hours earlier and 10
hours later in other species than in *Marchantia*, respectively.
706
eFP browser and CoNekT database for Marchantia
Bioinformatic data is only as useful as its accessibility. To make our data readily
accessible, we have constructed an eFP browser instance for Marchantia available at
https://bar.utoronto.ca/efp_marchantia/cgi-bin/efpWeb.cgi [49], and updated our CoNekT
database [50], with our stress and diurnal data, available at https://conekt.sbs.ntu.edu.sg/.
To exemplify how our data and these databases can be used, we provide an example
with phenylpropanoids, which contribute to all aspects of plant responses towards biotic
and abiotic stimuli. In flowering plants, phenylpropanoids were found to be highly
responsive to light or mineral treatment, important for resistance towards pests[51], and
invasion of new habitats and reproduction[52]. The biosynthesis of phenylpropanoids
begins with phenylalanine ammonia lyases (PAL) and tyrosine ammonia lyases (TAL)
that catalyze the non-oxidative deamination of phenylalanine to trans-cinnamate, which
directs the output from the shikimate pathway phenylpropanoid metabolism[51].
To study phenylpropanoid metabolism in Marchantia, we started with entering
'PAL1' into CoNekT's search box (https://conekt.sbs.ntu.edu.sg/), which took us to the
page of PAL1, a PAL gene *AT2G37040* from *Arabidopsis thaliana*. To identify PAL
genes in Marchantia, we clicked on the link of the Land Plants orthogroup (CoNekT
provides orthogroups for Archaeplastida, Land Plants, and Seed Plants), which
revealed that all 11 land plants in the database contain PALs, while the algae
Cyanophora paradoxa and *Chlamydomonas reinhardtii* do not. Marchantia contains a
surprisingly large number (13) of PALs given the low redundancy of its genome, which
is higher than Arabidopsis (4). CoNekT contains gene trees that also show the
expression of genes in major organ types, and the analysis revealed that the many PAL
genes in Marchantia are likely a result of a recent duplication within Marchantia (Figure
S9).
To gain insight into the function of the 13 PAL genes from Marchantia, we set out
to study the expression of the PALs during stress conditions. We copied the gene
identifiers into Tools/Create heatmap (https://conekt.sbs.ntu.edu.sg/heatmap/),
revealing that most PALs show a high expression during cold treatment combined with

nitrogen starvation (Figure 7A). We clicked on one of the highly responsive genes
Mp4g14110.1, which took us to the page dedicated to the gene. The page contains
various information, such as gene description, CDS/protein sequences, gene families,
found protein domains, co-expression networks, and others. The detailed expression
profile confirmed the high cold + nitrogen-specific expression of *Mp4g14110* by using
CoNekT expression profile plot (Figure 7B) and eFP viewer (Figure 7C, red square at
intersection of cold and nitrogen deficiency). To better understand the function of
Mp4g14110.1, we clicked on the cluster link that directed us to the co-expression cluster
link of *Mp4g14110.1*. The cluster page contains information about the genes found in
the cluster, Gene Ontology enrichment analysis, found protein domains, and other
functional information. Interestingly, the 'Similar Clusters' cluster page revealed similar
co-expression clusters in other species (similarity is based on ortholog membership and
defined by Jaccard Index), with another similar cluster in Marchantia. To study these
duplicated clusters, we clicked on the 'Compare' button, which revealed that the two
clusters contain several gene families involved in phenylpropanoid biosynthesis (yellow
rounded rectangles: PALs, brown rectangles: chalcone synthases), ABC and DTX
transporters implicated in metabolite transport across membranes (purple/gray/green
and red rectangles[53]), auresidin synthases that can hydroxylate or cyclize chalcones
(light blue rounded rectangles[54]) and glutathione transferases (red rounded square[55]).
Interestingly, both clusters contained WRKY transcription factors (salmon rectangles),
implicating these transcription factors in controlling the biosynthesis of the respective
metabolites.

Taken together, our tools revealed evidence of duplicated modules, likely
involved in the biosynthesis of related phenylpropanoids. The updated CoNekT platform
contains many additional tools to predict gene function and find relevant genes [40,50,56].
For example, the tool found many of the described genes by identifying genes highly
expressed during combined cold and nitrogen starvation (Table S16), by clicking on
Tools/Search Specific Profiles, selecting Marchantia and 'Cold-Nitrogen deficiency'.

Figure 7. Implementation of the Marchantia gene expression data in CoNekT and eFP browser. A) Heatmap showing expression of PALs in representative stresses. B) Expression profile of Mp4g14110.1 (*MpPAL7*) under single and combined stress in CoNekt. C) Expression of *MpPAL7* in different organs and under different abiotic stress conditions in eFP. D) Comparison of Cluster 102 and 50 bounded by green and blue box, respectively. The border of node MpPAL7 is coloured red and blue dashed lines indicate homology between genes from the two clusters.

772
Discussion
To study how plants perceive changes in their environment and how conserved their
responses are, we constructed a diurnal and stress expression atlas capturing gene
expression changes to single and combined stresses for *Marchantia polymorpha*. With
this data in hand, we set out to investigate how conserved the responses to the
environmental changes are between model plants and which genes and pathways are
responding to single or combined stress.
To determine the response of *M. polymorpha* towards various stresses, we
tested a range of severity for heat, cold, salt, osmotic, light, dark, and nitrogen
deficiency on *M. polymorpha* gemmae. As expected in most stresses, the size of the
plants decreased proportionally to the severity of stresses   (Figure 1B). Certain
stresses, such as $3^{o}C$ cold and 535 $uEm^{-2}s^{-1}$ light, caused only minor growth
phenotypes, suggesting that *Marchantia* can survive under even lower temperatures
and higher light intensities. Conversely, stresses such as osmotic (100mM mannitol),
salt (40mM NaCl), and carbon starvation (3 days of darkness) produce strong growth
phenotypes resulting in reduced growth (osmotic, salt, darkness) and discoloration
(yellow-green for mannitol > 150mM, darker plants for salt > 60mM salt) (Figure 1B).
Death of plants occurred for heat stress at $36^{o}C$ for 24 hours and at higher salt
concentrations (>200mM). Interestingly, the plants grew at 0mM KNO_3, albeit slower
and with reddish discoloration likely caused by auronidin, a flavonoid shown to
accumulate during phosphate deficiency[44].
In most cases, the combination of stresses resulted in additive phenotypes (e.g.,
salt+mannitol stress results in smaller plants than the two stresses separately, Figure
1C-E). While heat ($33^{o}C$) and high light intensity (435 $uEm^{-2}s^{-1}$) resulted in death, this
was caused by temperature built up in the sealed plates, causing the temperature to
rise to the lethal $38^{o}C$. Consequently, this stress combination should be performed
using an open plate setup in the future. The only exception to this was salt+nitrogen
deficiency (SN) (40 mM NaCl, 0 mM KNO_3), which was significantly larger than plants
exposed only to salt stress and was observed to have extensive and dense rhizoids
(Supplemental Data 1). Curiously, we did not observe any dramatic phenotypes when
combining carbon/energy starvation (3 days of darkness) with any other stresses
(Figure 1C). This is counterintuitive as, e.g., heat stress acclimation is a costly process
requiring the biosynthesis of new transcripts and proteins to repair, replace and
rebalance the affected cellular machinery[57]. A hypothesis for the unexpectedly mild
phenotype for combined dark stress could be due to the lack of photosynthetic
processes in the absence of light, which increases the plant's capacity to cope with
increased ROS, a common response to most abiotic stresses[58]. This is also exemplified
by the significant downregulation of oxidoreductases and chloroplast redox homeostasis
in combined dark stress (Figure 5F). These phenotypes demonstrate that Marchantia is

able to survive various adverse growth conditions and can serve as an excellent model
to study stress acclimation.

Worryingly, we observed no evidence for conservation of global stress
acclimation responses between any of the five species of algae (*Cyanophora* and
Chlamydomonas) and land plants (*Marchantia*, rice and *Arabidopsis)*, when comparing
the responses using biological pathways and gene families (Figure 2E, Figure S5,
respectively). While surprising, this is not entirely unexpected as massive changes such
as genome and gene duplications have occurred over time, rendering a lack of
homology in genes across species and differences in gene families and regulation. This
suggests that each of the model plants uses a different strategy to acclimatize to stress,
making it uncertain to what degree knowledge gained from model species such as
Arabidopsis can be used to improve our crops. This lack of conservation of responses
to abiotic stresses has been observed by us at the species level in *Cyanophora*[20], and
at the intraspecies level by a salt stress study in six Lotus accessions, where only 1% of
genes showed a conserved response[59], in seven *Arabidopsis* accessions, which
showed a divergent response to treatment by salicylic acid[60] and by two strawberry
cultivars, which displayed modest conservation of DEGs to the same pathogen[61].
However, since mRNA levels are only capturing one layer of gene regulation, the lack of
conservation of responses to the abiotic stresses should be further investigated with
other omic technologies measuring, e.g., protein levels, translation rate, transcription
rate, and others.

Our analysis revealed that gene families comprising various enzymes (MapMan
bin Enzyme classification), mRNA expression control (RNA biosynthesis), protein
degradation, nutrient uptake, various membrane channels (Solute transport), cell wall
and photosynthesis are among some of the most responsive to up to seven stresses
(Figure 3A). Conversely, more basal processes, such as protein
biosynthesis/translocation, polyamine/amino acid/nucleotide metabolism, DNA damage
response/chromatin organization/cell cycle, were markedly less stress-responsive
(Figure 3A). These results indicate that *Marchantia* adjusts the physiology, energy, and
nutrient uptake via various gene expression regulatory mechanisms to ensure the
homeostasis of basic biological functions. Since younger gene families show a higher
degree of organ-specific gene expression[40], we speculated that younger gene families
could have evolved to provide adaptations to specialized environments and their abiotic
stresses. However, we observed similar stress responsiveness patterns for the oldest
gene families (Archaeplastida, Figure 3B, red line) and the much younger *Marchantia*-
specific gene families (Mpo, green line), suggesting that younger gene families are not
more active in stress responses, which is in line with our previous observation in
Cyanophora[20].

Since plants are often exposed to stress combinations in nature and our
understanding of how plants respond to these combinations is lacking[29,62], we have

exposed *Marchantia* to 19 possible combinations of the seven abiotic stresses
(darkness+high light and cold+heat being mutually exclusive). Interestingly, purely
physical stresses (cold, heat, darkness, and high light) showed a higher number of
DEGs than chemical stresses (salt, mannitol, and nitrogen deficiency) (Figure 4A). We
speculate that the physical stresses cause more DEGs because these stresses are able
to permeate every cell and affect every protein (heat, darkness) and/or have a dramatic
effect on the energy levels that have consequences on all processes (darkness, high
light). Conversely, stresses such as mannitol and salt can be effectively contained by
the action of ion transporters and osmolyte accumulation[63]. However, we did not
observe any correlation between the number of DEGs and the effect on plant growth
(Figure 4B); while salt and mannitol treatments caused the most dramatic growth
defects, these two stresses also showed the lowest number of DEGs (Figure 4B). This
observation is in line with our study in alga *Cyanophora*[20], suggesting that this
phenomenon is likely to be found in other organisms.
We observed a complex interplay of the stress combinations, where, e.g., the
combination of heat and prolonged darkness caused a unique upregulation of ~500
DEGs (Figure 4C, first column), but we also observed that certain stresses produce a
core set of DEGs present in any stress combination (e.g., Figure 4C, fourth column of
darkness combinations). Further analysis revealed that darkness-containing stresses
show a similar pattern of DEGs, indicating that darkness is perceived as more severe
stress by the plant and is thus able to dominate the DEG responses over other stresses
(Figure 5D, E, dominance plots are red in darkness stresses). This dominance is
caused by darkness-containing stresses having a similar pattern of downregulated
biological pathways (Figure 5F, blue cells indicate downregulation). Taken together, the
analysis allowed us to rank the strength of dominance of abiotic stresses as darkness
(six stress combinations), heat and light (four combinations each), nitrogen deficiency,
and cold (three each).
Interestingly, while salt and mannitol were the least dominant stresses and at the
same time most similar in terms of DEGs (Figure 5D-E, SM is dark blue for similarity),
when combined, they produced the most novel response (Figure 5D, E, dark blue for
'novel interaction' for SM, Figure 5F, SM is an outgroup). Other novel responses
comprise cold+high light and cold+nitrogen deficiency (Figure 5F). The basis for the
novel responses is unclear, but the unique DEG response found in the combination of
cold and high light (Figure 5D, CL is dark blue; Figure 5F, CL is an outlier) and the
enrichment of hydrogen peroxide removal pathway (Figure 5F) suggests that certain
stress combinations can enhance one another. High light can lead to the production of
oxidative stress by ROS, and plants regulate the photosynthetic ROS production by
adjusting light-harvesting and electron consumption pathways[64]. However, since cold is
known to slow down most metabolic reactions, the electron consumption pathways are
likely unable to absorb the excess electrons, leading to ROS accumulation. Since ROS

can cause oxidative damage to a wide range of lipids, proteins, and nucleic acids[65] and
is itself a signalling molecule (for recent reviews, see [66–68]), this resulted in the many
novel response that we observed for cold+high light.

Since the diurnal cycle regulates a significant number of genes in most
Archaeplastida [43], we set to augment our stress expression atlas with diurnal gene
expression data. Similarly to other land plants and in contrast to algae [43], Marchantia's
cell division is not synchronized by the diurnal cycle (Figure 6C). Interestingly,
Marchantia seems to have two populations of genes that peak at the end of the day
(phase 10) and the end of the night (phase 20, Figure 6B), which is also reflected in the
expression patterns of numerous biological pathways (Figure 6D). We observed this
previously in *Arabidopsis thaliana* and to lesser extent in rice, but not in *Physcomitrium*
patens, Selaginella moellendorffii or *Picea abie*s, suggesting that this pattern is not
dictated by the evolutionary distance. However, we observed that the diurnal gene
expression is conserved between most members of Archaeplastida (Figure 6F),
indicating that *Marchantia*, with its simplified regulatory network, is an excellent model to
study diurnal gene expression.

To make our data more readily accessible, we provide an eFP browser for
Marchantia, a popular tool allowing the visualization of gene expression by an
'electronic Fluorescent Pictograph'. Furthermore, we provide an updated CoNekT
database with expression atlases of 13 species comprising various algae and land
plants. The database provides tools to study gene expression, functional enrichment
analyses of co-expression networks, and other comparative tools. These valuable tools
will help further dissect the gene regulatory networks behind abiotic stress responses in
Marchantia and other species Figure 7).

Through our analyses, we have demonstrated the similarity of *Marchantia* to
other members of the Archaeplastida in terms of the responsiveness of gene families to
stress, the dominance, and the novel response of *Marchantia* to combined stress, and
the conserved diurnal expression of genes. We have validated the observations that
when presented with more than one stress, plants have a more dominant response to
the more severe stressor and that novel responses are adopted when plants are
exposed to combined stress. This strengthens the call for more emphasis on the study
of combined biotic and abiotic stresses in light of future challenges posed by climate
change[33,62].

Regrettably, while our study suggests a lack of conservation in gene expression
across stress and species, this might be convoluted by the difference in sample type
(shoot, leaves, and whole organisms), the analysis of few representative species, and
the sheer complexity of how plants respond to stress at the epigenetic (methylation of
genes), transcriptomic (mRNA, microRNA, lncRNA) and proteomic (posttranslational
modifications and activation) levels[69–73]. In addition, the broad level comparison in this
study may not have been sufficient to sift out the similarities between basal and higher

plants due to the increased complexity of plants due to gene and genome duplications.
Conservation of stress response in plants has been observed in other studies which
looked at transcription factors and hormones[74–79]. Taking the various limitations and
considerations outlined above, future studies of abiotic stress in plants should involve a
multi-omics approach with the inclusion of more species for us to delve into the
intricacies of the adaptations plants have developed over millions of years to respond to
stress.

Acknowledgments
We want to thank the Nanyang Technological University start-up grant for funding. We
want to thank the members of the Mutwil lab for their comments on the manuscript. We
thank D. Maizels (http://www.scientific-art.com/) for the illustrations used for the eFP
browser.

Author contributions
M.M. conceived the project, wrote the manuscript, Q.W.T. performed the single stress
experiments, data analysis, updated the CoNekT database and wrote the manuscript,
P.K.L performed the combined stress experiments, N.P. and A.P. provided the eFP
browser, Z.C., provided the Marchantia cultures and feedback on growing Marchantia.

Other competing interests
The authors declare no competing interests.

References

1. Taylor, L. L., Banwart, S. A., Valdes, P. J., Leake, J. R. & Beerling, D. J. Evaluating

the effects of terrestrial ecosystems, climate and carbon dioxide on weathering over

geological time: a global-scale process-based approach. *Philos Trans R Soc Lond B*

Biol Sci **367**, 565–582 (2012).

2. Wickett, N. J. *et al.* Phylotranscriptomic analysis of the origin and early

diversification of land plants. *Proc Natl Acad Sci U S A* **111**, E4859-4868 (2014).

3. Graham, L. E., Cook, M. E. & Busse, J. S. The origin of plants: body plan changes

contributing to a major evolutionary radiation. *Proc Natl Acad Sci U S A* **97**, 4535–

4540 (2000).

4. Bowman, J. L. Walkabout on the long branches of plant evolution. *Curr Opin Plant*

Biol **16**, 70–77 (2013).

5. Rubinstein, C. V., Gerrienne, P., de la Puente, G. S., Astini, R. A. & Steemans, P.

Early Middle Ordovician evidence for land plants in Argentina (eastern Gondwana).

New Phytol **188**, 365–369 (2010).

6. Wellman, C. H., Osterloff, P. L. & Mohiuddin, U. Fragments of the earliest land

plants. *Nature* **425**, 282–285 (2003).

7. Wheeler, J. A. Molecular Phylogenetic Reconstructions of the Marchantioid

Liverwort Radiation. *bryo* **103**, 314–333 (2000).

8. Bowman, J. L. *et al.* Insights into Land Plant Evolution Garnered from the

Marchantia polymorpha Genome. *Cell* **171**, 287-304.e15 (2017).

9. Ishizaki, K., Nishihama, R., Yamato, K. T. & Kohchi, T. Molecular Genetic Tools and

Techniques for Marchantia polymorpha Research. *Plant Cell Physiol* **57**, 262–270

(2016).

10. Cesarino, I. *et al.* Plant science's next top models. *Ann Bot* **126**, 1–23 (2020).

11. Lampugnani, E. R. *et al.* Cellulose Synthesis – Central Components and Their

Evolutionary Relationships. *Trends in Plant Science* **24**, 402–412 (2019).

12. Shimamura, M. Marchantia polymorpha: Taxonomy, Phylogeny and Morphology of a

Model System. *Plant Cell Physiol* **57**, 230–256 (2016).

13. Eklund, D. M. *et al.* Auxin Produced by the Indole-3-Pyruvic Acid Pathway

Regulates Development and Gemmae Dormancy in the Liverwort Marchantia

polymorpha. *Plant Cell* **27**, 1650–1669 (2015).

14. Tanaka, D., Ishizaki, K., Kohchi, T. & Yamato, K. T. Cryopreservation of Gemmae

from the Liverwort Marchantia polymorpha L. *Plant and Cell Physiology* **57**, 300–306

(2016).

15. Togawa, T. *et al.* Cryopreservation of Marchantia polymorpha spermatozoa. *J Plant*

Res **131**, 1047–1054 (2018).

16. Borghi, M., Perez de Souza, L., Yoshida, T. & Fernie, A. R. Flowers and climate

change: a metabolic perspective. *New Phytol* **224**, 1425–1441 (2019).

17. Bailey-Serres, J., Parker, J. E., Ainsworth, E. A., Oldroyd, G. E. D. & Schroeder, J. I.

Genetic strategies for improving crop yields. *Nature* **575**, 109–118 (2019).

18. Walters, D., Walsh, D., Newton, A. & Lyon, G. Induced resistance for plant disease

control: maximizing the efficacy of resistance elicitors. *Phytopathology* **95**, 1368–

1373 (2005).

19. Abdelrahman, M., Jogaiah, S., Burritt, D. J. & Tran, L.-S. P. Legume genetic

resources and transcriptome dynamics under abiotic stress conditions. *Plant, Cell &*

Environment **41**, 1972–1983 (2018).

20. Ferrari, C. & Mutwil, M. Gene expression analysis of Cyanophora paradoxa reveals

conserved abiotic stress responses between basal algae and flowering plants. *New*

Phytologist (2019) doi:10.1111/nph.16257.

21. Zandalinas, S. I. *et al.* Systemic signaling during abiotic stress combination in plants.

PNAS **117**, 13810–13820 (2020).

22. Thoen, M. P. M. *et al.* Genetic architecture of plant stress resistance: multi-trait

genome-wide association mapping. *New Phytol* **213**, 1346–1362 (2017).

23. Cairns, J. E. *et al.* Identification of Drought, Heat, and Combined Drought and Heat

Tolerant Donors in Maize. *Crop Science* **53**, 1335–1346 (2013).

24. Davila Olivas, N. H. *et al.* Genome-wide association analysis reveals distinct genetic

architectures for single and combined stress responses in Arabidopsis thaliana. *New*

Phytologist **213**, 838–851 (2017).

25. Nilson, S. E. & Assmann, S. M. The control of transpiration. Insights from

Arabidopsis. *Plant Physiol* **143**, 19–27 (2007).

26. Sun, Z., Jin, X., Albert, R. & Assmann, S. M. Multi-level Modeling of Light-Induced

Stomatal Opening Offers New Insights into Its Regulation by Drought. *PLOS*

Computational Biology **10**, e1003930 (2014).

27. Zandalinas, S. I. *et al.* ABA is required for the accumulation of APX1 and MBF1c

during a combination of water deficit and heat stress. *J Exp Bot* **67**, 5381–5390

(2016).

28. Zhou, R. *et al.* Screening and validation of tomato genotypes under heat stress

using Fv/Fm to reveal the physiological mechanism of heat tolerance. *Environmental*

and Experimental Botany **118**, 1–11 (2015).

29. Rivero, R. M., Mittler, R., Blumwald, E. & Zandalinas, S. I. Developing climate-

resilient crops: improving plant tolerance to stress combination. *Plant J* (2021)

doi:10.1111/tpj.15483.

30. Jiang, J. & Dehesh, K. Plastidial retrograde modulation of light and hormonal

signaling: an odyssey. *New Phytologist* **230**, 931–937 (2021).

31. Martínez-Ruiz, A. *et al.* Specificity in S-Nitrosylation: A Short-Range Mechanism for

NO Signaling? *Antioxidants & Redox Signaling* **19**, 1220–1235 (2013).

32. Stulemeijer, I. J. E. & Joosten, M. H. a. J. Post-translational modification of host

proteins in pathogen-triggered defence signalling in plants. *Molecular Plant*

Pathology **9**, 545–560 (2008).

33. Atkinson, N. J., Jain, R. & Urwin, P. E. The Response of Plants to Simultaneous

Biotic and Abiotic Stress. in *Combined Stresses in Plants: Physiological, Molecular,*

and Biochemical Aspects (ed. Mahalingam, R.) 181–201 (Springer International

Publishing, 2015). doi:10.1007/978-3-319-07899-1_9.

34. Pandey, P., Ramegowda, V. & Senthil-Kumar, M. Shared and unique responses of

plants to multiple individual stresses and stress combinations: physiological and

molecular mechanisms. *Frontiers in Plant Science* **6**, 723 (2015).

35. Gamborg, O. L., Miller, R. A. & Ojima, K. Nutrient requirements of suspension

cultures of soybean root cells. *Experimental Cell Research* **50**, 151–158 (1968).

36. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and

dispersion for RNA-seq data with DESeq2. *Genome Biology* **15**, (2014).

37. Schwacke, R. *et al.* MapMan4: A Refined Protein Classification and Annotation

Framework Applicable to Multi-Omics Data Analysis. *Molecular Plant* **12**, 879–892

(2019).

38. Jin, J. *et al.* PlantTFDB 4.0: Toward a central hub for transcription factors and

regulatory interactions in plants. *Nucleic Acids Research* **45**, D1040–D1045 (2017).

39. Zheng, Y. *et al.* iTAK: A Program for Genome-wide Prediction and Classification of

Plant Transcription Factors, Transcriptional Regulators, and Protein Kinases. *Mol*

Plant **9**, 1667–1670 (2016).

40. Julca, I. *et al.* Comparative transcriptomic analysis reveals conserved programmes

underpinning organogenesis and reproduction in land plants. *Nat Plants* **7**, 1143–

1159 (2021).

41. Benjamini, Y. & Hochberg, Y. Controlling the False Discovery Rate: A Practical and

Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society:*

Series B (Methodological) **57**, 289–300 (1995).

42. Hughes, M. E., Hogenesch, J. B. & Kornacker, K. JTK-CYCLE: An efficient

nonparametric algorithm for detecting rhythmic components in genome-scale data

sets. *Journal of Biological Rhythms* **25**, 372–380 (2010).

43. Ferrari, C. *et al.* Kingdom-wide comparison reveals the evolution of diurnal gene

expression in Archaeplastida. *Nature Communications* **10**, (2019).

44. Rico-Reséndiz, F. *et al.* Transcriptional and Morpho-Physiological Responses of

Marchantia polymorpha upon Phosphate Starvation. *International Journal of*

Molecular Sciences **21**, 8354 (2020).

45. Buti, M. *et al.* Transcriptome profiling of short-term response to chilling stress in

tolerant and sensitive Oryza sativa ssp. Japonica seedlings. *Funct Integr Genomics*

18, 627–644 (2018).

46. Huang, L. *et al.* Comparative transcriptome sequencing of tolerant rice introgression

line and its parents in response to drought stress. *BMC Genomics* **15**, 1026 (2014).

47. Domazet-Lošo, T., Brajković, J. & Tautz, D. A phylostratigraphy approach to

uncover the genomic history of major adaptations in metazoan lineages. *Trends in*

Genetics **23**, 533–539 (2007).

48. Das, P., Nutan, K. K., Singla-Pareek, S. L. & Pareek, A. Oxidative environment and

redox homeostasis in plants: dissecting out significant contribution of major cellular

organelles. *Frontiers in Environmental Science* **2**, 70 (2015).

49. Winter, D. *et al.* An 'electronic fluorescent pictograph' Browser for exploring and

analyzing large-scale biological data sets. *PLoS ONE* **2**, (2007).

50. Proost, S. & Mutwil, M. CoNekT: An open-source framework for comparative

genomic and transcriptomic network analyses. *Nucleic Acids Research* **46**, W133–

W140 (2018).

51. Vogt, T. Phenylpropanoid Biosynthesis. *Molecular Plant* **3**, 2–20 (2010).

52. Bais, H. P., Vepachedu, R., Gilroy, S., Callaway, R. M. & Vivanco, J. M. Allelopathy

and Exotic Plant Invasion: From Molecules and Genes to Species Interactions.

Science **301**, 1377–1380 (2003).

53. Gräfe, K. & Schmitt, L. The ABC transporter G subfamily in Arabidopsis thaliana.

Journal of Experimental Botany **72**, 92–106 (2021).

54. Nakayama, T. *et al.* Specificity analysis and mechanism of aurone synthesis

catalyzed by aureusidin synthase, a polyphenol oxidase homolog responsible for

flower coloration. *FEBS Lett* **499**, 107–111 (2001).

55. Pégeot, H. *et al.* The poplar Phi class glutathione transferase: expression, activity

and structure of GSTF1. *Frontiers in Plant Science* **5**, 712 (2014).

56. Ferrari, C. *et al.* Expression Atlas of Selaginella moellendorffii Provides Insights into

the Evolution of Vasculature, Secondary Metabolism, and Roots. *Plant Cell* **32**, 853–

870 (2020).

57. Bokszczanin, K. *et al.* Perspectives on deciphering mechanisms underlying plant

heat stress response and thermotolerance. *Frontiers in Plant Science* **4**, 315 (2013).

58. You, J. & Chan, Z. ROS Regulation During Abiotic Stress Responses in Crop

Plants. *Front Plant Sci* **6**, 1092 (2015).

59. Sanchez, D. H. *et al.* Comparative functional genomics of salt stress in related

model and cultivated plants identifies and overcomes limitations to translational

genomics. *PLoS ONE* **6**, (2011).

60. Van Leeuwen, H. *et al.* Natural variation among Arabidopsis thaliana accessions for

transcriptome response to exogenous salicylic acid. *Plant Cell* **19**, 2099–2110

(2007).

61. Wang, F. *et al.* Comparative transcriptomics reveals differential gene expression

related to Colletotrichum gloeosporioides resistance in the octoploid strawberry.

Frontiers in Plant Science **8**, (2017).

62. Mahalingam, R. Consideration of Combined Stress: A Crucial Paradigm for

Improving Multiple Stress Tolerance in Plants. in *Combined Stresses in Plants:*

Physiological, Molecular, and Biochemical Aspects (ed. Mahalingam, R.) 1–25

(Springer International Publishing, 2015). doi:10.1007/978-3-319-07899-1_1.

63. Hanin, M., Ebel, C., Ngom, M., Laplaze, L. & Masmoudi, K. New Insights on Plant

Salt Tolerance Mechanisms and Their Potential Use for Breeding. *Frontiers in Plant*

Science **7**, 1787 (2016).

64. Lima-Melo, Y. *et al.* Photoinhibition of Photosystem I Provides Oxidative Protection

During Imbalanced Photosynthetic Electron Transport in Arabidopsis thaliana.

Frontiers in Plant Science **10**, 916 (2019).

65. McCord, J. M. The evolution of free radicals and oxidative stress. *Am J Med* **108**,

652–659 (2000).

66. Mittler, R. ROS Are Good. *Trends in Plant Science* **22**, 11–19 (2017).

67. Czarnocka, W. & Karpiński, S. Friend or foe? Reactive oxygen species production,

scavenging and signaling in plant response to environmental stresses. *Free Radical*

Biology and Medicine **122**, 4–20 (2018).

68. Mullineaux, P. M., Exposito-Rodriguez, M., Laissue, P. P. & Smirnoff, N. ROS-

dependent signalling pathways in plants and algae exposed to high light:

Comparisons with other eukaryotes. *Free Radical Biology and Medicine* **122**, 52–64

(2018).

69. Chang, Y.-N. *et al.* Epigenetic regulation in plant abiotic stress responses. *Journal of*

Integrative Plant Biology **62**, 563–580 (2020).

70. Mazzucotelli, E. *et al.* Abiotic stress response in plants: When post-transcriptional

and post-translational regulations control transcription. *Plant Science* **174**, 420–431

(2008).

71. Franke, K. R. *et al.* Analysis of Brachypodium miRNA targets: evidence for diverse

control during stress and conservation in bioenergy crops. *BMC Genomics* **19**, 547

(2018).

72. Barrero-Gil, J. & Salinas, J. Post-translational regulation of cold acclimation

response. *Plant Science* **205–206**, 48–54 (2013).

73. Schütze, K., Harter, K. & Chaban, C. Post-translational regulation of plant bZIP

factors. *Trends in Plant Science* **13**, 247–255 (2008).

74. Castro, P. H. *et al.* Phylogenetic analysis of F-bZIP transcription factors indicates

conservation of the zinc deficiency response across land plants. *Sci Rep* **7**, 3806

(2017).

75. Wu, T.-Y. *et al.* Evolutionarily conserved hierarchical gene regulatory networks for

plant salt stress response. *Nat. Plants* **7**, 787–799 (2021).

76. Balderas-Hernández, V. E., Alvarado-Rodríguez, M. & Fraire-Velázquez, S.

Conserved versatile master regulators in signalling pathways in response to stress

in plants. *AoB Plants* **5**, plt033 (2013).

77. Chen, W. J. & Zhu, T. Networks of transcription factors with roles in environmental

stress response. *Trends in Plant Science* **9**, 591–596 (2004).

78. Ju, C. *et al.* Conservation of ethylene as a plant hormone over 450 million years of

evolution. *Nature Plants* **1**, 1–7 (2015).

79. Lieberman-Lazarovich, M., Yahav, C., Israeli, A. & Efroni, I. Deep Conservation of

cis-Element Variants Regulating Plant Hormonal Responses. *The Plant Cell* **31**,

2559–2572 (2019).

1154

1155

1156   **Supplementary figures**

1157

1158   **Figure S1. Area of plants at 15 days (day of harvest) and day 21 (6 days post-**
1159   **harvest) for the single and combined stresses.** Each bar represents measurements
1160   from at least 4 plants. Error bars are represented by standard deviation and significance
1161   was determined by Student's two-tailed, t-test, p < 0.05.

1162
Figure S2. Hierarchical clustering of the Marchantia stress experiments. Transcript
per million (TPM) gene expression values were scaled with standard scaler and
clustered with seaborn.clustermap() function.

1166

Figure S3. Volcano plots of stress experiments. The x-axis represents $\log_2$-fold change, the y-axis indicates the $-\log_{10}$ of adjusted p-value. Each point represents a gene that has an absolute $\log_2$-fold change more than 1 and is significantly (adjusted p-value<0.05, orange color) or not significantly (p-value>0.05, blue point) differentially expressed.

1172

Figure S4. Comparison of found differentially expressed genes for the single and combined stressed. For each stress combination, we estimated significant DEGs (adjusted p-value<0.05) by using control D2 (red circle) and control H2 (green circle). The orange areas indicate the overlap between the found DEGs of the two controls. The overlapping DEGs were used for further study. The following abbreviations are used for the description of stresses: cold (C), dark (D), heat (H), light (L), mannitol (M), nitrogen deficiency (N) and salt (S). Combination of stresses is indicated by two letters, where e.g., HD indicates heat+darkness treatment.

1181

Figure S5. Clustering of stress/species similarities by using differentially expressed gene families. Gene families were classified according to the conservation of differential gene expression and similarities between the differential gene families were calculated by jaccard index. The color of each cell indicates the Jaccard Distance between any two stress/species. Species are colour-coded blue, light blue, red, yellow and green on the left column for *C. paradoxa, C. reinhardtii, M. polymorpha, O. sativa and A. thaliana* respectively while labels for heat, cold, salt, mannitol, light, dark, nitrogen and other stresses are coloured in red, blue, purple, magenta, yellow, black, green and grey respectively. The following abbreviations are used for the description of stresses: cold (C), dark (D), heat (H), light (L), mannitol (M), nitrogen deficiency (N), salt (S), iron deficiency (minFe), sulfur deficiency (minS) and phosphorus deficiency (minP).

1193

1194

Figure S6. Venn diagrams of upregulated genes in single and combined stresses.
The abbreviates are cold (C), dark (D), heat (H), light (L), mannitol (M), nitrogen deficiency (N) and salt (S). Combination of stresses is indicated by two letters and purple circles. The first and second stress in a pair is colored red and green (e.g., cold is red in CD), while the combined stress is colored purple. The sizes of the circles and intersections and the numbers within indicate the number of significant DEGs.

1201

Figure S7. Venn diagrams of downregulated genes in single and combined stresses. The abbreviates are cold (C), dark (D), heat (H), light (L), mannitol (M), nitrogen deficiency (N) and salt (S). Combination of stresses is indicated by two letters and purple circles. The first and second stress in a pair is colored red and green (e.g., cold is red in CD), while the combined stress is colored purple. The sizes of the circles and intersections and the numbers within indicate the number of significant DEGs.

1208

A

B

1209

1210 **Figure S8. Analysis of diurnal data.** A) Principal Component Analysis (PCA) of the
1211 diurnal gene expression data. Each point represents one RNA-seq sample. The
1212 different zeitgeber (ZT) time points are color coded. The data was scaled with standard
1213 scaler. B) PCA analysis of genes. The genes were color-coded by their lag value. Gray
1214 genes are not rhythmic.

Figure S9. Phylogenetic Tree of PAL OG_02_0000119 (Land Plants).

Supplementary Tables

Supplementary Table 1. Summary of experiment label to experiment conditions and mapping statistics from kallisto.

Supplementary Table 2. TPM normalised gene expression matrix for *Marchantia* experiments

Supplementary Table 3. Raw gene expression matrix for *Marchantia* experiments

Supplementary Table 4. PCC values within *Marchantia* experiment replicates

Supplementary Table 5. Details on source, sample and stress conditions of other species in interspecies comparison

Supplementary Table 6. Summary of DESeq2 output of *Marchantia* stress experiments

Supplementary Table 7. List of genes that are identified to be differentially expressed in both controls with corresponding mapman bins and annotation. 'UP' and 'DOWN' in L2FC_D2 and L2FC_H2 columns indicate significant up and downregulation when compared against control D2 and H2 respectively.

Supplementary Table 8. Summary of DESeq2 output of *Oryza sativa* stress experiments

Supplementary Table 9. List of genes that are identified to be differentially expressed in *Oryza sativa*.

Supplementary Table 10. Annotation of transcription factors for Cpa, Cre, Mpo, Osa and Ath

Supplementary Table 11. Annotation of kinases for Cpa, Cre, Mpo, Osa and Ath

Supplementary Table 12. Status of orthogroups in corresponding species and stress condition. 'UP', 'DOWN', 'AMB', 'NC' and 'None' indicates that all differentially expressed genes are upregulated, downregulated, up and downregulated, no differentially expressed genes and no genes in orthogroup respectively.

Supplementary Table 13. Summary of age classification of orthogroups, combined status of response to stress in various organisms for non-species specific orthogroups and response to stress in species-specific orthogroups.

Supplementary Table 14. Orthogroup file used for orthogroup comparisons, obtained from Julca et al., 2021

Supplementary Table 15. JTK_Cycle output of diurnal gene expression for *Marchantia*

Supplementary Table 16. Cold-Nitrogen deficiency specific genes as obtained from conekt.plant.tools