1

Generalized harmonic analysis reveals a frequency modulated timer regulates mammalian hibernation

Shingo Gibo[1], Yoshifumi Yamaguchi[2,3,4,]*, and Gen Kurosawa[1,]*

[1]RIKEN Interdisciplinary Theoretical and Mathematical Sciences Program (iTHEMS); Wako, 351-0198, Japan

[2]Institute of Low Temperature Science, Hokkaido University; Kita-19, Nishi-8, Kita-ku, Sapporo, 060-0819, Japan

[3]Global Station for Biosurfaces and Drug Discovery, Global Institution for Collaborative Research and Education (GI-CoRE), Hokkaido University; Kita-12, Nishi-6, Kita-ku, Sapporo 060-0812, Japan.

[4]Inamori Research Institute for Science Fellowship (InaRIS); 620 Suiginya-cho, Shimogyo-ku, Kyoto 600-8411, Japan.

*Corresponding authors. Emails: g.kurosawa@riken.jp; bunbun@lowtem.hokudai.ac.jp.

Abstract

Mammalian hibernators decrease basal metabolism and body temperature (Tb) to minimize energy expenditure in harsh seasons. During hibernation, Tb drops to low temperature (<10 °C) and remains constant for days, known as deep torpor. Spontaneous interbout arousals interrupt torpor bouts, when Tb recovers to euthermic state ~37 °C. Torpor-interbout arousal event repeats during hibernation. Little is known about mechanisms governing Tb fluctuation during hibernation. Here, we analyzed Tb fluctuation across Syrian hamsters' hibernation cycle using generalized harmonic analysis and discovered a model with frequency modulation quantitatively reproducing Tb fluctuation. This analysis identified that an unexpectedly longer period of 120–430 days modulates period of several days, generating Tb fluctuation. We propose that concerted action of two endogenous periods governs torpor-interbout arousal cycles during hibernation.

Introduction

Hibernation is a strategy for organisms to survive in the environment with limited food and water availability[1,2]. During a season with little or no food, small mammalian hibernators drastically decrease their basal metabolism and core body temperature (Tb) to 10 °C and become immobile. This hypometabolic, hypothermic, and immobile state is called deep torpor. Interbout arousal (IBA) interrupt deep torpor, during which Tb rapidly arise to euthermic state. Thus, during hibernation period, Tb does not remain constant at neither low nor high values but shows fluctuation between euthermia and hypothermia with an interval of several days. This multiday-scale-phenomena, known as torpor-IBA cycles, is a conserved property of hibernation across mammalian hibernators.

IBA is also called periodic arousal[3], for patterns of Tb fluctuation in the middle of hibernation season attempted one to find certain periodicity in torpor-IBA cycles. However, length of torpor bout gradually changes in the early and late period of hibernation even under a constant condition in a laboratory, casting questions whether the torpor-IBA cycle is explicitly defined as "periodic" and what biological processes are behind it. Several hypotheses have long been proposed to explain the regulation and significance of the torpor-IBA cycle[4]. There are two non-mutually exclusive hypotheses currently proposed. First is that the timing of IBA is regulated by accumulation or consumption of certain proteins, protein modifications, or metabolites during torpor period. And the second one is that its timing is a reflection of certain innate endogenous rhythm(s) in the animal, such as circadian or circannual rhythms[5]. In nature and human societies, some systems exhibit a gradual change in the period of oscillations[6]. For instance, timing of sleep onset for some non-24 h patients with sleep-wake disorder is delayed every day and fluctuates several times a month[7]. Theoretically, this phenomenon can be understood as the desynchrony between circadian rhythms and 24 h environmental cycles; however, it has not been tested whether such desynchronization is responsible for producing Tb patterns during hibernation. Now that recent technology development enables to monitor

Tb for more than hundred days with high precision, quantitative analysis and phenomenological modeling of Tb time-series data can be used to address the principle governing hibernation.

The frequency change in biological time series is often quantified by short-time Fourier transform (STFT) and Wavelet transform. In STFT, the time series, which is multiplied by short interval window function, is analyzed, instead of analyzing the original time series. In Wavelet transform, the basis function is localized in time and frequency, called Wavelet function, which is distinct from trigonometric function. These two methods have been widely accepted in the field of time-series analysis. However, it is often difficult for the two methods to estimate the period of the signal within short time intervals (such as analyzing the patterns of Tb fluctuation during hibernation that can be used to uncover certain periodicity in torpor-IBA cycles) because of the fundamental tradeoff between time and frequency resolutions. To circumvent this problem, Generalized Harmonic Analysis (GHA), a methodology usually applied to acoustics for characterization of irregularities in music or circadian rhythms, can be applied to Tb fluctuation during hibernation. In contrast to STSF and Wavelet transforms, GHA is advantageous because it simply fits the data by the summation of trigonometric functions based on least square, and there is no trade-off between time and frequency resolutions.

Mammalian hibernators are roughly classified into two types: (i) obligate (or strongly seasonal) and (ii) facultative (or opportunistic) types[1,8]. Obligate hibernators, such as ground squirrels, marmots, and bears, undergo fall transition and enter hibernation spontaneously even without environmental cues[9-11]. For example, 13-lined ground squirrels and chipmunks exhibit hibernation iteratively with a period of ~1 year under conditions of constant cold and continuous darkness[12,13]. These lines of evidence suggest that endogenous circannual rhythms underlie hibernation in these species, although mechanisms for the circannual rhythms are unknown. In contrast, facultative hibernators, such as Syrian hamsters (*Mesocricetus auratus*), enter hibernation at any time of year, at least in a laboratory condition, when they are exposed to cold and short photoperiodic conditions[14,15]. Consistent with the successful induction of hibernation irrespective of seasons in appropriate laboratory conditions, little evidence has been found so far that circannual control of hibernation exists in this species. Nevertheless, this species awakens from hibernation without any external factors[14,15], implying the existence of mechanisms for estimation of the hibernation length. However, few clues about this phenomenon exist, not only at molecular level but also the underlying mathematical principle in both obligate and facultative hibernators.

In this study, we performed theoretical analysis of Tb across hibernation cycle (>50 days) in Syrian hamsters[15]. Typically, the Tb time series during hibernation is noisy and fluctuates irregularly, which complicates analysis. Therefore, to uncover temporal changes of torpor-IBA cycle, we used GHA, a methodology usually employed for acoustics for characterization of irregularity in music.

Results

Generalized Harmonic Analysis of changes of dominant period behind torpor-IBA cycle

To find rules for Tb fluctuation during hibernation, we applied GHA to analyze Tb from 13

hibernating Syrian hamsters[15] (Fig. 1a, b, Supplementary Fig. 1). GHA enables us to accurately

quantify periodic components underlying several torpor-IBA cycles (Fig. 1c-f) and determine

the strongest periodic component (Fig. 1g, h), the so-called dominant frequency ("Methods"[16]).

Because the torpor-IBA cycles take several days (Fig. 1a-c), we extracted dominant frequency

of the torpor-IBA cycles ranging from 0.1 to 0.3 per day (Fig. 1e, f), which corresponds to 3.3–

10 days dominant period calculated as the inverse of the dominant frequency (Fig. 1g, h) to

reveal the temporal changes of periodic components during hibernation. The extracted

dominant periods underlying Tb time series were 3.5–8.5 days during the first 16 days after the

onset of first torpor bout and 4.5–8.0 days during 68–84 days. This GHA analysis quantitatively

revealed that dominant period of torpor-IBA cycle gradually changed (Fig. 1i, j). The dominant

period increases or decreases over time during hibernation depending on the individual

(Supplementary Fig. 2). To compare patterns of Tb time series, we quantified the change of

torpor-IBA cycle period at the initial 0–48 days of hibernation using linear regression

("Methods", Fig. 1k, l, Supplementary Fig. 2). For 12 of 13 individuals (except #3 in

Supplementary Fig. 2), the slope of regressed line was positive, and its average was 0.0087 per

day, which indicates that the period of torpor-IBA cycle increased 0.87% per day on average

with time during the initial 0–48 days (Fig. 1m). Particularly, for five of thirteen individuals,

who hibernated for more than 120 days (#1, 2, and #8–10), the dominant period of torpor-IBA

cycle initially increased and decreased later toward the end of hibernation (Fig. 1g-l,

Supplementary Fig. 2i, l-n, p, q). Although one of thirteen individuals (#3) exhibited strongly

fluctuated Tb pattern and its slope of the linear regression was negative, it was due to the

occurrence of shallow/daily torpor and long IBA during first 20 days of hibernation because

the dominant period of torpor-IBA cycle exhibited an increasing tendency thereafter

(Supplementary Fig. 1a, Supplementary Fig. 2a, d). Taken together, this analysis demonstrated

that the period of torpor-IBA cycle changes at hundreds-days scale, suggesting that it is

possibly governed by as-yet-unknown physiological temporal process.

In search for a model reproducing the pattern of torpor-arousal cycles

To understand biological processes behind gradual change of the torpor-IBA cycle, we tested

two theoretical models that possibly reproduce the experimental data, (a) frequency modulation

model (FM) and (b) desynchrony model (Fig. 2a, b). FM assumes that the shorter frequency

(period) may be modulated by another slower frequency (longer period). Its typical example is

FM radio exhibiting gradual change in period over time[17], although few biological processes

other than auditory and vocalization systems were proposed to exhibit such features. In contrast,

the desynchrony model assumes that one period, torpor-IBA cycle period in this case, changes

over time due to the failure of synchronization with another period. A typical example of this

is desynchronization of internal circadian rhythm and external photoperiod (Fig. 2a, b,
"Methods").

By varying the parameters in both models, we first investigated which model would
better reproduce the observed Tb fluctuation using maximum likelihood estimation. The FM
model with the best parameter set, yielding maximum likelihood, reproduced most of the
timing for the transitions between deep torpor and IBA for all individual data (Fig. 2c, d). In
contrast, the desynchrony model did not reproduce most transitions between deep torpor and
IBA for some individuals (Fig. 2g, h). Indeed, likelihood values of the FM model with the best
parameter set were always larger for each individual's data than those of the desynchrony
model, suggesting that the FM model is the more proper model for reproducing Tb fluctuation
of Syrian hamsters during hibernation (Fig. 2k, Supplementary Fig. 3, Supplementary Fig. 4 in
detail).

We next evaluated the agreement between theoretical models and Tb experimental
data by other statistical method, the so-called "Itakura–Saito (IS) divergence." IS divergence
may be a better statistical method to identify the model and its parameter set that reproduces
the timing of IBA than widely-used maximum likelihood estimation[18] from the following
points; IS divergence, originally a speech recognition method for mixed sound, has axial
asymmetry while the Euclidean distance has axial symmetry. Thus, IS divergence imposes
more penalties if the value of the model is smaller than that of experimental data[19]. Indeed, FM
model with the best parameter set, yielding minimum IS divergence, reproduced most of the
timing of IBAs (Fig. 2e, f, Supplementary Fig. 5). Quantification of the maximum likelihood
estimation using IS divergence demonstrated that the FM model is much better than the
desynchrony model in reproducing Tb fluctuation (Fig. 2l, for examples compare #5, #8, and
#12 in Supplementary Fig. 5). For an individual wherein the desynchrony model showed a
better score than the FM model (#3 in Fig. 2l), it was evident that FM model fits the Tb data
much better than desynchrony model for the most part in the time series (Fig. 2f, j), suggesting
that the judgment with IS divergence is inappropriate for the individual. Taken together, both
maximum likelihood estimation and IS divergence statistically justified that the FM model is
a proper model for reproducing periodic change of torpor-IBA cycles.

Quantifying rhythms that underlie hibernation
The above result indicates that the main property of Tb fluctuation during hibernation can be
described by the two key parameters in the FM model, shorter (day/ω_1) and longer (day/ ω_2)
period. In this model, shorter period is modulated by longer period ($\omega_1 > \omega_2$). By determining
the two periods with the theoretical model, we can now quantify and explain individual
variation in complex patterns of Tb fluctuation. Because the dominant period estimated by
GHA was 3.5–9 days and the longer period at the timescale of approximately 100–500 days,
we first rigorously varied the shorter period at the timescale of approximately 1–10 days to
precisely determine the two periods (see "Methods"). To this end, we compared estimations by

maximum likelihood and IS divergence as follows.

Based on the best maximum likelihood estimation values for the FM model that
reproduced experimental data (Fig. 3a-c, Supplementary Fig. 4), we determined the faster and
slower period (ω_1 and ω_2); the shorter period of the FM model, yielding the maximum
likelihood estimation, was between 4.0 and 9.2 days, which covers the dominant period
extracted by GHA (Fig. 3d). Meanwhile, the estimated longer period was between 119 and 430
days (Fig. 3d). Conversely, when the parameter set of the FM model was chosen by IS
divergence, the distribution of the shorter period of the FM model (day/ω_1) was narrower such
that it was between 3.9 and 6.5 days (Fig. 3e). The longer period by IS divergence (day/ ω_2)
was also between 115 and 430 days (Fig. 3e). Therefore, there was a discrepancy between the
maximum likelihood and IS divergence estimation. It was probably because the FM model
yielding the maximum likelihood estimation did not reproduce shallow/daily torpor of some
individuals, whereas the FM model yielding minimum IS divergence did reproduce them (see
#5 and #7 in Supplementary Fig. 3, Supplementary Fig. 5). Taken together, analysis with the
FM model enabled us to quantify a period of several days and another longer period of 115–
430 days that modulates the former period during hibernation.

Forecasting body temperature fluctuation during hibernation
The validity of the FM model for Tb fluctuation can be assessed by its predictability. To test
this, we estimated the two parameters (day/ω_1 and ω_2) from three-quarters (Fig. 4a, b) of the
whole time series of Tb data for each individual's data by the FM model using IS divergence
(Fig. 4, Supplementary Fig. 9, Supplementary Fig. 10) or maximum likelihood (Supplementary
Fig. 7, Supplementary Fig. 8). Simulation with the parameters predicted the last quarter of the
time series because the estimated shorter and longer period (day/ω_1 and day/ω_2) from the three-
quarters of the Tb time series were close to those estimated from the whole time series (Fig.
4a, b). When we simulated the FM model with the parameters estimated from the first half of
the whole time series of the Tb data, it also predicted the first few cycles of the second half of
the original whole Tb time series but gradually deviated from the original (Fig. 4c, v). Indeed,
the longer periods (day/ω_2) predicted from the first half were distant from those estimated from
the original whole Tb time series (Fig. 4e, g), implying that the estimation of the parameter
needs sufficiently long experimental Tb time series (Fig. 4e, g). In contrast, when the shorter
frequency (ω_1) was predicted from the first half of the whole time-series data, it ranged from
3.4 to 5.9 days and was close to that estimated from the whole data (Fig. 4e, f). These results
suggest that estimation of longer frequency (ω_2) is affected by and that of shorter frequency
(ω_1) and is robust to the length of Tb time series used for statistical analysis.

Discussion
In this study, we applied GHA to quantify the time series of Tb fluctuation during hibernation
and defined theoretical models that reproduce it. This study demonstrates the effectiveness of

GHA to quantify Tb fluctuation, the patterns of which are complex and rich in variation among
animals, and is an improvement over using STFT and Wavelet transform to explore and model
Tb oscillations.

Through statistical analysis of the FM model and desynchrony model, we conclude
that the FM model is better than the desynchrony model to realize the pattern of Tb fluctuation
during hibernation of Syrian hamsters. To the best of our knowledge, only a few studies have
applied FM to explain biological processes, such as circadian desynchronization in rats and
changes in oscillatory frequency of brain waves in visual perception[6,20]. The FM model
hypothesizes that the frequency of oscillation (ω_1) gradually changes over time with another
longer frequency (ω_2). Our statistical analysis derives realistic values for ω_1 and ω_2. The
derived ranges of both values in the case of analysis with IS divergence were 3.9–6.5 day period
for ω_1 and 115–430 day period for ω_2, well within the 2–3 year lifespan of Syrian hamsters.
This result raises the question of what biological processes are reflected in these values.

The faster frequency ω_1, a period with a few days, could be responsible for
determining the period of torpor-IBA cycles. In many small hibernators, length of each torpor
bout gradually increases from the beginning, reaches its peak in the middle, and then decreases
at the end of the hibernation season, even in constant ambient temperature conditions[13,21-24].
Ambient and core body temperatures also affect duration of each deep torpor bout, suggesting
involvement of a temperature-sensitive process in the regulation of torpor-IBA cycles[13,25-27].
Metabolic rate contributes to the determination of torpor-IBA timings to only a minor extent
(about 30%) in golden-mantled ground squirrels[26]. The exact process responsible for
determination of torpor-IBA timing is not yet clear. Most chemical reactions are temperature-
sensitive. It should be noted that the temperature-sensitive nature of torpor-IBA cycle is
apparently different from temperature-compensation of circadian rhythm[28,29]. In fact, several
studies suggested that clock genes involved in circadian rhythm have little or small contribution
to determining timing of deep torpor-IBA cycles during hibernation[30,31]. These lines of
evidence suggest that mechanisms other than circadian rhythm governed by transcription-
translation feedback loop underlying the timing of torpor onset and arousal during hibernation.
ω_1 may correspond to the period generated by such mechanisms. In contrast, timing of
shallow/daily torpor, an adaptive response to winter condition or food shortage, during which
basal metabolisms and Tb drops for a shorter period, typically several hours and within a day,
is related to circadian rhythm in some species, including Djungarian hamster (*Phodopus
sungorus*), which exhibits seasonal shallow/daily torpor, and mice entering fasting-induced
torpor[10,27,30-33]. Thus, the situation is still complex and further research is needed to determine
whether ω_1 is sensitive to ambient temperature during hibernation.

Another frequency ω_2, a period of a few hundred days, implies endogenous circannual
rhythm. Although we recognize the limitation of our study that our model does not integrate
the timing of the start and end of the hibernation period, it is surprising that only Tb time-series
data during hibernation period could derive such a long period. In some species including

mammalian hibernators and migratory birds, the circannual rhythm is regulated by unknown
endogenous mechanisms independent of environmental triggers, such as photoperiod and
ambient temperature[13,34]. Ground squirrels kept in a constant condition change their body
weight and exhibit torpor phenotypes in a circannual cycle[13,35,36]. Additionally, recordings over
several years in ground squirrels or eastern chipmunks kept under constant photoperiods and
cold temperatures demonstrated that the periods from the onset of hibernation to that in next
year range from 5 months to 16 months or 5 months to 13 months in those animals,
respectively[12,13,35,36]. Thus, the endogenous period underlying circannual phenomena could
vary among individuals, implying that coordination of the endogenous circannual period with
phenological changes would be necessary for proper adaptation to animal habitat. Although
Syrian hamsters are facultative hibernators that do not depend on or may not have an
endogenous circannual rhythm, they spontaneously quit hibernation after several months of
hibernation period under a constant cold and photoperiodic condition. This strongly suggests
that Syrian hamsters have an unknown endogenous timer for measuring the length of
hibernation. In fact, a phenomenon called refractoriness has long been known in Siberian and
Syrian hamsters[37-39]. The hamsters regress or regrow their gonads in response to chronic short
or long photoperiod, respectively. But if the animals were kept further in short or long
photoperiod conditions for several months after the gonadal regression, the gonads then start
to regrow or regress without any environmental changes, suggesting the existence of yet-
unidentified endogenous timer measuring seasonal length in hamsters. Thus, ω_2 may
correspond to possible endogenous period governed by such an endogenous timer.

We anticipate that our theoretical analysis of Tb fluctuation can be a starting point for
quantitative comparison of hibernation patterns across close and distantly related hibernating
species exhibiting various Tb patterns. Furthermore, quantification across two or more
consecutive hibernation seasons may allow prediction of the timing of torpor and IBA and will
also foster the understanding of molecular mechanism of hibernation by searching for
biological processes that operate within those periods.

Fig. 1. GHA analysis quantifies change of periods in Tb fluctuation during hibernation.

(**a**, **b**) Time series of Tb fluctuation during hibernation under a short day and cold

temperature (8L:16D cycle, ambient temperature = 4 °C). Two representative hibernating
Syrian hamsters (from n = 13) are shown in Fig. 1a, b (see also Supplementary Fig. 1 for data
of the others). (**c, d**) Scheme of GHA analysis for Tb data (see text). (**e, f**) Sequence of
estimated frequency by analysis of data from two representative individuals during
hibernation (#1, 2). The heatmaps show the magnitude of spectrogram as logarithmic
compression of power defined by log (1 + |amplitude|^2). (**g-i**) Estimated dominant period (i.e.,
day/frequency) for #1 (**g**), #2 (**h**), and all 13 individual hamsters (**i**) changed over time. (**j-l**)
Dominant periods normalized by the initial dominant period. (**k, l**) The change of dominant
period for individuals #1 and #2 at 0–48 days was quantified using liner regression. (**m**)
Distribution of quantified slopes of regression line per day for the change in normalized
dominant period.

Fig. 2. Determine a model reproducing the torpor-IBA cycles of Syrian hamster. (**a, b**)
Schematic representation of the proposed FM model (**a**) and desynchrony model (**b**) with
multiple frequencies of ω_1 and ω_2. (**c-i**) Comparison of simulation done by the two models.
(**c-f**) FM model simulation (red) with the best parameter set for Tb time series recorded in
two representative animals (black, #1, 3). The best parameter was chosen by maximum
likelihood estimation (**c, d**) and IS divergence (**e, f**). RMSE is the value of root mean squared
error. (**g-j**) Desynchrony model simulation (red) with the best parameter set for the Tb data
from same animals used for FM simulation (#1, 3). The best parameter was chosen by

maximum likelihood estimation (**g, h**) or IS divergence (**i, j**). (**k, l**) Maximum likelihood

estimation comparison of the two models to realize each individual time series (#1–13).

Likelihood of FM (black) and desynchrony model (gray) was compared using maximum

likelihood estimation (**k**) and IS divergence (**l**). Time series of fixed minimum of Tb was set

as the reference model. Note that as the model becomes closer to the experimental data,

maximum likelihood estimation becomes larger and IS divergence becomes smaller.

Fig. 3. Quantification of rhythms behind hibernation. (**a-c**) Distribution of likelihood

values for the set of faster and slower period (ω_1 and ω_2 in FM model) underlying Tb

fluctuation was estimated by maximum likelihood. (**d, e**) Individual variation in the set of

faster and slower period estimated using maximum likelihood (**d**) and IS divergence (**e**).

Fig. 4. Forecasting Tb fluctuation during hibernation. Tb time series was reconstructed (magenta) and predicted (cyan) using the three-quarters (**a**, **b**) and the first half (**c**, **d**) of the whole time series of original Tb data in animals (black). (**e**) Predicted faster and slower period (ω_1 and ω_2 in FM model) using partial time series based on IS divergence estimation. (**f**) The ratio of ω_1 period for half time series to that for whole time series. (**g**) The ratio of ω_2 period for half time series to that for whole time series.

Methods

Animal housing and Tb measurement

Animal housing and Tb measurement were done as described previously[15]. Briefly, female Syrian hamsters (Mesocricetus auratus) were purchased from SLC, Inc., Japan and reared under LD-Warm conditions (light condition = 16L:8D cycle, lights on 05:00–21:00, ambient temperature = 24–25ºC) until most animals weighed over 100–120 g. Then the animals were subjected to surgical operation under inhalation anesthesia with 4% isoflurane (DS Pharma Animal Health, Japan) and intraperitoneal injection of pentobarbital sodium (65 mg/kg, diluted with phosphate-buffered saline; Kyoritu Seiyaku, Japan) for intraperitoneally implanting core body temperature (Tb) loggers (iButton®, Maxim Integrated, USA, #DS1992 L-F5 model) coated with rubber (Plasti Dip, Performix®). After one to two weeks of recovery, animals were transferred to SD-Cold conditions (8L:16D cycle, lights on 10:00–18:00, ambient temperature = 5ºC) for hibernation induction. Animals were individually housed in polypropylene cages, and the Tb of animals were measured every 90 min with an accuracy of 0.5ºC. The cage replacement was done every two weeks and skipped when animals were hibernating at deep torpor to avoid disturbing it. The Tb loggers were recovered from animals sacrificed by decapitation after they were subjected to 10–15 min anesthesia with intraperitoneal injection of pentobarbital sodium (65 mg/kg) and inhalation of 4% isoflurane.

Proposed theoretical models for Tb fluctuation

To simulate Tb fluctuation, we tested two models, frequency modulation and desynchrony models. In the FM model, Tb during hibernation is expressed by the following equation:

$$S[\cos(2\pi\omega_1(1 + A_2\cos(2\pi\omega_2 t + \phi_2))t + \phi_1)] \tag{1}$$

wherein oscillation with frequency of oscillation (ω_1) is assumed to change over time with frequency (ω_2). (b) In the desynchrony model, Tb is expressed as

$$S[\cos(2\pi\omega_1 t + \phi_1) + A_2\cos(2\pi\omega_2 t + \phi_2)]. \tag{2}$$

Although the desynchrony model can yield synchronous limit cycle oscillation when $\omega_1 = \omega_2$, it also yields quasiperiodic oscillations for certain parameter choices with $\omega_1 \napprox \omega_2$. Step function S was used in both models, which realize sharp change in Tb time series. Function $S[x]$ is defined as $S[x]$ = maximum value of Tb if $x > \theta$ and $S[x]$ = minimum value of Tb otherwise. In search of the models and parameters that reproduce each data, all the parameters were rigorously varied (variation is detailed in Supplementary Tables 1, 2).

Quantification of body temperature fluctuation

GHA was conducted as previously described for the studies of music and circadian rhythms[16,40]. GHA is a method in signal processing to quantify periodic components within a certain frequency range, so-called "dominant period" by estimating the most fitted summation of trigonometric functions for given time series[16]. The changes of torpor-IBA period over time were quantified by (1) setting a time window; (2) estimating dominant torpor frequency of each

time window by using GHA; (3) shifting time window by 4 days; and (4) repeating (2) and (3)

until the end of time series. Here, the time-window length was set to be 16 days so that it is

sufficient for covering 2–3 cycles of DT-PA cycles. In the experiment, 13 individual Syrian

golden hamsters continued to hibernate under laboratory conditions for more than 50 days. In

this analysis, the onset of hibernation is defined to be the point such that Tb is lower than 15 °C.

Time series of body temperature within the time window of 16 days for the 13 hamsters is

modeled as

$$\sum_{j=1}^{jmax}\{a_j\cos(2\pi f_j t) + b_j\sin(2\pi f_j t)\} \tag{3}$$

where f_j is the frequency and a_j and b_j are the amplitudes. The resolution of frequency was

1/500 per day, and j_{max} is the maximum number of frequency component, which was set to be

4000 (i.e., $f_{jmax} = 8$ /day). The frequency, called dominant frequency (text) and the amplitude

within each time window were quantified by repeatedly minimizing the square residuals:

$$\int_0^L \left[x(t) - \sum_{j=1}^{jmax}\{a_j\cos(2\pi f_j t) + b_j\sin(2\pi f_j t)\}\right]^2 dt \tag{4}$$

where L is time-window length. Then, the estimated dominant period during hibernation, which

is the inverse of the dominant frequency was plotted over time (Fig. 1).

After quantifying changes in the dominant period of torpor-IBA cycle during

hibernation, the change in the dominant period during the initial 0–48 days of hibernation was

specifically measured by linear regression. The positive (negative) slope of the regression line

indicates that the period of torpor-IBA cycle increases (decreases) with time.

Statistical analysis for model and parameter selection

In search of theoretical models and their parameters reproducing the experimental Tb data

during hibernation, we used the following two statistical quantities: maximum likelihood and

Itakura–Saito divergence (IS), defined as follows, respectively:

$$\text{Log likelihood} = -\frac{N}{2}\log(2\pi\sigma^2) - \frac{1}{2\sigma^2}\sum_{t=1}^{N}\left(x(t) - x_{model}(t)\right)^2, \tag{5}$$

$$\text{IS divergence} = \sum_{t=1}^{N}\left(\frac{x(t)}{x_{model}(t)} - \log\left(\frac{x(t)}{x_{model}(t)}\right) - 1\right), \tag{6}$$

where $x(t)$ is original time series, $x_{model}(t)$ is time series of model, N is length of time series,

and σ^2 is variance of $x(t)$-$x_{model}(t)$. Analysis using IS divergence was conducted as previously

described in the field of speech recognition[18]. In fact, IS divergence is small when the model

reproduces the timing of periodic arousal of experimental data.

To find the best model and the best parameter for each model to reproduce the

experimental data, we varied parameters as shown in Supplementary Tables 1 and 2. When

maximum likelihood becomes larger, IS divergence becomes smaller; the model becomes

closer to the experimental data. The best parameter for each model to reproduce the

experimental data was computationally identified. Maximum likelihood and IS divergence of

each model using the best parameter is shown for each individual's data in Fig. 2k, l. There was a discrepancy between maximum likelihood and IS divergence estimation. This is because maximum likelihood often imposes more penalties than IS divergence does if the difference of $x(t)$ and $x_{model}(t)$ is large; maximum likelihood assumes that the difference of $x(t)$ and $x_{model}(t)$ follows normal distribution while IS divergence assumes that it follows asymmetrical distribution.

To obtain the precise values of ω_1 and ω_2 in FM model, the best parameter was searched within a narrow range of ω_1 and ω_2, after it was searched within various parameter values (Table 1). Also, in the analysis of prediction of Tb fluctuation (Fig. 4), the lower limit of ω_1 was set to be 0.5358 /days (1.83 days) for computation cost because the likelihood of the model with the period around 24h was always small.

Acknowledgments

We thank Y. Chayama and H. Taii for their help in collecting Tb data; R. Enoki, E. Gracheva, and Y. Kawahara for comments on the manuscript; and A. Mochizuki for encouragement. We would like to thank Enago for the English language review. This work was supported by grants from Japan Science and Technology Agency (JPMJCR1913 to G.K.), and from the Japanese Society for the Promotion of Science, and the Ministry of Education, Culture, Sports, Science, and Technology in Japan (JP20H05766 to Y.Y. and JP21K06105 to G.K.), Toray Science Foundation (to Y.Y.), the Takeda Science Foundation (to Y.Y.), and Inamori Foundation (to Y.Y.).

Authors contribution

G.K., Y.Y, and S.G. designed the work and wrote the manuscript. Y.Y. collected Tb data, and S.G. and G.K. developed theoretical models and analyzed the data.

Competing interests

The authors declare no competing interests.

Data availability

Original time-series data are presented in Fig.1, Supplementary Fig. 1. All other data are available upon reasonable request to the corresponding authors G.K. and Y.Y.

Code availability

All code are available upon reasonable request to the corresponding authors G.K. and Y.Y.

References

1. Geiser, F. Hibernation. *Curr. Biol.* **23**, R188-R193 (2013).

2. Mohr, S.M., Bagriantsev, S.N. & Gracheva, E.O. Cellular, Molecular, and Physiological Adaptations of Hibernation: The Solution to Environmental Challenges. *Annu. Rev. Cell and Dev. Biol.* **36**, 315-338 (2020).

3. Lyman, C.P., Willis, J.S., Malan, A. & Wang, L.C.H. *Hibernation and Torpor in Mammals and Birds*. (Academic Press, 1982).

4. Andrews, M.T. Molecular interactions underpinning the phenotype of hibernation in mammals. *J. Exp. Biol.* **222**, jeb160606 (2019).

5. Hampton M. & Andrews M.T. A simple molecular mathematical model of mammalian hibernation. *J Theor Biol* **247**:297-302 (2007).

6. Granada, A.E., Cambras, T., Díez-Noguera, A. & Hanspeter, H. Circadian desynchronization. *Interface Focus* **1**, 153-166 (2011).

7. Uchiyama, M., Okawa, M., Ozaki, S., Shirakawa, S. & Takahashi, K. Delayed phase jumps of sleep onset in a patient with non-24-hour sleep-wake syndrome. *Sleep* **19**, 637-640 (1996).

8. Giroud, S. et al. The Torpid State: Recent advances in metabolic adaptations and Protective Mechanisms. *Front. Physiol.* **11**, 623665 (2021).

9. Zucker, I. Pineal gland influences period of circannual rhythms of ground squirrels. *Am. J. Physiol.* **249**, R111-R115 (1985).

10. Körtner, G. & Geiser, F. The temporal organization of daily torpor and hibernation: circadian and circannual rhythms. *Chronobiol. Int* **17**, 103-128 (2000).

11. Schwartz, C. & Andrews, M.T. Circannual transitions in gene expression: lessons from seasonal adaptations. *Curr. Top. Dev. Biol.* **105**, 247-273 (2013).

12. Kondo, N. et al. Circannual control of hibernation by HP complex in the brain. *Cell* **125**, 161-72 (2006).

13. MacCannell, A.D.V. & Staples, J.F. Elevated ambient temperature accelerates aspects of torpor phenology in an obligate hibernator. *J. Therm. Biol.* **96**, 102839 (2021).

14. Janský, L., Haddad G., Kahlerová, Z. & Nedoma, J. Effect of external hibernation of golden hamsters. *J. Comp. Physiol.* **154**, 427-433 (1984).

15. Chayama, Y., Ando, L., Tamura, Y., Miura, M. & Yamaguchi, Y. Decreases in body temperature and body mass constitute pre-hibernation remodeling in the Syrian golden hamster, a facultative mammalian hibernator. *R. Soc. Open Sci.* **3**, 160002 (2016).

16. Terada, T., Nakajima, H., Tohyama, M. & Hirata, Y. Nonstationary waveform analysis and synthesis using generalized harmonic analysis. In *Proceeding of the IEEE-SP International Symposium on Time-Frequency and Time-Scale Analysis* (IEEE), 429-432 (1994).

17. Armstrong, E.H. Frequency modulation multiplex system. *United States Patent Office* 2,630,497 (1953).

18. Itakura, K. & Saito, S. Analysis synthesis telephony based on the maximum likelihood
method. *In Proc 6th International Congress on Acoustics* C17-C20 (1968).

19. Hashimoto, N., Nakano, S., Yamamoto, K. & Nakagawa, S. Speech recognition based on
Itakura-Saito divergence and dynamics / sparseness constraints from mixed sound of
speech and music by non-negative matrix factorization. In *Interspeech-2014* 2749-2753
(2014).

20. Wutz, A., Melcher, D. & Samaha, J. Frequency modulation of neural oscillations according
to visual task demands. *Proc. Natl. Acad. Sci. USA* **115**, 1346-1351 (2018).

21. Ortmann, S. & Heldmaier, G. Regulation of body temperature and energy requirements of
hibernating Alpine marmots (Marmota marmota). *Am. J. Physiol. Regul. Integr. Comp.*
Physiol. **278**, R698-704 (2000).

22. Arnold, W., Ruf, T., Frey-Roos, F. & Bruns, U. Diet-independent remodeling of cellular
membranes precedes seasonally changing body temperature in a hibernator. *PLoS One* **6**,
e18641 (2011).

23. Sheriff, M.J., Richter, M.M., Buck, C.L. & Barnes, B.M. Changing seasonality and
phenological responses of free-living male arctic ground squirrels: the importance of sex.
Phylos. Trans. R. Soc. Lond. B Biol. Sci. **368**, 20120480 (2013).

24. Siutz, C., Valent, M., Ammann, V., Niebauer, A. & Millesi, E. Sex-specific effects of food
supplementation on hibernation performance and reproductive timing in free-ranging
common hamsters. *Sci. Rep.* **8**, 13082 (2018).

25. Twente, J.W. & Twente, J.A. Regulation of hibernating periods by temperature. *Proc. Natl.*
Acad. Sci. USA **54**, 1058-1061 (1965).

26. Geiser, F. & Kenagy, G.J. Torpor duration in relation to temperature and metabolism in
hibernating ground squirrels. *Physiol. Zool.* **61**, 442-449 (1988).

27. Malan, A., Ciocca, D., Challet, E. & Pévet, P. Implicating a temperature-dependent clock
in the regulation of torpor bout duration in classic hibernation. *J. Biol. Rhythms* **33**, 626-
636 (2018).

28. Pittendrigh, C.S. On temperature independence in the clock system controlling emergence
time in Drosophila. *Proc. Natl. Acad. Sci. USA* **40**, 1018-1029 (1954).

29. Kurosawa, G., Fujioka, A., Koinuma, S., Mochizuki, A. & Shigeyoshi, Y. Temperature—
amplitude coupling for stable biological rhythms at different temperatures. *PLoS Comput.*
Biol. **13**, e1005501 (2017).

30. Oklejewicz, M., Daan, S. & Strijkstra, A.M. Temporal organisation of hibernation in wild-
type and tau mutant Syrian hamsters. *J. Comp. Physiol. B* **171**, 431-439 (2001).

31. van der Vinne, V., Bingaman, M.J., Weaver, D.R. & Swoap, S.J. Clocks and meals keep
mice from being cool. *J. Exp. Biol.* **221**, jeb179812 (2018).

32. Kirsch, R., Ouarour, A. & Pévet, P. Daily torpor in the Djungarian hamster (Phodopus
sungorus): photoperiodic regulation, characteristics and circadian organization. *J. Comp.*
Physiol. **168**, 121-128 (1991).

33. Ruf, T. & Geiser, F. Daily torpor and hibernation in birds and mammals. *Biol. Rev. Camb.*
Philos. Soc. **90**, 891–926 (2015).

34. Gwinner, E. Circannual rhythms in birds. *Curr. Opin. Neurobiol.* **13**, 770-778 (2003).

35. Mrosovsky, N. Hibernation and the Hypothalamus. In: *Hibernation and the*
Hypothalamus. Neuroscience Series. (Springer, 1971).

36. Pengelley, E.T., Aloia, R.C. & Barnes, B.M. Circannual rhythmicity in the hibernating
ground squirrel Citellus lateralis under constant light and hyperthermic ambient
temperature. *Comp. Biochem. Physiol. A Physiol.* **61**, 599-603 (1978).

37. Carr, A.J.F. et al. Photoperiod differentially regulates circadian oscillators in central and
peripheral tissues of the syrian hamster. *Curr. Biol.* **13**, 1543–1548 (2003).

38. Johnston, J.D. et al. Evidence for an endogenous per1- and ICER-independent seasonal
timer in the hamster pituitary gland. *FASEB J.* **17**, 810–815 (2003).

39. Herwig, A. et al. Hypothalamic ventricular ependymal thyroid hormone deiodinases are
an important element of circannual timing in the Siberian hamster (Phodopus sungorus).
PLoS ONE **8**, e62003 (2013).

40. Gibo, S. & Kurosawa, G. Theoretical study on the regulation of circadian rhythms by
RNA methylation. *J. Theor. Biol.* **490**, 110140 (2020).

Supplementary Information

Supplementary Figs. 1 to 10

Supplementary Tables 1 to 2