Universal COVID-19 vaccine with updated spike antigen confers full protection against all
SARS-CoV-2 variants of concern.

Sapna Sharma [a], Thomas Vercruysse [a, b], Lorena Sanchez-Felipe [a], Winnie Kerstens [a, b], Rana Abdelnabi

[a], Caroline Foo [a], Viktor Lemmens [a], Dominique Van Looveren [a, b], Piet Maes [c], Guy Baele [d], Birgit

Weynand [e], Philippe Lemey [d], Johan Neyts [a], Hendrik Jan Thibaut [a, b, &], and Kai Dallmeier [a, &, #]

[a] KU Leuven Department of Microbiology, Immunology and Transplantation, Rega Institute,
Laboratory of Virology, Molecular Vaccinology and Vaccine Discovery, Leuven, Belgium.

[b] KU Leuven Department of Microbiology, Immunology and Transplantation, Rega Institute,
Laboratory of Virology and Chemotherapy, Translational Platform Virology and Chemotherapy,
Leuven, Belgium.

[c] KU Leuven Department of Microbiology, Immunology and Transplantation, Rega Institute,
Laboratory of Clinical and Epidemiological Virology, Zoonotic Infectious Diseases Unit, Leuven,
Belgium.

[d] KU Leuven Department of Microbiology, Immunology and Transplantation, Rega Institute,
Laboratory of Clinical and Epidemiological Virology, Evolutionary and Computational Virology,
Leuven, Belgium.

[e] KU Leuven Department of Imaging and Pathology, Translational Cell and Tissue Research, B-3000
Leuven, Belgium.
20

21   & contributed equally

22   # correspondence should be addressed to: kai.dallmeier@kuleuven.be
23

Abstract

All currently used first-generation COVID-19 vaccines are based on prototypic spike sequences from

ancestral 2019 SARS-CoV-2 strains. However, it remains unclear to which extent vaccination protects

against variants of concern (VOC) which fuel the ongoing pandemic. Here we show in a stringent

hamster challenge model that immunization using prototypic spike expressed form a potent YF17D viral

vector (Sanchez-Felipe et al. 2021) provides vigorous protection against infection with ancestral virus

and VOC Alpha (B.1.1.7), however, is insufficient to provide optimal protection against the Beta

(B.1.351) variant. To improve vaccine efficacy, a revised vaccine candidate was created that carries a

modified spike antigen designed to cover the entire VOC spectrum. Vaccination of hamsters with this

updated vaccine candidate provides full protection against intranasal challenge with all four VOC Alpha,

Beta, Gamma (P.1) and Delta (B.1.617.2) resulting in complete elimination of infectious virus from the

lungs and a marked improvement in lung pathology. Vaccinated hamsters did also no longer transmit
the Delta variant to non-vaccinated sentinels. Overall, our data indicate that current first-generation
COVID-19 vaccines need to be urgently updated to cover emerging sequence diversity of VOCs to
maintain vaccine efficacy and to impede virus spread at the community level.
40
41 **Key Words:** SARS-CoV-2, variants of concern (VOC), vaccine efficacy, antigenic cartography, virus
42 transmission, hamster model
43

Introduction

Severe Acute Respiratory Syndrome Corona Virus 2 (SARS-CoV-2) emerged as zoonosis likely from a limited number of spill-over events into the human population (Holmes et al., 2021). Nevertheless, the ongoing COVID-19 pandemic is entirely driven by variants that evolved during subsequent large-scale human-to-human transmission. In particular, mutations within the viral spike protein are under continuous surveillance (ECDC, 2021) considering their role in viral pathogenesis and as target for virus-neutralizing antibodies (nAb). Following early diversification, the D614G SARS-CoV-2 variant (B.1 lineage) became dominant in March 2020. Consecutively, Variants of Concern (VOC) were identified in many countries with increased transmissibility, virulence and evidence for escape from naturally acquired and vaccine-induced immunity (Tian et al., 2021). Each of the four currently recognized VOCs harbor a unique set of partially convergent, partially unique spike mutations as compared to prototypic (Wuhan) or early European D614G (B.1) lineages of SARS-CoV-2, namely VOC Alpha (B.1.1.7; N501Y D614G), Beta (B.1.351; K417N E484K N501Y D614G), Gamma (P.1; K417T E484K N501Y D614G) and Delta (B.1.617.2; K417T L452R T478K D614G P681R)(Cella et al., 2021). N501Y first detected in VOC Alpha has been linked to an enhanced transmissibility due to an increased affinity for the human ACE-2 receptor (Liu et al., 2021a; Moyo-Gwete et al., 2021). Subsequent emergence of E484K within this lineage hampers the activity of nAb suggestive for immune escape (Graham et al., 2021; Muik et al., 2021; Wu et al., 2021). Likewise, a combination of K417N E484K (Greaney et al., 2021) may explain a marked reduction in vaccine efficacy (VE) of some vaccines such as ChAdOx1 nCoV-19 (AstraZeneca, Vaxzevria) in clinical trials in South Africa during high prevalence of VOC Beta (Madhi et al., 2021). Similarly, sera from vaccinees immunized with first-generation mRNA (Pfizer-BioNTech, Cormirnaty; Moderna, mRNA-1273) or nanoparticle subunit vaccines (Novavax) showed a substantial drop in neutralizing capacity for VOC Beta (Wang et al., 2021). Furthermore VOC Gamma harboring K417T E484K emerged in regions of Brazil with high seroprevalence, hence despite naturally acquired immunity against prototypic SARS-CoV-2 (Sabino et al., 2021). VOC Delta was first identified in October 2020 in India (Cherian et al., 2021; Hoffmann et al., 2021; Yadav et al., 2021) and has since then become the predominant SARS-CoV-2 lineage worldwide, driven by a substantially increased transmissibility (Liu et al., 2021b).

All currently licensed COVID-19 vaccines and vaccine candidates in advanced clinical development are based on antigen sequences of early SARS-CoV-2 isolates that emerged in 2019 (Kyriakidis et al., 2021). We also reported on a YF17D-vectored SARS-CoV-2 vaccine candidate using prototypic spike as vaccine antigen (YF-S0; S0) that had an outstanding preclinical efficacy against homologous challenge (Sanchez-Felipe et al., 2021). However, we now demonstrate to what extent VE of S0 and hence first-generation spike vaccines in general, may decline when trialed against VOC Beta in a stringent hamster model (Abdelnabi et al., 2021). Therefore, a second-generation vaccine candidate (YF-S0*) was designed by (*i*) modifying its antigen sequence to catch up with the evolving spike variant

spectrum, in combination with (*ii*) a further stabilized protein conformation (Juraszek et al., 2021).

Furthermore, we here demonstrate that this new S0* vaccine candidate provides full protection against

all current VOCs (Alpha, Beta, Gamma and Delta). Finally, hamsters vaccinated with S0* do no longer

transmit the virus to non-vaccinated sentinels during close contact, even under conditions of prolonged

co-housing and exposure to a high infectious dose of VOC Delta.

Our findings suggest that first-generation COVID-19 vaccines may need to be adapted to follow the

evolution of SARS-CoV-2 variants fueling the ongoing pandemic. This is important as new variants

may emerge that contain critical combinations of driver mutations responsible for both nAb escape (e.g.,

E484K) (Greaney et al., 2021) and enhanced transmission (e.g., N501Y; P681R/H) (Collier et al., 2021)

already observed in variant of interest (VOI) Mu (B.1.621) currently in surge in some regions of Latin

America (ECDC, 2021).

Results and Discussion

Reduced efficacy of first-generation spike vaccine against VOCs Alpha and Beta

To assess VE of prototypic spike antigen against VOCs, hamsters were vaccinated twice with each 10^4
PFU of YF-S0 (S0) or sham at day 0 and 7 via the intraperitoneal route (Sanchez-Felipe et al., 2021)
(**Fig. 1A**). Serological analysis at day 21 confirmed that 30/32 (94%) vaccinated hamsters had
seroconverted to high levels of nAbs against prototypic SARS-CoV-2 with geometric mean titre (GMT)
of 2.3 $\log_{10}$ (95% CI 2.0-2.6) (**Fig. 1B**). Next, animals were challenged intranasally with 1×10^3 $TCID_{50}$
of either prototypic SARS-CoV-2, VOC Alpha or Beta as established and characterized before in the
hamster model (Abdelnabi et al., 2021). At day four after infection (4 dpi), viral replication was
determined in lung tissue by qPCR and virus titration (**Fig. 1C, D**). In line with what was originally
described for S0 (Sanchez-Felipe et al., 2021), a marked reduction in viral RNA and infectious virus
loads down to undetectable levels (up to $6\log_{10}$ reduction) was observed in the majority of animals
challenged with either prototypic SARS-CoV-2 (8/10; 86% VE) or VOC Alpha (9/10; 88% VE). In
those animals (2/10 and 1/10, respectively) that were not completely protected, virus loads were at least
100 times lower than in infected sham controls. By contrast and despite full immunization, S0
vaccination proved to be less effective against VOC Beta, with only 4/12 hamsters without detectable
infectious virus (60% VE). Nonetheless, in the remaining 8/12 animals with breakthrough infection by
VOC Beta, viral replication was tempered as vaccination still resulted in a 10 to 100-fold reduction in
infectious virus titres relative to sham.

Logistic regression used to define immune correlates of protection (van der Lubbe et al., 2021)
confirmed that comparable nAb levels were required for protection against prototypic SARS-CoV-2
(1.5 $\log_{10}$ for 50% and 2.9 $\log_{10}$ for 90% protection) and VOC Alpha (1.2 $\log_{10}$ for 50% and 2.5 $\log_{10}$ for
90% protection) (**Fig. 1E**). Intriguingly, for VOC Beta a markedly (up to 25x) higher nAb threshold (2.6
$\log_{10}$) was required for 50% protection. Importantly, no 90% protective nAb threshold could be defined
anymore for VOC Beta infection, considering the high number of S0 vaccinated animals with viral
breakthrough ($>10^2$ $TCID_{50}$/100mg lung tissue) (van der Lubbe et al., 2021). Overall, these data suggest
that first-generation vaccines employing prototypic spike as antigen may generally suffer from a
markedly reduced efficacy against emerging SARS-CoV-2 variants, such as VOC Beta.

Updated spike antigen offers complete protection against full range of VOCs

Although prototype S0 showed induction of high titres of nAb against prototypic SARS-CoV-2 (**Fig.**
1B) and protective immunity against prototypic SARS-CoV-2 and VOC Alpha (**Fig. 1C-E**), the
prototypic spike antigen failed to induce consistent nAb responses against remaining VOCs (**Fig. 2A**).
Most importantly, YF-S0 vaccination resulted only in poor seroconversion and low nAb titres against
VOC Beta (seroconversion rate 15/32; GMT 1.0 $\log_{10}$, 95% CI of 0.6-1.3;) and Gamma (19/32; GMT

1.3 log$_{10}$, 95% CI 0.9-1.8). Intriguingly, also a pool of human convalescent serum used as benchmark
(WHO standard NIBSC 20/130) originating from 2020 prior to the surge of VOC (**Fig. 2A-B**) showed
a similar loss of activity against VOC Beta, in line with what was observed in our hamster sera (**Fig.
2A**).

It is not clear if the full spectrum of antigenic variability of current VOCs and emerging variants can be
covered by a COVID-19 vaccine that is based on a single antigen (Lopez Bernal et al., 2021; Rubin,
2021). In an attempt to generate a more universal SARS-CoV-2 vaccine (YF-S0*, S0*), we adapted the
spike sequence in our original YF-S0 construct to include the full amino acid spectrum from VOC
Gamma, plus three extra proline residues (A892P, A942P and V987P) to stabilize spike in conformation
favorable for immunogenicity (Hsieh et al., 2020; Juraszek et al., 2021) (**Fig. 2C**). YF-S0* proved to be
highly immunogenic against prototypic SARS-CoV-2, with nAb levels reaching GMT of 2.2 log$_{10}$ (95%
CI 1.8-2.6) and a seroconversion rate of 21/24 (**Fig. 2D**), comparable to original YF-S0 (GMT 2.3 log$_{10}$,
95% CI 2.0-2.6; 30/32 seroconversion rate) (**Fig. 2A**). Also, for both constructs, seroconversion rates
and nAb levels against VOC Delta were similar (YF-S0: 30/32; GMT 2.0 log$_{10}$, 95% CI 1.7-2.2; YF-
S0*: 22/24; GMT 2.0 log$_{10}$, 95% CI 1.6-2.3). Notably, for YF-S0*, nAb levels and seroconversion rates
against VOC Beta (GMT 2.9 log$_{10}$, 95% CI 2.6-3.2; seroconversion rate 23/24) and Gamma (GMT 3.0
log$_{10}$, 95% CI 2.8-3.2; seroconversion rate 24/24) were markedly increased (by 50 to 80-fold for GMT;
1.7 to 2-times more frequent seroconversion) (**Fig. 2A, D**).

We further studied the pattern of cross-reactivity of the sera raised by the original (YF-S0) and updated
(YF-S0*) vaccine antigen against four different virus variants (prototype; VOCs Beta, Gamma and
Delta) using antigenic cartography (Smith et al., 2004). VOC Alpha was not considered since it did not
differ from the prototype virus, neither regarding VE of S0 nor nAb titre as correlate protection (**Fig. 1**).
Specifically, we constructed a two-dimensional projection that geometrically maps median serum
neutralization titres (SNT$_{50}$) between sera and respective antigens as antigenic distances. This revealed
a pattern of antigenic diversification between prototype virus on the one hand and VOCs Beta and
Gamma on the other hand, with VOC Delta being mapped closer to the prototype virus as compared to
Beta and Gamma. This is consistent with recently described patterns of convergent evolution in spike
for VOCs Beta and Gamma, and Delta climbing a different fitness peak (Martin et al., 2021). In line
with the visual pattern of clustering, antigenic distances for S0* sera were significantly larger to
prototype and VOC Delta as compared to Beta and Gamma (t-test; p<0.001). Intriguingly, this obvious
antigenic drift did not reduce the overall higher potency of S0*, which included an equally strong
humoral response to prototypic spike and VOC Delta (**Fig. 2A, D**).

S0*-vaccinated animals were subsequently challenged with each 10^3 TCID$_{50}$ of either of the four VOCs
Alpha, Beta, Gamma or Delta, and sacrificed 4 dpi for assessment of viral loads in the lung (**Fig. 2F, G**)
and associated lung pathology (**Fig. 2H, I**). In S0*-vaccinated hamsters, viral RNA loads were uniformly

reduced compared to matched sham controls by ~3 (VoC Delta) up to ~6 log$_{10}$ (VoC Gamma) depending
on the respective challenge virus under study (**Fig. 2F**). Importantly, no infectious virus could be
detected anymore (~6log$_{10}$ reduction) in any of the animals vaccinated with S0*, irrespective of which
VOC they had been exposed to (**Fig. 2G**), confirming 100% VE conferred by S0* against all four VOCs.

Protection from infection also translated in a markedly reduced pathology (**Fig. 2H, I**). Non-vaccinated
sham animals developed characteristic signs of bronchopneumonia with perivascular and peribronchial
infiltrations, edema and consolidation of lung tissues (Abdelnabi et al., 2021; Boudewijns et al., 2020).
In contrast, lungs of S0*-vaccinated hamsters remained markedly less affected with a clear reduction in
overall histological scores, irrespectively of the VOC used (**Fig. 2H, I**). In conclusion, second-
generation YF-S0* expressing an updated S0* antigen induced consistently high levels of broadly
neutralizing antibodies (**Fig. 2D**) which translated into efficient protection from lower respiratory tract
infection and COVID-19-like pathology by the entire spectrum of circulating VOCs (**Fig. 2G, H**). VE
of S0* covered VOC Beta and Gamma, i.e. variants harbouring key mutations K417N/T and E484K
escaping original spike-specific nAb activity (**Fig. 2B**), and may therefore offer protection against other
emerging variants such as VOI Mu (E484K) with a similar signature.

Blocking of viral transmission

An added benefit of vaccination at the population level would be an efficient reduction in viral shedding
and transmission by vaccinated people (Eyre et al., 2021), ideally from single-dose vaccination. For
experimental assessment, two groups of hamsters (N=6 each) were either vaccinated once with 10^4 PFU
of S0* or sham (Sanchez-Felipe et al., 2021), and were three weeks later intranasally infected with a
high dose comprising 10^5 TCID$_{50}$ of VOC Delta to serve as index (donor) animals for direct contact
transmission (**Fig. 3A**). At 2 dpi, i.e. at onset of increasing viral loads and shedding (Kaptein et al., 2020;
Sia et al. 2020), index animals were each co-housed with one non-vaccinated sentinel for two
consecutive days. At 4 dpi, index hamsters were sacrificed, and lungs were assessed for viral RNA,
infectious virus and histopathology. Sentinels were sacrificed another two days later and analyzed
accordingly.

As expected from previous experiments, viral loads in S0*-vaccinated index animals were much lower
than in non-vaccinated index animals, or than in sentinels that had been in close contact with non-
vaccinated donors (**Fig. 3B, C**). Importantly, only very low levels of viral RNA and no infectious virus
was observed in non-vaccinated sentinels that had been co-housed with S0*-vaccinated donors. Also,
lung pathology was reduced significantly in vaccinated index and co-housed sentinels as compared to
sham vaccinated index and respective co-housed sentinels (**Fig. 3D**). To our knowledge, this is first
experimental evidence for full protection from SARS-CoV-2 transmission by any vaccine. The block

conferred by S0* appears to be more complete than that observed in humans by current vaccines (Siddle

et al., 2021).

Discussion

Little is known about how well current first-generation vaccines protect against the full spectrum of VOCs. While likely protecting from severe COVID-19 caused by any SARS-CoV-2 strain, a clear drop in VE was observed during clinical trials conducted in regions with high circulation of VOC Beta as paradigm of an E484K Spike variant and others known to escape nAb recognition (Sadoff et al., 2021). Experimentally, such a drop in protective immunity is confirmed by higher viral loads in macaques vaccinated with an Adenovirus-vectored prototype spike antigen (Ad26.COV2.S) and challenged with VOC Beta (Yu et al., 2021). Likewise, in the more stringent hamster model, immunity acquired during previous SARS-CoV-2 (prototype) infection, or by Ad26.COV2.S vaccination, led only to partial restraint of heterologous VOC Beta replication (Tostanoski et al.). In the latter case, replicative viral RNA was still detectable two weeks after challenge (< 2 $\log_{10}$ reduction compared to sham), which is completely in line with the observed failure of prototypic YF-S0 to confer full protection against VOC Beta (**Fig. 1F-H**). By contrast, viral replication was reduced to undetectable levels for all four VOCs by YF-S0* vaccination using an updated spike antigen (**Fig. 2G**). Finally, S0* blocked transmission of VOC Delta (**Fig. 3**). In summary, our findings strongly suggest that first-generation COVID-19 vaccines will need to be adapted to keep up with the evolution of variants driving the ongoing global SARS-CoV-2 pandemic, including variants in surge that contain critical combinations of driver mutations responsible for both nAb escape and enhanced transmission. The stringent hamster model is particularly well suited to assess both aspects of preclinical VE, individual protection and transmission (Abdelnabi et al., 2021).

Overall combined experimental and abundant clinical evidence suggests that first-generation COVID-19 vaccines employing the prototypic spike (from 2019/early 2020) as antigen may not suffice to cover current circulating and emerging SARS-CoV-2 variants anymore; in particular variants carrying mutations in key epitopes (e.g., those containing K417 and E484) targeted by nAb (Greaney et al., 2021).

Methods

Viruses and animals

All virus-related work was conducted in the high-containment BSL3 facilities of the KU Leuven Rega

Institute (3CAPS) under licenses AMV 30112018 SBB 219 2018 0892 and AMV 23102017 SBB 219

2017 0589 according to institutional guidelines. All SARS-CoV-2 strains used throughout this study

were isolated in house (University Hospital Gasthuisberg, Leuven) and characterized by direct

sequencing using a MinION as described before (Boudewijns et al., 2020). Strains representing

prototypic SARS-CoV-2 (Wuhan; EPI_ISL_407976) (Boudewijns et al., 2020), VOC Alpha (B.1.117;

EPI_ISL_791333) and Beta (B.1.351; EPI_ISL_896474) have been described (Abdelnabi et al., 2021).

Strains representing VOC Gamma (P.1; EPI_ISL_1091366) and Delta (B.1.617.2; EPI_ISL_2425097)

were local Belgian isolates from March and April 2021, respectively. All virus stocks were grown on

Vero E6 cells and used for experimental infections at low *in vitro* passage (P) number, P3 for prototype

and P2 for all four VOCs. Absence of furin cleavage site mutations was confirmed by deep sequencing.

Median tissue culture infectious doses ($TCID_{50}$) were defined by titration as described (Abdelnabi et al.,

2021; Boudewijns et al., 2020) using Vero E6 cells as substrate, except for VOC Delta, for which A549

cells were used for a more pronounced virus induced cytopathic effect (CPE).

Housing and experimental infections of hamsters have been described (Boudewijns et al., 2020; Kaptein

et al., 2020; Sanchez-Felipe et al., 2021) and conducted under supervision of the ethical committee of

KU Leuven (license P050/2020 and P055/2021). In brief, 6 to 8 weeks old female Syrian hamsters

(*Mesocricetus auratus*) were sourced from Janvier Laboratories and kept per two in individually

ventilated isolator cages. Animals were anesthetized with ketamine/xylazine/atropine and intranasally

infected with 50 µL of virus stock (25 µL in each nostril) containing either 10^3 or 10^5 $TCID_{50}$ as specified

in the text and euthanized 4 days post infection (dpi) for sampling of the lungs and further analysis.

Animals were monitored daily for signs of disease (lethargy, heavy breathing, or ruffled fur).

Vaccine Candidate

The general methodology for the design and construction of a first YF17D-based SARS-CoV-2 vaccine

candidate (YF-S0) has been described (Sanchez-Felipe et al., 2021). Several mutations were introduced

into original YF-S0 to generate second-generation vaccine candidate YF-S0*. The first series of

mutations is based on the spike sequence of VOC Gamma: L18F, T20N, P26S, D138Y, R190S, K417T,

E484K, N501Y, D614G, H655Y, T1027I, V1176F. A second series of mutations is based on a locked

spike variant described by Juraszek et al. (2021), stabilizing the protein in a more immunogenic

prefusion confirmation: A892P, A942P, V987P (Juraszek et al., 2021).

Production of spike-pseudotyped virus and serum neutralization test (SNT)

Virus-neutralizing antibodies (nAb) were determined using a set of VSV spike-pseudotype viruses

essentially as described (Sanchez-Felipe et al., 2021). For this purpose, four different pseudotypes were

generated using expression plasmids of respective spike variants: for prototype B.1/D614G as before

(Sanchez-Felipe et al., 2021) or sourced from Invivogen for VOC Beta (Cat. No. plv-spike-v3), Gamma

(Cat. No. plv-spike-v5) and Delta (Cat. No. plv-spike-v8). Briefly, depending on the plasmid

background, BHK-21J cells (variant B.1/D614G) or HEK-293T cells (Beta, Gamma and Delta) were

transfected with the respective SARS-CoV-2 protein expression plasmids, and one day later infected

(MOI = 2) with GFP-encoding VSVΔG backbone virus (Whitt, 2010). Two hours later, the medium was

replaced by medium containing anti-VSV-G antibody (I1-hybridoma, ATCC CRL-2700) to neutralize

residual VSV-G input. After 24h incubation at 32℃, the supernatants were harvested. To quantify

nAb, serial dilutions of serum samples were incubated for 1 hour at 37 °C with an equal volume of S-

pseudotyped VSV particles and inoculated on Vero E6 cells for 18 hours.

The resulting number of GFP expressing cells was quantified on a Cell Insight CX5/7 High Content

Screening platform (Thermo Fischer Scientific) with Thermo Fisher Scientific HCS Studio (v.6.6.0)

software. Median serum neutralization titres (SNT_{50}) were determined by curve fitting

in Graphpad Prism after normalization to virus (100%) and cell controls (0%) (inhibitor *vs.* response,

variable slope, four parameters model with top and bottom constraints of 100% and 0%,

respectively).The research reagent for SARS-CoV-2 RNA (NIBSC 20/130) was obtained from the

National Institute for Biological Standards and Control, UK (Mattiuzzo et al., 2020).

Antigenic cartography

We used the antigenic cartography approach developed for influenza hemagglutination inhibition assay

data to study the antigenic characteristics of the SARS-CoV-2 Spikes (Smith et al., 2004). This approach

transforms SNT_{50} data to a matrix of immunological distances. Immunological distance d_{ij} is defined as

$d_{ij} = s_j - H_{ij}$, where H_{ij} is the $\log_2$ titre of virus i against serum j and s_j is the maximum observed titre to

the antiserum from any antigen ($s_j = \max(H_{1j}, \ldots, H_{nj})$). Subsequently, a multidimensional scaling

algorithm was used to position points representing antisera and antigens in a two-dimensional space

such that their distances best fit their respective immunological distances. Even though distances are

measured between sera raised by vaccination using specific Spike antigens (and the pooled NIBSC

serum) and antigens, such an antigenic map also provides estimates of antigenic distances between the

antigens themselves.

Vaccination and challenge

COVID-19 vaccine candidate YF-S0 (Sanchez-Felipe et al., 2021) was used to vaccinate hamsters at

day 0 and day 7 (N=32) with a dose of 10^4 PFU via the intraperitoneal route and control animals (N=18)

were dosed with MEM (Modified Earl's Minimal) medium containing 2% bovine serum as sham

controls. Blood was drawn at day 21 for serological analysis and infection was done on the same day

with prototype (N=10 vaccinated; N=6 sham), VOC Alpha (N=10 vaccinated; and N=6 sham), and Beta

variant (N=12 vaccinated; N=6 sham) with the inoculum of 10^3 TCID$_{50}$ intranasally. Protective nAb

levels were calculated using logistic regression analysis in GraphPad Prism (version 9) as described (van

der Lubbe et al., 2021)

Similarly, hamsters were vaccinated twice with 10^4 YF-S0* (N=24) or sham (N=16) at day 0 and day 7.

Blood was collected at day 21 to analyze nAb in serum, and animals were infected on day 24 with

different variants, including VOC Alpha, Beta, Gamma and Delta with the inoculum of 10^3 TCID50

intranasally (N=6 vaccinated and N=4 sham vaccinated infected against each variant). Lungs were

collected for analysis of viral RNA, infectious virus and for histopathological examination as described

in (Sanchez-Felipe et al., 2021). Resulting vaccine efficacy (VE) was calculated as [1 – (number of

vaccinated animals with detectable virus) / (number of all infected animals)] x 100% per group of

hamsters infected with the same virus strain, whereby a lung viral load $>10^2$ TCID$_{50}$/100mg was set as

cutoff for infection (van der Lubbe et al., 2021).

Viral load and viral RNA quantification

307 Virus loads were determined by titration and RT-qPCR from lung homogenates was performed exactly

308 as previously described in detail (Boudewijns et al., 2020; Kaptein et al., 2020; Sanchez-Felipe et al.,

309 2021).

Histopathology

For histological examination, the lungs were fixed overnight in 4% formaldehyde, embedded in paraffin

and tissue sections (5 μm) after staining with H&E scored blindly for lung damage (cumulative score of

1 to 3 each for congestion, intra-alveolar hemorrhage, apoptotic bodies in bronchial epithelium,

necrotizing bronchiolitis, perivascular edema, bronchopneumonia, perivascular inflammation,

peribronchial inflammation, and vasculitis) as previously established (Abdelnabi et al., 2021;

Boudewijns et al., 2020)

Blocking of viral transmission

Hamsters (N=6) were vaccinated with 10^4 PFU of vaccine once, were bled at day 21 and infected with

delta variant with 1x10^5 TCID50, intranasally. Another group of non-vaccinated hamsters (N=6) were

also infected. Two days post infection index animals were co-housed with sentinels for two days and

separated after two days of exposure. All the index animals were euthanized on day four post infection

and sentinels were sacrificed after 4 days of exposure. Lungs were analyzed for viral RNA and infectious

virus and subjected to histopathology.

Statistical analysis

All statistical analyses were performed using GraphPad Prism 9 software (GraphPad, San Diego, CA, USA). Results are presented as GM± IQR or medians ± IQR as indicated. Data were analyzed using uncorrected Kruskal-Wallis test and considered statistically significant at p-values ≤0.05.

Acknowledgements

We are grateful to Prof. Michael A. Whitt (University of Tennessee Health Science Center) for generously sharing of plasmids for rescue of VSVΔG, and Dr. Maya Imbrechts and Dr. Nick Geukens (PharmAbs, KU Leuven) for help with hybridoma culture. We thank Carolien De Keyzer and Lindsey Bervoets (Rega) for excellent technical assistance with animal experimentation as well as the staff of the Rega animalium for strong support. We thank Jasper Rymenants, Tina van Buyten, Dagmar Buyst and Thibault Francken for steady and timely help with analysing of tissue samples and virus titrations. We finally thank Jasmine Paulissen, Madina Rasulova, Catherina Coun, Céline Sablon, Jolien Timmermans and Nathalie Thys (TPVC) for diligent serology assessment, vaccine stock generation and skilled generation of plasmid constructs.

Current work was supported by the Flemish Research Foundation (FWO) emergency COVID-19 fund (G0G4820N) and the FWO Excellence of Science (EOS) program (No. 30981113; VirEOS project), the European Union's Horizon 2020 research and innovation program (No 101003627; SCORE project and No 733176; RABYD-VAX consortium), the Bill and Melinda Gates Foundation (INV-00636), KU Leuven Internal Funds (C24/17/061) and the KU Leuven/UZ Leuven Covid-19 Fund (COVAX-PREC project). G.B. acknowledges support from the KU Leuven Internal Funds (Grant No. C14/18/094) and the Research Foundation - Flanders ("Fonds voor Wetenschappelijk Onderzoek - Vlaanderen," G0E1420N, G098321N). P.L. acknowledges funding from the European Research Council under the European Union's Horizon 2020 research and innovation program (grant agreement no. 725422-ReservoirDOCS). K.D. acknowledges grant support from KU Leuven Internal Funds (C3/19/057 Lab of Excellence).

Contributions

S.S. and K.D. conceptualization; S.S. animal experimentation; S.S., T.V., W.K. and H.J.T. data generation, analysis and curation; S.S., H.J.T. and K.D. original manuscript draft; S.S. and H.J.T. visualization; T.V. and L.S.F. construct design; T.V., W.K. and D.V.L. serological analysis; R.A. and C.S.F. VoC hamster models; B.W. histological analysis; P.L. and G.B. antigenic cartography; L.S.F., V.L., T.V., W.K., and P.M. vaccine stocks and virus isolation; J.N., H.J.T., and K.D. supervision, writing and project administration; J.N. and K.D. funding acquisition. All authors read, edited and approved the final version of the manuscript.

References

Abdelnabi, R., Boudewijns, R., Foo, C.S., Seldeslachts, L., Sanchez-Felipe, L., Zhang, X., Delang,
L., Maes, P., Kaptein, S.J.F., Weynand, B*., et al.* (2021). Comparing infectivity and virulence of
emerging SARS-CoV-2 variants in Syrian hamsters. EBioMedicine *68*, 103403.

Boudewijns, R., Thibaut, H.J., Kaptein, S.J.F., Li, R., Vergote, V., Seldeslachts, L., Van
Weyenbergh, J., De Keyzer, C., Bervoets, L., Sharma, S*., et al.* (2020). STAT2 signaling restricts
viral dissemination but drives severe pneumonia in SARS-CoV-2 infected hamsters. Nat
Commun *11*, 5838.

Cella, E., Benedetti, F., Fabris, S., Borsetti, A., Pezzuto, A., Ciotti, M., Pascarella, S., Ceccarelli,
G., Zella, D., Ciccozzi, M*., et al.* (2021). SARS-CoV-2 Lineages and Sub-Lineages Circulating
Worldwide: A Dynamic Overview. Chemotherapy *66*, 3-7.

Cherian, S., Potdar, V., Jadhav, S., Yadav, P., Gupta, N., Das, M., Rakshit, P., Singh, S., Abraham,
P., Panda, S*., et al.* (2021). Convergent evolution of SARS-CoV-2 mutations, L452R, E484Q and
P681R, in the second wave of COVID-19 in Maharashtra, India. bioRxiv,
2021.2004.2022.440932.

Collier, D.A., De Marco, A., Ferreira, I., Meng, B., Datir, R.P., Walls, A.C., Kemp, S.A., Bassi, J.,
Pinto, D., Silacci-Fregni, C*., et al.* (2021). Sensitivity of SARS-CoV-2 B.1.1.7 to mRNA vaccine-
elicited antibodies. Nature *593*, 136-141.

ECDC (2021). SARS-CoV-2 variants of concern as of 7 October 2021.

Eyre, D.W., Taylor, D., Purver, M., Chapman, D., Fowler, T., Pouwels, K.B., Walker, A.S., and
Peto, T.E. (2021). The impact of SARS-CoV-2 vaccination on Alpha & Delta variant transmission.
medRxiv, 2021.2009.2028.21264260.

Graham, M.S., Sudre, C.H., May, A., Antonelli, M., Murray, B., Varsavsky, T., Kläser, K., Canas,
L.S., Molteni, E., Modat, M*., et al.* (2021). Changes in symptomatology, reinfection, and
transmissibility associated with the SARS-CoV-2 variant B.1.1.7: an ecological study. Lancet
Public Health *6*, e335-e345.

Greaney, A.J., Loes, A.N., Crawford, K.H.D., Starr, T.N., Malone, K.D., Chu, H.Y., and Bloom, J.D.
(2021). Comprehensive mapping of mutations in the SARS-CoV-2 receptor-binding domain
that affect recognition by polyclonal human plasma antibodies. Cell Host Microbe *29*, 463-
476.e466.

Hoffmann, M., Arora, P., Groß, R., Seidel, A., Hörnich, B.F., Hahn, A.S., Krüger, N., Graichen, L.,
Hofmann-Winkler, H., Kempf, A*., et al.* (2021). SARS-CoV-2 variants B.1.351 and P.1 escape
from neutralizing antibodies. Cell *184*, 2384-2393.e2312.

Holmes, E.C., Goldstein, S.A., Rasmussen, A.L., Robertson, D.L., Crits-Christoph, A., Wertheim,
J.O., Anthony, S.J., Barclay, W.S., Boni, M.F., Doherty, P.C*., et al.* (2021). The origins of SARS-
CoV-2: A critical review. Cell *184*, 4848-4856.

Hsieh, C.L., Goldsmith, J.A., Schaub, J.M., DiVenere, A.M., Kuo, H.C., Javanmardi, K., Le, K.C.,
Wrapp, D., Lee, A.G., Liu, Y*., et al.* (2020). Structure-based design of prefusion-stabilized SARS-
CoV-2 spikes Science *369*, 1501-1505.

Juraszek, J., Rutten, L., Blokland, S., Bouchier, P., Voorzaat, R., Ritschel, T., Bakkers, M.J.G.,
Renault, L.L.R., and Langedijk, J.P.M. (2021). Stabilizing the closed SARS-CoV-2 spike trimer.
Nat Commun *12*, 244.

Kaptein, S.J.F., Jacobs, S., Langendries, L., Seldeslachts, L., Ter Horst, S., Liesenborghs, L., Hens,
B., Vergote, V., Heylen, E., Barthelemy, K.*, et al.* (2020). Favipiravir at high doses has potent
antiviral activity in SARS-CoV-2-infected hamsters, whereas hydroxychloroquine lacks activity.
Proc Natl Acad Sci U S A *117*, 26955-26965.

Kyriakidis, N.C., López-Cortés, A., González, E.V., Grimaldos, A.B., and Prado, E.O. (2021).
SARS-CoV-2 vaccines strategies: a comprehensive review of phase 3 candidates. NPJ Vaccines
6, 28.

Liu, J., Liu, Y., Xia, H., Zou, J., Weaver, S.C., Swanson, K.A., Cai, H., Cutler, M., Cooper, D., Muik,
A.*, et al.* (2021a). BNT162b2-elicited neutralization of B.1.617 and other SARS-CoV-2 variants.
Nature *596*, 273-275.

Liu, Y., Liu, J., Johnson, B.A., Xia, H., Ku, Z., Schindewolf, C., Widen, S.G., An, Z., Weaver, S.C.,
Menachery, V.D.*, et al.* (2021b). Delta spike P681R mutation enhances SARS-CoV-2 fitness over
Alpha variant. bioRxiv.

Lopez Bernal, J., Andrews, N., Gower, C., Gallagher, E., Simmons, R., Thelwall, S., Stowe, J.,
Tessier, E., Groves, N., Dabrera, G.*, et al.* (2021). Effectiveness of Covid-19 Vaccines against
the B.1.617.2 (Delta) Variant. N Engl J Med *385*, 585-594.

Madhi, S.A., Baillie, V., Cutland, C.L., Voysey, M., Koen, A.L., Fairlie, L., Padayachee, S.D.,
Dheda, K., Barnabas, S.L., Bhorat, Q.E.*, et al.* (2021). Efficacy of the ChAdOx1 nCoV-19 Covid-
19 Vaccine against the B.1.351 Variant. N Engl J Med *384*, 1885-1898.

Martin, D.P., Weaver, S., Tegally, H., San, J.E., Shank, S.D., Wilkinson, E., Lucaci, A.G.,
Giandhari, J., Naidoo, S., Pillay, Y.*, et al.* (2021). The emergence and ongoing convergent
evolution of the SARS-CoV-2 N501Y lineages. Cell *184*, 5189-5200.e5187.

Mattiuzzo, G., Bentley, E.M., Hassall, M., Routley, S., Richardson, S., Bernasconi, V.,
Kristiansen, P., Harvala, H. , Roberts, D., Semple, M.G., Turtle, L.C.W., Openshaw P.J.M., *et al.*
(2020). Establishment of the WHO International Standard and Reference Panel for anti-SARS-
CoV-2antibody. WHO/BS/2020.2403.

Moyo-Gwete, T., Madzivhandila, M., Makhado, Z., Ayres, F., Mhlanga, D., Oosthuysen, B.,
Lambson, B.E., Kgagudi, P., Tegally, H., Iranzadeh, A.*, et al.* (2021). SARS-CoV-2 501Y.V2
(B.1.351) elicits cross-reactive neutralizing antibodies. bioRxiv, 2021.2003.2006.434193.

Muik, A., Wallisch, A.K., Sänger, B., Swanson, K.A., Mühl, J., Chen, W., Cai, H., Maurus, D.,
Sarkar, R., Türeci, Ö.*, et al.* (2021). Neutralization of SARS-CoV-2 lineage B.1.1.7 pseudovirus
by BNT162b2 vaccine-elicited human sera. Science *371*, 1152-1153.

Rubin, R. (2021). COVID-19 Vaccines vs Variants-Determining How Much Immunity Is Enough.
Jama *325*, 1241-1243.

Sabino, E.C., Buss, L.F., Carvalho, M.P.S., Prete, C.A., Jr., Crispim, M.A.E., Fraiji, N.A., Pereira,
R.H.M., Parag, K.V., da Silva Peixoto, P., Kraemer, M.U.G.*, et al.* (2021). Resurgence of COVID-
19 in Manaus, Brazil, despite high seroprevalence. Lancet *397*, 452-455.

Sadoff, J., Gray, G., Vandebosch, A., Cárdenas, V., Shukarev, G., Grinsztejn, B., Goepfert, P.A.,
Truyers, C., Fennema, H., Spiessens, B., *et al.* (2021). Safety and Efficacy of Single-Dose
Ad26.COV2.S Vaccine against Covid-19. N Engl J Med *384*, 2187-2201.

Sanchez-Felipe, L., Vercruysse, T., Sharma, S., Ma, J., Lemmens, V., Van Looveren, D.,
Arkalagud Javarappa, M.P., Boudewijns, R., Malengier-Devlies, B., Liesenborghs, L., *et al.*
(2021). A single-dose live-attenuated YF17D-vectored SARS-CoV-2 vaccine candidate. Nature
590, 320-325.

Siddle, K.J., Krasilnikova, L.A., Moreno, G.K., Schaffner, S.F., Vostok, J., Fitzgerald, N.A.,
Lemieux, J.E., Barkas, N., Loreth, C., Specht, I., *et al.* (2021). Evidence of transmission from fully
vaccinated individuals in a large outbreak of the SARS-CoV-2 Delta variant in Provincetown,
Massachusetts. medRxiv.

Smith, D.J., Lapedes, A.S., de Jong, J.C., Bestebroer, T.M., Rimmelzwaan, G.F., Osterhaus, A.D.,
and Fouchier, R.A. (2004). Mapping the antigenic and genetic evolution of influenza virus.
Science *305*, 371-376.

Tian, D., Sun, Y., Zhou, J., and Ye, Q. (2021). The global epidemic of SARS-CoV-2 variants and
their mutational immune escape. J Med Virol. 2021 Oct 5. doi: 10.1002/jmv.27376.

Tostanoski, L.H., Yu, J., Mercado, N.B., McMahan, K., Jacob-Dolan, C., Martinot, A.J., Piedra-
Mora, C., Anioke, T., Chang, A., Giffin, V.M., *et al.* Immunity elicited by natural infection or
Ad26.COV2.S vaccination protects hamsters against SARS-CoV-2 variants of concern. Science
Translational Medicine *0*, eabj3789.

van der Lubbe, J.E.M., Rosendahl Huber, S.K., Vijayan, A., Dekking, L., van Huizen, E.,
Vreugdenhil, J., Choi, Y., Baert, M.R.M., Feddes-de Boer, K., Izquierdo Gil, A., *et al.* (2021).
Ad26.COV2.S protects Syrian hamsters against G614 variant SARS-CoV-2 and does not
enhance respiratory disease. NPJ Vaccines *6*, 39.

Wang, P., Nair, M.S., Liu, L., Iketani, S., Luo, Y., Guo, Y., Wang, M., Yu, J., Zhang, B., Kwong,
P.D., *et al.* (2021). Antibody resistance of SARS-CoV-2 variants B.1.351 and B.1.1.7. Nature *593*,
130-135.

Whitt, M.A. (2010). Generation of VSV pseudotypes using recombinant ΔG-VSV for studies on
virus entry, identification of entry inhibitors, and immune responses to vaccines. J Virol
Methods *169*, 365-374.

Wu, K., Werner, A.P., Moliva, J.I., Koch, M., Choi, A., Stewart-Jones, G.B.E., Bennett, H.,
Boyoglu-Barnum, S., Shi, W., Graham, B.S., *et al.* (2021). mRNA-1273 vaccine induces
neutralizing antibodies against mutants from global SARS-CoV-2 variants. bioRxiv,
2021.2001.2025.427948.

Yadav, P.D., Sapkal, G.N., Abraham, P., Ella, R., Deshpande, G., Patil, D.Y., Nyayanit, D.A.,
Gupta, N., Sahay, R.R., Shete, A.M., *et al.* (2021). Neutralization of variant under investigation
B.1.617 with sera of BBV152 vaccinees. Clin Infect Dis.

Yu, J., Tostanoski, L.H., Mercado, N.B., McMahan, K., Liu, J., Jacob-Dolan, C., Chandrashekar,
A., Atyeo, C., Martinez, D.R., Anioke, T., *et al.* (2021). Protective efficacy of Ad26.COV2.S
against SARS-CoV-2 B.1.351 in macaques. Nature *596*, 423-427.

486
487
Figure 1. Immunogenicity and protective efficacy of first-generation Spike vaccine against
prototype SARS-CoV-2 and VOCs Alpha and Beta. A, Vaccination scheme with prototypic
YF17D-based vaccine candidate YF-S0 (S0). Syrian hamsters were immunized twice
intraperitoneally with 10^4 PFU of S0 on day 0 and 7 and inoculated intranasally on day 21 with
10^3 median tissue-culture infectious dose (TCID$_{50}$) of either prototype SARS-CoV-2 (*grey*
circles), VOC Alpha (*green triangles*) or VOC Beta (*blue squares*). **B**, nAb titers against
prototypic spike (D614G) pseudotyped virus on day 21 after vaccination. Red datapoint
indicates the NIBSC 20/130 human reference sample included as benchmark. **C, D**, Viral loads
in hamster lungs four days after infection quantified by quantitative RT-PCR (**C**) and virus
titration (**D**). **E**, correlates of protection against prototype SARS-CoV-2, VOC Alpha and VOC
Beta. Logistic regression model to calculate nAb titers correlating with 50% and 90%
probability for protection. '*Protected'* was defined by a viral load $<10^2$ TCID$_{50}$/100mg lung
tissue and '*infected'* by a viral load $>10^2$ TCID$_{50}$/100mg lung tissue (van der Lubben et al.,
2021). Shaded areas indicate 95% CI. LLOQ is lower limit of quantification. Error bars denote
median ± IQR. Differences between groups were analyzed using non-parametric Kruskal
Wallis test uncorrected for ties.

504
505
Figure 2. A vaccine based on the updated Spike antigen S0* offers complete protection
against all four VOCs. A, nAb titers against prototypic (*grey*), VOC Beta (*blue*), VOC Gamma
(*orange*) and VOC Delta (*purple*) spike pseudotyped virus on day 21 after vaccination with
prototype YF-S0. Red datapoint indicates the NIBSC 20/130 human reference sample (see Fig.
1B). **B**, Neutralization curves for NIBSC 20/130 human reference sample against same set of
pseudotyped viruses. **C**, Schematic of the updated YF-S0* (S0*) vaccine candidate based on
VOC Gamma, plus three extra stabilizing proline residues. Vaccination scheme with YF-S0*.
Syrian hamsters were immunized twice intraperitoneally with 10^4 PFU of S0* on day 0 and 7
and inoculated intranasally on day 24 with 10^3 median tissue-culture infectious dose ($TCID_{50}$)
of either VOC Alpha (*green*), VOC Beta (*blue*), VOC Gamma (*orange*) and VOC Delta
(*purple*). **D**, nAb titers against prototypic, VOC Beta, VOC Gamma and VOC Delta spike
pseudotyped virus on day 21 after vaccination with YF-S0*. Red datapoint indicates the NIBSC
20/130 human reference sample. **E**, antigenic cartography. Cross-reactivity of the sera raised
by original S0 (*grey squares*) and updated S0* (*orange squares*) vaccine antigen against four
different virus variants (circles: prototype, *grey*; VOC Beta, *blue*; Gamma, *orange*, and Delta,
purple) plotted on two-dimensional distance map (Smith et al., 2004). **F, G,** Viral loads in
hamster lungs four days after infection quantified by quantitative PCR with reverse
transcription (RT–qPCR) (**F**) and virus titration (**G**). **H**, cumulative lung pathology scores from
H&E-stained slides of lungs for signs of damage. **I**, Representative H&E-stained images of

sham- or S0*-vaccinated hamster lungs after challenge. Perivascular inflammation (*black*
arrows) with focal endothelialitis (*green arrows*); peri-bronchial inflammation (*blue arrows*);
patches of bronchopneumonia (*red arrows*). Error bars denote median ± IQR. Differences
between groups were analyzed using non-parametric Kruskal Wallis test uncorrected for ties.
529

530
531
Figure 3. A vaccine based on the updated Spike antigen S* completely prevents
transmission of the Delta variant. Effect of YF-S0* vaccination on viral transmission to non-
vaccinated contact hamsters. Index hamsters were either sham-vaccinated or vaccinated with a
single dose of 10^4 PFU of YF-S0* and infected intranasally on day 21 with 10^5 TCID$_{50}$ of VOC
Delta. Two days after infection, index animals were paired and co-housed with each one naïve
sentinel. Index and sentinel animals were sacrificed each 4 days after infection or exposure,
respectively. **B, C,** Viral loads in hamster lungs four days after infection quantified by
quantitative RT-qPCR (**B**) and virus titration (**C**). **D**, cumulative lung pathology scores from
H&E-stained slides of lungs for signs of damage. Error bars denote median $\pm$ IQR. Differences
between groups were analyzed using non-parametric Kruskal Wallis test uncorrected for ties.
542