Phage-encoded sigma factors alter bacterial dormancy

Schwartz DA[*], Lehmkuhl BK, Lennon JT[*]

Department of Biology, Indiana University, Bloomington, IN 47405, USA

* Corresponding authors: danschw@iu.edu and lennonj@iu.edu

ABSTRACT

By entering a reversible state of reduced metabolic activity, dormant microorganisms are able

to tolerate suboptimal conditions that would otherwise reduce their fitness. Dormancy may also

benefit bacteria by serving as a refuge from parasitic infections. Here we focus on dormancy in

the *Firmicutes*, where endospore development is transcriptionally regulated by the expression of

sigma factors. A disruption of this process could influence the survivorship and reproduction of

phages that infect spore-forming hosts with implications for coevolutionary dynamics. Here, we

characterized the distribution and diversity of sigma factors in nearly 3,500 phage genomes.

Homologs of sporulation-specific sigma factors were identified in phages that infect spore-forming

hosts. Unlike sigma factors required for phage reproduction, the sporulation-like sigma factors

were non-essential for lytic infection. However, when expressed in the spore-forming *Bacillus*

subtilis, sigma factors from phages activated the bacterial sporulation gene network and reduced

spore yield. Our findings suggest that the acquisition of host-like transcriptional regulators may

allow phages to manipulate a complex and ancient trait in one of the most abundant cell types on

Earth.

INTRODUCTION

Dormancy is a life history strategy that allows individuals to enter a reversible state of reduced metabolic activity. An example of convergent evolution, it has independently arisen throughout the tree of life as a means of coping with fluctuating and unpredictable environments[1]. Dormancy is particularly prevalent among microbial life forms where it contributes to the persistence and fitness of populations in environments where variables like pH, oxygen, and resource availability are suboptimal for growth and reproduction[2]. In addition to buffering populations against abiotic features of the environment, dormancy may be reinforced through dynamics that arise from species interactions. For example, dormancy diminishes the strength of competition, which in turn can promote species coexistence[3]. In addition, dormancy may benefit populations by serving as a refuge against predator consumption or parasite infection[4,5].

Among microorganisms, dormancy can protect hosts from viral parasites in a number of ways. As cells transition into an inactive state, they often undergo morphological changes that affect how viruses physically interact with their host. For example, the formation of dormant cells, such as cysts and spores, often involves the development of a thick exterior coating[6-8] that masks the surface molecules used by viruses for attachment[9-12]. Even if a virus is able to gain entry into a dormant cell, parasite productivity will be low owing to constraints imposed by the host's reduced metabolism[13-16]. Furthermore, viral defense genes are often located in proximity to genes that regulate dormancy and cell suicide, suggesting that dormancy may contribute to multilayered protection against viral infection[17,18]. For example, virus-induced dormancy has been linked to CRISPR-Cas systems in bacteria[20] and archaea[19,20]. As a physiological refuge[21], dormancy can

confer herd immunity and diminish the spread of epidemics[22], which may ultimately shape host-

virus coevolutionary dynamics.

A take-home lesson from studies on antagonistic coevolution is that host defenses are prone to

being overcome by viruses[23,24]. One general mode of virus adaptation involves the acquisition of

host genes. Viral genomes commonly encode homologs of genes that are involved in host

metabolism. These so-called "auxiliary" genes can alter cellular processes in ways that affect virus

fitness[25]. Originally motivated by the discovery of photosynthesis genes in marine

cyanophages[26,27], auxiliary genes have been implicated in host nutrition (e.g., N and P metabolism)

and energy acquisition (sulfur oxidation, fermentation, etc.), along with basic cellular functions

such as protein translation and bacterial communication via quorum sensing[28-30]. In addition, some

virus genomes contain host defense genes, which has led to speculation that auxiliary genes may

modify parasite infectivity and reproduction[31,32]. Similarly, some phage genomes have been

reported to have genes similar to those required for the development of endospore-forming bacteria

[33-42]. Phages might use sporulation homologs to inhibit their host from entering a dormancy refuge,

thereby enhancing the reproductive component of parasite fitness. Alternatively, phages might

exploit sporulation in a way that extends longevity and thereby enhance the survivorship

component of fitness. This could happen through a process known as entrapment whereby a phage

genome is translocated into the developing spore resulting in the production of a "virospore"[12,43-

46]. Analogous to pseudolysogeny, the phage genome is protected by the endospore from conditions

that would otherwise contribute to phage decay without it being integrated into the host

chromosome[45]. When environmental conditions improve, the dormant cell undergoes germination,

and the phage resumes its lytic reproductive cycle.

As a complex form of dormancy, sporulation presents phages with many opportunities for intervention. For proper development, sporulation requires the coordinated regulation of a large gene network[47-49]. The central regulatory module of sporulation relies on the activity of sigma factors, the exchangeable subunit of the transcriptional machinery which dictates promoter specificity of RNA polymerase[50]. Among bacteria, a primary sigma factor (*sigA* in *Bacillus subtilis)* is essential for growth, reproduction, and other housekeeping processes in a wide range of bacteria[51]. During *B. subtilis* spore development *sigA* is swapped out by a cascade of sporulation-specific sigma factors, each driving the expression of a subset of sporulation genes in distinct cellular compartments at specific times[52]. Following an asymmetrical cell division, gene expression in the maturing spore (i.e., forespore) is driven first by *sigF* and then *sigG*, while expression in the mother cell is driven by *sigE* and then *sigK*. Sigma factors are also encoded by some phage genomes, where they regulate phage gene expression during different stages of lytic development[52,53]. More recently, homologs of sporulation-specific sigma factors have been identified in phage genomes[34,41,42,54], yet their function has not been explored. As central nodes of the sporulation gene network, sigma factors could be coopted by phages to modify the outcome of sporulation.

Here, we use a combination of bioinformatics and laboratory experiments to test whether homologs of sporulation-specific sigma factors can be used by phages to manipulate host dormancy. Using sequence homology and phylogenetic analyses, we identify and classify hundreds of phage-encoded sigma factors. We find that phages capable of infecting spore-forming hosts preferentially encode sigma factors that are homologous to the forespore-specific sigma

92 factors, *sigF* and *sigG*. When homologs of sporulation-specific sigma factors are expressed in *B.*

93 *subtilis*, we observed that conserved phage-encoded homologs alter host gene expression and

94 reduce spore yield. Together, our findings have implications for understanding dormancy

95 dynamics in environmental, engineered, and host-associated ecosystems where endospores

96 constitute one of the most abundant cell types on Earth[55].

98   **RESULTS**

Distribution of sigma factors in phage genomes — We characterized the diversity and

distribution of sigma factors in more than 3,400 phage genomes in the viral orthologous groups

(VOG) database. From this, we found that 14% of all the genomes analyzed contained at least one

sigma factor gene while some genomes (0.7%) contained up to three different homologs (Fig. 1).

The distribution of phage-encoded sigma factors was not random with respect to virus or host

taxonomy (Fisher's Exact Test, $P_{simulated}$ < 0.0001, Fig. 1). Sigma factor containing phages were

most prevalent among those that infect the *Cyanobacteria* (65%), *Proteobacteria* (15%) and

Firmicutes (19%). Among the tailed phages (*Caudovirales*), sigma factors were recovered in 5 of

9 families. Three of these five viral families (*Siphoviridae*, *Podoviridae*, and *Myoviridae*) have

members that infect hosts from multiple bacterial phyla. However, only among phages that infect

Firmicutes do all three of these virus families have members with sigma factor genes (Fig. 1c).

Firmicutes phages were also notable in that many of them encode multiple sigma factors.

For phages that infect *Cyanobacteria* and *Proteobacteria*, 95% of the genomes with a sigma factor

only contained a single copy of such a gene (Fig. 1d). In contrast, for phages that infect *Firmicutes*,

41% of the genomes with a sigma factor possessed two or three gene copies. Sigma factor gene

multiplicity was detected in diverse *Firmicutes* phages, predominantly among strains that could

infect spore-forming genera (Fig. S1). Most of these phages belonged to the *Herelleviridae*, a

family of strictly lytic phages that was recently split from the *Myoviridae* and are thought to only

infect bacteria in the *Firmicutes*[56]. Of the 102 *Herelleviridae* phages in our dataset, half possessed

multiple sigma factors (n = 51) and only infected *Bacillus* hosts (Fig. S2). In contrast, none of the

49 phages that infect non spore-forming genera (*Enterococcus*, *Lactiplantibacillus*, *Lactobacillus*,

Listeria, *Staphylococcus*) possessed more than a single sigma factor gene. Additionally, a similar

trend was found among *Firmicutes* phages belonging to the *Siphoviridae*. Within this group, there

were five phages with multiple sigma factors (Fig. S2), four that infect hosts capable of forming

endospores (*Bacillus* and *Brevibacillus*), and one whose host range likely evolved from a *Bacillus*

to *Staphylococcus* host[57].

Sporulation-specific sigma factors are encoded by *Firmicutes* phages — Many of the sigma

factors recovered in *Firmicutes* phages were similar to the bacterial-encoded sigma factors

involved in sporulation. To interpret this finding in an evolutionary context, we constructed a

phylogeny of the VOG sigma factors together with sigma factors from diverse bacteria.

Sporulation-specific sigma factors from bacteria belonged to a monophyletic clade that also

contained *sigB*, a gene that *B. subtilis* uses to regulate its stress response (Fig. 2a,b). While some

phage-encoded sigma factors were found in phage-specific clades, others clustered with bacterial-

encoded sigma factors, including those that are known to regulate sporulation. All phage-encoded

sigma factors in the sporulation clade were from *Firmicutes* phages, apart from two cyanophage

genes that grouped with the *sigB* sub-clade.

To complement the phylogenetic analyses, we classified phage-encoded sigma factors by

estimating homology to bacterial sigma factor protein families (TIGRFAMs) using hidden Markov

models. Phage-encoded homologs of sporulation-specific sigma factors of *Firmicutes*, henceforth

"sporulation-like" sigma factors, were found nearly exclusively in phages that infect hosts of this

phylum (Fig. 2c). Homologs of three of the four sporulation-specific sigma factors (*sigF*, *sigG* and

sigE) were only detected in *Firmicutes* phages, with the majority resembling *sigF*. Furthermore,

these homologs were only recovered from phages that infect *Bacillus*, *Clostridium*, and

Brevibacillus, which are genera that commonly engage in endosporulation (Fig. S4). However, we

did identify homologs of the sporulation-specific *sigK* in a few phages that infect *Cyanobacteria*

and *Actinobacteria.*

The sporulation-like sigma factors recovered from phage genomes were distinct from phage-

encoded sigma factors that regulate expression during lytic replication (Fig. 2a). In our

phylogenetic analysis, sigma factors that are essential for phage development clustered in one of

three other groups. First, there was a phage-specific clade of genes from phages that infect

Proteobacteria and *Cyanobacteria*. This clade included gp55 from phage T4, which is known to

control transcription during late infection stages[50,53]. Second, we identified a phage-specific clade

of genes from *Firmicutes* phages, which included gp34, a sigma factor that controls late gene

expression in *B. subtilis* phage SPO1[36]. Last, there was a paraphyletic cluster that included phage-

and bacterial-derived sigma factors. This group contained gp28, which regulates the expression of

middle infection genes in phage SPO1[36]. In addition, the latter group contained alternative sigma

factors that are involved in bacterial regulation of motility (*sigD*), stationary phase (*sigH*), and heat

stress response (*sigI*).

The pattern of gene multiplicity among *Firmicutes* phages (Fig. S1) reflects that these viruses

contain sigma factors with divergent functions. They either cluster with genes known to regulate

the lytic cycle or with bacterial sporulation-specific sigma factors (Fig. 2b). For example, *B.*

subtilis phage SPO1 encodes two sigma factors that are essential for phage replication (gp28 and

gp34)[36]. Likewise, phage SP10 has two sigma factors that cluster with the essential SPO1 genes

which are known to regulate phage genes[58]. Each of these phages also encode a third homolog that

clusters with sporulation-specific sigma factors. Similarly, all but two of the 58 *Firmicutes* phages

exhibiting gene multiplicity contain one or two sporulation-like sigma factors (Fig. S3).

Additionally, in several *Bacillus* phages the sole sigma factor is sporulation-like. This group of

phages was enriched with representatives from the *Podoviridae* and the *Siphoviridae*, which

includes some temperate phages such as Wβ which infects *B. anthracis*[42]. In sum, our

bioinformatic analyses revealed that homologs of sporulation-specific sigma factors are found in

diverse phages with contrasting lifestyles (i.e., lytic and temperate), often alongside sigma factors

that phages use to regulate lytic reproduction.

A sporulation-like sigma factor is non-essential for phage reproduction — Over time, a host-

derived sigma factor could be repurposed by phages for regulating phage genes that are essential

for reproductive functions like genome replication and capsid assembly. To test this hypothesis,

we used CRISPR-Cas9 to delete the sporulation-like sigma factor g120 from SP10, a phage that

infects *B. subtills* (see Figs. 2b and Table S1) and has overall 3 sigma factors (see above). After

removing the entire coding sequence of the sporulation-like sigma factor g120, this phage could

still productively infect its host, demonstrating that g120 was non-essential for phage reproduction

under standard lab conditions. In fact, there was no detectable reduction of virulence when

infecting *B. subtilis* with the mutant phage ($t_{12.9}$ = 1.22, P = 0.25; Fig. S4).

Phage-encoded sigma factors alter expression of host sporulation genes — To evaluate the

ability of phage-encoded homologs to activate transcription of host sporulation genes, we cloned

four diverse sporulation-like sigma factors (Fig. S5) from three *Bacillus* phages (Table S1) under

an IPTG inducible promoter into the chromosome of a spore-forming strain of *B. subtilis*. As a

control, we independently cloned the host-encoded sigma factors *sigF* and *sigG* into the same

strain of *B. subtilis* in a similar manner. We then induced the expression of each of the cloned

sigma factors during exponential growth, a time when native sporulation-specific sigma factors

(i.e., *sigF*, *sigG, sigE* and *sigK*) and other sporulation genes are not typically expressed (Fig. 3a).

Using RNAseq, we found that there was a strong positive correlation (ρ = 0.76) between

differential gene expression following induction of the sigma factor g169 derived from phage

Eldridge (ELDg169) and host-derived sigma factors (*sigF* and *sigG*) suggesting conservation of

gene function (Fig. 3b, S6). Similarly, when ELDg169 was induced, we observed the upregulation

of genes involved in sporulation (P <0.0001; Figs. 3c, S7). However, not all phage-derived sigma

factors affected host expression equally. For example, induction of the other sporulation-like sigma

factor cloned from phage Eldridge (ELDg168) also resulted in differential expression of many host

genes. However, the genes affected by ELDg168 were significantly different than those observed

in populations where host-derived sigma factors were induced (P < 0.0001; Fig. S6), and they were

not enriched in sporulation genes (Figs. 3c, S7). Meanwhile, induction of the two other less

conserved sporulation-like sigma factors (SP10 g120 and Goe3 g157) had only modest effects on

gene expression, with less than 50 genes differentially expressed by each induced gene, and no

enrichment of sporulation genes (Figs. 3c, S6, S7, S8).

Phage-encoded sigma factors inhibit sporulation — Induction of the sporulation-like sigma

factors altered spore yield in populations of *B. subtills* (Figs. 4, S9). When the host-derived sigma

factors were induced in sporulating cultures, we observed a reduction in spore yield, a pattern that

likely resulted from the misregulation of the sporulation gene network. Compared to an empty

vector control with no cloned sigma factor, induction of *sigF* led to an 85% reduction in spore

yield ($t_{8.2} = 6.43$, $P < 0.001$) while *sigG* reduced spore yield by 50% ($t_{13.9} = 6.43$, $P = 0.015$).

Expression of phage-derived sigma factors also affected sporulation, in one case to a greater degree

than host-derived sigma factors (Fig. 4, Table S2). Induction of ELDg169 reduced the spore yield

by 99% compared to the empty vector control ($t_7 = 7.8$, $P < 0.0001$), while the other Eldridge-

derived gene, ELDg168, had a smaller (~33%) and marginal effect on spore yield ($t_{15.1} = 2.1$, $P =$

0.064). The sigma factor from phage Goe3 reduced spore yield by $> 50\%$ ($t_{8.9} = 4.39$, $P = 0.002$),

while expression of the sigma factor from phage SP10 had no effect on spore yield ($t_{10.4} = 0.46$, P

$= 0.65$). Because spore yield was calculated as a percentage of the total population, we compared

cell counts between induced and non-induced controls (Fig. S10). From this, we concluded that

the observed reductions in spore yield were not due to a significant reduction in vegetative cells

(Table S3).

226   **DISCUSSION**

It is well known that some phages encode for sigma factors that regulate transcription during

lytic development[50,52,53]. In fact, early investigations of phage-encoded sigma factors elucidated

how the swapping of sigma subunits controls gene transcription by RNA polymerase[50]. However,

those phage-encoded sigma factors are quite divergent from bacteria-encoded sigma factors[53,59].

More recently, genomic data have identified phage-encoded sigma factors that bear a greater

resemblance to bacterial sigma factors, especially those involved in the regulation of sporulation

in the *Firmicutes*. While the function of these homologs remains unexplored, it is possible that

sporulation-like sigma factors have been coopted by phages to manipulate host dormancy in ways

that could potentially enhance their reproduction and survival. In support of this hypothesis, our

bioinformatic and phylogenetic analyses revealed that sporulation-like sigma factors were found

primarily in phages that infect spore-forming hosts, where they could potentially act as analogs of

the host's own sporulation-specific sigma factors. We show that these phage-encoded sigma

factors are distinct from genes known to be essential for regulating lytic reproduction and that they

are non-essential for phage reproduction. When expressed in *Bacillus subtilis,* phage-encoded

sigma factors reduced spore yield, likely owing to their ability to alter the expression of host genes,

which in one case led to transcriptional activation of the sporulation network. Together our

findings highlight novel ways in which dormancy may influence antagonistic coevolution between

spore-forming bacteria and their phages.

Diversity and distribution of phage-encoded sigma factors — Phages are able to regulate

gene expression in different ways. For example, they can make use of host-encoded sigma factors,

encode sigma factor analogs (e.g., gp33 in T4), or use their own RNA polymerases that do not

require sigma factors[50,53,60,61]. In addition, some phages encode their own sigma factors. Of the

nearly 3,500 genomes examined, we found that 14% of the phages possess a sigma factor. The

distribution of sigma factors in phage genomes was strongly influenced by host and viral

taxonomy. While common among phages that infect *Cyanobacteria* and *Proteobacteria*, only

among the *Firmicutes* phages did we recover sigma factors that were encoded by representatives

from all three of the major tailed-phage families (*Siphoviridae*, *Podoviridae* and *Myoviridae*; Fig.

1). This suggests that sigma factors may be beneficial to diverse phages that infect *Firmicutes*

hosts. Furthermore, we identified a host-specific pattern of gene multiplicity. A diverse set of

Firmicutes phages carried two sigma factors with host ranges that were limited to spore-forming

genera, while phages containing three sigma factors were restricted to just two viral families

(*Herelleviridae* and *Siphoviridae*) that could only infect strains of *Bacillus*. This convergent

pattern of gene multiplicity is associated with the presence of sporulation-like sigma factors in

phages that infect *Firmicutes*. Given the high degree of sequence similarity between sigma factors

encoded by phages and those encoded by their hosts, it seems likely that viruses have acquired

these genes from their hosts. Furthermore, because the genes are widespread among diverse viral

lineages, it seems reasonable to hypothesize that there are functional consequences for phage that

encode sporulation-like sigma factors. Nevertheless, in the database used in our study, these genes

are not universal in phages that infect spore-forming *Firmicutes*. We expect that future

investigations of metagenomes from environmental, engineered, and host-associated ecosystems

will shed light on the eco-evolutionary factors that influence the distribution and abundance of

dormancy related auxiliary genes, including sporulation-like sigma factors.

Sporulation-like sigma factors appear to be non-essential for phages — In bacteria, sporulation

and other specialized functions are controlled by alternative sigma factors that replace the primary

housekeeping sigma factor[51]. Because alternative sigma factors are only transiently used, their

deletion is typically non-lethal and thus they are considered non-essential. Similarly, multiple lines

of evidence suggest that sporulation-like sigma factors are non-essential for fundamental aspects

of phage biology. Unlike gp28 and gp34, which are required by phage SPO1, a sporulation-like

sigma factor (gp2.21) is known to be non-essential for lytic infection under standard laboratory

conditions[36]. While it has been hypothesized that gp2.21 directs transcription of phage genes from

sigK-like promoters found in its genome[36], this has not been confirmed. In our study, we deleted

the sporulation-like sigma factor from phage SP10 (g120) with no observable effect on virulence

(Fig. S4). Even though g120 is expressed during SP10 infection of its host[58], available evidence

suggests that at least some sporulation-like sigma factors are non-essential for lytic infection. If

this pattern is generalizable, then non-essentiality could reflect that phage-encoded sigma factors

have alternate function, which could involve the control of genes that are required for processes

other than replication.

Phage-encoded sigma factors disrupt regulation of the sporulation network — The prevalence

of sporulation-like sigma factors among diverse phages suggests that these genes may have

consequences for phage performance. One possibility is that these sigma factors play the same role

in phages as they do in their host, that is, they regulate the expression of sporulation genes. In

support of this notion, previous experimental studies have documented that phage-encoded

sporulation-like sigma factors retain a function that is relevant to host sporulation. For example,

in lysogenic *B. anthracis*, expression of a sporulation-like sigma factor encoded by the Wβ

prophage is elevated during sporulation[42]. Additionally, *in vitro* reconstitution of RNA polymerase

with the sporulation-like sigma factor of *B. anthracis* phage Fah, a close relative of Wβ, revealed

patterns of transcriptional activity and inhibition that are similar to that of a bacterial-encoded

sigma factor (*sigF*)[54]. Finally, sporulation-like sigma factors cloned from two diverse *B. anthracis*

phages, Bcp1 (*Herelleviridae*) and Wip4 (*Siphoviridae*), were associated with phage-dependent

inhibition of host sporulation[41].

In our study, the induction of phage-derived sporulation-like sigma factors altered host gene

expression and disrupted sporulation in *B. subtilis*. The magnitude of this effect was contingent

upon the identity and phylogenetic distance between the phage- and host-encoded sigma factors

(Fig, S11). When we induced ELDg169, the most host-like gene in our experiments, we observed

the differential expression of hundreds of bacterial genes, including the upregulation of nearly 400

sporulation genes (Fig. S7), which led to a ~99% reduction in spore yield (Fig. 4b). In contrast, we

observed a less pronounced reduction in spore yield, and almost no effect on the sporulation gene

network when two less conserved phage-derived genes (ELDg168 and Goe3 g157) were induced.

Last, induction of g120 from phage SP10 resulted in a very mild transcriptional response and had

no effect on spore yield, in accordance with it being the most divergent gene from bacterial *sigF*

and *sigG*. Thus, our studies suggest that that the functionality of phage-encoded sigma factors is

influenced by the degree of similarity with host-encoded sigma factors. However, this does not

rule out that phage-encoded sigma factors may play other roles depending on host and

environmental conditions. For example, the manipulation of sporulation seems to be the likely

function of these genes in phage Eldridge, possibly also in phage Goe3, but not in phage SP10.

As is the case with many evolutionary investigations, carefully planned experiments are

required to make strong inferences about the adaptive significance of genetic variation in a

population. We hypothesize that sporulation-like sigma factors could influence phage fitness in

different ways. By controlling the timing of host sporulation, phages may increase their probability

of entrapment[46], which would increase the survivorship component of phage fitness. On the other

hand, by impeding sporulation, there is more opportunity for virus replication, which would

increase the reproductive component of phage fitness. In our control experiments, induced

expression of bacterial genes activated the sporulation network during exponential growth.

However, when induced during sporulation, these host genes, which are known to promote

sporulation, led to a reduction in spore yield. This pattern likely stems from the regulatory genes

being expressed in the wrong place and time. As a consequence, it is challenging to conclusively

deduce whether the reduction in spore-yield reflects the function of phage proteins in infected cells

or other aspects of our experiment. However, results from one particular phage provide some

additional insight. The two Eldridge-derived genes, which are genomic neighbors, both reduced

sporulation to some degree, even though they had very different transcriptional profiles. If these

adjacent genes work toward a common function, it seems likely that this results in the inhibition

of host sporulation. This interpretation is consistent with the observation that complete inhibition

of sporulation in *B. anthracis* is likely mediated by a pair of sporulation-like sigma factors found

in tandem in the genome of phage Bcp1[41]. Additional experiments are needed to understand the

evolutionary consequences of sporulation-like sigma factors for phage populations. For example,

comparisons between phages in which sigma factors are present or deleted (like the SP10 mutant

we constructed) will be useful in determining whether auxiliary dormancy genes enhance phage

fitness. Likewise, comparative transcriptomic analysis during infection with phages with and

without sigma factors may help identify the regulatory targets of phage-encoded sigma factors.

Looking beyond sporulation, studies are needed to better understand how bacterial-like sigma

factors may be used by phages to manipulate other survival strategies that are common in non-

growing bacteria[62].

Taken together, our findings reveal a pattern of genomic convergence. Phages from diverse families, with contrasting infection strategies (virulent vs. temperate) that infect spore-forming hosts also contain sporulation-specific sigma factor homologs. Importantly, these sigma factors are phylogenetically distinct from sigma factors that regulate phage genes during lytic replication. At least some of these genes are non-essential for phages, which further supports the view that alternate sigma factors were not acquired for the purpose of regulating lytic programs. Instead, our experiments demonstrate that phage-encoded sigma factors retain features of their ancestral function, which means that they can be viewed as auxiliary genes that can regulate the sporulation network with consequences for spore yield. That being said, our results also reveal that some sporulation-like sigma factors in phages have variable and yet unknown functions, which may reflect neutral or adaptive divergence.

Eco-evolutionary implications of auxiliary dormancy genes — As obligate parasites, phages are unavoidably dependent on the metabolism of their bacterial hosts. Bacteria are capable of responding to their dynamic environments by replacing the sigma subunit of RNA polymerase, which leads to changes in gene expression[50,51]. Some phages use a similar strategy to coordinate expression of their own genes during different stages of infection[53]. Our analysis points to the existence of a second class of sigma factors that phages may use to manipulate host metabolism. Whether phages promote or inhibit sporulation, such manipulation of host dormancy has the potential to modify the environmental conditions under which we should expect to find bacteria in an active vs. dormant state. This has implications for development of novel therapeutic treatments

that combine phage therapy with antimicrobials, which tend to target metabolically active

bacteria[63].

Viruses have evolved to utilize, maintain, and rearrange a variety of biochemical pathways in

the cells that they take over[25,28,64]. The discovery of auxiliary metabolic genes has revealed that

viruses can appropriate the cellular building blocks and protein translation machinery of their

hosts. Our findings highlight an additional aspect of phage-host coevolution involving the co-

option of gene networks used to coordinate a complex and ancient form of dormancy, which many

microorganisms use to contend with harsh and unpredictable environments. Manipulation of the

host's response to such conditions through the acquisition of host regulatory genes could represent

a strategy which buffers viruses from the dynamic cellular environment on which their survival

and reproduction is dependent.

379   **METHODS**

Phage sigma factor distribution and classification — We retrieved sigma factors from the

database of viral orthologous groups (VOG release vog202, vogdb.org) based on text searches of

the VOG descriptors (Table S4). We matched VOG proteins to host and viral taxonomy using the

virus-host database[65]. We classified phage-encoded sigma factors using hmmscan with default

parameter settings using HMMER v3.3[66], and queried each protein against hidden Markov model

(HMM) profiles of bacterial-encoded sigma factor families that were retrieved from TIGRFAM

(Table S5). We used the best hmmscan match (smallest sequence E-value) to classify proteins,

unless it was a general TIGRFAM ("sigma70-ECF" or "SigBFG"), in which case the next best

match was chosen, if available.

Phylogenetic analysis — For phylogenetic analysis of sigma factors, we aligned VOG proteins

from phage genomes along with sigma factor proteins belonging to 24 bacteria from diverse taxa[67].

We aligned sigma factor sequences using MAFFT (v.7.475)[68] with the E-INS-I strategy and

trimmed the alignment with trimAL (v1.4.rev22)[69] using the gappyout method. From the 180

amino acids in the trimmed alignment we then inferred 200 maximum likelihood phylogenetic

trees using RAxML-NG (v0.9.0-pthreads)[70] with the LG+G4 substitution model selected using

modeltest-NG (v0.1.6)[71] with default settings. We present the best scoring maximum likelihood

tree with Transfer Bootstrap Expectation supports[72] from 500 bootstrapped trees. We plotted the

tree using the ggtreeExtra R package[73].

Strains and media — Strains used are listed in Table S1. For routine culturing of bacteria, we

used LB medium with low salt (5 g/L NaCl). We amended this recipe with agar (15 g/L) for plating

and with $CaCl_2$ (10 mM) to facilitate virus adsorption. We used Difco sporulation media (DSM)

for sporulation assays[74]. For plaque assays, we used double-layer plating with 0.3% agar

overlays[75]. To amplify phages, we collected lysates from plate infections after flooding Petri dishes

with phage buffer (10 mM Tris, 10 mM $MgSO_4$, 4 g/L NaCl, 1 mM $CaCl_2$, pH 7.5). We then

cleared the phage-containing buffer from bacteria by centrifugation (7,200 Xg, 10 min) and

filtration (0.2 μm).

Deletion of phage-encoded sigma factor — We used the CRISPR-Cas9 system and the CutSPR

assay design-tool[76] to test whether sigma factors are essential for phage replication. Briefly, we

cloned a single-guide RNA and a deletion cassette into plasmid pJOE8999[77] (Tables S6, S7) and

transformed the resulting plasmid into *B. subtilis* TS01 (Table S3), which was made competent

with D-mannitol induction. We next infected the transformed culture with phage SP10 (Table S1)

and conducted a plaque assay with medium containing the Cas9-inducer D-mannose. Using

primers SP10_validF+R (Table S7), we screened multiple plaques for the deletion. To isolate the

mutant phages, we picked and replated PCR-positive plaques onto host *B. subtilis* Δ6[78] (Table S1).

We screened these secondary plaques as above and confirmed the deletion by Sanger sequencing

of the locus. We then quantified the virulence of the mutant and wild-type phages[79]. After

dispensing *B. subtilis* Δ6 host cultures (OD600 = 1) into microtiter wells, we infected cells with

serially diluted lysates of SP10 or SP10 Δg120 that were adjusted to an equal titer. We monitored

bacterial density during growth for 16 h by OD600 with a Synergy H1 plate reader (Biotek). From

this, we calculated the virulence index[79] based on change in bacterial growth and lysis as a function

of the phage:bacteria ratio (i.e., multiplicity of infection; Fig S4).

Inducible expression of sigma factors — We tested the effect of phage-derived sigma factors on

bacterial expression by cloning coding sequences under an inducible promoter into an ectopic site

(*amyE*) of the *B. subtilis* genome. As a control, we also cloned host-derived sporulation genes

(s*igF* and *sigG*) in the same manner, and a gene-less promoter as a negative control.

Strain construction. We amplified coding sequences by PCR from phage lysates or from extracted

bacterial genomic DNA as templates using primers adapted with restriction sites, and a ribosome

binding site on the forward primer (Table S7). We then cloned the PCR products into plasmid

pDR110 (Table S6) by restriction enzyme digestion (Table S6), gel purification, and ligation (T4

ligase). We selected for plasmids that were transformed into *E. coli* (One Shot TOP10, Fisher)

with ampicillin (100 μg/ml) and verified the insertion by PCR and Sanger sequencing using

primers oDAS9+10 (Table S7). We transformed purified plasmids (QIAgen mini prep) into *B.*

subtilis TS01, as described above, using spectinomycin selection (100 μg/ml). We verified the

insertion into the *amyE* locus by PCR and Sanger sequencing, and by the loss of erythromycin

resistance carried in the *amyE* locus by strain TS01.

Transcriptional response to phage-encoded sigma factor. We diluted overnight *B. subtilis* cultures

(OD600 = 0.1) in fresh LB and grew them (37 °C, 200 RPM) to mid exponential phase (OD600 =

0.5). We then split the cultures and added 1mM IPTG (final concentration) to one half to induce

expression of the cloned gene. We added an equal volume of water to the other half of the split

culture as a non-induced control. After induction, we incubated the cultures for 2 h before

harvesting cells. Upon sampling, we immediately treated bacteria with the RNAprotect Bacteria

reagent (Qiagen), and stored pellets at -80 °C for <1 week before RNA extraction using RNeasy

Protect Bacteria Mini Kit (Qiagen) according to the manufacturer's instructions (protocol #5),

including an on-column RNase-free DNase digestion. Library construction, sequencing, and

analysis of differential gene expression were all carried out at the Indiana University Center for

Genomics and Bioinformatics. Libraries were constructed using the Illumina TruSeq Stranded

mRNA HT kit following depletion of rRNA using Illumina Ribo-Zero Plus kit. Libraries were then

sequenced on an Illumina NextSeq 500 platform as paired end reads (2 x 38 bp). We trimmed

adapters and filtered reads using Trimmomatic 0.38[80] with the cutoff threshold for average base

quality score set at 20 over a window of 3 bases. Reads shorter than 20 bases post-trimming were

excluded. We mapped the cleaned reads to the reference genome (Deposited with sequencing data

to the Gene Expression Omnibus, see below) using bowtie2 version 2.3.2[81], and counted reads

mapping concordantly and uniquely to the annotated genes using featureCounts tool ver. 2.0.0 of

subread package[82]. Read alignments to antisense strand, or to multiple regions on the genome or

those overlapping with multiple genes were ignored (parameters: -s 2 -p -B -C). We performed

differential expression analysis using DESeq2 ver. 1.24.0[83] from normalized read counts by

comparing samples induced with IPTG to non-induced paired control samples, with multiple-

testing correction. We tested for the effects of gene enrichment and overlap of differentially

expressed genes using the hypergeometric distribution in R[84].

Sporulation of cells expressing cloned sigma factors. To test for the effects of induced sigma

factors on host sporulation, we diluted overnight *B. subtilis* cultures in fresh DSM (OD600 = 0.05)

and dispensed each culture into multiple wells of a 96-well plate that was then incubated in a

Biotek Synergy H1 plate reader (37 °C, fast and continuous shake setting). Under these conditions,

we determined that cells enter stationary phase after approximately 4.5 h, marking the onset of

sporulation. At this time, we induced expression of the cloned gene by adding IPTG (final

concentration 1 mM) to half the cultures in the plate. We added water to the rest of the wells, which

served as non-induced controls. At 24 h, we quantified the number of spores and vegetative cells

in each well using a flow-cytometry assay that distinguished spores from vegetative cells (non-

spores) based on differential uptake of the nucleic acid stain SYBR green[85]. We diluted each

sample in TE buffer (pH 8) and then fixed the cells in 0.5% glutaraldehyde for 15 min at 4 °C. We

stained the fixed samples with SYBR green (20,000x dilution of commercial stock, Lonza) for 10

min at room temperature in the dark. We then enumerated cells using a volumetric NovoCyte

2000R flow cytometer (Acea; ex 488 nm, em 530/30 nm) and an automatic gating pipeline.

Code and data availability — All code and data used in the analyses in this study are available

at github.com/LennonLab/sigma-spore-phage and github.com/LennonLab/sigma-spore-phage-

flow. RNA sequencing data are available at the Gene Expression Omnibus under accession

number GSE187004. In addition, prior to publication, all data and code will be made available

on Zenodo.

ACKNOWLEDGEMENTS

We acknowledge R Hertel, L Temple, AC Hernandez, M Liu, X Wang, David Rudner, D Zeigler,

and DB Kearns for suggestions and strains; E Long and C Chen for technical support; and J Bird

for assistance with preliminary computation. An earlier version of the manuscript was improved

based on critical feedback from FJ Fishman, C Karakoç, A Magalie, JG McMullen, RZ Moger-

Reischer, EA Muller, PG Wall and JS Weitz. Research was supported by the National Science

Foundation (DEB-1934554 JTL and DAS, DBI- 2022049 JTL), US Army Research Office Grant

(W911NF-14-1-0411 JTL), the National Aeronautics and Space Administration

(80NSSC20K0618 JTL). This research was also supported in part by Lilly Endowment, Inc.,

through its support for the Indiana University Pervasive Technology Institute.

REFERENCES

1.   Lennon, J. T., den Hollander, F., Wilke-Berenguer, M. & Blath, J. Principles of seed
banks and the emergence of complexity from dormancy. *Nat. Commun.* **12**, 1-16 (2021).
2.   Lennon, J. T. & Jones, S. E. Microbial seed banks: the ecological and evolutionary
implications of dormancy. *Nat. Rev. Microbiol.* **9**, 119-130 (2011).
3.   Cáceres, C. E. Temporal variation, dormancy, and coexistence: a field test of the storage
effect. *Proc. Natl. Acad. Sci. USA* **94**, 9171-9175 (1997).
4.   Klobutcher, L. A., Ragkousi, K. & Setlow, P. The *Bacillus subtilis* spore coat provides
"eat resistance" during phagocytic predation by the protozoan *Tetrahymena thermophila*.
Proc. Natl. Acad. Sci. USA **103**, 165-170 (2006).

5. Verin, M. & Tellier, A. Host-parasite coevolution can promote the evolution of seed
banking as a bet-hedging strategy. *Evolution* **72**, 1362-1372 (2018).
6. Berleman, J. E. & Bauer, C. E. Characterization of cyst cell formation in the purple
photosynthetic bacterium *Rhodospirillum centenum*. *Microbiology* **150**, 383-390 (2004).
7. Driks, A. & Eichenberger, P. in *The Bacterial Spore: From Molecules to Systems*   179-
200 (American Society of Microbiology, 2016).
8. Kaplan-Levy, R. N., Hadas, O., Summers, M. L., Rücker, J. & Sukenik, A. in *Dormancy*
and resistance in harsh environments Topics in Current Genetics (eds Esther Lubzens,
Joan Cerdà, & Melody Clark) 5-27 (Springer, Berlin, Heidelberg, 2010).
9. Burroughs, N. J., Marsh, P. & Wellington, E. M. H. Mathematical analysis of growth and
interaction dynamics of streptomycetes and a bacteriophage in soil. *Appl. Environ.*
Microbiol. **66**, 3868-3877 (2000).
10. Dowding, J. Characterization of a bacteriophage virulent for *Streptomyces coelicolor* A3
(2). *Microbiology* **76**, 163-176 (1973).
11. Singh, R. N. & Singh, P. K. Isolation of Cyanophages from India. *Nature* **216**, 1020-1021
(1967).
12. Gabiatti, N. *et al.* Bacterial endospores as phage genome carriers and protective shells.
Appl. Environ. Microbiol. **84**, e01186 (2018).
13. Hadas, H., Einav, M., Fishov, I. & Zaritsky, A. Bacteriophage T4 development depends
on the physiology of its host *Escherichia coli*. *Microbiology* **143 (Pt 1)**, 179-185 (1997).
14. Middelboe, M. Bacterial growth rate and marine virus–host dynamics. *Microb. Ecol.* **40**,
114-124 (2000).
15. Abedon, S. T. & Yin, J. in *Bacteriophages: Methods and Protocols, Volume 1: Isolation,*
Characterization, and Interactions   (eds Martha R. J. Clokie & Andrew M. Kropinski)
161-174 (Humana Press, 2009).
16. Bryan, D., El-Shibiny, A., Hobbs, Z., Porter, J. & Kutter, E. M. Bacteriophage T4
infection of stationary phase *E. coli*: life after log from a phage perspective. *Front*
Microbiol **7** (2016).
17. Makarova, K. S., Anantharaman, V., Aravind, L. & Koonin, E. V. Live virus-free or die:
coupling of antivirus immunity and programmed suicide or dormancy in prokaryotes.
Biol Direct **7**, 7-40 (2012).
18. Koonin, E. V. & Zhang, F. Coupling immunity and programmed cell suicide in
prokaryotes: Life-or-death choices. *Bioessays* **39**, e201600186 (2017).
19. Meeske, A. J., Nakandakari-Higa, S. & Marraffini, L. A. Cas13-induced cellular
dormancy prevents the rise of CRISPR-resistant bacteriophage. *Nature* **570**, 241-245
(2019).
20. Bautista, M. A., Zhang, C. & Whitaker, R. J. Virus-Induced dormancy in the archaeon
Sulfolobus islandicus. *mBio* **6**, e02565 (2015).
21. Lenski, R. E. in *Adv. Microb. Ecol.* Vol. 10 (ed K. C. Marshall) 1-44 (Springer, 1988).
22. Lopatina, A., Tal, N. & Sorek, R. Abortive infection: bacterial suicide as an antiviral
immune strategy. *Annu Rev Virol* **7**, 371-384 (2020).
23. van Houte, S., Buckling, A. & Westra, E. R. Evolutionary ecology of prokaryotic
immune mechanisms. *Microbiol. Mol. Biol. Rev.* **80**, 745-763 (2016).
24. Ofir, G. & Sorek, R. Contemporary phage biology: from classic models to new insights.
Cell **172**, 1260-1270 (2018).

25.   Thompson, L. R. *et al.* Phage auxiliary metabolic genes and the redirection of
554   cyanobacterial host carbon metabolism. *Proc. Natl. Acad. Sci. USA* **108**, E757-E764
555   (2011).
556   26.   Lindell, D. *et al.* Transfer of photosynthesis genes to and from *Prochlorococcus* viruses.
557   *Proc. Natl. Acad. Sci. USA* **101**, 11013-11018 (2004).
558   27.   Mann, N. H., Cook, A., Millard, A., Bailey, S. & Clokie, M. Bacterial photosynthesis
559   genes in a virus. *Nature* **424**, 741-741 (2003).
560   28.   Hargreaves, K. R., Kropinski, A. M. & Clokie, M. R. Bacteriophage behavioral ecology:
561   How phages alter their bacterial host's habits. *Bacteriophage* **4**, e29866-e29866 (2014).
562   29.   Silpe, J. E. & Bassler, B. L. Phage-encoded LuxR-type receptors responsive to host-
563   produced bacterial quorum-sensing autoinducers. *mBio* **10**, e00638-00619 (2019).
564   30.   Mizuno, C. M. *et al.* Numerous cultivated and uncultivated viruses encode ribosomal
565   proteins. *Nat. Commun.* **10**, 752 (2019).
566   31.   Seed, K. D., Lazinski, D. W., Calderwood, S. B. & Camilli, A. A bacteriophage encodes
567   its own CRISPR/Cas adaptive response to evade host innate immunity. *Nature* **494**, 489-
568   491 (2013).
569   32.   Murphy, J. *et al.* Methyltransferases acquired by lactococcal 936-type phage provide
570   protection against restriction endonuclease activity. *BMC Genomics* **15**, 1-11 (2014).
571   33.   Dragoš, A. *et al.* Pervasive prophage recombination occurs during evolution of spore-
572   forming Bacilli. *ISME J* **15**, 1344-1358 (2021).
573   34.   Reveille, A. M., Eldridge, K. A. & Temple, L. M. Complete genome sequence of *Bacillus
574   megaterium* bacteriophage Eldridge. *Genome Announc.* **4**, e0172815 (2016).
575   35.   Ritz, M. P., Perl, A. L., Colquhoun, J. M., Chamakura, K. R. & Everett, G. F. K.
576   Complete genome of *Bacillus subtilis* myophage CampHawk. *Microbiol Resour Announc*
577   **1**, e00984-00913 (2013).
578   36.   Stewart, C. R. *et al.* The genome of *Bacillus subtilis* bacteriophage SPO1. *J. Mol. Biol.*
579   **388**, 48-70 (2009).
580   37.   Van Goethem, M. W., Swenson, T. L., Trubl, G., Roux, S. & Northen, T. R.
581   Characteristics of wetting-induced bacteriophage blooms in biological soil crust. *mBio*
582   **10**, e02287-02219 (2019).
583   38.   Yuan, Y., Gao, M., Wu, D., Liu, P. & Wu, Y. Genome characteristics of a novel phage
584   from *Bacillus thuringiensis* showing high similarity with phage from *Bacillus cereus*.
585   *PLoS One* **7**, e37557 (2012).
586   39.   El-Arabi, T. F. *et al.* Genome sequence and analysis of a broad-host range lytic
587   bacteriophage that infects the *Bacillus cereus* group. *Virol J* **10**, 1-11 (2013).
588   40.   Zimmer, M., Scherer, S. & Loessner, M. J. Genomic analysis of *Clostridium perfringens*
589   bacteriophage φ3626, which integrates into guaA and possibly affects sporulation. *J.
590   Bacteriol.* **184**, 4359-4368 (2002).
591   41.   Schuch, R. & Fischetti, V. A. The secret life of the anthrax agent Bacillus anthracis:
592   bacteriophage-mediated ecological adaptations. *PLoS One* **4** (2009).
593   42.   Schuch, R. & Fischetti, V. A. Detailed genomic analysis of the Wβ and γ phages
594   infecting *Bacillus anthracis*: implications for evolution of environmental fitness and
595   antibiotic resistance. *J. Bacteriol.* **188**, 3037-3051 (2006).
596   43.   Sonenshein, A. L. Trapping of unreplicated phage DNA into spores of *Bacillus subtilis*
597   and its stabilization against damage by ^{32}P decay. *Virology* **42**, 488-495 (1970).

598   44.   Sonenshein, A. L. Bacteriophages: How bacterial spores capture and protect phage DNA.
599   *Curr. Biol.* **16**, R14-R16 (2006).
600   45.   Takahashi, I. Incorporation of bacteriophage genome by spores of *Bacillus subtilis*. *J.*
601   *Bacteriol.* **87**, 1499-1502 (1964).
602   46.   Sonenshein, A. L. & Roscoe, D. H. The course of phage Øe infection in sporulating cells
603   of *Bacillus subtilis* strain 3610. *Virology* **39**, 265-276 (1969).
604   47.   Galperin, M. Y. *et al.* Genomic determinants of sporulation in *Bacilli* and *Clostridia*:
605   towards the minimal set of sporulation-specific genes. *Environ. Microbiol.* **14**, 2870-2890
606   (2012).
607   48.   Meeske, A. J. *et al.* High-throughput genetic screens identify a large and diverse
608   collection of new sporulation genes in *Bacillus subtilis*. *PLoS Biol.* **14**, e1002341 (2016).
609   49.   Ramos-Silva, P., Serrano, M. & Henriques, A. O. From root to tips: sporulation evolution
610   and specialization in *Bacillus subtilis* and the intestinal pathogen *Clostridioides difficile*.
611   *Mol. Biol. Evol.* **36**, 2714-2736 (2019).
612   50.   Helmann, J. D. Where to begin? Sigma factors and the selectivity of transcription
613   initiation in bacteria. *Mol. Microbiol.* **112**, 335-347 (2019).
614   51.   Paget, M. S. Bacterial sigma factors and anti-sigma factors: structure, function and
615   distribution. *Biomolecules* **5**, 1245-1265 (2015).
616   52.   Losick, R. & Pero, J. Cascades of sigma factors. *Cell* **25**, 582-584 (1981).
617   53.   Nechaev, S. & Severinov, K. Bacteriophage-induced modifications of host RNA
618   polymerase. *Annu. Rev. Microbiol.* **57**, 301-322 (2003).
619   54.   Minakhin, L. *et al.* Genome sequence and gene expression of *Bacillus anthracis*
620   bacteriophage Fah. *J. Mol. Biol.* **354**, 1-15 (2005).
621   55.   Wörmer, L. *et al.* Microbial dormancy in the marine subsurface: Global endospore
622   abundance and response to burial. *Sci Adv* **5**, eaav1024 (2019).
623   56.   Barylski, J. *et al.* Analysis of spounaviruses as a case study for the overdue
624   reclassification of tailed phages. *Syst. Biol.* **69**, 110-123 (2020).
625   57.   Swanson, M. M. *et al.* Novel bacteriophages containing a genome of another
626   bacteriophage within their genomes. *PLoS One* **7**, e40683 (2012).
627   58.   Yee, L. M. *et al.* The genome of *Bacillus subtilis* phage SP10: a comparative analysis
628   with phage SPO1. *Biosci., Biotechnol., Biochem.* **75**, 944-952 (2011).
629   59.   Lonetto, M., Gribskov, M. & Gross, C. A. The sigma 70 family: sequence conservation
630   and evolutionary relationships. *J. Bacteriol.* **174**, 3843-3849 (1992).
631   60.   Twist, K.-A. F. *et al.* Crystal structure of the bacteriophage T4 late-transcription
632   coactivator gp33 with the β-subunit flap domain of *Escherichia coli* RNA polymerase.
633   *Proc. Natl. Acad. Sci. USA* **108**, 19961-19966 (2011).
634   61.   Sokolova, M. *et al.* A non-canonical multisubunit RNA polymerase encoded by the AR9
635   phage recognizes the template strand of its uracil-containing promoters. *Nucleic Acids*
636   *Res.* **45**, 5958-5967 (2017).
637   62.   Jaishankar, J. & Srivastava, P. Molecular basis of stationary phase survival and
638   applications. *Front Microbiol* **8**, 2000 (2017).
639   63.   Rodriguez-Gonzalez, R. A., Leung, C. Y., Chan, B. K., Turner, P. E. & Weitz, J. S.
640   Quantitative models of phage-antibiotic combination therapy. *mSystems* **5**, e0075619
641   (2020).
642   64.   Breitbart, M., Bonnain, C., Malki, K. & Sawaya, N. A. Phage puppet masters of the
643   marine microbial realm. *Nat Microbiol* **3**, 754-766 (2018).

65. Mihara, T. *et al.* Linking virus genomes with host taxonomy. *Viruses* **8**, 66 (2016).

66. Eddy, S. R. Accelerated profile HMM searches. *PLoS Comp. Biol.* **7**, e1002195 (2011).

67. Burton, A. T., DeLoughery, A., Li, G.-W. & Kearns, D. B. Transcriptional regulation and
mechanism of SigN (ZpdN), a pBS32-encoded sigma factor in *Bacillus subtilis*. *mBio* **10**,
e0189919 (2019).

68. Katoh, K. & Standley, D. M. MAFFT multiple sequence alignment software version 7:
improvements in performance and usability. *Mol. Biol. Evol.* **30**, 772-780 (2013).

69. Capella-Gutiérrez, S., Silla-Martínez, J. M. & Gabaldón, T. trimAl: a tool for automated
alignment trimming in large-scale phylogenetic analyses. *Bioinformatics* **25**, 1972-1973
(2009).

70. Kozlov, A. M., Darriba, D., Flouri, T., Morel, B. & Stamatakis, A. RAxML-NG: a fast,
scalable and user-friendly tool for maximum likelihood phylogenetic inference.
Bioinformatics **35**, 4453-4455 (2019).

71. Darriba, D. *et al.* ModelTest-NG: a new and scalable tool for the selection of DNA and
protein evolutionary models. *Mol. Biol. Evol.* **37**, 291-294 (2020).

72. Lemoine, F. *et al.* Renewing Felsenstein's phylogenetic bootstrap in the era of big data.
Nature **556**, 452-456 (2018).

73. Xu, S. *et al.* ggtreeExtra: Compact visualization of richly annotated phylogenetic data.
Mol. Biol. Evol. **38**, 4039-4042 (2021).

74. Harwood, C. R. & Cutting, S. M. *Molecular biological methods for Bacillus*. (Wiley,
1990).

75. Kauffman, K. M. & Polz, M. F. Streamlining standard bacteriophage methods for higher
throughput. *MethodsX* **5**, 159-172 (2018).

76. Schilling, T., Dietrich, S., Hoppert, M. & Hertel, R. A CRISPR-Cas9-based toolkit for
fast and precise in vivo genetic engineering of *Bacillus subtilis* phages. *Viruses* **10**, 241
(2018).

77. Altenbuchner, J. Editing of the *Bacillus subtilis* genome by the CRISPR-Cas9 system.
Appl. Environ. Microbiol. **82**, 5421-5427 (2016).

78. Westers, H. *et al.* Genome engineering reveals large dispensable regions in *Bacillus
subtilis*. *Mol. Biol. Evol.* **20**, 2076-2090 (2003).

79. Storms, Z. J., Teel, M. R., Mercurio, K. & Sauvageau, D. The virulence index: a metric
for quantitative analysis of phage virulence. *Phage (New Rochelle)* **1**, 27-36 (2020).

80. Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina
sequence data. *Bioinformatics* **30**, 2114-2120 (2014).

81. Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat. Methods*
9, 357-359 (2012).

82. Liao, Y., Smyth, G. K. & Shi, W. featureCounts: an efficient general purpose program for
assigning sequence reads to genomic features. *Bioinformatics* **30**, 923-930 (2014).

83. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion
for RNA-seq data with DESeq2. *Genome Biol* **15**, 1-21 (2014).

84. R: A Language and Environment for Statistical Computing (R Foundation for Statistical
Computing, Vienna, Austria, 2021).

85. Karava, M., Bracharz, F. & Kabisch, J. Quantification and isolation of *Bacillus subtilis*
spores using cell sorting and automated gating. *PLoS One* **14**, e0219892 (2019).

86.   Zhu, B. & Stülke, J. Subti Wiki in 2018: from genes and proteins to functional network
689   annotation of the model organism *Bacillus subtilis*. *Nucleic Acids Res.* **46**, D743-D748
690   (2018).
691

 FIGURE CAPTIONS

Fig 1. Sigma factors in viral genomes. a-c, The overall number of phage genomes in each

taxonomic group is depicted by light grey bars with nested darker bars depicting how many of

those genomes contained one or more sigma factors. Taxonomic groups are host phylum (**a**),

viral family (**b**), and viral families containing sigma factors separated by host phyla (**c**). **d**,

Multiplicity of sigma factors genes in phage genomes. The distribution of sigma factor gene

counts in phage genomes is shown for all phages in the Viral Orthologous Groups (VOG)

database (left facet), and for each of the host phyla as in **c**. Phage genomes from the VOG

database were mapped to hosts using the Virus-Host Database. Viral families are specified for

tailed phages (*Caudovirales*), and all non-tailed phages are shown in a single category of non-

Caudovirales, with an asterisk to indicate that this is not a true viral family. The two

Bacteroidetes phages with sigma factors indicated in **a**, each from a different viral family are

excluded from **c** for clarity.

Fig 2. Phylogeny and classification of phage-encoded sigma factors. a, Phylogenetic tree

representing proteins of phage-encoded sigma factor from the database of viral orthologous

groups. For reference, we included sigma factors from 24 genomes of diverse bacterial species.

Blue branches on the tree correspond with monophyletic clades that only contain phage-encoded

sigma factors. Black branches show bacterial-encoded sigma factors and the internal branches

leading to them. Phage proteins discussed in the text are labeled at branch tips (e.g., ELD

gp128). In circle 1 (C1) we identify the source (phage vs. bacteria) of a sigma factor protein. In

circle 2 (C2), we designate the taxonomy of the bacterial host (phyla). In circle 3 (C3) we

provide the best hit (hmmscan sequence E-value) of the homolog to bacterial-encoded sigma

factors families in TIGRfam. Outside of circle 3, we indicate the relative positions of *Bacillus*

subtilis sigma factor genes and depict the different sigma factor families (grey wedges). Branch

nodes are labeled with nonparametric bootstrap support values (n = 500). **b**, Clade of

sporulation-specific sigma factors depicted by red arc in **a**. Triangles point to genes cloned and

expressed in this study. **c**, Summarized proportions of phage-encoded sigma factors as a function

of host taxonomy and protein family as described for circles 2 and 3 of panel **a***,* respectively.

Fig 3. Bacterial gene expression following induction of phage-encoded sigma factors. a,

Illustration of the experimental design. Phage and host derived sigma factors were cloned in

Bacillus subtilis under an IPTG-inducible promoter (grey arrow) in a strain (Δ6) that contains a

fully functional sporulation gene network, including its native sporulation specific sigma factors

(black arrows show representative examples). To estimate differential expression, we compared

splits of *Bacillus subtilis* cultures. One half was induced to express a cloned sigma factor during

exponential growth, while the other half used as a non-induced control (n = 3). Differentially

expressed genes were defined as those where *P* values < 0.05 (horizonal dashed line in **b**) and

|fold change| > 2 (outside of vertical grey bar in **b**). **b***,* Correlation of differentially expressed

genes for cells induced to express phage-encoded sigma factors and cells induced to express

bacteria-encoded sigma factors (*sigF* and *sigG*). Spearman's correlation coefficient (ρ) is

displayed. **c**, Sporulation genes[86] were upregulated after induction of a phage-encoded sigma

factor (ELDg169) and host-encoded sigma factors that regulate sporulation (*sigF* and *sigG*). The

number of significantly upregulated ($\triangle$) and downregulated ($\triangledown$) genes are noted.

Fig. 4. Phage-derived sigma factors reduced sporulation. a, Illustration of the experimental design. Phage and host derived sigma factors were cloned in *Bacillus subtilis* under an IPTG-inducible promoter (grey arrow) in a strain (Δ6) that contains a fully functional sporulation gene network, including its native sporulation specific sigma factors (black arrows show representative examples). We compared sporulation in these strains to an empty-vector negative control strain that also had an IPTG-inducible promoter. For each data point in **b** replicate cultures (n = 6) of a single colony were grown in sporulation medium for 4.5 hours, a time at which sporulation was induced by nutrient exhaustion. We then induced expression of the cloned sigma factor by the addition of IPTG to half of the replicates, while leaving the remainder as controls. Spore and vegetative cells were quantified by flow-cytometry. Spore yield was calculated as the ratio of percent spores in induced cultures and their paired controls. **b**, Bars represent the mean ± SEM of independent clones (n ≥ 8). Shapes represent different experimental batches. Asterisks correspond to significance levels (adjusted for multiple testing) from Welch's *t*-tests used to evaluate the effect of each sigma factor compared to the empty-vector negative control (* = $P < 0.05$, ** = $P < 0.01$, *** = $P < 0.001$). ELD = phage Eldridge.

752   **Fig. 1**

Fig. 2

Fig. 3

759   **Fig. 4.**

760