An integrated resource for functional and structural connectivity of the marmoset brain

Xiaoguang Tian[1,*], Yuyan Chen[2], Piotr Majka[3,4], Diego Szczupak[1], Yonatan Sanz Perl[5,6], Cecil Chern-Chyi Yen[7], Chuanjun Tong[2], Furui Feng[2], Haiteng Jiang[8,9], Daniel Glen[10], Gustavo Deco[5,11,12,13], Marcello G. P. Rosa[4, *], Afonso C. Silva[1,*], Zhifeng Liang[2, *], Cirong Liu[2,14,15,16, *]

[1] Department of Neurobiology, University of Pittsburgh Brain Institute, University of Pittsburgh, Pittsburgh PA 15261, USA
[2] Center for Excellence in Brain Science and Intelligence Technology, Institute of Neuroscience, CAS Key Laboratory of Primate Neurobiology, Chinese Academy of Sciences, Shanghai, China
[3] Laboratory of Neuroinformatics, Nencki Institute of Experimental Biology of the Polish Academy of Sciences, 02-093 Warsaw, Poland
[4] Department of Physiology and Neuroscience Program, Biomedicine Discovery Institute, Monash University, Clayton, VIC 3800, Australia
[5] Center for Brain and Cognition, Computational Neuroscience Group, Department of Information and Communication Technologies, Universitat Pompeu Fabra, Roc Boronat 138, Barcelona, 08018, Spain
[6] Universidad de San Andrés, Vito Dumas 284 (B1644BID), Buenos Aires, Argentina
[7] Cerebral Microcirculation Section, Laboratory of Functional and Molecular Imaging, National Institute of Neurological Disorders and Stroke, National Institutes of Health (NINDS/NIH), Bethesda, MD 20892, USA
[8] Department of Neurobiology, Affiliated Mental Health Center & Hangzhou Seventh People's Hospital, Zhejiang University School of Medicine
[9] MOE Frontier Science Center for Brain Science and Brain-machine Integration, Zhejiang University, Hangzhou, China.
[10] Scientific and Statistical Computing Core, National Institute of Mental Health, National Institutes of Health (NIMH/NIH), Bethesda, MD 20892, USA
[11] Institució Catalana de la Recerca i Estudis Avançats (ICREA), Passeig Lluís Companys 23, Barcelona, 08010, Spain
[12] Department of Neuropsychology, Max Planck Institute for Human Cognitive and Brain Sciences, 04103 Leipzig, Germany
[13] School of Psychological Sciences, Monash University, Melbourne, Clayton VIC 3800, Australia
[14] Shanghai Center for Brain Science and Brain-Inspired Intelligence Technology, Shanghai, China
[15] Lingang Laboratory, Shanghai 200031, China
[16] University of Chinese Academy of Sciences, Beijing, China

* Co-Corresponding authors

Cirong Liu crliu@ion.ac.cn
Xiaoguang Tian txgxp88@gmail.com
Afonso Silva AFONSO@pitt.edu
Zhifeng Liang zliang@@ion.ac.cn
Marcello Rosa marcello.rosa@monash.edu

Abstract

Comprehensive integration of structural and functional connectivity data is required to model brain functions accurately. While resources for studying the structural connectivity of non-human primate brains already exist, their integration with functional connectivity data has remained unavailable. Here we present a comprehensive resource that integrates the largest awake marmoset resting-state fMRI data available to date (39 marmoset monkeys, 710 runs, 12117 mins) with previously published cellular-level neuronal tracing (52 marmoset monkeys, 143 injections) and multi-resolution diffusion MRI datasets. The combination of these data allowed us to (1) map the fine-detailed functional brain networks and cortical parcellations; (2) develop a deep-learning-based parcellation generator that preserves the topographical organization of functional connectivity and reflects individual variabilities, and (3) investigate the structural basis underlying functional connectivity by computational modeling. This resource will enable modeling structure-function relationships and facilitate future comparative and translational studies of primate brains.

Introduction

Mapping brain architecture is critical for decoding brain functions and understanding the mechanisms of brain diseases [1]. Non-human primate (NHP) neuroimaging provides a granular view of the evolution of the brain [2] and could overcome constraints of human neuroimaging by integration with "ground truth" data from cellular-resolution tracing [3].

As one of the few non-invasive imaging techniques capable of mapping whole-brain functional activity patterns, resting-state fMRI (rs-fMRI) provides insights into large-scale functional architecture [4]. However, data-sharing initiatives of NHP neuroimaging are still at an early stage, with existing open datasets of rs-fMRI data originating in different laboratories and collected for different purposes [5]. This leads to inconsistent imaging protocols and data quality, which hinder analyses across datasets. In addition, most presently available rs-fMRI datasets have been acquired in anesthetized animals, resulting in difficulties for cross-species studies, particularly relative to awake human brains [6]. The final barrier is the practical difficulty of training large numbers of NHPs to be fully awake during MRI scans [7, 8]. A platform for international collaborative research (PRIMatE RESOURCE EXCHANGE) was initiated to address these problems and promote open resource exchange and standards for NHP neuroimaging [5, 9].

The common marmoset monkey (Callithrix jacchus) has drawn considerable interest as an NHP species, offering many practical advantages for neuroscience research, including neuroimaging [10, 11, 12]. Previous work from our groups has contributed ultra-high-resolution ex-vivo diffusion MRI data [13], mesoscale neural tracing data [14], and structural atlases [15, 16, 17], which have enabled unprecedented precision in analyses of NHP brain anatomy. However, an essential component for understanding brain architecture has been missing: integrating these anatomical datasets with rs-fMRI. To address this limitation, and in alignment with a strategic plan developed by the NHP imaging community [8], we developed standardized protocols for imaging awake marmosets, which were implemented across two institutions, the National Institutes of Health (NIH), USA, and the Institute of Neuroscience (ION), China. This effort resulted

97 in the largest awake NHP rs-fMRI dataset to date, which is being made available

98 through an open-access platform. Furthermore, we integrated neuronal tracing and

99 different diffusion MRI datasets into the same MRI space, resulting in a comprehensive

100 resource that allows us to explore the relationships between the structural and

101 functional connectomes by computational modeling.

Results

The resource reported in this paper (summary in Fig. 1) is supported by a publicly

available standardized dataset. Following the same protocols for animal training and

MRI imaging, including the designs of the radiofrequency coil and MRI pulse sequences,

we acquired an extensive awake resting-state fMRI dataset to date from 39 marmosets

of two research institutes (13 from ION, age 3+/-1 years old; 26 from NIH, age 4+/-2

years old; 12117 mins in total scanning, Supplementary Table 1 for details). This is also

the same range of ages used in our previous studies of structural connectivity [13, 14]. For

test-retest evaluation, we scanned multiple runs (17 mins/run) for each marmoset,

resulting in an essentially similar data quantity of two institutes (346 ION runs and 364

NIH runs) and included two "flagship" marmosets with many runs (64 runs from the ION

and 40 runs from the NIH). Besides similar quantity, we comprehensively evaluated the

data quality across two datasets, which is essential for data harmonization [18]. We

compared different metrics in the SNR, temporal SNR (tSNR), spatial CNR, and head

motions across the sites/scanners (Supplementary Fig. 1-3, Supplementary Table 2).

We found no significant differences between the two datasets. We also compared the

similarity of functional connectivity across subjects (Supplementary Fig. 4-A) and

sessions (Supplementary Fig. 4-B), which still revealed no significant differences. The

quality assessments and quality control (QA/QC) measurements demonstrated

consistency and interpretability across datasets, which makes them suitable for further

analysis.

Using these datasets, we created a comprehensive map of resting-state brain networks
and a fine-grained functional cortical parcellation based on resting-state functional
connectivity. Furthermore, we developed a deep-learning-based approach to map the
population-based functional cortical parcellation onto individual brains. This allowed
investigation of the structural basis underlying functional connectivity. For this purpose,
we sampled the most extensive collection of NHP neuronal tracing data available (52
marmosets and 143 injections) onto the same MRI space at the voxel or vertex level
and integrated it with the same functional MRI data space mentioned above. In addition,
further enhancing the capacity of our resource, we also integrated extra high-resolution
ex-vivo diffusion MRI and in-vivo diffusion MRI data obtained at 25 marmosets from the
same cohort. On this basis, we investigated the relationship between structural and
functional connectivity using a whole-brain computational model. To facilitate the user to
explore connectomes reported in this paper and compare them with other connectomes
[13, 14, 15], we make online connectome viewers (connectome.marmosetbrainmapping.org).
140

Mapping functional brain networks

Identifying functional networks of areas showing highly correlated fMRI signals is critical
to characterizing the brain architecture. Using the independent component analysis
(ICA), a data-driven approach for separating independent patterns in multivariate data,
we identified 15 cortical networks from awake resting-state fMRI data (Fig. 2;
Supplementary Figs. 5-6 include the power spectrum). All specified components
showed clear neural-like patterns spatially (all peaks located in the cortical gray matter)
and temporally (no patterns of artifacts or noises), as shown in Supplementary Fig. 6.
We also conducted the ICA separately on each dataset. We found that both the ION
and the NIH data reveal these components with similar spatial patterns and temporal
CNR (tCNR, the definition of this metric is demonstrated in the method part of Data
harmonization, the results are in Supplementary Fig. 7), demonstrating the
reproducibility of networks and the consistency of the two datasets.
155

The details of the 15 cortical networks were as follows. Six functional networks were characterized by short-range connectivity, including the ventral somatomotor (Fig. 2A), the dorsal somatomotor (Fig. 2B), the premotor (Fig. 2C), the frontopolar (Fig. 2D), the orbitofrontal (Fig. 2E), and the parahippocampal/ temporopolar cortex (Fig. 2F) networks. The next two components are the auditory and salience-related networks, the first being primarily located in the auditory and insular cortices, and weakly coupled with the anterior cingulate cortex (Fig. 2G), and the second (Fig. 2H) encompassing the anterior cingulate cortex. In addition, we also identified two trans-modal networks (Fig. 2I-J), including association areas in the dorsolateral prefrontal cortex (dlPFC), rostral premotor cortex, lateral and medial parietal cortices, and temporal cortex. One of these is most likely the frontoparietal-like network (Fig. 2I), which has not been identified in previous studies [19, 20], and the other is the default-mode network (DMN, Fig. 2J), which we characterized extensively in a previous study [21]. The remaining five networks represent the first complete mapping of visual-related functional networks of the marmoset cortex (Fig. 2K-O). Three networks included the primary visual cortex and parts of extrastriate areas related to far peripheral vision (Fig. 2K), near-peripheral vision (Fig. 2L), and foveal vision (Fig. 2M). The other two networks involve hierarchically higher visual areas (Fig. 2N-O), such as V3, V4, the inferior temporal cortex, the adjacent polysensory temporal cortex, and vison-related frontal regions.

Based on their spatial overlap patterns and connectivity strengths (normalized Z-scores), we combined the 15 cortical networks into network-parcellation maps (Fig. 2P-Q). Due to local connectivity being stronger than long-range connectivity, the primary map (Fig. 2P-Q, top rows) is dominated by the short-range networks (i.e., Fig. 2G, I, J, K, L, I, and O). Thus, we created a second connectivity map (Fig. 2P-Q, bottom rows) to cover the long-range connectivity not captured by the primary map. The two network-parcellation maps characterized the entire cortical networks, and will likely be of value for future functional connectivity studies of the marmoset brain.

Mapping functional connectivity boundaries

The brain network maps provided a global view of cortical functional organization. Next,

we aimed to characterize the cortex at a finer local scale. For this, we used a functional

connectivity boundary mapping approach to identify putative borders for the functional

parcels [22, 23, 24] as an efficient way to map transitions in functional connectivity.

Population boundary maps based on the ION, the NIH, or combined datasets are

visually similar, presenting clear functional connectivity borders (Fig. 3A), and were

highly reproducible with average Dice's coefficients for both hemispheres: 0.7 (ION-NIH),

0.71 (ION-Both), and 0.69 (NIH-Both), respectively (Supplementary Fig. 8). However,

although consistent at the population level, boundary maps indicate variability across

individuals (Fig. 3B), with an average Dice's coefficient of 0.3842 for both hemispheres

(Fig. 3C-D), which is significantly lower than those for the population. We also found

across-session variability in the same individual, but more scanning runs efficiently

enhanced the reproducibility (Fig. 3E). Therefore, these results suggest that both

individual and across-session variability contribute to the low consistency of individual

boundary maps and that using test-retest data is essential for improving the reliability of

maps.

Generation of functional connectivity parcels (Marmoset Brain Mapping Atlas

Version 4, MBMv4)

Because the population boundary maps are more reproducible than individual maps, we

used the combined ION-NIH population boundary map to generate cortical functional

connectivity parcels. By the detection of the local-minima [22], "watershed-flood" region

growing [25], and semi-manual optimization of parcel boundaries (Fig. 4A), we obtained

96 parcels per hemisphere (Fig. 4B). Since we processed each hemisphere

independently, we compared the similarity of the parcellations of the two hemispheres.

The hemispherical parcellations are similar in the parcel sizes (Supplementary Fig. 9-A;

Wilcoxon paired signed-rank test, N=96, p=0.7981; Supplementary Fig. 9-C for the size

comparison in the subject's native space) and functional connectivity patterns between

vertices within the same parcel (Supplementary Fig. 9-B; Wilcoxon paired signed-rank

test, N=96, p=0.411). For continuity with previously released resources [13, 15, 16], we

named this functional connectivity-based parcellation of the cortex "Marmoset Brain
Mapping Atlas Version 4" (MBMv4). We also provide an online viewer to visualize
parcellations, including previous versions (atlasviewer.marmosetbrainmapping.org).

To estimate the validity of the generated functional parcels, we used the distance-
controlled boundary coefficient (DCBC) [26]. The basic idea of DCBC is that when a
boundary divides two functionally homogenous regions, for any equal distance on the
cortical surface, the functional connectivity pattern between vertices within the same
parcel should be higher than that between vertices in different parcels (Fig. 4C). In other
words, a higher DCBC (within - between) means higher within-parcel homogeneity and
higher between-parcel heterogeneity. We calculated the DCBC between the vertex
pairs using a range of spatial bins (0–4 mm) with a 0.5 mm step (the spatial resolution
of the rs-fMRI data). Here, we compared the fit of the functional map represented by
MBMv4 with existing structural cortical parcellations, including MBMv1 atlas [15], the
digital reconstruction of the Paxinos atlas [15, 27], and the RIKEN atlas [28]. The result of
DCBC in Fig. 4C demonstrates that MBMv4 has the best performance for the
presentation of functional connectivity (the average DCBC values were 0.0186, 0.0135,
0.0177, 0.0330 for RIKEN, MBMv1, Paxinos, and MBMv4 atlas; multiple comparisons
for One-Way ANOVA $F_{(3,8556)}=22.44$, $p=1.81 \times 10^{-14}$).

Mapping MBMv4 in Individual Brains by Deep Neural Networks

To overcome the limitation of variable individual boundary maps (see Fig. 3C-E), we
employed a deep-learning approach for the individual Mapping from MBMv4 (Fig. 5A).
First, based on the population-level whole-brain functional connectivity, we trained a
deep neural network classifier for each parcel to learn the associated fingerprint of
functional connectivity. Then, the trained networks distinguished the goal parcel for
every marmoset based on the corresponding functional connectivity of the searching
area, consisting of the goal parcel and its neighbors. Due to the overlap of searching
areas, vertices could belong to multiple parcels. Therefore, we only kept these vertices
attributed to a single parcel as the seeds for regional growth by the "watershed"

algorithm. This iterative region-growing procedure would assign all vertices to a parcel,
resulting in an individual cortical parcellation.

Since individual parcellations should be close to the population definition [29, 30], we
compared the population-based MBMv4 parcellation and the automatically generated
individual parcellations. By calculating the percentage of vertices sharing the same
labels from both hemispheres (a metric of concordance), we found that the individual
parcellations from all marmosets are similar to MBMv4 with an average of 90%
concordance (Fig. 5B, the violin/box plot on the left, the examples on the right). Using
the test-retest dataset, we revealed the consistency of the individual parcellations
across different sessions (Fig. 5C, the violin/box plot on the left, and examples on the
right). The across-session analysis yielded an average of 86.7% concordance, lower
than the average value of 91.3% across-individual similarities. Furthermore, we
observed that the lateral prefrontal cortex and occipital-temporal cortex had higher
across-individual and across-session mapping variabilities (Supplementary Fig. 10),
consistent with previous findings in humans [22, 31]. Thus, the deep-learning approach
efficiently adjusts the parcel borders to reflect the individual variabilities while
maintaining high consistency with the population parcellation.

We also used the DCBC to evaluate whether the border adjustment of the individual
parcellation captured the specific features of individual functional connectivity patterns.
We assumed that the deep learning-based method (DNN-reg) should result in a higher
DCBC than the direct spatial registration of MBMv4 (Spatial-reg). Figure 5D (Top panel)
presents the functional connectivity for the pairs of vertices within the same parcel
(average correlation values within the same surface length 0-4 mm were 0.8331 and
0.8172 for DNN-reg and Spatial-reg) and between different parcels (average correlation
values were 0.8256 and 0.8171 for DNN-reg and Spatial-reg). Thus, the DNN-reg had
higher DCBC (within-between) than the Spatial-reg (Fig. 5D, bottom panel; the average
DCBC values were 0.0167 and 0.0085 for the DNN-reg and the Spatial-reg, respectively;
multiple comparisons for One-Way ANOVA $F_{(1,2512)}=20.35$, $p=6.74\times10^{-6}$). In sum, the

border adjustment by the proposed deep learning network reflects individual functional
connectivity patterns.
281
MBMv4 reflects accurate functional and topographical organizations
283
As evaluated from functional connectivity, MBMv4 provides a more accurate reflection
of the MRI-based functional parcellation of the cortex than current histology-based
atlases. To further verify this reliability, we took a task activation map during the
presentation of movie [32] which encompassed 10 deg × 8 deg of the visual field. This
activation map was then registered onto the same individual MBMv4 map and the
histology-based Paxinos atlas to examine the spatial overlap between the activations
and functional parcels. As a result, we found that the MBMv4 has a good
correspondence with task activations by visual inspection, such as the co-activation of
foveal areas V1 and MT, and the temporal parcels (Fig. 6A, flat maps). Additionally, by
measuring the shortest distances from every vertex in the boundary of the activation
map to the atlas boundaries (MBMv4 or Paxinos boundaries), we found that the parcel
borders of MBMv4 have higher consistency with the activation map than the Paxinos
atlas (Fig. 6A, the scatterplots; Wilcoxon paired signed-rank test: Monkey ID 25, N=878,
$p=3.07\times10^{-40}$ for the left hemisphere; N=816, $p=6.11\times10^{-26}$ for the right hemisphere.
Monkey ID 15, N=826, $p=2.22\times10^{-25}$ for the left hemisphere; N=850, $p=2.95\times10^{-53}$ for
the right hemisphere). Thus, MBMv4 reflects functional differences that
cytoarchitectonics does not capture, possibly because the latter contains the full visual
field representations. Thus, the MBMv4 may provide functional-related contrast that can
help enhance the precision of cross-species studies [33].
303
Besides the clear functional boundaries, MBMv4 also preserved the topographical
organization of the functional connectivity. Recent methodological developments have
allowed complex brain features to be mapped to the low-dimensional representations as
gradients [34], and these gradients characterized the topographical organization of the
functional brain connectome from unimodal to trans-modal networks. If the atlas
complies with this topographical organization, it should be able to identify such

gradients. As shown in Fig. 6B left panel, MBMv4 results in a pattern of gradient
spectrum for functional connectivity. In contrast, we did not find a gradient pattern based
on the Paxinos et al. (2012) atlas (right panel in Fig. 6B). Therefore, MBMv4 offers an
alternative view to understanding the functional connectome of the marmoset brain by
reflecting the characteristics of functional connectivity.

316 MBMv4 is an essential link between the functional and structural
317 connectivity

Since MBMv4 offers a more comprehensive scheme to study the functional connectome,
it is worth linking it to structural information to investigate the relationships between
structural and functional connectivity. We have also provided online viewers to facilitate
the comparison between them (connectome.marmosetbrainmapping.org). Furthermore,
we used a whole-brain computational model [35, 36, 37] to explore structural and functional
connectivity relationships. The processing procedure is shown in Fig. 7A. First, we
established structural connectivity values using the in-vivo diffusion MRI, the ex-vivo
ultra-high-resolution diffusion MRI, or the neuronal tracing datasets fitted to either the
MBMv4 or the Paxinos et al. (2012) atlas parcellations [14]. Then, we used a Hopf
bifurcation hemodynamic model [36] to simulate the functional connectivity of every brain
parcel or region based on their structural connectivity and compare it with the empirical
functional connectivity from the actual resting-state fMRI data. We used Pearson's
correlation to measure the similarity between the simulated and the empirical functional
connectivity. Additionally, we used group-average functional connectivity as an
empirical observation for the ex-vivo diffusion MRI and neuronal tracing dataset and
individual functional connectivity for the individual in-vivo diffusion MRI.

The modeling results in Fig. 7B-C demonstrate that: (1) the MBMv4 parcellation
provides the best fitting value (R=0.721) compared to the Paxinos atlas (R=0.638) for
the extra high-resolution ex-vivo diffusion MRI dataset (the polygon in Fig. 7B and
results in Fig. 7C); (2) the MBMv4 parcellation fits the in-vivo diffusion MRI datasets

better than the Paxinos atlas in a variety of individual simulations (all circles in Fig. 7B:
the average fitting values from 25 animals were R=0.4707 for MBMv4 and R=0.3659 for
Paxinos atlas); and (3) both parcellations performed equally (R=0.525 for MBMv4 and
R=0.472 for Paxinos atlas) fitting the cellular connectivity data from the
aggregated neuronal tracing (the star in Fig. 7B and results in Fig. 7C). However, no
matter which structural data was used for estimating functional connectivity, we found
that the modeling predicted by MBMv4 fits the empirical functional connectivity data
better than the Paxinos atlas (Fig. 7B; Wilcoxon paired signed-rank test: N=27,
p=0.002947).

Due to its inherent limitation for tracking long connections, the diffusion tractography
was more affected by distance, with longer connections having lower FC fitting values
(Fig. 7D, blue lines). Based on studies demonstrating that cellular-resolution
connections follow an exponential distribution of projection lengths [38, 39, 40], we
introduced an exponential distance rule (EDR) to our model to compensate for the
distance effect (Fig. 7D, redlines). Notably, the EDR fully corrected the distance effect in
neural tracing data (Fig. 7D), suggesting that the neural tracing data might be a more
reliable bridge to link structural and functional connectivity of long-range connections.
Still, as shown in Fig. 7E-F, the EDR-constrained modeling based on MBMv4 fits the
empirical functional data better than the Paxinos atlas (Wilcoxon paired signed-rank test:
N=27, p=0.01947), similar to the modeling without EDR (Fig. 7B), showing that MBMv4
is better suited for investigating the relationship between structural and functional
connectivity.

Discussion

There are challenges in adapting well-established approaches for human neuroimaging
to NHP neuroimaging [5]. The present study used effective and practical animal training
and imaging protocols to scan a large cohort of marmosets. Despite the different MRI
scanners (7T and 9.4T) in two institutions (ION and NIH), the protocol produced similar

data quality, suggesting the compatibility of our approach (see Methods for details,

Supplementary Fig. S1-S4). Therefore, we pooled the in-vivo resting-state fMRI

datasets from both institutions (details in Supplementary Table 1) to create the most

comprehensive functional connectivity dataset of the NHP brain to date, which was

integrated with the in-vivo diffusion MRI of the same cohort, as well as the highest

resolution ex-vivo diffusion MRI [13] and the most extensive mesoscale retrograde

neuronal tracing [14] available. This resource expedites the mapping of marmoset brains

and will allow cross-species comparisons.

Like the human cortex, the marmoset cerebral cortex is composed of large-scale

functional networks. However, the first awake resting-state fMRI study of the marmosets

[19] found only 12 functional networks (10 cortical networks), and another ICA-based

study described 8 brain networks, possibly due to the influence of anesthesia [20]. Based

on a more extensive awake rs-fMRI dataset, the present study mapped the large-scale

functional networks and built the first network-based parcellation, providing a more

comprehensive description of functional networks in the marmoset brain, including a

total of 15 cortical networks. Moreover, based on functional connectivity boundary maps,

we created a population-based cortical parcellation on a fine scale (MBMv4; Fig. 4) with

192 distinct parcels (96 per hemisphere). The further verification of hemispheric

symmetry is also done by inverting warping the standard average surfaces to every

subject's native space in Supplementary Fig. 9-C. A previous study of the human

cerebral cortex identified 422 discrete functional connectivity parcels using the same

approach, 206 in the left hemisphere and 216 in the right hemisphere [22]. Therefore, our

results may reflect that the asymmetry in marmosets may be smaller than in humans, as

expected from previous analyses based on anatomical measurements [41] and the

evidence that the number of cortex subdivisions increases with brain volume [42].

It is important to emphasize that our functional parcels do not correspond to the

traditional cytoarchitectonic definition of the cortical areas [43, 44]. Consistent with many

brain parcellations generated by non-invasive neuroimaging [22, 29, 45, 46, 47], our functional

parcels most likely reflect a different type of computational sub-units, agreeing with the

idea that the brain is organized in multiple scales [48, 49]. Therefore, compared with other

available structural atlases, MBMv4 captures the organization of functional connectivity

accurately. For example, MBMv4 achieved better task correspondence (Fig. 6A) due to

a strong link between task-fMRI and rs-fMRI [50, 51, 52]. Another evidence for the accuracy

of the MBMv4 parcellation is the topographical gradient organization of functional

connectivity (Fig. 6B). Finally, MBMv4 better links the structural and functional

connectivity, as demonstrated by our modeling simulation (see Fig. 7).

Consistent with the previous findings in humans [22, 29, 47], the parcels defined in MBMv4

do not follow the boundaries of cytoarchitectonic areas, thus demonstrating an essential

difference between anatomical features and functional connectivity. For example, the

somatomotor cortex is parcellated into subregions that appear to correspond to

representations of the face, forelimbs, hindlimbs, and trunk musculatures across

multiple areas, and areas such as V1 and V2 are subdivided into several functional

parcels according to the representation of eccentricity in visual field representation,

which is contiguous across areas [53] but may include discontinuities [54]. Previous studies

revealed that some topographically organized cytoarchitectonic areas could be

dissociated from the resting-state functional responses [55, 56]. Therefore, the present

MBMv4 should be considered a functional connectivity description, providing

complementary information about a type of organization that cannot be observed via

classical anatomy.

An essential goal of this study was to reflect individual characteristics by creating

parcels from each individual's data. Although the boundary map-derived parcels could

be used for individual analysis, we found that the subject boundary maps had significant

variations and that the reproducibility became lower than the group-level map (Fig. 3).

This finding emphasizes the need to acquire large amounts of data for the reliable test-

retest of the individual boundary map. Accordingly, we developed a deep neural

network to map reliable population-level MBMv4 into every individual nonlinearly. As a

result, we demonstrated good reliability in the test-retest dataset (across sessions from

the same individual; see Fig. 5C) and the applications of task-fMRI activation mapping

from the same individuals (Fig. 6A). Importantly, the locations of the most variable

functional parcels are in the lateral prefrontal cortex and lateral temporal-occipital cortex

(see Supplementary Fig. 10), corresponding to previously reported exceptionally high

inter-subject variability resting-state functional connectivity patterns [31]. Moreover, these

regions co-locate, expanding preferentially in primate evolution [57] and maturing later in

postnatal development [58]. As resting-state functional connectivity can be altered by

many biological features, including behavioral context during development associated

with phenotypic correlations, a better understanding of the causes of inter-subject parcel

variation will be an important focus for our future work.

In addition to the functional connectome mapping, we integrated all currently available

structural connectome datasets, including the in-vivo diffusion MRI, the ex-vivo high-

resolution diffusion MRI [13], and the mesoscale tracing dataset [14]. Using multimodal data

allowed us to investigate the relationship between functional and structural connectivity

with unprecedented detail. Using whole-brain modeling [36, 37], we observed that the

structural connectivity simulated functional connectivity based on MBMv4 had a high

coherence with empirical data, no matter which types of structural connectivity were

used (Fig. 7B-C and examples Fig. 7E-F). The finding corroborates the conclusion that

MBMv4 reflects meaningful computational sub-units from the view of whole-brain

functional connectivity. Meanwhile, we also found room for modeling performance

improvement through the detailed estimation of structural connectivity. For example,

compared with in-vivo diffusion MRI data, the ultra-high-resolution ex-vivo diffusion MRI

data from a brain sample could provide more thorough structural information. Notably,

we should also be careful about the influence of the distance when we use the diffusion

dataset to simulate functional connectivity because the diffusion tract may generate the

fiber cut and so on in the long-distance tract (Fig. 7D, blue curves). To overcome this

shortcoming, we introduced a correction factor for structural connectivity based on the

exponential distance rule (Fig. 7D, red curves). Nevertheless, the simulation results in

Fig. 7B and 7E demonstrated the best fitting by the present MBMv4, no matter which

datasets were used. Furthermore, since our modeling is simple, with only two

parameters that avoid overfitting simulation (see our method description), the whole-

brain model could be an efficient tool with broad applications to link structure and
function for future studies.

Although we provided the most comprehensive multimodal data resource for mapping
the marmoset connectome, our current study still faced limitations that need to be
addressed by future work, which is essential for both experimenters and users of non-
human primate neuroimaging. First is the data collection. The population used to
generate the MBMv4 was sex-biased (31 males vs. 8 females) due to the priority of
colony expansion worldwide. The neuronal tracing data were also limited, not covering
all cortical regions and missing subcortical information. Including neuronal-tracing data
is critical for accurately mapping the future fully structural connectome. Finally, our
parcellation only used the resting-state functional connectivity information, as in many
human studies[22, 45]. Therefore, multimodal brain parcellation incorporated structural
contrasts, especially the T1w/T2w myelin map, multiple task-fMRI data, gene
expression data, etc., will be necessary to fully capture the anatomical and functional
architectures of the marmoset brain.

Methods

Data Collection and Preprocessing

Animals and MRI scanning

Experimental procedures followed policies established by the Chinese Laboratory
Animal – Guideline for Ethical Review of Animal Welfare (ION data) or the US Public
Health Service Policy on Humane Care and Use of Laboratory Animals (NIH data). All
procedures were approved by the Animal Care and Use Committee (ACUC) of the
Institute of Neuroscience, Chinese Academy of Sciences (ION data), or the ACUC of
the National Institute of Neurological Disorders and Stroke, National Institutes of Health
(NIH data). The respective ACUC-approved protocols specify group size numbers

based on a power analysis to detect differences between animals to ensure rigor and
reproducibility of the results while minimizing the number of animals used in the study.
Our studies are powered to detect inter-individual differences. The experimental designs
are typically two or 3-factor ANOVAs. Values of $p < 0.05$ are considered statistically
significant. The number of animals used is the minimum necessary to provide reliable
estimates of inter-individual effects based on power considerations. Typically, sample
size estimates are based on the number of animals needed to achieve a power of 0.80
for moderate effect size and 0.99 for large effect size. Both marmoset colonies are
socially housed to ensure psychosocial well-being and are offered a varied diet that
includes food treats. In addition, dedicated husbandry and veterinary teams interact with
the animals daily as part of the psychological enrichment plans approved by the ACUCs
of both institutions.

The data acquisition procedure from both centers followed the same animal training
protocol, 8-element radiofrequency (RF) coil design [59], and MRI scanning protocols.
Thirteen marmosets (12 males and 1 female) were recruited from the ION cohort, from
which we generated 62 awake resting fMRI sessions and 349 runs (17 min per run). As
three of the 349 runs had extensive head motions (> 10% time points were motion
censored based on the preprocessed pipeline described below), we excluded the three
runs from the analysis, resulting in a total of 346 runs (see Supplementary Table 2 for
the summary of the head-motion per run). Twenty-six marmosets (19 males and 7
females) were recruited from the NIH cohort to produce 51 awake resting-state fMRI
sessions and 364 runs. Therefore, the NIH and ION data had a comparable number of
valid runs. The two datasets included 39 marmosets with 113 sessions, 710 valid fMRI
runs, and 12117 mins of total scan time. Detailed demographic information is provided
in Supplementary Table 1. All marmosets underwent a 3- to 4-week acclimatization
protocol as previously described [60]. After completing the training, all marmosets were
properly acclimated to laying in the sphinx position in an MRI-compatible cradle. Their
heads were comfortably restrained with 3D-printed anatomically conforming helmets
that allowed the resting-state fMRI (rs-fMRI) data acquisition as the animals lay relaxed
in their natural resting position.

All 39 marmosets were imaged using identical rs-fMRI protocols and pulse sequences,

except for a minor adjustment in the echo time (TE) to accommodate hardware

differences between the ION and the NIH gradient sets. The ION marmosets were

scanned in a 9.4T/30cm horizontal MRI scanner (Bruker, Billerica, USA) equipped with

a 20 cm gradient set capable of 300 mT/m gradient strength and an 8-channel phased-

array RF coil [59] custom-built for marmosets (Fine Instrument Technology, Brazil).

Multiple runs of rs-fMRI data were collected in ParaVision 6.0.1 software using a 2D

gradient-echo (GE) EPI sequence with the following parameters: TR$\square$=$\square$2 s, TE$\square$=$\square$18

ms, flip angle$\square$=$\square$70.4°, FOV$\square$=$\square$28 ×$\square$36$\square$mm, matrix size$\square$=$\square$56$\square$×$\square$72, 38 axial

slices, slice thickness$\square$=$\square$0.5 mm, 512 volumes (17$\square$min) per run. The GE-EPI fMRI

data were collected using two opposite phase-encoding directions (LR and RL) to

compensate for EPI distortions and signal dropouts. Two sets of spin-echo EPI with

opposite phase-encoding directions (LR and RL) were also collected for EPI-distortion

correction (TR$\square$=$\square$3000$\square$ms, TE$\square$=$\square$37.69 ms, flip angle$\square$=$\square$90°, FOV$\square$=$\square$28 ×$\square$36

mm, matrix size$\square$=$\square$56$\square$×$\square$72, 38 axial slices, slice thickness$\square$=$\square$0.5 mm, 8 volumes

for each set). After each rs-fMRI session, a T2-weighted structural image (TR$\square$=$\square$8000

ms, TE$\square$=$\square$10 ms, flip angle$\square$=$\square$90°, FOV$\square$= 28 ×$\square$36$\square$mm, matrix size$\square$=$\square$112 ×$\square$144,

38 axial slices, slice thickness$\square$=$\square$0.5$\square$mm) was scanned for co-registration purposes.

The NIH marmosets were scanned in a 7T/30cm horizontal MRI (Bruker, Billerica, USA)

equipped with a 15 cm customized gradient set capable of 450 mT/m gradient strength

(Resonance Research Inc., Billerica, USA) and an 8-channel phased-array RF coil

custom-built for marmosets [59] with identical coil geometry to the one used by ION.

Multiple runs of rs-fMRI data were collected during each scanning session in ParaVision

6.0.1. software using a 2D gradient-echo (GE) EPI sequence with the following

parameters: TR$\square$=$\square$2s, TE$\square$=$\square$22.2$\square$ms, flip angle$\square$=$\square$70.4°, FOV$\square$=$\square$28$\square$×$\square$36$\square$mm,

matrix size$\square$=$\square$56$\square$×$\square$72, 38 axial slices, slice thickness$\square$=$\square$0.5$\square$mm, 512 volumes

(17$\square$min) per run. The GE-EPI fMRI data were collected using two opposite phase-

encoding directions (LR and RL) to compensate for EPI distortions and signal dropouts.

Two sets of spin-echo EPI with opposite phase-encoding directions (LR and RL) were

also collected for EPI-distortion correction (TR□=□3000□ms, TE□=□36□ms, flip angle

= 90°, FOV□=□28 × 36□mm, matrix size □= 56 × 72, 38 axial slices, slice

thickness□=□0.5□mm, 8 volumes for each set). After each rs-fMRI session, a T2-

weighted structural image (TR□=□6000□ms, TE□=□9□ms, flip angle□=□90°,

FOV□=□28□×□36□mm, matrix size = 112□×□144, 38 axis slices, slice

thickness□=□0.5□mm) was scanned for co-registration purposes. Furthermore,

multishell diffusion MRI (DTI) datasets were collected using a 2D diffusion-weighted

spin-echo EPI sequence with the following parameters: TR = 5.1 s, TE = 38 ms, number

of segments = 88, FOV = 36 × 28 mm, matrix size = 72 × 56, slice thickness = 0.5 mm,

a total of 400 DWI images for two-phase encodings (blip-up and blip-down) and each

has 3 b values (8 b = 0, 64 b =2400, and 128 b = 4800), and the scanning duration was

about 34 min. The multishell gradient sampling scheme was generated using the Q-

shell sampling method [61].

Data Preprocessing

The rs-fMRI datasets were preprocessed by the customized script involving AFNI [62],

FSL [63], ANTs [64], and Connectome Workbench [65]. In brief, the rs-fMRI data were slice-

timing-corrected and motion-corrected by the "3dTshift" and "3dvolreg" commands of

AFNI and corrected for EPI distortions by the "top-up" command of FSL (see our

examples in Supplementary Fig. 11). The rs-fMRI datasets were further preprocessed

by regressing linear and quadratic trends, motion parameters and their derivatives, and

motion-sensor regressors (any TRs and the previous TRs were censored if the

detection motion was > 0.2□mm and temporal outlier > 0.1). Note that, for the motion

measurements, we calculated the weighted euclidean norm of six motion parameters

with a 0.25 weight for the three rotation degrees (yaw, pitch, and roll) according to the

relative head radius of the marmosets compared to humans. White matter and

cerebrospinal fluid signals were removed, and the rs-fMRI datasets were band-pass

filtered (0.01–0.1□Hz). The above nuisance signal regression and band-passing filtering

were carried out by the "3dDeconvolve" and "3dTproject" commands in AFNI. Next, the

preprocessed data were spatially normalized to the template space of our Marmoset

Brain Atlas Version-3 (MBMv3) by the "antsRegistration" routine of ANTs [16]. The spatial

normalization concatenated multiple transformations, including 1) rigid-body

transformation of each fMRI run to the T2-weighted image acquired at the end of each

session, 2) rigid-body transformation of T2-weighted images from each session to a

cross-session averaged T2-weighted image from each animal, 3) affine and nonlinear

transformation of the averaged T2-weighted image from each animal to the T2w

template of our MBMv3 space. Finally, all preprocessed data were mapped to 3D brain

surfaces of the MBMv3 using the Connectome Workbench (wb_command -volume-to-

surface-mapping function and ribbon constrained mapping algorithm), normalized

(subtract mean and divide by standard deviation) and concatenated per session before

the boundary mapping described below. The preprocessed data were smoothed with

1mm FWHM using 3dBlurInMask (for volume data) and wb_command -cifti-smoothing

(for surface data), respectively, before the network analysis and cortical parcellation.

The in-vivo diffusion MRI dataset was preprocessed by the DIFF_PREP, DR_BUDDI,

and DR_TAMAS pipelines of TORTOISE [66]. The DIFF_PREP and DR_BUDDI routines

incorporated correction for eddy-currents- and EPI-induced distortions using pairs of

diffusion data acquired with opposite phase encoding (blip-up and blip-down) and the

T2-weighted image and merging the preprocessed pairs into one dataset. The nonlinear

spatial registration from the individual space to the DTI template of our MBMv3 space [16]

was carried out using the DR_TAMAS routine of TORTOISE. The registration

information was then used to transform multiple atlases into the individual space for

diffusion tractography.

All diffusion trackings were performed using the iFOD2 method of the software Mrtrix3 [67].

The response function of each preprocessed diffusion MRI data was calculated by the

"dhollander" method of the "dwi2response" command, and then the fiber orientation

distributions (FOD) were estimated using spherical deconvolution by the multishell

multi-tissue CSD method of the "dwi2fod" command. Finally, region-to-region

tractography was performed using the iFOD2 method of the "tckgen" command. For

each pair of cortical regions, diffusion tractography was conducted by using one region

as the seed and the other as the target, and vice-versa. Thus, each pair of regions

generated two sets of tracking probability maps, which were normalized by the total

streamlines selected, and the two probability maps were averaged into a single map to

represent the final map of the connection between the two regions. Finally, all pairs of

connections formed the whole cortical structural connectome for computational

modeling.

The neuronal tracing data were mapped onto the histological NM template from our

previous study [17]. The NM template is a population-based 3D cortical template

generated from Nissl-stained serial sections of 20 marmosets. Since the NM template

only covers the cortex and has Nissl-stain contrast and a 75μm isotropic high spatial

resolution, its direct spatial transformation to our in-vivo MBMv3 template is inaccurate.

Thus, we modified the 80μm isotropic ultra-high-resolution MTR template of our

Marmoset Brain Atlas Version-2 (MBMv2) atlas [13] to remove the parts of the brain that

were not covered in the NM template, including the cerebellum, brainstem, and parts of

subcortical structures. This step increased the accuracy of registration between the NM

template and the MBMv2 template. Then, the ex-vivo MTR template of the MBMv2 was

transformed into the in-vivo myelin-map template of our MBMv3. By concatenating the

two transformations (the NM-to-MBMv2 and the MBMv2-to-MBMv3), we accurately

converted the neuronal tracing data from the NM template to the MRI template. We then

mapped the neuronal tracing data onto the MBMv3 cortical surfaces. For the above

registrations, we used the CC similarity metric as the cost functions and three-stage

alignments (rigid alignment, affine alignment, and nonlinear SyN transformations), which

were also the default options antsRegistrationSyN.sh. An example of registration results

is shown in Supplementary Fig. 12.

The instruction and example code for the data preprocessing pipeline is provided via the

resource webpage https://marmosetbrainmapping.org/data.html (ReadMe and Codes

sections), allowing the user to replicate our protocols.

Data harmonization across NIH and ION datasets

We calculated a series of indices to test the data harmonization across different

datasets (NIH and ION). They include the single time points SNR, mean images

(average across time for one fMRI run), SNR, temporal Signal Noise Ratio (tSNR, from

one fMRI run), Contrast to Noise Ratio (CNR, the mean of the gray matter intensity

values minus the mean of the white matter intensity values divided by the standard

deviation of the values outside the brain), temporal contrast to Noise Ratio (tCNR, the

variance of optimal resting-state fMRI components after ICA contrast to the noisy

component), the Fiber (Foreground to Background Energy Ratio: the variance of voxels

inside the brain divided by the variance of voxels outside the brain), head motion and

the whole-brain functional connectivity across subjects and sessions.

Functional Networks, Cortical Parcellation and Network Modeling

Brain network identification by the Group-ICA

Independent Component Analysis (ICA) was performed by the Group-ICA routine of the

GIFTI software (https://trendscenter.org/software/gift/) to identify the brain networks

using a number of different component settings. First, preprocessed data without

bandpassing and regression of nuisance covariates were group-ICA analyzed with

increasing numbers of ICA components from 20 to 80 in a step of 10. We tested the

reliability of different ICA methods, including the default "Infomax" ICA algorithm or

"ICASSO" group-ICA method, on different datasets (the NIH dataset, the ION dataset,

or combined both datasets) and obtained consistent results regardless of the ICA

setting or dataset used. Finally, every resulting component from Group-ICA analyses

was visually inspected and sorted according to its neuroanatomical features. Since the

sorted elements were highly consistent across different settings of ICA-component

numbers, we selected the best component to represent every labeled network. We also

did manual correction before creating the final network parcellation; for example, when

the left and right parts of the same network were separated into two components, we

merged them into one network by averaging their maps. We identify 15 cortical resting-

state networks from the group-ICA analysis (Fig. 2A-O and Supplementary Fig. 5).

We combined the 15 cortical networks from Fig. 2A-O according to their normalized Z

scores from ICA to create a cortical-network parcellation. The details include 1) the

combination of networks according to their spatial locations; 2) if they have spatial

overlapping, we took the highest value according to their normalized Z scores from ICA;

3) short-range (local) connectivity is usually stronger than long-range connectivity, so

the single map cannot cover long-range connectivity due to the spatial overlapping.

Therefore, we created a second map to cover the components with long-range

connectivity that are missed in the first map. We repeat the above step but only applied

to networks with long-range connections (such as Fig. 2I-K) to obtain the second map.

The primary map **(**Fig. 2P-Q, top rows) is mostly contributed by the short-range

networks (i.e., Fig. 1G, L, I, and O), and the second one (Fig. 2P-Q, bottom rows) is to

cover the long-range connectivity that was not captured by the primary map.

Since the human connectome project released a pipeline for denoising by ICA-FIX, it

has recently been applied to some animal research studies. Therefore, we also

explored whether adopting this processing would affect the identification of brain

networks. Before using ICA-FIX, we had to create a training dataset as a standard. For

this, we ran first-level ICA on each fMRI run, randomly selected 24 runs from 24 animals

(12 runs from the ION dataset and 12 runs from the NIH dataset), and manually

classified the noise components based on their spatial patterns and power spectrums.

Since the recommendations for the use of ICA-FIX suggest a training dataset of at least

10, we created a total of 24 datasets (trainingMBMv4.RData) to improve significantly our

ICA-FIX classifier, which included both the ION dataset (trainingION.RData, using 12

ION training datasets) and the NIH (trainingNIH.RData, using 12 NIH training datasets)

respectively. Then, a trained-weighted file (trainingMBMv4.RData) was used to clean all

fMRI data based on three sensible-value thresholds (5, 10, and 20) since ICA-FIX

recommends a threshold in the range of 5-20. Thus, we created three different versions

of ICA-FIX cleaned datasets. In addition, we provide the mask files to allow ICA-FIX to

work on the marmoset data since ICA-FIX had several steps that use human-default

settings and files incompatible with the marmoset data. The ICA-FIX training dataset

can be accessed via https://marmosetbrainmapping.org/data_for_MBMv4_reviews.html.

Regardless of which training dataset was used, we still obtained the same results, as

shown in Fig. 2A-O. Here, it is worth noting the limitations of ICA-FIX compared to our

traditional ICA methods. 1). ICA-FIX is a subjective approach that requests the user to

classify the noise and signal. Therefore, a subjective sensitive-value threshold has to be

defined. On the contrary, although less flexible, the traditional regression approach is an

objective method and is easier for others to replicate. 2). ICA-FIX does not work for the

raw dataset, meaning the data should be preprocessed by spatially smooth in a large

window. To some extent, some signal information might be lost. 3). The research aim of

ICA-FIX is to be pipelined for human MRI imaging. Thus, many predefined parameters

are optimized for human data preprocessing, which may affect the results' reliability

unless widely tested with large-scale species-specific datasets and optimized by

developers.

Boundary map generation

Following similar procedures to those described previously in a human imaging study [22],

the boundary mapping of resting-state functional connectivity data was implemented in

the Connectome Workbench and using customized Matlab codes (Mathworks, Natick,

USA, Version 2019b; see the scripts in our open resource). First, the time course of

every surface vertex for each brain hemisphere of each subject was correlated with

every other surface vertex to make a correlation map. Then, a similarity map was

created for every vertex by calculating pairwise spatial correlations between all

correlation maps. Thirdly, the first spatial derivative was applied on the similarity map by

the Connectome Workbench's function "cifti-gradient" to generate gradient maps for

each brain hemisphere of each subject. Next, the gradient maps were averaged across

subjects to produce the group gradient maps for each brain hemisphere. Lastly, the

"watershed by flooding" algorithm was applied to identify boundaries in the gradient

maps.

Test-retest evaluation of the boundary map

To compare the reliability of the boundary maps between the ION and the NIH datasets (Fig. 3A and Supplementary Fig. 8), between the individuals (Fig. 3C-E), and between runs from the same individual (Fig. 3E), we first thresholded two resulting boundary maps for each hemisphere to retain the cortical vertices most likely to be boundaries (i.e., retaining the top quartile of boundary values for a cumulative probability of 0.75) and assessed the overlap of the two thresholded boundaries by calculating the Dice's coefficient. The Dice's similarity coefficient of two thresholded boundaries, A and B, is expressed as:

$$\text{dice}(A, B) = 2 * \left| \frac{\text{intersection}(A, B)}{|A| + |B|} \right| \tag{1}$$

The average Dice's similarity coefficient is the mean of Dice's similarity coefficients across hemispheres.

Cortical parcellation based on the population-level boundary map

The creation of parcels was implemented by the customized Matlab scripts (see our open resource). Firstly, based on the vertices with values smaller than their neighbors that were <5 vertices away, we identified all local minima of vertices on the boundary map as seeds for parcel creation. Then, the parcels were grown from these seeds using the "watershed algorithm" procedure as above, allowing them to expand outward from the seed until they met other parcels. Because the whole process depends on the number of seeds for parcel creation, this might result in a large number of parcels. Therefore, according to the performance, we manually defined a threshold for merging adjacent parcels, which is the 60th percentile of the values in the boundary map [22]. It means that any two adjacent parcels with an average value below this threshold were considered not sufficiently dissimilar and should be merged. Finally, according to the population-level boundary map, we visually examined the remaining parcels to identify

those needing further adjustment, including eliminating vertices and spatial smoothing.

The detailed manual processings for the post-optimization included 1) manually

adjusting the parcel borders, 2) manually correcting wrong areal attributions of the

region growing, and 3) spatial smoothing the parcel borders by 8-neighbor vertices. We

finally found the resulting cortical parcellation with 96 functional parcels in each

hemisphere as our Marmoset Brain Mapping Atlas Version-4 (MBMv4) in Fig. 4B.

Evaluation of cortical parcellation by the distance-controlled boundary coefficient

(DCBC)

Following a previous study [26], we used the distance-controlled boundary coefficient

(DCBC) as a metric to evaluate functional boundaries between our parcels. The

rationale for this method is that any two points belonging to any given parcel should

have more similar functional profiles than those belonging to different parcels.

Furthermore, because the functional organization varies smoothly, the correlation

between two points will weaken with increasing spatial distance. Thus, we calculated

the correlation coefficients for all pairs of points separated by a specific surface

Euclidean distance, using 0.5 mm spatial bins (same as fMRI imaging spatial resolution)

ranging from 0 to 4 mm for pairs of points residing within parcels or across different

parcels (between). The DCBC defines the difference between the within-parcel and

between-parcel pair correlations. A higher DCBC reflects that pairs within the same

region are more functional, serving as a global parcellation measure. For the group

comparison across atlases (Fig. 4C), the DCBC metrics were calculated for each

participant in each spatial bin and then averaged. For the same participant comparison

across atlases (Fig. 5D), the DCBC metrics were calculated for each session in each

spatial bin and then averaged.

Comparison with alternative atlases

We compared our parcellation against alternative digital parcellations created by various

approaches. These alternative parcellations included: (1) Paxinos atlas [68], the most

commonly used atlas in marmoset brain research, which is cytoarchitectonic

characterization by immunohistochemical sections, and here we used its 3D digital

version [15, 27]; (2) RIKEN atlas [28]: The atlas is cytoarchitecture based on Nissl-staining

contrast. (3) The first atlas version of Marmoset Brain Mapping (MBMv1) [15]: The borders

were delineated based on the high-resolution diffusion MRI contrast and parcellated by

a structural-connectivity-based approach.

Deep-learning-based individual parcellation generator

The group-average parcellation described in the preceding sections is desirable for

generating parcellations of individual animals. Although applying our group-level

parcellation to individual animals is feasible, as demonstrated in a previous human

study [22], we still found misalignments between individuals and cannot be highly

consistent with the tendency of the group-average parcellation (MBMv4) when the

scanning runs are limited (Fig. 3D-E). Therefore, inspired by previous works [29, 30], we

trained a multi-layer deep learning network to classify parcels based on the fingerprints

from MBMv4. There were two assumptions for this approach: (1) We assumed that

individual cortical parcels were close to the group definition after the feature-based

surface registration; (2) We assumed that every identified cortical parcel should be in a

single class which was the combination of the target parcel and its spatially adjacent

parcels (the "searchlight" for the candidate parcel). Thus, the setup of the classifier

network was straightforward. Its architecture was as follows (for the graphic reference,

see Fig. 5A): for each of the 96 parcels in each hemisphere, a multi-layer deep neural

network was designed, which comprised three layers (one input, five hidden, and one

output) and 384 hidden neurons (a reasonable compromise between accuracy and

training speed for the classification). The whole-brain fingerprint of the candidate parcel

from the MBMv4 worked as the training set for the network to classify whether or not

each vertex in an individual ROI contained the parcel plus all of its neighbor parcels.

Because of the spatial overlap of the "searchlight," we excluded the vertices belonging

to multiple parcels. Then, we applied the same procedure of parcel creation as above,

meaning that the borders of each identified parcel became the seeds to expand outward

until they met other parcels using the "watershed by flooding" procedure. The whole

process of individual parcellation was automatic and implemented using customized

MATLAB   codes   (example   codes   are   shared   via
www.marmosetbrainmapping.org/data.html) combined with the MATLAB Deep Learning
Network toolbox.

Evaluation by task-activation pattern

We examined the functional relevance of the borders by evaluating the parcels
contained within the fMRI activation pattern to a visual task (Fig. 6A) from our previous
study [32]. A subset of animals from the NIH dataset participated in the visual-choice task,
which consisted of watching 20-s-long movies (visual field is 10deg x 8deg) and 16s
resting periods (206 trials for marmoset-ID15 and 280 trials for marmoset-ID25). We
performed a contrast comparison between the movie-presentation blocks and the
resting blocks to generate visual-task activation statistical maps for each session. A
mixed-effects analysis was then applied to all statistical maps across sessions by the
3dMEMA command of AFNI to obtain a final statistical map. The map was thresholded
at a voxel-wise threshold of $p < 0.05$ and a cluster-wise threshold of $p < 0.05$ for
multiple comparison corrections. To compare the similarity of the activation map and the
parcellations in each hemisphere (for results, see the flat maps in Fig. 6A), we
calculated the shortest Euclidean distance of every vertex/voxel in the boundary of the
activation map to the vertexes/voxels in the boundary of parcels/regions from different
parcellations. We considered the parcellation with the overall shortest distances of
every vertex/voxel in the boundary of the activation map as the best border consistency
(for results, see the scatterplots in Fig. 6A).

Evaluation by functional connectivity gradient spectrum

It is widely accepted that the cerebral cortex of multiple species, including both human
and macaque primates, is organized along principal functional gradients that provide a
spatial framework for the co-existence of multiple large-scale networks operating in a
spectrum from unimodal to transmodal functional activity [34, 69]. Therefore, if the MBMv4
parcellation created here accurately represents the functional organization of the
marmoset cortex, we can presume that it will also reveal these principal functional
gradients. Thus, as in previous studies [34, 70], we followed a workflow for gradient

identification: we first computed the rs-fMRI functional connectivity (RSFC) based on

MBMv4. Next, the RSFC matrix $M_{x,y}$ with the same size as the atlas was made sparse

(to a 10% sparsity), and a similarity matrix $A_{x,y}$ with the normalized angle was computed

according to the following equation:

870
$$A(x,y) = 1 - \frac{\cos^{-1}(\text{cossim}(x,y))}{\pi} \tag{2}$$

872   Next, the similarity matrix was decomposed via Laplacian transformation into a set of

873   principal eigenvectors describing the axes of most significant variance using the

874   following equation:

876
$$Lg = \lambda Dg \tag{3}$$

878   Where $D_{x,y} = \sum_y A(x,y)$, L is the graph Laplacian matrix, and the eigenvectors g which

879   corresponds to the m smallest eigenvalues λ_k are used to build the new low-

880   dimensional representation:

882
$$\varsigma_{LE} = [g_1, g_2, \dots, g_m] \tag{4}$$

Finally, the first two axes g_1, g_2 of each parcel were plotted in 2D space. Meanwhile, we

used the scores g_1 to sort the functional connectivity matrix (for results, see the

heatmaps in Fig. 6B).

The whole-brain modeling for the link between structural connectivity and

functional connectivity

As we know, structural connectivity and functional connectivity are closely related to

each other. Therefore, the lack of structural evidence generally implies biological

implausibility for functional connections. Testing whether the cortical parcels created

above MBMv4 are accurate representations of the functional areas in the cerebral

cortex requires investigation of the underlying structural connectivity. A computational

model is a powerful approach to bridge structural and functional connectivity [69, 71, 72, 73, 74]

. In the present study, we implemented a whole-brain model with only two free

parameters from previous studies [35, 36], as outlined below (for a graphic reference, see

Fig. 7A, note that the fMRI data for the modeling part is frequently unfiltered, so the

model used the full band of frequency):

According to the whole cortical parcellations (192 total parcels, 96 per hemisphere from

MBMv4 or 232 total regions, 116 regions per hemisphere from the Paxinos atlas), the

structural connectivity between parcels/regions $C_{i,j}$ was estimated from the structural

datasets (see examples in Fig. 7B-C and 7E-F), either DTI data (in-vivo or ex-vivo) or

the neuronal tracing data. Then, the local dynamics for every parcel/region j can be

properly approximated to the normal form of a Hopf bifurcation:

907
$$\frac{dz_j}{dt} = [a_j + iw_j]z_j + z_j|z_j|^2 \tag{5}$$

In this equation, z_j is a complex-valued variable $z_j = x_j + y_j$, and w_j is the intrinsic signal

frequency of parcel/region j, which ranged from 0.04-0.07Hz and was determined by the

averaged peak frequency of the bandpass-filtered fMRI signals of the parcel/region j [36,

75, 76, 77, 78]. a_j is a bifurcation free parameter controlling the dynamics of the parcel/region

j . For $a_j < 0$, the phase space presents a unique stable and is governed by noise.

For $a_j > 0$, the phase space presents the stable state, giving rise to a self-sustained

oscillation. For $a_j \approx 0$ the phase presents an unstable state, switching back and forth

and giving rise to a mixture of oscillation and noise.

The coordinated dynamics of the resting state activity for parcel/region j could be

modeled by coupling determined by the above structural connectivity $C_{i,j}$. To ensure the

oscillatory dynamics for $a_j > 0$, the structural connectivity $C_{i,j}$ should be normalized and

scaled to 0.2 in a weak coupling condition before starting the simulation. The coupled

differential equations of the model are the following:

921
$$\frac{dx_j}{dt} = [a_j - x_j^2 - y_j^2]x_j - w_jy_j + G\sum_i C_{i,j}(x_i - x_j)\beta\eta_j(t) \tag{6}$$

922
$$\frac{dy_j}{dt} = \left[a_j - x_j^2 - y_j^2\right]y_j + w_jx_j + G\sum_i C_{i,j}(y_i - y_j)\beta\eta_j(t) \tag{7}$$

In this equation, G is another free parameter representing the fixed global coupling
factor that scales structural connectivity $C_{i,j}$. η_j represents additive Gaussian noise in
each parcel/region and is scaled by a factor β fixed at 0.04 according to previous
studies [36]. Euler-Maruyama algorithm integrated these equations with a time step of 0.1
seconds to accelerate simulation [79].

The free bifurcation parameter a_j for parcel/region j could be locally optimized based on
fitting the spectral information of the empirical BOLD signals. To achieve this, we filtered
raw empirical BOLD data in the 0.04–0.25Hz band and calculated the power spectrum
$p_j(f)$ for each parcel j as below:

932
$$p_j = \frac{\int_{0.04}^{0.07} p_j(f)\,df}{\int_{0.04}^{0.25} p_j(f)\,df} \tag{8}$$

933 and updated the local bifurcation parameter a_j by a gradient descendent strategy:

934
$$a_j = a_j + \eta\left(p_j^{empirical} - p_j^{simulated}\right) \tag{9}$$

We applied the above optimization process to receive the best bifurcation parameters a_j
of every parcel/region defined in the parcellations. Once we found the optimized set of
bifurcation parameters a_j, we adjusted the free parameter G within the range of 0-8 in
steps of 0.1 according to a reasonable compromise from previous studies [36, 79] to
simulate the same number of sessions for each animal and the same number of
animals. To compare the performance in different atlases, we just needed to compare
fitting (similarity) metrics, Pearson's correlation coefficient between the simulated
functional connectivity and the one used for the empirical data, when we fixed the same
value of parameter G.

Since the distributions of the optimal bifurcation parameter a are identical in different
parcellations MBMv4 or Paxinos atlas (see Supplementary Fig. 13, one way ANOVA
$F_{(1,11986)}=9.09$, p=0.26), we selected the best free parameter G from the Paxinos atlas

for comparison performance with our MBMv4 (see results in Fig. 7B and 7E, examples

in Fig. 7C and 7F). Moreover, we also selected the group-averaged functional

connectivity from all individuals as the empirical observable for the ultra-high resolution

diffusion MRI and neuronal tracing datasets and the individual functional connectivity for

the corresponding in-vivo diffusion MRI.

The resulting simulation of functional connectivity demonstrated the influence of the

distance, especially for the diffusion MRI (see Fig. 7D). To decrease this impact, we

added an Exponential Distance Rule (EDR) for a more accurate estimation of structural

connectivity before the simulation, which can be implemented as follows: we first

normalized the structural connectivity estimated by diffusion tractography to 0-1, then

calculated the probability of structural connectivity according to the EDR rule ($p(d)=ce^{-\lambda d}$,

$\lambda \approx 0.3$, $c \approx 0.94$ where d is the distance [40]) and normalized to 0-1. Finally, we transformed

the normalized structural connectivity to match the probability of structural connectivity.

For simplicity to help us to identify abnormal values, we defined a threshold as the

Median Absolute Deviation for the normalized structural connectivity.

964
$$\widehat{\sigma} = \frac{\text{Median}(|X_i - \widetilde{X}|)}{0.6745} \tag{10}$$

965 Where X_i = each value, $\widetilde{X}$ = average value

967 If any values were larger than this threshold, we considered them abnormal values, and

968 the corresponding probability of structural connectivity would replace them to reduce the

969 impact of distance.

Data availability

All of our datasets, including raw and preprocessed NIH and ION resting-state fMRI, diffusion MRI, and neuronal tracing datasets, are available on our specific webpage of the Marmoset Brain Mapping Project (www.marmosetbrainmapping.org/data.html). The volume data are in NIFTI format, and the surface data are in CIFTI format. The raw MRI data without processing is provided in the standard BIDS format for cross-platform sharing. The MBMv4 parcellations are also provided on the same webpage (www.marmosetbrainmapping.org/data.html), and the MBMv1 and Paxinos parcellations on our MRI template space are a part of the MBMv3 resource (marmosetbrainmapping.org/v3.html). The high-resolution ex-vivo diffusion MRI data are a part of the MBMv2 resource (marmosetbrainmapping.org/atlas.html#v2). The raw neuronal tracing data are from Marmoset Brain Connectivity project (https://www.marmosetbrain.org/reference). Note that the MBMv4 datasets are only available for scientific purposes and are licensed under Creative Commons Attribution-NonCommercial-ShareAlike (CC BY-NC-SA 4.0).

Code availability

The codes for analyzing pipelines used in this study could be available on our webpage (www.marmosetbrainmapping.org/data.html). The codes package with examples includes the generation of cortical functional boundary maps, functional parcellation, and their evaluation metrics, as well as computational modeling which could link with structural imaging and neuronal tracing. Note that the codes and analyzing pipelines are only available for scientific purposes and are licensed under Creative Commons Attribution-NonCommercial-ShareAlike (CC BY-NC-SA 4.0).

References

1. Buffalo EA, Movshon JA, Wurtz RH. From basic brain research to treating human brain disorders. Proc Natl Acad Sci U S A, (2019).

2. Thiebaut de Schotten M, Croxson PL, Mars RB. Large-scale comparative neuroimaging: Where are we and what do we need? Cortex **118**, 188-202 (2019).

3. Buckner RL, Margulies DS. Macroscale cortical organization and a default-like apex transmodal network in the marmoset monkey. Nat Commun **10**, 1976 (2019).

4. Smith SM, et al. Resting-state fMRI in the Human Connectome Project. Neuroimage **80**, 144-168 (2013).

5. Milham MP, et al. An Open Resource for Non-human Primate Imaging. Neuron **100**, 61-74 e62 (2018).

6. Hudetz AG. General anesthesia and human brain connectivity. Brain Connect **2**, 291-302 (2012).

7. Milham M, et al. Accelerating the Evolution of Nonhuman Primate Neuroimaging. Neuron **105**, 600-603 (2020).

8. Milham M, et al. Toward next-generation primate neuroscience: A collaboration-based strategic plan for integrative neuroimaging. Neuron, (2021).

9. Messinger A, et al. A collaborative resource platform for non-human primate neuroimaging. Neuroimage **226**, 117519 (2021).

10. Okano H. Current Status of and Perspectives on the Application of Marmosets in Neurobiology. Annu Rev Neurosci **44**, 27-48 (2021).

11. Walker J, MacLean J, Hatsopoulos NG. The marmoset as a model system for studying voluntary motor control. Dev Neurobiol **77**, 273-285 (2017).

12. Matsuzaki M, Ebina T. Common marmoset as a model primate for study of the motor control system. Curr Opin Neurobiol **64**, 103-110 (2020).

13. Liu C, et al. A resource for the detailed 3D mapping of white matter pathways in the marmoset brain. Nat Neurosci **23**, 271-280 (2020).

14. Majka P, et al. Open access resource for cellular-resolution analyses of corticocortical connectivity in the marmoset monkey. Nat Commun **11**, 1133 (2020).

1043

1044   15.   Liu C, et al. A digital 3D atlas of the marmoset brain based on multi-modal MRI.
1045   Neuroimage **169**, 106-116 (2018).

1046

1047   16.   Liu C, Yen CC, Szczupak D, Tian X, Glen D, Silva AC. Marmoset Brain Mapping V3:
1048   Population multi-modal standard volumetric and surface-based templates. Neuroimage
1049   **226**, 117620 (2021).

1050

1051   17.   Majka P, et al. Histology-Based Average Template of the Marmoset Cortex With
1052   Probabilistic Localization of Cytoarchitectural Areas. Neuroimage **226**, 117625 (2021).

1053

1054   18.   Autio JA, et al. Minimal specifications for non-human primate MRI: Challenges in
1055   standardizing and harmonizing data collection. Neuroimage **236**, 118082 (2021).

1056

1057   19.   Belcher AM, et al. Large-scale brain networks in the awake, truly resting marmoset
1058   monkey. J Neurosci **33**, 16796-16804 (2013).

1059

1060   20.   Ghahremani M, Hutchison RM, Menon RS, Everling S. Frontoparietal Functional
1061   Connectivity in the Common Marmoset. Cereb Cortex **27**, 3890-3905 (2017).

1062

1063   21.   Liu C, Yen CC, Szczupak D, Ye FQ, Leopold DA, Silva AC. Anatomical and functional
1064   investigation of the marmoset default mode network. Nat Commun **10**, 1975 (2019).

1065

1066   22.   Gordon EM, Laumann TO, Adeyemo B, Huckins JF, Kelley WM, Petersen SE. Generation
1067   and Evaluation of a Cortical Area Parcellation from Resting-State Correlations. Cereb
1068   Cortex **26**, 288-303 (2016).

1069

1070   23.   Cohen AL, et al. Defining functional areas in individual human brains using resting
1071   functional connectivity MRI. Neuroimage **41**, 45-57 (2008).

1072

1073   24.   Wig GS, Laumann TO, Petersen SE. An approach for parcellating human cortical areas
1074   using resting-state correlations. Neuroimage **93 Pt 2**, 276-291 (2014).

1075

1076   25.   Cousty J, Bertrand G, Najman L, Couprie M. Watershed Cuts: Minimum Spanning Forests
1077   and the Drop of Water Principle. Ieee T Pattern Anal **31**, 1362-1374 (2009).

1078

1079   26.   King M, Hernandez-Castillo CR, Poldrack RA, Ivry RB, Diedrichsen J. Functional
1080   boundaries in the human cerebellum revealed by a multi-domain task battery. Nat
1081   Neurosci **22**, 1371-1378 (2019).

1082

1083   27.   Majka P, et al. Towards a comprehensive atlas of cortical connections in a primate brain:
1084   Mapping tracer injection studies of the common marmoset into a reference digital
1085   template. J Comp Neurol **524**, 2161-2181 (2016).

1086

1087   28.   Woodward A, et al. The Brain/MINDS 3D digital marmoset brain atlas. Sci Data **5**,
1088   180009 (2018).
1089

1090   29.   Glasser MF, et al. A multi-modal parcellation of human cerebral cortex. Nature **536**, 171-
1091   178 (2016).
1092

1093   30.   Hacker CD, et al. Resting state network estimation in individual subjects. Neuroimage **82**,
1094   616-633 (2013).
1095

1096   31.   Mueller S, et al. Individual variability in functional connectivity architecture of the
1097   human brain. Neuron **77**, 586-595 (2013).
1098

1099   32.   Tian X, Silva AC, Liu C. The Brain Circuits and Dynamics of Curiosity-Driven Behavior in
1100   Naturally Curious Marmosets. Cereb Cortex **31**, 4220-4232 (2021).
1101

1102   33.   Russ BE, et al. Common functional localizers to enhance NHP & cross-species
1103   neuroscience imaging research. Neuroimage **237**, 118203 (2021).
1104

1105   34.   Margulies DS, et al. Situating the default-mode network along a principal gradient of
1106   macroscale cortical organization. Proc Natl Acad Sci U S A **113**, 12574-12579 (2016).
1107

1108   35.   Ipina IP, et al. Modeling regional changes in dynamic stability during sleep and
1109   wakefulness. Neuroimage **215**, 116833 (2020).
1110

1111   36.   Deco G, Kringelbach ML, Jirsa VK, Ritter P. The dynamics of resting fluctuations in the
1112   brain: metastability and its dynamical cortical core. Sci Rep **7**, 3095 (2017).
1113

1114   37.   Deco G, Jirsa V, McIntosh AR, Sporns O, Kotter R. Key role of coupling, delay, and noise
1115   in resting brain fluctuations. Proc Natl Acad Sci U S A **106**, 10302-10307 (2009).
1116

1117   38.   Horvat S, et al. Spatial Embedding and Wiring Cost Constrain the Functional Layout of
1118   the Cortical Network of Rodents and Primates. PLoS Biol **14**, e1002512 (2016).
1119

1120   39.   Ercsey-Ravasz M, et al. A predictive network model of cerebral cortical connectivity
1121   based on a distance rule. Neuron **80**, 184-197 (2013).
1122

1123   40.   Theodoni P, Majka P, Reser DH, Wojcik DK, Rosa MGP, Wang XJ. Structural Attributes
1124   and Principles of the Neocortical Connectome in the Marmoset Monkey. Cereb Cortex
1125   **32**, 15-28 (2021).
1126

1127   41.   Rilling JK, Insel TR. Differential expansion of neural projection systems in primate brain
1128   evolution. Neuroreport **10**, 1453-1459 (1999).
1129

1130   42.   Changizi MA, Shimojo S. Parcellation and area-area connectivity as a function of
1131   neocortex size. Brain Behav Evol **66**, 88-98 (2005).
1132

1133   43.   Kaas JH. The organization of neocortex in mammals: implications for theories of brain
1134   function. Annu Rev Psychol **38**, 129-151 (1987).
1135

1136   44.   Amunts K, Zilles K. Architectonic Mapping of the Human Brain beyond Brodmann.
1137   Neuron **88**, 1086-1107 (2015).
1138

1139   45.   Schaefer A, et al. Local-Global Parcellation of the Human Cerebral Cortex from Intrinsic
1140   Functional Connectivity MRI. Cereb Cortex **28**, 3095-3114 (2018).
1141

1142   46.   Van Essen DC, Glasser MF, Dierker DL, Harwell J, Coalson T. Parcellations and
1143   hemispheric asymmetries of human cerebral cortex analyzed on surface-based atlases.
1144   Cereb Cortex **22**, 2241-2262 (2012).
1145

1146   47.   Van Essen DC, Glasser MF. Parcellating Cerebral Cortex: How Invasive Animal Studies
1147   Inform Noninvasive Mapmaking in Humans. Neuron **99**, 640-663 (2018).
1148

1149   48.   van den Heuvel MP, Yeo BTT. A Spotlight on Bridging Microscale and Macroscale Human
1150   Brain Architecture. Neuron **93**, 1248-1251 (2017).
1151

1152   49.   Churchland PS, Sejnowski TJ. Perspectives on cognitive neuroscience. Science **242**, 741-
1153   745 (1988).
1154

1155   50.   Smith SM, et al. Correspondence of the brain's functional architecture during activation
1156   and rest. Proc Natl Acad Sci U S A **106**, 13040-13045 (2009).
1157

1158   51.   Mennes M, et al. Inter-individual differences in resting-state functional connectivity
1159   predict task-induced BOLD activity. Neuroimage **50**, 1690-1701 (2010).
1160

1161   52.   Tavor I, Parker Jones O, Mars RB, Smith SM, Behrens TE, Jbabdi S. Task-free MRI predicts
1162   individual differences in brain activity during task performance. Science **352**, 216-220
1163   (2016).
1164

1165   53.   Rosa MG, Tweedale R. Brain maps, great and small: lessons from comparative studies of
1166   primate visual cortical organization. Philos Trans R Soc Lond B Biol Sci **360**, 665-691
1167   (2005).
1168

1169   54.   Yu HH, Rowley DP, Price NSC, Rosa MGP, Zavitz E. A twisted visual field map in the
1170   primate dorsomedial cortex predicted by topographic continuity. Sci Adv **6**, (2020).
1171

1172   55.   Long X, Goltz D, Margulies DS, Nierhaus T, Villringer A. Functional connectivity-based
1173   parcellation of the human sensorimotor cortex. Eur J Neurosci **39**, 1332-1342 (2014).

1174
1175   56.   Rao SM, et al. Somatotopic mapping of the human primary motor cortex with functional
1176   magnetic resonance imaging. Neurology **45**, 919-924 (1995).
1177
1178   57.   Chaplin TA, Yu HH, Soares JG, Gattass R, Rosa MG. A conserved pattern of differential
1179   expansion of cortical areas in simian primates. J Neurosci **33**, 15120-15125 (2013).
1180
1181   58.   Sawiak SJ, et al. Trajectories and Milestones of Cortical and Subcortical Development of
1182   the Marmoset Brain From Infancy to Adulthood. Cereb Cortex **28**, 4440-4453 (2018).
1183
1184   59.   Papoti D, Yen CC, Hung CC, Ciuchta J, Leopold DA, Silva AC. Design and implementation
1185   of embedded 8-channel receive-only arrays for whole-brain MRI and fMRI of conscious
1186   awake marmosets. Magn Reson Med **78**, 387-398 (2017).
1187
1188   60.   Silva AC, et al. Longitudinal functional magnetic resonance imaging in animal models.
1189   Methods Mol Biol **711**, 281-302 (2011).
1190
1191   61.   Caruyer E, Lenglet C, Sapiro G, Deriche R. Design of multishell sampling schemes with
1192   uniform coverage in diffusion MRI. Magn Reson Med **69**, 1534-1540 (2013).
1193
1194   62.   Cox RW. AFNI: what a long strange trip it's been. Neuroimage **62**, 743-747 (2012).
1195
1196   63.   Jenkinson M, Beckmann CF, Behrens TE, Woolrich MW, Smith SM. Fsl. Neuroimage **62**,
1197   782-790 (2012).
1198
1199   64.   Tustison N, et al. Multivariate Analysis of Diffusion Tensor Imaging and Cortical
1200   Thickness Maps in a Traumatic Brain Injury (Tbi) Cohort Using Advanced Normalization
1201   Tools (Ants). J Neurotraum **28**, A111-A111 (2011).
1202
1203   65.   Marcus DS, et al. Informatics and data mining tools and strategies for the human
1204   connectome project. Front Neuroinform **5**, 4 (2011).
1205
1206   66.   Pierpaoli C. Quantitative brain MRI. Top Magn Reson Imaging **21**, 63 (2010).
1207
1208   67.   Tournier J-D, Calamante F, Connelly A. MRtrix: Diffusion tractography in crossing fiber
1209   regions. International Journal of Imaging Systems and Technology **22**, 53-66 (2012).
1210
1211   68.   Paxinos G, Watson C, Petrides M, Rosa M, Tokuno H. The marmoset brain in stereotaxic
1212   coordinates. Elsevier Academic Press (2012).
1213
1214   69.   Wang P, et al. Inversion of a large-scale circuit model reveals a cortical hierarchy in the
1215   dynamic resting human brain. Sci Adv **5**, eaat7854 (2019).
1216

1217 70. Vos de Wael R, et al. BrainSpace: a toolbox for the analysis of macroscale gradients in
1218 neuroimaging and connectomics datasets. Commun Biol **3**, 103 (2020).
1219
1220 71. Hansen EC, Battaglia D, Spiegler A, Deco G, Jirsa VK. Functional connectivity dynamics:
1221 modeling the switching behavior of the resting state. Neuroimage **105**, 525-535 (2015).
1222
1223 72. Breakspear M. Dynamic models of large-scale brain activity. Nat Neurosci **20**, 340-352
1224 (2017).
1225
1226 73. Schirner M, McIntosh AR, Jirsa V, Deco G, Ritter P. Inferring multi-scale neural
1227 mechanisms with brain network modelling. Elife **7**, (2018).
1228
1229 74. Honey CJ, et al. Predicting human resting-state functional connectivity from structural
1230 connectivity. Proc Natl Acad Sci U S A **106**, 2035-2040 (2009).
1231
1232 75. Ponce-Alvarez A, Deco G, Hagmann P, Romani GL, Mantini D, Corbetta M. Resting-state
1233 temporal synchronization networks emerge from connectivity topology and
1234 heterogeneity. PLoS Comput Biol **11**, e1004100 (2015).
1235
1236 76. Achard S, Salvador R, Whitcher B, Suckling J, Bullmore E. A resilient, low-frequency,
1237 small-world human brain functional network with highly connected association cortical
1238 hubs. J Neurosci **26**, 63-72 (2006).
1239
1240 77. Buckner RL, et al. Cortical hubs revealed by intrinsic functional connectivity: mapping,
1241 assessment of stability, and relation to Alzheimer's disease. J Neurosci **29**, 1860-1873
1242 (2009).
1243
1244 78. Glerean E, Salmi J, Lahnakoski JM, Jaaskelainen IP, Sams M. Functional magnetic
1245 resonance imaging phase synchronization as a measure of dynamic functional
1246 connectivity. Brain Connect **2**, 91-101 (2012).
1247
1248 79. Sanz Perl Y, et al. Perturbations in dynamical models of whole-brain activity dissociate
1249 between the level and stability of consciousness. PLoS Comput Biol **17**, e1009139 (2021).

Acknowledgments

We thank Kaiwei Zhang and Binshi Bo at the 9.4T core facility (CEBSIT) for assistance in the data collection of the ION data, Lisa Zhang for the assistance in the data collection of the NIH 7T data, the Marmoset Animal Facility of CEBSIT for animal care, Xiaojia Zhu for the assistance in organizing MRI data in BIDS format, and the NIH Fellows Editorial Board for the editorial assistance on the early version of the manuscript.

The study was supported by the grants from National Science and Technology Innovation 2030 Major Project of China (2022ZD0205000, 2021ZD0203900), the Pennsylvania Department of Health Commonwealth Universal Research Enhancement (CURE) Tobacco Settlement Appropriation – Phase 18 (Grant SAP4100083102 to ACS), the US National Institute on Aging (Grants R24AG073190 and U19AG074866 to ACS), the Shanghai Municipal Science and Technology Major Project (No. 2018SHZDZX05 to CL and ZF), the Lingang Laboratory (Grant No. LG-QS-202201-02 to CL), the National Natural Science Foundation of China (No. 32171088 to CL) , the Australian Research Council (DP110101200, DP140101968, CE140100007) and National Health and Medical Research Council (APP1194206) to MR, National Science Centre (2019/35/D/NZ4/03031 to PM), NIH Intramural Research Programs (ZIA NS003041 to AS and CY, ZICMH002888 to DG), International Neuroinformatics Coordinating Facility Seed Funding Grant (to PM and MR), a Spanish research project funded by the Spanish Ministry of Science, Innovation, and Universities (MCIU), State Research Agency (AEI), and European Regional Development Funds (FEDER) (ref. PID2019-105772GB-I00 AEI FEDER EU to GD), HBP SGA3 Human Brain Project Specific Grant Agreement 3 (grant agreement no. 945539 to GD), and the EU H2020 FET Flagship program and SGR Research Support Group support (ref. 2017 SGR 1545 to GD).

Author contributions

1294

C.L., X.T., A.C.S., and M.R. designed and supervised the study; Z.L. and Y.C. collected the ION MRI data; D.S., C.L., C.Y., and X.T. collected the NIH MRI data; M.R. and P.M. collected the neuronal tracing data; C.L., X.T., C.T., and Z.L. preprocessed and organized the MRI data; P.M., C.L., X.T., and H.J. preprocessed the neuronal tracing data; C.L. and X.T. constructed the functional-network maps; C.L., X.T., and M.R. constructed and evaluated the cortical parcellation maps; X.T., C.L., Y.P., and G.D. conducted the computational modeling; C.L. and F.F. developed the online web applications; D.G. and C.L. implemented the atlas into AFNI/SUMA. X.T. and C.L. wrote the original draft. C.L., X.T., A.C.S., and M.R. revised the manuscript.

Competing Interests

The Authors declare no competing interests

Figure Legends

Figure 1. Outline of Marmoset Brain Mapping Resource. The resource provides the awake test-retest resting-state fMRI data, in-vivo diffusion MRI data from the same marmoset cohorts, and the neuronal tracing data mapped onto the same MRI space at the voxel/vertex level. In addition to the datasets, it also supports the study of whole-brain functional networks and computational modelings, as well as functional connectivity-based parcellation of the cortex (Marmoset Brain Mapping Atlas Version 4) using a deep neural network for accurate individual mapping. As a comprehensive multimodal resource for marmoset brain research, we also provide an online platform to explore the relationship between structural and functional connectivity. This functionality is embodied in online interactive viewers.

Figure 2. Functional cortical networks and their parcellation maps. The identified networks include: (A) the ventral somatomotor, (B) the dorsal somatomotor, (C) the premotor, (D) the frontal pole, (E) the orbital frontal cortex, (F) the parahippocampus and temporal pole, (G-H) the auditory and salience-related network, (I-J) two trans-modal networks, including a putative frontoparietal network and the default-mode-network, and (K-O) visual-related networks from the primary visual cortex to higher-order functional regions. These networks were combined to form two network-parcellation maps (P-Q), which are dominated by the networks with short-range connectivity (P-Q, top rows) and with long-range connectivity (P-Q, bottom rows), respectively.

Figure 3. The functional connectivity boundary maps. (A). The population-based boundary maps from the ION, the NIH, and the combined datasets. These maps are highly consistent, with an average Dice coefficient of 0.7. (B). Boundary maps in the left hemisphere from four exemplar marmosets (two from the NIH cohort and two from the ION, including the flagship marmosets). (C-D). The heatmap of the average Dice's coefficients for both hemispheres between individuals and its distribution histogram. (E).

1339   The average Dice's coefficients change for both hemispheres with the number of runs in

1340   the same individuals.

1341

Figure 4. Marmoset Brain Mapping Atlas Version 4 (MBMv4). (A) The processing

procedure includes generating the population functional connectivity boundary maps,

defining the local minima for seeding, and generating parcels by the "watershed"

algorithm. (B) The resulting 96 functional connectivity parcels per hemisphere are

overlaid on the white matter surface and flat map of MBMv3 [16]. (C) Distance-controlled

boundary coefficient (DCBC) evaluation metric. All pairs of voxels/vertices were

categorized into "within" or "between" parcels (left panel) according to different brain

parcellations (MBMv1, MBMv4, Paxinos, and RIKEN atlases, right panel), and the

DCBC metric was calculated by the differences (within-between) in functional

connectivity as the function of distance on the surface (0-4 mm in steps of 0.5 mm).

Data are presented in mean +/- s.e.m.

1353

Figure 5. Mapping individual functional connectivity parcellation. (A) An overview of

individual Mapping based on the deep neural network approach. (B) MBMv4 Mapping of

each individual. Left panel: the concordance between the population MBMv4 and

individual parcellations (N=78, all hemispheres from 39 subjects). Data are presented

by the violin and the box plots (the 25th percentile: 0.9068; the 75 percentile: 0.92448),

in which the white point represents the average value 0.915 (the maximum value: 0.931;

minimum 0.900); Right panel: three examples of individual parcellations. The underlay

(color-coded) presents the population MBMv4, and the overlay (black border) shows the

individual parcellations. (C) Mapping of MBMv4 per session. Left panel: The

concordance between every individual parcellation and the corresponding parcellation

using one session data (N=78, all hemispheres from 39 subjects). Data are presented

by the violin and the box plots (the 25th percentile 0.865; the 75 percentile 0.877), in

which the white point represents the average value 0.874 (the maximum: 0.878; the

minimum 0.859); Right panel: representative parcellations of three sessions from one

marmoset. The color-coded underlay represents individual parcellation, while the black

border overlay shows the session-based parcellation. (D) The distance-controlled

boundary coefficient (DCBC) for the individual parcellation generated by the spatial registration (Spatial-reg, blue) and the deep neural network (DNN-reg, red). Top panel: the functional connectivity for all pairs of vertices within the same parcel and between parcels for DNN-reg and Spatial-reg, respectively. Bottom panel: the comparison of DNN-reg and Spatial-reg by DCBC. Data are presented in mean +/- s.e.m.

Figure 6. MBMv4 matches functional boundaries and preserves the topographical organization of the functional connectivity. (A) The visual activation maps from two monkeys are overlaid on the parcel boundaries from individual MBMv4 parcellation and Paxinos atlas (Left panel: monkey ID 15; Right panel: monkey ID 25). The scatter plots compare the boundary matching of the MBMv4 and the Paxinos atlas with the activation maps, measured by the shortest distance from every voxel in the borders of the activation maps to the parcel borders of the MBMv4 or the Paxinos atlas. The dashed black line represents the diagonal line, and the red line represents the linear fitting line. (B) The scatter plots in the left panel are the first two axes of gradients (the color scale of dots represents the scores of the first axis for every gradient), decomposed by the functional connectivities of the MBMv4 and the Paxinos atlas (the spectrum colors denote the gradient position in this 2D space). The heat maps of functional connectivity sorted by the scores of the first axis (gradient 1) are shown in the right panel.

Figure 7. A computational framework links the structural-functional connectivity according to different parcellations. (A) Application of whole-brain modeling, including an estimation of structural connectivity from neuronal tracing, diffusion MRI (in-vivo or ex-vivo) according to the Paxinos atlas or MBMv4, simulation of functional connectivity from structural connectivity by the Hopf bifurcation hemodynamic functions, and a similarity measure with empirical connectivity from resting-state fMRI. (B) Model fitting comparison in different spatial scales using the Paxinos versus the MBMv4 atlases. Individual examples using in-vivo diffusion MRI (round dots; an example animal marked in black dot), ex-vivo diffusion MRI (polygon), and neuronal tracing (star) show higher fitting values for MBMv4 versus the Paxinos atlas. (C) Examples of correlations between the simulated and empirical functional connectivity from (B); solid black lines

represent marginal regression lines. (D) Model fitting comparisons with (red) and
without (blue) an exponential distance rule (EDR) correction. From left to right, the plots
present the simulation results obtained with in-vivo diffusion MRI from the example
animal (the linear regression values R = 0.689, p = 9.27e-50 for the blue line; R = 0.413,
p = 1.28e-23 for the red); in-vivo diffusion MRI from all animals (R = 0.11, p = 6.05e-114
for the blue; R = 0.011, p = 4.17e-13 for the red); the ex-vivo diffusion MRI (R = 0.7, p =
2.69e-51 for the blue; R = 0.395, p = 2.19e-22 for the red); and the neuronal-tracing
dataset (R = 0.009, p=0.02). The line plots are presented as mean values +/- 95% C.I.
Dashed lines represent the 95% confidence interval. (E) Model fitting comparison with
EDR correction between the Paxinos atlas and MBMv4 in different spatial scales. (F)
Examples of correlation between the simulated and empirical functional connectivity
from (E).
1413

Marmoset Brain Mapping | Version 4

Data

New Data
Awake Resting-state fMRI

n = 39
710 runs
12117 mins

Major Updates
Voxel-level Neuro-tracing Data

n = 52
l = 143

New Data
In-vivo Diffusion MRI

n = 25

Mapping

ICA-based
Functional Network Parcellation

Gradient-boundary-map based
Group Cortical Parcellation

Deep-learning based
Individual Parcellation Generator

Neural Network

Evaluation

Compared with
Existing Cortical Parcellations

Distance-controlled boundary coefficient
— MBMv4   — MBMv1
— Paxinos   — RIKEN

Correlation value (Within - Between)

Distance (mm)
0 1 2 3 4

Evaluated by
Functional Gradients and Activations

Functional Gradient

Gradient 2

MBMv4

Paxinos atlas

Gradient 1
-0.3 -0.2 -0.1 0 0.1 0.2 0.3

Computational Modeling by
Structural and Functional Connectivity

Neural Tracing
Structural Connectivity

Simulated
Functional Connectivity

Cortical regions

Cortical regions

Online

Interactive Atlas Viewer
Parcellations and Atlases

Interactive Connectome Viewer
Connection Graphs

Interactive Connectome Viewer
Connectivity Matrices

A

Population Boundary Map → Local Minima → Region growing (Watershed algorithm)

B

Left   Right

C

Within

Between

Within

Within - Between

The distance-controlled boundary coefficient (DCBC)

Spatial distance (mm)

MBMv4
MBMv1
Paxinos
RIKEN

A

Individual Data

Vertices from Parcel-*i*

Vertices from Parcel-*i* and its neighbors

2 Classification for Parcel-*i*

3 Combination and remove the overlaps

4 Region growing (Watershed algorithm)

Individual Parcellation

Repeat for all parcels

Population Parcellation

Parcel-*i*

Vertices from Parcel-*i*

All Cortical Vertices

1

Deep neural network (DNN) training for Parcel-*i*

NOT

B

Consistent rate (%)

Across-Subjects

Monkey ID 4 (The flagship from ION)

Monkey ID 19 (The flagship from NIH)

Monkey ID 27

C

Consistent rate (%)

Across-Sessions

Monkey ID 4 - Session 1

Session 2

Session 3

D

Spatial-reg

DNN-reg

Within

Between

Vertex-Vertex correlation

Spatial distance (mm)

Within - Between

Difference in the correlation

Spatial distance (mm)

Spatial-reg
DNN-reg