An integrated resource for functional and structural connectivity of the marmoset brain

Xiaoguang Tian[1,*], Yuyan Chen[2], Piotr Majka[3,4], Diego Szczupak[1], Yonatan Sanz Perl[5,6], Cecil Chern-Chyi Yen[7], Chuanjun Tong[2], Furui Feng[2], Haiteng Jiang[8,9], Daniel Glen[10], Gustavo Deco[5,11,12,13], Marcello G. P. Rosa[4, *], Afonso C. Silva[1,*], Zhifeng Liang[2,14, *], Cirong Liu[2,14,15,16, *]

[1] Department of Neurobiology, University of Pittsburgh Brain Institute, University of Pittsburgh, Pittsburgh PA 15261, USA
[2] Center for Excellence in Brain Science and Intelligence Technology, Institute of Neuroscience, CAS Key Laboratory of Primate Neurobiology, Chinese Academy of Sciences, Shanghai, China
[3] Laboratory of Neuroinformatics, Nencki Institute of Experimental Biology of the Polish Academy of Sciences, 02-093 Warsaw, Poland
[4] Department of Physiology and Neuroscience Program, Biomedicine Discovery Institute, Monash University, Clayton, VIC 3800, Australia
[5] Center for Brain and Cognition, Computational Neuroscience Group, Department of Information and Communication Technologies, Universitat Pompeu Fabra, Roc Boronat 138, Barcelona, 08018, Spain
[6] Universidad de San Andrés, Vito Dumas 284 (B1644BID), Buenos Aires, Argentina
[7] Cerebral Microcirculation Section, Laboratory of Functional and Molecular Imaging, National Institute of Neurological Disorders and Stroke, National Institutes of Health (NINDS/NIH), Bethesda, MD 20892, USA
[8] Department of Neurobiology, Affiliated Mental Health Center & Hangzhou Seventh People's Hospital, Zhejiang University School of Medicine
[9] MOE Frontier Science Center for Brain Science and Brain-machine Integration, Zhejiang University, Hangzhou, China.
[10] Scientific and Statistical Computing Core, National Institute of Mental Health, National Institutes of Health (NIMH/NIH), Bethesda, MD 20892, USA
[11] Institució Catalana de la Recerca i Estudis Avançats (ICREA), Passeig Lluís Companys 23, Barcelona, 08010, Spain
[12] Department of Neuropsychology, Max Planck Institute for Human Cognitive and Brain Sciences, 04103 Leipzig, Germany
[13] School of Psychological Sciences, Monash University, Melbourne, Clayton VIC 3800, Australia
[14] Shanghai Center for Brain Science and Brain-Inspired Intelligence Technology, Shanghai, China
[15] Lingang Laboratory, Shanghai 200031, China
[16] University of Chinese Academy of Sciences, Beijing, China

* Co-Corresponding authors

Cirong Liu crliu@ion.ac.cn
Xiaoguang Tian txgxp88@gmail.com
Afonso Silva AFONSO@pitt.edu
Zhifeng Liang zliang@@ion.ac.cn
Marcello Rosa marcello.rosa@monash.edu

Abstract

Comprehensive integration of structural and functional connectivity data is required to model brain functions accurately. While resources for studying the structural connectivity of non-human primate brains already exist, their integration with functional connectivity data has remained unavailable. Here we present a comprehensive resource that integrates the most extensive awake marmoset resting-state fMRI data available to date (39 marmoset monkeys, 710 runs, 12117 mins) with previously published cellular-level neuronal tracing (52 marmoset monkeys, 143 injections) and multi-resolution diffusion MRI datasets. The combination of these data allowed us to (1) map the fine-detailed functional brain networks and cortical parcellations; (2) develop a deep-learning-based parcellation generator that preserves the topographical organization of functional connectivity and reflects individual variabilities, and (3) investigate the structural basis underlying functional connectivity by computational modeling. This resource will enable modeling structure-function relationships and facilitate future comparative and translational studies of primate brains.

Introduction

Mapping brain architecture is critical for decoding brain functions and understanding the mechanisms of brain diseases [1]. Non-human primate (NHP) neuroimaging provides a granular view of the evolution of the brain [2] and could overcome constraints of human neuroimaging by integration with "ground truth" data from cellular-resolution tracing [3].

As one of the few non-invasive imaging techniques capable of mapping whole-brain functional activity patterns, resting-state fMRI (rs-fMRI) provides insights into large-scale functional architecture [4]. However, data-sharing initiatives of NHP neuroimaging are still at an early stage, with existing open datasets of rs-fMRI data originating in different laboratories and collected for different purposes [5]. This leads to inconsistent imaging protocols and data quality, which hinder analyses across datasets. In addition, most presently available rs-fMRI datasets have been acquired in anesthetized animals, resulting in difficulties for cross-species studies, particularly relative to awake human brains [6]. The final barrier is the practical difficulty of training large numbers of NHPs to be fully awake during MRI scans [7, 8]. A platform for international collaborative research (PRIMatE RESOURCE EXCHANGE) was initiated to address these problems and promote open resource exchange and standards for NHP neuroimaging [5, 9].

The common marmoset monkey (Callithrix jacchus) has drawn considerable interest as an NHP species, offering many practical advantages for neuroscience research, including neuroimaging [10, 11, 12]. Previous work from our groups has contributed ultra-high-resolution ex-vivo diffusion MRI data [13], mesoscale neural tracing data [14], and structural atlases [15, 16, 17], which have enabled unprecedented precision in analyses of NHP brain anatomy. However, an essential component for understanding brain architecture has been missing: integrating these anatomical datasets with rs-fMRI. To address this limitation, and in alignment with a strategic plan developed by the NHP imaging community [8], we developed standardized protocols for imaging awake marmosets, which were implemented across two institutions, the National Institutes of Health (NIH), USA, and the Institute of Neuroscience (ION), China. This effort resulted in the largest awake NHP rs-

96 fMRI dataset to date, which is being made available through an open-access platform.

97 Furthermore, we integrated neuronal tracing and different diffusion MRI datasets into the

98 same MRI space, resulting in a comprehensive resource that allows us to explore the

99 relationships between the structural and functional connectomes by computational

100 modeling.

Results

The resource reported in this paper (summary in Fig. 1) is supported by a publicly

available standardized dataset. Following the same protocols for animal training and MRI

imaging, including the designs of the radiofrequency coil and MRI pulse sequences, we

acquired an extensive awake resting-state fMRI dataset to date from 39 marmosets of

two research institutes (13 from ION, age 3+/-1 years old; 26 from NIH, age 4+/-2 years

old; 12117 mins in total scanning, Supplementary Table 1 for details). This is also the

same range of ages used in our previous studies of structural connectivity [13, 14]. For test-

retest evaluation, we scanned multiple runs (17 mins/run) for each marmoset, resulting in

an essentially similar data quantity of two institutes (346 ION runs and 364 NIH runs) and

included two "flagship" marmosets with many runs (64 runs from the ION and 40 runs

from the NIH). Besides similar quantity, we comprehensively evaluated the data quality

across two datasets, which is essential for data harmonization [18]. We compared different

metrics in the SNR, temporal SNR (tSNR), spatial CNR, and head motions across the

sites/scanners (Supplementary Fig. 1-3, Supplementary Table 2). We found no significant

differences between the two datasets. We also compared the similarity of functional

connectivity across subjects (Supplementary Fig. 4-A) and sessions (Supplementary Fig.

4-B), which still revealed no significant differences. The quality assessments and quality

control (QA/QC) measurements demonstrated consistency and interpretability across

datasets, which makes them suitable for further analysis.

Using these datasets, we created a comprehensive map of resting-state brain networks

and a fine-grained functional cortical parcellation based on resting-state functional

connectivity. Furthermore, we developed a deep-learning-based approach to map the
population-based functional cortical parcellation onto individual brains. This allowed
investigation of the structural basis underlying functional connectivity. For this purpose,
we sampled the most extensive collection of NHP neuronal tracing data available (52
marmosets and 143 injections) onto the same MRI space at the voxel or vertex level and
integrated it with the same functional MRI data space mentioned above. In addition,
further enhancing the capacity of our resource, we also integrated extra high-resolution
ex-vivo diffusion MRI and in-vivo diffusion MRI data obtained at 25 marmosets from the
same cohort. On this basis, we investigated the relationship between structural and
functional connectivity using a whole-brain computational model. To facilitate the user to
explore connectomes reported in this paper and compare them with other connectomes
[13, 14, 15], we make online connectome viewers (connectome.marmosetbrainmapping.org).
138

139 **Mapping functional brain networks**
140

Identifying functional networks of areas showing highly correlated fMRI signals is critical
to characterizing the brain architecture. Using the independent component analysis (ICA),
a data-driven approach for separating independent patterns in multivariate data, we
identified 15 cortical networks from awake resting-state fMRI data (Fig. 2; Supplementary
Figs. 5-6 include the power spectrum). All specified components showed clear neural-like
patterns spatially (all peaks located in the cortical gray matter) and temporally (no patterns
of artifacts or noises), as shown in Supplementary Fig. 6. We also conducted the ICA
separately on each dataset. We found that both the ION and the NIH data reveal these
components with similar spatial patterns and temporal CNR (tCNR, the definition of this
metric is demonstrated in the method part of Data harmonization, the results are in
Supplementary Fig. 7), demonstrating the reproducibility of networks and the consistency
of the two datasets.
153

The details of the 15 cortical networks were as follows. Six functional networks were
characterized by short-range connectivity, including the ventral somatomotor (Fig. 2A),
the dorsal somatomotor (Fig. 2B), the premotor (Fig. 2C), the frontopolar (Fig. 2D), the

orbitofrontal (Fig. 2E), and the parahippocampal/ temporopolar cortex (Fig. 2F) networks.
The next two components are the auditory and salience-related networks, the first being
primarily located in the auditory and insular cortices, and weakly coupled with the anterior
cingulate cortex (Fig. 2G), and the second (Fig. 2H) encompassing the anterior cingulate
cortex. In addition, we also identified two trans-modal networks (Fig. 2I-J), including
association areas in the dorsolateral prefrontal cortex (dlPFC), rostral premotor cortex,
lateral and medial parietal cortices, and temporal cortex. One of these is most likely the
frontoparietal-like network (Fig. 2I), which has not been identified in previous studies [19,
20], and the other is the default-mode network (DMN, Fig. 2J), which we characterized
extensively in a previous study [21]. The remaining five networks represent the first
complete mapping of visual-related functional networks of the marmoset cortex (Fig. 2K-
O). Three networks included the primary visual cortex and parts of extrastriate areas
related to far peripheral vision (Fig. 2K), near-peripheral vision (Fig. 2L), and foveal vision
(Fig. 2M). The other two networks involve hierarchically higher visual areas (Fig. 2N-O),
such as V3, V4, the inferior temporal cortex, the adjacent polysensory temporal cortex,
and vison-related frontal regions.
173
Based on their spatial overlap patterns and connectivity strengths (normalized Z-scores),
we combined the 15 cortical networks into network-parcellation maps (Fig. 2P-Q). Due to
local connectivity being stronger than long-range connectivity, the primary map (Fig. 2P-
Q, top rows) is dominated by the short-range networks (i.e., Fig. 2G, I, J, K, L, I, and O).
Thus, we created a second connectivity map (Fig. 2P-Q, bottom rows) to cover the long-
range connectivity not captured by the primary map. The two network-parcellation maps
characterized the entire cortical networks, and will likely be of value for future functional
connectivity studies of the marmoset brain.
182

183 **Mapping functional connectivity boundaries**
184
The brain network maps provided a global view of cortical functional organization. Next,
we aimed to characterize the cortex at a finer local scale. For this, we used a functional
connectivity boundary mapping approach to identify putative borders for the functional

parcels [22, 23, 24] as an efficient way to map transitions in functional connectivity. Population

boundary maps based on the ION, the NIH, or combined datasets are visually similar,

presenting clear functional connectivity borders (Fig. 3A), and were highly reproducible

with average Dice's coefficients for both hemispheres: 0.7 (ION-NIH), 0.71 (ION-Both),

and 0.69 (NIH-Both), respectively (Supplementary Fig. 8). However, although consistent

at the population level, boundary maps indicate variability across individuals (Fig. 3B),

with an average Dice's coefficient of 0.3842 for both hemispheres (Fig. 3C-D), which is

significantly lower than those for the population. We also found across-session variability

in the same individual, but more scanning runs efficiently enhanced the reproducibility

(Fig. 3E). Therefore, these results suggest that both individual and across-session

variability contribute to the low consistency of individual boundary maps and that using

test-retest data is essential for improving the reliability of maps.

Generation of functional connectivity parcels (Marmoset Brain Mapping Atlas

Version 4, MBMv4)

Because the population boundary maps are more reproducible than individual maps, we

used the combined ION-NIH population boundary map to generate cortical functional

connectivity parcels. By the detection of the local-minima [22], "watershed-flood" region

growing [25], and semi-manual optimization of parcel boundaries (Fig. 4A), we obtained 96

parcels per hemisphere (Fig. 4B). Since we processed each hemisphere independently,

we compared the similarity of the parcellations of the two hemispheres. The

hemispherical parcellations are similar in the parcel sizes (Supplementary Fig. 9-A;

Wilcoxon paired signed-rank test, N=96, p=0.7981; Supplementary Fig. 9-C for the size

comparison in the subject's native space) and functional connectivity patterns between

vertices within the same parcel (Supplementary Fig. 9-B; Wilcoxon paired signed-rank

test, N=96, p=0.411). For continuity with previously released resources [13, 15, 16], we named

this functional connectivity-based parcellation of the cortex "Marmoset Brain Mapping

Atlas Version 4" (MBMv4). We also provide an online viewer to visualize parcellations,

including previous versions (atlasviewer.marmosetbrainmapping.org).

To estimate the validity of the generated functional parcels, we used the distance-
controlled boundary coefficient (DCBC) [26]. The basic idea of DCBC is that when a
boundary divides two functionally homogenous regions, for any equal distance on the
cortical surface, the functional connectivity pattern between vertices within the same
parcel should be higher than that between vertices in different parcels (Fig. 4C). In other
words, a higher DCBC (within - between) means higher within-parcel homogeneity and
higher between-parcel heterogeneity. We calculated the DCBC between the vertex pairs
using a range of spatial bins (0–4 mm) with a 0.5 mm step (the spatial resolution of the
rs-fMRI data). Here, we compared the fit of the functional map represented by MBMv4
with existing structural cortical parcellations, including MBMv1 atlas [15], the digital
reconstruction of the Paxinos atlas [15, 27], and the RIKEN atlas [28]. The result of DCBC in
Fig. 4C demonstrates that MBMv4 has the best performance for the presentation of
functional connectivity (the average DCBC values were 0.0186, 0.0135, 0.0177, 0.0330
for RIKEN, MBMv1, Paxinos, and MBMv4 atlas; multiple comparisons for One-Way
ANOVA $F_{(3,8556)}=22.44$, $p=1.81 \times 10^{-14}$).

Mapping MBMv4 in Individual Brains by Deep Neural Networks

To overcome the limitation of variable individual boundary maps (see Fig. 3C-E), we
employed a deep-learning approach for the individual Mapping from MBMv4 (Fig. 5A).
First, based on the population-level whole-brain functional connectivity, we trained a deep
neural network classifier for each parcel to learn the associated fingerprint of functional
connectivity. Then, the trained networks distinguished the goal parcel for every marmoset
based on the corresponding functional connectivity of the searching area, consisting of
the goal parcel and its neighbors. Due to the overlap of searching areas, vertices could
belong to multiple parcels. Therefore, we only kept these vertices attributed to a single
parcel as the seeds for regional growth by the "watershed" algorithm. This iterative region-
growing procedure would assign all vertices to a parcel, resulting in an individual cortical
parcellation.

Since individual parcellations should be close to the population definition [29, 30], we

compared the population-based MBMv4 parcellation and the automatically generated

individual parcellations. By calculating the percentage of vertices sharing the same labels

from both hemispheres (a metric of concordance), we found that the individual

parcellations from all marmosets are similar to MBMv4 with an average of 90%

concordance (Fig. 5B, the violin/box plot on the left, the examples on the right). Using the

test-retest dataset, we revealed the consistency of the individual parcellations across

different sessions (Fig. 5C, the violin/box plot on the left, and examples on the right). The

across-session analysis yielded an average of 86.7% concordance, lower than the

average value of 91.3% across-individual similarities. Furthermore, we observed that the

lateral prefrontal cortex and occipital-temporal cortex had higher across-individual and

across-session mapping variabilities (Supplementary Fig. 10), consistent with previous

findings in humans [22, 31]. Thus, the deep-learning approach efficiently adjusts the parcel

borders to reflect the individual variabilities while maintaining high consistency with the

population parcellation.

We also used the DCBC to evaluate whether the border adjustment of the individual

parcellation captured the specific features of individual functional connectivity patterns.

We assumed that the deep learning-based method (DNN-reg) should result in a higher

DCBC than the direct spatial registration of MBMv4 (Spatial-reg). Figure 5D (Top panel)

presents the functional connectivity for the pairs of vertices within the same parcel

(average correlation values within the same surface length 0-4 mm were 0.8331 and

0.8172 for DNN-reg and Spatial-reg) and between different parcels (average correlation

values were 0.8256 and 0.8171 for DNN-reg and Spatial-reg). Thus, the DNN-reg had

higher DCBC (within-between) than the Spatial-reg (Fig. 5D, bottom panel; the average

DCBC values were 0.0167 and 0.0085 for the DNN-reg and the Spatial-reg, respectively;

multiple comparisons for One-Way ANOVA $F_{(1,2512)}$=20.35, p=6.74×10^{-6}). In sum, the

border adjustment by the proposed deep learning network reflects individual functional

connectivity patterns.

MBMv4 reflects accurate functional and topographical organizations

As evaluated from functional connectivity, MBMv4 provides a more accurate reflection of the MRI-based functional parcellation of the cortex than current histology-based atlases. To further verify this reliability, we took a task activation map during the presentation of movie [32] which encompassed 10 deg × 8 deg of the visual field. This activation map was then registered onto the same individual MBMv4 map and the histology-based Paxinos atlas to examine the spatial overlap between the activations and functional parcels. As a result, we found that the MBMv4 has a good correspondence with task activations by visual inspection, such as the co-activation of foveal areas V1 and MT, and the temporal parcels (Fig. 6A, flat maps). Additionally, by measuring the shortest distances from every vertex in the boundary of the activation map to the atlas boundaries (MBMv4 or Paxinos boundaries), we found that the parcel borders of MBMv4 have higher consistency with the activation map than the Paxinos atlas (Fig. 6A, the scatterplots; Wilcoxon paired signed-rank test: Monkey ID 25, N=878, $p=3.07×10^{-40}$ for the left hemisphere; N=816, $p=6.11×10^{-26}$ for the right hemisphere. Monkey ID 15, N=826, $p=2.22×10^{-25}$ for the left hemisphere; N=850, $p=2.95×10^{-53}$ for the right hemisphere). Thus, MBMv4 reflects functional differences that cytoarchitectonics does not capture, possibly because the latter contains the full visual field representations. Thus, the MBMv4 may provide functional-related contrast that can help enhance the precision of cross-species studies [33].

Besides the clear functional boundaries, MBMv4 also preserved the topographical organization of the functional connectivity. Recent methodological developments have allowed complex brain features to be mapped to the low-dimensional representations as gradients [34], and these gradients characterized the topographical organization of the functional brain connectome from unimodal to trans-modal networks. If the atlas complies with this topographical organization, it should be able to identify such gradients. As shown in Fig. 6B left panel, MBMv4 results in a pattern of gradient spectrum for functional connectivity. In contrast, we did not find a gradient pattern based on the Paxinos et al. (2012) atlas (right panel in Fig. 6B). Therefore, MBMv4 offers an alternative view to

310 understanding the functional connectome of the marmoset brain by reflecting the
311 characteristics of functional connectivity.

313 MBMv4 is an essential link between the functional and structural
314 connectivity

Since MBMv4 offers a more comprehensive scheme to study the functional connectome,
it is worth linking it to structural information to investigate the relationships between
structural and functional connectivity. We have also provided online viewers to facilitate
the comparison between them (connectome.marmosetbrainmapping.org). Furthermore,
we used a whole-brain computational model [35, 36, 37] to explore structural and functional
connectivity relationships. The processing procedure is shown in Fig. 7A. First, we
established structural connectivity values using the in-vivo diffusion MRI, the ex-vivo ultra-
high-resolution diffusion MRI, or the neuronal tracing datasets fitted to either the MBMv4
or the Paxinos et al. (2012) atlas parcellations [14]. Then, we used a Hopf bifurcation
hemodynamic model [36] to simulate the functional connectivity of every brain parcel or
region based on their structural connectivity and compare it with the empirical functional
connectivity from the actual resting-state fMRI data. We used Pearson's correlation to
measure the similarity between the simulated and the empirical functional connectivity.
Additionally, we used group-average functional connectivity as an empirical observation
for the ex-vivo diffusion MRI and neuronal tracing dataset and individual functional
connectivity for the individual in-vivo diffusion MRI.

The modeling results in Fig. 7B-C demonstrate that: (1) the MBMv4 parcellation provides
the best fitting value (R=0.721) compared to the Paxinos atlas (R=0.638) for the extra
high-resolution ex-vivo diffusion MRI dataset (the polygon in Fig. 7B and results in Fig.
7C); (2) the MBMv4 parcellation fits the in-vivo diffusion MRI datasets better than the
Paxinos atlas in a variety of individual simulations (all circles in Fig. 7B: the average fitting
values from 25 animals were R=0.4707 for MBMv4 and R=0.3659 for Paxinos atlas); and
(3) both parcellations performed equally (R=0.525 for MBMv4 and R=0.472 for Paxinos

atlas) fitting the cellular connectivity data from the aggregated neuronal tracing (the star

in Fig. 7B and results in Fig. 7C). However, no matter which structural data was used for

estimating functional connectivity, we found that the modeling predicted by MBMv4 fits

the empirical functional connectivity data better than the Paxinos atlas (Fig. 7B; Wilcoxon

paired signed-rank test: N=27, p=0.002947).

Due to its inherent limitation for tracking long connections, the diffusion tractography was

more affected by distance, with longer connections having lower FC fitting values (Fig.

7D, blue lines). Based on studies demonstrating that cellular-resolution connections

follow an exponential distribution of projection lengths [38, 39, 40], we introduced an

exponential distance rule (EDR) to our model to compensate for the distance effect (Fig.

7D, redlines). Notably, the EDR fully corrected the distance effect in neural tracing data

(Fig. 7D), suggesting that the neural tracing data might be a more reliable bridge to link

structural and functional connectivity of long-range connections. Still, as shown in Fig.

7E-F, the EDR-constrained modeling based on MBMv4 fits the empirical functional data

better than the Paxinos atlas (Wilcoxon paired signed-rank test: N=27, p=0.01947),

similar to the modeling without EDR (Fig. 7B), showing that MBMv4 is better suited for

investigating the relationship between structural and functional connectivity.

Discussion

There are challenges in adapting well-established approaches for human neuroimaging

to NHP neuroimaging [5]. The present study used effective and practical animal training

and imaging protocols to scan a large cohort of marmosets. Despite the different MRI

scanners (7T and 9.4T) in two institutions (ION and NIH), the protocol produced similar

data quality, suggesting the compatibility of our approach (see Methods for details,

Supplementary Fig. S1-S4). Therefore, we pooled the in-vivo resting-state fMRI datasets

from both institutions (details in Supplementary Table 1) to create the most

comprehensive functional connectivity dataset of the NHP brain to date, which was

integrated with the in-vivo diffusion MRI of the same cohort, as well as the highest

resolution ex-vivo diffusion MRI [13] and the most extensive mesoscale retrograde neuronal

tracing [14] available. This resource expedites the mapping of marmoset brains and will

allow cross-species comparisons.

Like the human cortex, the marmoset cerebral cortex is composed of large-scale

functional networks. However, the first awake resting-state fMRI study of the marmosets

[19] found only 12 functional networks (10 cortical networks), and another ICA-based study

described 8 brain networks, possibly due to the influence of anesthesia [20]. Based on a

more extensive awake rs-fMRI dataset, the present study mapped the large-scale

functional networks and built the first network-based parcellation, providing a more

comprehensive description of functional networks in the marmoset brain, including a total

of 15 cortical networks. Moreover, based on functional connectivity boundary maps, we

created a population-based cortical parcellation on a fine scale (MBMv4; Fig. 4) with 192

distinct parcels (96 per hemisphere). The further verification of hemispheric symmetry is

also done by inverting warping the standard average surfaces to every subject's native

space in Supplementary Fig. 9-C. A previous study of the human cerebral cortex identified

422 discrete functional connectivity parcels using the same approach, 206 in the left

hemisphere and 216 in the right hemisphere [22]. Therefore, our results may reflect that the

asymmetry in marmosets may be smaller than in humans, as expected from previous

analyses based on anatomical measurements [41] and the evidence that the number of

cortex subdivisions increases with brain volume [42].

It is important to emphasize that our functional parcels do not correspond to the traditional

cytoarchitectonic definition of the cortical areas [43, 44]. Consistent with many brain

parcellations generated by non-invasive neuroimaging [22, 29, 45, 46, 47], our functional parcels

most likely reflect a different type of computational sub-units, agreeing with the idea that

the brain is organized in multiple scales [48, 49]. Therefore, compared with other available

structural atlases, MBMv4 captures the organization of functional connectivity accurately.

For example, MBMv4 achieved better task correspondence (Fig. 6A) due to a strong link

between task-fMRI and rs-fMRI [50, 51, 52]. Another evidence for the accuracy of the MBMv4

parcellation is the topographical gradient organization of functional connectivity (Fig. 6B).

Finally, MBMv4 better links the structural and functional connectivity, as demonstrated by
our modeling simulation (see Fig. 7).

Consistent with the previous findings in humans [22, 29, 47], the parcels defined in MBMv4 do
not follow the boundaries of cytoarchitectonic areas, thus demonstrating an essential
difference between anatomical features and functional connectivity. For example, the
somatomotor cortex is parcellated into subregions that appear to correspond to
representations of the face, forelimbs, hindlimbs, and trunk musculatures across multiple
areas, and areas such as V1 and V2 are subdivided into several functional parcels
according to the representation of eccentricity in visual field representation, which is
contiguous across areas [53] but may include discontinuities [54]. Previous studies revealed
that some topographically organized cytoarchitectonic areas could be dissociated from
the resting-state functional responses [55, 56]. Therefore, the present MBMv4 should be
considered a functional connectivity description, providing complementary information
about a type of organization that cannot be observed via classical anatomy.

An essential goal of this study was to reflect individual characteristics by creating parcels
from each individual's data. Although the boundary map-derived parcels could be used
for individual analysis, we found that the subject boundary maps had significant variations
and that the reproducibility became lower than the group-level map (Fig. 3). This finding
emphasizes the need to acquire large amounts of data for the reliable test-retest of the
individual boundary map. Accordingly, we developed a deep neural network to map
reliable population-level MBMv4 into every individual nonlinearly. As a result, we
demonstrated good reliability in the test-retest dataset (across sessions from the same
individual; see Fig. 5C) and the applications of task-fMRI activation mapping from the
same individuals (Fig. 6A). Importantly, the locations of the most variable functional
parcels are in the lateral prefrontal cortex and lateral temporal-occipital cortex (see
Supplementary Fig. 10), corresponding to previously reported exceptionally high inter-
subject variability resting-state functional connectivity patterns [31]. Moreover, these
regions co-locate, expanding preferentially in primate evolution [57] and maturing later in
postnatal development [58]. As resting-state functional connectivity can be altered by many

biological features, including behavioral context during development associated with

phenotypic correlations, a better understanding of the causes of inter-subject parcel

variation will be an important focus for our future work.

In addition to the functional connectome mapping, we integrated all currently available

structural connectome datasets, including the in-vivo diffusion MRI, the ex-vivo high-

resolution diffusion MRI [13], and the mesoscale tracing dataset [14]. Using multimodal data

allowed us to investigate the relationship between functional and structural connectivity

with unprecedented detail. Using whole-brain modeling [36, 37], we observed that the

structural connectivity simulated functional connectivity based on MBMv4 had a high

coherence with empirical data, no matter which types of structural connectivity were used

(Fig. 7B-C and examples Fig. 7E-F). The finding corroborates the conclusion that MBMv4

reflects meaningful computational sub-units from the view of whole-brain functional

connectivity. Meanwhile, we also found room for modeling performance improvement

through the detailed estimation of structural connectivity. For example, compared with in-

vivo diffusion MRI data, the ultra-high-resolution ex-vivo diffusion MRI data from a brain

sample could provide more thorough structural information. Notably, we should also be

careful about the influence of the distance when we use the diffusion dataset to simulate

functional connectivity because the diffusion tract may generate the fiber cut and so on in

the long-distance tract (Fig. 7D, blue curves). To overcome this shortcoming, we

introduced a correction factor for structural connectivity based on the exponential distance

rule (Fig. 7D, red curves). Nevertheless, the simulation results in Fig. 7B and 7E

demonstrated the best fitting by the present MBMv4, no matter which datasets were used.

Furthermore, since our modeling is simple, with only two parameters that avoid overfitting

simulation (see our method description), the whole-brain model could be an efficient tool

with broad applications to link structure and function for future studies.

Although we provided the most comprehensive multimodal data resource for mapping the

marmoset connectome, our current study still faced limitations that need to be addressed

by future work, which is essential for both experimenters and users of non-human primate

neuroimaging. First is the data collection. The population used to generate the MBMv4

was sex-biased (31 males vs. 8 females) due to the priority of colony expansion worldwide.

The neuronal tracing data were also limited, not covering all cortical regions and missing

subcortical information. Including neuronal-tracing data is critical for accurately mapping

the future fully structural connectome. Finally, our parcellation only used the resting-state

functional connectivity information, as in many human studies[22, 45]. Therefore, multimodal

brain parcellation incorporated structural contrasts, especially the T1w/T2w myelin map,

multiple task-fMRI data, gene expression data, etc., will be necessary to fully capture the

anatomical and functional architectures of the marmoset brain.

Methods

Data Collection and Preprocessing

Animals and MRI scanning

Experimental procedures followed policies established by the Chinese Laboratory Animal

– Guideline for Ethical Review of Animal Welfare (ION data) or the US Public Health

Service Policy on Humane Care and Use of Laboratory Animals (NIH data). All

procedures were approved by the Animal Care and Use Committee (ACUC) of the

Institute of Neuroscience, Chinese Academy of Sciences (ION data), or the ACUC of the

National Institute of Neurological Disorders and Stroke, National Institutes of Health (NIH

data). The respective ACUC-approved protocols specify group size numbers based on a

power analysis to detect differences between animals to ensure rigor and reproducibility

of the results while minimizing the number of animals used in the study. Our studies are

powered to detect inter-individual differences. The experimental designs are typically two

or 3-factor ANOVAs. Values of $p<0.05$ are considered statistically significant. The number

of animals used is the minimum necessary to provide reliable estimates of inter-individual

effects based on power considerations. Typically, sample size estimates are based on

the number of animals needed to achieve a power of 0.80 for moderate effect size and

0.99 for large effect size. Both marmoset colonies are socially housed to ensure

psychosocial well-being and are offered a varied diet that includes food treats. In addition,

dedicated husbandry and veterinary teams interact with the animals daily as part of the

psychological enrichment plans approved by the ACUCs of both institutions.

The data acquisition procedure from both centers followed the same animal training

protocol, 8-element radiofrequency (RF) coil design [59], and MRI scanning protocols.

Thirteen marmosets (12 males and 1 female) were recruited from the ION cohort, from

which we generated 62 awake resting fMRI sessions and 349 runs (17 min per run). As

three of the 349 runs had extensive head motions (> 10% time points were motion

censored based on the preprocessed pipeline described below), we excluded the three

runs from the analysis, resulting in a total of 346 runs (see Supplementary Table 2 for the

summary of the head-motion per run). Twenty-six marmosets (19 males and 7 females)

were recruited from the NIH cohort to produce 51 awake resting-state fMRI sessions and

364 runs. Therefore, the NIH and ION data had a comparable number of valid runs. The

two datasets included 39 marmosets with 113 sessions, 710 valid fMRI runs, and 12117

mins of total scan time. Detailed demographic information is provided in Supplementary

Table 1. All marmosets underwent a 3- to 4-week acclimatization protocol as previously

described [60]. After completing the training, all marmosets were properly acclimated to

laying in the sphinx position in an MRI-compatible cradle. Their heads were comfortably

restrained with 3D-printed anatomically conforming helmets that allowed the resting-state

fMRI (rs-fMRI) data acquisition as the animals lay relaxed in their natural resting position.

All 39 marmosets were imaged using identical rs-fMRI protocols and pulse sequences,

except for a minor adjustment in the echo time (TE) to accommodate hardware

differences between the ION and the NIH gradient sets. The ION marmosets were

scanned in a 9.4T/30cm horizontal MRI scanner (Bruker, Billerica, USA) equipped with a

20 cm gradient set capable of 300 mT/m gradient strength and an 8-channel phased-

array RF coil [59] custom-built for marmosets (Fine Instrument Technology, Brazil). Multiple

runs of rs-fMRI data were collected in ParaVision 6.0.1 software using a 2D gradient-echo

(GE) EPI sequence with the following parameters: TR = 2 s, TE = 18 ms, flip angle = 70.4°,

FOV = 28 × 36 mm, matrix size = 56 × 72, 38 axial slices, slice thickness = 0.5 mm, 512

volumes (17 min) per run. The GE-EPI fMRI data were collected using two opposite

phase-encoding directions (LR and RL) to compensate for EPI distortions and signal

dropouts. Two sets of spin-echo EPI with opposite phase-encoding directions (LR and

RL) were also collected for EPI-distortion correction (TR = 3000 ms, TE = 37.69 ms, flip

angle = 90°, FOV = 28 × 36 mm, matrix size = 56 × 72, 38 axial slices, slice thickness = 0.5

mm, 8 volumes for each set). After each rs-fMRI session, a T2-weighted structural image

(TR = 8000 ms, TE = 10 ms, flip angle = 90°, FOV = 28 × 36 mm, matrix size = 112 × 144,

38 axial slices, slice thickness = 0.5 mm) was scanned for co-registration purposes.

The NIH marmosets were scanned in a 7T/30cm horizontal MRI (Bruker, Billerica, USA)

equipped with a 15 cm customized gradient set capable of 450 mT/m gradient strength

(Resonance Research Inc., Billerica, USA) and an 8-channel phased-array RF coil

custom-built for marmosets [59] with identical coil geometry to the one used by ION. Multiple

runs of rs-fMRI data were collected during each scanning session in ParaVision 6.0.1.

software using a 2D gradient-echo (GE) EPI sequence with the following parameters:

TR = 2s, TE = 22.2 ms, flip angle = 70.4°, FOV = 28 × 36 mm, matrix size = 56 × 72, 38

axial slices, slice thickness = 0.5 mm, 512 volumes (17 min) per run. The GE-EPI fMRI

data were collected using two opposite phase-encoding directions (LR and RL) to

compensate for EPI distortions and signal dropouts. Two sets of spin-echo EPI with

opposite phase-encoding directions (LR and RL) were also collected for EPI-distortion

correction (TR = 3000 ms, TE = 36 ms, flip angle = 90°, FOV = 28 × 36 mm, matrix size =

56 × 72, 38 axial slices, slice thickness = 0.5 mm, 8 volumes for each set). After each rs-

fMRI session, a T2-weighted structural image (TR = 6000 ms, TE = 9 ms, flip angle = 90°,

FOV = 28 × 36 mm, matrix size = 112 × 144, 38 axis slices, slice thickness = 0.5 mm) was

scanned for co-registration purposes. Furthermore, multishell diffusion MRI (DTI)

datasets were collected using a 2D diffusion-weighted spin-echo EPI sequence with the

following parameters: TR = 5.1 s, TE = 38 ms, number of segments = 88, FOV = 36 × 28

mm, matrix size = 72 × 56, slice thickness = 0.5 mm, a total of 400 DWI images for two-

phase encodings (blip-up and blip-down) and each has 3 b values (8 b = 0, 64 b =2400,

and 128 b = 4800), and the scanning duration was about 34 min. The multishell gradient

sampling scheme was generated using the Q-shell sampling method [61].

Data Preprocessing

The rs-fMRI datasets were preprocessed by the customized script involving AFNI [62], FSL
[63], ANTs [64], and Connectome Workbench [65]. In brief, the rs-fMRI data were slice-timing-
corrected and motion-corrected by the "3dTshift" and "3dvolreg" commands of AFNI and
corrected for EPI distortions by the "top-up" command of FSL (see our examples in
Supplementary Fig. 11). The rs-fMRI datasets were further preprocessed by regressing
linear and quadratic trends, motion parameters and their derivatives, and motion-sensor
regressors (any TRs and the previous TRs were censored if the detection motion was >
0.2 mm and temporal outlier > 0.1). Note that, for the motion measurements, we
calculated the weighted euclidean norm of six motion parameters with a 0.25 weight for
the three rotation degrees (yaw, pitch, and roll) according to the relative head radius of
the marmosets compared to humans. White matter and cerebrospinal fluid signals were
removed, and the rs-fMRI datasets were band-pass filtered (0.01–0.1 Hz). The above
nuisance signal regression and band-passing filtering were carried out by the
"3dDeconvolve" and "3dTproject" commands in AFNI. Next, the preprocessed data were
spatially normalized to the template space of our Marmoset Brain Atlas Version-3
(MBMv3) by the "antsRegistration" routine of ANTs [16]. The spatial normalization
concatenated multiple transformations, including 1) rigid-body transformation of each
fMRI run to the T2-weighted image acquired at the end of each session, 2) rigid-body
transformation of T2-weighted images from each session to a cross-session averaged
T2-weighted image from each animal, 3) affine and nonlinear transformation of the
averaged T2-weighted image from each animal to the T2w template of our MBMv3 space.
Finally, all preprocessed data were mapped to 3D brain surfaces of the MBMv3 using the
Connectome Workbench (wb_command -volume-to-surface-mapping function and ribbon
constrained mapping algorithm), normalized (subtract mean and divide by standard
deviation) and concatenated per session before the boundary mapping described below.
The preprocessed data were smoothed with 1mm FWHM using 3dBlurInMask (for volume
data) and wb_command -cifti-smoothing (for surface data), respectively, before the
network analysis and cortical parcellation.

The in-vivo diffusion MRI dataset was preprocessed by the DIFF_PREP, DR_BUDDI, and DR_TAMAS pipelines of TORTOISE [66]. The DIFF_PREP and DR_BUDDI routines incorporated correction for eddy-currents- and EPI-induced distortions using pairs of diffusion data acquired with opposite phase encoding (blip-up and blip-down) and the T2-weighted image and merging the preprocessed pairs into one dataset. The nonlinear spatial registration from the individual space to the DTI template of our MBMv3 space [16] was carried out using the DR_TAMAS routine of TORTOISE. The registration information was then used to transform multiple atlases into the individual space for diffusion tractography.

All diffusion trackings were performed using the iFOD2 method of the software Mrtrix3 [67]. The response function of each preprocessed diffusion MRI data was calculated by the "dhollander" method of the "dwi2response" command, and then the fiber orientation distributions (FOD) were estimated using spherical deconvolution by the multishell multi-tissue CSD method of the "dwi2fod" command. Finally, region-to-region tractography was performed using the iFOD2 method of the "tckgen" command. For each pair of cortical regions, diffusion tractography was conducted by using one region as the seed and the other as the target, and vice-versa. Thus, each pair of regions generated two sets of tracking probability maps, which were normalized by the total streamlines selected, and the two probability maps were averaged into a single map to represent the final map of the connection between the two regions. Finally, all pairs of connections formed the whole cortical structural connectome for computational modeling.

The neuronal tracing data were mapped onto the histological NM template from our previous study [17]. The NM template is a population-based 3D cortical template generated from Nissl-stained serial sections of 20 marmosets. Since the NM template only covers the cortex and has Nissl-stain contrast and a 75µm isotropic high spatial resolution, its direct spatial transformation to our in-vivo MBMv3 template is inaccurate. Thus, we modified the 80µm isotropic ultra-high-resolution MTR template of our Marmoset Brain Atlas Version-2 (MBMv2) atlas [13] to remove the parts of the brain that were not covered

in the NM template, including the cerebellum, brainstem, and parts of subcortical
structures. This step increased the accuracy of registration between the NM template and
the MBMv2 template. Then, the ex-vivo MTR template of the MBMv2 was transformed
into the in-vivo myelin-map template of our MBMv3. By concatenating the two
transformations (the NM-to-MBMv2 and the MBMv2-to-MBMv3), we accurately converted
the neuronal tracing data from the NM template to the MRI template. We then mapped
the neuronal tracing data onto the MBMv3 cortical surfaces. For the above registrations,
we used the CC similarity metric as the cost functions and three-stage alignments (rigid
alignment, affine alignment, and nonlinear SyN transformations), which were also the
default options antsRegistrationSyN.sh. An example of registration results is shown in
Supplementary Fig. 12.

The instruction and example code for the data preprocessing pipeline is provided via the
resource webpage https://marmosetbrainmapping.org/data.html (ReadMe and Codes
sections), allowing the user to replicate our protocols.

Data harmonization across NIH and ION datasets

We calculated a series of indices to test the data harmonization across different datasets
(NIH and ION). They include the single time points SNR, mean images (average across
time for one fMRI run), SNR, temporal Signal Noise Ratio (tSNR, from one fMRI run),
Contrast to Noise Ratio (CNR, the mean of the gray matter intensity values minus the
mean of the white matter intensity values divided by the standard deviation of the values
outside the brain), temporal contrast to Noise Ratio (tCNR, the variance of optimal resting-
state fMRI components after ICA contrast to the noisy component), the Fiber (Foreground
to Background Energy Ratio: the variance of voxels inside the brain divided by the
variance of voxels outside the brain), head motion and the whole-brain functional
connectivity across subjects and sessions.

Functional Networks, Cortical Parcellation and Network Modeling

Brain network identification by the Group-ICA

Independent Component Analysis (ICA) was performed by the Group-ICA routine of the GIFTI software (https://trendscenter.org/software/gift/) to identify the brain networks using a number of different component settings. First, preprocessed data without bandpassing and regression of nuisance covariates were group-ICA analyzed with increasing numbers of ICA components from 20 to 80 in a step of 10. We tested the reliability of different ICA methods, including the default "Infomax" ICA algorithm or "ICASSO" group-ICA method, on different datasets (the NIH dataset, the ION dataset, or combined both datasets) and obtained consistent results regardless of the ICA setting or dataset used. Finally, every resulting component from Group-ICA analyses was visually inspected and sorted according to its neuroanatomical features. Since the sorted elements were highly consistent across different settings of ICA-component numbers, we selected the best component to represent every labeled network. We also did manual correction before creating the final network parcellation; for example, when the left and right parts of the same network were separated into two components, we merged them into one network by averaging their maps. We identify 15 cortical resting-state networks from the group-ICA analysis (Fig. 2A-O and Supplementary Fig. 5).

We combined the 15 cortical networks from Fig. 2A-O according to their normalized Z scores from ICA to create a cortical-network parcellation. The details include 1) the combination of networks according to their spatial locations; 2) if they have spatial overlapping, we took the highest value according to their normalized Z scores from ICA; 3) short-range (local) connectivity is usually stronger than long-range connectivity, so the single map cannot cover long-range connectivity due to the spatial overlapping. Therefore, we created a second map to cover the components with long-range connectivity that are missed in the first map. We repeat the above step but only applied to networks with long-range connections (such as Fig. 2I-K) to obtain the second map. The primary map (Fig. 2P-Q, top rows) is mostly contributed by the short-range networks (i.e., Fig. 1G, L, I, and

O), and the second one (Fig. 2P-Q, bottom rows) is to cover the long-range connectivity

that was not captured by the primary map.

Since the human connectome project released a pipeline for denoising by ICA-FIX, it has

recently been applied to some animal research studies. Therefore, we also explored

whether adopting this processing would affect the identification of brain networks. Before

using ICA-FIX, we had to create a training dataset as a standard. For this, we ran first-

level ICA on each fMRI run, randomly selected 24 runs from 24 animals (12 runs from the

ION dataset and 12 runs from the NIH dataset), and manually classified the noise

components based on their spatial patterns and power spectrums. Since the

recommendations for the use of ICA-FIX suggest a training dataset of at least 10, we

created a total of 24 datasets (trainingMBMv4.RData) to improve significantly our ICA-

FIX classifier, which included both the ION dataset (trainingION.RData, using 12 ION

training datasets) and the NIH (trainingNIH.RData, using 12 NIH training datasets)

respectively. Then, a trained-weighted file (trainingMBMv4.RData) was used to clean all

fMRI data based on three sensible-value thresholds (5, 10, and 20) since ICA-FIX

recommends a threshold in the range of 5-20. Thus, we created three different versions

of ICA-FIX cleaned datasets. In addition, we provide the mask files to allow ICA-FIX to

work on the marmoset data since ICA-FIX had several steps that use human-default

settings and files incompatible with the marmoset data. Regardless of which training

dataset was used, we still obtained the same functional network results shown in Fig. 2A-

O.

Boundary map generation

Following similar procedures to those described previously in a human imaging study [22],

the boundary mapping of resting-state functional connectivity data was implemented in

the Connectome Workbench and using customized Matlab codes (Mathworks, Natick,

USA, Version 2019b; see the scripts in our open resource). First, the time course of every

surface vertex for each brain hemisphere of each subject was correlated with every other

surface vertex to make a correlation map. Then, a similarity map was created for every

vertex by calculating pairwise spatial correlations between all correlation maps. Thirdly,

the first spatial derivative was applied on the similarity map by the Connectome

Workbench's function "cifti-gradient" to generate gradient maps for each brain

hemisphere of each subject. Next, the gradient maps were averaged across subjects to

produce the group gradient maps for each brain hemisphere. Lastly, the "watershed by

flooding" algorithm was applied to identify boundaries in the gradient maps.

Test-retest evaluation of the boundary map

To compare the reliability of the boundary maps between the ION and the NIH datasets

(Fig. 3A and Supplementary Fig. 8), between the individuals (Fig. 3C-E), and between

runs from the same individual (Fig. 3E), we first thresholded two resulting boundary maps

for each hemisphere to retain the cortical vertices most likely to be boundaries (i.e.,

retaining the top quartile of boundary values for a cumulative probability of 0.75) and

assessed the overlap of the two thresholded boundaries by calculating the Dice's

coefficient. The Dice's similarity coefficient of two thresholded boundaries, A and B, is

expressed as:

727
$$\text{dice}(A, B) = 2 * \left| \frac{\text{intersection}(A,B)}{|A|+|B|} \right| \tag{1}$$

The average Dice's similarity coefficient is the mean of Dice's similarity coefficients across

hemispheres.

Cortical parcellation based on the population-level boundary map

The creation of parcels was implemented by the customized Matlab scripts (see our open

resource). Firstly, based on the vertices with values smaller than their neighbors that were

<5 vertices away, we identified all local minima of vertices on the boundary map as seeds

for parcel creation. Then, the parcels were grown from these seeds using the "watershed

algorithm" procedure as above, allowing them to expand outward from the seed until they

met other parcels. Because the whole process depends on the number of seeds for parcel

creation, this might result in a large number of parcels. Therefore, according to the

performance, we manually defined a threshold for merging adjacent parcels, which is the

60th percentile of the values in the boundary map [22]. It means that any two adjacent

parcels with an average value below this threshold were considered not sufficiently

dissimilar and should be merged. Finally, according to the population-level boundary map,

we visually examined the remaining parcels to identify those needing further adjustment,

including eliminating vertices and spatial smoothing. The detailed manual processings for

the post-optimization included 1) manually adjusting the parcel borders, 2) manually

correcting wrong areal attributions of the region growing, and 3) spatial smoothing the

parcel borders by 8-neighbor vertices. We finally found the resulting cortical parcellation

with 96 functional parcels in each hemisphere as our Marmoset Brain Mapping Atlas

Version-4 (MBMv4) in Fig. 4B.

Evaluation of cortical parcellation by the distance-controlled boundary coefficient

(DCBC)

Following a previous study [26], we used the distance-controlled boundary coefficient

(DCBC) as a metric to evaluate functional boundaries between our parcels. The rationale

for this method is that any two points belonging to any given parcel should have more

similar functional profiles than those belonging to different parcels. Furthermore, because

the functional organization varies smoothly, the correlation between two points will

weaken with increasing spatial distance. Thus, we calculated the correlation coefficients

for all pairs of points separated by a specific surface Euclidean distance, using 0.5 mm

spatial bins (same as fMRI imaging spatial resolution) ranging from 0 to 4 mm for pairs of

points residing within parcels or across different parcels (between). The DCBC defines

the difference between the within-parcel and between-parcel pair correlations. A higher

DCBC reflects that pairs within the same region are more functional, serving as a global

parcellation measure. For the group comparison across atlases (Fig. 4C), the DCBC

metrics were calculated for each participant in each spatial bin and then averaged. For

the same participant comparison across atlases (Fig. 5D), the DCBC metrics were

calculated for each session in each spatial bin and then averaged.

Comparison with alternative atlases

We compared our parcellation against alternative digital parcellations created by various approaches. These alternative parcellations included: (1) Paxinos atlas [68], the most commonly used atlas in marmoset brain research, which is cytoarchitectonic characterization by immunohistochemical sections, and here we used its 3D digital version [15, 27]; (2) RIKEN atlas [28]: The atlas is cytoarchitecture based on Nissl-staining contrast. (3) The first atlas version of Marmoset Brain Mapping (MBMv1) [15]: The borders were delineated based on the high-resolution diffusion MRI contrast and parcellated by a structural-connectivity-based approach.

Deep-learning-based individual parcellation generator

The group-average parcellation described in the preceding sections is desirable for generating parcellations of individual animals. Although applying our group-level parcellation to individual animals is feasible, as demonstrated in a previous human study [22], we still found misalignments between individuals and cannot be highly consistent with the tendency of the group-average parcellation (MBMv4) when the scanning runs are limited (Fig. 3D-E). Therefore, inspired by previous works [29, 30], we trained a multi-layer deep learning network to classify parcels based on the fingerprints from MBMv4. There were two assumptions for this approach: (1) We assumed that individual cortical parcels were close to the group definition after the feature-based surface registration; (2) We assumed that every identified cortical parcel should be in a single class which was the combination of the target parcel and its spatially adjacent parcels (the "searchlight" for the candidate parcel). Thus, the setup of the classifier network was straightforward. Its architecture was as follows (for the graphic reference, see Fig. 5A): for each of the 96 parcels in each hemisphere, a multi-layer deep neural network was designed, which comprised three layers (one input, five hidden, and one output) and 384 hidden neurons (a reasonable compromise between accuracy and training speed for the classification). The whole-brain fingerprint of the candidate parcel from the MBMv4 worked as the training set for the network to classify whether or not each vertex in an individual ROI contained the parcel plus all of its neighbor parcels. Because of the spatial overlap of the

"searchlight," we excluded the vertices belonging to multiple parcels. Then, we applied

the same procedure of parcel creation as above, meaning that the borders of each

identified parcel became the seeds to expand outward until they met other parcels using

the "watershed by flooding" procedure. The whole process of individual parcellation was

automatic and implemented using customized MATLAB codes (example codes are

shared via www.marmosetbrainmapping.org/data.html) combined with the MATLAB

Deep Learning Network toolbox.

Evaluation by task-activation pattern

We examined the functional relevance of the borders by evaluating the parcels contained

within the fMRI activation pattern to a visual task (Fig. 6A) from our previous study [32]. A

subset of animals from the NIH dataset participated in the visual-choice task, which

consisted of watching 20-s-long movies (visual field is 10deg x 8deg) and 16s resting

periods (206 trials for marmoset-ID15 and 280 trials for marmoset-ID25). We performed

a contrast comparison between the movie-presentation blocks and the resting blocks to

generate visual-task activation statistical maps for each session. A mixed-effects analysis

was then applied to all statistical maps across sessions by the 3dMEMA command of

AFNI to obtain a final statistical map. The map was thresholded at a voxel-wise threshold

of $p < 0.05$ and a cluster-wise threshold of $p < 0.05$ for multiple comparison corrections.

To compare the similarity of the activation map and the parcellations in each hemisphere

(for results, see the flat maps in Fig. 6A), we calculated the shortest Euclidean distance

of every vertex/voxel in the boundary of the activation map to the vertexes/voxels in the

boundary of parcels/regions from different parcellations. We considered the parcellation

with the overall shortest distances of every vertex/voxel in the boundary of the activation

map as the best border consistency (for results, see the scatterplots in Fig. 6A).

Evaluation by functional connectivity gradient spectrum

It is widely accepted that the cerebral cortex of multiple species, including both human

and macaque primates, is organized along principal functional gradients that provide a

spatial framework for the co-existence of multiple large-scale networks operating in a

spectrum from unimodal to transmodal functional activity [34, 69]. Therefore, if the MBMv4

parcellation created here accurately represents the functional organization of the marmoset cortex, we can presume that it will also reveal these principal functional gradients. Thus, as in previous studies [34, 70], we followed a workflow for gradient identification: we first computed the rs-fMRI functional connectivity (RSFC) based on MBMv4. Next, the RSFC matrix $M_{x,y}$ with the same size as the atlas was made sparse (to a 10% sparsity), and a similarity matrix $A_{x,y}$ with the normalized angle was computed according to the following equation:

$$A(x,y) = 1 - \frac{\cos^{-1}(\text{cossim}(x,y))}{\pi} \tag{2}$$

Next, the similarity matrix was decomposed via Laplacian transformation into a set of principal eigenvectors describing the axes of most significant variance using the following equation:

$$Lg = \lambda Dg \tag{3}$$

Where $D_{x,y} = \sum_y A(x,y)$, L is the graph Laplacian matrix, and the eigenvectors g which corresponds to the m smallest eigenvalues λ_k are used to build the new low-dimensional representation:

$$\varsigma_{LE} = [g_1, g_2, ..., g_m] \tag{4}$$

Finally, the first two axes g_1, g_2 of each parcel were plotted in 2D space. Meanwhile, we used the scores g_1 to sort the functional connectivity matrix (for results, see the heatmaps in Fig. 6B).

The whole-brain modeling for the link between structural connectivity and functional connectivity

As we know, structural connectivity and functional connectivity are closely related to each other. Therefore, the lack of structural evidence generally implies biological implausibility

for functional connections. Testing whether the cortical parcels created above MBMv4

are accurate representations of the functional areas in the cerebral cortex requires

investigation of the underlying structural connectivity. A computational model is a powerful

approach to bridge structural and functional connectivity [69, 71, 72, 73, 74]. In the present study,

we implemented a whole-brain model with only two free parameters from previous studies

[35, 36], as outlined below (for a graphic reference, see Fig. 7A, note that the fMRI data for

the modeling part is frequently unfiltered, so the model used the full band of frequency):

According to the whole cortical parcellations (192 total parcels, 96 per hemisphere from

MBMv4 or 232 total regions, 116 regions per hemisphere from the Paxinos atlas), the

structural connectivity between parcels/regions $C_{i,j}$ was estimated from the structural

datasets (see examples in Fig. 7B-C and 7E-F), either DTI data (in-vivo or ex-vivo) or the

neuronal tracing data. Then, the local dynamics for every parcel/region j can be properly

approximated to the normal form of a Hopf bifurcation:

$$877 \qquad \frac{dz_j}{dt} = [a_j + iw_j]z_j + z_j|z_j|^2 \qquad\qquad (5)$$

In this equation, z_j is a complex-valued variable $z_j = x_j + y_j$, and w_j is the intrinsic signal

frequency of parcel/region j, which ranged from 0.04-0.07Hz and was determined by the

averaged peak frequency of the bandpass-filtered fMRI signals of the parcel/region j [36,

75, 76, 77, 78]. a_j is a bifurcation free parameter controlling the dynamics of the parcel/region

j . For $a_j < 0$, the phase space presents a unique stable and is governed by noise.

For $a_j > 0$, the phase space presents the stable state, giving rise to a self-sustained

oscillation. For $a_j \approx 0$ the phase presents an unstable state, switching back and forth and

giving rise to a mixture of oscillation and noise.

The coordinated dynamics of the resting state activity for parcel/region j could be modeled

by coupling determined by the above structural connectivity $C_{i,j}$. To ensure the oscillatory

dynamics for $a_j > 0$, the structural connectivity $C_{i,j}$ should be normalized and scaled to 0.2

in a weak coupling condition before starting the simulation. The coupled differential

equations of the model are the following:

891
$$\frac{dx_j}{dt} = \left[a_j - x_j^2 - y_j^2\right]x_j - w_j y_j + G \sum_i C_{i,j}(x_i - x_j)\beta\eta_j(t) \qquad (6)$$

892
$$\frac{dy_j}{dt} = \left[a_j - x_j^2 - y_j^2\right]y_j + w_j x_j + G \sum_i C_{i,j}(y_i - y_j)\beta\eta_j(t) \qquad (7)$$

In this equation, G is another free parameter representing the fixed global coupling factor

that scales structural connectivity $C_{i,j}$. η_j represents additive Gaussian noise in each

parcel/region and is scaled by a factor β fixed at 0.04 according to previous studies [36].

Euler-Maruyama algorithm integrated these equations with a time step of 0.1 seconds to

accelerate simulation [79].

The free bifurcation parameter a_j for parcel/region j could be locally optimized based on

fitting the spectral information of the empirical BOLD signals. To achieve this, we filtered

raw empirical BOLD data in the 0.04–0.25Hz band and calculated the power spectrum

$p_j(f)$ for each parcel j as below:

902
$$p_j = \frac{\int_{0.04}^{0.07} p_j(f)df}{\int_{0.04}^{0.25} p_j(f)df} \qquad (8)$$

903 and updated the local bifurcation parameter a_j by a gradient descendent strategy:

904
$$a_j = a_j + \eta\left(p_j^{empirical} - p_j^{simulated}\right) \qquad (9)$$

We applied the above optimization process to receive the best bifurcation parameters a_j

of every parcel/region defined in the parcellations. Once we found the optimized set of

bifurcation parameters a_j, we adjusted the free parameter G within the range of 0-8 in

steps of 0.1 according to a reasonable compromise from previous studies [36, 79] to simulate

the same number of sessions for each animal and the same number of animals. To

compare the performance in different atlases, we just needed to compare fitting (similarity)

metrics, Pearson's correlation coefficient between the simulated functional connectivity

and the one used for the empirical data, when we fixed the same value of parameter G.

Since the distributions of the optimal bifurcation parameter a are identical in different

parcellations MBMv4 or Paxinos atlas (see Supplementary Fig. 13, one way ANOVA

$F_{(1,11986)}$=9.09, p=0.26), we selected the best free parameter G from the Paxinos atlas for

comparison performance with our MBMv4 (see results in Fig. 7B and 7E, examples in Fig.

7C and 7F). Moreover, we also selected the group-averaged functional connectivity from

all individuals as the empirical observable for the ultra-high resolution diffusion MRI and

neuronal tracing datasets and the individual functional connectivity for the corresponding

in-vivo diffusion MRI.

The resulting simulation of functional connectivity demonstrated the influence of the

distance, especially for the diffusion MRI (see Fig. 7D). To decrease this impact, we

added an Exponential Distance Rule (EDR) for a more accurate estimation of structural

connectivity before the simulation, which can be implemented as follows: we first

normalized the structural connectivity estimated by diffusion tractography to 0-1, then

calculated the probability of structural connectivity according to the EDR rule ($p(d)=ce^{-\lambda d}$,

$\lambda \approx 0.3$, $c \approx 0.94$ where d is the distance [40]) and normalized to 0-1. Finally, we transformed

the normalized structural connectivity to match the probability of structural connectivity.

For simplicity to help us to identify abnormal values, we defined a threshold as the Median

Absolute Deviation for the normalized structural connectivity.

$$\hat{\sigma} = \frac{\text{Median}(|X_i - \widetilde{X}|)}{0.6745} \tag{10}$$

Where X_i = each value, $\widetilde{X}$ = average value

If any values were larger than this threshold, we considered them abnormal values, and

the corresponding probability of structural connectivity would replace them to reduce the

impact of distance.

Data availability

All of our datasets, including raw and preprocessed NIH and ION resting-state fMRI,
diffusion MRI, and neuronal tracing datasets, are available on our specific webpage of
the Marmoset Brain Mapping Project (www.marmosetbrainmapping.org/data.html). The
volume data are in NIFTI format, and the surface data are in CIFTI format. The raw MRI
data without processing is provided in the standard BIDS format for cross-platform
sharing. The MBMv4 parcellations are also provided on the same webpage
(www.marmosetbrainmapping.org/data.html), and the MBMv1 and Paxinos parcellations
on our MRI template space are a part of the MBMv3 resource
(marmosetbrainmapping.org/v3.html). The high-resolution ex-vivo diffusion MRI data are
a part of the MBMv2 resource (marmosetbrainmapping.org/atlas.html#v2). The raw
neuronal tracing data are from Marmoset Brain Connectivity project
(https://www.marmosetbrain.org/reference). Note that the MBMv4 datasets are only
available for scientific purposes and are licensed under Creative Commons Attribution-
NonCommercial-ShareAlike (CC BY-NC-SA 4.0).

Code availability

The codes for analyzing pipelines used in this study could be available on our webpage
(www.marmosetbrainmapping.org/data.html). The codes package with examples
includes the generation of cortical functional boundary maps, functional parcellation, and
their evaluation metrics, as well as computational modeling which could link with structural
imaging and neuronal tracing. Note that the codes and analyzing pipelines are only
available for scientific purposes and are licensed under Creative Commons Attribution-
NonCommercial-ShareAlike (CC BY-NC-SA 4.0).

References

1. Buffalo EA, Movshon JA, Wurtz RH. From basic brain research to treating human brain disorders. Proc Natl Acad Sci U S A, (2019).

2. Thiebaut de Schotten M, Croxson PL, Mars RB. Large-scale comparative neuroimaging: Where are we and what do we need? Cortex **118**, 188-202 (2019).

3. Buckner RL, Margulies DS. Macroscale cortical organization and a default-like apex transmodal network in the marmoset monkey. Nat Commun **10**, 1976 (2019).

4. Smith SM, et al. Resting-state fMRI in the Human Connectome Project. Neuroimage **80**, 144-168 (2013).

5. Milham MP, et al. An Open Resource for Non-human Primate Imaging. Neuron **100**, 61-74 e62 (2018).

6. Hudetz AG. General anesthesia and human brain connectivity. Brain Connect **2**, 291-302 (2012).

7. Milham M, et al. Accelerating the Evolution of Nonhuman Primate Neuroimaging. Neuron **105**, 600-603 (2020).

8. Milham M, et al. Toward next-generation primate neuroscience: A collaboration-based strategic plan for integrative neuroimaging. Neuron, (2021).

9. Messinger A, et al. A collaborative resource platform for non-human primate neuroimaging. Neuroimage **226**, 117519 (2021).

10. Okano H. Current Status of and Perspectives on the Application of Marmosets in Neurobiology. Annu Rev Neurosci **44**, 27-48 (2021).

11. Walker J, MacLean J, Hatsopoulos NG. The marmoset as a model system for studying voluntary motor control. Dev Neurobiol **77**, 273-285 (2017).

12. Matsuzaki M, Ebina T. Common marmoset as a model primate for study of the motor control system. Curr Opin Neurobiol **64**, 103-110 (2020).

13. Liu C, et al. A resource for the detailed 3D mapping of white matter pathways in the marmoset brain. Nat Neurosci **23**, 271-280 (2020).

14. Majka P, et al. Open access resource for cellular-resolution analyses of corticocortical connectivity in the marmoset monkey. Nat Commun **11**, 1133 (2020).

1012
1013 15. Liu C, et al. A digital 3D atlas of the marmoset brain based on multi-modal MRI.
1014   Neuroimage **169**, 106-116 (2018).
1015
1016 16. Liu C, Yen CC, Szczupak D, Tian X, Glen D, Silva AC. Marmoset Brain Mapping V3:
1017   Population multi-modal standard volumetric and surface-based templates. Neuroimage
1018   **226**, 117620 (2021).
1019
1020 17. Majka P, et al. Histology-Based Average Template of the Marmoset Cortex With
1021   Probabilistic Localization of Cytoarchitectural Areas. Neuroimage **226**, 117625 (2021).
1022
1023 18. Autio JA, et al. Minimal specifications for non-human primate MRI: Challenges in
1024   standardizing and harmonizing data collection. Neuroimage **236**, 118082 (2021).
1025
1026 19. Belcher AM, et al. Large-scale brain networks in the awake, truly resting marmoset
1027   monkey. J Neurosci **33**, 16796-16804 (2013).
1028
1029 20. Ghahremani M, Hutchison RM, Menon RS, Everling S. Frontoparietal Functional
1030   Connectivity in the Common Marmoset. Cereb Cortex **27**, 3890-3905 (2017).
1031
1032 21. Liu C, Yen CC, Szczupak D, Ye FQ, Leopold DA, Silva AC. Anatomical and functional
1033   investigation of the marmoset default mode network. Nat Commun **10**, 1975 (2019).
1034
1035 22. Gordon EM, Laumann TO, Adeyemo B, Huckins JF, Kelley WM, Petersen SE. Generation
1036   and Evaluation of a Cortical Area Parcellation from Resting-State Correlations. Cereb
1037   Cortex **26**, 288-303 (2016).
1038
1039 23. Cohen AL, et al. Defining functional areas in individual human brains using resting
1040   functional connectivity MRI. Neuroimage **41**, 45-57 (2008).
1041
1042 24. Wig GS, Laumann TO, Petersen SE. An approach for parcellating human cortical areas
1043   using resting-state correlations. Neuroimage **93 Pt 2**, 276-291 (2014).
1044
1045 25. Cousty J, Bertrand G, Najman L, Couprie M. Watershed Cuts: Minimum Spanning Forests
1046   and the Drop of Water Principle. Ieee T Pattern Anal **31**, 1362-1374 (2009).
1047
1048 26. King M, Hernandez-Castillo CR, Poldrack RA, Ivry RB, Diedrichsen J. Functional
1049   boundaries in the human cerebellum revealed by a multi-domain task battery. Nat
1050   Neurosci **22**, 1371-1378 (2019).
1051
1052 27. Majka P, et al. Towards a comprehensive atlas of cortical connections in a primate brain:
1053   Mapping tracer injection studies of the common marmoset into a reference digital
1054   template. J Comp Neurol **524**, 2161-2181 (2016).
1055

1056   28.   Woodward A, et al. The Brain/MINDS 3D digital marmoset brain atlas. Sci Data **5**,
1057   180009 (2018).
1058

1059   29.   Glasser MF, et al. A multi-modal parcellation of human cerebral cortex. Nature **536**, 171-
1060   178 (2016).
1061

1062   30.   Hacker CD, et al. Resting state network estimation in individual subjects. Neuroimage
1063   **82**, 616-633 (2013).
1064

1065   31.   Mueller S, et al. Individual variability in functional connectivity architecture of the
1066   human brain. Neuron **77**, 586-595 (2013).
1067

1068   32.   Tian X, Silva AC, Liu C. The Brain Circuits and Dynamics of Curiosity-Driven Behavior in
1069   Naturally Curious Marmosets. Cereb Cortex **31**, 4220-4232 (2021).
1070

1071   33.   Russ BE, et al. Common functional localizers to enhance NHP & cross-species
1072   neuroscience imaging research. Neuroimage **237**, 118203 (2021).
1073

1074   34.   Margulies DS, et al. Situating the default-mode network along a principal gradient of
1075   macroscale cortical organization. Proc Natl Acad Sci U S A **113**, 12574-12579 (2016).
1076

1077   35.   Ipina IP, et al. Modeling regional changes in dynamic stability during sleep and
1078   wakefulness. Neuroimage **215**, 116833 (2020).
1079

1080   36.   Deco G, Kringelbach ML, Jirsa VK, Ritter P. The dynamics of resting fluctuations in the
1081   brain: metastability and its dynamical cortical core. Sci Rep **7**, 3095 (2017).
1082

1083   37.   Deco G, Jirsa V, McIntosh AR, Sporns O, Kotter R. Key role of coupling, delay, and noise
1084   in resting brain fluctuations. Proc Natl Acad Sci U S A **106**, 10302-10307 (2009).
1085

1086   38.   Horvat S, et al. Spatial Embedding and Wiring Cost Constrain the Functional Layout of
1087   the Cortical Network of Rodents and Primates. PLoS Biol **14**, e1002512 (2016).
1088

1089   39.   Ercsey-Ravasz M, et al. A predictive network model of cerebral cortical connectivity
1090   based on a distance rule. Neuron **80**, 184-197 (2013).
1091

1092   40.   Theodoni P, Majka P, Reser DH, Wojcik DK, Rosa MGP, Wang XJ. Structural Attributes
1093   and Principles of the Neocortical Connectome in the Marmoset Monkey. Cereb Cortex
1094   **32**, 15-28 (2021).
1095

1096   41.   Rilling JK, Insel TR. Differential expansion of neural projection systems in primate brain
1097   evolution. Neuroreport **10**, 1453-1459 (1999).
1098

1099 42. Changizi MA, Shimojo S. Parcellation and area-area connectivity as a function of
1100   neocortex size. Brain Behav Evol **66**, 88-98 (2005).
1101

1102 43. Kaas JH. The organization of neocortex in mammals: implications for theories of brain
1103   function. Annu Rev Psychol **38**, 129-151 (1987).
1104

1105 44. Amunts K, Zilles K. Architectonic Mapping of the Human Brain beyond Brodmann.
1106   Neuron **88**, 1086-1107 (2015).
1107

1108 45. Schaefer A, et al. Local-Global Parcellation of the Human Cerebral Cortex from Intrinsic
1109   Functional Connectivity MRI. Cereb Cortex **28**, 3095-3114 (2018).
1110

1111 46. Van Essen DC, Glasser MF, Dierker DL, Harwell J, Coalson T. Parcellations and
1112   hemispheric asymmetries of human cerebral cortex analyzed on surface-based atlases.
1113   Cereb Cortex **22**, 2241-2262 (2012).
1114

1115 47. Van Essen DC, Glasser MF. Parcellating Cerebral Cortex: How Invasive Animal Studies
1116   Inform Noninvasive Mapmaking in Humans. Neuron **99**, 640-663 (2018).
1117

1118 48. van den Heuvel MP, Yeo BTT. A Spotlight on Bridging Microscale and Macroscale Human
1119   Brain Architecture. Neuron **93**, 1248-1251 (2017).
1120

1121 49. Churchland PS, Sejnowski TJ. Perspectives on cognitive neuroscience. Science **242**, 741-
1122   745 (1988).
1123

1124 50. Smith SM, et al. Correspondence of the brain's functional architecture during activation
1125   and rest. Proc Natl Acad Sci U S A **106**, 13040-13045 (2009).
1126

1127 51. Mennes M, et al. Inter-individual differences in resting-state functional connectivity
1128   predict task-induced BOLD activity. Neuroimage **50**, 1690-1701 (2010).
1129

1130 52. Tavor I, Parker Jones O, Mars RB, Smith SM, Behrens TE, Jbabdi S. Task-free MRI predicts
1131   individual differences in brain activity during task performance. Science **352**, 216-220
1132   (2016).
1133

1134 53. Rosa MG, Tweedale R. Brain maps, great and small: lessons from comparative studies of
1135   primate visual cortical organization. Philos Trans R Soc Lond B Biol Sci **360**, 665-691
1136   (2005).
1137

1138 54. Yu HH, Rowley DP, Price NSC, Rosa MGP, Zavitz E. A twisted visual field map in the
1139   primate dorsomedial cortex predicted by topographic continuity. Sci Adv **6**, (2020).
1140

1141 55. Long X, Goltz D, Margulies DS, Nierhaus T, Villringer A. Functional connectivity-based
1142   parcellation of the human sensorimotor cortex. Eur J Neurosci **39**, 1332-1342 (2014).

1143
1144 56. Rao SM, et al. Somatotopic mapping of the human primary motor cortex with functional
1145   magnetic resonance imaging. Neurology **45**, 919-924 (1995).
1146
1147 57. Chaplin TA, Yu HH, Soares JG, Gattass R, Rosa MG. A conserved pattern of differential
1148   expansion of cortical areas in simian primates. J Neurosci **33**, 15120-15125 (2013).
1149
1150 58. Sawiak SJ, et al. Trajectories and Milestones of Cortical and Subcortical Development of
1151   the Marmoset Brain From Infancy to Adulthood. Cereb Cortex **28**, 4440-4453 (2018).
1152
1153 59. Papoti D, Yen CC, Hung CC, Ciuchta J, Leopold DA, Silva AC. Design and implementation
1154   of embedded 8-channel receive-only arrays for whole-brain MRI and fMRI of conscious
1155   awake marmosets. Magn Reson Med **78**, 387-398 (2017).
1156
1157 60. Silva AC, et al. Longitudinal functional magnetic resonance imaging in animal models.
1158   Methods Mol Biol **711**, 281-302 (2011).
1159
1160 61. Caruyer E, Lenglet C, Sapiro G, Deriche R. Design of multishell sampling schemes with
1161   uniform coverage in diffusion MRI. Magn Reson Med **69**, 1534-1540 (2013).
1162
1163 62. Cox RW. AFNI: what a long strange trip it's been. Neuroimage **62**, 743-747 (2012).
1164
1165 63. Jenkinson M, Beckmann CF, Behrens TE, Woolrich MW, Smith SM. Fsl. Neuroimage **62**,
1166   782-790 (2012).
1167
1168 64. Tustison N, et al. Multivariate Analysis of Diffusion Tensor Imaging and Cortical
1169   Thickness Maps in a Traumatic Brain Injury (Tbi) Cohort Using Advanced Normalization
1170   Tools (Ants). J Neurotraum **28**, A111-A111 (2011).
1171
1172 65. Marcus DS, et al. Informatics and data mining tools and strategies for the human
1173   connectome project. Front Neuroinform **5**, 4 (2011).
1174
1175 66. Pierpaoli C. Quantitative brain MRI. Top Magn Reson Imaging **21**, 63 (2010).
1176
1177 67. Tournier J-D, Calamante F, Connelly A. MRtrix: Diffusion tractography in crossing fiber
1178   regions. International Journal of Imaging Systems and Technology **22**, 53-66 (2012).
1179
1180 68. Paxinos G, Watson C, Petrides M, Rosa M, Tokuno H. The marmoset brain in stereotaxic
1181   coordinates. Elsevier Academic Press (2012).
1182
1183 69. Wang P, et al. Inversion of a large-scale circuit model reveals a cortical hierarchy in the
1184   dynamic resting human brain. Sci Adv **5**, eaat7854 (2019).
1185

1186   70.   Vos de Wael R, et al. BrainSpace: a toolbox for the analysis of macroscale gradients in
1187   neuroimaging and connectomics datasets. Commun Biol **3**, 103 (2020).
1188
1189   71.   Hansen EC, Battaglia D, Spiegler A, Deco G, Jirsa VK. Functional connectivity dynamics:
1190   modeling the switching behavior of the resting state. Neuroimage **105**, 525-535 (2015).
1191
1192   72.   Breakspear M. Dynamic models of large-scale brain activity. Nat Neurosci **20**, 340-352
1193   (2017).
1194
1195   73.   Schirner M, McIntosh AR, Jirsa V, Deco G, Ritter P. Inferring multi-scale neural
1196   mechanisms with brain network modelling. Elife **7**, (2018).
1197
1198   74.   Honey CJ, et al. Predicting human resting-state functional connectivity from structural
1199   connectivity. Proc Natl Acad Sci U S A **106**, 2035-2040 (2009).
1200
1201   75.   Ponce-Alvarez A, Deco G, Hagmann P, Romani GL, Mantini D, Corbetta M. Resting-state
1202   temporal synchronization networks emerge from connectivity topology and
1203   heterogeneity. PLoS Comput Biol **11**, e1004100 (2015).
1204
1205   76.   Achard S, Salvador R, Whitcher B, Suckling J, Bullmore E. A resilient, low-frequency,
1206   small-world human brain functional network with highly connected association cortical
1207   hubs. J Neurosci **26**, 63-72 (2006).
1208
1209   77.   Buckner RL, et al. Cortical hubs revealed by intrinsic functional connectivity: mapping,
1210   assessment of stability, and relation to Alzheimer's disease. J Neurosci **29**, 1860-1873
1211   (2009).
1212
1213   78.   Glerean E, Salmi J, Lahnakoski JM, Jaaskelainen IP, Sams M. Functional magnetic
1214   resonance imaging phase synchronization as a measure of dynamic functional
1215   connectivity. Brain Connect **2**, 91-101 (2012).
1216
1217   79.   Sanz Perl Y, et al. Perturbations in dynamical models of whole-brain activity dissociate
1218   between the level and stability of consciousness. PLoS Comput Biol **17**, e1009139
1219   (2021).

1230

Acknowledgments

1232

1233 We thank Kaiwei Zhang and Binshi Bo at the 9.4T core facility (CEBSIT) for assistance
1234 in the data collection of the ION data, Lisa Zhang for the assistance in the data collection
1235 of the NIH 7T data, the Marmoset Animal Facility of CEBSIT for animal care, Xiaojia Zhu
1236 for the assistance in organizing MRI data in BIDS format, and the NIH Fellows Editorial
1237 Board for the editorial assistance on the early version of the manuscript.

1238

The study was supported by the grants from National Science and Technology Innovation
2030 Major Project of China (2022ZD0205000, 2021ZD0203900), the Pennsylvania
Department of Health Commonwealth Universal Research Enhancement (CURE)
Tobacco Settlement Appropriation – Phase 18 (Grant SAP4100083102 to ACS), the US
National Institute on Aging (Grants R24AG073190 and U19AG074866 to ACS), the
Shanghai Municipal Science and Technology Major Project (No. 2018SHZDZX05 to CL
and ZF), the Lingang Laboratory (Grant No. LG-QS-202201-02 to CL), the National
Natural Science Foundation of China (No. 32171088 to CL) , the Australian Research
Council (DP110101200, DP140101968, CE140100007) and National Health and Medical
Research Council (APP1194206) to MR, National Science Centre (2019/35/D/NZ4/03031
to PM), NIH Intramural Research Programs (ZIA NS003041 to AS and CY, ZICMH002888
to DG), International Neuroinformatics Coordinating Facility Seed Funding Grant (to PM
and MR), a Spanish research project funded by the Spanish Ministry of Science,
Innovation, and Universities (MCIU), State Research Agency (AEI), and European
Regional Development Funds (FEDER) (ref. PID2019-105772GB-I00 AEI FEDER EU to
GD), HBP SGA3 Human Brain Project Specific Grant Agreement 3 (grant agreement no.
945539 to GD), and the EU H2020 FET Flagship program and SGR Research Support
Group support (ref. 2017 SGR 1545 to GD).

1257

1258

1259

1260

1261

Author contributions

1263

C.L., X.T., A.C.S., and M.R. designed and supervised the study; Z.L. and Y.C. collected
the ION MRI data; D.S., C.L., C.Y., and X.T. collected the NIH MRI data; M.R. and P.M.
collected the neuronal tracing data; C.L., X.T., C.T., and Z.L. preprocessed and organized
the MRI data; P.M., C.L., X.T., and H.J. preprocessed the neuronal tracing data; C.L. and
X.T. constructed the functional-network maps; C.L., X.T., and M.R. constructed and
evaluated the cortical parcellation maps; X.T., C.L., Y.P., and G.D. conducted the
computational modeling; C.L. and F.F. developed the online web applications; D.G. and
C.L. implemented the atlas into AFNI/SUMA. X.T. and C.L. wrote the original draft. C.L.,
X.T., A.C.S., and M.R. revised the manuscript.

1273

Competing Interests
1275

1276 The Authors declare no competing interests
1277

Figure Legends

Figure 1. Outline of Marmoset Brain Mapping Resource. The resource provides the awake test-retest resting-state fMRI data, in-vivo diffusion MRI data from the same marmoset cohorts, and the neuronal tracing data mapped onto the same MRI space at

the voxel/vertex level. In addition to the datasets, it also supports the study of whole-brain

functional networks and computational modelings, as well as functional connectivity-

based parcellation of the cortex (Marmoset Brain Mapping Atlas Version 4) using a deep

neural network for accurate individual mapping. As a comprehensive multimodal resource

for marmoset brain research, we also provide an online platform to explore the

relationship between structural and functional connectivity. This functionality is embodied

in online interactive viewers.

1290

1291

Figure 2. Functional cortical networks and their parcellation maps. The identified networks

include: (A) the ventral somatomotor, (B) the dorsal somatomotor, (C) the premotor, (D)

the frontal pole, (E) the orbital frontal cortex, (F) the parahippocampus and temporal pole,

(G-H) the auditory and salience-related network, (I-J) two trans-modal networks, including

a putative frontoparietal network and the default-mode-network, and (K-O) visual-related

networks from the primary visual cortex to higher-order functional regions. These

networks were combined to form two network-parcellation maps (P-Q), which are

dominated by the networks with short-range connectivity (P-Q, top rows) and with long-

range connectivity (P-Q, bottom rows), respectively.

1301

1302

Figure 3. The functional connectivity boundary maps. (A). The population-based
boundary maps from the ION, the NIH, and the combined datasets. These maps are
highly consistent, with an average Dice coefficient of 0.7. (B). Boundary maps in the left
hemisphere from four exemplar marmosets (two from the NIH cohort and two from the
ION, including the flagship marmosets). (C-D). The heatmap of the average Dice's
coefficients for both hemispheres between individuals and its distribution histogram. (E).
The average Dice's coefficients change for both hemispheres with the number of runs in
the same individuals.

1311

Figure 4. Marmoset Brain Mapping Atlas Version 4 (MBMv4). (A) The processing procedure includes generating the population functional connectivity boundary maps, defining the local minima for seeding, and generating parcels by the "watershed" algorithm. (B) The resulting 96 functional connectivity parcels per hemisphere are overlaid on the white matter surface and flat map of MBMv3 [16]. (C) Distance-controlled boundary coefficient (DCBC) evaluation metric. All pairs of voxels/vertices were categorized into "within" or "between" parcels (left panel) according to different brain parcellations (MBMv1, MBMv4, Paxinos, and RIKEN atlases, right panel), and the DCBC metric was calculated

1321   by the differences (within-between) in functional connectivity as the function of distance

1322   on the surface (0-4 mm in steps of 0.5 mm). Data are presented in mean +/- s.e.m.

1323

1324

Figure 5. Mapping individual functional connectivity parcellation. (A) An overview of

individual Mapping based on the deep neural network approach. (B) MBMv4 Mapping of

each individual. Left panel: the concordance between the population MBMv4 and

individual parcellations (N=78, all hemispheres from 39 subjects). Data are presented by

the violin and the box plots (the 25th percentile: 0.9068; the 75 percentile: 0.92448), in

which the white point represents the average value 0.915 (the maximum value: 0.931;

minimum 0.900); Right panel: three examples of individual parcellations. The underlay

(color-coded) presents the population MBMv4, and the overlay (black border) shows the

individual parcellations. (C) Mapping of MBMv4 per session. Left panel: The concordance

between every individual parcellation and the corresponding parcellation using one

session data (N=78, all hemispheres from 39 subjects). Data are presented by the violin

and the box plots (the 25th percentile 0.865; the 75 percentile 0.877), in which the white

point represents the average value 0.874 (the maximum: 0.878; the minimum 0.859);

Right panel: representative parcellations of three sessions from one marmoset. The color-

coded underlay represents individual parcellation, while the black border overlay shows

the session-based parcellation. (D) The distance-controlled boundary coefficient (DCBC)

for the individual parcellation generated by the spatial registration (Spatial-reg, blue) and

the deep neural network (DNN-reg, red). Top panel: the functional connectivity for all pairs

of vertices within the same parcel and between parcels for DNN-reg and Spatial-reg,

respectively. Bottom panel: the comparison of DNN-reg and Spatial-reg by DCBC. Data

are presented in mean +/- s.e.m.

1346

Figure 6. MBMv4 matches functional boundaries and preserves the topographical organization of the functional connectivity. (A) The visual activation maps from two monkeys are overlaid on the parcel boundaries from individual MBMv4 parcellation and Paxinos atlas (Left panel: monkey ID 15; Right panel: monkey ID 25). The scatter plots compare the boundary matching of the MBMv4 and the Paxinos atlas with the activation maps, measured by the shortest distance from every voxel in the borders of the activation maps to the parcel borders of the MBMv4 or the Paxinos atlas. The dashed black line represents the diagonal line, and the red line represents the linear fitting line. (B) The scatter plots in the left panel are the first two axes of gradients (the color scale of dots represents the scores of the first axis for every gradient), decomposed by the functional connectivities of the MBMv4 and the Paxinos atlas (the spectrum colors denote the gradient position in this 2D space). The heat maps of functional connectivity sorted by the scores of the first axis (gradient 1) are shown in the right panel.

1362

1363 Figure 7. A computational framework links the structural-functional connectivity according

1364 to different parcellations. (A) Application of whole-brain modeling, including an estimation

1365 of structural connectivity from neuronal tracing, diffusion MRI (in-vivo or ex-vivo)

1366 according to the Paxinos atlas or MBMv4, simulation of functional connectivity from

structural connectivity by the Hopf bifurcation hemodynamic functions, and a similarity measure with empirical connectivity from resting-state fMRI. (B) Model fitting comparison in different spatial scales using the Paxinos versus the MBMv4 atlases. Individual examples using in-vivo diffusion MRI (round dots; an example animal marked in black dot), ex-vivo diffusion MRI (polygon), and neuronal tracing (star) show higher fitting values for MBMv4 versus the Paxinos atlas. (C) Examples of correlations between the simulated and empirical functional connectivity from (B); solid black lines represent marginal regression lines. (D) Model fitting comparisons with (red) and without (blue) an exponential distance rule (EDR) correction. From left to right, the plots present the simulation results obtained with in-vivo diffusion MRI from the example animal (the linear regression values R = 0.689, p = 9.27e-50 for the blue line; R = 0.413, p = 1.28e-23 for the red); in-vivo diffusion MRI from all animals (R = 0.11, p = 6.05e-114 for the blue; R = 0.011, p = 4.17e-13 for the red); the ex-vivo diffusion MRI (R = 0.7, p = 2.69e-51 for the blue; R = 0.395, p = 2.19e-22 for the red); and the neuronal-tracing dataset (R = 0.009, p=0.02). The line plots are presented as mean values +/- 95% C.I. Dashed lines represent the 95% confidence interval. (E) Model fitting comparison with EDR correction between the Paxinos atlas and MBMv4 in different spatial scales. (F) Examples of correlation between the simulated and empirical functional connectivity from (E).