Maternal diet disrupts the placenta-brain axis in a sex-specific manner

Alexis M Ceasrine[1], Benjamin A Devlin[1], Jessica L. Bolton[2], Lauren A. Green[1], Young Chan Jo[1], Carolyn Huynh[1], Bailey Patrick[1], Kamryn Washington[1], Cristina L. Sanchez[4], Faith Joo[1], A. Brayan Campos-Salazar[3], Elana R. Lockshin[3], Cynthia Kuhn[4], Susan K. Murphy[5], Leigh Ann Simmons[6], Staci D. Bilbo*[1,3,7]

[1]Department of Psychology and Neuroscience, Duke University, Durham, NC, USA
[2]Neuroscience Institute, Georgia State University, Atlanta, GA, USA
[3]Department of Neurobiology, Duke University Medical Center, Durham, NC, USA
[4]Department of Pharmacology and Cancer Biology, Duke University Medical Center, Durham, NC, USA
[5]Department of Obstetrics and Gynecology, Duke University Medical Center, Durham, NC, USA
[6]Department of Human Ecology, Perinatal Origins of Disparities Center, University of California, Davis, California, USA
[7]Lurie Center for Autism, Massachusetts General Hospital, Boston, MA

SUMMARY: High maternal weight is associated with a number of detrimental outcomes in offspring, including increased susceptibility to neurological disorders such as anxiety, depression, and communicative disorders (e.g. autism spectrum disorders)[1–8]. Despite widespread acknowledgement of sex-biases in the prevalence, incidence, and age of onset of these disorders, few studies have investigated potential sex-biased mechanisms underlying disorder susceptibility. Here, we use a mouse model to demonstrate how maternal high-fat diet, one contributor to overweight, causes endotoxin accumulation in fetal tissue, and subsequent perinatal inflammation influences sex-specific behavioral outcomes in offspring. In male high-fat diet offspring, increased macrophage toll like receptor 4 signaling results in excess phagocytosis of serotonin neurons in the developing dorsal raphe nucleus, decreasing serotonin bioavailability in the fetal and adult brain. Bulk sequencing from a large cohort of matched first trimester human fetal brain, placenta, and maternal decidua samples reveals sex-specific transcriptome-wide changes in placenta and brain tissue in response to maternal triglyceride accumulation (a proxy for dietary fat content). Further, we find that fetal brain serotonin levels decrease as maternal dietary fat intake increases in males only. These findings uncover a microglia-dependent

mechanism through which maternal diet may impact offspring susceptibility for

neuropsychiatric disorder development in a sex-specific manner.

MAIN: In the United States, more than 50% of women are overweight or obese when they

become pregnant[9], and elevated maternal weight is associated with adverse outcomes

in offspring, particularly increased incidence of neuropsychiatric disorders such as

anxiety, and depression[1–8]. Diets high in saturated fats are one major contributor to

overweight and obesity[10]. Thus, we sought to dissect the contribution of a maternal high-

fat diet (mHFD) to offspring risk for neuropsychiatric disorder development.

Rodent and non-human primate models have implicated perinatal high-fat diet in

the establishment of offspring anxiety[11–13]. Decades of research into anxiety and

depression have strongly linked disruptions in the serotonin (5-HT) system as causal to

neuropsychiatric diseases such as anxiety and depression. Although women

disproportionately bear the burden of these disorders[14,15], little has been done to

investigate the contribution of 5-HT to anxiety/depression on the basis of sex.

Understanding the developmental mechanisms through which sex contributes to

neuropsychiatric disorder development, susceptibility, and severity in response to

environmental exposures such as diet is of utmost importance when considering clinical

implications such as treatment efficacy.

Maternal high-fat diet imparts sex-specific offspring behavioral outcomes

To interrogate the contribution of maternal dietary fat content to offspring

neurological development, female wild-type (*C57BL/6J*) mice were placed on either a

high-fat (45% kcal from fat; HFD) or low-fat (10% kcal from fat, sucrose matched; LFD)

diet at approximately 4 weeks of age (juvenile) for six weeks prior to mating (Fig 1a).

Starting females on HFD during the juvenile period best mimics trends observed in fat

consumption in human populations – juveniles tend to get more calories from high-fat

foods, and their eating patterns as adults tend to mimic their adolescent eating patterns[16–]

[20]. HFD females gained significantly more weight than LFD females before pregnancy

and their weight remained elevated throughout gestation (Extended Data Fig 1a-b). No

significant differences were observed in litter size, composition, or maternal care (during

either the light or dark phase; Extended Data Fig 1c-e). Both male and female maternal

high-fat diet (mHFD) offspring weighed significantly more than maternal low-fat diet

(mLFD) offspring throughout life, though placental weight at embryonic day 14.5 (e14.5)

Fig 1. Maternal high-fat diet imparts sex-specific offspring behavioral outcomes.

a, Schematic of maternal low-fat and high-fat diet paradigm (10% and 45% represent %kcal from fat in diet). **b,** Schematic of neonatal maternal separation-induced ultrasonic vocalization (USV) recording and analysis. **c-f,** mHFD decreases neonatal USV number and syllable repertoire similarity in male and female offspring (n=9 mLFD, 8 mHFD male offspring; 18 mLFD, 14 mHFD female offspring from 5 mLFD and 4 mHFD litters). **g,** Schematic of 3-chambered social preference test. **h,** Male mLFD and mHFD offspring display a strong social preference (n=13 mLFD, 15 mHFD male offspring from 5 mLFD and 6 mHFD litters). **i,** Female mHFD offspring have a reduced social preference versus mLFD offspring (n=17 mLFD, 16 mHFD female offspring from 6 mLFD and 6 mHFD litters). **j,** Schematic of sucrose preference test. **k,** Male mHFD offspring display a decreased preference for sucrose versus mLFD offspring (n=10 mLFD, 15 mHFD male offspring from 5 mLFD and 5 mHFD litters). **l,** Female mLFD and mHFD offspring display a strong sucrose preference (n=14 mLFD, 12 mHFD female offspring from 5 mLFD and 5 mHFD litters). Data are mean ± s.e.m.; asterisk denoted p-values are derived from unpaired two-tailed t-tests (**c, e, i, k**) or paired two-tailed t-tests (**d, f**). other p-values (# p<0.0001, † p<0.001, & p<0.01) are derived from one-sample t-tests assessing difference from chance (50%) (**h, i, k, l**).

was not affected by mHFD (Extended Data Fig 1f-g). Offspring behavior was assessed

at neonatal, juvenile, and adult time-points (Fig 1a).

Neonatal maternal separation-induced ultrasonic vocalizations (USVs) provide a

readout of early communicative behaviors in mice, and peak at postnatal day 8 in

C57BL/6J mice[21,22]. USV analyses (Fig 1b) revealed a significant decrease in the number

of USVs emitted, total call time, mean call length, and mean syllable number along with

a significant increase in inter-syllable interval in both male (Fig 1c, Extended Data Fig 2a-

e) and female (Fig 1e, Extended Data Fig 2g-k) mHFD offspring compared to same sex

mLFD offspring. Female, but not male, mHFD offspring USVs had a higher mean

frequency (kHz) compared to mLFD offspring (Extended Data Fig 2d, j). To rule out a

potential shift in USV peak in mHFD offspring, we quantified USVs at P7 and P10 in

addition to P8, but mHFD offspring called less across the assessed neonatal window

(Extended Data Fig 2f, l). In-depth characterization of overall USV composition

(integrating alterations in frequency (kHz), energy (dB), and syllable complexity [e.g.

number of syllables, frequency of calls]) using Mouse Ultrasonic Profile ExTraction

(MUPET[23]) revealed a significant decrease in USV similarity (an integrated measure of

call parameters, see methods for more detail) between mLFD and mHFD male (Fig 1d)

and female (Fig 1f) offspring. While less is known about what USV complexity may

represent in neonates, adult USVs are context-dependent, and specific USV parameters

such as frequency (kHz) are known to vary in response to a stressful environment[24]. To

parse out whether the USV changes were reflective of an overall lack of social

engagement in mHFD offspring, we assessed juvenile sociability using a 3-chamber

social preference test (Fig 1g). Male mHFD offspring showed no change in sociability,

time investigating the social stimulus, or time in the social chamber when compared to

controls (Fig 1h; Extended Data Fig 2m). Female mHFD offspring had a significantly

decreased sociability score, driven by female mHFD offspring spending significantly less

time investigating the social stimulus as compared to controls (Fig 1i; Extended Data Fig

2o). Interestingly, USV emission (number of calls) at P8 significantly positively correlated

with social behavior in females only (r^2=0.39, p=0.006 for females, r^2=0.004, p=0.78 for

males), suggesting that decreased USVs may predict impaired social behavior in females,

but not in males. In a complementary social novelty preference (SNP) assay, male mHFD

showed no change in SNP, time investigating the novel social stimulus, or time in the

novel social chamber compared to mLFD controls (Extended Data Fig 2n). Female mHFD

offspring had a significantly decreased SNP score, driven by female mHFD offspring

spending significantly less time investigating the novel social stimulus and less time in the

novel social stimulus chamber compared to mLFD controls (Extended Data Fig 2p). To

better understand the behavioral changes affecting male mHFD offspring, we determined

average sucrose preference over a 3-day free-choice test (Fig 1j). Mice normally display

a strong preference for sucrose, and diminished sucrose preference is one measure of

anhedonia, or lack of reward/pleasure. Average sucrose preference was significantly

decreased in male, but not female, mHFD offspring (Fig 1k-l). Neither male nor female

mHFD offspring showed a change in total liquid consumption (water + sucrose) over the

3-day testing period (Extended Data Fig 3a,f), and there was no correlation between body

weight and sucrose preference within either mLFD or mHFD groups. We additionally

assessed mobility behavior in a forced swim test (FST; Extended Data Fig 3b). Male

mHFD offspring spent significantly less time immobile, moved significantly further, and

with a higher velocity than mLFD offspring (Extended Data Fig 3c-e). No changes were

observed in FST behavior in female mHFD offspring compared to female mLFD offspring

(Extended Data Fig 3g-i). Lastly, neither male nor female mHFD offspring had any

changes in general activity or anxiety-like behavior in an open field test (OFT; Extended

Data Fig 3j-r). This suggests that the social deficits observed in females are not due to

impaired motor skills or an anxiety-like phenotype, and that the increased velocity and

distance moved by males in the FST are not due to general hyperactivity. In sum, despite

some similar behavioral phenotypes in USVs in early life, by adolescence and into

adulthood the behavior patterns diverge based on sex, with mHFD female offspring

showing social deficits, and mHFD male offspring showing diminished non-social reward

behavior and increased activity in the FST.

mHFD decreases 5-HT in male offspring and increasing 5-HT levels is sufficient to

prevent behavioral phenotypes

We observed altered neonatal USVs, anhedonia, and increased activity during the

FST in male mHFD offspring. These behaviors are associated with diminished brain 5-

HT (5-HT) in mice[25–29]. Previous research has demonstrated that brain 5-HT levels during

discrete windows of development are dependent on placental 5-HT synthesis[30]. The

placenta is a temporary organ that develops during early gestation and surrounds the

fetus to both restrict and facilitate macromolecule and nutrient transport. Further, the

placenta acts as a critical interface between the maternal environment and the developing

fetal brain and responds to maternal diet/environment in a fetal sex-specific manner[31–33].

However, a potential role for fetal sex in placental 5-HT production is unknown.

Assessment of 5-HT levels in mLFD and mHFD placenta and fetal forebrain revealed a

Fig 2. mHFD decreases 5-HT in male offspring and increasing 5-HT levels is sufficient to prevent behavioral phenotypes

a, Male mHFD offspring have decreased fetal forebrain serotonin (n=12 mLFD, 3 mHFD male offspring from 3 mLFD and 3 mHFD litters). **b,** Male mHFD offspring have decreased placenta serotonin (n=18 mLFD, 20 mHFD offspring from 4 mLFD and 6 mHFD litters). **c-d,** mHFD does not impact fetal serotonin levels in female offspring (forebrain: n=11 mLFD, 13 mHFD from 3 mLFD and 3 mHFD litters; placenta: n=11 mLFD, 24 mHFD female offspring from 4 mLFD and 6 mHFD litters) **e-f,** Fetal serotonin levels are significantly correlated between brain and placenta in mLFD males only (n=17 male, 10 female mLFD offspring from 4 mLFD litters; 3 male, 9 female mHFD offspring from 3 mHFD litters). **g,** Maternal dietary tryptophan enrichment increases serotonin levels in mHFD fetal male forebrain (n=14 mLFD, 15 mLFD+Trp, 12 mHFD, 17 mHFD+Trp offspring from 5 mLFD, 5 mLFD+Trp, 3 mHFD, and 5 mHFD+Trp litters) **h-i,** Maternal dietary tryptophan enrichment rescues mHFD-induced decrease in USV number and syllable repertoire similarity (n= 13 mLFD, 21 mLFD+Trp, 23 mHFD, 19 mHFD+Trp offspring from 5 mLFD, 7 mLFD+Trp, 8 mHFD, and 7 mHFD+Trp litters). **j-k,** Maternal dietary tryptophan enrichment rescues sucrose preference in mHFD male offspring and does not affect total consumption (n= 12 mLFD, 9 mLFD+Trp, 11 mHFD, 7 mHFD+Trp male offspring from 6 mLFD, 5 mLFD+Trp, 5 mHFD, and 6 mHFD+Trp litters). Data are mean ± s.e.m. asterisk denoted p-values are derived from unpaired two-tailed t-tests (**a, b**), Pearson's correlation (**e, f**), or two-way ANOVA (fat content x tryptophan content; **g, h, i, j**). other p-values (# p<0.0001, ^p<0.05) are derived from one-sample t-tests assessing difference from chance (50%) (**j**). n.s. not significant.

male-specific decrease in tissue 5-HT in mHFD offspring (Fig 2a-d, also see Extended

Data Fig 4d-g). Circulating maternal 5-HT levels were not changed by HFD, suggesting

that this regulation is within fetal tissue (Extended Data Fig 4a). In agreement with

previous studies demonstrating that placental 5-HT correlates with fetal brain 5-HT[30], we

observed a significant positive correlation between placental and fetal forebrain 5-HT in

mLFD offspring (Fig 2e, see also Extended Data Fig 5f). Interestingly, the correlation was

exclusively driven by male 5-HT levels, but mHFD disrupted the correlation between

placental and brain 5-HT levels in males, potentially a floor effect due to low 5-HT levels

(Fig 2f, see also Extended Data Fig 5g).

144   Quantitative real-time PCR (qPCR) showed a decrease in both the 5-HT synthesis

enzyme tryptophan hydroxylase 2 (*Tph2*) and the 5-HT transporter (*5HTT*) in male mHFD

placenta (Extended Data Fig 4b-c). To better understand the dynamics of placental 5-HT

in response to mHFD, we additionally quantified levels of 5-HT, the 5-HT precursor

tryptophan, and the 5-HT metabolite 5-hydroxyindoleacetic acid (5-HIAA), in the placenta

using high-performance liquid chromatography (HPLC; Extended Data Fig 4d-g and

ExtendedDataFig4_RawData_Stats). In agreement with decreased *Tph2* levels, 5-HT

synthesis (ratio of 5-HT to tryptophan) was significantly decreased in male, but not

female, mHFD offspring placenta (Extended Data Fig 4d, f). We did not observe any

changes in the expression of the alternate 5-HT synthesis enzyme tryptophan

hydroxylase 1 (*Tph1*), monoamine oxidase A (*MAOA*), an enzyme responsible for the

oxidative deamination of neurotransmitters, including 5-HT, or the rate-limiting enzyme

indoleamine 2,3-dioxygenase (*Ido1*) that promotes the degradation of tryptophan through

the kynurenine pathway in male placentas (Extended Data Fig 4b). In line with this, we

also did not observe any changes in the protein levels of quinolinic acid, and kynurenic

acid levels were undetectable in e14.5 brain and placental tissue, suggesting that

tryptophan is not being diverted into the kynurenine pathway in mHFD offspring

(Extended Data Fig4h-i). Comparing the ratio of 5-HIAA to 5-HT (5-HT turnover) revealed

a modest but significant increase in 5-HT turnover in male mHFD offspring placenta,

suggesting that in addition to diminished 5-HT synthesis, increasing turnover may

contribute to total decreased placental 5-HT levels in male mHFD offspring (Extended

Data Fig 4d). In the fetal forebrain, we did not see changes in 5-HT synthesis or turnover,

but we did see a decrease in tryptophan in both male and female mHFD offspring

(Extended Data Fig 4e, g). Lastly, *MAOA* expression was significantly decreased in

female mHFD placenta, suggesting that an alternate neurotransmitter may be affected in

female mHFD offspring, and that the placenta may play a critical role as a source for brain

neurotransmitters other than 5-HT (Extended Data Fig 4c).

To determine if increasing 5-HT in male mHFD offspring could rescue the mHFD-

induced behavioral outcomes, we sought to rescue 5-HT by supplementing the maternal

low- and high-fat diets with tryptophan. Tryptophan is the precursor to 5-HT, and placental

5-HT levels are largely dependent on maternal tryptophan levels[30,32]. Tryptophan

supplementation (+Trp; provided in LFD or HFD chow for the same duration as in Fig 1a)

did not affect maternal weight gain, litter composition (i.e., size and sex ratio), or offspring

weights (Extended Data Fig 5a-d).

Maternal tryptophan supplementation increased placental 5-HT levels in male

mLFD offspring (Extended Data Fig 5e) in support of previous literature demonstrating

that placental 5-HT levels are dependent on maternal tryptophan levels. In male mHFD

offspring, tryptophan supplementation was unable to substantially increase placental 5-

HT levels (Extended Data Fig 5e). Conversely, maternal tryptophan supplementation did

increase fetal forebrain 5-HT levels in male mHFD, but not mLFD offspring (Fig 2g).

Further, Pearson correlation analyses demonstrated that maternal tryptophan enrichment

disrupted the significant positive correlation between brain and placental 5-HT levels in

male mLFD offspring (Extended Data Fig 5f). No significant correlation was detected in

mHFD (in agreement with our previous finding) or mHFD+Trp offspring (Extended Data

Fig 5g). This suggests that in mHFD+Trp offspring, the fetal brain produces 5-HT

independently of the placenta. Together these data demonstrate that in males, placental

5-HT levels are dependent on maternal tryptophan levels and can be disrupted by

maternal diet, but fetal forebrain 5-HT levels are not solely dependent on placental 5-HT

levels. Further, mHFD led to decreased 5-HT levels in adult male midbrain – where the

majority of 5-HT neuronal cell bodies reside – levels that were also rescued by maternal

tryptophan enrichment (Extended Data Fig 5h).

To determine if mHFD-induced behavioral deficits in male offspring were rescued

by maternal tryptophan supplementation, we assessed USVs and sucrose preference in

male mLFD and mHFD offspring. Tryptophan supplementation did not impact USV or

sucrose preference behavior in male mLFD offspring (Fig 2h-k, Extended Data Fig5 i-m).

However, in mHFD offspring, maternal tryptophan supplementation fully rescued the

decreased USV call number and similarity, as well as the increased inter-syllable interval

(Fig 2h-i, Extended Data Fig 5k), partially rescued the deficits seen in total USV call time,

mean call length, and mean syllable number (Extended Data Fig5 i-m), as well as the

diminished sucrose preference (Fig 2j-k). In all cases, mHFD+Trp offspring behavior was

indistinguishable from that of mLFD or mLFD+Trp offspring. Of note, maternal tryptophan

supplementation rescued male mHFD offspring USV phenotypes despite the observation

that male mHFD+Trp offspring still weigh more than mLFD and mLFD+Trp offspring (see

Extended Data Fig 5c), suggesting that changes in offspring weight is not driving the USV

phenotypes. Further, maternal tryptophan supplementation did not alter male offspring

social behavior or open field behavior (Extended Data Fig 5n-u).

Although we did not expect to see a rescue of behavioral phenotypes in female

mHFD+Trp offspring because mHFD did not impact 5-HT levels in females, we assessed

midbrain 5-HT and behavior outcomes. Adult female offspring midbrain 5-HT levels were

unaffected by mHFD or tryptophan supplementation (Extended Data Fig 6a). Maternal

tryptophan enrichment did not rescue female mHFD offspring USV behavior, but

interestingly did decrease USV call similarity and call length in mLFD offspring (Extended

Data Fig 6b-h). Maternal tryptophan supplementation also increased USV frequency

(kHz) in female offspring regardless of maternal dietary fat content (Extended Data Fig

6g). This suggests that despite similar neonatal behavioral outcomes in male and female

mHFD offspring, the mechanisms driving USV behavior in males and females may be

distinct. Interestingly, maternal tryptophan enrichment increased sucrose preference in

female offspring regardless of maternal dietary fat intake (Extended Data Fig 6i). Maternal

tryptophan enrichment did not rescue female mHFD social behavior deficits (Extended

Data Fig 6j-m). In sum, these data demonstrate 1) a developmental role for 5-HT in males

that is perturbed by maternal high-fat diet, and 2) that maternal tryptophan enrichment is

sufficient to rescue mHFD-induced brain 5-HT deficiency and associated 5-HT-

dependent behaviors in males.

mHFD induces Tlr4-dependent inflammation driving offspring behavior changes

Maternal HFD creates an environment of chronic inflammation that affects the

developing fetus as well as the placenta[34]. Growing literature has demonstrated complex

relationships between inflammation and 5-HT[35–37], and we hypothesized that the mHFD-

induced inflammatory environment is translated to the fetus, and responsible for

decreased 5-HT in male mHFD offspring. At e14.5, we observed increased macrophage

(F4/80+ cells located on the fetal side of the placenta) density in male and female mHFD

placenta (Extended Data Fig 7a), as well as increased Iba1 and CD68 immunoreactivity

(labeling microglia and phagosomes, respectively) in both male and female fetal dorsal

raphe nuclei (DRN), where the majority of serotonergic cell bodies reside (Fig 3a).

Microglia influence central circuits in numerous ways, such as by pruning excess

synapses or phagocytosing excess or dying progenitor cells[38–41]. Interestingly, qPCR for

genes important for phagocytosis from isolated midbrain microglia at e14.5 did not reveal

any significant changes in gene expression, although the levels of some genes were near

or below the limit of detection (see Methods, Extended Data Fig 7b). Given the increased

immunoreactivity seen by immunohistochemistry (IHC), we further investigated potential

interactions between microglia and the developing central 5-HT system by 3D

reconstructing microglia from e14.5 DRN. IMARIS reconstructions revealed that 5-HT

signal comprised a higher percentage of phagosomes (CD68) in male mHFD microglia

versus mLFD (Fig 3b), demonstrating that microglial phagocytosis of serotonergic

neurons is increased in males in the context of mHFD, leading to decreased central 5-HT

bioavailability. Postnatal microglia express the 5-HT receptor 5HT2B[42] and decrease their

phagocytic capacity in response to 5-HT[43]. Further, mice lacking microglial 5HT2B display

prolonged neuroinflammation in response to lipopolysaccharide (LPS), a classic agonist

of macrophage toll-like receptor signaling[44]. Thus, we sought to investigate potential

feedback between 5-HT levels and microglial phagocytosis of 5-HT neurons. Microglial

5-HT phagocytosis was rescued in mHFD+Trp offspring, in parallel with 5-HT levels (Fig

3b; see Figure 2g), suggesting that in the context of mHFD, enrichment of 5-HT levels is

sufficient to prevent further aberrant microglia phagocytosis. Consistent with our

observations that mHFD does not decrease 5-HT levels in females, we did not see

increased 5-HT signal within phagosomes in mHFD female microglia (Extended Data Fig

7c). Interestingly, there was a significant increase in 5-HT signal within phagosomes in

reconstructed microglia from mHFD+Trp female microglia (Extended Data Fig 7c),

possibly contributing to the trend towards lower midbrain 5-HT we observed in the adult

female mHFD+Trp offspring (see Extended Data Fig 6a). This trend parallels the open

field behavior, with mHFD+Trp female offspring showing increased activity in the open

field test (see Extended Data Fig 6n-o), suggesting that fetal microglial interactions with

5-HT neurons may dictate different behaviors in males and females.

The presence of 5-HT receptors on placental macrophages, or the potential

influence of 5-HT on placental macrophages is unknown, although recent studies have

demonstrated the presence of functional 5-HT transporters in placental tissue[45,46]. Thus,

we sought to investigate crosstalk between placental 5-HT and placental macrophage

density as well as fetal brain 5-HT and microglial activity. Maternal tryptophan

supplementation did not significantly influence placental macrophage density in either

mLFD or mHFD male or female offspring (Extended Data Fig 7d). Considering our finding

that male mLFD+Trp offspring have increased placental 5-HT (see Extended Data Fig

5e), this suggests that placental macrophage density is not significantly influenced by 5-

HT.

To further investigate the potential pro-inflammatory pathways leading to

increased placental macrophage and fetal brain microglia density, we assessed a panel

of pro-inflammatory markers with qPCR (Extended Data Fig 7e). This revealed a specific

increase in Toll-like receptor 4 (Tlr4) in both male and female mHFD placenta (Extended

Data Fig 7e). Tlr4 is a pattern recognition receptor known to initiate pro-inflammatory

signaling cascades in response to infectious pathogens (e.g. bacterial, viral), and whose

expression is increased in response to saturated fats/obesity[47–50]. Isolated fetal microglia

and placental macrophages from obese dams have exaggerated cytokine production in

response to lipopolysaccharide (LPS), a classic Tlr4 agonist[33].

284   To determine if increased *Tlr4* signaling in macrophages was causal to decreased

285 5-HT and 5-HT-dependent behaviors in males, or to behavioral changes in females, we

Fig 3. mHFD induces Tlr4-dependent inflammation driving offspring behavior changes

a, Increased microglia and phagosome immunoreactivity in male and female mHFD DRN at e14.5 (scale=50μm; n=4 male and 3 female/diet from 3mLFD and 3mHFD litters). **b,** IMARIS reconstructions of DRN microglia reveals increased microglial phagocytosis of serotonin in males at e14.5. Statistics are displayed for animal average (large circles), small circles are individual microglia reconstructions (n=6 mLFD, 5mLFD+Trp, 5 mHFD, 6 mHFD+Trp mice from 3 mLFD, 2 mLFD+Trp, 3 mHFD, and 3mHFD+Trp litters). **c,** IMARIS reconstruction of DRN microglia in male control and *Tlr4* cKO e14.5 offspring (n=6 mLFD control, 6 mLFD cKO, 5 mHFD control, and 3 mHFD cKO male offspring from 7 mLFD and 5 mHFD litters). **d,** Decreased sucrose preference is rescued in male mHFD offspring lacking macrophage-Tlr4 signaling. (n=11 mLFD, 10 mLFD cKO, 11 mHFD, and 10 mHFD cKO from 5 mLFD and 6 mHFD litters). **e,** Decreased social preference is rescued in female mHFD offspring lacking macrophage-Tlr4 signaling. (n=9 mLFD, 6 mLFD cKO, 7 mHFD, and 6 mHFD cKO from 5 mLFD and 5 mHFD litters). **f-g,** Tlr4-reporter activity in response to e14.5 brain (f) and placenta (g) lysate from mLFD and mHFD offspring (n=6 offspring/sex/diet from 5 mLFD and 5 mHFD litters). **h-i,** Endotoxin levels in mLFD and mHFD brain (h) and placenta (i) tissue lysate (n=3-5 mLFD male, 4-5 mLFD female, 5 mHFD male, 4-5 mHFD female from 5 mLFD and 5 mHFD litters). Data are mean ± s.e.m.; asterisk denoted p-values are derived from two-way ANOVA (**a-i**). other p-values (# p<0.0001, † p<0.001, & p<0.01) are derived from one-sample t-tests assessing difference from chance (50%) (**d, e**). n.s. not significant

generated a conditional Tlr4 knock-out mouse (cKO) line using Cx3cr1-Cre BAC-

transgenic[51] and Tlr4[f/f] mice[52] (Extended Data Fig 7f). Female control (*Tlr4[f/f]*) mice were

transitioned to HFD for six weeks, then subsequently mated to male *Cx3cr1-Cre;Tlr4[f/f]*

mice to generate control and cKO mHFD offspring. qPCR revealed significant (~95%)

knockdown of *Tlr4* in cKO microglia (Extended Data Fig 7f). 5-HT was increased in male

fetal forebrain and adult midbrain in mHFD cKO offspring (Extended Data Fig 7g-i).

IMARIS reconstructions of mLFD and mHFD microglia from control and *Tlr4* cKO e14.5

DRN revealed a significant rescue of 5-HT phagocytosis by microglia in mHFD *Tlr4* cKO

male offspring (Fig 3c). No significant changes were detected in females (Extended Data

Fig 7j). Loss of *Tlr4* decreased USV similarity in male mLFD offspring while increasing it

in male mHFD offspring (Extended Data Fig 8a), suggesting a role for macrophage *Tlr4*

signaling in USV behavior independent of maternal dietary fat composition. In male mHFD

offspring, loss of *Tlr4* completely rescued mHFD-induced anhedonia (Fig 3d, Extended

Data Fig 8c), and did not influence male social preference (Extended Data Fig 8d).

Interestingly, female mHFD *Tlr4* cKO offspring did not display a preference for sucrose

(Extended Data Fig 8e), suggesting that increased Tlr4 signaling in female mHFD

offspring may be preventing anhedonia. *Tlr4* knockdown partially rescued the number of

USVs, but did not rescue total call time, or mean call length in male mHFD cKO offspring

(Extended Data Fig 8a), suggesting that different USV metrics may be driven by different

mechanisms. Corroborating this, in female mHFD cKO offspring, we saw significant

improvement of USV similarity (Extended Data Fig 8b) but no change in the number of

USVs. *Tlr4* cKO resulted in a partial rescue of total call time, inter-syllable interval, and

mean syllable number in female mHFD cKO mice, but no rescue of mean call length or

mean frequency (Extended Data Fig 8b). Juvenile female mHFD cKO offspring showed

a complete rescue of social preference (Fig 3e, Extended Data Fig 8f), suggesting that

mHFD-induced *Tlr4*-dependent inflammation in females is responsible for most

behavioral outcomes, through a non-5-HT mechanism.

To understand how mHFD results in activated Tlr4 signaling in fetal tissue, we

used a Tlr4-reporter cell line (HEK-Dual™ mTLR4 (NF/IL8), Invivogen) to determine if

mHFD tissue lysate stimulates Tlr4 signaling. Application of lysate from mLFD and mHFD

placenta and brain tissue revealed robust Tlr4 activation in response to mHFD versus

mLFD tissue (Fig 3f-g). As dietary fats reach the fetus through the placenta, and some

literature suggests that saturated fatty acids (SFAs) stimulate Tlr4 signaling[53], we

determined if the increased SFAs in our model could directly activate Tlr4 signaling by

stimulating the reporter cell line with six SFAs that were most abundant (and most

increased) in our HFD model. While stearic acid was able to significantly activate Tlr4

signaling, as previously described[54], the magnitude of the response was minimal

(Extended Data Fig 7k).

Previous literature has demonstrated that pre-gravid obesity and high-fat diet

increase circulating endotoxin levels[55–57], and that LPS injected into pregnant dams can

reach fetal tissue[58]. As Tlr4 is classically activated by bacterial endotoxins, we

hypothesized that mHFD offspring tissue may be exposed to and accumulate endotoxin.

Thus, we quantified bacterial endotoxin presence by limulus amebocyte lysate (LAL)

assay. We detected significantly more endotoxin in mHFD placenta and fetal brain tissue

from male and female offspring (Fig 3h-i), suggesting that mHFD increases endotoxin

load in fetal tissue and activates Tlr4 signaling.

Maternal overnutrition causes decreased prenatal 5-HT in human male brain tissue

To determine to what degree our findings are translatable to the human population, we obtained a large (37 individuals) cohort of matched tissues (fetal brain, fetal placenta, and maternal placenta (decidua)) from fetuses collected via elective termination at 72-82 days post conception (d.p.c.), a timeframe that closely matches the embryonic developmental window used in our mouse studies (14.5-16.5 d.p.c.; Fig 4a). PCR for the Y chromosome (*SRY;* Supplementary Table 2) identified 17 male and 20 female fetal tissue sets, with the average age of 78.2 d.p.c. for both sexes (Extended Data Fig 9a). No maternal data were available, so we assessed if triglyceride accumulation in the placenta could serve as a proxy for maternal dietary fat consumption, given that in humans, placental triglyceride/lipid levels have been shown to be increased in the context of maternal obesity[59–61]. To show proof of principle in our mouse model, triglyceride measurements from mLFD and mHFD placenta revealed a significant increase in triglyceride levels in mHFD offspring placenta, in agreement with previous literature[62] (Extended Data Fig 9b), and placental triglyceride levels were significantly negatively correlated with fetal brain 5-HT in males (Extended Data Fig 9c). Thus, in the human samples, we assessed decidual triglyceride accumulation to use as a proxy for maternal dietary fat consumption. Triglyceride levels trended higher in pregnancies with female fetuses, but there was no significant difference between triglyceride accumulation in decidua associated with male or female pregnancies (Extended Data Fig 9d). Correlation of gene expression with maternal triglyceride accumulation from bulk RNA-seq from 16 matched male and 19 matched female brain and placenta samples revealed a strong dimorphic trend in gene expression. In male placenta, differentially correlated genes were

predominantly over-expressed with increasing maternal triglycerides (i.e. gene

expression increased as maternal triglycerides increased); notably, the opposite effect

was seen in female placenta (Fig 4b). In the male brain, only 45 genes significantly

correlated with maternal triglyceride accumulation (either positive or negative), whereas

781 genes were significantly correlated in female brain tissue (Fig 4b). At collection, the

entire fetal brain tissue was collected and thus we could not control for region-specificity.

To address this, we used the Allen Brain Atlas Developmental Human Transcriptome

(https://www.brainspan.org/) to identify marker genes for major brain regions in human

development (8-12 weeks post conception) and we plotted their gene expression

(Extended Data Fig 9e). The majority of our samples showed consistent gene expression

across all brain regions (i.e. one region was not oversampled). Gene Ontology (GO)

Enrichment Analysis of genes significantly positively correlated with maternal triglyceride

accumulation revealed enrichment of inflammatory signaling pathways in both male and

female placenta (Fig 4c; full GO enrichment data is available in Supplementary Table 1).

Correlation analyses of targeted genes reinforced these pro-inflammatory phenotypes;

assessment of common macrophage, trophoblast and syncytiotrophoblast markers[63]

revealed specific gene associations predominantly within macrophages (Extended Data

Fig 9f, Fig 4b). In agreement with our findings in mice, *TLR* expression was positively

correlated with maternal triglyceride accumulation in males, along with several genes

associated with MyD88-independent TLR4 signaling (Extended Data Fig 9f). In males,

GO enrichment from placental genes that were negatively correlated with maternal

triglycerides revealed a pattern of repressive cell differentiation, particularly of immune

cells (e.g. RUNX1 differentiation and myeloid cell differentiation). That is, with increasing

maternal triglyceride accumulation, gene expression of factors important for cell

differentiation decreased (Fig 4c). In females, placental genes that were negatively

correlated with maternal triglycerides were often involved in vascular growth and

development, suggesting that maternal triglyceride accumulation may decrease placental

vasculogenesis/angiogenesis (Fig 4c), an important finding given that changes in the

placental vascular network are associated with numerous pathologies, such as

preeclampsia[64,65]. Inflammatory responses were additionally enriched in female brain

(e.g. response to TGF-β), but too few genes were correlated in males to determine if this

association was true independent of sex. These inflammatory transcriptomic patterns

support our findings in mice that male mHFD offspring primarily accumulate endotoxin in

placental tissue, whereas female mHFD offspring primarily accumulate endotoxin in brain

tissue (see Fig 3h-i). Lastly, the top GO enrichment categories from female brain

transcripts negatively correlated with maternal triglyceride accumulation involved

neuronal development (Fig 4c), suggesting that maternal high-fat diet and accompanying

fetal inflammation may impede brain development.

Finally, to determine if 5-HT levels were impacted by maternal triglyceride

accumulation, we assessed 5-HT in human male and female fetal brain and placenta.

Fig 4. Maternal triglycerides negatively correlate with male fetal brain 5-HT

a, Human fetal development at 63-81 days post conception (d.p.c.) is roughly equivalent to mouse embryonic development at 14-16 d.p.c. b, Scatter plots showing transcript (TPM) correlation with maternal triglycerides versus -Log$_{10}$(p-value) in male and female placenta and brain bulk sequencing data (n=16 matched male tissue sets, 19 female). Red dots indicate a significant p-value (<0.05). c, GO Enrichment analysis of transcripts significantly correlated with maternal triglyceride accumulation demonstrates robust and distinct upregulation in immune responses in male and female placenta in response to increasing maternal triglyceride accumulation. GO enrichment of transcripts significantly correlated with maternal triglyceride accumulation shows downregulation of neuronal development and function in female brain tissue in response to increasing maternal triglyceride accumulation. d, Multivariate dot plots show Pearson correlations between brain and placental serotonin from male and female human fetal tissues. Dot color represents the age of the matched tissue set, and dot size represents the maternal decidual triglyceride level (mg/dL), a proxy for maternal dietary fat intake. Brain and placental serotonin levels are significantly positively correlated in male tissues between 72 and 82 d.p.c. Brain and placental serotonin levels are not correlated in female tissues (n=17 matched male tissue sets, 20 female). e, Brain serotonin levels are significantly negatively correlated with decidual triglyceride levels in males only. Placental serotonin levels trend towards a negative correlation with decidual triglycerides in males only. Data are mean ± s.e.m.; n represents biologically independent matched human tissue set. asterisk denoted p-values are derived from Pearson's correlations.

Correlation analyses revealed a significant positive correlation between placental and

brain 5-HT only in male fetal tissues (Fig 4d). This is consistent with previous literature

(sex not reported)[30] and our findings in male mice (see Fig 2e, Extended Data Fig 5f).

Importantly, brain region marker gene expression did not correlate with 5-HT levels, and

the top samples that most overrepresented midbrain gene expression profiles did not

drive the correlation. Correlation analyses further revealed a negative correlation between

brain or placental 5-HT levels and maternal triglyceride accumulation in male tissue only

(Fig 4e). This is in agreement with our findings in mice that mHFD results in decreased

brain and placental 5-HT levels in male fetal tissue (see Fig 2a-b, Extended Data Fig 4d-

e). In sum, in both mice and humans, maternal high-fat diet results in the perpetuation of

inflammation to the fetal brain and placenta in both sexes. In males, this inflammatory

response results in aberrant phagocytosis of 5-HT neurons and long-lasting decreased

levels of brain 5-HT and decreased reward behavior.

Discussion

Here, we demonstrate a fundamental, sex-biased mechanism through which maternal

high-fat diet increases offspring susceptibility to neuropsychiatric disorder development.

Our results demonstrate that, in the context of maternal high-fat diet, endotoxin

accumulation mediates *in utero* inflammation through the pattern recognition receptor Tlr4

in males and females, leading to increased macrophage reactivity in both the placenta

and the fetal brain. In female mice, Tlr4-dependent inflammation causes diminished social

preference through a 5-HT-independent mechanism. In human females, maternal

decidual triglyceride accumulation is negatively associated with brain development,

suggesting that Tlr4-dependent inflammation impacts neuronal development in females,

though the target neuronal population is not yet known. In male mHFD offspring,

embryonic microglia aberrantly phagocytose 5-HT neurons in the DRN, leading to

diminished brain 5-HT from embryonic stages through adulthood, and offspring

anhedonia. In human males, maternal decidual triglyceride accumulation is associated

with pro-inflammatory signaling pathways in both sexes, and negatively correlated with

brain 5-HT levels in males only, reinforcing our findings in mice.

Seminal work reported the placenta as a transient source of 5-HT for the fetal

brain[30], though recent studies suggest that the fetal placenta lacks the enzymes required

for 5-HT synthesis[46]. No studies have investigated whether fetal sex plays a role in 5-HT

synthesis or transfer in the placenta. Further, whether developmental 5-HT levels are

reflective of adult 5-HT levels, and if disruptions to 5-HT levels during this window impact

the brain long-term are unknown. Here, we show that fetal placenta and brain 5-HT levels

are susceptible to perinatal inflammation in males only, and that decreased fetal 5-HT

levels are reflective of decreased adult brain 5-HT levels in the context of maternal high-

fat diet. We further found that transcript levels of both the neuronal 5-HT synthesis

enzyme and 5-HT transporter (*Tph2* and *5HTT*, respectively) were decreased in male

mHFD placenta, but we did not distinguish between the maternal and fetal compartments

of the placenta. By manipulating fetal 5-HT levels through maternal dietary tryptophan

enrichment, we can shed light on the relationship between placental and fetal brain 5-HT.

While at baseline there is a significant positive correlation between placental and fetal

brain 5-HT in males only, maternal tryptophan supplementation disrupted the correlation.

This suggests that either the placenta is not the sole source of fetal brain 5-HT, or that

the fetal brain is capable of sensing and limiting the influx of placental 5-HT in cases of

excess. Alternatively, maternal dietary changes may alter the timing of the switch from

placental 5-HT dependence to fetal midbrain 5-HT dependence. Careful investigation of

the spatiotemporal expression of placental 5-HT transport and synthesis enzymes in male

and female placenta and decidua is warranted to further clarify this topic.

5-HT has been postulated to influence macrophage phagocytic activity and cytokine

production for decades[66,67]. Here, we demonstrate a novel male-specific potential

feedback loop through which microglial phagocytosis influences central 5-HT levels,

which in turn influence microglial phagocytic activity. mHFD results in increased

inflammation dependent on macrophage Tlr4 signaling, which results in microglial over-

phagocytosis of 5-HT neurons in the DRN of males only. Considering 5-HT is thought to

decrease phagocytic capacity[43], this resulting decrease in 5-HT thus propagates the over-

phagocytosis of the mHFD microglia. Both abrogation of overactive Tlr4-signaling in

macrophages and increased 5-HT bioavailability via maternal tryptophan

supplementation were sufficient to rescue central 5-HT levels, demonstrating the

reciprocal relationship between inflammation and 5-HT in the male fetal brain.

In closing, mechanisms through which chronic maternal inflammation impact

offspring neurodevelopment are scarce. In our mouse model, we demonstrate that fetal

inflammation is propagated by macrophage-specific Tlr4 in both male and female mHFD

offspring, along with sex-specific neurodevelopmental and behavioral outcomes. Further,

we generated a large sequencing dataset of human fetal tissue from both male and

female pregnancies, which demonstrated 1) a high degree of fetal inflammation in both

male and female placenta, and particularly in female brain in response to increasing

maternal triglyceride accumulation, and 2) sex-specific inflammatory responses. These

data will shed light on which other fetal inflammatory pathways might be responsive to

maternal triglyceride accumulation and will provide direction for future studies.

Methods

Animals: All procedures relating to animal care and treatment conformed to Duke

Institutional Animal Care and Use Committee (IACUC) and NIH guidelines. Animals were

group housed in a standard 12:12 light-dark cycle. The following mouse lines were used

in this study: *C57BL/6J* (Jackson Laboratory, stock no. 000664), *C57BL/6NCrl* (Charles

River, #027), Cx3cr1-Cre BAC-transgenic[51], *Tlr4^{f/f}* (JAX 024872[52]). Cx3cr1-Cre mice were

developed in *C57BL/6N* mice but have been backcrossed with *C57BL/6J* mice for multiple

generations. The Cre transgene was maintained in males for all experimental studies.

Jackson Labs maintains the *Tlr4^{f/f}* mice on a *C57BL/6J* background, though single

nucleotide polymorphism (SNP) analysis suggests that the original mice provided to JAX

were a mixed *C57BL/6N;C57BL/6J* strain. Our in-house breeding has backcrossed them

with *C57BL/6J* for multiple generations.

Diets: Female *C57BL/6J* mice were placed on randomly assigned diet at 4-weeks old

and given *ad libitum* access to their assigned diet chow and water. Weight gain was

assessed three times/week. After 6-weeks on diet, females were mated with *C57BL/6J*

males. Pregnancy was determined by the presence of a copulation plug (gestational day

0.5) and maternal weight was assessed daily. Dams were maintained on their assigned

diet chow throughout gestation and until the pups were weaned. At weaning, offspring

were group housed with sex-matched littermates and provided *ad libitum* access to rodent

chow 5001 (standard rodent chow, 13% kcal from fat). Low-fat, high-fat, low-

fat+tryptophan, and high-fat+tryptophan diets were purchased from Research Diets (low

fat: D12450Hi, high-fat: D12451i). Tryptophan enriched diets were custom formulations

resulting in 1% final tryptophan content.

Tissue Collection and Analyses: <u>*Murine Tissue collection*</u>*:* Pregnant dams were

euthanized using CO_2 following Duke University Animal Care and Use guidelines. Uterine

horns were dissected and placed on ice in sterile 1X PBS. Individual embryos were

separated, and placenta, brain, and tail tissue (for genotyping) were collected. Placenta

were dissected into three equal pieces. One piece was flash-frozen in liquid nitrogen at

stored at -80°C until ELISA. One piece was placed into 4% paraformaldehyde in PBS

(PFA, Sigma) for immunohistochemistry (IHC). The final piece was placed in TriZol and

stored at -80°C. Dissected fetal brains were cut in half coronally; fetal forebrain tissue

was flash frozen for ELISA while the remainder of the brain was placed into 4% PFA for

IHC. For adult brain collections, offspring were anesthetized with CO_2 and transcardially

perfused with saline. Further dissections were done to isolate the midbrain, and brain

tissues were immediately flash frozen in liquid nitrogen and stored at -80°C until

processing.

Microglia isolation: Microglia were isolated using CD11b-beads as previously described[68].

Human tissue collection: Human placenta (maternal decidua separated from fetal

placenta) and fetal brain tissues were obtained from the NIH-supported (5R24HD000836)

Laboratory for the Study of Embryology at the University of Washington, Seattle between

the years 1999 and 2010 and used under a protocol approved by the Duke University

Institutional Review Board (Pro00014066). Fetuses with known chromosomal

abnormalities were not collected. At the time of collection, tissue specimens were placed

in cryogenic tubes and immediately frozen on dry ice and stored at -80°C until required

for RNA or protein extraction. Prior to collection, informed consent was obtained from all

individuals undergoing the elective termination.

Immunohistochemistry: Placenta and brain tissue were fixed in 4% PFA overnight at 4°C,

cryoprotected in 30% sucrose + 0.1% sodium azide in PBS (Sigma), and embedded in

OCT (Sakura Finetek, Torrance, CA) before being cryo-sectioned directly onto Superfrost

slides (Fisher) at 40µm. Sections were frozen at -20°C for storage. For IHC, sections were

permeabilized in 1% Triton X-100 in PBS for 12 minutes, blocked at room temperature (RT) with 5% goat serum (GS) in PBS + 0.1% Tween-20, then incubated for 2 nights at 4°C with chicken anti-Iba1 (Synaptic Systems, 234 006) or goat anti-Iba1 (Novus, NB100-1028), rat anti-CD68 (Biolegend, 137002), and rabbit anti-5-HT (Sigma, S5545) or rat anti-F4/80 (Abcam, ab6640). Sections were then washed with PBS and incubated in the following secondary antibodies: anti-rabbit Alexa-488, anti-chicken Alexa-647 or anti-goat Alexa-647, and anti-rat Alexa-568 (brain), or anti-rat Alexa-568 (placenta; 1:200; ThermoFisher), and DAPI (100µg/mL). Ten z-stacks of 0.5µm optical thickness were taken at 20X using a Zeiss AxioImager.M2 (with ApoTome.2) from at least 5 sections (each 400µm apart) from the fetal compartment of each placenta for F4/80+ cell quantification. Ten z-stacks of 0.5µm optical thickness were also taken at 20X from at least 3 fetal brain sections/animal (each section being 200µm apart) for Iba1 and CD68 immunoreactivity. For IMARIS reconstructions, z-stacks of 0.25µm optical thickness were taken to capture entire microglia. Surface reconstructions of Iba1, CD68, 5-HT, and DAPI were done for at least 19 microglia/diet from 6 mLFD, 5mLFD+Trp, 5 mHFD, and 6 mHFD+Trp male offspring and at least 12 microglia/diet from 3 mLFD, 5 mLFD+Trp, 6 mHFD, and 5 mHFD+Trp female offspring (3 litters/diet). Reconstructions from were done for at least 16 microglia/diet from 6 mLFD control, 6 mLFD cKO, 5 mHFD control, and 3 mHFD cKO male offspring and 7 mLFD control, 5 mLFD cKO, 5 mHFD control, and 4 mHFD cKO female offspring from 7 mLFD and 5 mHFD litters. Slides with cryosectioned tissue were de-identified and the individual performing the IHC, imaging, and image analyses was blind to sex, genotype, and maternal diet at all times. Statistics were run on animal average values, not individual microglia values.

5-HT ELISA: Mouse placenta and brain tissue were thawed on ice and homogenized in

freshly prepared lysis buffer (final concentrations: 1mg/mL ascorbic acid, 0.147% NP-40,

1X HALT Protease Inhibitor (Thermo 78443) in 1X PBS). Samples were then centrifuged

at 13,000 xg for 20 minutes; supernatant was diluted 1:2 for placenta, or undiluted for

fetal brain preparations, and loaded and processed according to manufacturer

instructions (Enzo, ADI-900-175). Dam blood was collected in heparinized capillary tubes

and immediately transferred to Eppendorf tubes on ice before being centrifuged at 1.6 x

1000 rpm for 20 minutes at 4°C. Plasma was stored at -80°C for a maximum of 2 weeks

before assessment for 5-HT according to manufacturer's instructions (1:16 dilution).

Quinolinic Acid/Kynurenic Acid ELISA: Mouse placenta and brain tissue were thawed on

ice and homogenized with Dounce homogenizers in 500µL 1X PBS with 1X HALT

Protease Inhibitor. Samples were then centrifuged at 13,000 xg for 10 minutes and

supernatant processed according to manufacturer instructions (ImmuSmol IS I-0200, IS

I-0100).

Decidual Triglycerides: Human decidua samples (stored at -80°C) were kept on dry ice

to remove ~150mg tissue which was then thawed on ice, and homogenized in 1mL NP40

Substitute Assay Reagent (Cayman Chemical, 10010303). Homogenized samples were

centrifuged at 10,000 xg for 10 minutes at 4°C and the supernatant was processed

(undiluted) according to manufacturer instructions at room temperature (Cayman

Chemical, 10010303). Mouse placenta samples (stored at -80°C) were thawed on ice and

~40mg of tissue was homogenized in 500µL NP40 Substitute Assay Reagent.

Homogenized samples were centrifuged at 10,000 xg for 10 minutes at 4°C and the

supernatant was processed (undiluted) according to manufacturer instructions at room

temperature (Cayman Chemical, 10010303).

High-performance liquid chromatography (HPLC) with electrochemical detection: Female

C57BL/6NCrl (Charles River #027) mice were placed on LFD or HFD as described above,

with the exception that these diets were not irradiated. Pregnant females were euthanized

at gd14.5 by lethal i.p. injection with ketamine/xylazine. Placentae were separated from

each fetus, the amniotic sac and maternal membranes were removed, and the resulting

discoid placenta was cut in half sagitally. One half was snap-frozen in liquid nitrogen for

HPLC analysis. Fetal brains were dissected and snap-frozen in liquid nitrogen as well.

Placenta and brain samples were analyzed for 5-HT, the 5-HT precursor tryptophan (Trp),

and the 5-HT metabolite 5-hydroxyindoleacetic acid (5-HIAA). On the day of analysis,

500µl of ice-cold standard buffer (0.5 mM sodium metabisulfite, 0.2 N perchloric acid, and

0.5 mM EDTA) was added to thawed (on ice) placenta tissue and 250µl was added to

thawed brain tissue. Tissues were then disrupted by sonication until completely

homogenized, then centrifuged at 16,000 x g for 10 min at 4°C. The supernatant was

collected and filtered through a 0.45µm membrane via centrifugation (Durapore PVDF

centrifugal filters, Millipore, Billerica, MA, USA). The filtrate was kept on ice until analysis.

Processed samples were separated using a 100 x 4.6 mm Kinetex (C18 5 µm 100 Å,

Phenomenex) column on a reverse-phase HPLC system with a BAS LC-4B

electrochemical detector with dual 3mm carbon electrode (MF-1000) and reference

electrode (MF-2021) as previously described[69]. An external standard curve of all analytes

was run each day. Trp quantification was performed using a mobile phase consisting of

8% acetonitrile (v/v), 0.05 M citric acid, 0.05 M Na2HPO4•H2O, and 0.1 mM EDTA. No

correction for pH was needed. The detector was set to 0.875 V versus Ag/AgCl reference

electrode, sensitivity at 20 nA, and a flow rate of 1.0 ml/min. 5-HT and 5-HIAA were

separated with a mobile phase consisting of 18% methanol (v/v), 0.1M sodium phosphate,

0.8 mM octanesulfonic acid (anhydrous), and 0.1 mM EDTA (final pH adjusted to 3.1).

The detector was set to 0.70 V, sensitivity at 20 nA, and a flow rate of 1.0 ml/min.

Limulus amebocyte lysate (LAL) assay for endotoxin detection: Fetal placenta and brain

tissue from e14.5 mLFD and mHFD offspring were collected in a sterile environment and

frozen in the vapor phase above liquid nitrogen before being transferred to 2mL

Eppendorf tubes and being stored at -80°C to prevent endotoxin leaching into the tubes.

The day before sampling for endotoxin, the samples were rapidly transferred to glass

Dounce homogenizers and homogenized in 750µL of endotoxin free water (Lonza W50-

100). Homogenized samples were then diluted 1:5 in endotoxin free water in pyrogen-

free glass tubes (Lonza N207) and tested for endotoxin in duplicate using a Kinetic-

QCL™ LAL Assay (Lonza, 50-650U) according to manufacturer's instructions and

including a third 'spiked' sample from each tissue (5EU/mL endotoxin added) to ensure

acceptable recovery from every sample. Plates were run on a kinetic SPECTROstar Nano

(BMG Labtech) plate reader and analyzed with MARS data analysis software (BMG

Labtech) with a baseline correction of ΔO.D.=0.2[70]. Samples in which the 'spiked' sample

fell outside the acceptable recovery range were excluded (two samples in total).

Behavior: Behaviors are presented sex-stratified as they are often sex-dependent in their

presentation[71,72].

Ultrasonic Vocalizations (USVs): At P8, pups were brought in their home cage (with the

dam being the only adult in the cage) to the testing room. Each pup was removed from

their home cage and placed directly in a small cup in an insulated chamber with an AviSoft

Condenser ultrasound microphone (Avisoft-Bioacoustics CM16/CMPA) suspended four

inches above the contained pup. The dam and any remaining littermates were removed

from the testing room in the home cage. After 3 minutes of recording, pups were weighed

and sexed before being returned to their home cage. After the dam attended to each

returned pup, the home cage was returned to the colony room. USV analysis was done

with Avisoft and MUPET[23] (Mouse Ultrasonic Vocalization Extraction Tool). Default

MUPET parameters were used with the exception of: noise-reduction = 8.8, minimum-

syllable-duration = 2.0, as we have previously optimized[39]. 80-unit repertoires were

generated for each dataset (e.g. all male mLFD calls from one cohort were combined to

generate ranked syllable units by frequency of occurrence) and manually inspected

before USV similarity was determined using MUPET. USV similarity represents an overall

Pearson's rank order comparison of individual repertoire units (RUs) across datasets

based on their spectral shape[23].

Open Field Test: Adult offspring were placed in a 50cm x 50cm square enclosure with

38cm high walls and allowed to explore freely for ten minutes[73]. Behaviors were recorded

and analyzed using EthoVision video tracking software (Noldus). Center-avoidance was

assessed by comparing time spent at the periphery to time spent centrally/number of

central entries (middle third of chamber).

Sociability/Social Novelty Preference: Mice were assessed for social preference or social

novelty preference using a classic 3-chambered social preference test[74]. One day prior

to testing, subject and stimulus animals were habituated to the testing room for a minimum

of 1 hour before being individually habituated to the 3-chambered social preference boxes

for 5 minutes each. On test day, subject and stimulus animals were habituated to the testing room for a minimum of one hour before testing. Juvenile offspring were tested for preference to interact with either a novel age- and sex-matched conspecific or an inanimate novel object (rubber duck). Mice were placed in the middle chamber with the inanimate object confined in a clear plexiglass cylindrical cage (plexiglass rods for thorough cleaning) on one side of the test and a novel conspecific confined in an identical cage on the other side for 5 minutes. The inanimate object was then replaced with a new novel age- and sex-matched conspecific and experimental animals were allowed to freely investigate either the new animal or the familiar animal. The entire 10-minute test was recorded with a Logitech webcam, and all behavior was manually quantified using Solomon Coder by an observer blind to sex/treatment. The time spent, latency to first instance, and frequency of each of the following behaviors was coded: in chamber, investigating (nose poking into the cage), and climbing (front paws on cup and back fully extended/arched or on top of the cage). A mouse was considered "in" a chamber after its head crossed the threshold from one chamber to the next. Mice that spent >20% of the test climbing were excluded from analysis. Mice were also excluded from analysis if they did not enter any one of the three chambers, or if the latency to enter any one chamber was greater than 150s (half the test). Social preference is represented by a sociability score significantly higher than chance (50%). Sociability score formula: (time investigating social stimulus/ (total investigation time (social + object))*100). Social Novelty Preference score formula: (time investigating novel social stimulus/ (time investigating novel + familiar social stimuli))*100).

Sucrose Preference: Adult offspring were tested for their preference for a 1% sucrose

solution in water or plain water. Mice were singly housed and provided with 2 drinking

bottles and extra enrichment (toy block or bone) for a total of 6 days. After 3 days

(acclimation to 2 bottles with drinking water in both), one bottle was filled with a 1%

sucrose solution while the other was filled with drinking water. Sucrose and water intake

were measured daily (by weight) over the next 3 days. The positions of the bottles were

switched daily to reduce any bias in side-preference. Sucrose preference is calculated as

((sucrose intake)/(sucrose + water intake))*100) averaged over 3 days.

Forced Swim Test: Adult offspring were assessed for mobility behavior in a forced-swim

test (FST). Individual mice were placed in a cylindrical container (20cm in diameter x

50cm high) filled 2/3 with water (26°C-30°C) for 7 minutes. Mice were monitored via

webcam for the testing period, and mice were removed from the test early (and not used

for analysis) if their head dipped fully below the water at any point. After the test period,

animals were moved to an enclosure with a small towel and heating pad to dry. Time

spent time immobile (not swimming or climbing) was quantified using Solomon Coder by

an observer blind to sex/treatment. Velocity and distance moved were quantified by

EthoVision XT tracking software (Noldus).

qRT-PCR: RNA was isolated using TRIzol RNA extraction reagent chloroform extraction.

Either 1µg (placenta) or 200-250ng (isolated microglia) RNA was reverse-transcribed into

cDNA using a Qiagen QuantiTect Reverse Transcription kit (205311). qPCR was

performed on an Eppendorf Realplex ep Mastercycler using Sybr/Rox amplification with

a QuantiFAST PCR kit (204056). Primer sequences are listed in Supplementary Data

Table 2. Fold change was calculated using the $2^{(-\Delta\Delta CT)}$ method with 18S as an

endogenous control. When transcripts were not detected a CT value of 40 (max cycle

number) was assigned to that sample. *Trem2* was largely undetected in e14.5 microglia

(~40% of samples equally between male and female mLFD and mHFD microglia) and is

thus not represented in Extended Data Fig 7b).

Sequencing: RNA from human tissue was prepared for sequencing using an Illumina

TrueSeq RNA Exome kit and sequenced on an Illumina NovaSeq 6000 configured for a

S2 flow cell at 50bp PE. Brain and placenta count matrices were processed separately.

Reads were aligned to the human genome using Bowtie2[75]. Transcripts per million (TPM)

was calculated on the raw counts and used for all subsequent analyses. Gene inclusion

was determined using EgeR[76] (threshold of 1). Pearson's correlation coefficients and p-

values were calculated for each gene (TPM correlated with log10(maternal triglyceride

accumulation)), and data was plotted in a volcano plot using custom R script. All code for

processing is currently hosted and publicly available

(https://github.com/bendevlin18/human-fetal-RNASeq). Gene Ontology (GO) enrichment

analysis (Metascape[77]) was performed on genes with significant positive and negative

correlations separately (p<0.05).

All data is publicly available (GEO Accession number: GSE188872).

Statistics: Statistical tests, excluding RNA-sequencing analyses, were performed using

Graphpad Prism 9. Raw data as well as a description of the tests and results (including

multiple comparison corrections and post-hoc analyses) are provided. Unless otherwise

noted, data are mean ± s.e.m except for box plots (whiskers are min. to max., hinges of

boxes are 25th and 75th percentiles, middle line is the median). All analyses were

performed by an individual blind to animal sex, condition (maternal diet), and/or genotype.

In addition to correlation analyses in human tissues (see Fig 4e), we additionally ran a

global ANCOVA to address the variance between the linear regressions by sex and tissue

type. ANCOVA analysis confirms that the correlation between male fetal brain 5-HT levels

and maternal triglyceride accumulation is significantly different than the correlations

between male placental 5-HT levels and maternal triglyceride accumulation. It is also

different than the correlations between female 5-HT levels and maternal triglyceride

accumulation.

Acknowledgements: We thank the Duke University School of Medicine for the use of

the Sequencing and Genomic Technologies Shared Resource, which provided library

preparation and sequencing service. We also thank Kristina Sakers for guidance and R

scripts for sequencing analysis.

711
712   **Funding**: Research reported in this publication was supported by the Eunice Kennedy

713   Shriver National Institute of Child Health & Human Development (F32HD104430 to

714   A.M.C.), the National Institute of Environmental Health Sciences (R01 ES025549 to

715   S.D.B.), the Robert and Donna Landreth Family Foundation, and the Charles Lafitte

716   Foundation.

717
718   **Author Contributions**: S.D.B, L.S., and J.B. conceived the study and, together with

719   A.M.C, designed the experiments. A.M.C, B.A.D, J.B., L.A.G., YC.J, C.H., B.P., K.W.,

720   C.L.S., F.J., A.B.C-S, and E.R.L. performed experiments and data analysis. A.M.C. wrote

721   the manuscript with contributions from all of the authors.

722
723   The authors declare no competing interests.

Supplementary Information is available for this paper.

Supplementary Table 1. Raw GO enrichment analysis results

Figure1-4_RawData_Stats. Raw data and detailed statistical information for Figures 1

through 4

Extended_Data_Fig1-9_RawData_Stats. Raw data and detailed statistical information

for Extended Data Figures 1 through 9

Supplementary Table 2. PCR Primer sequences

*Correspondence and requests for materials should be addressed to Staci D. Bilbo

(staci.bilbo@duke.edu)

1. Vickers, M. H., Breier, B. H., Cutfield, W. S., Hofman, P. L. & Gluckman, P. D. Fetal origins of

hyperphagia, obesity, and hypertension and postnatal amplification by hypercaloric nutrition. *Am J*

Physiol Endocrinol Metab **279**, E83-7 (2000).

2. Edlow, A. G. Maternal obesity and neurodevelopmental and psychiatric disorders in offspring. *Prenat*

Diagn **37**, 95–110 (2017).

3. Kott, J. & Brummelte, S. Trick or treat? Evaluating contributing factors and sex-differences for

developmental effects of maternal depression and its treatment. *Horm Behav* **111**, 31–45 (2019).

4. Krakowiak, P. *et al.* Maternal metabolic conditions and risk for autism and other neurodevelopmental

disorders. *Pediatrics* **129**, e1121-8 (2012).

5. Mina, T. H. *et al.* Prenatal exposure to very severe maternal obesity is associated with adverse

neuropsychiatric outcomes in children. *Psychological Medicine* **47**, 353–362 (2017).

6. Rivera, H., Christiansen, K. & Sullivan, E. The role of maternal obesity in the risk of neuropsychiatric

disorders. *Frontiers in Neuroscience* **9**, (2015).

7. Rodriguez, A. Maternal pre-pregnancy obesity and risk for inattention and negative emotionality in

children. *Journal of Child Psychology and Psychiatry* **51**, 134–143 (2010).

8. Robinson, M. *et al.* Pre-pregnancy maternal overweight and obesity increase the risk for affective

disorders in offspring. *Journal of Developmental Origins of Health and Disease* **4**, 42–48 (2013).

9. Deputy, N., Dub, B. & Sharma, A. J. Prevalence and Trends in Prepregnancy Normal Weight — 48

States, New York City, and District of Columbia, 2011–2015. *MMWR Morb Mortal Wkly Rep* **66**, 1402–

1407 (2019).

10.   Hariri, N., Gougeon, R. & Thibault, L. A highly saturated fat-rich diet is more obesogenic than

diets with lower saturated fat content. *Nutr Res* **30**, 632–643 (2010).

11.   Thompson, J. R. *et al.* Exposure to a High-Fat Diet during Early Development Programs Behavior

and Impairs the Central Serotonergic System in Juvenile Non-Human Primates. *Front Endocrinol*

(Lausanne) **8**, (2017).

12.   Peleg-Raibstein, D., Luca, E. & Wolfrum, C. Maternal high-fat diet in mice programs emotional

behavior in adulthood. *Behavioural Brain Research* **233**, 398–404 (2012).

13.   Bilbo, S. D. & Tsang, V. Enduring consequences of maternal obesity for brain inflammation and

behavior of offspring. *The FASEB Journal* **24**, 2104–2115 (2010).

14.   Altemus, M., Sarvaiya, N. & Epperson, C. N. Sex differences in anxiety and depression clinical

perspectives. *Front Neuroendocrinol* **35**, 320–30 (2014).

15.   McLean, C. P., Asnaani, A., Litz, B. T. & Hofmann, S. G. Gender Differences in Anxiety Disorders:

Prevalence, Course of Illness, Comorbidity and Burden of Illness. *J Psychiatr Res* **45**, 1027–35 (2011).

16.   Cutler, G. J., Flood, A., Hannan, P. & Neumark-Sztainer, D. Major Patterns of Dietary Intake in

Adolescents and Their Stability over Time. *The Journal of Nutrition* **139**, 323–328 (2009).

17.   Li, L. *et al.* Fast food consumption among young adolescents aged 12–15 years in 54 low- and

middle-income countries. *Glob Health Action* **13**, 1795438.

18.   Fryar, C. D. & Ogden, C. L. Fast Food Intake Among Children and Adolescents in the United

States, 2015–2018. 8 (2020).

19.   Cruz, F., Ramos, E., Lopes, C. & Araújo, J. Tracking of food and nutrient intake from adolescence

into early adulthood. *Nutrition* **55–56**, 84–90 (2018).

20.   Hu, T. *et al.* Higher Diet Quality in Adolescence and Dietary Improvements Are Related to Less

Weight Gain During the Transition From Adolescence to Adulthood. *The Journal of Pediatrics* **178**, 188-

193.e3 (2016).

21.   Scattoni, M. L., Crawley, J. & Ricceri, L. Ultrasonic vocalizations: a tool for behavioural

phenotyping of mouse models of neurodevelopmental disorders. *Neurosci Biobehav Rev* **33**, 508–515

(2009).

22.   Yin, X. *et al.* Maternal Deprivation Influences Pup Ultrasonic Vocalizations of C57BL/6J Mice.

PLoS One **11**, (2016).

23.   Van Segbroeck, M., Knoll, A. T., Levitt, P. & Narayanan, S. MUPET-Mouse Ultrasonic Profile

ExTraction: A Signal Processing Tool for Rapid and Unsupervised Analysis of Ultrasonic Vocalizations.

Neuron **94**, 465-485.e5 (2017).

24.   Mun, H.-S., Lipina, T. V. & Roder, J. C. Ultrasonic Vocalizations in Mice During Exploratory

Behavior are Context-Dependent. *Frontiers in Behavioral Neuroscience* **9**, 316 (2015).

25.   Mosienko, V., Beis, D., Alenina, N. & Wöhr, M. Reduced isolation-induced pup ultrasonic

communication in mouse pups lacking brain serotonin. in *Mol Autism* vol. 6 (2015).

26.   Sachs, B. D., Ni, J. R. & Caron, M. Sex differences in response to chronic mild stress and

congenital serotonin deficiency. *Psychoneuroendocrinology* **40**, 123–9 (2014).

27.   Angoa-Pérez, M. *et al.* Mice Genetically Depleted of Brain Serotonin Do Not Display a

Depression-like Behavioral Phenotype. *ACS Chem. Neurosci.* **5**, 908–919 (2014).

28.   Savelieva, K. V. *et al.* Genetic Disruption of Both Tryptophan Hydroxylase Genes Dramatically

Reduces Serotonin and Affects Behavior in Models Sensitive to Antidepressants. *PLoS One* **3**, e3301

(2008).

29.   Osipova, D. V., Kulikov, A. V. & Popova, N. K. C1473G polymorphism in mouse tph2 gene is

linked to tryptophan hydroxylase-2 activity in the brain, intermale aggression, and depressive-like

behavior in the forced swim test. *Journal of Neuroscience Research* **87**, 1168–1174 (2009).

30.   Bonnin, A. *et al.* A transient placental source of serotonin for the fetal forebrain. *Nature* **472**,

347–350 (2011).

31.   Song, L. *et al.* Prenatal high-fat diet alters placental morphology, nutrient transporter

expression, and mtorc1 signaling in rat. *Obesity (Silver Spring)* **25**, 909–919 (2017).

32.   Goeden, N. *et al.* Maternal Inflammation Disrupts Fetal Neurodevelopment via Increased

Placental Output of Serotonin to the Fetal Brain. *J. Neurosci.* **36**, 6041–6049 (2016).

33.   Edlow, A. G. *et al.* Placental Macrophages: A Window Into Fetal Microglial Function in Maternal

Obesity. *International Journal of Developmental Neuroscience* S0736574818302508 (2018)

doi:10.1016/j.ijdevneu.2018.11.004.

34.   Roberts, K. A. *et al.* Placental structure and inflammation in pregnancies associated with obesity.

Placenta **32**, 247–54 (2011).

35.   Etienne, F. *et al.* Two-photon Imaging of Microglial Processes' Attraction Toward ATP or

Serotonin in Acute Brain Slices. *JoVE (Journal of Visualized Experiments)* e58788 (2019)

doi:10.3791/58788.

36.   Turkin, A., Tuchina, O. & Klempin, F. Microglia Function on Precursor Cells in the Adult

Hippocampus and Their Responsiveness to Serotonin Signaling. *Frontiers in Cell and Developmental*

Biology **9**, (2021).

37.   Vetreno, R. P., Patel, Y., Patel, U., Walter, T. J. & Crews, F. T. Adolescent intermittent ethanol

reduces serotonin expression in the adult raphe nucleus and upregulates innate immune expression that

is prevented by exercise. *Brain, Behavior, and Immunity* **60**, 333–345 (2017).

38.   Cunningham, C. L., Martinez-Cerdeno, V. & Noctor, S. C. Microglia regulate the number of neural

precursor cells in the developing cerebral cortex. *J Neurosci* **33**, 4216–33 (2013).

39.   Block, C. L. *et al. Prenatal Environmental Stressors Impair Postnatal Microglia Function and Adult*

Behavior in Males. 2020.10.15.336669 https://www.biorxiv.org/content/10.1101/2020.10.15.336669v2

(2020) doi:10.1101/2020.10.15.336669.

40.   Kopec, A. M., Smith, C. J., Ayre, N. R., Sweat, S. C. & Bilbo, S. D. Microglial dopamine receptor

elimination defines sex-specific nucleus accumbens development and social behavior in adolescent rats.

Nature Communications **9**, 1–16 (2018).

41.   VanRyzin, J. W. *et al.* Microglial Phagocytosis of Newborn Cells Is Induced by Endocannabinoids

and Sculpts Sex Differences in Juvenile Rat Social Play. *Neuron* **102**, 435-449.e6 (2019).

42.   Kolodziejczak, M. *et al.* Serotonin Modulates Developmental Microglia via 5-HT2B Receptors:

Potential Implication during Synaptic Refinement of Retinogeniculate Projections. *ACS Chem Neurosci* **6**,

1219–1230 (2015).

43.   Krabbe, G. *et al.* Activation of serotonin receptors promotes microglial injury-induced motility

but attenuates phagocytic activity. *Brain Behav Immun* **26**, 419–28 (2012).

44.   Béchade, C. *et al.* The serotonin 2B receptor is required in neonatal microglia to limit

neuroinflammation and sickness behavior in adulthood. *Glia* **69**, 638–654 (2021).

45.   Baković, P. *et al.* Differential Serotonin Uptake Mechanisms at the Human Maternal–Fetal

Interface. *Int J Mol Sci* **22**, 7807 (2021).

46.   Kliman, H. J. *et al.* Pathway of Maternal Serotonin to the Human Embryo and Fetus.

Endocrinology **159**, 1609–1629 (2018).

47.   Medzhitov, R. Toll-like receptors and innate immunity. *Nat Rev Immunol* **1**, 135–145 (2001).

48.   Reyna, S. M. *et al.* Elevated Toll-Like Receptor 4 Expression and Signaling in Muscle From Insulin-

Resistant Subjects. *Diabetes* **57**, 2595–2602 (2008).

49.   Shi, H. *et al.* TLR4 links innate immunity and fatty acid–induced insulin resistance. *J Clin Invest*

116, 3015–3025 (2006).

50.   Milanski, M. *et al.* Saturated Fatty Acids Produce an Inflammatory Response Predominantly

through the Activation of TLR4 Signaling in Hypothalamus: Implications for the Pathogenesis of Obesity.

Journal of Neuroscience **29**, 359–370 (2009).

51.   Rivera, P. D. *et al.* Removal of microglial-specific MyD88 signaling alters dentate gyrus

doublecortin and enhances opioid addiction-like behaviors. *Brain Behav Immun* **76**, 104–115 (2019).

52.   McAlees, J. W. *et al.* Distinct Tlr4-expressing cell compartments control neutrophilic and

eosinophilic airway inflammation. *Mucosal Immunol* **8**, 863–873 (2015).

53.   Wang, Z. *et al.* Saturated fatty acids activate microglia via Toll-like receptor 4/NF-κB signalling.

Br J Nutr **107**, 229–241 (2012).

54.   Li, M., Fu, W. & Li, X.-A. Differential fatty acid profile in adipose and non-adipose tissues in obese

mice. *Int J Clin Exp Med* **3**, 303–307 (2010).

55.   Basu, S. *et al.* Pre-gravid obesity associates with increased maternal endotoxemia and metabolic

inflammation. *Obesity (Silver Spring)* **19**, 476–482 (2011).

56.   Bowser, S. M. *et al.* Serum endotoxin, gut permeability and skeletal muscle metabolic

adaptations following a short term high fat diet in humans. *Metabolism* **103**, 154041 (2020).

57.   Pendyala, S., Walker, J. M. & Holt, P. R. A High-Fat Diet Is Associated With Endotoxemia That

Originates From the Gut. *Gastroenterology* **142**, 1100-1101.e2 (2012).

58.   Brown, A. G., Maubert, M. E., Anton, L., Heiser, L. M. & Elovitz, M. A. The tracking of

lipopolysaccharide through the feto-maternal compartment and the involvement of maternal TLR4 in

inflammation-induced fetal brain injury. *American Journal of Reproductive Immunology* **82**, e13189

(2019).

59.   Hirschmugl, B. *et al.* Maternal obesity modulates intracellular lipid turnover in the human term

placenta. *Int J Obes* **41**, 317–323 (2017).

60.   Saben, J. *et al.* Maternal obesity is associated with a lipotoxic placental environment. *Placenta*

35, 171–177 (2014).

61.   Calabuig-Navarro, V. *et al.* Effect of Maternal Obesity on Placental Lipid Metabolism.

Endocrinology **158**, 2543–2555 (2017).

62.   Qiao, L. *et al.* Maternal High-Fat Feeding Increases Placental Lipoprotein Lipase Activity by

Reducing SIRT1 Expression in Mice. *Diabetes* **64**, 3111–3120 (2015).

63.   Liu, Y. *et al.* Single-cell RNA-seq reveals the diversity of trophoblast subtypes and patterns of

differentiation in the human placenta. *Cell Res* **28**, 819–832 (2018).

64.   Wadsack, C., Desoye, G. & Hiden, U. The feto-placental endothelium in pregnancy pathologies.

Wien Med Wochenschr **162**, 220–224 (2012).

65.   VanWijk, M. J., Kublickiene, K., Boer, K. & VanBavel, E. Vascular function in preeclampsia.

Cardiovascular Research **47**, 38–48 (2000).

66.   Sternberg, E. M., Trial, J. & Parker, C. W. Effect of serotonin on murine macrophages:

suppression of Ia expression by serotonin and its reversal by 5-HT2 serotonergic receptor antagonists. *J*

Immunol **137**, 276–282 (1986).

67.   Sternberg, E. M., Wedner, H. J., Leung, M. K. & Parker, C. W. Effect of serotonin (5-HT) and other

monoamines on murine macrophages: modulation of interferon-gamma induced phagocytosis. *J*

Immunol **138**, 4360–4365 (1987).

68.   Bordt, E. A., Block, C. L. ,. Petrozziello, T. ,. Sadri-Vakili, G. ,. Smith, C. J. ,. Edlow, A. G. ,. Bilbo, S.

D. Isolation of Microglia from Mouse or Human Tissue. *STAR Protocols* (2020)

doi:10.1016/j.xpro.2020.100035.

69.   Sánchez, C. L., Van Swearingen, A. E. D., Arrant, A. E., Kuhn, C. M. & Zepf, F. D. Dietary

manipulation of serotonergic and dopaminergic function in C57BL/6J mice with amino acid depletion

mixtures. *J Neural Transm (Vienna)* **121**, 153–162 (2014).

70.   Quinlan, C. & Peters, C. Endotoxin detection using a colorimetric, kinetic microplate reader assay

and integrated data analysis. 2.

71.   Cox, K. H. & Rissman, E. F. Sex differences in juvenile mouse social behavior are influenced by

sex chromosomes and social context. *Genes Brain Behav* **10**, 465–472 (2011).

72.   An, X.-L. *et al.* Strain and sex differences in anxiety-like and social behaviors in C57BL/6J and

BALB/cJ mice. *Exp Anim* **60**, 111–123 (2011).

73.   Seibenhener, M. L. & Wooten, M. C. Use of the Open Field Maze to Measure Locomotor and

Anxiety-like Behavior in Mice. in *J Vis Exp* (2015). doi:10.3791/52434.

74.   Moy, S. S. *et al.* Sociability and preference for social novelty in five inbred strains: an approach

to assess autistic-like behavior in mice. *Genes, Brain and Behavior* **3**, 287–302 (2004).

75.   Langmead, B. Aligning short sequencing reads with Bowtie. *Current protocols in bioinformatics /*

editoral board, Andreas D. Baxevanis ... [et al.] **CHAPTER**, Unit (2010).

76.   Robinson, M. D., McCarthy, D. J. & Smyth, G. K. edgeR: a Bioconductor package for differential

expression analysis of digital gene expression data. *Bioinformatics* **26**, 139–140 (2010).

77.   Zhou, Y. *et al.* Metascape provides a biologist-oriented resource for the analysis of systems-level

datasets. *Nat Commun* **10**, 1523 (2019).

918