Genomic hallmarks and therapeutic implications of G0 cell cycle arrest in cancer

Anna J. Wiecek[1], Stephen J. Cutty[2], Daniel Kornai[1], Mario Parreno-Centeno[1], Lucie E. Gourmet[1], Guidantonio Malagoli Tagliazucchi[1], Daniel H. Jacobson[1,3], Ping Zhang[4], Lingyun Xiong[4], Gareth L. Bond[5], Alexis R. Barr[2,6], Maria Secrier[1,*]

[1] UCL Genetics Institute, Department of Genetics, Evolution and Environment, University College London, London, UK

[2] Institute of Clinical Sciences, Faculty of Medicine, Imperial College London, London, UK

[3] UCL Cancer Institute, Paul O'Gorman Building, University College London, London, UK

[4] Wellcome Centre for Human Genetics, University of Oxford, Oxford, UK

[5] Institute of Cancer and Genomic Sciences, University of Birmingham, Edgbaston, Birmingham, UK

[6] Cell Cycle Control Team, MRC London Institute of Medical Sciences (LMS), London, UK

* To whom correspondence should be addressed (m.secrier@ucl.ac.uk)

Email addresses:

Anna J Wiecek - anna.wiecek.18@ucl.ac.uk

Stephen J. Cutty - s.cutty@lms.mrc.ac.uk

Daniel Kornai - daniel.kornai.18@ucl.ac.uk

Mario Parreno-Centeno - m.parreno-centeno@ucl.ac.uk

Lucie E. Gourmet - lucie.gourmet.17@ucl.ac.uk

Guidantonio Malagoli Tagliazucchi - guidantonio.mt@gmail.com

Daniel H. Jacobson - dan.jacobson.15@ucl.ac.uk

Ping Zhang - ping.zhang@well.ox.ac.uk

Lingyun Xiong - lingyun.xiong@dtc.ox.ac.uk

Gareth L. Bond - G.Bond.1@bham.ac.uk

Alexis R. Barr - a.barr@lms.mrc.ac.uk

Maria Secrier – m.secrier@ucl.ac.uk

ABSTRACT

Therapy resistance in cancer is often driven by a subpopulation of cells that are temporarily arrested in a non-proliferative G0 state, which is difficult to capture and whose mutational drivers remain largely unknown. We developed methodology to robustly identify this state from transcriptomic signals and characterised its prevalence and genomic constraints in solid primary tumours. We show that G0 arrest preferentially emerges in the context of more stable, less mutated genomes which maintain *TP53* integrity and lack the hallmarks of DNA damage repair deficiency, while presenting increased APOBEC mutagenesis. We employ machine learning to uncover novel genomic dependencies of this process and validate the role of the centrosomal gene *CEP89* as a modulator of proliferation/G0 arrest capacity. Lastly, we demonstrate that G0 arrest underlies unfavourable responses to various therapies exploiting cell cycle, kinase signalling and epigenetic mechanisms in single cell data, and propose a G0 arrest transcriptional signature that is linked with therapeutic resistance and can be used to further study and clinically track this state.

Keywords: cell cycle arrest, G0, cancer, persister cells, genomic dependencies, machine learning, data integration, bulk/single cell sequencing

BACKGROUND

Tumour proliferation is one of the main hallmarks of cancer development(1), and has been extensively studied. While most of the cells within the tumour have a high proliferative capacity, occasionally under stress conditions some cells will become arrested temporarily in the G0 phase of the cell cycle, in a reversible state often referred to as 'quiescence', 'dormancy', 'diapause-like', or a potentially irreversible state called 'senescence', where they maintain minimal basal activity(2, 3, 4, 5). It has been proposed that G0 arrest enables cells to become resistant to anti-cancer compounds that target actively dividing cells, such as chemotherapy(5, 6, 7). Moreover, a drug-tolerant 'persister' cell state represented by slow cycling, entirely quiescent or even senescent cells(4, 8, 9, 10, 11) has been observed in a variety of pre-existing or acquired resistance scenarios, also in the context of targeted therapies(12, 13). As neoplastic cells evolve, G0 arrest can also be employed as a mechanism to facilitate immune evasion(14, 15) or adaptation to new environmental niches during metastatic seeding(16, 17). In the context of disseminated tumour cells, these G0 cycle arrest states can facilitate minimal residual disease, a major cause of relapse in the clinic(18).

Although G0 arrest is a widely conserved cellular state, essential for the normal development and homeostasis of eukaryotes(2, 19), and has been extensively studied in a variety of organisms including bacteria and yeast(20, 21), its role and different facets in cancer are still poorly defined. Hampering our understanding is the fact that it represents a number of heterogeneous states(19, 22). Canonically, cells can be forced into G0 arrest through serum starvation, mitogen withdrawal or contact inhibition(19). Cells can also undergo G0 arrest spontaneously in response to cell-intrinsic factors like replication stress(23, 24, 25). This process is controlled by p53(26), which triggers the inhibition of cyclin-CDK complexes by activating p21(24). This in turn allows the assembly of the DREAM complex - a key effector responsible for repression of cell-cycle dependent gene expression(27). Min and Spencer(28) recently demonstrated a much broader systemic coordination of 198 genes underlying distinct types of G0 arrest by profiling the

transcriptomes of cells that entered this state either spontaneously or upon different stimuli.

Additionally, proliferation-G0 decisions can be impacted by oncogenic changes such as *MYC*

amplification(29) or altered p38/ERK signalling(30).

Despite these advances, the identification of G0-arrested cells within tumours presents an ongoing

challenge due to their scarcity and lack of universal, easily measurable markers for the activation

and maintenance of this state. As they are often defined by a lack of proliferative markers(31, 32),

different forms of G0 arrest such as quiescence, senescence, dormancy and (to a lesser extent)

stemness, might sometimes be used interchangeably (4, 33). Quiescent and dormant cells can

readily resume their proliferative state, senescent cells are irreversibly arrested(28) while cancer

stem cells have a high capacity for self-renewal and sit at the top of the differentiation

hierarchy(34). Even though the same cell cycle arrest programme underlies all of these states, they

are linked with distinct environmental stimuli, and drive cancer progression and therapeutic

resistance in different ways (11, 12, 35, 36). Increasing evidence from the literature points towards

the rapid adaptation of tumour cells to drug treatments being enabled by a slow dividing or a

quiescent state that persists for a short period of time before the cells start reproliferating(37).

Thus, quiescence could more frequently be encountered at the early stages of therapeutic resistance

compared to other cell cycle arrest phenotypes, although senescence and stemness are also often

discussed in this context. Biomarkers of cell cycle arrest and persistence that are sufficiently

specific and robust to be clinically useful are clearly needed.

Furthermore, our understanding of how cancer evolution is shaped by proliferation and G0 arrest

decisions is limited. The proliferative heterogeneity of cancer cell populations has been previously

described and linked with FAK/AKT1 signalling(38), but the constraints and consequences of

these cell state switches have not been systematically profiled across cancer tissues. The extent to

which G0 arrest in cancer is enacted through transcriptional or genetic control is unknown(5, 39),

and neither are the mutational processes and genomic events modulating this state. Understanding

the evolutionary triggers and molecular mechanisms that enable cancer cells to enter and maintain

G0 arrest would enable us to develop pharmacological strategies to selectively eradicate these

arrested cancer cells or prevent them from re-entering proliferative cycles.

To address these challenges, we have developed a new method to reliably quantify G0 arrest in

cancer using transcriptomic data, and employed it to characterise this phenomenon in bulk and

single cell datasets from a variety of solid tumours. We describe the spectrum of proliferation and

G0 arrest decisions in primary tumours, which reflects a range of stress adaptation mechanisms

during the course of cancer development from early to advanced disease. We identify and validate

mutational constraints for the emergence of G0 arrest, hinting at potential new therapeutic targets

that could exploit this mechanism. We also demonstrate the relevance of G0 arrest to responses to a

range of compounds targeting cell cycle, kinase signalling and epigenetic mechanisms in single cell

datasets, and propose an expression signature that could be employed to detect treatment resistance

induced by G0 arrested tumour cells.

81
RESULTS

Evaluating G0 arrest in cancer from transcriptomic data

We hypothesised that primary tumours contain varying numbers of cells temporarily or

permanently arrested in the cell cycle, which reflect evolutionary adaptations to cellular stress and

may determine their ability to overcome antiproliferative therapies. To capture this elusive

phenotype, we developed a computational framework that would allow us to quantify G0 arrest

signals in bulk and single cell sequenced cancer samples (Figure 1a). To define a signature of G0

arrest, we focused on genes that have been shown by Min and Spencer(28) to be specifically

activated or inactivated during quiescence that arises spontaneously or as a response to serum

starvation, contact inhibition, MEK inhibition or CDK4/6 inhibition. The activity of 139 of these

genes changed in a coordinated manner across all these five distinct forms of quiescence, likely

representing generic transcriptional consequences of G0 arrest. The expression levels of these

markers were used to derive a score reflecting the relative abundance of G0 arrested cells within

individual tumours (see Methods, Supplementary Table 1).

To validate this signature and select the optimal method to score G0 arrest in individual samples

amongst different enrichment/rank-based scoring methodologies(40, 41, 42, 43), we used seven

single-cell and bulk datasets(12, 44, 45, 46, 47, 48, 49) where actively proliferating and

quiescent/dormant cells had been independently isolated and sequenced (Supplementary Table 2,

Methods). We tested the performance of our signature and scoring methodology, as well as that of

other commonly used gene signatures, in distinguishing between the truly quiescent/dormant and

truly proliferating cells in these seven datasets while varying the expression cut-offs for labelling

cells as G0 arrested or proliferating based on the respective signature. A combined Z-score

approach had the highest accuracy in detecting signals of G0 arrest, with a 91% mean performance

in classifying cells as G0 arrested or cycling (Figure 1b, Supplementary Figure 1a-b). Indeed, the

individual cells that had been identified as arrested in G0 in the experiments showed a significantly

higher Z-score than the dividing cells across all datasets (Supplementary Figure 1c). Our signature

reflected an expected increase in p27 protein levels, which are elevated in during G0 arrest(50)

(Supplementary Figure 1d). It also outperformed classical cell cycle and arrest markers, such as the

expression of targets of the DREAM complex, CDK2, Ki67 and of mini-chromosome replication

maintenance (MCM) protein complex genes - which are involved in the initiation of eukaryotic

genome replication, as well as recently defined G1/S and G2/M signatures (51) (Figure 1c).

Importantly, our approach provided a good separation between G0 and proliferating samples across

a variety of cancer types and models including cancer cell lines, 3D organoid cultures, circulating

tumour cells and patient-derived xenografts (Supplementary Table 2), thereby demonstrating its

broad applicability. Furthermore, the strength of the score appeared to reflect the duration of G0

arrest (52) (Figure 1d).

We further experimentally validated our methodology in nine lung adenocarcinoma cell lines. We

estimated the fraction of G0 cells in each of these cell lines using quantitative, single-cell imaging

of phospho-Ser807/811-Rb (phospho-Rb, which labels proliferative cells[53]) and 24 hour EdU

proliferation assays (Figure 1e-h). In these assays, cells were pulse-labelled with the nucleotide

analogue, EdU, for 24h before fixation and immunostaining. Only cells which have proliferated in

the last 24h will be labelled by EdU. EdU negative cells are classed as G0. Cells that were negative

for phospho-Rb were also defined as G0, and not G1, since they have not yet passed the restriction

point (phospho-Rb negative; see Methods, Figure 1e-f). This G0 fraction was further validated in

A549 and NCI-H1944 cells where endogenous PCNA has been labelled with an mRuby

fluorophore to enable tracking of cell cycle phase lengths by live-cell imaging (Zerjatke et al[54],

Methods). By quantifying the G0/G1 length in individual cells (i.e. time taken to enter S-phase

after mitotic exit) over a 48h period, we could see that these cells were quiescent and not senescent

(or in deep quiescence), as all G0/G1 cells did eventually enter S-phase, albeit with variable timing

(Supplementary Figure 1e).

There was a remarkably good correlation between our predicted G0 arrest levels based on the

expression of these cell lines from the Cancer Cell Line Encyclopedia (CCLE) and the fraction of

G0 cells in the experiment as assessed by lack of EdU incorporation over a 24h period (EdU

incorporation only occurs during S phase) but particularly by lack of Rb phosphorylation.

Phosphorylation and inactivation of the retinoblastoma protein is often used to define the boundary

between G0 and G1, and was specifically shown to distinguish the G0 state recently by Stallaert et

al[53]. Furthermore, a G1 signature (Methods) was not associated with these experimental

measurements, suggesting our method recovers a state more similar to G0 arrest rather than a

prolonged G1 state (Figure 1g-h). The G0 arrest correlations appeared robust to random removal of

individual genes from the signature, with no single gene having an inordinate impact on the score

(Supplementary Figures 1f-h). This provided further reassurance that our Z-score based

methodology is successful in capturing G0 arrest signals from bulk tumour data.

The spectrum of G0 arrest capacity in solid primary tumours

Having established a robust framework for quantifying G0 cell cycle arrest in cancer, we next

profiled 8,005 primary tumour samples across 31 solid cancer tissues from The Cancer Genome

Atlas (TCGA). After accounting for potential confounding signals of non-cycling non-tumour cells

from the microenvironment by correcting for tumour purity (see Methods, Supplementary Figures

1i-j), we observed an entire spectrum of fast proliferating to slowly cycling tumours, with the latter

presenting stronger G0-linked signals (Figure 2a). While we acknowledge that no tumour would be

entirely quiescent/senescent and we cannot identify individual G0 arrested cells within the tumour,

this analysis does capture a broad range of phenotypes reflecting varying proliferation and cell

cycle arrest rates, which suggests that G0 arrest is employed to different extents by tumours as an

adaptive mechanism to various extrinsic and intrinsic stress factors. Cancers known to be

frequently dormant, such as glioblastoma(6, 44), were amongst the highest ranked in terms of G0

arrest levels, along with kidney and adrenocortical carcinomas (Figure 2b). This is likely explained

by the innate proliferative capacity of the respective tissues. Indeed, tissues with lower stem cell

division rates presented a greater propensity for G0 arrest (Figure 2c)(55).

Our score showed strong negative correlations with the expression of proliferation markers (Figure

2d), suggesting that it captures a cellular state that could potentially act as a baseline for all major

forms of cell cycle arrest, including quiescence, senescence, stemness and clinical dormancy.

Indeed, we found that our signature could to a certain extent also separate senescent cells from

proliferating ones in single cell data from Hernandez-Segura et al(56), which is unsurprising since

these cells are also in the G0 phase (Supplementary Figure 2a). Indeed, the authors of this resource

highlight that some of the pathways uncovered in these senescent cells may be shared with

quiescence, which is also backed up by a study from Fujimaki and Yao(57) suggesting similarities

between deep quiescence and senescence. Our score did not show strong correlations with other

markers of senescence such as the Senescence-Associated Secretory Phenotype (SASP) and β-

galactosidase activity(58, 59, 60) in single cells or TCGA samples (Figure 2d, Supplementary

Figures 2b-d), although we cannot exclude the possibility of a senescent state being captured

occasionally given that neither β-galactosidase nor the SASP are obligatory for maintaining

senescence(61). However, the underpinning programme appears to be distinct from that of cancer

stem cells, marked by signatures associated with high telomerase activity and an undifferentiated

state(62, 63) (Supplementary Figure 2e-f).

Lastly, we confirmed expected dependencies on the p53/p21/DREAM activation axis: tumours that

were proficient in *TP53* or the components of the DREAM complex, as well as those with higher

p21 expression, had elevated G0 arrest levels across numerous tissues (Figure 2e, Supplementary

Figures 2g-h), although only 8 out of 139 genes in our signature are directly transcriptionally

regulated by p53(64). Nevertheless, p53 proficiency appears to be a non-obligatory dependency of

G0 arrest, which is also observed to arise in p53 mutant scenarios in 21% of cases. p53 has also

been shown to play a role in preventing the occurrence of larger structural events and

polyploidy(65, 66, 67), potentially explaining the lower G0 arrest levels we observed in tumours

that had undergone whole genome duplication (Figure 2f).

The genomic background of G0 arrest in cancer

Cancer evolution is often driven by a variety of genomic events, ranging from single base

substitutions to larger scale copy number variation and rearrangements of genomic segments. It is

reasonable to expect that certain mutations accumulated by the cancer cells might enable a more

proliferative phenotype, impairing the ability of cells to enter G0 arrest, or – on the contrary –

might favour cell cycle exit as a temporary adaptive mechanism to extreme levels of stress. Having

obtained G0 arrest estimates for primary tumour samples, we set out to identify potential genomic

triggers or constraints that may shape proliferation versus G0 arrest decisions in cancer. We

identified 285 cancer driver genes that were preferentially altered (via mutations or copy number

alterations) either in slow cycling or fast proliferating tumours (Figure 3a). Reassuringly, this list

included genes previously implicated in driving cell cycle exit decisions such as *TP53* and

MYC(26, 29). We also investigated associations with mutagenic footprints of carcinogens (termed

"mutational signatures"), which can be identified as trinucleotide substitution patterns in the

genome(68, 69). 15 mutational signatures were linked with G0 arrest levels either within individual

cancer studies or pan-cancer (Supplementary Figure 2i).

Following the initial prioritisation of putative genomic constraints of G0 arrest, we employed

machine learning to identify those events that could best distinguish slow cycling tumours with

higher abundance of G0 arrested cells from fast proliferating ones, while accounting for tissue

effects. An ensemble elastic net selection approach similar to the one described by Pich et al(70)

was applied for this purpose (Figure 3b, Methods). Our pan-cancer model identified tissue type to

be a major determinant of G0 arrest levels (Supplementary Figure 3a). It also uncovered a reduced

set of 57 genomic events linked with proliferation/G0 arrest switches, including SNVs and copy

number losses in 17 cancer genes, as well as amplifications of 10 cancer genes (Figure 3c). These

events could then be successfully employed to predict G0 arrest in a separate test dataset, thus

internally validating our model (Supplementary Figure 3b). Thus, while these events are not

necessarily causative, the link is strong enough to be identifying G0 arrest states from genomic data

alone. Such events may also pinpoint cellular vulnerabilities that could be exploited

therapeutically.

Overall, the genomic dependencies of G0 arrest mainly comprised genes involved in cell cycle

pathways, p53 regulation and ubiquitination (most likely of cell cycle targets), and RUNX3

regulation, which have previously been shown to play a role in controlling proliferation and cell

cycle entry(71) (Supplementary Figure 3c). Invariably, this analysis has captured several events

that are well known to promote cellular proliferation in cancer: this is expected and confirms the

validity of our model. It was reassuring that a functional *TP53*, lack of *MYC* amplification and

lower mutation rates (Figure 3c) were amongst the top ranked characteristics of tumours with high

levels of G0 arrest, which also displayed less aneuploidy. However, our analysis has also

uncovered novel dependencies of G0 arrest-proliferation decisions that have not been reported

previously, such as *CEP89* and *LMNA* amplifications observed in fast cycling tumours, or *ZMYM2*

deletions prevalent in samples with high levels of G0 arrest. ZMYM2 has recently been described

as a novel binding partner of B-MYB and has been shown to be important in facilitating the G1/S

cell cycle transition(72). p16 (*CDKN2A*) deletions, one of the frequent early events during cancer

evolution(73, 74), were enriched in tumours with high proportions of cells in G0. *RB1* deletions

and amplifications were both associated with a reduction in G0 arrest, which might reflect the dual

role of RB1 in regulating proliferation and apoptosis(75).

Our model also calls to attention to the broader mutational processes associated with this cellular

state. Such processes showed fairly weak and heterogeneous correlations with G0 arrest within

individual cancer tissues (Supplementary Figure 2g), but their contribution becomes substantially

clearer pan-cancer once other genomic sources are accounted for. In particular, we identified an

association between G0 arrest and mutagenesis induced by the AID/APOBEC family of cytosine

deaminases as denoted by signature SBS2(68) (Figure 3c). As highlighted by Mas-Ponte and

Supek(76), APOBEC/AID driven mutations tend to be directed towards early-replicating, gene-rich

regions of the genome, inducing deleterious events on several genes including *ZMYM2*, which our

pan-cancer model has linked with G0 arrest.

In turn, defective DNA mismatch repair, as evidenced by signatures SBS44, SBS20, SBS15,

SBS14 and SBS6(68), was prevalent in fast cycling tumours (Figure 3c). Mismatch repair

deficiencies lead to hypermutation in a phenomenon termed "microsatellite instability" (MSI),

which has been linked with increased immune evasion(77). Cancers particularly prone to MSI

include colon, stomach and endometrial carcinomas(78), where this state was indeed linked with

reduced G0 arrest (Figure 3d). Furthermore, tumours with high proportions of cells in G0 were

depleted of alterations across all DNA damage repair pathways (Figure 3e).

Our measurements of G0 arrest also reflected expected cycling patterns across 27 MCF7

strains(79): cell lines with longer doubling times exhibited increased G0 arrest (Figure 3f). This

coincided with a depletion of *PTEN* mutations, a dependency highlighted by the pan-cancer model.

When checking for dependencies in individual cancer tissues, 24 out of the 25 genes identified by

the model were significantly associated with G0 arrest or proliferation decisions in at least one

tissue, most prominently in breast, lung and liver cancers which also represent the largest studies

within TCGA (Figure 3g, top panel). Most of these genomic insults were linked with a decrease in

G0 arrest. In external validation datasets these associations, including deletions in *PTEN* and

LRP1B or amplifications of *MYC*, *CEP89* and *ETV6*, featured most prominently in the largest

cohort of breast cancer samples (Figure 3g, bottom panel). These results highlight the fact that

although a pan-cancer approach is suited to capture genomic events that are universally associated

with cell cycle exit, certain genetic alterations may facilitate a higher or lower propensity of G0

arrest in a single tissue only.

Indeed, when building a tissue-specific breast cancer model of G0 arrest using a combined

ANOVA and random forest classification approach (Supplementary Figure 4a), we not only

recovered the associations with the *TP53, MYC*, *LMNA* and *ETV6* events already seen in the pan-

cancer model (Supplementary Figure 4b), but identified additional events which validated in the

METABRIC cohort and were also seen in several other cancers, e.g. bladder, lung and lower grade

glioma (Supplementary Figure 4c). Notably, the APOBEC mutational signature SBS2 was the

strongest genomic signal linked with G0 arrest in breast cancer (Supplementary Figures 4b,d) and

was most prevalent in Her2+ tumours, although the Luminal A subtype showed the highest levels

of G0 arrest overall, as expected given its well-known lower proliferative capacity(80)

(Supplementary Figures 4e-f).

Validation of CEP89 as a modulator of G0 arrest capacity

To gain more insight into the underlying biology of G0 arrest in cancer, we sought to

experimentally validate associations highlighted by the pan-cancer model. We focused on the

impact of *CEP89* activity on proliferation/arrest decisions due to the high ranking of this putative

oncogene in the model, the relatively unexplored links between *CEP89* and cell cycle control, as

well as its negative association with G0 arrest across a variety of cancer cell lines (Supplementary

Figures 5a-c). The function of *CEP89* is not well characterised, however, the encoded protein has

been proposed to function as a centrosomal-associated protein(81, 82). Centrosomes function as

major microtubule-organising centres in cells, playing a key role in mitotic spindle assembly(83)

and the mitotic entry checkpoint(84). Moreover, centrosomes act as sites of ubiquitin-mediated

proteolysis of cell cycle targets(85), and members of several growth signalling pathways, such as

Wnt and NF-kB, localise at these structures(86, 87). Several genetic interactions have also been

reported between *CEP89* and key cell cycle proteins, including cyclin D2(88) (Figure 4a).

Our model linked *CEP89* amplification with fast cycling tumours (Figure 3c). Centrosome

amplification is a common feature of tumours with high proliferation rates and high genomic

instability(89), and overexpression of centrosomal proteins can alter centriole structure(90, 91).

Indeed, *CEP89* amplified tumours presented elevated expression of a previously reported

centrosome amplification signature (CA20)(89) (Figure 4b), which was strongly anticorrelated

with G0 arrest levels (Figure 4c). Furthermore, *CEP89* expression was prognostic across multiple

cancer tissues (Figure 4d) and linked with toxicity of several cancer compounds in cell line models

(Supplementary Figure 5d).

We validated this target in the lung adenocarcinoma cell line NCI-H1299 showing high levels of

CEP89 amplification. Cep89 depletion via siRNA knockdown caused a consistent decrease in cell

number, in the absence of any detectable cell death, and an increase in the fraction of G0 cells as

measured by phospho-Rb and EdU assays (Figure 4e-f). Thus, we propose *CEP89* as a novel cell

proliferation regulator that may be exploited in certain scenarios to control tumour growth.

293 Characterisation of individual stress response programmes of G0 arrest

While we had previously examined a generic programme of G0 arrest, cancer cells can enter this

state due to different stimuli(19) and this may inform its aetiology and manifestation. To explore

this, we re-scored tumours based on gene expression programmes specific to serum starvation,

contact inhibition, MEK inhibition, CDK4/6 inhibition or spontaneously occurring quiescence as

defined by Min and Spencer(28) (see Methods). We observed a good correlation between our

estimates representing individual stress response programmes and the expression of genes

associated with the corresponding form of G0 arrest in the literature (Figure 5a-e, Supplementary

Figure 6, Methods). Specifically, strong inverse correlations were seen between our CDK4/6

inhibition scores and the mean expression of *CDK4* and *CDK6*, or between our MEK inhibition

scores and the expression of genes involved in the MAPK pathway(92). Spontaneous quiescence

and serum starvation scores were most correlated with the activity of p21, or of genes involved in

the cellular response to starvation, respectively. The contact inhibition programme was also

captured, but with lesser specificity.

CDK4/6 inhibition-induced G0 arrest levels were further validated using external RNA-seq

datasets from cancer cell lines and xenograft mice sequenced before and after treatment with the

CDK4/6 inhibitor Palbociclib(93, 94) (Figure 5f, Supplementary Table 3). The fact that the

estimates for a generic G0 arrest phenotype were equally or, in some cases, more discriminative of

cells treated with Palbociclib confirms the generalisability of this score, which may be more

broadly applicable to different tissues and/or model systems, as shown previously in the single cell

validation data. The CDK4/6 inhibition scores outperformed all the other stress response subtype

scores, suggesting that a combination of individual programmes and the generic score might best

identify a specific stimulus driving G0 arrest. Interestingly, we also observed significant

differences in spontaneous quiescence scores before and after treatment. Indeed, p21 activity has

been linked with the Palbociclib mechanism of action(95, 96), and this analysis suggests potential

similarities between CDK4/6 inhibition and p21-dependent G0 arrest phenotypes.

Having validated our framework for quantifying stimulus-specific G0 arrest programmes, we

proceeded to estimate the dominant form of stress that may induce cell cycle arrest in different

cancer types (Figure 5g). We found a range of G0 arrest aetiologies across most tissues, while a

minority of cancers were dominated by a single form of stress response, e.g. serum starvation in all

G0 arrested pheochromocytomas and paragangliomas, contact inhibition in 88% of head and neck

carcinomas and CDK4/6 inhibition in 80% of adrenocortical carcinomas. While we do not wish to

claim that the state of cell cycle arrest will have necessarily been induced by the actual predicted

stimulus (impossible in the case of CDK4/6 or MEK inhibition, as the analysed samples are all

treatment-naïve), we suggest that the downstream signalling cascade may resemble that triggered

by such stimuli, e.g. via CDK4/6 or MEK loss of function mutations.

Some of the differences observed might be explained by the dependency between p53 activity and

the form of stress response that is enacted. Amongst the five different forms, spontaneous G0 arrest

appeared most strongly dependent on p53 functionality, with a nearly two-fold enrichment of p53

proficient tumours in this group (Supplementary Figure 6f). Indeed, significantly higher levels of

spontaneous G0 arrest were observed in the majority of cancers (56%) when p53 was functional

rather than mutated. The second most dependent state was that of CDK4/6 inhibition, with

increased levels in 36% of cancer types displaying p53 proficiency (Supplementary Figure 6g).

Overall, these analyses of stress response states point to common transcriptional features of drug-

tolerant G0 arrested cells in different cancer settings that could be employed in designing ways to

eradicate these cells in the future.

Role of G0 arrest in driving therapeutic resistance in cancer uncovered from single cell data

Overall, G0 arrest appears to be beneficial for the long-term outcome of cancer patients, even when

accounting for potential confounders such as stage, sex and tissue (Figure 6a, Supplementary

Figure 7a). No clear relation was observed between G0 arrest levels within the primary tumour and

risk of relapse, although higher G0 arrest was occasionally deemed favourable to avoiding disease

recurrence or progression (Supplementary Figure 7b-e). Indeed, such slow cycling, indolent

tumours would have higher chances of being eradicated earlier in the disease, which is consistent

with reported worse prognosis of patients with higher tumour cell proliferation rates (97). As

expected, G0 arrest levels were increased in stage 1 tumours, although later stages also exhibited

this phenotype occasionally (Supplementary Figure 7f). However, outcomes do vary depending on

the stress source, with worse survival observed upon contact inhibition (Figure 6b). The outcomes

also vary by tissue: when the cut-offs between increased G0 arrest and high proliferation were

defined on an individual cancer basis rather than pan-cancer, we found that lung, colon or

esophageal carcinoma patients displayed significantly worse prognosis in the context of high

proportions of G0 arrested cells in the tumour (Figure 6c, Methods). Indeed, p53 wild type

colorectal cancers expressing a quiescence-linked fetal phenotype have been recently associated

with metastasis and poor prognosis(98). In contrast, adrenocortical and kidney papillary cell

carcinoma ranked in the top of cancers with improved survival. It is noticeable that the cancers in

the former, worse prognosis group are also amongst the ones displaying lower than average G0

arrest (Figure 2b), so the observed inferior outcomes could in part be linked to these cancers being

intrinsically faster progressing. It is possible there is a lower limit below which G0 arrest stops

being useful for delaying growth and becomes detrimental instead, perhaps in conjunction with

treatment. Indeed, other factors such as the type of therapy received could play a role too. While

we are limited in the investigation of such factors in TCGA due to the incomplete records

available, these discrepancies should be subject to future research.

While G0 arrest may confer an overall survival advantage in most cancers, it can also provide a

pool of cells that are capable of developing resistance to therapy(12, 99). Using our methodology,

we indeed observed an increase in G0 arrest levels in cell lines following treatment with EGFR,

BRAF and CDK4/6 inhibitors, as well as conventionally used chemotherapies such as 5-

Fluorouracil (5-FU) in multiple bulk RNA-seq datasets (Figure 6d).

Furthermore, the recent widespread availability of single-cell transcriptomics offers the opportunity

to investigate the impact of G0 arrest on such therapies with much greater granularity than is

allowed by bulk data. Using our G0 arrest signature and single-cell data from RKO and SW480

colon cancer cell lines treated with 5-FU(100), we could observe G0 arrest and proliferation

decisions following conventional chemotherapy treatment. Within the p53 proficient cell line

RKO, the fraction of G0 arrested cells increased from 41% to 93% after treatment with a low dose

(10 μM) of 5-FU and persisted at higher doses (Figure 6e-f). In contrast, a comparable increase in

G0 arrest was not observed in *TP53* mutant SW480 cells, further emphasizing the key role of p53

as a regulator of cell cycle exit (Figure 6g-h). This implies that although *TP53* mutations confer a

more aggressive tumour phenotype and may drive resistance via other mechanisms, *TP53* wild-

type tumour cells are more likely to be capable of entering a G0 "persistent" state associated with

drug resistance. SW480 cells showed higher apoptotic activity following treatment compared to

RKO cells, particularly within actively cycling cells, further corroborating that cells capable of

entering G0 arrest may be intrinsically less vulnerable to this therapy (Supplementary Figures 8a-

b).

Similarly, using single cell data from an *EGFR* mutant Non-Small Cell Lung Cancer (NSCLC) cell

line treated with the EGFR inhibitor Erlotinib(101), we predicted that 40% of cells were likely to

exist in a G0 arrest state prior to treatment. EGFR inhibition led to a massive decrease in cell

numbers immediately after treatment, mostly due to proliferating cells dying off (Supplementary

Figures 8c-d), while the proportion of arrested cells increased to 96% at day 1, indicating an

immediate selective advantage for such cells (Figure 6i-j). These cells appear to gradually start

proliferating again in the following days during continuous treatment, with the percentage of

proliferating cells approaching pre-treatment levels by day 11 (Figure 6j). The same trend captured

by our signature could be observed upon KRAS and BRAF inhibition in different cell line models

(Supplementary Figure 8e-h, Supplementary Table 3)(12, 101). Furthermore, during the first days

of treatment the NSCLC cells that survived EGFR inhibition appeared to reside in a state most

resembling that induced by serum starvation (Figures 6k-l). Both EGFR kinase inhibitors and

serum starvation have been shown to trigger autophagy(102), which may explain the convergence

between this inhibitory trigger and the type of stress response. At day 11 most of the remaining

arrested cells appeared in a state similar to that preceding the treatment (Figure 6l).

Thus, G0 arrest appears to explain resistance to broad acting chemotherapy agents as well as

targeted molecular inhibitors of the Ras/MAPK signalling pathway, being either selected for, or

induced immediately upon treatment, and gradually waning over time as cells start re-entering the

cell cycle. Using massively multiplexed chemical transcriptomic data, we also analysed responses

to 188 small molecule inhibitors in cell lines at single-cell resolution(103) (Supplementary Figure

9). We observed a large increase in G0 arrest following treatment with compounds targeting cell

cycle regulation and tyrosine kinase signalling, consistent with our previous results, but also for

compounds modulating epigenetic regulation, e.g. histone deacetylase inhibitors – thus

highlighting the broad relevance of G0 arrest.

While links between G0 arrest and therapeutic resistance are prevalently observed in cell lines, one

would question whether this translates to similar pathology in cancer patients. While we observed

significantly higher G0 arrest levels in pre-treated tumours of non-responders to neoadjuvant

chemotherapy in a breast cancer study by Hatzis et al[104] (Supplementary Figure 10a), surveying

various targeted therapy datasets from the SELECT study[105] and the TCGA data for links with

response to various therapies (single agent and combinations) showed little to no evidence for a

bulk signature of G0 being useful for _predicting_ resistance in these studies (Supplementary Figures

10b-c). Although the studies available for inspection are rather sparse, evidence from all the

analyses presented here suggests there is no universal a priori role that G0 arrest has within the pre-

treated primary tumour in determining response to treatment, with more favourable overall

outcomes observed occasionally due to slower progressing malignancy, but also resistance

observed in the case of chemotherapy in breast cancer. Instead, the role of G0 arrest in enabling

therapeutic resistance as a short-lived acquired phenotype as demonstrated in single cell datasets

appears more consistent.

Tumour cell G0 arrest signature for use in single-cell transcriptomics data

Our ability to probe the nature of G0 arrest phenotypes in single cell RNA-seq data using a defined

G0 signature could aid the development of methods to selectively target G0 arrested drug-resistant

persister cells. However, a major challenge of single cell RNA-seq data analysis is the high

percentage of gene dropout, which could impact our ability to evaluate G0 arrest using the full 139

gene signature. The single cell RNA-seq datasets we analysed exhibited an average drop-out of 8.5

genes out of the full gene signature. While our scoring method remains robust to such levels of

dropout (Supplementary Figures 1d-f), we also employed machine learning to reduce our initial list

of 139 markers of quiescence to a robust 35-gene signature, comprised mainly of RNA metabolism

and splicing-regulating factors, but also of genes involved in cell cycle progression, ageing and

senescence, which could be applied to sparser datasets with larger levels of gene dropout (Methods,

Figure 7a-b, Supplementary Table 4). The optimised signature of G0 arrest performed similarly to

the initial broadly defined programme in distinguishing fast cycling tumours from those containing

high proportions of G0 cells (Figure 7c), it showed an average dropout of only 0.5 genes across the

single cell RNA-seq datasets used in this study (Figure 7d), was similarly prognostic (p=0.004) and

showed comparable profiles of resistance to treatment (Figure 7e, Supplementary Figure 11). This

minimal expression signature could be employed to track and further study emerging G0 arrest-

enabled resistance in a variety of therapeutic scenarios.

DISCUSSION

Despite its crucial role in cancer progression and resistance to therapies, G0 arrest in all its forms

remains poorly characterised due to the scarcity of suitable models and biomarkers for large-scale

tracking in the tissue or blood. The lack of proliferative markers such as Ki67 or CDK2(31, 106)

does not uniquely distinguish G0 arrest from other cell cycle phases, e.g. G1. Miller et al(32) have

shown that the Ki67 is expressed at the mRNA level but the protein is degraded continuously both

in G0 and G1, and it rather acts as a graded marker of S/G2/M. Similarly, CDK2 activity is low in

G0, G1, builds up at the restriction point, is high in the S phase and is then replaced by CDK1 in

mitosis. Reduced CDK2 expression can manifest due to quiescence, mitosis, but also to DNA

damage(107), and thus cut-offs to uniquely distinguish its activity in G0 would be difficult to

define. Furthermore, these and other reliable markers of G0 arrest such as p27 or p130(50) are best

captured at protein level, which is much more sparsely measured, and expression does not

accurately reflect their activity. This study overcame this limitation by employing genes active in

different forms of quiescence whose patterns of expression are distinct from markers of a longer

G1 phase, and capture cell cycle arrest as also observed in senescence, stemness or dormancy. We

have extensively validated our method and signature in single cell datasets and cancer cell lines,

and have demonstrated that it can reliably and robustly capture signals of G0 arrest both in bulk

tissue as well as in single cells.

Within bulk tissue, we are limited in our capacity to distinguish between large fractions of cells

residing in short-lived G0 arrest and a smaller fraction of cells that are in deeper G0 arrest, as our

score seems to reflect both parameters to a certain extent. Bearing in mind this limitation, our score

could potentially also be used in a single cell setting to capture longer-lived cell cycle arrest states

such as ones demonstrated in senescence or dormancy, and could assist in identifying such states,

but only with the help of additional cell-state specific, immune or secretory biomarkers. Indeed,

gene activity linked to cell cycle arrest is not exclusive to quiescence, but can be shared with

senescence or dormancy in certain scenarios, as also demonstrated in some of our analyses of

senescent cells. This makes it difficult to clearly distinguish states like dormancy, senescence,

quiescence (particulary deep quiescence), as even their definitions can be contentious at times both

in the context of human cancers (4, 57, 108, 109, 110) as well as in physiological conditions in

other organisms (111, 112). Since our signature was derived and validated in experiments that were

tailored specifically to induce and/or measure quiescence, we believe the signature proposed in this

study best reflects a quiescent-like, reversible G0 cell cycle arrest state. While senescent and

dormant cells could be distinguished from their quiescent counterparts simply based on additional

senescence and dormancy markers, further research is nevertheless required in the future to

delineate signatures that are both necessary and sufficient to unambiguously discriminate all three

states. In the meantime, future studies utilising our G0 signature should also test for such additional

markers like β-galactosidase activity, the SASP and other senescence markers, or *NRF2*, *NR2F1*,

SOX9, RARβ(113, 114) and other dormancy markers to ascertain the type of G0 arrest that is being

captured.

The versatility of our signature is evidenced by high classification accuracies across a variety of

solid cancer datasets. More variable performance was observed when applied to hematopoietic

stem cells as it was not designed to capture signals in this context (Supplementary Figure 1b).

While we cannot exclude that the patterns captured may also occasionally reflect cell cycle arrest

in G1 or G2, this broad signature would still capture phenotypes resulting from intrinsic or

extrinsic cellular stress that reflect temporary tumour adaptation during the course of cancer

evolution or upon treatment with drugs. Thus, studying such states is relevant for identifying

vulnerabilities that could be exploited at different time points during the course of cancer

treatment.

We show that G0 arrest is pervasive across different solid cancers and generally associated with

more stable, less mutated genomes with intact DNA damage repair pathways. We also find a link

between APOBEC mutagenesis and higher levels of G0 arrest. Some neoplastic events enriched in

tumours with increased G0 arrest, such as p16 or *ZMYM2* deletions, could mark elevated genomic

stress that renders cells more prone to cell cycle exit. We also identified mutational events affecting

a variety of genes such as *PTEN*, *CEP89, CYLD*, *LMNA* that appear unfavourable to cell cycle

arrest, thus potentially implicating them in influencing G0 arrest-proliferation decisions. Among

these, we propose and validate *CEP89* as a novel modulator of G0 arrest capacity in non-small cell

lung cancer. A recent paper describes how increased CEP89 copy number and expression

correlates with a worse prognosis in ovarian cancer(115), which we hypothesise could be linked to

Cep89's role in modulating G0 arrest. Although we do not yet know how Cep89 regulates G0

arrest, two roles have been ascribed to Cep89 which could be significant. First, Cep89 is required

for primary cilium assembly(116, 117). The primary cilium acts as a signalling hub, transducing

extracellular signals to intracellular signalling networks, many of which regulate growth and

proliferation(118). Cep89 deficiency also leads to defects in Complex IV assembly in the electron

transport chain in mitochondria, leading to decreased mitochondrial function and ATP

production(119). Decreased ATP would impair the ability of cells to proliferate. Since Cep89 is a

coiled-coil protein with no obviously targetable regulatory domains, it will be important to

ascertain which Cep89 function is key to regulating the balance between proliferation and arrest in

cancer cells to be able to potentially target that process, rather than Cep89 itself, to induce or

maintain G0 arrest.

These large-scale genomic associations with G0 arrest phenotypes are only currently feasible in

bulk datasets. However, bulk sequenced data has a major limitation in capturing an average signal

across all cells within the tumour, which prevents individual cell state identification and counting.

Our subsequent exploration of single cell datasets across 193 therapeutic scenarios complements

this analysis and illustrates the power of applying our signature in single cells.

Our signature of G0 arrest is prognostic and marks primary tumours with a lower proliferative

capacity before treatment, but we also clearly demonstrate that it can be employed to track

resistance to multiple cell cycle, kinase signalling and epigenetic targeting regimens, where it often

appears as a short-lived phenotype. While this discrepancy may appear incompatible at first glance,

it is not unlike other cellular processes that have been shown to present dual roles in a cancer

setting, such as reactive oxygen species(120), but also p38α(121) or NRF2(122), both of which

have been implicated in quiescence or dormancy(114, 123). It is possible that there is a tipping

point between G0 arrest acting beneficially or detrimentally during tumour development and

treatment. Furthermore, this is likely influenced by a myriad of other complex factors that we have

not had the chance to analyse in depth here, and in some cases it may just be the baseline for

acquiring cancer cell stemness or senescent properties. While we acknowledge this conundrum

requires further study, we believe this phenotype also offers a unique opportunity to further

understand mechanisms of tumour resistance. A key open question remains: if G0 arrest drives

resistance, does it do so in a Darwinian fashion, as a pre-existing population that is selected for

upon drug treatment, or is it instead an acquired phenotype? Our single cell analyses cannot

exclude either scenario. Given the variable links to treatment response and lack of clear evidence

for relapse when surveying G0 arrest in primary tumours *before* treatment, it is likely our G0 arrest

signature in its current form cannot be used to predict resistance to chemotherapy or targeted

therapy, and we would not recommend it for this purpose unless further validated in a specific

cancer setting. We have also not inspected the role of G0 arrest in the context of immunotherapy,

which remains an area of future study. However, we believe our signature has high value for the

study of emerging resistance in an *in vitro*/*in vivo* setting, as a short-lived enabler of drug tolerance.

The optimised signature we propose for single cell data makes it tractable to a variety of future

studies in this area.

In a treatment setting, vulnerabilities of G0 arrested cells could be exploited for combination

therapies. Cells which have exited the cell cycle utilise several mechanisms to achieve drug

resistance, including upregulation of stress-induced pathways such as anti-apoptotic BCL-2

signalling(124), anti-ROS programmes(28) or immune evasion(15). Further studies are needed to

elucidate which of these mechanisms are specifically employed on a case-by-case basis.

Our findings contribute to the understanding of the aetiology and genetic context of G0 arrest in

cancer. This is particularly relevant to identifying new anti-proliferative targets, but also for the

detection and eradication of drug tolerant persister cells, which have been frequently, although not

always, observed to be slow cycling or entirely quiescent/senescent(8, 9). Importantly, the state of

G0 arrest that we have studied here is distinct from that of disseminated tumour cells causing

clinical dormancy and cancer relapse, often after many years from the treatment of the primary

tumour(4, 125). Here, we have focused on understanding how tumours make proliferation and G0

arrest decisions during the earlier stages of cancer development, within the treatment-naïve primary

tumour and as an immediate response to anti-cancer therapies. However, since the dormancy of

disseminated tumour cells is fundamentally enabled through a long but temporary cell cycle arrest,

we believe our findings of the fundamental processes linked with G0 arrest could in the future help

inform a better characterisation of dormant tumour cells when combined with specific

microenvironmental signatures that are critical for enabling that process.

CONCLUSIONS

Overall, our study provides, for the first time, a pan-cancer view of G0 arrest and its evolutionary

constraints, underlying novel mutational dependencies which could be exploited in the clinic. We

propose a G0 arrest signature which can be robustly measured in bulk tissue or single cells and

could potentially inform therapeutic strategies in the longer term. This signature could be assessed

in the clinic to track rapidly emerging resistance, e.g. through liquid biopsies or targeted gene

panels. We hope these insights can be used as building blocks for future studies into the different

regulators of G0 arrest, including epigenetics and microenvironmental interactions, as well as the

mechanisms by which it enables therapeutic resistance both in solid and haematological

malignancies.

METHODS

Selection of G0 arrest marker genes

Generic G0 arrest markers:

Differential expression analysis results comparing cycling immortalised, non-transformed human

epithelial cells and cells in five different forms of quiescence (spontaneous quiescence, contact

inhibition, serum starvation, CDK4/6 inhibition and MEK inhibition) were obtained from Min and

Spencer(28). A total of 195 genes were differentially expressed in all five forms of quiescence

under an adjusted p-value cut-off of 0.05. This gene list, reflective of a generic G0 arrest

phenotype, was subjected to the following refinement and filtering steps: (1) selection of genes

with a unidirectional change of expression across all five forms of quiescence; (2) removal of

genes involved in other cell cycle stages included in the "KEGG_CELL_CYCLE" gene list

deposited at MSigDB; (3) removal of genes showing low standard deviation and low levels of

expression within the TCGA dataset, or which showed low correlation with the pan-cancer

expression of the transcriptional targets of the DREAM complex, the main effector of quiescence,

in TCGA. The resulting 139-gene signature is presented in Supplementary Table 1.

G0 arrest stress response-specific markers:

Gene lists representing spontaneous quiescence, contact inhibition, serum starvation, CDK4/6

inhibition and MEK inhibition programmes were obtained using genes differentially expressed in

each individual quiescence form using an adjusted p-value cutoff of 0.05. The gene lists were

subjected to filtering steps 2 and 3 described above. Following the refinement steps, 10 upregulated

and 10 downregulated genes with highest log2 fold changes were selected for each stress response

type.

Quantification of G0 arrest in tumours

The *GSVA* R package was used to implement the combined z-score(40), ssGSEA(41) and

GSVA(42) gene set enrichment methods. For the above three methods a separate score was

obtained for genes upregulated in quiescence and genes downregulated in quiescence, following

which a final G0 arrest score was obtained by subtracting the two scores. The singscore single-

sample gene signature scoring method(43) was implemented using the *singscore* R package. In

addition to these, we also calculated a mean scaled G0 arrest score based on the refined list of

genes upregulated and downregulated in quiescence, as well as a curated housekeeping genes from

the "HSIAO_HOUSEKEEPING_GENES" list deposited at MSigDB, as follows:

$$\text{G0}m = \frac{\frac{1}{n}\sum G_U - \frac{1}{n}\sum G_D}{\frac{1}{n}\sum G_H}$$

G0m = mean scale G0 arrest score

G$_U$ = expression of genes upregulated in quiescence

G_D = expression of genes downregulated in quiescence

G_H = expression of housekeeping genes

n = number of genes in each gene set

G0 arrest scores for the TCGA cohort were derived from expression data scaled by tumour purity

estimates. The pan-cancer TCGA samples were also classified into groups with "high" or "low"

levels of G0 arrest based on k-means clustering (k=2) on the expression data of 139 G0 biomarker

genes, following the removal of tissue-specific expression differences using the *ComBat* function

from the *sva* R package(126).

Measuring the duration of G0 arrest

We employed the GSE124109 dataset from Fujimaki et al(52) where rat embryonic fibroblasts

were transcriptomically profiled as they moved from short to long-term quiescence in the absence

of growth signals. The derived G0 arrest scores using our combined z-score methodology increased

from short to longer-term quiescence.

Validation of G0 arrest scoring methodologies

Single-cell RNA-sequencing validation datasets:

Datasets were obtained from the ArrayExpress and Gene Expression Omnibus (GEO) databases

though the following GEO Series accession numbers: GSE83142, GSE75367, GSE137912,

GSE139013, GSE90742 and E-MTAB-4547. Quality control analysis was standardised using the

SingleCellExperiment(127) and *scater*(128) R packages. Normalisation was performed using the

scran(129) R package.

Bulk RNA-sequencing validation datasets:

Datasets were obtained from the GEO database through the following GEO Series accession

numbers: GSE93391, GSE114012, GSE131594, GSE152699, GSE124854, GSE135215,

GSE99116, GSE124109 , GSE61130, GSE64553 and GSE63577. GSE114012 count data were

normalised to TPM values using the *GeoTcgaData* R package. All normalised datasets were log-

transformed before further analysis.

The accuracy with which the G0 arrest scoring methods could separate proliferating and quiescent

samples within the validation datasets was determined by calculating the area under the curve of

the receiver operating characteristic (ROC) curves, using the *plotROC* R package.

Experimental validation in lung adenocarcinoma cell lines

The average fraction of cancer cells spontaneously entering quiescence was estimated for nine lung

adenocarcinoma cell lines (NCIH460, A549, NCIH1666, NCIH1944, NCIH1563, NCIH1299,

NCIH1650, H358, L23) using EdU and phospho-Rb staining proliferation assays.

Cell lines were obtained from ATCC or Sigma and regularly checked for mycoplasma. A549 and

NCIH460 were cultured in DMEM (Gibco). NCIH358, NCIH1299 and NCIH1563 were

maintained in RPMI-1640 (Gibco) supplemented with 5mM sodium pyruvate and 0.5% glucose.

NCIH1944, NCIH1666, NCIH1650 and L23 were grown in RPMI-1640 ATCC formulation

(Gibco). A427 were cultured in EMEM (ATCC). A549, NCIH460, H358, NCIH1299, NCIH1563,

A427 were supplemented with 10% heat inactivated FBS. NCIH1666 with 5% heat-inactivated

FBS and all other cell lines with 10% non-heat inactivated FBS. All cell lines had penicillin-

streptomycin (Gibco) added to 1%. Cells were maintained at 37°C and 5% CO_2. To calculate the

quiescent fraction, A549 and NCIH460 cells were plated at a density of 500 cells/well, and all

other cell lines at a density of 1000/well, in 384well CellCarrier Ultra plates (PerkinElmer) in the

relevant media. 24h later, 5 μM EdU was added and cells were incubated for a further 24h before

fixing in a final concentration of 4% formaldehyde (15 min, RT), permeabilization with PBS/0.5%

Triton X-100 (15 min, RT) and blocking with 2% BSA in PBS (60 min, RT). The EdU signal was

detected using Click-iT chemistry, according to the manufacturer's protocol (ThermoFisher). Cells

were also labelled for phospho-Ser807/811 Rb (phospho-Rb) using Rabbit mAb 8516 (CST) at

1:2000 in blocking solution, overnight at 4°C. Unbound primary antibody was washed three times

in PBS and secondary Alexa-conjugated antibodies were used to detect the signal (ThermoFisher,

1:1000, 1h at RT). Finally nuclei were labelled with Hoechst 33258 (1 μg/ml, 15 min RT) before

imaging on a high-content widefield Operetta microscope, 20x N.A. 0.8. Automated image

analysis (Harmony, PerkinElmer) was used to segment and quantify nuclear signals in imaged

cells. Quiescent cells were defined by the absence of EdU or phospho-Rb staining, determined by

quantification of their nuclear expression (Figure 1e-f).

Endogenous PCNA was labelled at the N-terminus with a cDNA encoding mRuby in both A549

and NCI-H1944 cells, using AAV-mediated gene-targeting, according to methods described in

Zerjatke et al(54). mRuby-expressing cells were sorted into 50:50 conditioned:fresh media at

single-cell density into 96-well plates by FACS and single-cell clones expanded. For live-cell

imaging, 500 cells in phenol-red free media were plated per well of a 384 CellCarrierUltra plate

(PerkinElmer) the day before imaging. Prior to imaging, a breathable membrane was applied to the

plate and cells were imaged on the Operetta HCS microscope (PerkinElmer) at 37 °C, 5% CO2

using the 20x N.A. 0.8 objective and at 10-12 min intervals for 48h. Images were then exported

and G0/G1 length (time from mitotic exit to S-phase entry) was analysed manually in FIJI.

The G0 arrest scores for cancer cell lines were calculated using corresponding log-transformed

RPKM normalised bulk RNA-seq data from the Cancer Cell Line Encyclopedia (CCLE)

database(130).

CEP89 was depleted by ON-Target siRNA Pool from Horizon. NCI-H1299 cells were reverse

transfected in 384 well plates with 20nM of Non-targeting control (NTC) or CEP89-targeting

siRNA using Lipofectamine RNAiMax (ThermoFisher), according to the manufacturer's

instructions. Cells were left for 24h, before 5 µM EdU was added for the final 24h and then cells

were processed as above to determine the quiescent fraction. To determine the level of Cep89

depletion by western blot, cells were reverse transfected with siRNA in 24 well plates. 48h after

transfection, cells were lysed directly in 1x SDS sample buffer with 1mM DTT (ThermoFisher).

Samples were separated on pre-cast 4-20% Tris-Glycine gels, transferred to PVDF using the iBlot2

system and membranes blocked in blocking buffer (5% milk in TBS) for 1h at RT. The membrane

was then cut and the upper half was incubated in 1:1000 Cep89 antibody (Sigma, HPA040056), the

bottom half in B-actin antibody 1:2000 (CST; 3700S) diluted in blocking buffer overnight at 4'C.

Membranes were washed three times in TBS-0.05% TritonX-100 before being incubated in

secondary anti-rabbit (Cep89) or anti-mouse (B-actin) HRP conjugated antibodies (CST 7074P2

and CST 7076P2, respectively) diluted 1:2000 in blocking buffer for 1h at RT. Membranes were

washed three times again and signal detected using Clarity ECL solution (BioRad) and scanned on

an Amersham ImageQuant 800 analyser.

Multi-omics discovery cohort

FPKM normalised RNA-sequencing expression data, copy number variation gene-level data,

RPPA levels for p27 as well as mutation annotation files aligned against the GRCh38 human

reference genome from the Mutect2 pipeline were downloaded using the *TCGABiolinks* R package

(131) for 9,712 TCGA primary tumour samples across 31 solid cancer types. Haematological

malignancies were excluded as the G0 markers were derived in epithelial cells and might not be

equally suited to capture this phenotype in blood. For patients with multiple samples available, one

RNA-seq barcode entry was selected for each individual patient resulting in 9,631 total entries. All

expression data were log-transformed for downstream analysis. During G0 arrest score calculation,

expression data for the primary tumour samples was scaled according to tumour purity estimates

reported by Hoadley et al(132) to account for potential confounding cell cycle arrest signals

coming from non-tumour cells in the microenvironment. Samples with purity estimates lower than

30% were removed, leaving 8,005 samples for downstream analysis.

The mutation rates of all TCGA primary tumour samples were determined by log-transforming the

total number of mutations in each sample divided by the length of the exome capture (38Mb).

TP53 functional status was assessed based on somatic mutation and copy number alterations as

described in Zhang et al(133). *TP53* mutation and copy number for the TCGA tumours were

downloaded from cBioPortal (http://www.cbioportal.org). Tumours with *TP53* oncogenic

mutations (annotated by OncoKB) and copy-number alterations (GISTIC score ≤ -1) were assigned

as *TP53* mutant and CNV loss. Tumours without these *TP53* alterations were assigned as TP53

wild type. The effects of the *TP53* mutation status on G0 arrest were then determined with a linear

model approach with the G0 arrest score as a dependent variable and mutational status as an

independent variable. The *P* values were FDR-adjusted.

APOBEC mutagenesis enriched samples were determined through pan-cancer clustering of

mutational signature contributions as described in Wiecek et al(134). The APOBEC mutagenesis

cluster was defined as the cluster with highest mean SBS2 and SBS13 contribution. This was

repeated 100 times and only samples which appeared in the APOBEC cluster at least 50 times were

counted as being APOBEC enriched.

Aneuploidy scores and whole genome duplication events across TCGA samples were obtained

from Taylor et al(135). Microsatellite instability status for uterine corpus endometrial carcinoma,

as well as stomach and colon adenocarcinoma samples were obtained from Cortes-Ciriano et

al(78). Telomerase enzymatic activity "EXTEND" scores were obtained from Noureen et al(62).

Expression-based cancer cell stemness indices were obtained from Malta et al(63). Centrosome

amplification transcriptomic signature (CA20) scores were obtained from Almeida et al(89).

PHATE dimensionality reduction

The *phateR* R package(136) was used to perform the dimensionality reduction with a constant seed for reproducibility. The *ComBat* function from the *sva* R package(137) was used to remove tissue-specific expression patterns from the TCGA RNA-seq data.

Cancer stem cell division estimates

The mean stem cell division estimates for different cancer types used in this study were obtained from Tomasetti and Vogelstein(55).

Mutational signature estimation

Mutational signature contributions were inferred as described in Wiecek et al(134).

Machine learning of G0 arrest-linked features via ensemble elastic net regression models

The COSMIC database was used to source a list of 723 known drivers of tumorigenesis (Tiers 1+2). 285 oncogenes and tumour suppressors from a curated list showed a significant enrichment or depletion of mutations or copy number variants in samples with high levels of G0 arrest either pan-cancer or within individual TCGA studies.

To classify G0 arrest-prone from fast proliferating tumours, the 285 genes were used as input features for an ensemble elastic net regression model along the tumour mutational rate, whole-genome doubling estimates, ploidy, aneuploidy scores and 15 mutational signatures, which showed a significant correlation with G0 arrest scores either pan-cancer or within individual TCGA studies. The *caret* R package was used to build an elastic net regression model 1000 times on the training dataset of 3,753 TCGA primary tumour samples (80% of the total dataset). Only samples with at least 50 mutations were used in the model, for which mutational signatures could be reliably estimated. For each of the 1000 iterations, we randomly selected 90% of the samples from the

training dataset to build the model. Only features which were included in all 1000 model iterations

were selected for further analysis. To test the performance of our approach, a linear regression

model was built using the reduced list of genomic features and their corresponding coefficients

averaged across the 1000 elastic net regression model iterations. When applying the resulting linear

regression model on the internal validation dataset of 936 samples, we found a strong correlation

between the observed and predicted G0 arrest scores (R = 0.73, p < 2.2e-16).

SHAP values for the linear regression model used to predict G0 arrest scores were obtained using

the *fastshap* R package.

Gene enrichment and network analysis

Gene set enrichment analysis was carried out using the *ReactomePA* R package, as well as

GeneMania(138) and ConsensusPathDB(139). Interactions between *CEP89* and other cell cycle

components were inferred using the list of cell cycle genes provided by cBioPortal and GeneMania

to reconstruct the expanded network with direct interactors (*STAG1*, *CCND2*, *STAT3*). Networks

were visualised using Cytoscape(140).

Gene lists

Genes associated with the G1 phase of the cell cycle were obtained from the curated

"REACTOME_G1_PHASE " list deposited at MSigDB. Genes associated with the G1/S and

G2/M phases of the cell cycle were obtained from Tirosh et al (51).

Genes associated with apoptosis were obtained from the curated "HALLMARK_APOPTOSIS" list

deposited at MSigDB.

Genes associated with the senescence-associated secretory phenotype were obtained from Basisty

et al(60). Lists of genes making up the various DNA damage repair pathways were derived from

Pearl et al(141).

Genes associated with contact inhibition were obtained from the curated "contact inhibition" gene

ontology term. Genes associated with serum starvation were obtained from the curated

"REACTOME_CELLULAR_RESPONSE_TO_STARVATION" list deposited at MSigDB. MEK

inhibition was assessed based on the activity of the MAPK pathway as determined using an

expression signature (MPAS) consisting of 10 downstream MAPK transcripts(92).

Validation of the genomic constraints of G0 arrest

For elastic net model feature validation, RNA-seq data was downloaded for six cancer studies from

cBioPortal(142), along with patient-matched whole-genome, whole-exome and targeted

sequencing data. The 6 datasets used comprise breast cancer (SMC(143) and METABRIC(144)),

paediatric Wilms' tumor (TARGET(145)), bladder cancer, prostate adenocarcinoma and sarcoma

(MSKCC(146, 147, 148)) studies. The data were processed and analysed in the same manner as the

TCGA data. RNA-seq data for 27 MCF7 cell line strains, alongside cell line growth rates and

targeted mutational sequencing data were obtained from Ben-David et al(79).

Genomic dependency modelling in breast cancer

An ANOVA-based feature importance classification was used and identified 30 genomic features

most discriminative of samples with lower and higher than average G0 arrest scores. A random

forest model was then built using the identified features and correctly classified samples according

to their G0 arrest state with a mean accuracy of 74% across five randomly sampled test datasets

from the cohort.

Survival analysis

Multivariate Cox Proportional Hazards analysis was carried out using the *coxph* function from the

survival R package. The optimal quiescence score cut-off value of 2.95 was determined pan-cancer

using the *surv_cutpoint* function. We also used this function to determine optimal cut-offs for

individual cancer types, as presented in Figure 6c.

Treatment response single cell and bulk RNA-seq datasets

Datasets have been obtained from the GEO database through the following GEO Series accession

numbers for the cell line experiments: GSE134836, GSE134838, GSE134839, GSE137912,

GSE149224, GSE124854, GSE135215, GSE99116, GSE152699, GSE178839, GSE139944, and

the following accession numbers for the patient sample datasets: GSE191127, GSE109211,

GSE50509, GSE65185, GSE66399, GSE68871, GSE99898 (Supplementary Table 3). Unified

treatment response data for TCGA was obtained from Moiso, medRxiv 2021(149). The *umap* R

package was used for dimensionality reduction with constant seed for reproducibility.

Stress response subtype determination

TCGA cohort studies:

Samples with evidence of cell cycle arrest characterised by a generic G0 score > 0 were further

subclassified based on the most likely form of stress response, among CDK4/6 inhibition, contact

inhibition, MEK inhibition, spontaneous quiescence or serum starvation, using stress-specific

expression signatures. We opted for a conservative approach and classed each sample with a high

level of G0 arrest into a specific stress response subtype if the arrest score for the corresponding

programme was higher than one standard deviation of the distribution across the TCGA cohort, and

if the score was significantly higher than for the remaining programmes when assessed using a

Student's t test. Samples which could not be classified into any of the five stress response states

characterised in this study were classified as "uncertain".

Single-cell RNA seq treatment response datasets:

The stress response subtype of individual single cells was inferred by mapping such individual

cells onto the reference dataset of MCF10A cells reflecting different forms of G0 arrest obtained

from Min and Spencer(28). The *ComBat* R package was used to remove the study batch effect

between the expression data to be classified and the reference bulk RNA-seq data. PCA

dimensionality reduction analysis was then used on the combined datasets using the *prcomp* R

function. For each patient sample or single-cell expression data entry, a k-nearest neighbour

algorithm classification was performed using the *knn* function from the *class* R package. During

the classification the three nearest reference bulk RNA-seq data points were considered, with two

nearest neighbours with identical class needed for classification.

Optimisation of the G0 arrest signature

We investigated if a subset of the 139 G0 phase-related genes could act as a more reliable marker

of cell cycle arrest that would bypass dropout issues in single cell data. This was performed in three

steps:

(1) Assessment of individual importance as G0 arrest marker for a given gene

We collected three high confidence single cell expression datasets separating arrested from

proliferating cells. A random forest model was trained on each dataset separately to predict the

state (G0 arrest/cycling) of a given cell based on the expression levels of the 139 genes in the

signature. The Gini indices corresponding to each gene in the model were normalised to a range

of values between 0 and 1, which would reflect how important an individual gene was for

determining G0 arrest state relative to the other 138 genes. The procedure was repeated 1000

times for each of the three datasets, and the average Gini coefficients across iterations were

stored.

(2) Prioritisation of gene subsets based on cumulative importance in the model

Genes were placed in the candidate subset if their importance metric was above a given

threshold in at least one of the datasets. By gradually increasing the threshold from 0 to 1,

different gene combinations were produced.

(3) External validation of candidate subsets

The gene combinations in (2) were tested for their ability to predict G0 arrest. For this, a

separate validation dataset was utilized, which contained gene expression levels for the 139

genes in the 10 lung cancer cell lines previously employed for experimental validation, along

with the quiescence state of the lines as inferred by phospho-Rb and EdU staining. For each

gene subset, a combined z-score of G0 arrest was calculated from the expression levels as

described previously. The correlations between this z-score and the two experimental

measurements of quiescence were used to establish the ability of a gene combination to predict

quiescence. Among the top performing subsets, a 35 gene signature with a mean correlation of

78% between predicted and measured G0 arrest levels in the test data (p=0.016) showed the

highest correlation with phospho-Rb measurements capturing short-lived G0 arrest, the more

common state observed in single cell treatment datasets. Therefore, this signature was deemed

to achieve the best trade-off between gene numbers and signal capture.

The optimised gene signature is provided in Supplementary Table 4.

Statistical analysis

Groups were compared using a two-sided Student's t test, Wilcoxon rank-sum test or ANOVA, as appropriate. P-values were adjusted for multiple testing where appropriate using the Benjamini-Hochberg method. Graphs were generated using the ggplot2 and ggpubr R packages.

DECLARATIONS

Ethics approval and consent to participate

All data employed in this study comply with ethical regulations, with approval and informed consent for collection and sharing already obtained by the relevant consortia where the data were obtained from (TCGA, METABRI, MSK-IMPACT).

Consent for publication

Not applicable.

Availability of data and materials

The results published here are in part based upon data generated by the TCGA Research Network: https://www.cancer.gov/tcga, METABRIC (https://ega-archive.org/studies/EGAS00000000083), MSK-IMPACT (https://www.mskcc.org/msk-impact), or deposited at cBioPortal (https://www.cbioportal.org/) and GEO (https://www.ncbi.nlm.nih.gov/geo/). The specific GEO datasets employed in the analyses are summarised in Supplementary Tables 2 and 4.

All code developed for the purpose of this study can be found at the following repository: https://github.com/secrierlab/CancerG0Arrest

Competing interests

The authors declare that they have no competing interests.

Funding

AJW and DHJ were supported by MRC DTP grants (MR/N013867/1). MPC was supported by an Academy of Medical Science Springboard award (SBF004\1042). GMT was supported by a Wellcome Seed Award in Science (215296/Z/19/Z). MS was supported by a UKRI Future Leaders Fellowship (MR/T042184/1). Work in MS's lab was supported by a BBSRC equipment grant (BB/R01356X/1) and a Wellcome Institutional Strategic Support Fund (204841/Z/16/Z). ARB and SC are supported by a CRUK CDF (C63833/A25729) and work in ARB's lab is supported by MRC core-funding to the London Institute of Medical Sciences (MC-A658-5TY60).

Authors' contributions

MS designed the study and supervised the computational analyses. ARB designed and supervised the experimental validation in cell lines. GB supervised the analysis of p53 functional association. AJW developed the quiescence scoring methodology and performed all computational analyses to validate and apply it in bulk and single cell datasets, as well as link it to genomic features. SC performed the experimental validation of G0 arrest prevalence and CEP89 association with G0 arrest in cell lines. DK performed the inference of the minimal signature of G0 arrest applicable in single cell data. MPC performed the random forest modelling and feature selection in breast cancer. LG helped with the gene prioritisation. GMT wrote the code for batch effect correction and for PCA mapping of single cell data on a reference dataset. DHJ performed the APOBEC enrichment classification. PZ and LX performed the G0 arrest comparison of p53 wild type and mutated cancers. MS, AJW, ARB and SC wrote the manuscript, with contributions from all other authors. All authors read and approved the manuscript.

Acknowledgements

We would like to thank Prof Chris Barnes for the very helpful discussions and input on the findings of the study.

REFERENCES

1.   Hanahan D, Weinberg RA. The hallmarks of cancer. Cell. 2000;100(1):57-70.
2.   van Dijk D, Dhar R, Missarova AM, Espinar L, Blevins WR, Lehner B, et al. Slow-growing cells within isogenic populations have increased RNA polymerase error rates and DNA damage. Nat Commun. 2015;6:7972.
3.   Davis JE, Kirk J, Ji Y, Tang DG. Tumor Dormancy and Slow-Cycling Cancer Cells. Adv Exp Med Biol. 2019;1164:199-206.
4.   Phan TG, Croucher PI. The dormant cancer cell life cycle. Nat Rev Cancer. 2020;20(7):398-411.
5.   Rehman SK, Haynes J, Collignon E, Brown KR, Wang Y, Nixon AML, et al. Colorectal Cancer Cells Enter a Diapause-like DTP State to Survive Chemotherapy. Cell. 2021;184(1):226-42.e21.
6.   Chen J, Li Y, Yu TS, McKay RM, Burns DK, Kernie SG, et al. A restricted cell population propagates glioblastoma growth after chemotherapy. Nature. 2012;488(7412):522-6.
7.   Puig I, Tenbaum SP, Chicote I, Arqués O, Martínez-Quintanilla J, Cuesta-Borrás E, et al. TET2 controls chemoresistant slow-cycling cancer cell survival and tumor recurrence. J Clin Invest. 2018;128(9):3887-905.
8.   Cabanos HF, Hata AN. Emerging Insights into Targeted Therapy-Tolerant Persister Cells in Cancer. Cancers (Basel). 2021;13(11).
9.   Sharma SV, Lee DY, Li B, Quinlan MP, Takahashi F, Maheswaran S, et al. A chromatin-mediated reversible drug-tolerant state in cancer cell subpopulations. Cell. 2010;141(1):69-80.
10.   Swayden M, Chhouri H, Anouar Y, Grumolato L. Tolerant/Persister Cancer Cells and the Path to Resistance to Targeted Therapy. Cells. 2020;9(12).
11.   Hsu CH, Altschuler SJ, Wu LF. Patterns of Early p21 Dynamics Determine Proliferation-Senescence Cell Fate after Chemotherapy. Cell. 2019;178(2):361-73.e12.
12.   Xue JY, Zhao Y, Aronowitz J, Mai TT, Vides A, Qeriqi B, et al. Rapid non-uniform adaptation to conformation-specific KRAS(G12C) inhibition. Nature. 2020;577(7790):421-5.
13.   Aissa AF, Islam ABMM, Ariss MM, Go CC, Rader AE, Conrardy RD, et al. Single-cell transcriptional changes associated with drug tolerance and response to combination therapies in cancer. Nat Commun. 2021;12(1):1628.
14.   Malladi S, Macalinao DG, Jin X, He L, Basnet H, Zou Y, et al. Metastatic Latency and Immune Evasion through Autocrine Inhibition of WNT. Cell. 2016;165(1):45-60.
15.   Ribas A. Adaptive Immune Resistance: How Cancer Protects from Immune Attack. Cancer Discov. 2015;5(9):915-9.
16.   Sosa MS, Bragado P, Aguirre-Ghiso JA. Mechanisms of disseminated cancer cell dormancy: an awakening field. Nat Rev Cancer. 2014;14(9):611-22.

17. Barkan D, El Touny LH, Michalowski AM, Smith JA, Chu I, Davis AS, et al. Metastatic growth
from dormant cells induced by a col-I-enriched fibrotic environment. Cancer Res.
2010;70(14):5706-16.
18. Masago K, Fujita S, Yatabe Y. Targeting minimal residual disease after surgery with
molecular targeted therapy: the real path to a cure? J Thorac Dis. 2018;10(Suppl 17):S1982-S5.
19. Coller HA, Sang L, Roberts JM. A new description of cellular quiescence. PLoS Biol.
2006;4(3):e83.
20. Rittershaus ES, Baek SH, Sassetti CM. The normalcy of dormancy: common themes in
microbial quiescence. Cell Host Microbe. 2013;13(6):643-51.
21. Miles S, Bradley GT, Breeden LL. The budding yeast transition to quiescence. Yeast.
2021;38(1):30-8.
22. Marescal O, Cheeseman IM. Cellular Mechanisms and Regulation of Quiescence. Dev Cell.
2020;55(3):259-71.
23. Arora M, Moser J, Phadke H, Basha AA, Spencer SL. Endogenous Replication Stress in
Mother Cells Leads to Quiescence of Daughter Cells. Cell Rep. 2017;19(7):1351-64.
24. Barr AR, Cooper S, Heldt FS, Butera F, Stoy H, Mansfeld J, et al. DNA damage during S-
phase mediates the proliferation-quiescence decision in the subsequent G1 via p21 expression.
Nat Commun. 2017;8:14728.
25. Heldt FS, Barr AR, Cooper S, Bakal C, Novák B. A comprehensive model for the
proliferation-quiescence decision in response to endogenous DNA damage in human cells. Proc
Natl Acad Sci U S A. 2018;115(10):2532-7.
26. Itahana K, Dimri GP, Hara E, Itahana Y, Zou Y, Desprez PY, et al. A role for p53 in
maintaining and establishing the quiescence growth arrest in human cells. J Biol Chem.
2002;277(20):18206-14.
27. Sadasivam S, DeCaprio JA. The DREAM complex: master coordinator of cell cycle-
dependent gene expression. Nature Reviews Cancer. 2013;13(8):585-95.
28. Min M, Spencer SL. Spontaneously slow-cycling subpopulations of human cells originate
from activation of stress-response pathways. PLoS Biol. 2019;17(3):e3000178.
29. García-Gutiérrez L, Delgado MD, León J. MYC Oncogene Contributions to Release of Cell
Cycle Brakes. Genes (Basel). 2019;10(3).
30. Aguirre-Ghiso JA, Estrada Y, Liu D, Ossowski L. ERK(MAPK) activity as a determinant of
tumor growth and dormancy; regulation by p38(SAPK). Cancer Res. 2003;63(7):1684-95.
31. Spencer SL, Cappell SD, Tsai FC, Overton KW, Wang CL, Meyer T. The proliferation-
quiescence decision is controlled by a bifurcation in CDK2 activity at mitotic exit. Cell.
2013;155(2):369-83.
32. Miller I, Min M, Yang C, Tian C, Gookin S, Carter D, et al. Ki67 is a Graded Rather than a
Binary Marker of Proliferation versus Quiescence. Cell Rep. 2018;24(5):1105-12.e5.
33. Reya T, Morrison SJ, Clarke MF, Weissman IL. Stem cells, cancer, and cancer stem cells.
Nature. 2001;414(6859):105-11.
34. Kleinsmith LJ, Pierce GB, Jr. MULTIPOTENTIALITY OF SINGLE EMBRYONAL CARCINOMA
CELLS. Cancer Res. 1964;24:1544-51.
35. Ashraf HM, Moser J, Spencer SL. Senescence Evasion in Chemotherapy: A Sweet Spot for
p21. Cell. 2019;178(2):267-9.
36. Basile KJ, Aplin AE. Resistance to chemotherapy: short-term drug tolerance and stem cell-
like subpopulations. Adv Pharmacol. 2012;65:315-34.
37. Basu S, Dong Y, Kumar R, Jeter C, Tang DG. Slow-cycling (dormant) cancer cells in therapy
resistance, cancer relapse and metastasis. Semin Cancer Biol. 2022;78:90-103.

38.   Dey-Guha I, Alves CP, Yeh AC, Salony, Sole X, Darp R, et al. A mechanism for asymmetric
cell division resulting in proliferative asynchronicity. Mol Cancer Res. 2015;13(2):223-30.
39.   Turati VA, Guerra-Assunção JA, Potter NE, Gupta R, Ecker S, Daneviciute A, et al.
Chemotherapy induces canalization of cell state in childhood B-cell precursor acute lymphoblastic
leukemia. Nature Cancer. 2021.
40.   Lee E, Chuang HY, Kim JW, Ideker T, Lee D. Inferring pathway activity toward precise
disease classification. PLoS Comput Biol. 2008;4(11):e1000217.
41.   Barbie DA, Tamayo P, Boehm JS, Kim SY, Moody SE, Dunn IF, et al. Systematic RNA
interference reveals that oncogenic KRAS-driven cancers require TBK1. Nature.
2009;462(7269):108-12.
42.   Hänzelmann S, Castelo R, Guinney J. GSVA: gene set variation analysis for microarray and
RNA-seq data. BMC Bioinformatics. 2013;14:7.
43.   Foroutan M, Bhuva DD, Lyu R, Horan K, Cursons J, Davis MJ. Single sample scoring of
molecular phenotypes. BMC Bioinformatics. 2018;19(1):404.
44.   Aulestia FJ, Néant I, Dong J, Haiech J, Kilhoffer MC, Moreau M, et al. Quiescence status of
glioblastoma stem-like cells involves remodelling of Ca(2+) signalling and mitochondrial shape. Sci
Rep. 2018;8(1):9731.
45.   Kurppa KJ, Liu Y, To C, Zhang T, Fan M, Vajdi A, et al. Treatment-Induced Tumor Dormancy
through YAP-Mediated Transcriptional Reprogramming of the Apoptotic Pathway. Cancer Cell.
2020;37(1):104-22.e12.
46.   Ebinger S, Özdemir EZ, Ziegenhain C, Tiedt S, Castro Alves C, Grunert M, et al.
Characterization of Rare, Dormant, and Therapy-Resistant Cells in Acute Lymphoblastic Leukemia.
Cancer Cell. 2016;30(6):849-62.
47.   Jordan NV, Bardia A, Wittner BS, Benes C, Ligorio M, Zheng Y, et al. HER2 expression
identifies dynamic functional states within circulating breast cancer cells. Nature.
2016;537(7618):102-6.
48.   Buczacki SJA, Popova S, Biggs E, Koukorava C, Buzzelli J, Vermeulen L, et al. Itraconazole
targets cell cycle heterogeneity in colorectal cancer. J Exp Med. 2018;215(7):1891-912.
49.   Grigore F, Yang H, Hanson ND, VanBrocklin MW, Sarver AL, Robinson JP. BRAF inhibition in
melanoma is associated with the dysregulation of histone methylation and histone
methyltransferases. Neoplasia. 2020;22(9):376-89.
50.   Vairo G, Soos TJ, Upton TM, Zalvide J, DeCaprio JA, Ewen ME, et al. Bcl-2 retards cell cycle
entry through p27(Kip1), pRB relative p130, and altered E2F regulation. Mol Cell Biol.
2000;20(13):4745-53.
51.   Tirosh I, Izar B, Prakadan SM, Wadsworth MH, Treacy D, Trombetta JJ, et al. Dissecting the
multicellular ecosystem of metastatic melanoma by single-cell RNA-seq. Science.
2016;352(6282):189-96.
52.   Fujimaki K, Li R, Chen H, Della Croce K, Zhang HH, Xing J, et al. Graded regulation of cellular
quiescence depth between proliferation and senescence by a lysosomal dimmer switch. Proc Natl
Acad Sci U S A. 2019;116(45):22624-34.
53.   Stallaert W, Kedziora KM, Taylor CD, Zikry TM, Ranek JS, Sobon HK, et al. The structure of
the human cell cycle. Cell Syst. 2022;13(1):103.
54.   Zerjatke T, Gak IA, Kirova D, Fuhrmann M, Daniel K, Gonciarz M, et al. Quantitative Cell
Cycle Analysis Based on an Endogenous All-in-One Reporter for Cell Tracking and Classification.
Cell Rep. 2017;19(9):1953-66.
55.   Tomasetti C, Vogelstein B. Cancer etiology. Variation in cancer risk among tissues can be
explained by the number of stem cell divisions. Science. 2015;347(6217):78-81.

56.   Hernandez-Segura A, de Jong TV, Melov S, Guryev V, Campisi J, Demaria M. Unmasking
Transcriptional Heterogeneity in Senescent Cells. Curr Biol. 2017;27(17):2652-60.e4.
57.   Fujimaki K, Yao G. Cell dormancy plasticity: quiescence deepens into senescence through a
dimmer switch. Physiol Genomics. 2020;52(11):558-62.
58.   Dimri GP, Lee X, Basile G, Acosta M, Scott G, Roskelley C, et al. A biomarker that identifies
senescent human cells in culture and in aging skin in vivo. Proceedings of the National Academy of
Sciences. 1995;92(20):9363.
59.   Debacq-Chainiaux F, Erusalimsky JD, Campisi J, Toussaint O. Protocols to detect
senescence-associated beta-galactosidase (SA-betagal) activity, a biomarker of senescent cells in
culture and in vivo. Nat Protoc. 2009;4(12):1798-806.
60.   Basisty N, Kale A, Jeon OH, Kuehnemann C, Payne T, Rao C, et al. A proteomic atlas of
senescence-associated secretomes for aging biomarker development. PLOS Biology.
2020;18(1):e3000599.
61.   Lee BY, Han JA, Im JS, Morrone A, Johung K, Goodwin EC, et al. Senescence-associated
beta-galactosidase is lysosomal beta-galactosidase. Aging Cell. 2006;5(2):187-95.
62.   Noureen N, Wu S, Lv Y, Yang J, Alfred Yung WK, Gelfond J, et al. Integrated analysis of
telomerase enzymatic activity unravels an association with cancer stemness and proliferation.
Nature Communications. 2021;12(1):139.
63.   Malta TM, Sokolov A, Gentles AJ, Burzykowski T, Poisson L, Weinstein JN, et al. Machine
Learning Identifies Stemness Features Associated with Oncogenic Dedifferentiation. Cell.
2018;173(2):338-54.e15.
64.   Fischer M. Census and evaluation of p53 target genes. Oncogene. 2017;36(28):3943-56.
65.   Andreassen PR, Lohez OD, Lacroix FB, Margolis RL. Tetraploid state induces p53-
dependent arrest of nontransformed mammalian cells in G1. Mol Biol Cell. 2001;12(5):1315-28.
66.   Di Leonardo A, Khan SH, Linke SP, Greco V, Seidita G, Wahl GM. DNA rereplication in the
presence of mitotic spindle inhibitors in human and mouse fibroblasts lacking either p53 or pRb
function. Cancer Res. 1997;57(6):1013-9.
67.   Ganem NJ, Cornils H, Chiu SY, O'Rourke KP, Arnaud J, Yimlamai D, et al. Cytokinesis failure
triggers hippo tumor suppressor pathway activation. Cell. 2014;158(4):833-48.
68.   Alexandrov LB, Kim J, Haradhvala NJ, Huang MN, Tian Ng AW, Wu Y, et al. The repertoire
of mutational signatures in human cancer. Nature. 2020;578(7793):94-101.
69.   Alexandrov LB, Nik-Zainal S, Wedge DC, Aparicio SAJR, Behjati S, Biankin AV, et al.
Signatures of mutational processes in human cancer. Nature. 2013;500(7463):415-21.
70.   Pich O, Muiños F, Lolkema MP, Steeghs N, Gonzalez-Perez A, Lopez-Bigas N. The
mutational footprints of cancer therapies. Nat Genet. 2019;51(12):1732-40.
71.   Miyazono K, Maeda S, Imamura T. Coordinate regulation of cell growth and differentiation
by TGF-beta superfamily and Runx proteins. Oncogene. 2004;23(24):4232-7.
72.   Cibis H, Biyanee A, Dörner W, Mootz HD, Klempnauer KH. Characterization of the zinc
finger proteins ZMYM2 and ZMYM4 as novel B-MYB binding proteins. Sci Rep. 2020;10(1):8390.
73.   Maley CC, Reid BJ. Natural selection in neoplastic progression of Barrett's esophagus.
Semin Cancer Biol. 2005;15(6):474-83.
74.   Bardeesy N, DePinho RA. Pancreatic cancer biology and genetics. Nat Rev Cancer.
2002;2(12):897-909.
75.   Indovina P, Pentimalli F, Casini N, Vocca I, Giordano A. RB1 dual role in proliferation and
apoptosis: cell fate control and implications for cancer therapy. Oncotarget. 2015;6(20).
76.   Mas-Ponte D, Supek F. DNA mismatch repair promotes APOBEC3-mediated diffuse
hypermutation in human cancers. Nat Genet. 2020;52(9):958-68.

77.   Kloor M, von Knebel Doeberitz M. The Immune Biology of Microsatellite-Unstable Cancer.
Trends Cancer. 2016;2(3):121-33.
78.   Cortes-Ciriano I, Lee S, Park W-Y, Kim T-M, Park PJ. A molecular portrait of microsatellite
instability across multiple cancers. Nature Communications. 2017;8(1):15180.
79.   Ben-David U, Siranosian B, Ha G, Tang H, Oren Y, Hinohara K, et al. Genetic and
transcriptional evolution alters cancer cell line drug response. Nature. 2018;560(7718):325-30.
80.   Yersal O, Barutca S. Biological subtypes of breast cancer: Prognostic and therapeutic
implications. World J Clin Oncol. 2014;5(3):412-24.
81.   Jakobsen L, Vanselow K, Skogs M, Toyoda Y, Lundberg E, Poser I, et al. Novel
asymmetrically localizing components of human centrosomes identified by complementary
proteomics methods. Embo j. 2011;30(8):1520-35.
82.   Sillibourne JE, Specht CG, Izeddin I, Hurbain I, Tran P, Triller A, et al. Assessing the
localization of centrosomal proteins by PALM/STORM nanoscopy. Cytoskeleton (Hoboken).
2011;68(11):619-27.
83.   Bettencourt-Dias M, Glover DM. Centrosome biogenesis and function: centrosomics brings
new understanding. Nature Reviews Molecular Cell Biology. 2007;8(6):451-63.
84.   Grallert A, Chan KY, Alonso-Nuñez ML, Madrid M, Biswas A, Alvarez-Tabarés I, et al.
Removal of centrosomal PP1 by NIMA kinase unlocks the MPF feedback loop to promote mitotic
commitment in S. pombe. Curr Biol. 2013;23(3):213-22.
85.   Wigley WC, Fabunmi RP, Lee MG, Marino CR, Muallem S, DeMartino GN, et al. Dynamic
association of proteasomal machinery with the centrosome. J Cell Biol. 1999;145(3):481-90.
86.   Itoh K, Jenny A, Mlodzik M, Sokol SY. Centrosomal localization of Diversin and its relevance
to Wnt signaling. J Cell Sci. 2009;122(Pt 20):3791-8.
87.   Kfoury Y, Nasr R, Favre-Bonvin A, El-Sabban M, Renault N, Giron ML, et al. Ubiquitylated
Tax targets and binds the IKK signalosome at the centrosome. Oncogene. 2008;27(12):1665-76.
88.   Lin A, Wang RT, Ahn S, Park CC, Smith DJ. A genome-wide map of human genetic
interactions inferred from radiation hybrid genotypes. Genome Res. 2010;20(8):1122-32.
89.   de Almeida BP, Vieira AF, Paredes J, Bettencourt-Dias M, Barbosa-Morais NL. Pan-cancer
association of a centrosome amplification gene expression signature with genomic alterations and
clinical outcome. PLOS Computational Biology. 2019;15(3):e1006832.
90.   Kohlmaier G, Loncarek J, Meng X, McEwen BF, Mogensen MM, Spektor A, et al. Overly
long centrioles and defective cell division upon excess of the SAS-4-related protein CPAP. Curr
Biol. 2009;19(12):1012-8.
91.   Tang C-JC, Fu R-H, Wu K-S, Hsu W-B, Tang TK. CPAP is a cell-cycle regulated protein that
controls centriole length. Nature Cell Biology. 2009;11(7):825-31.
92.   Wagle MC, Kirouac D, Klijn C, Liu B, Mahajan S, Junttila M, et al. A transcriptional MAPK
Pathway Activity Score (MPAS) is a clinically relevant biomarker in multiple cancer types. NPJ
Precis Oncol. 2018;2(1):7.
93.   Hafner M, Mills CE, Subramanian K, Chen C, Chung M, Boswell SA, et al. Multiomics
Profiling Establishes the Polypharmacology of FDA-Approved CDK4/6 Inhibitors and the Potential
for Differential Clinical Activity. Cell Chem Biol. 2019;26(8):1067-80.e8.
94.   Salvador-Barbero B, Álvarez-Fernández M, Zapatero-Solana E, El Bakkali A, Menéndez
MDC, López-Casas PP, et al. CDK4/6 Inhibitors Impair Recovery from Cytotoxic Chemotherapy in
Pancreatic Adenocarcinoma. Cancer Cell. 2020;37(3):340-53.e6.
95.   Guiley KZ, Stevenson JW, Lou K, Barkovich KJ, Kumarasamy V, Wijeratne TU, et al. p27
allosterically activates cyclin-dependent kinase 4 and antagonizes palbociclib inhibition. Science.
2019;366(6471).

96.   Pack LR, Daigh LH, Chung M, Meyer T. Clinical CDK4/6 inhibitors induce selective and
immediate dissociation of p21 from cyclin D-CDK4 to inhibit CDK2. Nat Commun.
2021;12(1):3356.
97.   Zhu X, Chen L, Huang B, Wang Y, Ji L, Wu J, et al. The prognostic and predictive potential of
Ki-67 in triple-negative breast cancer. Scientific Reports. 2020;10(1):225.
98.   Solé L, Lobo-Jarne T, Álvarez-Villanueva D, Alonso-Marañón J, Guillén Y, Guix M, et al. p53
wild-type colorectal cancer cells that express a fetal gene signature are associated with metastasis
and poor prognosis. Nat Commun. 2022;13(1):2866.
99.   Brown JA, Yonekubo Y, Hanson N, Sastre-Perona A, Basin A, Rytlewski JA, et al. TGF-β-
Induced Quiescence Mediates Chemoresistance of Tumor-Propagating Cells in Squamous Cell
Carcinoma. Cell Stem Cell. 2017;21(5):650-64.e8.
100.   Park SR, Namkoong S, Friesen L, Cho CS, Zhang ZZ, Chen YC, et al. Single-Cell
Transcriptome Analysis of Colon Cancer Cell Response to 5-Fluorouracil-Induced DNA Damage.
Cell Rep. 2020;32(8):108077.
101.   Aissa AF, Islam A, Ariss MM, Go CC, Rader AE, Conrardy RD, et al. Single-cell transcriptional
changes associated with drug tolerance and response to combination therapies in cancer. Nat
Commun. 2021;12(1):1628.
102.   Tan X, Thapa N, Sun Y, Anderson RA. A kinase-independent role for EGF receptor in
autophagy initiation. Cell. 2015;160(1-2):145-60.
103.   Srivatsan SR, McFaline-Figueroa JL, Ramani V, Saunders L, Cao J, Packer J, et al. Massively
multiplex chemical transcriptomics at single-cell resolution. Science. 2020;367(6473):45-51.
104.   Hatzis C, Pusztai L, Valero V, Booser DJ, Esserman L, Lluch A, et al. A genomic predictor of
response and survival following taxane-anthracycline chemotherapy for invasive breast cancer.
JAMA. 2011;305(18):1873-81.
105.   Lee JS, Nair NU, Dinstag G, Chapman L, Chung Y, Wang K, et al. Synthetic lethality-
mediated precision oncology via the tumor transcriptome. Cell. 2021;184(9):2487-502.e13.
106.   Moser J, Miller I, Carter D, Spencer SL. Control of the Restriction Point by Rb and p21. Proc
Natl Acad Sci U S A. 2018;115(35):E8219-e27.
107.   Bačević K, Lossaint G, Achour TN, Georget V, Fisher D, Dulić V. Cdk2 strengthens the intra-S
checkpoint and counteracts cell cycle exit induced by DNA damage. Sci Rep. 2017;7(1):13429.
108.   Terzi MY, Izmirli M, Gogebakan B. The cell fate: senescence or quiescence. Mol Biol Rep.
2016;43(11):1213-20.
109.   Kleffel S, Schatton T. Tumor dormancy and cancer stem cells: two sides of the same coin?
Adv Exp Med Biol. 2013;734:145-79.
110.   Miller AK, Brown JS, Enderling H, Basanta D, Whelan CJ. The Evolutionary Ecology of
Dormancy in Nature and in Cancer. Frontiers in Ecology and Evolution. 2021;9.
111.   Considine MJ, Considine JA. On the language and physiology of dormancy and quiescence
in plants. J Exp Bot. 2016;67(11):3189-203.
112.   Velappan Y, Signorelli S, Considine MJ. Cell cycle arrest in plants: what distinguishes
quiescence, dormancy and differentiated G1? Ann Bot. 2017;120(4):495-509.
113.   Sosa MS, Parikh F, Maia AG, Estrada Y, Bosch A, Bragado P, et al. NR2F1 controls tumour
cell dormancy via SOX9- and RARβ-driven quiescence programmes. Nat Commun. 2015;6:6170.
114.   Fox DB, Garcia NMG, McKinney BJ, Lupo R, Noteware LC, Newcomb R, et al. NRF2
activation promotes the recurrence of dormant tumour cells through regulation of redox and
nucleotide metabolism. Nat Metab. 2020;2(4):318-34.
115.   Berkel C, Cacan E. Copy number and expression of CEP89, a protein required for
ciliogenesis, are increased and predict poor prognosis in patients with ovarian cancer. Cell
Biochem Funct. 2022;40(3):298-309.

116.   Tanos BE, Yang HJ, Soni R, Wang WJ, Macaluso FP, Asara JM, et al. Centriole distal
appendages promote membrane docking, leading to cilia initiation. Genes Dev. 2013;27(2):163-8.
117.   Sillibourne JE, Hurbain I, Grand-Perret T, Goud B, Tran P, Bornens M. Primary ciliogenesis
requires the distal appendage component Cep123. Biol Open. 2013;2(6):535-45.
118.   Wheway G, Nazlamova L, Hancock JT. Signaling through the Primary Cilium. Front Cell Dev
Biol. 2018;6:8.
119.   van Bon BW, Oortveld MA, Nijtmans LG, Fenckova M, Nijhof B, Besseling J, et al. CEP89 is
required for mitochondrial metabolism and neuronal function in man and fly. Hum Mol Genet.
2013;22(15):3138-51.
120.   Huang R, Chen H, Liang J, Li Y, Yang J, Luo C, et al. Dual Role of Reactive Oxygen Species
and their Application in Cancer Therapy. J Cancer. 2021;12(18):5543-61.
121.   Gupta J, del Barco Barrantes I, Igea A, Sakellariou S, Pateras IS, Gorgoulis VG, et al. Dual
function of p38α MAPK in colon cancer: suppression of colitis-associated tumor initiation but
requirement for cancer cell survival. Cancer Cell. 2014;25(4):484-500.
122.   Moon EJ, Giaccia A. Dual roles of NRF2 in tumor prevention and progression: possible
implications in cancer treatment. Free Radic Biol Med. 2015;79:292-9.
123.   Sosa MS, Avivar-Valderas A, Bragado P, Wen HC, Aguirre-Ghiso JA. ERK1/2 and p38α/β
signaling in tumor cell quiescence: opportunities to control dormant residual disease. Clin Cancer
Res. 2011;17(18):5850-7.
124.   Minassian LM, Cotechini T, Huitema E, Graham CH. Hypoxia-Induced Resistance to
Chemotherapy in Cancer. Adv Exp Med Biol. 2019;1136:123-39.
125.   Aguirre-Ghiso JA, Sosa MS. Emerging Topics on Disseminated Cancer Cell Dormancy and
the Paradigm of Metastasis. Annual Review of Cancer Biology. 2018;2(1):377-93.
126.   Johnson WE, Li C, Rabinovic A. Adjusting batch effects in microarray expression data using
empirical Bayes methods. Biostatistics. 2007;8(1):118-27.
127.   Amezquita RA, Lun ATL, Becht E, Carey VJ, Carpp LN, Geistlinger L, et al. Orchestrating
single-cell analysis with Bioconductor. Nature Methods. 2020;17(2):137-45.
128.   McCarthy DJ, Campbell KR, Lun ATL, Wills QF. Scater: pre-processing, quality control,
normalization and visualization of single-cell RNA-seq data in R. Bioinformatics. 2017;33(8):1179-
86.
129.   Lun AT, McCarthy DJ, Marioni JC. A step-by-step workflow for low-level analysis of single-
cell RNA-seq data with Bioconductor. F1000Res. 2016;5:2122.
130.   Barretina J, Caponigro G, Stransky N, Venkatesan K, Margolin AA, Kim S, et al. The Cancer
Cell Line Encyclopedia enables predictive modelling of anticancer drug sensitivity. Nature.
2012;483(7391):603-7.
131.   Colaprico A, Silva TC, Olsen C, Garofano L, Cava C, Garolini D, et al. TCGAbiolinks: an
R/Bioconductor package for integrative analysis of TCGA data. Nucleic Acids Res. 2016;44(8):e71.
132.   Hoadley KA, Yau C, Hinoue T, Wolf DM, Lazar AJ, Drill E, et al. Cell-of-Origin Patterns
Dominate the Molecular Classification of 10,000 Tumors from 33 Types of Cancer. Cell.
2018;173(2):291-304.e6.
133.   Zhang P, Kitchen-Smith I, Xiong L, Stracquadanio G, Brown K, Richter PH, et al. Germline
and Somatic Genetic Variants in the p53 Pathway Interact to Affect Cancer Risk, Progression, and
Drug Response. Cancer Res. 2021;81(7):1667-80.
134.   Wiecek AJ, Jacobson DH, Lason W, Secrier M. Pan-Cancer Survey of Tumor Mass Dormancy
and Underlying Mutational Processes. Frontiers in Cell and Developmental Biology. 2021;9:1820.
135.   Taylor AM, Shih J, Ha G, Gao GF, Zhang X, Berger AC, et al. Genomic and Functional
Approaches to Understanding Cancer Aneuploidy. Cancer Cell. 2018;33(4):676-89.e3.

136.   Moon KR, van Dijk D, Wang Z, Gigante S, Burkhardt DB, Chen WS, et al. Visualizing
structure and transitions in high-dimensional biological data. Nat Biotechnol. 2019;37(12):1482-
92.
137.   Leek JT, Johnson WE, Parker HS, Jaffe AE, Storey JD. The sva package for removing batch
effects and other unwanted variation in high-throughput experiments. Bioinformatics.
2012;28(6):882-3.
138.   Warde-Farley D, Donaldson SL, Comes O, Zuberi K, Badrawi R, Chao P, et al. The
GeneMANIA prediction server: biological network integration for gene prioritization and
predicting gene function. Nucleic Acids Res. 2010;38(Web Server issue):W214-20.
139.   Kamburov A, Pentchev K, Galicka H, Wierling C, Lehrach H, Herwig R. ConsensusPathDB:
toward a more complete picture of cell biology. Nucleic Acids Res. 2011;39(Database issue):D712-
7.
140.   Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D, et al. Cytoscape: a software
environment for integrated models of biomolecular interaction networks. Genome Res.
2003;13(11):2498-504.
141.   Pearl LH, Schierz AC, Ward SE, Al-Lazikani B, Pearl FMG. Therapeutic opportunities within
the DNA damage response. Nature Reviews Cancer. 2015;15(3):166-80.
142.   Cerami E, Gao J, Dogrusoz U, Gross BE, Sumer SO, Aksoy BA, et al. The cBio cancer
genomics portal: an open platform for exploring multidimensional cancer genomics data. Cancer
Discov. 2012;2(5):401-4.
143.   Kan Z, Ding Y, Kim J, Jung HH, Chung W, Lal S, et al. Multi-omics profiling of younger Asian
breast cancers reveals distinctive molecular signatures. Nat Commun. 2018;9(1):1725.
144.   Curtis C, Shah SP, Chin SF, Turashvili G, Rueda OM, Dunning MJ, et al. The genomic and
transcriptomic architecture of 2,000 breast tumours reveals novel subgroups. Nature.
2012;486(7403):346-52.
145.   Gadd S, Huff V, Walz AL, Ooms AHAG, Armstrong AE, Gerhard DS, et al. A Children's
Oncology Group and TARGET initiative exploring the genetic landscape of Wilms tumor. Nat
Genet. 2017;49(10):1487-94.
146.   Iyer G, Al-Ahmadie H, Schultz N, Hanrahan AJ, Ostrovnaya I, Balar AV, et al. Prevalence and
co-occurrence of actionable genomic alterations in high-grade bladder cancer. J Clin Oncol.
2013;31(25):3133-40.
147.   Taylor BS, Schultz N, Hieronymus H, Gopalan A, Xiao Y, Carver BS, et al. Integrative
genomic profiling of human prostate cancer. Cancer Cell. 2010;18(1):11-22.
148.   Barretina J, Taylor BS, Banerji S, Ramos AH, Lagos-Quintana M, Decarolis PL, et al. Subtype-
specific genomic alterations define new targets for soft-tissue sarcoma therapy. Nat Genet.
2010;42(8):715-21.
149.   Moiso E. Manual curation of TCGA treatment data and identification of potential markers
of therapy response. medRxiv. 2021:2021.04.30.21251941.
1255

FIGURES

1257

Figure 1: Methodology for quantifying G0 arrest in cancer. (a) Workflow for evaluating G0

arrest from RNA-seq data. 139 genes differentially expressed in multiple forms of quiescence were

employed to score G0 arrest across cancer tissues. **(b)** Receiver operating characteristic (ROC)

curves illustrating the performance of the z-score methodology on separating actively proliferating

and G0 arrested cells in seven single-cell (continuous curves) and bulk RNA-seq (dotted curves)

datasets. AUC = area under the curve. **(c)** Compared classification accuracies of the G0 arrest z-

score approach and classic cell proliferation markers across the seven single-cell/bulk RNA-seq

validation datasets. **(d)** G0 arrest levels of embryonic fibroblast cells under serum starvation for

various amounts of time. Replicates are depicted in the same colour. **(e)** Representative images of

lung cancer cell lines immunostained and analysed to detect the G0 arrest fraction. Hoechst (labels

all nuclei) is in blue, phospho-Rb in green and EdU in red in merged image. White dashed circles

highlight G0 arrested cells that are negative for both phospho-Rb and EdU signals. Scale bar: 100

μm. **(f)** Graphs show single cell quantification of phospho-Rb and EdU intensities taken from

images and used to define the cut-off to calculate the G0 arrest fraction (green boxes). Images in

(e) and graphs in **(f)** are taken from the A549 cell line. **(g-h)** Correlation between theoretical

estimates of a G0 or G1 state and the fraction of cells entering G0 arrest in nine lung

adenocarcinoma cell lines, as assessed through **(g)** phospho-Rb assays and **(h)** EdU assays. Mean

of n=3 is shown for the average percentage of G0 arrested cells.

1276

1279 Figure 2: Pan-cancer evaluation of proliferative heterogeneity and linked tumour hallmarks.

(a) PHATE plot illustrating the wide spectrum of proliferative to slow cycling/arrested states

across 8,005 primary tumour samples from TCGA. Each sample is coloured according to the

relative G0 arrest level. **(b)** Variation in tumour G0 arrest levels across different cancer tissues. **(c)**

Correlation between mean G0 arrest capacity and stem cell division estimates for various tissue

types. **(d)** Correlating tumour G0 arrest scores with cancer cell stemness (Stemness Index),

telomerase activity (EXTEND score), p21 activity (CDKN1A) and the expression of several

commonly used proliferation markers. The Pearson correlation coefficient is displayed. RC –

replication complex. **(e)** Consistently higher levels of G0 arrest are detected in samples with

functional p53. **(f)** Lower G0 arrest scores are observed in tumours with one or two whole genome

duplication events. Wilcoxon rank-sum test p-values are displayed in boxplots, *p<0.05; **p<0.01;

***p<0.001; ****p<0.0001.

1291

Figure 3: Genomic landscape of G0 arrest decisions in cancer. (a) Cancer drivers with

1292

mutations or copy number alterations depleted pan-cancer in a G0 arrest context. Features further

selected by the pan-cancer model are highlighted. **(b)** Schematic of the ensemble elastic net

modelling employed to prioritise genomic changes associated with G0 arrest. **(c)** Genomic events

significantly associated with G0 arrest, ranked according to their importance in the model (highest

to lowest). Each point depicts an individual tumour sample, coloured by the value of the respective

feature. For discrete variables purple indicates the presence of the feature and green its absence.

The Shapley values indicate the impact of individual feature values on the G0 arrest score

prediction. **(d)** G0 arrest levels are significantly reduced in microsatellite unstable (MSI) samples

in stomach adenocarcinoma (STAD) and uterine corpus endometrial carcinoma (UCEC), with the

same trend (albeit not significant) shown in colon adenocarcinoma (COAD). Wilcoxon rank-sum

test *p<0.05; **p<0.01. **(e)** Genomic alterations are depleted across DNA repair pathways during

G0 arrest. Odds ratios of mutational load on pathway in G0 arrest are depicted, along with

confidence intervals. CS=chromosome segregation; p53=p53 pathway; UR=ubiquitylation

response; CPF=checkpoint factors; TM=telomere maintenance; CR=chromatin remodelling;

TLS=translesion synthesis; NHEJ=non-homologous end joining; NER=nucleotide excision repair;

MMR=mismatch repair; FA=Fanconi Anaemia; BER=base excision repair. **(f)** G0 arrest scores are

increased in cell lines with slow doubling time across MCF7 strains, which also show lower

prevalence of PTEN mutations. **(g)** Tissue-specific changes in G0 arrest between samples

with/without quiescence-associated deletions (blue), amplifications (red) and SNVs (brown) within

the TCGA cohort (top) and external validation datasets (bottom).

1313

Figure 4: CEP89 amplification is associated with lower G0 arrest capacity. (a) Network

illustrating CEP89 interactions with cell cycle genes (from GeneMania). The edge colour indicates

the interaction type, with green representing genetic interactions, orange representing predicted

interactions and purple indicating pathway interactions. The edge width illustrates the interaction

weight. **(b)** CA20 scores are significantly increased in TCGA primary tumours containing a *CEP89*

amplification. **(c)** Pan-cancer relationship between CA20 and G0 arrest scores across the TCGA

cohort. **(d)** Cox proportional hazards analysis estimates of the log hazards ratio for the impact of

CEP89 expression on patient prognosis within individual cancer studies, after adjusting for tumour

stage. Patients with high expression of *CEP89* show significantly worse prognosis within ACC,

LUSC, LIHC, KIRC and STAD, but significantly better prognosis within HNSC, PAAD and KIRP

studies. **(e)** Western blot showing depletion of Cep89 protein 48h after siRNA transfection of NCI-

H1299 cells. Mock is lipofectamine only, NTC is Non-targeting control siRNA. B-actin is used as

a loading control. **(f)** Graphs show that Cep89 depletion in NCI-H1299 cells leads to a reduction in

nuclear number and an increase in the fraction of G0 arrested cells, measured by an increase in the

percentage of EdU negative (24h EdU pulse) and Phospho-Ser 807/811 Rb negative cells. One-

Way ANOVA, *p<0.05, **p<0.01. Mean of n=3.

1330

1331

1332

Figure 5: Pan-cancer characterisation of individual G0 stress response programmes. (a-e)

Comparison of correlation coefficients between stress response programme scores and **(a)** mean

expression of CDK4 and CDK6, **(b)** mean expression of curated contact inhibition genes, **(c)** a

transcriptional MAPK Pathway Activity Score (MPAS), **(d)** mean expression of curated serum

starvation genes, and **(e)** CDKN1A expression, across TCGA cancers. The correlations expected to

be strongest (either negative or positive) are denoted by an asterisk. The generic G0 arrest score

refers to scores calculated using the original list of 139 genes differentially expressed across all 5

forms of G0 arrest. **(f)** Comparison of stress response programme scores measured in cancer cell

lines before (grey) and after (red) Palbociclib treatment across three validation studies. Datasets

used for validation are denoted by their corresponding GEO series accession number. **(g)** Predicted

stress response diversity in samples with high levels of G0 arrest across individual cancer types.

1344   The same colour legend as in **(a)** is applied. Gray bars represent the proportion of samples for

1345   which the G0 arrest inducer could not be estimated.

1346

Figure 6: Impact of G0 arrest on patient prognosis and treatment response. (a) Disease-

specific survival based on proliferation/G0 arrest levels for patients from TCGA within 15 years of

follow up. Patients with increased levels of G0 arrest in primary tumours showed significantly

better prognosis than patients with fast proliferating tumours. **(b-c)** Hazard ratio ranges illustrating

the impact of different forms of G0 induction **(b)** and different tissues **(c)** on patient prognosis,

after taking into account potential confounding factors. Values above 0 indicate significantly better

prognosis when tumours contain high proportions of cells arrested in G0. **(d)** Change in G0 arrest

scores inferred from bulk RNA-seq across breast, pancreatic, colorectal and skin cancer cells in

response to treatment with the CDK4/6 inhibitor Palbociclib, 5-FU or the BRAF inhibitor

Vemurafenib. **(e-f)** UMAP plot illustrating the response of the TP53-proficient RKO colorectal

cancer cell line to various 5-FU doses and the corresponding proportions of cells predicted to be

arrested/proliferating. Each dot is an individual cell, coloured according to its G0 arrest level. **(g-h)**

The same as previous, but for the TP53-deficient SW480 cell line. **(i-j)** UMAP plot illustrating the

response of individual PC9 NSCLC cells to the EGFR inhibitor Erlotinib across several time points

and the corresponding proportion of cells predicted to be arrested/proliferating. **(k)** Principal

component analysis illustrating the superimposition of single cell RNA-seq profiles (circles) of G0

arrested NSCLC cells before/after EGFR inhibition onto the bulk RNA-seq reference data

(triangles) for MCF10A cells occupying various stress response states. **(l)** The proportion of

NSCLC cells in **(k)** predicted to occupy different stress response states across several time points.

1367

1368

Figure 7: Optimisation of the G0 arrest signature for use in single cell RNA-seq data. (a)

Methodology for refining the gene signature of G0 arrest: random forest classifiers are trained to

distinguish arrested from cycling tumours on three high confidence datasets; Gini index

thresholding is optimised to prioritise a final list of 35 genes. **(b)** Gini index variation, correlation

with experimentally measured quiescence via EdU and phospho-Rb staining assays, and

corresponding p-values are plotted as the number of genes considered in the model is increased.

The red dotted line indicates the threshold chosen for the final solution of 35 genes. The black

dotted line indicates the threshold for p-value significance. **(c)** Additional external validation of the

35 gene signature acting as a classifier of G0 arrested and proliferating cells in single cell and bulk

datasets. **(d)** Dropout in single cell data by gene signature. The percentage of genes out of the 35

(red) and 139 (grey) gene lists with reported expression across the single-cell RNA-seq datasets

analysed in this study. **(e)** Proportion of cycling and G0 arrested cells estimated in single cell

datasets of p53 wild type and mutant lines treated with 5FU, as well as cells treated with EGFR

inhibitors. Data as in Figure 6.

1383