Genomic hallmarks of cellular dormancy in cancer and therapeutic implications

Anna J. Wiecek[1], Stephen J. Cutty[2], Daniel Kornai[1], Mario Parreno-Centeno[1], Lucie E. Gourmet[1], Guidantonio Malagoli Tagliazucchi[1], Daniel H. Jacobson[1,3], Ping Zhang[4], Lingyun Xiong[4], Gareth L. Bond[5], Alexis R. Barr[2,6], Maria Secrier[1,*]

[1] UCL Genetics Institute, Department of Genetics, Evolution and Environment, University College London, London, UK

[2] Institute of Clinical Sciences, Faculty of Medicine, Imperial College London, London, UK

[3] UCL Cancer Institute, Paul O'Gorman Building, University College London, London, UK

[4] Wellcome Centre for Human Genetics, University of Oxford, Oxford, UK

[5] Institute of Cancer and Genomic Sciences, University of Birmingham, Edgbaston, Birmingham, UK

[6] Cell Cycle Control Team, MRC London Institute of Medical Sciences (LMS), London, UK

* To whom correspondence should be addressed (m.secrier@ucl.ac.uk)

ABSTRACT

Therapy resistance in cancer is often driven by a subpopulation of cells that are temporarily arrested in a non-proliferative, quiescent or 'dormant' state, which is difficult to capture and whose mutational drivers remain largely unknown. We developed methodology to uniquely identify this state from transcriptomic signals and characterised its prevalence and genomic constraints in solid primary tumours. We show dormancy preferentially emerges in the context of more stable, less mutated genomes which maintain *TP53* integrity and lack the hallmarks of DNA damage repair deficiency, while presenting increased APOBEC mutagenesis. We uncover novel genomic dependencies of this process, including the amplification of the centrosomal gene *CEP89* as a driver of dormancy impairment. Lastly, we demonstrate that dormancy underlies unfavourable responses to various therapies exploiting cell cycle, kinase signalling and epigenetic mechanisms in single cell data, and propose a signature of dormancy-linked therapeutic resistance to further study and clinically track this state.

1 INTRODUCTION

Quiescence is a widely conserved state of reversible proliferative arrest, essential for the normal

development and homeostasis of eukaryotic organisms[1,2]. During quiescence, cells are halted in

the G0 phase of the cell cycle, where they maintain minimal basal activity[2]. In cancer, this

temporary cell cycle arrest, also termed cellular dormancy, has been proposed as one of the

contributing factors of resistance to treatments that target actively dividing cells, such as

chemotherapy[3-5]. Beyond these, a variety of other pre-existing or acquired resistance scenarios

have been explained by a drug-tolerant 'persister' cell state represented by slow cycling or

entirely quiescent cells[6,7]. Their presence is associated with minimal residual disease, a major

cause of relapse in the clinic[8]. As the neoplastic cells evolve, dormancy can also be employed

as a mechanism to facilitate immune evasion[9,10] or adaptation to new environmental niches

during metastatic seeding[11,12].

Despite increasing evidence for the importance of dormancy in cancer progression, it remains a

poorly defined state. Hampering our understanding is the fact that quiescence represents a

number of heterogeneous states[1]. Canonically, cells can be forced into quiescence through

serum starvation, mitogen withdrawal or contact inhibition[1]. Cells can also enter dormancy

spontaneously in response to cell-intrinsic factors like replication stress[13-15]. This process is

controlled by p53[16], which triggers the inhibition of cyclin-CDK complexes by activating p21[14].

This in turn allows the assembly of the DREAM complex - a key effector responsible for

repression of cell-cycle dependent gene expression[17]. Min and Spencer[18] recently demonstrated

a much broader systemic coordination of 198 genes underlying distinct types of quiescence by

profiling the transcriptomes of cells that entered this state either spontaneously or upon different

stimuli. Additionally, proliferation-quiescence decisions can be impacted by oncogenic changes

such as *MYC* amplification[19] or altered p38/ERK signalling[20].

Despite these advances, the identification of dormant cells within tumours presents an ongoing

challenge due to their scarcity and lack of universal, easily measurable markers for the activation

and maintenance of this state. As they are often defined by a lack of proliferative markers[21,22],

quiescent cells can be mistaken for other therapy resistant cell types such as senescent or cancer

stem cells[6,23]. Unlike quiescent cells, which can readily resume their proliferative state, senescent

cells are irreversibly arrested[18] while cancer stem cells have a high capacity for self-renewal and

sit at the top of the differentiation hierarchy[24]. Biomarkers of quiescence that are sufficiently

specific and robust to be clinically useful are clearly needed.

Furthermore, the extent to which dormancy is enacted through transcriptional or genetic control

is unknown[5,25], as the mutational processes and genomic changes driving quiescence have not

been systematically profiled across cancer tissues. Understanding the evolutionary triggers and

molecular mechanisms that enable cancer cells to enter and maintain dormancy would enable

us to develop pharmacological strategies to selectively eradicate quiescent cancer cells or

prevent them from re-entering proliferative cycles.

To address these challenges, we have developed a new method to uniquely and reliably quantify

tumour dormancy using transcriptomic data, and employed it to characterise this phenomenon

in bulk and single cell datasets from a variety of solid tumours. We identify and validate mutational

constraints for dormancy emergence, hinting at potential new therapeutic targets that could

exploit this mechanism. We also demonstrate the relevance of dormancy to responses to a range

of compounds targeting cell cycle, kinase signalling and epigenetic mechanisms in single cell

datasets, and propose an expression signature that could be employed to detect tumour

dormancy-induced treatment resistance.

47
RESULTS

Evaluating tumour dormancy from transcriptomic data

We hypothesised that primary tumours contain varying numbers of quiescent cells, and this in

turn may determine their ability to overcome therapies targeting cycling cells. To capture this

elusive phenotype, we developed a computational framework that would allow us to quantify

dormancy signals in bulk and single cell sequenced cancer samples (Figure 1a). We employed

139 differentially expressed genes across five distinct forms of quiescence resulting from serum

starvation, contact inhibition, MEK inhibition, CDK4/6 inhibition and spontaneous quiescence, as

described by Min and Spencer[18], as markers of a generic quiescence state. The expression

levels of these markers were used to derive a score reflecting the relative level of dormancy

within individual tumours (see Methods).

We validated multiple scoring methodologies[26-29] using seven single-cell and bulk datasets[30-36]

where actively proliferating and quiescent cells had been independently profiled (Supplementary

Figure 1a, Supplementary Table 1, Methods). A combined Z-score approach had the highest

accuracy in detecting signals of dormancy, with a 91% mean performance in classifying cells as

quiescent or cycling (Figure 1b, Supplementary Figure 1b). It also outperformed classical cell

cycle and dormancy markers, such as the expression of targets of the DREAM complex, CDK2,

Ki67 and of mini-chromosome replication maintenance (MCM) protein complex genes - which

are involved in the initiation of eukaryotic genome replication (Figure 1c). Importantly, our

approach provided a good separation between quiescent and proliferating samples across a

variety of cancer types and models including cancer cell lines, 3D organoid cultures, circulating

tumour cells and patient-derived xenografts (Supplementary Table 1), thereby demonstrating its

broad applicability. Furthermore, the strength of the score appeared to reflect the quiescence

depth[37] (Figure 1d).

We experimentally validated our methodology in nine lung adenocarcinoma cell lines. There was

a remarkably good correlation between our predicted quiescence levels and the fraction of

quiescent cells, estimated using quantitative, single-cell imaging of phospho-Ser807/811-Rb

(PRb, which labels proliferative cells[38]) and 24 hour EdU proliferation assays (Figure 1e-h). Cells

that were negative for either PRb or EdU were defined as quiescent (see Methods, Figure 1e-f).

Importantly, a G1 signature (Methods) was not associated with these experimental

measurements, proving our method indeed captures quiescence rather than a prolonged G1

state (Figure 1g-h). The quiescence correlations appeared robust to random removal of individual

genes from the signature, with no single gene having an inordinate impact on the score

(Supplementary Figure 1c-e). This provided further reassurance that our Z-score based

methodology is successful in capturing dormancy signals from bulk tumour data.

Having established a robust framework for quantifying tumour dormancy, we next profiled 8,005

primary tumour samples across 31 solid cancer tissues from The Cancer Genome Atlas (TCGA).

After accounting for potential confounding signals from the microenvironment by correcting for

tumour purity (see Methods, Supplementary Figure 1f-g), we observed an entire spectrum of

dormancy across tumours that ranged from highly dormant to highly proliferative (Figure 2a).

Cancers well known to be frequently dormant, such as glioblastoma[3,30], were amongst the

highest ranked in terms of quiescence, along with kidney and adrenocortical carcinomas (Figure

2b).

As expected, tissues with lower stem cell division rates presented a greater propensity for

quiescence (Figure 2c) [39]. Furthermore, we confirmed that our quiescence scores capture a

cellular state that is distinct from that of cancer stem cells, marked by signatures associated with

high telomerase activity and an undifferentiated state[40,41], and that of senescent cells, marked by

the Senescence-Associated Secretory Phenotype (SASP) and β-galactosidase activity[42-44]

(Figure 2d, Supplementary Figures 2a-d). Our scores also showed strong negative correlations

with the expression of proliferation markers (Figure 2d).

Lastly, we confirmed dependencies on the p53/p21/DREAM activation axis: tumours that were

proficient in *TP53* or the components of the DREAM complex, as well as those with higher p21

expression, had elevated dormancy levels across numerous tissues (Figure 2e, Supplementary

Figures 2e-f), although only 8 out of 139 dormancy genes are directly transcriptionally regulated

by p53[45]. Nevertheless, p53 proficiency appears to be a non-obligatory dependency of dormancy,

which is also observed to arise in p53 mutant scenarios in 21% of cases. p53 has also been

shown to play a role in preventing the occurrence of larger structural events and polyploidy[46-48],

potentially explaining the lower dormancy levels we observed in tumours that had undergone

whole genome duplication (Figure 2f).

The genomic background of cancer dormancy

After obtaining dormancy estimates for primary tumour samples, we set out to identify potential

genomic triggers which might create a favourable background for tumour dormancy. We

identified 285 cancer driver genes that were preferentially altered (via mutations or copy number

alterations) either in dormant or fast proliferating tumours (Figure 3a). Reassuringly, this list

included genes previously implicated in driving cellular quiescence-proliferation decisions such

as *TP53* and *MYC*[16,19]. An additional 10 cancer genes showed signals of positive selection in the

context of high dormancy/proliferation (Supplementary Table 2). We also investigated

associations with mutagenic footprints of carcinogens (termed "mutational signatures"), which

can be identified as trinucleotide substitution patterns in the genome[49,50]. 15 mutational

signatures were linked with dormancy either within individual cancer studies or pan-cancer

(Supplementary Figure 2g).

Following the initial prioritisation of putative genomic triggers of dormancy, we sought to identify

those that could best distinguish dormant from fast proliferating tumours, while accounting for

tissue effects. An ensemble elastic net selection approach similar to the one described by Pich

et al[51] was applied for this purpose (Figure 3b, Methods). Our pan-cancer model identified tissue

type to be a major determinant of quiescence levels (Supplementary Figure 3a). It also uncovered

a reduced set of 60 genomic events linked with tumour dormancy levels, including SNVs and

copy number losses in 17 cancer genes, as well as amplifications of 13 cancer genes (Figure

3c). These events could then be successfully employed to predict quiescence in a separate test

dataset, thus internally validating our model (Supplementary Figure 3b).

Reassuringly, a functional *TP53*, lack of *MYC* amplification and lower mutation rates (Figure 3c)

were amongst the top ranked characteristics of dormant tumours, which also displayed less

aneuploidy. Additionally, we uncovered novel dependencies, such as *CEP89* and *LMNA*

amplifications observed in fast cycling tumours, or *ZMYM2* deletions prevalent in dormant

samples. p16 (*CDKN2A*) deletions, one of the frequent early events during cancer evolution[52,53],

were enriched in dormant tumours. *RB1* deletions and amplifications were both associated with

a reduction in tumour dormancy, which might reflect the dual role of RB1 in regulating proliferation

and apoptosis[54]. Overall, the genomic dependencies of dormancy mainly comprised genes

involved in cell cycle pathways, p53 regulation and ubiquitination (most likely of cell cycle

targets), TGF-β signalling mediators and RUNX2/3 regulation which have previously been shown

to play a role in controlling proliferation and cell cycle entry (Supplementary Figure 3c).

Our model also calls attention to the broader mutational processes associated with this cellular

state. Such processes showed fairly weak and heterogeneous correlations with dormancy within

individual cancer tissues (Supplementary Figure 2g), but their contribution becomes substantially

clearer pan-cancer once other genomic sources are accounted for. In particular, we identified an

association between dormancy and mutagenesis induced by the AID/APOBEC family of cytosine

deaminases as denoted by signature SBS2[49] (Figure 3c). As highlighted by Mas-Ponte and

Supek[55], APOBEC/AID driven mutations are likely to have deleterious functional consequences

as they tend to be directed towards early-replicating, gene-rich regions of the genome. The

authors reported an enrichment of APOBEC-generated mutations in several genes including

ZMYM2, which our pan-cancer model has linked with higher dormancy.

In turn, defective DNA mismatch repair, as evidenced by signatures SBS44, SBS20, SBS15,

SBS14 and SBS6[49], was enriched in fast cycling tumours (Figure 3c). Mismatch repair

deficiencies lead to hypermutation of single nucleotides and short tandem repeats due to

unrepaired DNA polymerase slippage in a phenomenon termed "microsatellite instability" (MSI).

Cancers that are particularly prone to MSI include colon, stomach and endometrial carcinomas[56],

where we could indeed observe differences in quiescence levels based on the MSI status (Figure

3d). Furthermore, dormant tumours were depleted of alterations across all DNA damage repair

pathways (Figure 3e).

Our measurements of dormancy also reflected expected cycling patterns across 27 MCF7

strains[57]: cell lines with longer doubling times exhibited higher quiescence (Figure 3f). This also

coincided with a depletion of *PTEN* mutations, a dependency highlighted by the pan-cancer

model.

When checking for dependencies in individual cancer tissues, 27 out of the 28 genes identified

by the model were significantly associated with dormancy in at least one tissue, most prominently

in breast, lung and liver cancers which also represent the largest studies within TCGA (Figure

3g, top panel). Most of these genomic insults were linked with a decrease in dormancy. In

external validation datasets these associations, including deletions in *PTEN* and *LRP1B* or

amplifications of *MYC, CEP89* or *STAG1,* featured most prominently in the largest cohort of

breast cancer samples (Figure 3g, bottom panel).

CEP89 amplification contributes to high tumour proliferation rates

To gain more insight into the underlying biology of cellular dormancy, we sought to experimentally

validate associations highlighted by the pan-cancer model. We focused on the impact of *CEP89*

activity on quiescence-proliferation decisions due to the high ranking of this putative oncogene

in the model, the relatively unexplored links between *CEP89* and cell cycle control, as well as its

association with quiescence across a variety of cancer cell lines (Supplementary Figure 4a-c).

The function of CEP89 is not well characterised, however, the encoded protein has been

proposed to function as a centrosomal-associated protein[58,59]. Centrosomes function as major

microtubule-organising centres in cells, playing a key role in mitotic spindle assembly[60] as well

as in the mitotic entry checkpoint by providing a platform for concentrating and amplifying signals

of cell cycle proteins such as cyclin B/Cdk1[61]. Moreover, centrosomes function as sites of

ubiquitin-mediated proteolysis of cell cycle targets[62] and members of several growth signalling

pathways, such as Wnt and NF-kB, have been shown to localise at these structures[63,64]. Several

genetic interactions have also been reported between CEP89 and key cell cycle proteins,

including cyclin D2[65] (Figure 4a).

Our model linked *CEP89* amplification with fast cycling tumours (Figure 3c). Centrosome

amplification is a common feature of tumours with high proliferation rates and high genomic

instability[66]. This is reflected by the strong negative correlation we found between quiescence

levels and a previously reported centrosome amplification transcriptomic signature (CA20) in

TCGA (Figure 4b). Previous studies have shown that overexpression of centrosomal proteins

can lead to alterations in centriole structure[67,68] and we observed significantly higher CA20 scores

in tumours containing *CEP89* amplifications (Figure 4c). Furthermore, *CEP89* expression was

prognostic across multiple cancer tissues (Figure 4d) and linked with toxicity of several cancer

compounds in cell line models (Supplementary Figure 4d).

We validated this target in the lung adenocarcinoma cell line NCI-H1299 showing high levels of

CEP89 amplification. Cep89 depletion via siRNA knockdown caused a consistent increase in the

fraction of quiescent cells as measured by PRb and EdU assays (Figure 4e-f). Thus, we propose

CEP89 as a novel cell proliferation regulator that may be exploited in certain scenarios to control

tumour growth.

Dormancy in breast cancer

Due to the variation of dormancy levels across tissues and to increase power, we have taken a

pan-cancer approach to capture genomic features that are universally associated with dormancy.

However, we acknowledge that the extremely diverse genomic background for cancer

development in different tissues might mean that certain genetic alterations may lead to higher

or lower propensity of dormancy in a single tissue only. To investigate this further we focused on

breast cancer, the largest cancer dataset available from TCGA, where we still have adequate

power and balance of dormant/cycling phenotypes to uncover tissue-specific genomic markers

of dormancy. A combined ANOVA and random forest classification approach identified 30

genomic events that could distinguish dormant from fast cycling tumours with a mean accuracy

of 74% (Figure 5a). This model recovered the associations with the APOBEC mutational

signature SBS2, *TP53* mutation as well as *MYC*, *KLF6*, *LMNA*, *ETV6* and *RAD21* amplification

already seen in the pan-cancer model (Figure 5b). All additional associations validated in the

METABRIC cohort and were also seen in several other cancers, particularly bladder, lung and

lower grade glioma (Figure 5c).

Since APOBEC/AID activity has been linked with dormancy in both the pan-cancer and the

tissue-specific models, we further investigated its role in breast cancer, where the associated

mutation signature SBS2 is relatively prevalent. A significant correlation (Pearson R = 0.27, p =

3.5e-07) was observed between the fraction of mutations contributed by the SBS2 process and

dormancy in this cohort. Breast cancer samples enriched in APOBEC-associated mutations,

defined as mutations attributed to SBS2 and SBS13, presented significantly higher quiescence

levels (Figure 5d). Across breast cancer subtypes, the Luminal A subtype showed the highest

levels of dormancy, however, Her2+ tumours had the highest incidence of APOBEC associated

mutational events, suggesting this subtype may be the most susceptible to APOBEC-linked

quiescence (Figure 5e-f).

Characterisation of individual quiescence programmes

While we had previously examined a generic dormancy programme, cancer cells can enter

quiescence due to different stimuli[1] and this may inform its aetiology and manifestation. To

explore this, we re-scored tumours based on gene expression programmes specific to serum

starvation, contact inhibition, MEK inhibition, CDK4/6 inhibition or spontaneously occurring

quiescence as defined by Min and Spencer[18] (see Methods). We observed a good correlation

between the dormancy estimates representing individual quiescence programmes and the

expression of genes associated with the corresponding form of quiescence in the literature

(Figure 6a-e, Supplementary Figure 5). CDK4/6 inhibition-induced quiescence levels were further

validated using external RNA-seq datasets from cancer cell lines and xenograft mice sequenced

before and after treatment with the CDK4/6 inhibitor Palbociclib[69,70] (Figure 6f, Supplementary

Table 4). Interestingly, we also observed significant differences in spontaneous quiescence

scores before and after treatment. Indeed, p21 activity has been linked with the Palbociclib

mechanism of action[71,72], and this analysis suggests potential similarities between CDK4/6

inhibition and p21-dependent quiescence phenotypes.

Having validated our framework for quantifying stimulus-specific quiescence programmes, we

proceeded to estimate the dominant form of quiescence in different cancer types (Figure 6g). We

found a range of quiescence aetiologies across most tissues, while a minority of cancers were

dominated by a single form of quiescence, e.g. serum starvation in all dormant

pheochromocytomas and paragangliomas, contact inhibition in 88% of head and neck

carcinomas and CDK4/6 inhibition in 80% of dormant adrenocortical carcinomas. While we do

not wish to claim that the dormancy state will have necessarily been induced by the actual

predicted stimulus (impossible in the case of CDK4/6 or MEK inhibition, as the analysed samples

are all treatment-naïve), we suggest that the downstream signalling cascade may resemble that

triggered by such stimuli, e.g. via CDK4/6 or MEK loss of function mutations. Amongst these

states, spontaneous quiescence appeared most strongly dependent on p53 functionality

(Supplementary Table 3). This points to common transcriptional features of drug-tolerant

dormant cells in different cancer settings that could be employed in designing ways to eradicate

these cells in the future.

Clinical relevance of tumour dormancy

Overall, dormancy appears to be beneficial for the long-term outcome of cancer patients (Figure

7a, Supplementary Figure 6a). As expected, quiescence levels were increased in stage 1

tumours, although later stages also exhibited occasional elevated dormancy (Supplementary

Figure 6b). Indeed, such dormant, indolent tumours could have higher chances of being

eradicated earlier in the disease, which is consistent with reported worse prognosis of patients

with higher tumour cell proliferation rates and less stable, more mutated genomes[73]. However,

outcomes do vary depending on the quiescence subtype, with worse survival observed upon

contact inhibition (Figure 7b). The outcomes also vary by tissue, with lung, colon or esophageal

carcinoma patients displaying significantly worse prognosis in the context of dormancy (Figure

7c).

While dormancy may confer an overall survival advantage in most cancers, it can also provide a

pool of cells that are capable of developing resistance to therapy, as demonstrated in several

studies[34,74]. Using our methodology, we indeed observed an increase in quiescence levels in cell

lines following treatment with EGFR, BRAF and CDK4/6 inhibitors, as well as conventionally used

chemotherapies such as 5-Fluorouracil (5-FU) in multiple bulk RNA-seq datasets (Figure 7d).

Furthermore, the recent widespread availability of single-cell transcriptomics offers the

opportunity to investigate the impact of quiescence on such therapies with much greater

granularity than is allowed by bulk data. Using our quiescence signature and single-cell data from

RKO and SW480 colon cancer cell lines treated with 5-FU[75], we could observe quiescence-

proliferation decisions following conventional chemotherapy treatment. Within the p53 proficient

cell line RKO, the fraction of quiescent cells increased from 41% to 93% after treatment with a

low dose (10 µM) of 5-FU and persisted at higher doses (Figure 7e-f). In contrast, a comparable

increase was not observed in *TP53* mutant SW480 cells, further emphasizing the key role of p53

as a regulator of quiescence (Figure 7g-h). SW480 cells showed higher apoptotic activity

following treatment compared to RKO cells, particularly within actively cycling cells, further

corroborating that cells capable of entering quiescence may be less vulnerable to this therapy

(Supplementary Figure 6c-d).

Similarly, using single cell data from an *EGFR* mutant Non-Small Cell Lung Cancer (NSCLC) cell

line treated with the EGFR inhibitor Erlotinib[76], we predicted that 40% of cells were likely to exist

in a dormant state prior to treatment. EGFR inhibition led to a massive decrease in cell numbers

immediately after treatment, mostly due to proliferating cells dying off (Supplementary Figures

6e-f), while the proportion of quiescent cells increased to 96% at day 1, indicating an immediate

selective advantage for such cells (Figure 7i-j). Interestingly, these cells appear to gradually start

proliferating again in the following days during continuous treatment, with the percentage of

proliferating cells approaching pre-treatment levels by day 11 (Figure 7j). The same trend

captured by our signature could be observed upon KRAS and BRAF inhibition in different cell

line models (Supplementary Figure 6g-j, Supplementary Table 4)[34,76]. Furthermore, during the

first days of treatment the NSCLC cells that survived EGFR inhibition appeared to reside in a

state most resembling that induced by serum starvation (Figures 7k-l). Both EGFR kinase

inhibitors and serum starvation have been shown to trigger autophagy[77], which may explain the

convergence between this inhibitory trigger and the type of quiescence response. At day 11 most

of the remaining dormant cells appeared in a state similar to that preceding the treatment (Figure

7l).

Thus, quiescence appears to explain resistance to broad acting chemotherapy agents as well as

targeted molecular inhibitors of the Ras/MAPK signalling pathway, being either selected for, or

induced immediately upon treatment, and gradually waning over time as cells start re-entering

the cell cycle. Using massively multiplexed chemical transcriptomic data, we also analysed

responses to 188 small molecule inhibitors in cell lines at single-cell resolution[78] (Supplementary

Figure 7). We observed a large increase in quiescence following treatment with compounds

targeting cell cycle regulation and tyrosine kinase signalling, consistent with our previous results,

but also for compounds modulating epigenetic regulation, e.g. histone deacetylase inhibitors -

thus highlighting the broad relevance of quiescence.

These analyses showcase our ability to probe the nature of quiescent phenotypes using a defined

dormancy signature, which could aid the development of methods to selectively target quiescent

drug-resistant persister cells. However, a major challenge of scRNA-seq data analysis is the high

percentage of gene dropout, which could impact our ability to evaluate dormancy using the full

139 gene signature. The scRNA-seq datasets we employed to investigate responses to common

therapeutic agents exhibited an average drop-out of 10.6 genes out of the full gene signature.

While our scoring method remains robust to such levels of dropout (Supplementary Figure 1c-

e), we also employed machine learning to reduce our initial list of 139 markers of dormancy to a

robust 35-gene signature, comprised mainly of RNA metabolism and splicing-regulating factors,

but also of genes involved in cell cycle progression, ageing and senescence, which could be

applied to sparser datasets with larger levels of gene dropout (Methods, Figure 8a-b). The

optimised signature of dormancy performed similarly to the initial broadly defined programme in

distinguishing highly dormant and fast cycling tumours (Figure 8c), showed no dropout across

the scRNAseq datasets used in this study (Figure 8d), was similarly prognostic (p=0.006) and

showed comparable profiles of resistance to treatment (Figure 8e, Supplementary Figure 8). This

minimal expression signature could be employed to track and further study emerging

quiescence-induced resistance in a variety of therapeutic scenarios.

322
DISCUSSION

Despite its crucial role in cancer progression and resistance to therapies, dormancy has

remained poorly characterised due to the scarcity of suitable models and biomarkers for large-

scale tracking in the tissue or blood. Lack of proliferative markers such as Ki67 or CDK2[21,79] does

not uniquely distinguish quiescence from other cell cycle phases, e.g. G1 or S. This study

overcame this limitation by employing genes active in different forms of quiescence whose

patterns of expression reflect this specific state and are distinct from markers of a longer G1

phase, senescence or stemness. We demonstrate a robust method of profiling them to identify

dormancy in bulk and single cell RNA-seq data. Since our list of quiescence biomarkers is defined

using expression data from human epithelial cells, our approach could be limited in terms of

applicability to other cell types. However, carcinomas originating from epithelial tissues make up

85% of all malignancies. Moreover, our approach had high classification accuracy in a variety of

datasets encompassing solid as well as haematological cancers, thus demonstrating the

versatility of our method. More variable performance was observed when applied to

hematopoietic stem cells as it was not designed to capture signals in this context (Supplementary

Figure 1b).

We show that dormancy is pervasive across different solid cancers and generally associated with

more stable, less mutated genomes with intact DNA damage repair pathways. We also find a link

between APOBEC mutagenesis and higher levels of dormancy. We identify mutational events

affecting a variety of genes such as *PTEN*, *CEP89, CYLD*, *LMNA* that appear unfavourable to

dormancy development, thus potentially implicating them in influencing quiescence-proliferation

decisions. Among these, we propose *CEP89* amplification as novel biomarker of non-small cell

lung primary tumours with low quiescence capacity. Genomic alterations enriched in dormant

tumours, such as p16 or *ZMYM2* deletions, were more rare and predominantly captured pan-

cancer. Such targets should be further validated and could be exploited to either counteract

dormancy or induce it.

Our finding that primary tumour dormancy confers a favourable prognosis could raise a potential

limitation of our approach in distinguishing between slow-dividing and persister cell phenotypes[6]

in bulk transcriptome data. Nevertheless, our methodology is directly extendable to single cell

data, where we demonstrate a clear association between existent/emerging quiescence and

resistance to multiple cell cycle, kinase signalling and epigenetic targeting regimens in a variety

of cancers. Quiescent cells utilise a variety of mechanisms to achieve drug resistance, including

upregulation of stress-induced pathways such as anti-apoptotic BCL-2 signalling[80], anti-ROS

programmes[18] or immune evasion[10]. Further studies are needed to elucidate if these

mechanisms are specific to individual cancer tissues or individual quiescence types.

Overall, our study provides a pan-cancer view of dormancy and its evolutionary constraints,

underlying novel mutational dependencies which could be exploited in the clinic. We propose a

tumour dormancy signature which can be measured in bulk tissue or single cells, and could

inform therapeutic strategies or risk of relapse. This signature could be assessed in the clinic to

track rapidly emerging resistance, e.g. through liquid biopsies or targeted gene panels. We hope

these insights can be used as building blocks for future studies into the different regulators of

dormancy, including epigenetics and microenvironmental interactions, as well as the

mechanisms by which it enables therapeutic resistance both in solid and haematological

malignancies.

367
METHODS

Selection of quiescence marker genes

Generic quiescence markers:

Differential expression analysis results comparing cycling immortalised, non-transformed human

epithelial cells and cells in five different forms of quiescence (spontaneous quiescence, contact

inhibition, serum starvation, CDK4/6 inhibition and MEK inhibition) were obtained from Min and

Spencer[18]. A total of 195 genes were differentially expressed in all five forms of quiescence under

an adjusted p-value cut-off of 0.05. This gene list, reflective of a generic quiescence phenotype,

was subjected to the following refinement and filtering steps: (1) selection of genes with a

unidirectional change of expression across all five forms of quiescence; (2) removal of genes

involved in other cell cycle stages included in the "KEGG_CELL_CYCLE" gene list deposited at

MSigDB; (3) removal of genes showing low standard deviation and low levels of expression within

the TCGA dataset, or which showed low correlation with the pan-cancer expression of the

transcriptional targets of the DREAM complex, the main effector of quiescence, in TCGA.

Quiescence subtype-specific markers:

Gene lists representing spontaneous quiescence, contact inhibition, serum starvation, CDK4/6

inhibition and MEK inhibition programmes were obtained using genes differentially expressed in

each individual quiescence form using an adjusted p-value cutoff of 0.05. The gene lists were

subjected to filtering steps 2 and 3 described above. Following the refinement steps, 10

upregulated and 10 downregulated genes with highest log2 fold changes were selected for each

quiescence type.

Dormancy quantification

The *GSVA* R package was used to implement the combined z-score[26], ssGSEA[27] and GSVA[28]

gene set enrichment methods. For the above three methods a separate score was obtained for

genes upregulated in quiescence and genes downregulated in quiescence, following which a

final quiescence score was obtained by subtracting the two scores. The singscore single-sample

gene signature scoring method[29] was implemented using the *singscore* R package. In addition

to these, we also calculated a mean scaled quiescence score based on the refined list of genes

upregulated and downregulated in quiescence, as well as a curated housekeeping genes from

the "HSIAO_HOUSEKEEPING_GENES" list deposited at MSigDB, as follows:

$$QS = \frac{\frac{1}{n}\sum G_U - \frac{1}{n}\sum G_D}{\frac{1}{n}\sum G_H}$$

QS = mean scale quiescence score

G_U = expression of genes upregulated in quiescence

G_D = expression of genes downregulated in quiescence

G_H = expression of housekeeping genes

n = number of genes in each gene set

Quiescence scores for the TCGA cohort were derived from expression data scaled by tumour

purity estimates. The pan-cancer TCGA samples were also classified into "high" or "low"

quiescence groups based on k-means clustering (k=2) on the expression data of 139 quiescence

biomarker genes, following the removal of tissue-specific expression differences using the

ComBat function from the *sva* R package[81].

Measuring the depth of quiescence

We employed the GSE124109 dataset from Fujimaki et al[37] where rat embryonic fibroblasts were

transcriptomically profiled as they moved from shallow to deep quiescence over time in the

absence of growth signals. The derived quiescence scores using our combined z-score

methodology increased from shallow to deep quiescence.

Validation of quiescence scoring methodologies

Single-cell RNA-sequencing validation datasets:

Datasets were obtained from the ArrayExpress and Gene Expression Omnibus (GEO) databases

though the following GEO Series accession numbers: GSE83142, GSE75367, GSE137912,

GSE139013, GSE90742 and E-MTAB-4547. Quality control analysis was standardised using the

SingleCellExperiment[82] and *scater*[83] R packages. Normalisation was performed using the *scran*[84]

R package.

Bulk RNA-sequencing validation datasets:

Datasets were obtained from the GEO database through the following GEO Series accession

numbers: GSE93391, GSE114012, GSE131594, GSE152699, GSE124854, GSE135215,

GSE99116 and GSE124109. GSE114012 count data were normalised to TPM values using the

GeoTcgaData R package. All normalised datasets were log-transformed before further analysis.

The accuracy with which the quiescence scoring methods could separate proliferating and

quiescent samples within the validation datasets was determined by calculating the area under

the curve of the receiver operating characteristic (ROC) curves, using the *plotROC* R package.

Experimental validation in lung adenocarcinoma cell lines

The average fraction of cancer cells spontaneously entering quiescence was estimated for nine

lung adenocarcinoma cell lines (NCIH460, A549, NCIH1666, NCIH1944, NCIH1563, NCIH1299,

NCIH1650, H358, L23) using EdU and phospho-Rb staining proliferation assays.

Cell lines were obtained from ATCC or Sigma and regularly checked for mycoplasma. A549 and

NCIH460 were cultured in DMEM (Gibco). NCIH358, NCIH1299 and NCIH1563 were maintained

in RPMI-1640 (Gibco) supplemented with 5mM sodium pyruvate and 0.5% glucose. NCIH1944,

NCIH1666, NCIH1650 and L23 were grown in RPMI-1640 ATCC formulation (Gibco). A427 were

cultured in EMEM (ATCC). A549, NCIH460, H358, NCIH1299, NCIH1563, A427 were

supplemented with 10% heat inactivated FBS. NCIH1666 with 5% heat-inactivated FBS and all

other cell lines with 10% non-heat inactivated FBS. All cell lines had penicillin-streptomycin

(Gibco) added to 1%. Cells were maintained at 37°C and 5% CO_2. To calculate the quiescent

fraction, A549 and NCIH460 cells were plated at a density of 500 cells/well, and all other cell

lines at a density of 1000/well, in 384well CellCarrier Ultra plates (PerkinElmer) in the relevant

media. 24h later, 5μM EdU was added and cells were incubated for a further 24h before fixing in

a final concentration of 4% formaldehyde (15 min, RT), permeabilization with PBS/0.5% Triton

X-100 (15 min, RT) and blocking with 2% BSA in PBS (60 min, RT). The EdU signal was detected

using Click-iT chemistry, according to the manufacturer's protocol (ThermoFisher). Cells were

also labelled for phospho-Ser807/811 Rb (PRb) using Rabbit mAb 8516 (CST) at 1:2000 in

blocking solution, overnight at 4°C. Unbound primary antibody was washed three times in PBS

and secondary Alexa-conjugated antibodies were used to detect the signal (ThermoFisher,

1:1000, 1h at RT). Finally nuclei were labelled with Hoechst 33258 (1 μg/ml, 15 min RT) before

imaging on a high-content widefield Operetta microscope, 20x N.A. 0.8. Automated image

analysis (Harmony, PerkinElmer) was used to segment and quantify nuclear signals in imaged

cells. Quiescent cells were defined by the absence of EdU or PRb staining, determined by

quantification of their nuclear expression (Figure 1e-f).

The quiescence scores for cancer cell lines were calculated using corresponding log-transformed

RPKM normalised bulk RNA-seq data from the Cancer Cell Line Encyclopedia (CCLE)

database[85].

CEP89 was depleted by ON-Target siRNA Pool from Horizon. NCI-H1299 cells were reverse

transfected in 384 well plates with 20nM of Non-targeting control (NTC) or CEP89-targeting

siRNA using Lipofectamine RNAiMax (ThermoFisher), according to the manufacturer's

instructions. Cells were left for 24h, before 5μM EdU was added for the final 24h and then cells

were processed as above to determine the quiescent fraction. To determine the level of Cep89

depletion by western blot, cells were reverse transfected with siRNA in 24 well plates. 48h after

transfection, cells were lysed directly in 1x SDS sample buffer with 1mM DTT (ThermoFisher).

Samples were separated on pre-cast 4-20% Tris-Glycine gels, transferred to PVDF using the

iBlot2 system and membranes blocked in blocking buffer (5% milk in TBS) for 1h at RT. The

membrane was then cut and the upper half was incubated in 1:1000 Cep89 antibody (Sigma,

HPA040056), the bottom half in B-actin antibody 1:2000 (CST; 3700S) diluted in blocking buffer

overnight at 4'C. Membranes were washed three times in TBS-0.05% TritonX-100 before being

incubated in secondary anti-rabbit (Cep89) or anti-mouse (B-actin) HRP conjugated antibodies

(CST 7074P2 and CST 7076P2, respectivelt) diluted 1:2000 in blocking buffer for 1h at RT.

Membranes were washed three times again and signal detected using Clarity ECL solution

(BioRad) and scanned on an Amersham ImageQuant 800 analyser.

Multi-omics discovery cohort

FPKM normalised RNA-sequencing expression data, copy number variation gene-level data, as

well as mutation annotation files aligned against the GRCh38 human reference genome from the

Mutect2 pipeline were downloaded using the *TCGABiolinks* R package [86] for 9,712 TCGA

primary tumour samples across 31 solid cancer types. Haematological malignancies were

excluded as the quiescence markers were derived in epithelial cells and might not be equally

suited to capture this phenotype in blood. For patients with multiple samples available, one RNA-

seq barcode entry was selected for each individual patient resulting in 9,631 total entries. All

expression data were log-transformed for downstream analysis. During quiescence score

calculation, expression data for the primary tumour samples was scaled according to tumour

purity estimates reported by Hoadley et al[87] to account for potential confounding quiescence

signals coming from non-tumour cells in the microenvironment. Samples with purity estimates

lower than 30% were removed, leaving 8,005 samples for downstream analysis.

The mutation rates of all TCGA primary tumour samples were determined by log-transforming

the total number of mutations in each sample divided by the length of the exome capture (38Mb).

TP53 functional status was assessed based on somatic mutation and copy number alterations

as described in Zhang et al[88]. *TP53* mutation and copy number for the TCGA tumours were

downloaded from cBioPortal (http://www.cbioportal.org). Tumours with *TP53* oncogenic

mutations (annotated by OncoKB) and copy-number alterations (GISTIC score ≤ -1) were

assigned as *TP53* mutant and CNV loss. Tumours without these *TP53* alterations were assigned

as TP53 wild type. The effects of the *TP53* mutation status on quiescence score were then

determined with a linear model approach with the quiescence score as a dependent variable and

mutational status as an independent variable. The *P* values were FDR-adjusted.

APOBEC mutagenesis enriched samples were determined through pan-cancer clustering of

mutational signature contributions as described in Wiecek et al[89]. The APOBEC mutagenesis

cluster was defined as the cluster with highest mean SBS2 and SBS13 contribution. This was

repeated 100 times and only samples which appeared in the APOBEC cluster at least 50 times

were counted as being APOBEC enriched.

Aneuploidy scores and whole genome duplication events across TCGA samples were obtained

from Taylor et al[90]. Microsatellite instability status for uterine corpus endometrial carcinoma, as

well as stomach and colon adenocarcinoma samples were obtained from Cortes-Ciriano et al[56].

Telomerase enzymatic activity "EXTEND" scores were obtained from Noureen et al[40].

Expression-based cancer cell stemness indices were obtained from Malta et al[41]. Centrosome

amplification transcriptomic signature (CA20) scores were obtained from Almeida et al[66].

PHATE dimensionality reduction

509 The *phateR* R package[91] was used to perform the dimensionality reduction with a constant seed

510 for reproducibility. The *ComBat* function from the *sva* R package[92] was used to remove tissue-

511 specific expression patterns from the TCGA RNA-seq data.

Cancer stem cell division estimates

513 The mean stem cell division estimates for different cancer types used in this study were obtained

514 from Tomasetti and Vogelstein[39].

Positive selection analysis

516 Genes positively selected specifically in samples clustered into low or high dormancy groups

517 were identified based on dN/dS analysis using the *dNdScv* R package[93], run with default

518 parameters. Genes showing signals of positive selection in either the high dormancy or low

519 dormancy samples which encoded olfactory receptors were discarded from downstream

520 analysis.

Mutational signature estimation

Mutational signature contributions were inferred as described in Wiecek et al[89].

Ensemble elastic net regression model

The COSMIC database was used to source a list of 723 known drivers of tumorigenesis (Tiers 1+2). 285 oncogenes and tumour suppressors from a curated list showed a significant enrichment or depletion of mutations or copy number variants in dormant samples either pan-cancer or within individual TCGA studies.

To classify dormant from fast proliferating tumours, the 286 genes were used as input features for an ensemble elastic net regression model along the tumour mutational rate, whole-genome doubling estimates, ploidy, aneuploidy scores, 10 positively selected genes and 15 mutational signatures, which showed a significant correlation with quiescence scores either pan-cancer or within individual TCGA studies. The *caret* R package was used to build an elastic net regression model 1000 times on the training dataset of 3,753 TCGA primary tumour samples (80% of the total dataset). Only samples with at least 50 mutations were used in the model, for which mutational signatures could be reliably estimated. For each of the 1000 iterations, we randomly selected 90% of the samples from the training dataset to build the model. Only features which were included in all 1000 model iterations were selected for further analysis. To test the performance of our approach, a linear regression model was built using the reduced list of genomic features and their corresponding coefficients averaged across the 1000 elastic net regression model iterations. When applying the resulting linear regression model on the internal validation dataset of 936 samples, we found a strong correlation between the observed and predicted quiescence scores (R = 0.73, p < 2.2e-16).

SHAP values for the linear regression model used to predict quiescence scores were obtained using the *fastshap* R package.

Gene enrichment and network analysis

Gene set enrichment analysis was carried out using the *ReactomePA* R package, as well as GeneMania[94] and ConsensusPathDB[95]. Interactions between CEP89 and other cell cycle components were inferred using GeneMania. Networks were visualised using Cytoscape[96].

Gene lists

Genes associated with the G1 phase of the cell cycle were obtained from the curated "REACTOME_G1_PHASE " list deposited at MSigDB.

Genes associated with apoptosis were obtained from the curated "HALLMARK_APOPTOSIS" list deposited at MSigDB.

Genes associated with the senescence-associated secretory phenotype were obtained from Basisty et al[44]. Lists of genes making up the various DNA damage repair pathways were derived from Pearl et al[97].

Genes associated with contact inhibition were obtained from the curated "contact inhibition" gene ontology term. Genes associated with serum starvation were obtained from the curated "REACTOME_CELLULAR_RESPONSE_TO_STARVATION" list deposited at MSigDB. MEK inhibition was assessed based on the activity of the MAPK pathway as determined using an expression signature (MPAS) consisting of 10 downstream MAPK transcripts[98].

Validation of genomic features of dormancy

For elastic net model feature validation, RNA-seq data was downloaded for 6 cancer studies from cBioPortal[99], along with patient-matched whole-genome, whole-exome and targeted sequencing data. The 6 datasets used comprise breast cancer (SMC[100] and METABRIC[101]), paediatric Wilms' tumor (TARGET[102]), bladder cancer, prostate adenocarcinoma and sarcoma (MSKCC[103-105]) studies. The data were processed and analysed in the same manner as the TCGA data. RNA-

seq data for 27 MCF7 cell line strains, alongside cell line growth rates and targeted mutational

sequencing data were obtained from Ben-David et al[57].

Genomic dependency modelling in breast cancer

An ANOVA-based feature importance classification was used and identified 30 genomic features

most discriminative of samples with lower and higher than average quiescence scores. A random

forest model was then built using the identified features and correctly classified samples

according to their quiescence state with a mean accuracy of 74% across five randomly sampled

test datasets from the cohort.

Survival analysis

Multivariate Cox Proportional Hazards analysis was carried out using the *coxph* function from the

survival R package. The optimal quiescence score cut-off value of 2.95 was determined using

the *surv_cutpoint* function.

Treatment response scRNA-seq and bulk RNA-seq data

Datasets have been obtained from the GEO database through the following GEO Series

accession numbers: GSE134836, GSE134838, GSE134839, GSE137912, GSE149224,

GSE124854, GSE135215, GSE99116, GSE152699, GSE178839 and GSE139944. The *umap*

R package was used for dimensionality reduction with constant seed for reproducibility.

Quiescence subtype determination

TCGA cohort studies:

Samples with evidence of dormancy characterised by a generic quiescence score > 0 were

further subclassified based on the most likely form of quiescence exhibited, among CDK4/6

inhibition, contact inhibition, MEK inhibition, spontaneous quiescence or serum starvation, using

subtype-specific expression signatures. We opted for a conservative approach and classed each

dormant sample into a specific quiescence subtype if the quiescence score for the corresponding

programme was higher than one standard deviation of the distribution across the TCGA cohort,

and if the score was significantly higher than for the remaining programmes when assessed using

a Student's t test. Samples which could not be classified into any of the five quiescence states

characterised in this study were classified as "uncertain".

Single-cell RNA seq treatment response datasets:

The quiescence subtype of individual single cells was inferred by mapping such individual cells

onto the reference dataset of MCF10A cells reflecting different forms of quiescence obtained

from Min and Spencer[18]. The *ComBat* R package was used to remove the study batch effect

between the expression data to be classified and the reference bulk RNA-seq data. PCA

dimensionality reduction analysis was then used on the combined datasets using the *prcomp* R

function. For each patient sample or single-cell expression data entry, a k-nearest neighbour

algorithm classification was performed using the *knn* function from the *class* R package. During

the classification the three nearest reference bulk RNA-seq data points were considered, with

two nearest neighbours with identical class needed for classification.

Optimisation of dormancy signature

We investigated if a subset of the 139 quiescence-related genes could act as a more reliable

marker of dormancy that would bypass dropout issues in single cell data. This was performed in

three steps:

(1) Assessment of individual importance as quiescence marker for a given gene

We collected three high confidence single cell expression datasets separating quiescent from

dormant cells. A random forest model was trained on each dataset separately to predict the

quiescence state of a given cell based on the expression levels of the 139 genes. The Gini

indices corresponding to each gene in the model were normalised to a range of values

between 0 and 1, which would reflect how important an individual gene was for determining

quiescence state relative to the other 138 genes. The procedure was repeated for each of

the three datasets.

(2) Prioritisation of gene subsets based on cumulative importance in the model

Genes were placed in the candidate subset if their importance metric was above a given

threshold in at least one of the datasets. By gradually increasing the threshold from 0 to 1,

40 different gene combinations were produced.

(3) External validation of candidate subsets

The gene combinations in (2) were tested for their ability to predict quiescence. For this, a

separate validation dataset was utilized, which contained gene expression levels for the 139

genes in the 10 lung cancer cell lines previously employed for experimental validation, along

with the quiescence state of the lines as inferred by PRb and EdU staining. For each gene

subset, a combined z-score of quiescence was calculated from the expression levels as

described previously. The correlations between this z-score and the three experimental

measurements of quiescence were used to establish the ability of a gene combination to

predict quiescence. The best performing subset with a mean correlation of 80% between

predicted and measured quiescence levels in the test data (p=0.01) comprised 35 genes.

The optimised gene signature is available at https://github.com/secrierlab/CancerDormancy.

Data visualization and basic statistics

Graphs were generated using the ggplot2 and ggpubr R packages. Groups were compared using

the Student's t test, Wilcoxon rank-sum test or ANOVA, as appropriate.

Code

All code developed for the purpose of this analysis can be found at the following repository:

https://github.com/secrierlab/CancerDormancy

FUNDING

AJW and DHJ were supported by MRC DTP grants (MR/N013867/1). MPC was supported by an Academy of Medical Science Springboard award (SBF004\1042). GMT was supported by a Wellcome Seed Award in Science (215296/Z/19/Z). MS was supported by a UKRI Future Leaders Fellowship (MR/T042184/1). Work in MS's lab was supported by a BBSRC equipment grant (BB/R01356X/1) and a Wellcome Institutional Strategic Support Fund (204841/Z/16/Z). ARB and SC are supported by a CRUK CDF (C63833/A25729) and work in ARB's lab is supported by MRC core-funding to the London Institute of Medical Sciences (MC-A658-5TY60).

ACKNOWLEDGEMENTS

We would like to thank Prof Chris Barnes for the very helpful discussions and input on the findings of the study.

The results published here are in part based upon data generated by the TCGA Research Network: https://www.cancer.gov/tcga.

AUTHOR CONTRIBUTIONS

MS designed the study and supervised the computational analyses. ARB designed and supervised the experimental validation in cell lines. GB supervised the analysis of p53 functional association. AJW developed the dormancy scoring methodology and performed all computational analyses to validate and apply it in bulk and single cell datasets, as well as link it to genomic features. SC performed the experimental validation of quiescence prevalence and CEP89 association with quiescence in cell lines. DK performed the inference of the minimal signature of dormancy applicable in single cell data. MPC performed the random forest modelling and feature selection in breast cancer. LG performed the positive selection analysis. GMT wrote the code for batch effect correction and for PCA mapping of single cell data on a reference dataset. DHJ performed the APOBEC enrichment classification. PZ and LX performed the quiescence comparison of p53 wild type and mutated cancers. MS, AJW, ARB and SC wrote the

manuscript, with contributions from all other authors. All authors read and approved the

manuscript.

COMPETING INTEREST STATEMENT

None declared.

REFERENCES

1.   Coller, H.A., Sang, L. & Roberts, J.M. A new description of cellular quiescence. *PLoS Biol*

4, e83 (2006).

2.   van Dijk, D. *et al.* Slow-growing cells within isogenic populations have increased RNA

polymerase error rates and DNA damage. *Nat Commun* **6**, 7972 (2015).

3.   Chen, J. *et al.* A restricted cell population propagates glioblastoma growth after

chemotherapy. *Nature* **488**, 522-6 (2012).

4.   Puig, I. *et al.* TET2 controls chemoresistant slow-cycling cancer cell survival and tumor

recurrence. *J Clin Invest* **128**, 3887-3905 (2018).

5.   Rehman, S.K. *et al.* Colorectal Cancer Cells Enter a Diapause-like DTP State to Survive

Chemotherapy. *Cell* **184**, 226-242.e21 (2021).

6.   Phan, T.G. & Croucher, P.I. The dormant cancer cell life cycle. *Nat Rev Cancer* **20**, 398-

411 (2020).

7.   Cabanos, H.F. & Hata, A.N. Emerging Insights into Targeted Therapy-Tolerant Persister

Cells in Cancer. *Cancers (Basel)* **13**(2021).

8.   Masago, K., Fujita, S. & Yatabe, Y. Targeting minimal residual disease after surgery with

molecular targeted therapy: the real path to a cure? *J Thorac Dis* **10**, S1982-S1985

(2018).

9.   Malladi, S. *et al.* Metastatic Latency and Immune Evasion through Autocrine Inhibition of

WNT. *Cell* **165**, 45-60 (2016).

10.   Ribas, A. Adaptive Immune Resistance: How Cancer Protects from Immune Attack.

Cancer Discov **5**, 915-9 (2015).

11.   Sosa, M.S., Bragado, P. & Aguirre-Ghiso, J.A. Mechanisms of disseminated cancer cell

692   dormancy: an awakening field. *Nat Rev Cancer* **14**, 611-22 (2014).

693   12.   Barkan, D. *et al.* Metastatic growth from dormant cells induced by a col-I-enriched fibrotic

694   environment. *Cancer Res* **70**, 5706-16 (2010).

695   13.   Arora, M., Moser, J., Phadke, H., Basha, A.A. & Spencer, S.L. Endogenous Replication

696   Stress in Mother Cells Leads to Quiescence of Daughter Cells. *Cell Rep* **19**, 1351-1364

697   (2017).

698   14.   Barr, A.R. *et al.* DNA damage during S-phase mediates the proliferation-quiescence

699   decision in the subsequent G1 via p21 expression. *Nat Commun* **8**, 14728 (2017).

700   15.   Heldt, F.S., Barr, A.R., Cooper, S., Bakal, C. & Novák, B. A comprehensive model for the

701   proliferation-quiescence decision in response to endogenous DNA damage in human

702   cells. *Proc Natl Acad Sci U S A* **115**, 2532-2537 (2018).

703   16.   Itahana, K. *et al.* A role for p53 in maintaining and establishing the quiescence growth

704   arrest in human cells. *J Biol Chem* **277**, 18206-14 (2002).

705   17.   Sadasivam, S. & DeCaprio, J.A. The DREAM complex: master coordinator of cell cycle-

706   dependent gene expression. *Nature Reviews Cancer* **13**, 585-595 (2013).

707   18.   Min, M. & Spencer, S.L. Spontaneously slow-cycling subpopulations of human cells

708   originate from activation of stress-response pathways. *PLoS Biol* **17**, e3000178 (2019).

709   19.   García-Gutiérrez, L., Delgado, M.D. & León, J. MYC Oncogene Contributions to Release

710   of Cell Cycle Brakes. *Genes (Basel)* **10**(2019).

711   20.   Aguirre-Ghiso, J.A., Estrada, Y., Liu, D. & Ossowski, L. ERK(MAPK) activity as a

712   determinant of tumor growth and dormancy; regulation by p38(SAPK). *Cancer Res* **63**,

713   1684-95 (2003).

714   21.   Spencer, S.L. *et al.* The proliferation-quiescence decision is controlled by a bifurcation in

715   CDK2 activity at mitotic exit. *Cell* **155**, 369-83 (2013).

716   22.   Miller, I. *et al.* Ki67 is a Graded Rather than a Binary Marker of Proliferation versus

717   Quiescence. *Cell Rep* **24**, 1105-1112.e5 (2018).

23.   Reya, T., Morrison, S.J., Clarke, M.F. & Weissman, I.L. Stem cells, cancer, and cancer
stem cells. *Nature* **414**, 105-11 (2001).

24.   Kleinsmith, L.J. & Pierce, G.B., Jr. MULTIPOTENTIALITY OF SINGLE EMBRYONAL
CARCINOMA CELLS. *Cancer Res* **24**, 1544-51 (1964).

25.   Turati, V.A. *et al.* Chemotherapy induces canalization of cell state in childhood B-cell
precursor acute lymphoblastic leukemia. *Nature Cancer* (2021).

26.   Lee, E., Chuang, H.Y., Kim, J.W., Ideker, T. & Lee, D. Inferring pathway activity toward
precise disease classification. *PLoS Comput Biol* **4**, e1000217 (2008).

27.   Barbie, D.A. *et al.* Systematic RNA interference reveals that oncogenic KRAS-driven
cancers require TBK1. *Nature* **462**, 108-12 (2009).

28.   Hänzelmann, S., Castelo, R. & Guinney, J. GSVA: gene set variation analysis for
microarray and RNA-seq data. *BMC Bioinformatics* **14**, 7 (2013).

29.   Foroutan, M. *et al.* Single sample scoring of molecular phenotypes. *BMC Bioinformatics*
19, 404 (2018).

30.   Aulestia, F.J. *et al.* Quiescence status of glioblastoma stem-like cells involves remodelling
of Ca(2+) signalling and mitochondrial shape. *Sci Rep* **8**, 9731 (2018).

31.   Kurppa, K.J. *et al.* Treatment-Induced Tumor Dormancy through YAP-Mediated
Transcriptional Reprogramming of the Apoptotic Pathway. *Cancer Cell* **37**, 104-122.e12
(2020).

32.   Ebinger, S. *et al.* Characterization of Rare, Dormant, and Therapy-Resistant Cells in
Acute Lymphoblastic Leukemia. *Cancer Cell* **30**, 849-862 (2016).

33.   Jordan, N.V. *et al.* HER2 expression identifies dynamic functional states within circulating
breast cancer cells. *Nature* **537**, 102-106 (2016).

34.   Xue, J.Y. *et al.* Rapid non-uniform adaptation to conformation-specific KRAS(G12C)
inhibition. *Nature* **577**, 421-425 (2020).

35.   Buczacki, S.J.A. *et al.* Itraconazole targets cell cycle heterogeneity in colorectal cancer.
J Exp Med **215**, 1891-1912 (2018).

36.   Grigore, F. *et al.* BRAF inhibition in melanoma is associated with the dysregulation of

histone methylation and histone methyltransferases. *Neoplasia* **22**, 376-389 (2020).

37.   Fujimaki, K. *et al.* Graded regulation of cellular quiescence depth between proliferation

and senescence by a lysosomal dimmer switch. *Proc Natl Acad Sci U S A* **116**, 22624-

22634 (2019).

38.   Stallaert, W. *et al.* The structure of the human cell cycle. *bioRxiv*, 2021.02.11.430845

(2021).

39.   Tomasetti, C. & Vogelstein, B. Cancer etiology. Variation in cancer risk among tissues

can be explained by the number of stem cell divisions. *Science* **347**, 78-81 (2015).

40.   Noureen, N. *et al.* Integrated analysis of telomerase enzymatic activity unravels an

association with cancer stemness and proliferation. *Nature Communications* **12**, 139

(2021).

41.   Malta, T.M. *et al.* Machine Learning Identifies Stemness Features Associated with

Oncogenic Dedifferentiation. *Cell* **173**, 338-354.e15 (2018).

42.   Dimri, G.P. *et al.* A biomarker that identifies senescent human cells in culture and in aging

skin in vivo. *Proceedings of the National Academy of Sciences* **92**, 9363 (1995).

43.   Debacq-Chainiaux, F., Erusalimsky, J.D., Campisi, J. & Toussaint, O. Protocols to detect

senescence-associated beta-galactosidase (SA-betagal) activity, a biomarker of

senescent cells in culture and in vivo. *Nat Protoc* **4**, 1798-806 (2009).

44.   Basisty, N. *et al.* A proteomic atlas of senescence-associated secretomes for aging

biomarker development. *PLOS Biology* **18**, e3000599 (2020).

45.   Fischer, M. Census and evaluation of p53 target genes. *Oncogene* **36**, 3943-3956 (2017).

46.   Andreassen, P.R., Lohez, O.D., Lacroix, F.B. & Margolis, R.L. Tetraploid state induces

p53-dependent arrest of nontransformed mammalian cells in G1. *Molecular biology of the*

cell **12**, 1315-1328 (2001).

47.   Di Leonardo, A. *et al.* DNA rereplication in the presence of mitotic spindle inhibitors in

human and mouse fibroblasts lacking either p53 or pRb function. *Cancer Res* **57**, 1013-

9 (1997).

48.   Ganem, N.J. *et al.* Cytokinesis failure triggers hippo tumor suppressor pathway activation.

774   *Cell* **158**, 833-848 (2014).

775   49.   Alexandrov, L.B. *et al.* The repertoire of mutational signatures in human cancer. *Nature*

776   **578**, 94-101 (2020).

777   50.   Alexandrov, L.B. *et al.* Signatures of mutational processes in human cancer. *Nature* **500**,

778   415-421 (2013).

779   51.   Pich, O. *et al.* The mutational footprints of cancer therapies. *Nat Genet* **51**, 1732-1740

780   (2019).

781   52.   Maley, C.C. & Reid, B.J. Natural selection in neoplastic progression of Barrett's

782   esophagus. *Semin Cancer Biol* **15**, 474-83 (2005).

783   53.   Bardeesy, N. & DePinho, R.A. Pancreatic cancer biology and genetics. *Nat Rev Cancer*

784   **2**, 897-909 (2002).

785   54.   Indovina, P., Pentimalli, F., Casini, N., Vocca, I. & Giordano, A. RB1 dual role in

786   proliferation and apoptosis: cell fate control and implications for cancer therapy.

787   *Oncotarget* **6**(2015).

788   55.   Mas-Ponte, D. & Supek, F. DNA mismatch repair promotes APOBEC3-mediated diffuse

789   hypermutation in human cancers. *Nat Genet* **52**, 958-968 (2020).

790   56.   Cortes-Ciriano, I., Lee, S., Park, W.-Y., Kim, T.-M. & Park, P.J. A molecular portrait of

791   microsatellite instability across multiple cancers. *Nature Communications* **8**, 15180

792   (2017).

793   57.   Ben-David, U. *et al.* Genetic and transcriptional evolution alters cancer cell line drug

794   response. *Nature* **560**, 325-330 (2018).

795   58.   Jakobsen, L. *et al.* Novel asymmetrically localizing components of human centrosomes

796   identified by complementary proteomics methods. *Embo j* **30**, 1520-35 (2011).

797   59.   Sillibourne, J.E. *et al.* Assessing the localization of centrosomal proteins by

798   PALM/STORM nanoscopy. *Cytoskeleton (Hoboken)* **68**, 619-27 (2011).

799   60.   Bettencourt-Dias, M. & Glover, D.M. Centrosome biogenesis and function: centrosomics

800   brings new understanding. *Nature Reviews Molecular Cell Biology* **8**, 451-463 (2007).

61. Grallert, A. *et al.* Removal of centrosomal PP1 by NIMA kinase unlocks the MPF feedback

loop to promote mitotic commitment in S. pombe. *Curr Biol* **23**, 213-22 (2013).

62. Wigley, W.C. *et al.* Dynamic association of proteasomal machinery with the centrosome.

The Journal of cell biology **145**, 481-490 (1999).

63. Itoh, K., Jenny, A., Mlodzik, M. & Sokol, S.Y. Centrosomal localization of Diversin and its

relevance to Wnt signaling. *J Cell Sci* **122**, 3791-8 (2009).

64. Kfoury, Y. *et al.* Ubiquitylated Tax targets and binds the IKK signalosome at the

centrosome. *Oncogene* **27**, 1665-1676 (2008).

65. Lin, A., Wang, R.T., Ahn, S., Park, C.C. & Smith, D.J. A genome-wide map of human

genetic interactions inferred from radiation hybrid genotypes. *Genome Res* **20**, 1122-32

(2010).

66. de Almeida, B.P., Vieira, A.F., Paredes, J., Bettencourt-Dias, M. & Barbosa-Morais, N.L.

Pan-cancer association of a centrosome amplification gene expression signature with

genomic alterations and clinical outcome. *PLOS Computational Biology* **15**, e1006832

(2019).

67. Kohlmaier, G. *et al.* Overly long centrioles and defective cell division upon excess of the

SAS-4-related protein CPAP. *Curr Biol* **19**, 1012-8 (2009).

68. Tang, C.-J.C., Fu, R.-H., Wu, K.-S., Hsu, W.-B. & Tang, T.K. CPAP is a cell-cycle

regulated protein that controls centriole length. *Nature Cell Biology* **11**, 825-831 (2009).

69. Hafner, M. *et al.* Multiomics Profiling Establishes the Polypharmacology of FDA-Approved

CDK4/6 Inhibitors and the Potential for Differential Clinical Activity. *Cell Chem Biol* **26**,

1067-1080.e8 (2019).

70. Salvador-Barbero, B. *et al.* CDK4/6 Inhibitors Impair Recovery from Cytotoxic

Chemotherapy in Pancreatic Adenocarcinoma. *Cancer Cell* **37**, 340-353.e6 (2020).

71. Guiley, K.Z. *et al.* p27 allosterically activates cyclin-dependent kinase 4 and antagonizes

palbociclib inhibition. *Science* **366**(2019).

72.   Pack, L.R., Daigh, L.H., Chung, M. & Meyer, T. Clinical CDK4/6 inhibitors induce selective

and immediate dissociation of p21 from cyclin D-CDK4 to inhibit CDK2. *Nat Commun* **12**,

3356 (2021).

73.   Zhu, X. *et al.* The prognostic and predictive potential of Ki-67 in triple-negative breast

cancer. *Scientific Reports* **10**, 225 (2020).

74.   Brown, J.A. *et al.* TGF-β-Induced Quiescence Mediates Chemoresistance of Tumor-

Propagating Cells in Squamous Cell Carcinoma. *Cell Stem Cell* **21**, 650-664.e8 (2017).

75.   Park, S.R. *et al.* Single-Cell Transcriptome Analysis of Colon Cancer Cell Response to 5-

Fluorouracil-Induced DNA Damage. *Cell Rep* **32**, 108077 (2020).

76.   Aissa, A.F. *et al.* Single-cell transcriptional changes associated with drug tolerance and

response to combination therapies in cancer. *Nat Commun* **12**, 1628 (2021).

77.   Tan, X., Thapa, N., Sun, Y. & Anderson, R.A. A kinase-independent role for EGF receptor

in autophagy initiation. *Cell* **160**, 145-60 (2015).

78.   Srivatsan, S.R. *et al.* Massively multiplex chemical transcriptomics at single-cell

resolution. *Science* **367**, 45-51 (2020).

79.   Moser, J., Miller, I., Carter, D. & Spencer, S.L. Control of the Restriction Point by Rb and

p21. *Proc Natl Acad Sci U S A* **115**, E8219-e8227 (2018).

80.   Minassian, L.M., Cotechini, T., Huitema, E. & Graham, C.H. Hypoxia-Induced Resistance

to Chemotherapy in Cancer. *Adv Exp Med Biol* **1136**, 123-139 (2019).

81.   Johnson, W.E., Li, C. & Rabinovic, A. Adjusting batch effects in microarray expression

data using empirical Bayes methods. *Biostatistics* **8**, 118-27 (2007).

82.   Amezquita, R.A. *et al.* Orchestrating single-cell analysis with Bioconductor. *Nature*

Methods **17**, 137-145 (2020).

83.   McCarthy, D.J., Campbell, K.R., Lun, A.T.L. & Wills, Q.F. Scater: pre-processing, quality

control, normalization and visualization of single-cell RNA-seq data in R. *Bioinformatics*

33, 1179-1186 (2017).

84.   Lun, A.T., McCarthy, D.J. & Marioni, J.C. A step-by-step workflow for low-level analysis

of single-cell RNA-seq data with Bioconductor. *F1000Res* **5**, 2122 (2016).

85.   Barretina, J. *et al.* The Cancer Cell Line Encyclopedia enables predictive modelling of
anticancer drug sensitivity. *Nature* **483**, 603-7 (2012).

86.   Colaprico, A. *et al.* TCGAbiolinks: an R/Bioconductor package for integrative analysis of
TCGA data. *Nucleic Acids Res* **44**, e71 (2016).

87.   Hoadley, K.A. *et al.* Cell-of-Origin Patterns Dominate the Molecular Classification of
10,000 Tumors from 33 Types of Cancer. *Cell* **173**, 291-304.e6 (2018).

88.   Zhang, P. *et al.* Germline and Somatic Genetic Variants in the p53 Pathway Interact to
Affect Cancer Risk, Progression, and Drug Response. *Cancer Res* **81**, 1667-1680 (2021).

89.   Wiecek, A.J., Jacobson, D.H., Lason, W. & Secrier, M. Pan-Cancer Survey of Tumor
Mass Dormancy and Underlying Mutational Processes. *Frontiers in Cell and*
Developmental Biology **9**, 1820 (2021).

90.   Taylor, A.M. *et al.* Genomic and Functional Approaches to Understanding Cancer
Aneuploidy. *Cancer Cell* **33**, 676-689.e3 (2018).

91.   Moon, K.R. *et al.* Visualizing structure and transitions in high-dimensional biological data.
Nature biotechnology **37**, 1482-1492 (2019).

92.   Leek, J.T., Johnson, W.E., Parker, H.S., Jaffe, A.E. & Storey, J.D. The sva package for
removing batch effects and other unwanted variation in high-throughput experiments.
Bioinformatics **28**, 882-3 (2012).

93.   Martincorena, I. *et al.* Universal Patterns of Selection in Cancer and Somatic Tissues.
Cell **171**, 1029-1041.e21 (2017).

94.   Warde-Farley, D. *et al.* The GeneMANIA prediction server: biological network integration
for gene prioritization and predicting gene function. *Nucleic Acids Res* **38**, W214-20
(2010).

95.   Kamburov, A. *et al.* ConsensusPathDB: toward a more complete picture of cell biology.
Nucleic Acids Res **39**, D712-7 (2011).

96.   Shannon, P. *et al.* Cytoscape: a software environment for integrated models of
biomolecular interaction networks. *Genome Res* **13**, 2498-504 (2003).

97.   Pearl, L.H., Schierz, A.C., Ward, S.E., Al-Lazikani, B. & Pearl, F.M.G. Therapeutic
opportunities within the DNA damage response. *Nature Reviews Cancer* **15**, 166-180
(2015).

98.   Wagle, M.C. *et al.* A transcriptional MAPK Pathway Activity Score (MPAS) is a clinically
relevant biomarker in multiple cancer types. *NPJ Precis Oncol* **2**, 7 (2018).

99.   Cerami, E. *et al.* The cBio cancer genomics portal: an open platform for exploring
multidimensional cancer genomics data. *Cancer Discov* **2**, 401-4 (2012).

100.   Kan, Z. *et al.* Multi-omics profiling of younger Asian breast cancers reveals distinctive
molecular signatures. *Nat Commun* **9**, 1725 (2018).

101.   Curtis, C. *et al.* The genomic and transcriptomic architecture of 2,000 breast tumours
reveals novel subgroups. *Nature* **486**, 346-52 (2012).

102.   Gadd, S. *et al.* A Children's Oncology Group and TARGET initiative exploring the genetic
landscape of Wilms tumor. *Nat Genet* **49**, 1487-1494 (2017).

103.   Iyer, G. *et al.* Prevalence and co-occurrence of actionable genomic alterations in high-
grade bladder cancer. *J Clin Oncol* **31**, 3133-40 (2013).

104.   Taylor, B.S. *et al.* Integrative genomic profiling of human prostate cancer. *Cancer Cell* **18**,
11-22 (2010).

105.   Barretina, J. *et al.* Subtype-specific genomic alterations define new targets for soft-tissue
sarcoma therapy. *Nat Genet* **42**, 715-21 (2010).

FIGURES

Figure 1: Methodology for quantifying cancer dormancy. (a) Workflow for evaluating

dormancy in RNA-seq cancer samples. 139 genes differentially expressed in multiple forms of quiescence were employed to score dormancy across TCGA cancers. **(b)** Receiver operating characteristic (ROC) curves illustrating the performance of the z-score methodology on separating actively proliferating and quiescent cells in seven publicly available single-cell (continuous curves) and bulk RNA-seq (dotted curves) datasets. AUC = area under the curve. **(c)** Compared classification accuracies of the z-score approach and classic cell proliferation markers across the seven single-cell/bulk RNA-seq validation datasets. **(d)** Quiescence levels of embryonic fibroblast cells under serum starvation for various amounts of time. Replicates are depicted in the same colour. **(e)** Representative images of lung cancer cell lines immunostained and analysed to detect the quiescent fraction. Hoechst (labels all nuclei) is in blue, PRb in green and EdU in red in merged image. White dashed circles highlight quiescent cells that are negative for both PRb and EdU signals. Scale bar: 100 µm. **(f)** Graphs show single cell quantification of PRb and EdU intensities taken from images and used to define the cut-off to calculate the quiescent fraction (QF; green boxes). Images in **(e)** and graphs in **(f)** are taken from the A549 cell line. **(g-h)** Correlation between theoretical estimates of a quiescence or G1 state and the fraction of cells entering quiescence in nine lung adenocarcinoma cell lines, as assessed through **(g)** phospho-Rb assays and **(h)** EdU assays. Mean of n=3 is shown for the average percentage of quiescent cells.

Figure 2: Pan-cancer evaluation of dormancy. (a) PHATE plot illustrating the wide spectrum

of dormancy-proliferation states across 9,712 primary tumour samples from TCGA. Each sample

is coloured according to the relative dormancy level. **(b)** Variation in tumour dormancy levels

across different cancer tissues. **(c)** Correlation between mean quiescence scores and stem cell

division estimates for various tissue types. **(d)** Correlating tumour dormancy scores with cancer

cell stemness (Stemness Index), telomerase activity (EXTEND score) and the expression of

several commonly used proliferation markers. The Pearson correlation coefficient is displayed.

RC – replication complex. **(e)** Consistently higher levels of dormancy are detected in samples

with functional p53. **(f)** Lower quiescence scores are observed in tumours with one or two whole

genome duplication events. Wilcoxon rank-sum test p-values are displayed in boxplots, *p<0.05;

**p<0.01; ***p<0.001; ****p<0.0001.

Figure 3: Pan-cancer genomic landscape of dormancy. (a) Cancer drivers with mutations or

copy number alterations depleted pan-cancer in a dormancy context. Features further selected

by the pan-cancer model are highlighted. **(b)** Schematic of the ensemble elastic net modelling

employed to prioritise genomic changes associated with tumour dormancy. **(c)** Genomic events

significantly associated with dormancy, ranked according to their importance in the model (from

highest to lowest contribution). Each point represents an individual primary tumour sample of the

training dataset, coloured by the corresponding value of each feature. For discrete variables

purple indicates the presence of the feature and yellow indicates the absence of the feature. The

Shapley values indicate the impact of individual feature values on the quiescence score

prediction. **(d)** Dormancy levels are significantly reduced in microsatellite unstable (MSI) samples

in stomach adenocarcinoma (STAD) and uterine corpus endometrial carcinoma (UCEC), with

the same trend (albeit not significant) shown in colon adenocarcinoma (COAD). Wilcoxon rank-

sum test *$p < 0.05$; **$p < 0.01$. **(e)** Genomic alterations are depleted across DNA repair pathways

during dormancy. Odds ratios of mutational load on pathway in dormancy are depicted, along

with confidence intervals. CS, chromosome segregation; p53, p53 pathway; UR, ubiquitylation

response; CPF, checkpoint factor; TM, telomere maintenance; CR, chromatin remodelling; TLS,

translesion synthesis; NHEJ, non-homologous end joining; NER, nucleotide excision repair;

MMR, mismatch repair; FA, Fanconi Anaemia pathway; BER, base excision repair. **(f)** Dormancy

is increased in cell lines with slow doubling time across MCF7 strains, which also show lower

prevalence of PTEN mutations. **(g)** Tissue-specific changes in quiescence between samples

with/without dormancy-associated deletions (blue), amplifications (red) and SNVs (green) within

the TCGA cohort (top) and external validation datasets (bottom).

Figure 4: CEP89 amplification is associated with lower tumour dormancy. (a) Network illustrating CEP89 interactions with cell cycle genes (from GeneMania). The edge colour indicates the interaction type, with green representing genetic interactions, orange representing predicted interactions and purple indicating pathway interactions. The edge width illustrates the interaction weight. **(b)** Pan-cancer relationship between CA20 scores and tumour quiescence across the TCGA cohort. **(c)** CA20 scores are significantly increased in TCGA primary tumours containing a CEP89 amplification. **(d)** Cox proportional hazards analysis estimates of the log hazards ratio for the impact of CEP89 expression on patient prognosis within individual cancer studies, after considering tumour stage. Patients with high expression of CEP89 show significantly worse prognosis within ACC, LUSC, LIHC, KIRC and STAD, but significantly better prognosis within HNSC, PAAD and KIRP studies. **(e)** Western blot showing depletion of Cep89

protein 48h after siRNA transfection of NCI-H1299 cells. Mock is lipofectamine only, NTC is Non-

targeting control siRNA. B-actin is used as a loading control. **(f)** Graphs show that Cep89

depletion in NCI-H1299 cells leads to a reduction in nuclear number and an increase in the

fraction of quiescent cells, measured by an increase in the percentage of EdU negative (24h EdU

pulse) and Phospho-Ser 807/811 Rb negative cells. One-Way ANOVA, $*p<0.05$, $**p<0.01$. Mean

of n=3.

Figure 5: Genomic landscape of breast cancer dormancy. (a) ROC curve illustrating the

mean performance of the random forest model for dormancy prediction in breast cancer based

on 9 randomisations. **(b)** Impurity criterion information gain for the top ranked genomic events

predictive of dormancy in the TCGA BRCA cohort, modelled using random forests. Features also

appearing in the pan-cancer model are highlighted in blue. **(c)** Tissue-specific changes in

dormancy estimates between samples with and without dormancy-associated amplifications

(red) and SNVs (purple), identified by the breast cancer specific random forest model, within the

TCGA cohort (top panel) and external validation datasets (bottom panel). **(d)** Dormancy scores

are significantly higher in APOBEC mutation-enriched breast cancer samples. **(e)** APOBEC-

linked SBS2 mutational burden compared between breast cancer subtypes. **(f)** Association

between dormancy scores, APOBEC enrichment and breast cancer subtypes in TCGA.

Figure 6: Pan-cancer characterisation of individual quiescence programmes. (a-e)
Comparison of correlation coefficients between quiescence programme scores and **(a)** mean
expression of CDK4 and CDK6, **(b)** mean expression of curated contact inhibition genes, **(c)** a
transcriptional MAPK Pathway Activity Score (MPAS), **(d)** mean expression of curated serum
starvation genes, and **(e)** CDKN1A expression, across TCGA cancers. The correlations expected
to be strongest (either negative or positive) are denoted by an asterisk. The generic quiescence
score refers to scores calculated using the original list of 139 genes differentially expressed
across all 5 forms of quiescence. **(f)** Comparison of quiescence programme scores measured in
cancer cell lines before (red) and after (blue) Palbociclib treatment across three validation
studies. Datasets used for validation are denoted by their corresponding GEO series accession
number. **(g)** Predicted quiescence type composition of samples estimated to be dormant across
individual cancer types. The same colour legend as in **(a)** is applied. Gray bars represent the
proportion of samples for which the quiescence type could not be estimated.

1014

Figure 7: Impact of cellular dormancy on patient prognosis and treatment response. (a)

Disease-specific survival based on tumour dormancy for patients from TCGA within 15 years of

follow up. Patients with high primary tumour dormancy showed significantly better prognosis than

patients with low dormancy. **(b-c)** Hazard ratio ranges illustrating the impact of different forms of

quiescence **(b)** and different tissues **(c)** on patient prognosis, after taking into account potential

cofounding factors. Values above 0 indicate significantly better prognosis in the context of high

tumour dormancy. **(d)** Change in quiescence scores inferred from bulk RNA-seq across breast,

pancreatic, colorectal and skin cancer cells in response to treatment with the CDK4/6 inhibitor

Palbociclib, 5-FU or the BRAF inhibitor Vemurafenib. **(e-f)** UMAP plot illustrating the response of

the TP53-proficient RKO colorectal cancer cell line to various 5-FU doses and the corresponding

proportions of cells predicted to be quiescent/proliferating. Each dot is an individual cell, coloured

according to its dormancy level. **(g-h)** The same as previous, but for the TP53-deficient SW480

cell line. **(i-j)** UMAP plot illustrating the response of individual PC9 NSCLC cells to the EGFR

inhibitor Erlotinib across several timepoints and the corresponding proportion of cells predicted

to be quiescent/proliferating. **(k)** Principal component analysis illustrating the superimposition of

scRNA-seq profiles (circles) of quiescent NSCLC cells before and after EGFR inhibition onto the

bulk RNA-seq reference data (triangles) for MCF10A cells occupying various quiescence states.

(l) The proportion of NSCLC cells in **(k)** predicted to occupy different quiescence states across

several timepoints.

1034

1035

Figure 8: Optimisation of dormancy signature for use in scRNA-seq data. (a) Methodology for reducing the gene signature of dormancy: random forest classifiers are trained to distinguish dormant from cycling tumours on three high confidence datasets; Gini index thresholding is optimised to prioritise a final list of 35 genes. **(b)** Gini index variation, correlation with experimentally measured quiescence via EdU and PRb staining assays, and corresponding p-values are plotted as the number of genes considered in the model is increased. The red dotted line indicates the threshold chosen for the final solution of 35 genes. The black dotted line

indicates the threshold for p-value significance. **(c)** Additional external validation of the 35 gene

signature acting as a classifier of quiescent and proliferation cells in single cell and bulk datasets.

(d) Dropout in single cell data by gene signature. The percentage of genes out of the 35 (blue)

and 139 (red) gene lists with reported expression across five single-cell RNA-seq datasets. **(e)**

Proportion of cycling and quiescent cells estimated in single cell datasets of p53 wild type and

mutant lines treated with 5FU, as well as cells treated with EGFR inhibitors. Data as in Figure 7.