1 **Pathways to polar adaptation in fishes revealed by long-read sequencing**
2

3 Scott Hotaling[1], Thomas Desvignes[2], John S. Sproul[3], Luana S.F. Lins[4], and Joanna L. Kelley[1]
4
5 **Affiliations:**
6 [1] School of Biological Sciences, Washington State University, Pullman, WA, USA
7 [2] Institute of Neuroscience, University of Oregon, Eugene, OR, USA
8 [3] Department of Biology, University of Nebraska Omaha, Omaha, NE, USA
9 [4] CSIRO, New South Wales, Canberra, Australia
10
11 **Correspondence:**
12 Scott Hotaling, School of Biological Sciences, Washington State University, Pullman, WA,
13 99164, USA; Email: scott.hotaling@wsu.edu; Phone: (828) 507-9950; ORCID: 0000-0002-5965-
14 0986
15
16 Joanna L. Kelley, School of Biological Sciences, Washington State University, Pullman, WA,
17 99164, USA; Email: joanna.l.kelley@wsu.edu; Phone: (509) 335-0037; ORCID: 0000-0002-
18 7731-605X

19
20 **Running head:** Long-read sequencing of a polar fish
21
22 **Keywords:** polar fish, genome biology, extremophile, Southern Ocean, cold adaptation,
23 antifreeze proteins
24

Abstract:
Long-read sequencing is driving a new reality for genome science where highly contiguous
assemblies can be produced efficiently with modest resources. Genome assemblies from long-
read sequencing are particularly exciting for understanding the evolution of complex genomic
regions that are often difficult to assemble. In this study, we leveraged long-read sequencing to
generate a high-quality genome assembly for an Antarctic eelpout, *Opthalmolycus amberensis*,
the first for the globally distributed family Zoarcidae. We used this assembly to understand how
O. amberensis has adapted to the harsh Southern Ocean and compared it to another group of
Antarctic fishes: the notothenioids. We showed that from a genome-wide perspective, selection
has largely acted on different targets in eelpouts relative to notothenioids. However, we did find
some overlap; in both groups, selection has acted on genes involved in membrane structure
and DNA repair. We found evidence for historical shifts of transposable element activity in *O.*
amberensis and other polar fishes, perhaps reflecting a response to environmental change. We
were specifically interested in the evolution of two complex genomic regions known to underlie
key adaptations to polar seas: hemoglobin and antifreeze proteins (AFPs). We observed unique
evolution of the hemoglobin MN cluster in eelpouts and related fishes in the suborder Zoarcoidei
relative to other teleosts. For AFPs, we identified the first species in the suborder with no
evidence of *afpIII* sequences (*Cebidichthys violaceus*), potentially reflecting a lineage-specific
loss of this gene cluster. Beyond polar fishes, our results highlight the power of long-read
sequencing to understand genome evolution.

45
Introduction:
Long-reads are revolutionizing genome sequencing and allowing for improved resolution of
genotypes underlying phenotypes across the tree of life (Hotaling, Kelley, & Frandsen, in press;
Hotaling, Sproul, et al., 2021). This new potential is particularly valuable for understanding
phenotypes that stem from complex, difficult to assemble regions of the genome (e.g., highly
repetitive regions). For instance, the evolution of antifreeze proteins (AFPs)—proteins that

adsorb to ice and limit its growth (Davies, Baardsnes, Kuiper, & Walker, 2002; X. Li et al.,
1985)—has allowed a wide array of teleost fishes to thrive in marine habitats that are commonly
below freezing and ice-laden (Hobbs, Hall, Graham, Davies, & Fletcher, 2020). AFPs are
commonly found in highly repetitive, tandem arrays in the genome and have experienced a
large amount of duplication, rearrangement, and modification (e.g., Deng, Cheng, Ye, He, &
Chen, 2010; Scott, Hew, & Davies, 1985) Thus, the evolution of AFPs and phenotypes with
similarly complex genomic underpinnings has been difficult to understand despite their key role
in shaping ecological communities worldwide.
60
Extreme environments provide insight into the nature of adaptive evolution due to their strong,
and often long-term, selective pressures (Kim et al., 2019; S. Xu, Wang, Guo, He, & Shi, 2020).
Extreme habitats are an even more powerful tool when distantly related lineages are subjected
to the same selective pressures and their responses are compared. Under this framework, it is
possible to disentangle lineage-specific evolutionary responses from more general patterns
(e.g., convergent evolution). The Southern Ocean surrounding Antarctica is one of the most
contiguous extreme environments on Earth. It is generally demarcated by latitudes south of the
Antarctic Convergence (also known as Antarctic Polar Front), where extremely cold (sea
surface temperature < 0°C), northward-flowing water from Antarctica meets the comparatively
warmer waters of the subantarctic (sea surface temperature > 5.5°C; Moore, Abbott, &
Richman, 1999). The extreme nature of the Southern Ocean largely stems from three factors:
chronic, subfreezing temperatures, high levels of dissolved oxygen, and dramatic seasonal
shifts in light availability (DeVries & Steffensen, 2005). Such extreme conditions greatly
influence the biological diversity of the Southern Ocean (Griffiths, 2010) and have driven the
evolution of an array of extreme phenotypes (e.g., sea spider gigantism, A. L. Moran & Woods,
2012).
77
Among fishes, the most well-known outcome of the long-term selection at both poles is the
evolution of AFPs. Based on protein structure, multiple different AFP types have evolved in
fishes, including antifreeze glycoproteins (AFGPs) in cod and Antarctic notothenioids and Type
III antifreeze proteins in eelpouts (Davies et al., 2002; X. Li et al., 1985). AFPs appear
ubiquitous in polar fishes but understanding of their molecular evolution remains incomplete (but
see Chen, DeVries, & Cheng, 1997a; Chen, DeVries, & Cheng, 1997b; Deng et al., 2010;
Zhuang, Yang, Murphy, & Cheng, 2019). Another peculiarity of life in the Southern Ocean are
the white-blooded Antarctic icefishes (family Channichthyidae, Ruud, 1954), which are the only
vertebrates known to live without hemoglobin (Beck et al., 2021; Kim et al., 2019). To date,
practical limitations have inhibited our understanding of AFP and hemoglobin evolution,
particularly the challenge of assembling these repeat-rich, complex loci with short-read
sequence data (e.g., Ahn et al., 2017).
90
Thus far, genomic investigation of Antarctic fishes has been restricted to the notothenioid
adaptive radiation (e.g., Kim et al., 2019; Near et al., 2012). These studies have yielded high-
resolution understanding of how cryonotothenioids (Antarctic notothenioids) have adapted to the
Southern Ocean (Chen et al., 2019) and how the unique white-blooded icefishes evolved
following the loss of hemoglobin (Near, Parker, & Detrich III, 2006). Still, many questions
remain. Namely, how generalizable is the evolutionary trajectory of cryonotothenioids relative to
other fishes in the Southern Ocean? Eelpouts (family Zoarcidae) are common fishes in the
Southern Ocean. With roughly ~300 described species distributed around the world and many
in Antarctica (Anderson, 1994; Hotaling, Borowiec, Lins, Desvignes, & Kelley, 2021; Møller,
Nielsen, & Anderson, 2005), eelpouts are the most speciose family within the suborder
Zoarcoidei (also referred to as the infraorder Zoarcales). Similar to cryonotothenioids, eelpouts
have evolved AFPs to mitigate freezing stress, specifically Type III AFPs (*afpIII*; X. Li et al.,

1985). The most recent common ancestor of eelpouts and cryonotothenioids diverged ~80
million years ago (Rabosky et al., 2018), making these groups an ideal system for studying
potential convergent evolution to the harsh Southern Ocean.
106
In this study, we used long-read sequencing to generate a high-quality genome assembly for an
Antarctic eelpout, *Ophthalmolycus amberensis*, the first for the family Zoarcidae. We then
leveraged our new genomic resource to better understand how fishes have adapted to the
Southern Ocean through whole-genome perspectives on genome evolution and fine-scale
investigation of synteny and copy number variation in key adaptive regions (i.e., hemoglobin
and AFPs). Collectively, our results highlight convergent and species-specific mechanisms
underlying the adaptation of fishes to the harsh conditions of the Southern Ocean. Our study
also provides practical evidence for the power and promise of long-read sequencing for
understanding genome biology, particularly in highly repetitive and/or complex genomic regions.
116
117

118
119
Figure 1. (A) An Antarctic eelpout, *Ophthalmolycus amberensis*, sampled from the Gerlache Strait on the
West Antarctic Peninsula. (B) Depth records (*n* = 168 records) for *O. amberensis* from GBIF
(https://www.gbif.org/). (C) The distribution of *O. amberensis* from AquaMaps
(https://www.aquamaps.org/). Red circles represent sampling localities and the dark line indicates the
Antarctic Convergence (or Antarctic Polar Front) where subfreezing northward flowing Antarctic waters
meet the comparatively warmer waters of the sub-Antarctic. Sea surface temperatures south of the
Antarctic Convergence are commonly below 0°C. (D) Assembly and BUSCO (Benchmarking Universal
Single-Copy Orthologs; Simão, Waterhouse, Ioannidis, Kriventseva, & Zdobnov, 2015) for the *O.*
amberensis genome assembly presented in this study.
129
Materials and methods:
Species description and sample collection
Ophthalmolycus amberensis is a coastal Antarctic species, likely with a circumpolar distribution
(Fig. 1). *Ophthalmolycus amberensis* individuals were collected from 360-380 m in May 2016
using a 5-feet Blake trawl deployed from the Antarctic Research and Supply Vessel (ARSV)

Laurence M. Gould in the Gerlache Strait (64°44'S, 63°01'W) on the West Antarctic Peninsula.
Live fish were transferred immediately from the trawl net to the ship aquaria for transport to the
aquatic facilities at Palmer Station, Antarctica, where they were maintained in flow-through
seawater tanks at ~0°C. Following euthanasia with a lethal dose of MS-222 (Syndel, Ferndale,
WA, USA), individuals were dissected and sampled with skeletal muscle flash frozen in liquid
nitrogen and stored at -80°C, fin clips stored in 80% ethanol at room temperature, and liver
preserved in RNALater until it could be moved to longer term -80°C storage. All procedures
were performed according to protocols approved by the Institutional Animal Care and Use
Committees (IACUC) of the University of Oregon (#13-27RRAA). We confirmed visual species
identification through sequencing of the *mitochondrial cytochrome c oxidase I* (*mt-co1*) locus
and comparison to sequences on GenBank.
146
Sequence data collection
We collected three sequence data sets for this study: long-read Pacific Bioscience (PacBio)
data, short-read Illumina whole-genome (WGS) data, and short-read Illumina transcriptomic
(RNA-seq) data. Our sequence data were collected from three individuals and tissue types:
PacBio (EP06, muscle), WGS (EP10, fin clip), and RNA-seq (EP01, liver). PacBio Sequel
SMRTbell libraries were prepared and sequenced at the Oregon State University Center for
Quantitative Life Sciences on a PacBio Sequel platform with eight SMRT cells. For WGS library
preparation, genomic DNA was extracted using a MagAttract HMW DNA kit and the library was
prepared following an Illumina PCR-free protocol with a 400-bp insert size. The resulting WGS
library was sequenced at MedGenome, Inc. on an Illumina HiSeq 4000 platform with 150-bp
paired-end sequencing chemistry. Total RNA was extracted from ~30 mg of liver tissue using
the Macherey-Nagel NucleoSpin® kit and the manufacturer protocol. The tissue was flash
frozen with liquid nitrogen and crushed to homogenize it before extraction. An RNA-seq library
was prepared by the University of Utah Genomics Core Facility and sequenced on an Illumina
HiSeq 2500 with 125-bp paired-end sequencing chemistry. We assessed sequence quality for
both short-read data sets (WGS and RNA-seq) with FastQC v0.11.4 (Andrews, 2010). We
trimmed the raw WGS reads with Trim Galore! v0.4.2 (Krueger, 2015) with stringency of 5,
quality of 28, and a minimum length of 50 bp.
165
Genome assembly
We first estimated the genome size and heterozygosity of the *O. amberensis* genome for the
WGS data using the online portal of GenomeScope (Vurture et al., 2017). We produced the
input for GenomeScope, a histogram of 21-mer counts, with jellyfish v2.2.5 (Melsted &
Pritchard, 2011). We also estimated genome size from the WGS data with sga preQC v0.10.14
(Simpson & Durbin, 2010). For genome assembly, we first combined our separate subread BAM
files into a combined BAM using 'pbmerge', from the PacBio toolkit. Next, using 'bam2fastx',
another PacBio tool, we converted the raw, combined subread BAM into FASTQ format for
downstream analyses. Scripts and commands used in this study are provided GitHub
(http://github.io/oamberensis_genome).
176
We assembled the genome from PacBio sequences only using Canu v1.8 (Koren et al., 2017),
a dedicated long-read assembler designed for PacBio data. We changed one parameter,
cnsErrorRate = 0.25, all other parameters were left as defaults. For all iterations of the genome
assembly, we generated summary statistics using the Assemblathon2 summary script
(assemblathon_stats.pl, Bradnam et al., 2013). We also assessed completeness by calculating
the number of observed single-copy orthologs (BUSCOs) in the assembly using BUSCO v3 and
the 4,584 "actinopterygii_odb9" reference gene set (Simão et al., 2015).
184

According to our BUSCO results, the initial Canu assembly (v1) of the *O. amberensis* genome
was ~40% larger than expected at 1.02 Gb with many duplicated regions (*n* = 1180 complete
and duplicated BUSCOs), likely due to heterozygosity between haplotypes. To resolve duplicate
contigs we used Purge Haplotigs (Roach et al. 2018). We first mapped raw PacBio reads to the
v1 assembly with Minimap2 v2.14-r883 (Li 2018) and the options: -ax -map-pb. Next, we sorted
the resulting BAM file of aligned reads with samtools v1.9 (Li et al. 2009) and default settings.
We used the 'readhist' function of Purge Haplotigs to generate a coverage histogram of the
mapped PacBio reads. We set low (-l), middle (-m), and high (-h) in the coverage distribution of
10, 140, 195, respectively, and ran the second step in the Purge Haplotigs pipeline, 'contigcov',
to analyze coverage on a contig-by-contig basis. Finally, we purged duplicate contigs using the
function 'purge' to create a de-duplicated assembly. The resulting assembly (v2) was
substantially smaller at 679 Mb with 84.5% fewer duplicated genes.
197
We then polished the v2 assembly using our raw PacBio reads with Arrow, an algorithm in the
variantCaller program within the Falcon software (Chin et al. 2013, 2016). Since this approach is
not deterministic (i.e., repeated runs will always find changes), we performed two rounds of
polishing with Arrow (resulting in assemblies v3 and v4). For each round, we first re-mapped our
raw PacBio reads to the latest assembly version (e.g., to make v3, we mapped raw PacBio
reads to the v2 assembly) with Minimap2 v2.14-r883 (Li 2018) and the same flags as above.
Next, we converted the BAM file of mapped reads to PacBio format with 'pbbamify' (part of the
PacBio toolkit) with default flags. We sorted the re-formatted BAM file with samtools v1.9 as
above (Li et al. 2009). We created an index of the mapped, sorted BAM with another PacBio
tool, 'pbindex'. This process was used to create the polished v3 assembly and repeated for v4.
208
Transcriptome assembly
To aid in genome annotation, we assembled an *O. amberensis* liver transcriptome from raw
RNA-seq reads with Trinity v2.6.6 (Grabherr et al. 2011) using default settings and read
trimming with Trimmomatic (Bolger et al. 2014) with flag "--trimmomatic." We assessed
transcriptome assembly completeness with BUSCO v3 and the 4,584 "actinopterygii_odb9"
reference gene set (Simao et al. 2015).
215
Genome annotation
We annotated the *O. amberensis* genome assembly (v4) using Maker2 v31.10 (Holt & Yandell,
2011). To support the annotation pipeline, we generated a repeat library using RepeatModeler
v1.0.11 with default settings (A. F. Smit & Hubley, 2008). We also included gene evidence from
the *O. amberensis* transcriptome and proteins for two other species, three-spined stickleback
(*Gasterosteus aculeatus*) and Atlantic cod (*Gadus morhua*), which were both downloaded from
Ensembl release 97 (http://ftp.ensembl.org/pub/release-97/). We first performed a *de novo*
annotation of the *O. amberensis* genome assembly with our transcriptome and protein evidence
used with options est2genome = 1 and protein2genome = 1 in the Maker2 control file. Next, we
used the gene finding program, SNAP (Korf, 2004), to generate *ab initio* gene predictions for our
assembly. We first merged all of our gene models from our *de novo* Maker2 together into a
single GFF3 using the Maker2 tool "gff3_merge" and converted them to ZFF format with
"maker2zff." We trained SNAP on these data by first breaking up our sequences into one gene
per locus (with 1000 bp on either side of the predicted gene) using "fathom" (part of SNAP). We
estimated parameters for our genes using "forge" (also included with SNAP) before building a
Hidden Markov Model for our annotations with the script "hmm-assembler.pl". After SNAP
training, we re-ran Maker2 with our trained SNAP models by adding them in the control file
("snaphmm" option) and setting est2genome and protein2genome both to 0. We performed a
second round of SNAP training using the same approach followed by running Maker2 for a third
time. Next, we prepared to train another gene predictor, Augustus v2.5.5 (Stanke & Waack,

2003), on our gene models by first merging the gene models produced in the second round the
SNAP training using "gff3_merge", converting them to ZFF with "maker2zff", and then to GBK
format with a custom perl script (https://github.com/hyphaltip/genome-
scripts/blob/master/gene_prediction/zff2augustus_gbk.pl). As Augustus is a machine learning
approach, we first split our data set into a training and test set using the script "randomSplit.pl."
We then created a new "species" for Augustus with the script "new_species.pl" then performed
training by running the "etraining" function on our training data set followed by evaluating its
performance with the "augustus" tool. We optimized our prediction parameters with the script
"optimize_augustus.pl." After Augustus training, we re-ran Maker2 as above with the addition of
the control file option "augustus_species." Finally, we ran GeneMark for the *O. amberensis*
genome assembly with the self-training algorithm using flag -ES. We then performed a final run
of Maker2 with all of the gene model evidence included by adding the path to our GeneMark
trained predictions using "gmhmm" to our control file along with the above evidence for
Augustus ("augustus_species") and SNAP ("snaphmm"). We kept est2genome and
protein2genome as 0 for this final run.
251
Repetitive element classification and annotation
We identified and classified repetitive elements (REs) in the genome assembly of *O.*
amberensis using RepeatModeler v2.0.1 (Flynn et al., 2020). We then used the
"queryRepeatDatabase.pl" script included with RepeatMasker v4.1.0 (A. Smit, Hubley, & Green,
2015) to output RepeatMasker's internal library for ray-finned fishes (-species "actinopterygii")
and merged it with the *O. amberensis* custom repeat library generated by RepeatModeler. We
used this combined Actinopterygii + *O. amberensis* library to annotate REs in the *O. amberensis*
assembly using RepeatMasker v4.1.0 with the search engine set to "ncbi" and the -xsmall
option. We repeated the above steps to classify and annotate REs in each of five additional fish
genome assemblies—*Gasterosteus aculeatus* (BROAD S1, Ensembl release 97), *Gadus*
morhua (gadMor1, Ensembl release 97), *Takifugu rubripes* (FUGU5, Ensembl release 98),
Chaenocephalus aceratus (Chaenocephalus aceratus V1.0), and *Parachaenicthys charcoti*
(Parachaenichthys charcoti V1.0, GigaDB dataset #100321)—such that each assembly was
annotated using a combined library of Actinopterygii + the custom library from RepeatModeler
for each species. We summarized repeat abundance by parsing output from RepeatMasker and
generating plots in R v3.5.1 (R Core Team, 2021) using custom scripts. To investigate patterns
of transposable element (TE) activity through time we generated TE copy number divergence
plots. We calculated Kimura substitution level (CpG adjusted) for transposable elements using
the RepeatMasker utility script "calcDivergenceFromAlign.pl", removed non-transposable
element and unclassified repeats from the resulting ".divisum" file, and generated age
distribution landscapes using RepeatMasker's "createRepeatLandscape.pl" script.
273
Selection on protein-coding genes
To assess selection on protein-coding genes in the *O. amberensis* genome, we calculated the
ratio of non-synonymous to synonymous substitution (*dN/dS*) using CodeML with the PAML
package (Yang, 2007). We first identified orthologous groups of single-copy coding sequences
using the OrthoVenn2 web server (L. Xu et al., 2019). Coding sequences for the same six fish
genomes as the repeat analyses above were included: *G. aculeatus*, *G. morhua*, *O.*
amberensis, *T. rubripes*, *C. aceratus*, and *P. charcoti*. To identify orthologous gene groups, we
translated coding sequences (CDS) to protein using the function "transeq" in EMBOSS v.6.6.0
(Rice, Longden, & Bleasby, 2000). We then extracted corresponding CDS for each group of
single-copy orthologs for each taxon. We aligned each of the orthologous CDS with PRANK
v.170427 (Löytynoja, 2014) under a codon model. Before performing *dN/dS* analyses, we
generated a species tree with RAxML v.8.2.12 (Stamatakis, 2014) for a concatenated alignment

of all of the single-copy orthologous nucleotide sequences identified above with 100 bootstrap
replicates using a GTR+Gamma model.
288
Using CodeML, we calculated *dN/dS* for aligned single-copy orthologous coding sequences with
branch-site models A1 (model = 2, NSsites = 2, fix_omega = 1) and A (model = 2, NSsites = 2,
fix_omega = 0) and F3X4 codon frequency and cleandata = 1 for both. Using the same models,
we varied the foreground lineage(s) by first focusing only on *O. amberensis* with all other
lineages as background and second with the two notothenioids (*C. aceratus* and *P. charcoti*) as
foreground lineages. To statistically compare our test model (A) to our null neutral model (A1)
for each group of orthologs, we performed a Chi-Square Test. To obtain *P*-values for all chi-
square values, we used the pchisq() function in R v.3.6.3 (R Core Team, 2021). To control for
multiple tests, we performed false discovery rate (FDR) correction in R using the p.adjust()
function and the "Benjamini-Hochberg" method (Benjamini & Hochberg, 1995) with a cutoff of
0.05. We identified Gene Ontology (GO) terms for genes that exhibited evidence of positive
selection using Ensembl IDs for *G. aculeatus* and tested for GO term enrichment among target
lists using ShinyGO v0.74 (Ge, Jung, & Yao, 2020). For GO term enrichment analyses, we
compared lists of genes with evidence of positive selection against the background of all single-
copy orthologs. We assessed GO term enrichment for three categories of terms: biological
process, cellular component, and molecular function.
305
Annotating hemoglobin and AFP regions
To explore the evolution of hemoglobin genes in *O. amberensis* and across the suborder
Zoarcoidei, hemoglobin gene clusters were analyzed and compared to other species
representing major perciformes lineages: *Cebidichthys violaceus* (Cvio_1.0, Heras,
Chakraborty, Emerson, & German, 2020), *Anarrhichthys ocellatus* (GSC_Weel_1.0), *P.
gunnellus* (fPhoGun1.1), *Perca flavescens* (PFLA_1.0, Feron et al., 2020), *Etheostoma
spectabile* (UIUC_Espe_1.0, R. L. Moran, Catchen, & Fuller, 2020), *G. aculeatus* (BROAD S1),
Pseudoliparis sp. (ASM433547v1, Mu et al., 2021), *Cottoperca gobio* (fCotGob3.1, Bista et al.,
2020), *Ophiodon elongatus* (NWFSC_Oelon_r3.mflye, Longo et al., 2020), *Cyclopterus lumpus*
(fCycLum1.pri), and *Taurus bubalis* (fTauBub2.1). In teleost fish, hemoglobin genes occur in two
distinct, unlinked clusters (LA and MN) which were first located in each assembly by searching
for flanking genes (i.e., *rhbdf1b* and *aqp8.2* for the LA cluster and *kank2* and *nprl3* for the MN
cluster). Sequence, length, position, and strand of individual exons of each alpha and beta gene
were manually retrieved from each assembly by performing BLASTN searches and using the
yellow perch *P. flavescens* as a reference for hemoglobin gene exon boundaries. Exons of each
gene were concatenated and translated into protein sequence to verify their accuracy. If a gene
was incomplete (e.g., missing an exon) or displaying a premature stop codon, the gene was
considered pseudogenized.
324
To explore the evolution of AFPs across the Zoarcoidei (and an outgroup, *G. aculeatus*), region
associated with *afpIII* genes and their progenitor in eelpouts, sialic acid synthase (*nans*; Deng
et al., 2010), were annotated by hand via a combination of BLASTN searches and focused
alignments. We focused on six species (five from Zoarcoidei) with high-quality genomic
resources—five genome assemblies [*G. aculeatus* (BROAD S1), *O. amberensis*, *C. violaceous*
(Cvio_1.0), *A. ocellatus* (GSCB_Weel_1.0), *P. gunnellus* (fPhoGun1.1)] and one set of bacterial
artificial chromosomes (BACs) which targeted the same regions in another Antarctic eelpout,
Lycodicthys dearborni (Deng et al., 2010). For all genes except *afpIII*, we used all of the known
exons for *G. aculeatus* from Ensembl 104. For *afpIII* genes, we used a combination of the two
exons described by Deng et al. (2010) for *L. dearborni* and Hobbs et al. (2020) for *P. gunnellus*.
We first performed a BLASTN v.2.2.31 search with default settings against a reference
database (created with "makeblastdb" for each genome or BAC group) and each exon as a

separate query. We further investigated regions of interest by extracting them with samtools (H.
Li et al., 2009) using the function "faidx" and aligning sequences with Clustal Omega (Sievers et
al., 2011).
340
Where appropriate, we assessed the quality of the assembly at target regions within the
hemoglobin and AFP loci by mapping whole-genome, short-read Illumina data from a different
O. amberensis individual to the *O. amberensis* PacBio genome assembly using bwa v.0.7.12
(Heng Li, 2013). We indexed the PacBio assembly using the "index" function then mapped
reads with the "bwa-mem" algorithm with default settings. We converted the output from a BAM
to SAM file with samtools (H. Li et al., 2009) and visualized regions with the Integrated
Genomics Viewer (Robinson et al., 2011).
348
Results:
Sequence data
We generated 5,536,887 long-read PacBio sequences (65.7 Gb, mean = 13.2 Kb, read N50 =
20 kb) and 237,098,976 Illumina short-reads (WGS = 226,420,980; RNA-seq = 10,677,996). We
estimated the *O. amberensis* genome to be 689.8 Mb from the WGS data. Based on this
estimate, our sequencing coverage for the PacBio data was 95x and 49x for the WGS data. We
estimated the genome heterozygosity to be 0.36%. The genome assembly, raw sequence data,
and associated files for *O. amberensis* are deposited under NCBI BioProject PRJNA701078.
357
O. amberensis *genome assembly and annotation*
Our final assembly of the *O. amberensis* genome (v4) was 680.7 Mb in 1,828 scaffolds with an
N50 of 1.00 Mb. The longest scaffold was 6.98 Mb. Based on the k-mer based estimate of
genome size, our assembly represents 98.7% of the genome. The assembly contains 95.7% of
the single-copy, conserved orthologous genes (BUSCOs) in the 4,584 gene reference set for
Actinopterygii (Fig. 1D). We annotated 22,572 genes in the *O. amberensis* genome assembly.
According to the same BUSCO gene set used for assessing the genome assembly
completeness, the *O. amberensis* liver transcriptome used for annotation was 67.1% complete
with 9.1% of genes fragmented and 23.8% missing.
367
Repetitive element content
Repetitive elements (REs) comprised 30.8% (209.6 Mb) of the *O. amberensis* genome
assembly (Fig. 2). Of all repetitive sequences masked by RepeatMasker, 38.6% (78.8 Mb) were
classified as repeats. DNA transposons accounted for the largest fraction of classified REs
(33.8Mb or 16.2% of total REs), followed by long-interspersed nuclear elements (LINEs)
(25.2Mb or 12.0%) and long-terminal repeats (LTRs; 15.0 Mb or 7.2%), respectively.
Unclassified REs accounted for the remaining repetitive component (130.1 Mb or 62.4% total
REs) of the *O. amberensis* assembly.
376
Opthalmolycus amberensis and the two notothenioid species (*C. aceratus* and *P. charcoti*)
showed greater RE abundance in both genomic proportion and total repeats (range = 30.8–
49.4% or 209.6–525.6 Mb in *O. amberensis* and *C. aceratus*, respectively), relative to the
remaining non-Antarctic species (range = 11.9–24.6% or 52.7–205.0 Mb in *T. rubripes* and *G.*
morhua, respectively; Fig. 2). Although satellite DNAs make up a small fraction of REs in all
species, satellites were at least 10 times more abundant (in terms of genomic proportion) in the
O. amberensis assembly compared to all other assemblies except *C. aceratus*, for which they
were approximately five times more abundant. For all species either DNA transposons or LINEs
comprised the largest proportion of classified repeats.
386

Figure 2. Repetitive element abundance in focal fishes. (A) Abundance of major repetitive element (RE) categories shown as genomic proportions for *O. amberensis* and five other species, *C. aceratus, P. charcoti*, *G. aculeatus, T. rubripes,* and *G. morhua*. Yellow and blue boxes indicate the Antarctic notothenioids and eelpout shown in Figure 3, respectively. (B) RE abundance shown as total bases in the assembly such that the height of each bar is proportional to the assembly length for that species, with the assembly fraction not masked by RepeatMasker shown in gray. (C) Transposable element (TE) copy divergence plots. The y-axis shows TE abundance as a proportion of the genome (e.g., 1.0 = 1% of the genome). The x-axis shows sequence divergence (CpG adjusted Kimura distance) relative to consensus sequences for TE superfamilies. Copy number peaks with abundance skewed toward the left (i.e., low sequence divergence) may represent TE copies with a recent history of diversification relative to peaks with right-skewed abundance which may comprise remnants of more ancient bursts of TE activity. To simplify visual display, TE superfamilies within DNA transposons, Helitrons, LTRs, LINEs, and SINEs were grouped by the same color. A similar plot with all superfamilies separated by color is shown in Fig. S1. Snowflakes indicate species with distributions that include polar seas.

Transposable element (TE) copy divergence plots in *O. amberensis* suggested evidence of dynamic shifts in transposable activity through time (Fig. 2). LINEs dominated recent and

ongoing TE activity driven by recent expansion of *L2* and *Rex/Babar* elements, accounting for
much of the LINE abundance (blue peak between $K = 0–5$; Fig. 2C, Fig. S1). DNA transposons
showed a relatively older history of proliferation dominated by *hAT* elements (broad peak
centered at $K = 10$; Fig. 2C, Fig. S1), though some more recent peaks are also evident. Prior to
this putative increase in DNA transposon activity, Helitrons showed evidence of an ancient burst
with abundant sequences centered over $K = 35–40$ (Fig. 2C, Fig. S1). LTRs appear to have
maintained modest activity through time, although *ERV1* and *Gypsy* elements show some
evidence of recent expansion (Fig. 2C, Fig. S1). Similar trends of recent LINE expansions were
visible in the other two Antarctic species, *C. aceratus* and *P. charcoti*, as well as *T. rubripes*.
Recent DNA transposon activity was evident in all species, though least pronounced in *O.
amberensis*. *Gasterosteus aculeatus* was the only species for which LTRs dominated recent TE
activity (mostly *Gypsy*, *ERV1*, and *Pao* elements; Fig. 2C, Fig. S1). The three other polar
species, *C. aceratus, P. charcoti*, and *G. morhua*, all have evidence for ancient TE expansions
similar to the Helitron expansion in *O. amberensis*, however in these species LINEs and LTRs
drove the expansion (Fig. 2C, Fig. S1).

Figure 3. (A) The guide tree generated for *dN/dS* analyses from 3,461 protein-coding genes. Yellow and
blue boxes indicate the foreground lineages in respective analyses (top, yellow: notothenioids only;
bottom, blue: *O. amberensis* only). (B) The number of genes with evidence of positive selection in each
independent analysis and in both analyses (middle of the Venn diagram). (C) The 10 genes that were
shared between both analyses. When applicable, gene names and IDs from the ortholog in stickleback
(*Gasterosteus aculeatus*, BROAD S1) are given. The FDR cutoff used in all analyses was 0.05.
Snowflakes indicate species with distributions that include polar seas (Antarctic: *C. aceratus*, *P. charcoti*,
O. amberensis; Arctic: *G. morhua*).

Selection on protein-coding genes
We identified 3,461 single-copy orthologous genes that were present in all six focal species.
When only *O. amberensis* was the foreground lineage in *dN/dS* analyses, we identified 123
protein-coding genes with evidence of positive selection with FDR < 0.05 (Fig. 3). When the two
notothenioids (*P. charcoti* and *C. aceratus*) were the foreground lineages, 289 protein-coding
genes exhibited evidence of positive selection. When these two lists were intersected, just 10
protein-coding genes exhibited evidence of positive selection in both groups, leaving 113
(91.9%) and 279 (96.5%) genes that were uniquely under positive selection in *O. amberensis* or
the notothenioids, respectively (Fig. 3). The gene that exhibited the strongest signature of
positive selection in *O. amberensis* (Ensembl ID #ENSGACG00000014975; FDR = 2.02e-10) is
orthologous to human *MPV17L*, a component of the mitochondrial membrane. For the
notothenioids, the strongest signature of positive selection was *eif4enif1* (FDR = 5.06e-30), a

eukaryotic translation initiation factor which has been linked to thermal stress in salmonids
(Akbarzadeh et al., 2018).
447
For the 10 positively selected genes shared between *O. amberensis* and the two notothenioids,
there were not enough genes to assess higher level groupings of GO terms for biological
processes. For cellular components, however, several genes (*snx18*, *sprtn*, and *esrp1*) were
identified as playing roles in organelle and/or membrane processes. Only one pathway was
significantly enriched—Phosphatidylinositol-4,5-bisphosphate binding (FDR = 0.046)—and it
included one positively selected gene (*snx18*). For both the *O. amberensis* and the
notothenioid-specific analyses, there was no evidence of significant GO term enrichment.
However, here we report the top GO terms associated with positively selected genes in each
analysis. For the 124 genes detected in the *O. amberensis* only analysis, the most common GO
categories for biological processes were "regulation of biological process" (n = 26 genes),
"regulation of cellular process" (n = 24), and "response to stimulus" (n = 24). For cellular
components, the most common GO categories were "organelle" (n = 30), "membrane-bounded
organelle" (n = 28), and "intracellular organelle" (n = 28). For molecular function, "protein
binding" (n = 29), "ion binding (n = 27), and "organic cyclic compound binding" (n = 25) were the
most common. Finally, for the 290 genes detected in notothenioids, the most common GO
categories were very similar to those of *O. amberensis.* For biological processes, "regulation of
biological process" (n = 63), "regulation of cellular process" (n = 59), and "multicellular
organismal process" (n = 51) were the most common. For cellular components, the most
common were "organelle" (n = 83), "intracellular organelle" (n =81), and "membrane-bounded
organelle" (n = 75). For molecular function, "protein binding" (n = 79), "ion binding (n = 66),
"organic cyclic compound binding" (n = 66), and "heterocyclic compound binding" (n = 66) were
most common.
470
Hemoglobin and AFPs
In *O. amberensis* and across Zoarcoidei, the LA cluster remained stable with the typical
organization of two alpha-globin and one beta-globin genes (Opazo et al. 2013). However,
among perciformes, the beta-gene of the LA cluster, *hbbla*, was repeatedly lost or
pseudogenized in some species, including in the Zorcoidei monkeyface prickleback
Cebidichthys violaceus. In contrast, the MN cluster in perciformes underwent dynamic changes
in a lineage specific fashion and harbored up to 13 alpha-globin and 15 beta-globin genes in the
Longspined bullhead sculpin *Taurulus bubalis* and numerous instances of pseudogenization,
especially frequent in the Orangethroat darter *Etheostoma spectabile*. While the MN cluster
organization could not be analyzed in the wolf-eel *A. ocellatus* due to the lack of contiguity in the
assembly at this locus, within Zoarcoidei, however, the MN cluster composition remained
relatively stable among the three studied species with three alpha-globin genes and two or three
functional beta-genes. A specificity of Zoarcoidei compared to the other species is the presence
of a *hbbmn2* pseudogene in *C. violaceus* and *P. gunnellus* and the loss of this gene in *O.*
amberensis.
486

Figure 4. Evolution of the hemoglobin clusters in Perciformes. (A) The phylogenetic relationship of the studied perciform species was extracted from Rabosky et al. (2018). Snowflakes indicate species with distributions that overlap polar regions. (B) Genomic organization of the hemoglobin LA cluster flanked by the genes *rhbdf1b* and *aqp8.2* in all studied species. (C) Genomic organization of the hemoglobin MN cluster flanked by the genes *kank2* and *nprl3* in all studied species. Genes flanking the clusters are represented by dark-grey squares, alpha-globin genes (*hba*) are represented with red squares, beta-globin genes (*hbb*) are represented with light green squares, and pseudogenes are labelled with an asterisk (*). The arrowheads indicate the direction of the clusters in each assembly and the position of the square above or below the line denotes the position of the gene on the forward or reverse strand, respectively when the arrow points to the right, and the opposite when the arrow points to the left. "N" in *A. ocellatus* denote that the assembly contains gaps between exon1 of *hbamn1* and *nprl3*, therefore preventing the study of this cluster in this species. Basic metrics for the genome assemblies included in this analysis are provided in Table S1.

The sialic acid synthase (*nans*) region is well-conserved across and beyond Zoarcoidei (Fig. 5a). For *afpIII*, we found no evidence of AFPs in *G. aculeatus* and *C. violaceus* (family Stichaeidae; Fig. 5b). Among the four species with copies of some portion of an *afpIII* gene (either exon 1, exon 2, or complete copies of both exons 1 and 2 in order), copy number varied widely. *Lycodicthys dearborni* has at least 25 complete copies of *afpIII* with a number of partial copies (either exon 1 or exon 2 in isolation) scattered throughout the same region. Similarly, *O. amberensis* also had a high number of complete *afpIII copies* (at least 13) with partial copies in the same region. *Pholis gunnellus* has at least 15 complete *afpIII* copies as we all as partial copies. Finally, *A. ocellatus* has evidence of *afpIII* genes but with four complete *afpIII* copies spread over four of the six scaffolds including syntenic components of the *afpIII*, it is unclear

513 how many copies are truly present (Fig. 5b). This is primarily because, unlike *L. dearborni* for
514 instance, no *A. ocellatus* scaffold has evidence of both *afpIII* genes and flanking regions.
515

516
517
Figure 5. Synteny of the (A) sialic acid synthase (*nans*) and (B) *afpIII* region in the suborder Zoarcoidei
with three-spined stickleback (*G. aculeatus*) as the outgroup. Asterisks in (A) indicate evidence for
additional *nans* gene copies in *C. violaceus* and *P. gunnellus* (1 and 8 extra copies, respectively). In (B),
the *P. gunnellus* assembly has one scaffold that contains the *afpIII* flanking genes as well as evidence of
afpIII genes between them as well as a second scaffold that also has evidence of *afpIII* genes. This is
denoted by a second *afpIII* region (red line) directly below the primary scaffold. Snowflakes indicate
species with distributions that overlap polar regions and stars indicate breakpoints between scaffolds. As
in Figure 4, arrowheads indicate the direction of the clusters in each assembly and the position of the
square above or below the line denotes the position of the gene on the forward or reverse strand,
respectively when the arrow points to the right, and the opposite when the arrow points to the left. Basic
metrics for the genome assemblies included in this analysis (all species except for *L. dearborni* which is
not a genome assembly) are provided in Table S1.
530

Discussion:
The rapid accumulation of genomic resources across the tree of life is empowering a new era of
comparative genome science (Hotaling et al., in press; Marks, Hotaling, Frandsen, & VanBuren,
in press). Moreover, the rise of long-read sequencing has improved the contiguity of difficult to
assemble, repeat-rich portions of genomes. Indeed, genome assemblies that integrate long-
reads are on average 48x more contiguous than those that do not (Hotaling, Sproul, et al.,
2021). In this study, we used long-reads to generate a highly contiguous assembly for an
Antarctic eelpout, *Ophthalmolycus amberensis*, the first for the family Zoarcidae. We compared
the genome of *O. amberensis* to other high-quality assemblies for polar and non-polar fishes
with an emphasis on aspects that are difficult or impossible to assess with short-reads alone
(repetitive elements and repeat-rich, dynamic loci underlying key phenotypes). Collectively, our
results provide new understanding of adaptation to the extreme conditions of the Southern
Ocean and highlight the power of long reads for new levels of genomic insight in non-model
species.
545
Repetitive element dynamics
Repetitive elements (REs) are major drivers of genome evolution in fish and their abundance
varies widely across species. Transposable elements (TEs) make up the largest fraction of REs
in fish genomes ranging from 5–56% (Shao, Han, & Peng, 2019). *Ophthalmolycus amberensis*

is intermediate among fish lineages with a genomic proportion of ~31%, of which nearly 11%
are classified as TEs. Our findings generally corroborate previous studies that show DNA
transposons and LINEs are consistently abundant in fish genomes (e.g., Chalopin et al. 2015)
and that RE abundance roughly correlates with genome size. The age distribution of TEs in *O.*
amberensis revealed dynamic shifts in its TE landscape. While LINEs dominated present TE
activity, DNA transposons and Helitrons both showed earlier bursts of activity.
556
Recent studies have highlighted a potential link between environmental stress and bursts of TE
activity in fish (Chénais, Caruso, Hiard, & Casse, 2012). In Antarctic notothenioid fish, *DIRS1*
retrotransposons form hotspots of chromosome rearrangements within lineages, potentially
driving speciation (Auvinet et al., 2018; Auvinet et al., 2019). Heating and cooling cycles in polar
seas have also been hypothesized to induce environmental stress which in turn drives bursts of
TE activity (Auvinet et al., 2018). Consistent with this, we found evidence of recent major *DIRS*
expansion in the notothenioid, *C. aceratus*, but not in other polar fish lineages (Fig. S1). And,
more broadly, greater RE abundance and more evidence of historical activity bursts in the four
polar fishes included in our analysis relative to the temperate fishes, despite their lack of
phylogenetic (and spatial) overlap (Fig. 2, Fig. S1). It is possible that these signatures reflect
historical, climate-induced shifts that affected fishes at both poles. However, this hypothesis is
undermined by the fact that different fish groups adapted to cold environments at different times
in history. Future efforts to integrate dense taxonomic sampling will allow for contextualizing
links between TE activity, environmental stress, and genome evolution, particularly as they
relate to ancient and contemporary environmental change.
572
Selection on protein-coding genes
Across thousands of shared orthologs, only a small fraction exhibited signatures of selection in
either polar fish group (notothenioids or *O. amberensis*), and of those, just 10 genes were
shared between the two. The split between Zoarcoidei and Notothenioidei occurred roughly 80
million years ago (Rabosky et al., 2018). Thus, with a moderate amount of shared ancestry, if
adaptation to cold environments relies on universal biological and physiological mechanisms,
we might expect a signature of parallel evolution in both groups evidenced by significant overlap
between the targets of selection. This was not the case. However, among the subset that did
overlap, two common themes emerged: selection has acted on genes involved in (i) membrane
structure and (ii) DNA repair. Indeed, 50% of the shared genes have functional links to
membrane structure in vertebrates. Membranes play an obvious role in organismal and cell
survival as they delimit cellular structures, mediate transmembrane movement of solutes and
other materials, and play key roles in many other key processes and structures (Rothfield,
2014). Moreover, because environmental conditions play a key role in determining how lipids—a
key component of biological membranes—arrange themselves (Rothfield, 2014), it is
unsurprising that membranes are important for thermal adaptation in a wide variety of
organisms (Hazel, 1995). It is perhaps also unsurprising that fishes living in the most
consistently cold marine habitats on Earth have experienced positive selection on genes related
to membrane structure and function.
592
Generally speaking, membrane fluidity is maintained as temperature changes, a phenomenon
known as homeoviscous adaptation (HVA; Hazel, 1995). However, the extreme, but stable, cold
temperatures of polar seas means that maintenance of an HVA response to changing
conditions should not hold any fitness benefit (Cossins & Prosser, 1978). Still, polar fish
membranes must function at low temperature and directional selection may have acted on this
trait. Another target of selection in polar fishes, *snx18*, in concert with *fip5*, is involved in the
morphogenesis of the epithelial lumen including the formation of membrane tubules (Willenborg
et al., 2011). While the exact significance of selection on membrane-associated genes in polar

fish remains to be determined, it is plausible that at least one major driver is functioning in the
cold.
603
High amounts of dissolved oxygen in the Southern Ocean can boost the production of reactive
oxygen species (ROS) in marine organisms (Pischedda et al., 2018). ROS damage cellular
molecules including DNA (Van Houten, Santa-Gonzalez, & Camargo, 2018). Intact and high-
functioning DNA repair pathways may be particularly crucial for the survival of Antarctic species.
We found evidence for selection on two DNA repair-associated genes in polar fishes, *msh3* and
sprtn. A heterodimer between the proteins encoded by *msh3* and a related gene, *msh2*, binds to
the DNA helix and slides down it until mis-paired bases or other lesions are recognized (Pećina-
Šlaus, Kafka, Salamon, & Bukovac, 2020). The other gene, *sprtn*, is a key component of the
DNA replication machinery and is required for replication, thereby playing an essential role in
the maintenance of genomic stability (Halder et al., 2019). While evidence of positive selection
on DNA repair pathways has also been observed for deep sea fishes (Mu et al., 2021), and *O.*
amberensis commonly occurs in cold and deep waters, the notothenioids included in our
analysis are shallower water species. Our results suggest that evolution to mitigate elevated
oxidative stress may have influenced polar fish evolution. In the case of *O. amberensis*, and
other species living at depth in polar seas, a dual-benefit from selection on DNA repair
pathways may be occurring.
620
Evolution of hemoglobin genes in Zoarcoidei
Hemoglobin is a major trait for adaptation to different environments in fishes (Powers, 1980;
Verde, Giordano, di Prisco, & Andersen, 2012). Because the capacity of hemoglobin to bind
oxygen is sensitive to environmental conditions such as temperature and pH, the molecular
evolution of hemoglobin genes has been proposed to be necessary for fish to maintain
respiratory homeostasis in new or changing environments (Powers, 1980). Vertebrate
hemoglobin is a tetrameric protein composed of two alpha-globin and two beta-globin subunits.
Combinations of different alpha and beta subunits can result in multiple hemoglobin tetramers
with different functions. This "hemoglobin multiplicity" may also be a source of adaptive potential
(Brix, Clements, & Wells, 1999; Houston & Gingras-Bedard, 1994; Verde et al., 2012).
Hemoglobin multiplicity may be, at least in part, linked to hemoglobin gene copy number. For
instance, cod fishes living in shallow and unstable environments have more hemoglobin gene
copies than species living in more stable deep-sea environments (Baalsrud, Voje, et al., 2017).
The Southern Ocean is a chronically cold and hyper-oxygenated environment, yet stable marine
habitat. In this environment, the suborder Notothenioidei, which includes the white-blooded
Antarctic icefishes lacking functional hemoglobin (Kim et al., 2019; Near et al., 2006), are the
most common fish species. A reduction of hemoglobin multiplicity prior to cold adaptation
followed by a decline in oxygen affinity has been documented in cold-adapted notothenioids
(Verde, Parisi, & di Prisco, 2006). In contrast, studies of hemoglobin evolution in other polar
species, as well as other Zoarcoidei, remain scarce (di Prisco et al., 1990; Verde et al., 2002;
Weber, Hourdez, Knowles, & Lallier, 2003). Eelpouts and other species in the suborder
Zoarcodei, however, inhabit diverse environments from pole to pole and from the surface to
hadal depths, making them excellent models for understanding hemoglobin evolution across
disparate environments.
645
Across ray-finned fishes in the order Perciformes living in a wide range of habitats, the two
teleost hemoglobin clusters, LA and MN (Opazo, Butts, Nery, Storz, & Hoffmann, 2013), have
followed dramatically different evolutionary paths (Fig. 4). For the most part, the LA cluster has
remained stable across species, with two alpha and one beta-globin genes (Opazo et al. 2013),
and only a few instances of pseudogenization or loss of the beta-globin gene (Fig. 4a). In
contrast, the MN cluster showed dramatic lineage-specific evolution, in line with other teleosts

(Opazo et al., 2013). In most species in our comparison, the MN cluster expanded by multiple
tandem duplications with frequent pseudogenization. In contrast, the genetic organization of the
MN cluster has been relatively stable within Zoarcoidei, with three alpha-globin and two or three
beta-globin genes (Fig. 4b). However, all studied Zoarcoidei share either the pseudogenization
or loss of the *hbbmn2* gene, which is predicted to be functional in all other perciformes included.
Interestingly, all studied Zoarcoidei appear to have three to five hemoglobins (di Prisco et al.,
1990; Verde et al., 2002; Weber et al., 2003) despite a relatively low number of hemoglobin
genes in comparison to other perciformes. Thus, at least for Zoarcoidei, globin gene copy
number does not seem to correlate with hemoglobin multiplicity. The relative stability of the
Zoarcoidei MN cluster is also remarkable as the studied species inhabit similar environments to
other species with highly variable MN clusters. This suggests that adaptation in Zoarcoidei
hemoglobin may have been focused at the gene sequence level rather than duplication as seen
in these other perciformes. The study of hemoglobin should be pursued in more Zoarcoidei
species, including in the globally distributed eelpouts (Hotaling, Borowiec, et al., 2021), to better
understand the molecular mechanisms underlying adaptation to environmental conditions.
667
Evolution of antifreeze proteins
The evolution of antifreeze proteins (AFP) is a hallmark of adaptation to cold environments
across a wide range of taxa (Atıcı & Nalbantoğlu, 2003; Davies et al., 2002). Multiple routes to
AFP evolution have been documented in polar fishes, including the rise of *afpIII* from
neofunctionalization of a duplicated sialic acid synthase b (*nansb*) gene in the Zoarcoidei (Deng
et al., 2010) and the evolution of AFGPs in both notothenioids and Arctic cod (Chen et al.,
1997a, 1997b; Zhuang et al., 2019) via two distinct molecular routes (Baalsrud, Tørresen, et al.,
2017). Independent evolution of AFPs through non-overlapping pathways is one of the most
well-known differences between the adaptation of eelpouts and notothenioids to the Southern
Ocean. However, understanding of AFP evolution within the suborder Zoarcoidei—for which
eelpouts comprise the most speciose family (Zoarcidae)—remains incomplete, largely due to
the complexity of its history with many species and isoforms evolving at both poles and the
complexity of the genomic region underlying it. Recent phylogenetic evidence suggests that
afpIII arose only once, early in the evolution of the Zoarcoidei approximately 18 million years
ago (Hobbs et al., 2020). The lack of *afpIII* evidence in the genome of the monkeyface
prickleback *C. violaceus* (family Stichaeidae) poses an additional question about the evolution
of AFPs in Zoarcoidei. To our understanding, every previously surveyed species in the suborder
has exhibited *apfIII* evidence at the sequence level (at least 14 species from six families, Hobbs
et al., 2020). Given that *C. violaceus* is part of the Zoarcoidei "in group" (Hotaling, Borowiec, et
al., 2021) and is more closely related to other species in the suborder than the Alaskan ronquil
Bathymaster caeruleofasciatus, which has *afpIII* sequence evidence (Hobbs et al., 2020), its
lack of *apfIII* evidence at the sequence level suggests that either the *afpIII* gene did not arise
only once at the base of the clade, it has been horizontally transferred within the clade (e.g.,
Graham, Lougheed, Ewart, & Davies, 2008), or most likely, it has been secondarily lost in *C.*
violaceus. It is also possible that this finding could be due to assembly error, but this is unlikely
given the assembly's high-quality (scaffold N50 = 6.72 Mb, Table S1), lack of N's in the
predicted *afpIII* region, and contiguity of flanking regions (Fig. 5).
695
Gene duplication, rearrangement, and modification has long obscured the study of AFP
evolution (e.g., Deng et al., 2010). *afpIII* copy number varies considerably between closely
related species (Scott, Fletcher, & Davies, 1986) and even within species across their
geographic range (Hew et al., 1988). Short-read genome assemblies limit *afpIII* copy number
inference. For instance, the recent short-read assembly of *A. ocellatus* has *afpIII* sequences on
four different scaffolds which are all ~2 kb in length with identical sequences (including for *afpIII*
genes). However, long-read assemblies can largely ameliorate this issue. Indeed, following the

massive improvements of long-read sequencing for genome assembly, a complete sequence of the complex *afpIII* genome has now been produced (for *P. gunnellus*) with a second, near complete sequence reported in this study. For both Antarctic eelpouts, *O. amberensis* and *L. dearborni*, and the rock gunnell, *P. gunnellus*, a large number of similar but not identical *afpIII* copies (at least 13, 25, and 15 respectively, Fig. 5) may represent an evolutionarily tuned system for depressing the serum freezing point (Kelley, Aagaard, MacCoss, & Swanson, 2010).

Adaptation to the Southern Ocean: new insights from long-read sequencing
We have entered an exciting time for genome biology. It is now possible to characterize complex regions of genomes to better understand evolution across virtually any group of interest. In this study, we used long-read sequencing to generate a high-quality reference assembly for an Antarctic eelpout, *O. amberensis*, the first for the group. We used this new genomic resource to better understand the evolution of *O. amberensis* in the Southern Ocean, particularly as it relates to other Antarctic fish specialists (i.e., notothenioids) and the rest of the suborder Zoarcoidei. We showed evidence of historical shifts in TE activity in *O. amberensis* and other polar fishes, perhaps reflecting an ancient burst of TE activity in response to environmental change. Moreover, we found little overlap between the targets of natural selection in *O. amberensis* and in notothenioids inhabiting the same habitats, indicating that adaptation of both groups to the Southern Ocean appear to have followed largely different trajectories. However, we did find overlap in a few targets of selection, many of which were genes involved in DNA repair or membrane structure. We also observed the relative evolutionary stability of the hemoglobin MN cluster in the Zoarcoidei relative to other teleosts despite these species inhabiting similar environments to species with much more diverse MN clusters. For AFPs, we identified the first species in the suborder Zoarcoidei with no evidence of *afpIII* genes (*C. violaceus*), likely reflecting a lineage-specific loss of the cluster. To further understand the evolution of polar fishes, both within and across groups, it is clear that more dense taxonomic sampling is needed. Snailfishes (order Scorpaeniformes; family Liparidae) occur around the world, often at hadal depths, and are the most species-rich deep-water family in the Southern Ocean, making them another exceptional candidate for comparisons to understand the general rules of Antarctic adaptation.

Beyond polar fish adaptation, the utility of long-read sequencing for understanding complex regions of genomes is clear. The power of these data will increase as accuracy continues to improve (e.g. circular consensus sequencing (i.e., PacBio HiFi, Wenger et al., 2019)) and reads lengthen. The application of long-read technologies to eelpouts and other zoarcids will yield additional high-quality, contiguous sequences of the *afpIII* locus. The limiting factor for understanding questions similar to those posed in this study has likely shifted from generating high-quality assemblies to obtaining samples compatible with high-molecular weight DNA extraction and long-read sequencing. We echo previous studies (e.g., Buckner, Sanders, Faircloth, & Chakrabarty, 2021; Hotaling et al., in press) that the field needs curated specimen vouchers with properly stored tissues and associated metadata to empower the future of genome biology in polar fishes and beyond.

Acknowledgements:
We thank the captain and crew of the ARSV *Laurence M. Gould*, the personnel of the U.S. Antarctic Program Support Contractors for assistance in Chile, at sea, and at Palmer Station, as well as the logistics in Denver, CO for their support related to the Antarctic fieldwork required for this study. We also thank Blair Perry, Kerry McGowan, and Ellie Armstrong for comments that improved the manuscript. We acknowledge funding from NSF awards OPP-1543383 and OPP-1947040 (supporting T.D.), and OPP-1906015 (to J.L.K.) as well as funding from the Antarctic Science Bursary. The RNA sequencing data used in this study was provided as part of a

754 workshop sponsored by the NSF (OPP-1744877) and a Grant in Aid of Research through Long
755 Island University-Post (both to Scott Santagata). We acknowledge computing resources from
756 the Center for Institutional Research Computing at Washington State University and the NSF
757 Extreme Science and Engineering Discovery Environment (XSEDE).
758
References:
Ahn, D.-H., Shin, S. C., Kim, B.-M., Kang, S., Kim, J.-H., Ahn, I., . . . Park, H. (2017). Draft
genome of the Antarctic dragonfish, Parachaenichthys charcoti. *GigaScience, 6*(8),
gix060.
Akbarzadeh, A., Günther, O. P., Houde, A. L., Li, S., Ming, T. J., Jeffries, K. M., . . . Miller, K. M.
(2018). Developing specific molecular biomarkers for thermal stress in salmonids. *BMC*
Genomics, 19(1), 1-28.
Anderson, M. E. (1994). Systematics and osteology of the Zoarcidae (Teleostei: Perciformes).
Ichthyological Bulletin of the J.L.B. Smith Institute of Ichthyology, 60, 1-120.
Andrews, S. (2010). FastQC: a quality control tool for high throughput sequence data.
Atıcı, Ö., & Nalbantoğlu, B. (2003). Antifreeze proteins in higher plants. *Phytochemistry, 64*(7),
1187-1196.
Auvinet, J., Graça, P., Belkadi, L., Petit, L., Bonnivard, E., Dettaï, A., . . . Higuet, D. (2018).
Mobilization of retrotransposons as a cause of chromosomal diversification and rapid
speciation: the case for the Antarctic teleost genus Trematomus. *BMC Genomics, 19*(1),
1-18.
Auvinet, J., Graça, P., Ghigliotti, L., Pisano, E., Dettaï, A., Ozouf-Costaz, C., & Higuet, D.
(2019). Insertion hot spots of DIRS1 Retrotransposon and chromosomal diversifications
among the Antarctic Teleosts Nototheniidae. *International journal of molecular sciences,*
20(3), 701.
Baalsrud, H. T., Tørresen, O. K., Solbakken, M. H., Salzburger, W., Hanel, R., Jakobsen, K. S.,
& Jentoft, S. (2017). De novo gene evolution of antifreeze glycoproteins in codfishes
revealed by whole genome sequence data. *Mol Biol Evol, 35*(3), 593-606.
Baalsrud, H. T., Voje, K. L., Tørresen, O. K., Solbakken, M. H., Matschiner, M., Malmstrøm, M.,
. . . Jentoft, S. (2017). Evolution of hemoglobin genes in codfishes influenced by ocean
depth. *Sci Rep, 7*(1), 1-10.
Beck, E. A., Healey, H. M., Small, C. M., Currey, M. C., Desvignes, T., Cresko, W. A., &
Postlethwait, J. H. (2021). Advancing human disease research with fish evolutionary
mutant models. *Trends in Genetics*.
Benjamini, Y., & Hochberg, Y. (1995). Controlling the false discovery rate: a practical and
powerful approach to multiple testing. *Journal of the Royal statistical society: series B*
(Methodological), 57(1), 289-300.
Bista, I., McCarthy, S. A., Wood, J., Ning, Z., Detrich Iii, H. W., Desvignes, T., . . . Torrance, J.
(2020). The genome sequence of the channel bull blenny, Cottoperca gobio (Günther,
1861). *Wellcome Open Research, 5*.
Bradnam, K. R., Fass, J. N., Alexandrov, A., Baranay, P., Bechner, M., Birol, I., . . . Chikhi, R.
(2013). Assemblathon 2: evaluating de novo methods of genome assembly in three
vertebrate species. *GigaScience, 2*(1), 2047-2217X-2042-2010.
Brix, O., Clements, K., & Wells, R. (1999). Haemoglobin components and oxygen transport in
relation to habitat distribution in triplefin fishes (Tripterygiidae). *Journal of Comparative*
Physiology B, 169(4), 329-334.
Buckner, J. C., Sanders, R. C., Faircloth, B. C., & Chakrabarty, P. (2021). Science Forum: The
critical importance of vouchers in genomics. *Elife, 10*, e68264.
Chen, L., DeVries, A. L., & Cheng, C.-H. C. (1997a). Convergent evolution of antifreeze
glycoproteins in Antarctic notothenioid fish and Arctic cod. *Proceedings of the National*
Academy of Sciences, 94(8), 3817-3822.

Chen, L., DeVries, A. L., & Cheng, C.-H. C. (1997b). Evolution of antifreeze glycoprotein gene
from a trypsinogen gene in Antarctic notothenioid fish. *Proceedings of the National*
Academy of Sciences, 94(8), 3811-3816. doi:10.1073/pnas.94.8.3811
Chen, L., Lu, Y., Li, W., Ren, Y., Yu, M., Jiang, S., . . . Bilyk, K. T. (2019). The genomic basis for
colonizing the freezing Southern Ocean revealed by Antarctic toothfish and Patagonian
robalo genomes. *GigaScience, 8*(4), giz016.
Chénais, B., Caruso, A., Hiard, S., & Casse, N. (2012). The impact of transposable elements on
eukaryotic genomes: from genome size increase to genetic adaptation to stressful
environments. *Gene, 509*(1), 7-15.
Cossins, A., & Prosser, C. (1978). Evolutionary adaptation of membranes to temperature.
Proceedings of the National Academy of Sciences, 75(4), 2040-2043.
Davies, P. L., Baardsnes, J., Kuiper, M. J., & Walker, V. K. (2002). Structure and function of
antifreeze proteins. *Philos Trans R Soc Lond B Biol Sci, 357*(1423), 927-935.
doi:10.1098/rstb.2002.1081
Deng, C., Cheng, C. H., Ye, H., He, X., & Chen, L. (2010). Evolution of an antifreeze protein by
neofunctionalization under escape from adaptive conflict. *Proc Natl Acad Sci U S A,*
107(50), 21593-21598. doi:10.1073/pnas.1007883107
DeVries, A. L., & Steffensen, J. F. (2005). The Arctic and Antarctic polar marine environments.
Fish Physiology, 22, 1-24. doi:10.1016/S1546-5098(04)22001-5
di Prisco, G., D'Avino, R., Camardella, L., Caruso, C., Romano, M., & Rutigliano, B. (1990).
Structure and function of hemoglobin in Antarctic fishes and evolutionary implications.
Polar Biology, 10(4), 269-274.
Feron, R., Zahm, M., Cabau, C., Klopp, C., Roques, C., Bouchez, O., . . . Haffray, P. (2020).
Characterization of a Y-specific duplication/insertion of the anti-Mullerian hormone type II
receptor gene based on a chromosome-scale genome assembly of yellow perch, Perca
flavescens. *Molecular Ecology Resources, 20*(2), 531-543.
Flynn, J. M., Hubley, R., Goubert, C., Rosen, J., Clark, A. G., Feschotte, C., & Smit, A. F.
(2020). RepeatModeler2 for automated genomic discovery of transposable element
families. *Proceedings of the National Academy of Sciences, 117*(17), 9451-9457.
Ge, S. X., Jung, D., & Yao, R. (2020). ShinyGO: a graphical gene-set enrichment tool for
animals and plants. *Bioinformatics, 36*(8), 2628-2629.
Graham, L. A., Lougheed, S. C., Ewart, K. V., & Davies, P. L. (2008). Lateral transfer of a lectin-
like antifreeze protein gene in fishes. *PLoS One, 3*(7), e2616.
doi:10.1371/journal.pone.0002616
Griffiths, H. J. (2010). Antarctic marine biodiversity–what do we know about the distribution of
life in the Southern Ocean? *PLoS One, 5*(8), e11683.
Halder, S., Torrecilla, I., Burkhalter, M. D., Popović, M., Fielden, J., Vaz, B., . . . Wiseman, K.
(2019). SPRTN protease and checkpoint kinase 1 cross-activation loop safeguards DNA
replication. *Nat Commun, 10*(1), 1-18.
Hazel, J. R. (1995). Thermal adaptation in biological membranes: is homeoviscous adaptation
the explanation? *Annual Review of Physiology, 57*(1), 19-42.
Heras, J., Chakraborty, M., Emerson, J., & German, D. P. (2020). Genomic and biochemical
evidence of dietary adaptation in a marine herbivorous fish. *Proceedings of the Royal*
Society B, 287(1921), 20192327.
Hew, C. L., Wang, N., Joshi, S., Fletcher, G., Scott, G., Hayes, P., . . . Davies, P. (1988).
Multiple genes provide the basis for antifreeze protein diversity and dosage in the ocean
pout, Macrozoarces americanus. *Journal of Biological Chemistry, 263*(24), 12049-12055.
Hobbs, R. S., Hall, J. R., Graham, L. A., Davies, P. L., & Fletcher, G. L. (2020). Antifreeze
protein dispersion in eelpouts and related fishes reveals migration and climate alteration
within the last 20 Ma. *PLoS One, 15*(12), e0243273. doi:10.1371/journal.pone.0243273

Holt, C., & Yandell, M. (2011). MAKER2: an annotation pipeline and genome-database
management tool for second-generation genome projects. *BMC Bioinformatics, 12*, 491.
doi:10.1186/1471-2105-12-491
Hotaling, S., Borowiec, M. L., Lins, L. S., Desvignes, T., & Kelley, J. L. (2021). The
biogeographic history of eelpouts and related fishes: linking phylogeny, environmental
change, and patterns of dispersal in a globally distributed fish group. *Molecular*
phylogenetics and evolution, 107211.
Hotaling, S., Kelley, J. L., & Frandsen, P. B. (in press). Towards a genome sequence for every
animal: where are we now? *Proceedings of the National Academy of Sciences*.
Hotaling, S., Sproul, J., Heckenhauer, J., Powell, A., Larracuente, A., Pauls, S., . . . Frandsen,
P. (2021). Long-reads are revolutionizing 20 years of insect genome sequencing.
Genome Biology and Evolution, evab138.
Houston, A., & Gingras-Bedard, J. H. (1994). Variable versus constant temperature acclimation
regimes: effects on hemoglobin isomorph profile in goldfish, Carassius auratus. *Fish*
physiology and biochemistry, 13(6), 445-450.
Kelley, J. L., Aagaard, J. E., MacCoss, M. J., & Swanson, W. J. (2010). Functional
diversification and evolution of antifreeze proteins in the antarctic fish Lycodichthys
dearborni. *J Mol Evol, 71*(2), 111-118. doi:10.1007/s00239-010-9367-6
Kim, B.-M., Amores, A., Kang, S., Ahn, D.-H., Kim, J.-H., Kim, I.-C., . . . Lee, J. (2019). Antarctic
blackfin icefish genome reveals adaptations to extreme environments. *Nature Ecology &*
Evolution, 3(3), 469-478.
Koren, S., Walenz, B. P., Berlin, K., Miller, J. R., Bergman, N. H., & Phillippy, A. M. (2017).
Canu: scalable and accurate long-read assembly via adaptive k-mer weighting and
repeat separation. *Genome research, 27*(5), 722-736.
Korf, I. (2004). Gene finding in novel genomes. *BMC Bioinformatics, 5*(1), 1-9.
Krueger, F. (2015). Trim Galore!: a wrapper tool around Cutadapt and FastQC to consistently
apply quality and adapter trimming to FastQ files. Babraham Bioinformatics, Cambridge,
United Kingdom. In.
Li, H. (2013). Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM.
arXiv, 1303.3997.
Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., . . . Durbin, R. (2009). The
sequence alignment/map format and SAMtools. *Bioinformatics, 25*(16), 2078-2079.
Li, X., Trinh, K.-Y., Hew, C. L., Buettner, B., Baenziger, J., & Davies, P. L. (1985). Structure of
an antifreeze polypeptide and its precursor from the ocean pout, Macrozoarces
americanus. *Journal of Biological Chemistry, 260*(24), 12904-12909.
Longo, G. C., Lam, L., Basnett, B., Samhouri, J., Hamilton, S., Andrews, K., . . . Nichols, K. M.
(2020). Strong population differentiation in lingcod (Ophiodon elongatus) is driven by a
small portion of the genome. *Evol Appl, 13*(10), 2536-2554.
Löytynoja, A. (2014). Phylogeny-aware alignment with PRANK. In *Multiple sequence alignment*
methods (pp. 155-170): Springer.
Marks, R. A., Hotaling, S., Frandsen, P. B., & VanBuren, R. (in press). Representation and
participation across 20 years of plant genome sequencing. *Nature Plants*.
Melsted, P., & Pritchard, J. K. (2011). Efficient counting of k-mers in DNA sequences using a
bloom filter. *BMC Bioinformatics, 12*(1), 1-7.
Møller, P. R., Nielsen, J. G., & Anderson, M. E. (2005). Systematics of polar fishes. *Fish*
Physiology, 22, 25-78. doi:10.1016/S1546-5098(04)22002-7
Moore, J. K., Abbott, M. R., & Richman, J. G. (1999). Location and dynamics of the Antarctic
Polar Front from satellite sea surface temperature data. *Journal of Geophysical*
Research: Oceans, 104(C2), 3059-3073.
Moran, A. L., & Woods, H. A. (2012). Why might they be giants? Towards an understanding of
polar gigantism. *Journal of Experimental Biology, 215*(12), 1995-2002.

Moran, R. L., Catchen, J. M., & Fuller, R. C. (2020). Genomic resources for darters (Percidae:
Etheostominae) provide insight into postzygotic barriers implicated in speciation. *Mol*
Biol Evol, 37(3), 711-729.
Mu, Y., Bian, C., Liu, R., Wang, Y., Shao, G., Li, J., . . . Ao, J. (2021). Whole genome
sequencing of a snailfish from the Yap Trench (~ 7,000 m) clarifies the molecular
mechanisms underlying adaptation to the deep sea. *PLoS Genet, 17*(5), e1009530.
Near, T. J., Dornburg, A., Kuhn, K. L., Eastman, J. T., Pennington, J. N., Patarnello, T., . . .
Jones, C. D. (2012). Ancient climate change, antifreeze, and the evolutionary
diversification of Antarctic fishes. *Proceedings of the National Academy of Sciences,*
109(9), 3434-3439. doi:10.1073/pnas.1115169109
Near, T. J., Parker, S. K., & Detrich III, H. W. (2006). A genomic fossil reveals key steps in
hemoglobin loss by the antarctic icefishes. *Mol Biol Evol, 23*(11), 2008-2016.
Opazo, J. C., Butts, G. T., Nery, M. F., Storz, J. F., & Hoffmann, F. G. (2013). Whole-genome
duplication and the functional diversification of teleost fish hemoglobins. *Mol Biol Evol,*
30(1), 140-153.
Pećina-Šlaus, N., Kafka, A., Salamon, I., & Bukovac, A. (2020). Mismatch repair pathway,
genome stability and cancer. *Frontiers in molecular biosciences, 7*, 122.
Pischedda, A., Ramasamy, K. P., Mangiagalli, M., Chiappori, F., Milanesi, L., Miceli, C., . . .
Lotti, M. (2018). Antarctic marine ciliates under stress: superoxide dismutases from the
psychrophilic Euplotes focardii are cold-active yet heat tolerant enzymes. *Sci Rep, 8*(1),
1-13.
Powers, D. A. (1980). Molecular ecology of teleost fish hemoglobins: strategies for adapting to
changing environments. *American Zoologist*, 139-162.
R Core Team. (2021). R: A language and environment for statistical computing.
Rabosky, D. L., Chang, J., Title, P. O., Cowman, P. F., Sallan, L., Friedman, M., . . . Coll, M.
(2018). An inverse latitudinal gradient in speciation rate for marine fishes. *Nature,*
559(7714), 392-395. doi:10.1038/s41586-018-0273-1
Rice, P., Longden, I., & Bleasby, A. (2000). EMBOSS: the European molecular biology open
software suite. *Trends in Genetics, 16*(6), 276-277.
Robinson, J. T., Thorvaldsdóttir, H., Winckler, W., Guttman, M., Lander, E. S., Getz, G., &
Mesirov, J. P. (2011). Integrative genomics viewer. *Nature Biotechnology, 29*(1), 24-26.
Rothfield, L. I. (2014). *Structure and function of biological membranes*: Academic Press.
Ruud, J. T. (1954). Vertebrates without erythrocytes and blood pigment. *Nature, 173*(4410),
848-850.
Scott, G. K., Fletcher, G. L., & Davies, P. L. (1986). Fish Antifreeze Proteins: Recent Gene
Evolution. *Canadian Journal of Fisheries and Aquatic Sciences, 43*(5), 1028-1034.
doi:10.1139/f86-128
Scott, G. K., Hew, C. L., & Davies, P. L. (1985). Antifreeze protein genes are tandemly linked
and clustered in the genome of the winter flounder. *Proc Natl Acad Sci U S A, 82*(9),
2613-2617.
Shao, F., Han, M., & Peng, Z. (2019). Evolution and diversity of transposable elements in fish
genomes. *Sci Rep, 9*(1), 1-8.
Sievers, F., Wilm, A., Dineen, D., Gibson, T. J., Karplus, K., Li, W., . . . Söding, J. (2011). Fast,
scalable generation of high-quality protein multiple sequence alignments using Clustal
Omega. *Molecular systems biology, 7*(1), 539.
Simão, F. A., Waterhouse, R. M., Ioannidis, P., Kriventseva, E. V., & Zdobnov, E. M. (2015).
BUSCO: assessing genome assembly and annotation completeness with single-copy
orthologs. *Bioinformatics, 31*(19), 3210-3212.
Simpson, J. T., & Durbin, R. (2010). Efficient construction of an assembly string graph using the
FM-index. *Bioinformatics, 26*(12), i367-373. doi:10.1093/bioinformatics/btq217
Smit, A., Hubley, R., & Green, P. (2015). RepeatMasker Open-4.0. 2013–2015. In.

Smit, A. F., & Hubley, R. (2008). RepeatModeler Open-1.0. In.
Stamatakis, A. (2014). RAxML version 8: a tool for phylogenetic analysis and post-analysis of
large phylogenies. *Bioinformatics, 30*(9), 1312-1313. doi:10.1093/bioinformatics/btu033
Stanke, M., & Waack, S. (2003). Gene prediction with a hidden Markov model and a new intron
submodel. *Bioinformatics, 19*(suppl_2), ii215-ii225.
Van Houten, B., Santa-Gonzalez, G. A., & Camargo, M. (2018). DNA repair after oxidative
stress: current challenges. *Current opinion in toxicology, 7*, 9-16.
Verde, C., Carratore, V., Riccio, A., Tamburrini, M., Parisi, E., & Di Prisco, G. (2002). The
functionally distinct hemoglobins of the Arctic spotted wolffish Anarhichas minor. *Journal*
of Biological Chemistry, 277(39), 36312-36320.
Verde, C., Giordano, D., di Prisco, G., & Andersen, Ø. (2012). The haemoglobins of polar fish:
evolutionary and physiological significance of multiplicity in Arctic fish. *Biodiversity, 13*(3-
4), 228-233.
Verde, C., Parisi, E., & di Prisco, G. (2006). The evolution of thermal adaptation in polar fish.
Gene, 385, 137-145.
Vurture, G. W., Sedlazeck, F. J., Nattestad, M., Underwood, C. J., Fang, H., Gurtowski, J., &
Schatz, M. C. (2017). GenomeScope: fast reference-free genome profiling from short
reads. *Bioinformatics, 33*(14), 2202-2204.
Weber, R. E., Hourdez, S., Knowles, F., & Lallier, F. (2003). Hemoglobin function in deep-sea
and hydrothermal-vent endemic fish: Symenchelis parasitica (Anguillidae) and
Thermarces cerberus (Zoarcidae). *Journal of Experimental Biology, 206*(15), 2693-2702.
Wenger, A. M., Peluso, P., Rowell, W. J., Chang, P.-C., Hall, R. J., Concepcion, G. T., . . .
Olson, N. D. (2019). Accurate circular consensus long-read sequencing improves variant
detection and assembly of a human genome. *Nature Biotechnology, 37*(10), 1155-1162.
Willenborg, C., Jing, J., Wu, C., Matern, H., Schaack, J., Burden, J., & Prekeris, R. (2011).
Interaction between FIP5 and SNX18 regulates epithelial lumen formation. *Journal of*
Cell Biology, 195(1), 71-86.
Xu, L., Dong, Z., Fang, L., Luo, Y., Wei, Z., Guo, H., . . . Xia, Q. (2019). OrthoVenn2: a web
server for whole-genome comparison and annotation of orthologous clusters across
multiple species. *Nucleic Acids Res, 47*(W1), W52-W58.
Xu, S., Wang, J., Guo, Z., He, Z., & Shi, S. (2020). Genomic convergence in the adaptation to
extreme environments. *Plant communications*, 100117.
Yang, Z. (2007). PAML 4: phylogenetic analysis by maximum likelihood. *Mol Biol Evol, 24*(8),
1586-1591.
Zhuang, X., Yang, C., Murphy, K. R., & Cheng, C.-H. C. (2019). Molecular mechanism and
history of non-sense to sense evolution of antifreeze glycoprotein gene in northern
gadids. *Proceedings of the National Academy of Sciences, 116*(10), 4400-4405.
994

995 **Data accessibility:**
996 The genome assembly, raw sequence data, and associated files for *O. amberensis* are
997 deposited under NCBI BioProject #PRJNA701078 and this study's GitHub
998 (http://github.io/oamberensis_genome).
999

1000 **Author contributions:**
1001 S.H. and J.L.K. conceived of the study. L.S.F.L. performed laboratory work. S.H., T.D., J.S.S.
1002 and J.L.K. performed analyses. S.H. led the writing of the manuscript with considerable help
1003 from T.D., J.S.S., and J.L.K. T.D. provided the *O. amberensis* samples. All authors read and
1004 approved the final version of the manuscript.