FoxP3 associates with enhancer-promoter loops

to regulate Treg-specific gene expression

Ricardo N. Ramirez, Kaitavjeet Chowdhary, Juliette Leon, Diane Mathis* and Christophe Benoist*

Department of Immunology, Harvard Medical School, Boston, MA 02115, USA

*Address correspondence to: Christophe Benoist and Diane Mathis
Department of Immunology
Harvard Medical School
77 Avenue Louis Pasteur, Boston, MA 02115
e-mail: cbdm@hms.harvard.edu
Phone: (617) 432-7741

SUMMARY

Gene expression programs are specified by higher-order chromatin structure and enhancer-promoter loops (EPL). T regulatory cells (Treg) identity is dominantly specified by the transcription factor FoxP3, whose mechanism of action is unclear. We applied proximity-ligation with chromatin immunoprecipitation (HiChIP) in Treg and closely related conventional CD4+ T cells (Tconv). EPL identified by H3K27Ac HiChIP showed a range of connection intensity, with some super-connected genes. TF-specific HiChIP showed that FoxP3 interacts with EPLs at a large number of genes, including some not differentially expressed in Treg vs Tconv, but enriched at the core Treg signature loci that it upregulates. FoxP3 association correlates with heightened H3H27Ac looping, as ascertained by analysis of FoxP3-deficient Treg-like cells. There was marked asymmetry in the loci where FoxP3 associated at the enhancer- or the promoter-side of EPLs, with enrichment for different transcriptional cofactors. FoxP3 EPL intensity distinguished gene clusters identified by single-cell ATAC-seq as co-varying between individual Tregs, supporting a direct transactivation model for FoxP3 in determining Treg identity.

One Sentence Summary:

FoxP3 is associated with enhancer-promoter loops in Treg cells, and correlates with heightened enhancer-promoter cross-talk

INTRODUCTION

Looping is the solution that allows the packaging of the gigantic strands of chromosomal DNA into tight nuclear spaces, while ensuring the spatial organization and connectivity essential for the orderly deployment of their coding capacity (*1-3*). Some of the resulting topologies are very large, on the >megabase scale, such as the higher-order compartment domains, and primarily relate to overall organization. More directly associated with the unfolding of the genome's coding potential are the enhancer-promoter loops (EPL) that bring distal enhancers into close proximity to transcriptional startsites (TSS) in the 3D space, enabling their interactions to regulate transcriptional programs (*4-6*). Rewiring of EPLs occurs during differentiation in several mammalian systems, ranging from early differentiation to olfactory receptor allelic choice (*7-10*). It has been proposed (*5*) that programmatic shifts associated with differentiation involve new enhancer-promoter connectivity, while the fast responses to stimuli or growth factors exploit pre-existing contacts, by modifying the activity of associated proteins (*11-13*). This dichotomy may not be as sharp as stated, and it is unclear how much EPL rewiring exists between closely related cells. Loops are dynamic structures, and the burst-like nature of transcriptional output may correspond to transient enhancer-promoter interactions (*14*). Finally, how sequence-specific transcription factors (TFs) coopt the EPL landscape remains mostly conjectural. Intuitively, fast-acting TFs whose mechanism of action relies on nuclear translocation (e.g. NF-AT) or activation at the cell membrane (e.g. STATs) likely operate on pre-existing EPLs. Other TFs might, instead, provoke the formation of new EPLs by dimerization (*15*) or macromolecular complex formation. A corollary question is whether an EPL is intrinsically sufficient for transactivation, or only a framework onto which activation mechanisms latch (*4, 5*). The regulatory function of TFs operating via EPLs has been demonstrated for both cell-specific and general TFs (e.g. Klf4 and Yy1, respectively) (*16, 17*).

T regulatory (Treg) cells are a branch of CD4+ T cells, dominant negative regulators of many facets of the immune system (*18, 19*). They also control extra-immunologic consequences of inflammation in several organs (*20*). Consistent with these pleiotropic functions, several phenotypic variants of FoxP3+ Tregs exist, with tuned transcriptomes that adapt to their tissue localization and effector functions (*21*). FoxP3, a member of the forkhead/winged-helix family of TFs, is quasi-exclusively expressed in Treg cells, and plays a central role in determining their identity and function, and their characteristic transcriptional signature. It is, however, neither completely necessary nor sufficient for Treg determinism (*22-25*), and requires synergistic action from a number of transcriptional cofactors with which it interacts (*26-31*). FoxP3 is not a pioneer factor, as many of the enhancers and open chromatin regions (OCR) to which it binds are already accessible in the T cell lineage prior to FoxP3 expression (*32, 33*), although a smaller group of FoxP3-binding OCRs are activated contemporaneously with Treg differentiation.

While FoxP3 can synergize with several sequence-specific TFs and with chromatin modifiers, its true mechanism of action remains poorly understood. There is debate as to it being an activator or a repressor, and even whether it activates Treg-specific transcription directly or by tuning intermediates. Some have argued that it functions primarily as a transcriptional repressor (*26, 27, 34-36*). The repression of cytokines produced upon activation by conventional CD4+ T cells (Tconv), and in particular interleukin (IL)-2, is a well-established function of FoxP3. On the other hand, results from several experimental systems suggest that FoxP3 also behaves as an activator for many of its transcriptional targets (*31, 37-40*), and that this is its principal mode of action. The "cofactor model" posits that this functional dichotomy depends on the identity of the cofactor(s) with which it interacts in controlling specific targets (*27, 31, 38*). Super-resolution microscopy does show FoxP3 associated with activating or repressive co-factors in different "transcriptional hubs" in the nucleus of the same Treg cell (*31*).

An important missing piece of the FoxP3 puzzle is an understanding of its interactions with nuclear loop structures, in particular enhancer-promoter connectivity. Chromatin

immunoprecipitation (ChIP-seq) data show FoxP3 binding in the vicinity of TSS and at distant enhancers (*31, 32, 41*), but how FoxP3 inter-relates with EP connectivity is unknown. Does FoxP3 enhance, or disrupt, the formation of specific EPLs in Tregs? Does it dock onto existing loop structures to nucleate regulatory (activating or inhibitory) complexes? Here, we tackle this question by using chromatin proximity ligation with immunoprecipitation (HiChIP), first with anti-H3K27Ac antibody to establish the comparative landscape of EPLs in primary Treg and Tconv cells, two closely related cell-types. We then map FoxP3 molecules onto Treg EPLs. Directly relevant to the mystery of FoxP3 mechanics, the results show that FoxP3-bearing EPLs correspond to strong H3K27Ac loops genomewide, and positively associate with differential transcription between Tconv and Treg cells, consistent with action as a direct transcriptional activator.

RESULTS

Mapping the enhancer-promoter connectome in Treg and Tconv cells

To understand how the short- and long-range enhancer-promoter contacts mediate the cell type-specific expression patterns among mouse CD4+ T cells, we first used the "HiChIP" method (*9*), which identifies chromatin loops associated with a histone mark or a particular TF by combining proximity ligation of DNA strands with immunoprecipitation, quantitated by high-throughput sequencing. Antibodies against H3K27Ac, the histone modification typical of active enhancers, focused the analysis on loops connecting active enhancers. We analyzed primary mouse Treg and Tconv cells directly purified *ex vivo* to avoid imprints from cell culture, aiming for deep libraries to support robust quantitative analysis (200-300 M reads/sample on average). This required sizeable numbers of cells, achieved by magnetic purification from *Foxp3-Thy1.1* reporter mice. In line with current practice (*42, 43*), we quantitated EPL signals as reads that connect the promoter region (+/-500 bp around the TSS) to 5 kb bins tiled +/-250 kb from the gene's TSS. To correct for proximity-biased background in HiChIP data, we compared the EPL signals to a null distribution generated from the interactions of 1000 random intragenic positions, which allowed estimates of significance corrected for relative distance to the TSS (similar to (*42-44*)). Signals were conserved among biological replicates (fig. S1A). A control HiChIP with an irrelevant isotype-matched antibody showed that the vast majority of H3K27Ac+ EPLs were indeed specific (p <0.0001, Fig. S1B).

Across both Treg and Tconv, 115,538 significant EPLs were identified (FDR 0.05) among 9,058 genes selected for reliable mRNA expression in Treg or Tconv cells (Table S1), as illustrated for *Lrrc32* in Fig1A. The median H3K27Ac EPL size was 49.5 kb (S1C), concordant with EPL size estimates in other mammalian cells (*7, 9*). Only 0.5% of H3K27Ac EPLs reached further than 1 Mb from the TSS, indicating that such very-long-range interactions can occur, but

are the exception. Cumulative EPL intensity for each gene was only marginally related to its mRNA levels (fig. S1D), fitting with recent reports (*5, 45*).

A gene's cumulative EPL intensity showed a linear relationship to its number of EPLs (Fig. 1B). At first blush this is mathematically obvious, but on further thought it implies that an active promoter does not simply allocate a fixed interaction potential by alternating between its potential enhancers, but that each enhancer has a given probability of interaction, and that these are cumulated for each gene (plausibly by co-recruitment into a transcriptional hub). This continuity suggests that the regulatory crosstalk of enhancers that loop to the same promoter operates through additive modalities (*46*).

Some genes had far more EPL connectivity than others, over a 10-fold range (Fig. 1C, Table S2). Such a spread was observed in non-immunologic lineages, where some authors proposed to distinguish a class of "Super Interactive Promoters" (*47, 48*). Here, such super-connected genes (523 in Treg, 520 in Tconv) include several Treg and Tconv hallmarks like *Entpd1, Lef1, Ccr7, Cd5,* or *Lrrc32*. These super-interactive genes tend to be associated with "Super Enhancer" elements: of the 65 super-enhancers nominated in Tregs (*41*), 13 included super-interactive promoters (chisq p <0.01).

Differences between Treg and Tconv were readily identified, as exemplified by several prototypical Treg signature transcripts. Several EPLs were observed at the *Ikzf2* promoter in Tregs, but none significantly above background in Tconv (Fig. 1D). ATAC-seq accessibility patterns were essentially identical in Treg and Tconv at these positions, however, suggesting that the locus is poised but not connected in Tconv (which do express *Ikzf2* upon activation). Reciprocal Treg or Tconv patterns were observed for Treg signature genes *Il2rb* and *Pde3b* (fig. S1E). Genome-wide, we identified 10,168 (9% of total EPLs) differentially active EPLs (at fold-change >2, p <0.05; Fig. 1E, fig. S1F). Signal intensities in HiChIP data integrate true loop frequency and stability ("3D signal") with the total abundance of the mark targeted by the immunoprecipitation ("1D signal") (*9*). Total EPL signal in HiChIP data could thus appear to

change, even if there was no actual increase in loop abundance *per se*. Analysis of the variation of total (1D) H3K27Ac signal showed that the Treg/Tconv differences in EPL intensity did represent increased looping, as they could not be accounted for by simple variation in H3K27Ac deposition, which varied very modestly (Fig. 1F). This degree of EPL variation between CD4+ T subsets is generally more extensive than the disparity observed by chromatin accessibility (less than 1% difference in accessibility) (*32*). Some genes that prototypically distinguish Treg and Tconv show differential EPLs (*Lef1, Satb1, Il2ra, Foxp3,* fig. S1G*)*, consistent with observations in human T cells (*9*). Accordingly, pairing of gene expression and cumulative EPL intensity revealed a positive relationship between differential EPL intensity and mRNA levels between the two cell-types (Fig. 1F).

FoxP3 preferentially targets hyperactive enhancer-promoter loops

Having established the overall landscape of EPLs in Treg and Tconv cells, we performed HiChIP with anti-FoxP3 antibody to identify enhancer-promoter interactions that involve FoxP3, for clues to understand how it performs its central role in shaping Treg identity. Data were generated in well-correlated biological duplicates (fig. S2A), with correction for proximity ligation and an irrelevant IgG control to ensure specificity (fig. S2B-C). To help identify the most reliable FoxP3 EPLs, we leveraged existing FoxP3 ChIP-seq data (*32, 41*) (fig. S2D), and filtered putative FoxP3 EPLs against this input. (fig. S2E). We thus identified 13,681 robust FoxP3-decorated EPLs, of ~ 20 kb median length (fig. S2F, 2A). FoxP3 EPLs connected enhancers to the promoters of a surprisingly large number of genes (4,797), confirming that FoxP3 relates to far more loci than just the ~500 genes of the Treg signature. Here again, a subset of genes showed super-connectivity (Fig. 2B), enriched for T cell-specific ontologies (T activation p=10^{-12}, cytokine signaling p=10^{-11}) and Treg signature genes (*Il2ra, Ikzf2, Entpd1*). As illustrated for *Entpd1* and *Il2ra*, prototypical Treg-Up signature genes, abundant FoxP3-decorated EPLs corresponded to FoxP3-binding enhancers identified by ChIP-seq (Fig. 2A, S2G) and overlapped with a subset of

the H3K27Ac EPLs. At both the *Lef1* and *Tcf7* loci, which are under-expressed in Tregs, FoxP3 EPLs also aligned with the positions of H3K27Ac EPLs, the latter being far less intense in Treg than in Tconv cells, consistent with an inhibitory impact of FoxP3 (Fig. 2C, S2H). At the *Il2* locus, two H3K27Ac EPLs were observed in Tconv (fig. S2I), but no loops whatsoever were detected in Treg, whether in association with H3K27Ac or with FoxP3, suggesting that the locus is closed into heterochromatin independently of FoxP3.

We then analyzed the distribution of FoxP3 EPLs, beyond these examples, in genesets that reflect differential expression in Treg cells and FoxP3-dependence. Classic "Treg signature genes", which are known to have FoxP3-dependent and independent components (*39, 49*), showed a modest but significant increase in cumulative FoxP3 EPL intensity compared to expression-matched "neutral" genes (Fig. 2D). This enrichment was accentuated in a set of FoxP3-dependent genes that are up-regulated after short-term FoxP3 transfection (*31*) (Fig. 2E). No such enrichment was seen for down-regulated targets (Fig. 2D,E). The strongest enrichment in FoxP3-associated EPLs was seen for genes that constitute a core signature of FoxP3-dependent genes identified in FOXP3-deficient mice and patients (*25*), Fig. 2F). These results are consistent with the notion that transcriptional upregulation by FoxP3 is directly achieved through enhancer-promoter connections at up-regulated loci.

Overall, plotting the relationship between FoxP3 and H3K27Ac EPLs showed that FoxP3 EPLs correspond to loops with the highest level of H3K27Ac-associated signals (Fig. 2G), suggesting that FoxP3 associates with the most active chromatin. To ensure that this observation was not an experimental artefact, we performed HiChIP for the Yy1 TF, which is considered a structural regulator broadly involved in enhancer-promoter topologies (*17, 50*). In Tregs, Yy1 represses *Foxp3* (*51*), and inhibits transactivation by FoxP3 protein (*31, 51*). The 12,426 Yy1 EPLs thus identified were also enriched in H3K27Ac HiChIP intensity when compared to non-Yy1 chromatin interactions (p<0.05), similar to FoxP3, and consistent with observations in stem cells (*17*). But FoxP3 and Yy1 EPLs showed differential EPL patterns (fig. S4A,B), most strikingly at

Il2 (fig. S2H). In a combined set of FoxP3/or/Yy1 EPLs (n=21,341), 54.3% were differentially represented (>2-fold difference in intensity). The genes associated with FoxP3 or Yy1 EPLs belonged to different functional classes: Yy1 EPLs were enriched in "General transcription" or "Cell Protein Catabolism" ontologies, while FoxP3-associated EPLs belonged to more immune-specific pathways ("Adaptive immune system", "MHC-I presentation"; fig. S4C).

Examination in Tconv of those H3K27Ac EPLs that bind FoxP3 in Tregs revealed that most of them also had high intensity in Tconv (Fig. 2H). Thus, FoxP3 binds to enhancer-promoter loops that are very active in CD4+ T cells in general, and not solely to Treg-specific ones, consistent with the notion that it homes to chromatin locales that are ubiquitously open in T cells (*32, 33*). If FoxP3 locates to Treg EPLs with the highest degree of activation, how does its presence affect them? To address this question most finely, we compared the H3K27Ac EPL intensities of the EPLs in Tregs, where FoxP3 is bound, versus in Tconv where FoxP3 is absent, across bins of FoxP3 EPL intensity. Increasing FoxP3 corresponded to increased H3K27Ac EPL signal at those EPLs in Tregs relative to Tconv (Fig. 2I), suggesting that FoxP3 may exert a facilitating influence on enhancer-promoter loops.

Direct tuning of FoxP3-dependent transcriptional programs through FoxP3 enhancer-promoter loops

The results are compatible with, but do not prove, the notion that transcriptional activation by FoxP3 is actuated directly through enhancer-promoter connections around its target genes. However, this notion is at odds with the recent claim that FoxP3 shapes Treg identity indirectly, by regulating the abundance of intermediary TFs like TCF1 (encoded by *Tcf7*) (*52*). While we did observe FoxP3-decorated EPLs across the *Tcf7* locus (fig. S2H), the sweeping conclusion that indirect regulation shapes Treg transcriptional identity is likely an overstatement, as discussed in detail elsewhere (*53*). *Some* indirect control probably occurs (since FoxP3 controls several other TFs), but a sizeable portion of the FoxP3-dependent transcriptome is likely to be directly

controlled by FoxP3, including key functional transcripts. It was important, however, to verify that the FoxP3-decorated EPL structures reported here are functionally relevant. To this end, we performed H3K27Ac HiChIP on Treg-like cells sorted from a novel FoxP3-deficient line (hereafter KO-GFP, constructed from the classic FoxP3-ires-GFP reporter mouse (*54*) by a CRISPR-engineered frameshift mutation that eliminates the Forkhead domain resulting in a full *scurfy*-like Treg deficiency phenotype – see Methods). As other FoxP3-deficient mice (see (*25*) for refs), KO-GFP mice contain a sizeable proportion of FoxP3-null Treg-like cells, identifiable by activity of the GFP reporter appended to the disarmed *Foxp3* locus. Heterozygous females with one copy of this locus are protected from disease by the WT *Foxp3* allele (provided by a *Foxp3-Thy1.1* reporter chromosome to facilitate identification). Genomic comparison to cells in which the intact Foxp3-GFP allele (hereafter WT-GFP) is active enables very precise determination of the effect of the FoxP3 deficiency in Treg-like cells, identified by the same reporter configuration. We sorted WT-GFP and KO-GFP Tregs from half-sib heterozygous females (Fig. 3A) and performed H3K27Ac HiChIP to evaluate FoxP3's effects on enhancer-promoter looping.

Lieberman-Aiden et al showed that HiC and HiChIP data, when analyzed by Principal Component Analysis, reveals spatial segregation of open and closed chromatin that forms two genome-wide compartments (*55*). By this measure, FoxP3-deficiency did not affect nuclear compartments on the macro scale, with no change in heterochromatin compartments identified by Principal Component Analysis (Fig. 3B,C). However, loci with rich FoxP3 EPLs, like *Ikzf2* and *Lrrc32,* showed significantly reduced H3K27Ac-decorated loops in FoxP3 KO Tregs (arrows in Fig. 3D, p<0.05); these differences were reproducible in biological replicates (Fig. S3F). As above, this reduction in EPL intensity in KO-GFP Tregs denoted a true loss of connectivity, not merely lower H3K27Ac decoration (Fig. 3E): the drop in H3K27Ac EPL intensity (x-axis) was not accompanied by a drop in total "1D" H3K27Ac signal at these loci. Conversely, FoxP3 down-regulated targets *Lef1* and *Add3* showed increased H3K27Ac EPL signal in KO-GFP Tregs (Fig. S3G). Comparing genomewide H3K27Ac EPL abundance in WT-GFP and KO-GFP Treg-like

cells showed a general concordance, with only a light bulge for the most highly represented EPLs (Fig. 3F). This deviation was confirmed by plotting the KO-GFP/WT-GFP ratio of H3K27Ac EPL intensity across FoxP3 EPL intensity bins (Fig. 3G, p<0.01), showing a clear reduction in signal resulting from the lack of FoxP3 in those EPLs with highest FoxP3 EPL intensity. The conclusion that FoxP3 directly shapes chromatin connectivity, and its transcriptional relevance, were confirmed by the cumulative density plot of KO/WT ratios for FoxP3 target genes defined as above (from direct transactivation studies or the core Treg set - Fig. 3H). Enhancer-promoter loops in FoxP3-upregulated genes were more sensitive than the norm to the absence of FoxP3 (and the converse was true of FoxP3-repressed loci). These results support FoxP3's association to epigenetically active loops, though which it directly influences the Treg-specific gene program.

Two different categories of FoxP3-dependent genes

These results show that, as a group, transcripts whose expression is upregulated by FoxP3 are enriched in FoxP3-adorned EPLs, but this characteristic does not necessarily apply to all such genes. To further discriminate between loci with FoxP3-enhanced expression in Tregs, we compared accessibility in single-cells: similarly regulated genes tend to have correlated activity across individual cells, and correlation analysis can uncover groups of genes with similar regulatory logic (*56, 57*). We used single-cell ATAC-seq (scATAC-seq), as chromatin accessibility is more stable than mRNA levels (*57-59*), exploiting a dataset generated for another study (KC, in preparation), with high-quality scATAC-seq profiles for 5,810 splenic Tregs, including rTreg and aTreg types (Fig. 4A). We first defined a FoxP3-dependent geneset from the expression data of van der Veeken at al (*52*), by contrasting Treg-like cells from heterozygous female mice in which FoxP3 deficient or proficient X-chromosomes were active (fig. S4D; 377 upregulated genes in FoxP3 proficient vs deficient at FC>2 and p<0.01). We generated a gene accessibility score from the scATAC-seq data and correlated these scores across all Treg cells in the scATAC dataset. This gene-gene correlation network resolved into two predominant gene clusters (Fig. 4A), which

exhibited a marked imbalance in FoxP3 EPL density (Fig. 4B,C; p<0.01, Table S3). Loci in Cluster1 had high FoxP3 EPL density (Fig. 4B-D), and included many key Treg transcripts (*Il2ra*, *Il2rb*, *Lrrc32, Izumo1r* and *Nrp1*), predominantly characteristic of rTregs; in contrast, genes of Cluster2 showed much lower FoxP3 EPL density, and corresponded mainly to transcripts enriched in aTregs (e.g. *Ccr8*, Fig. 4D). Accordingly, genes from Cluster1 showed higher binding to FoxP3 in published ChIPseq datasets (Fig. 4E), with H3K27Ac EPL intensity also biased for Cluster1 transcripts (Fig. 4F). Overall, Cluster1 genes tended to be expressed at somewhat higher levels than Cluster2 (median 341 tpm, 0.05-0.95 range [27-3576] vs 76, [26-1417]), but genes in both clusters were up-regulated by FoxP3, and over-expressed in WT relative to FoxP3-deficient Treg-like cells (Fig. 4G).

We then asked, across all single-cell, how the expression of individual genes of Clusters 1 and 2 (approximated by the "gene accessibility score") correlated with that of FoxP3. The rationale was that, while all these genes appear FoxP3 dependent (as defined by the "sledgehammer" effect of FoxP3 deletion) their instant expression may respond differently to fluctuations in FoxP3 between individual cells. Interestingly, Cluster 1 genes had some positive correlation with Foxp3 (Fig. 4H). In contrast, genes of Cluster2, that are largely devoid of FoxP3-decorated EPLs, lacked this correlation (was even slightly negative on average), This dichotomy might actually be what one would expect if Cluster1 were under direct and positive activation by FoxP3, while Cluster2 genes responded to indirect control by FoxP3. Thus, the partition of FoxP3 dependent genes into clusters independently marked by FoxP3 EPL density suggests that different regulatory mechanisms are at play in the different groups of FoxP3-dependent genes, with different levels of direct FoxP3 involvement.

Distinct FoxP3 interaction at promoters and enhancers

FoxP3 binding across the Treg genome occurs at distal enhancers, as well as in proximal promoter regions (*32, 41*). It is unknown whether FoxP3 functions similarly in both locations. We

assessed whether FoxP3 bound to the promoter or enhancer ends of the loops, based on the consensus ChIP-seq map described above. EPLs associated with FoxP3 on either promoter or enhancer sides, or both (Fig. 5A). This dichotomy was unrelated to the clusters of Fig. 4, and not merely due to sparse sampling or arbitrary thresholds, as FoxP3 binding was truly absent on the "other" side (Fig. 5B). Further, EPLs with FoxP3 on both sides were not a random occurrence, as they occurred more frequently than through chance association (chisq p<0.01). High chromatin accessibility and H3K27Ac signal intensity coincided with the side of the EPL bound by FoxP3 (fig. S5A), confirming FoxP3's association with the most highly active chromatin. A clue as to the significance of FoxP3 placement relative to EPL ends came from Gene Ontology analysis of the corresponding genes. Generic programs linked to nucleic acid metabolism (e.g. Cell Cycle, DNA repair) dominated for genes with promoter-side FoxP3 EPL, while more immune-specific ontologies (Cytokine signaling and adaptive immune system) were enriched among enhancer- and dual-sided modes (Fig. 5C, Table S4). FoxP3-dependent loci belonging to Cluster1 described above were predominantly associated with enhancers (only 3 of 24 were of the promoter-side category).

FoxP3 collaborates with other TFs, these interactions determining its transcriptional specificity (*26, 27, 31*), and their binding motifs are enriched in the vicinity of FoxP3 binding sites (*32, 41*). FoxP3 positioned at either the promoter or enhancer sides of EPLs appears to interact with distinct cofactors (Fig. 5D, S5B), as shown by motif enrichment analysis (using as background comparators all TSS and all non-TSS OCRs, respectively). TATA-box motifs were enriched at promoter positions, while Runx and Ctcf motifs were over-represented at enhancers, and motifs for SP1, Ets, Irf/Stat and the Nuclear Receptor family factors (Rorα/γ) were enriched on both sides. Together, these results delineate different aspects of FoxP3-mediated regulation, with distinctive transcription programs and TF cofactor preference across FoxP3 enhancer-promoter loops.

DISCUSSION

To better understand the program controlled by the lineage-identifying factor FoxP3, we explored enhancer-promoter connectivity in true Treg and Tconv cells *ex vivo*, and how FoxP3 is associated with this architecture. The results lay out the landscape of enhancer elements that are connected to promoters of different genes in CD4+ T cells, and should thus serve as a roadmap to support cis-regulation studies in these important cells. Most importantly, and reinforcing the identity of FoxP3 as a positive transactivator, they establish how FoxP3 associates with a wide range of active EPLs, and positively potentiates their representation for a set of genes that are central to Treg cell physiology.

Previous models of enhancer regulation (*46, 60-62*) have hypothesized additive or even multiplicative enhancer activity. In the primary cells analyzed here, there was a significant but modest correlation between enhancer-promoter connectivity and mRNA levels in these two closely related CD4+ T cells. The absence of a linear relationship is consistent with the notion that simply establishing a connection between an enhancer and a promoter is not sufficient for transactivation, but that secondary events (recruitment of additional factors, post-translational modification of these cofactors) are necessary to functionally potentiate the connection. Cumulative EPL intensities for individual genes distributed as a very broad range, with a subset of highly connected T-cell genes, in line with recent work in neurons which identified a subset of "super-interactive promoters" (*48*). Such highly connected loci may be rationalized by the need for more complex transcriptional hubs for variegated regulation.

A key goal was to understand whether and how FoxP3 is involved with enhancer-promoter connectivity. Does it promote loop formation, or tune the transactivation potential of pre-established loops - the latter in keeping with the observation that, unlike other TFs that specify immunocyte lineages such as PU.1 or Pax5, FoxP3 opportunistically exploits already open chromatin and has limited "pioneer" activity - or does it have little influence, only acting indirectly,

as suggested elsewhere (*52*). Generally, FoxP3 associated with the most epigenetically active chromatin, as did Yy1. Several lines of evidence established that, once associated with an EPL, FoxP3 mostly exerts an enhancing influence on enhancer-promoter connectivity, e.g. by stabilizing them: (i) H3K27Ac EPLs that are also associated with FoxP3 in Tregs are more intense in Tregs than they are in Tconvs; (ii) these same EPLs lose intensity when FoxP3 is genetically inactivated; in other words, when FoxP3 is absent, these enhancers are less connected to their cognate promoters. The functional relevance of these effects is established by the observation that these enhanced connections are found for genes whose expression is known to be positively transactivated by FoxP3.

Different generations of chromatin immunoprecipitation (*31, 32, 41, 52, 63, 64*) showed that FoxP3 associates with many genomic regions that are nowhere near the Treg-specific loci. In this, FoxP3 is not unlike many other TFs (*65*). Our results now show that these apparently inactive locations actually correspond to true enhancer-promoter connections, even at loci that have no clear Treg relevance (38% of Treg-neutral genes have a FoxP3-decorated EPL). An interpretation consistent with the "cofactor model" is that these positions simply lack the appropriate cofactors or epigenetic modifying enzymes, or that FoxP3 or its cofactors are not suitably modified at these locations in order to actually influence transcription. FoxP3 may be truly inert at these locations, or it might provide buffering or stabilizing influence (*61, 62, 65*). At FoxP3-dependent loci that are over-expressed in Treg cells, high-intensity FoxP3 EPLs observed at many core signature genes, which strongly suggest a positive and direct transactivation, in contrast to the indirect model recently proposed (*52*). In contrast, an indirect modality for FoxP3-dependent repression is consistent with the dearth of FoxP3 EPLs at the *Il2* locus. But indirect repression by FoxP3 is not a general characteristic of Treg-underexpressed loci, as *Lef1* and *Tcf7* genes themselves show abundant FoxP3-associated EPLs, consistent with the notion that FoxP3 can act as a direct repressor when paired with the appropriate cofactors (*31*).

Finally, the dichotomy between loci at which FoxP3 associates on the enhancer or the promoter sides of EPLs implies a geographical variation of its direct regulatory functions. This demarcation parallels recent results in B cells where distinct multi-TF hubs bind to promoter versus distal enhancer sites (*66*). The enrichment analysis (Fig. 5D) suggests that the factors with which FoxP3 associates on the enhancer or the promoter sides are different, suggesting different modes of operation. It is also possible that this repartition is dynamic (per (*12, 13*)), and one could speculate for instance that, upon rapidly-acting cell triggering, FoxP3 already bound to the promoter region serves to further stabilize/potentiate an EPL through dimerization with another FoxP3 molecule recruited to the enhancer side.

In terms of how FoxP3 determines Treg identity, parsing the group of FoxP3-dependent genes by correlation analysis of chromatin accessibility across single Tregs distinguished two groups of genes, among those positively controlled by FoxP3. The larger cluster was associated with abundant FoxP3-decorated EPLs, indicative of direct and positive transactivation. The second, smaller, cluster grouped genes associated with markedly less FoxP3-bound enhancer-promoter loops. Interestingly, genes of this second cluster include genes that are over-expressed in aTreg relative to rTreg, and these might correspond to indirect control (*52*).

These results, and those from (*25*) and (*52*), suggest a multimodal set of actions through which FoxP3 helps determine Treg identity and stability: (i) direct control, both activating and repressive, of a range of target loci, in particular a core set of transcripts essential for Treg homeostasis; (ii) repression of TFs that otherwise favor the Tconv program (iii) stabilization of a Treg program that is not directly FoxP3-dependent.

METHODS

Mice

C57BL/6J (B6), B6.*Foxp3*$^{Thy1.1}$ (*67*) mice were obtained from the Jackson Laboratory and bred in the SPF facility at Harvard Medical School (HMS). All experimentation was performed following animal protocols approved by the HMS Institutional Animal Use and Care Committee (protocols IS00000054). FoxP3-deficient mice (B6.*Foxp3*$^{fs327-gfp/Doi}$) were generated by CRISPR mutagenesis of the B6.*Foxp3*$^{ires-GFPKuch}$ line (*54*) as described elsewhere (Leon et al, in preparation). Briefly, the Foxp3-ires-GFP allele carries a 1 bp insertion in exon 11 (NM_054039.2), at the position encoding FoxP3 amino-acid 327, leading to a frameshift mutation that would produce a truncated protein terminating after 34 amino-acids, and thus lacking the essential carboxy-terminal forkhead DNA-binding domain (fig. S3A). Males carrying the mutant allele present with a typical *Scurfy*-like phenotype of FoxP3 deficiency, with a wasting syndrome starting at 12-15 days of age (failure to thrive, cachexia, extensive exfoliative dermatitis, tail necrosis, crusty eyelids and ears, marked lymphadenopathy and splenomegaly) leading to death around 30-35 days. - Heterozygous females are healthy and don't show any clinical signs of inflammation. The germline mutation was maintained onto the B6 background. For the experiments reported here, females carrying either the normal *Foxp3*$^{ires-GFPKuch}$ or the deficient *Foxp3*$^{fs327-gfp}$ alleles were crossed to B6.*Foxp3*$^{Thy1.1}$ father(the FoxP3-Thy1.1 reporter was used to identify in the offspring normal Treg cells that express the WT balancing allele in these crosses. To improve the comparability between parallel fs327 and WT offsprings in these experiments, the *Foxp3*$^{ires-GFPKuch}$ and *Foxp3*$^{fs327-gfp}$ carrier mothers were co-housed and bred to the same B6.*Foxp3*$^{Thy1.1}$ father.

T cell isolation

For analyses in FoxP3 proficient mice, splenic CD4 T cells from FoxP3-Thy1.1 hemizygous male mice were first enriched using CD4 untouched kit (Miltenyi), followed by secondary purification with anti-CD90.1 microbeads (Miltenyi) according to manufacturer's instructions. Tconv (FoxP3-Thy1.1-Tcrb+CD4+CD45+CD19-CD8-CD11b-CD11c-) and Treg cells (FoxP3-Thy1.1+Tcrb+CD4+, CD45+CD19-CD8-CD11b-CD11c-) were sorted on a FACS-Aria to 96% and 94% purity, respectively. For experiments in FoxP3-deficient Tregs (Fig. 3), CD4+ T cells from spleen and peripheral lymph nodes from 6 week-old heterozygous females were first enriched as above by negative magnetic selection (Stem Cell), and Treg-like cells sorted as CD19-TCRβ+CD4+Thy1.1-GFP+ on a Moflo Astrios (Beckman Coulter).

H3K27Ac, FoxP3, Yy1, and IgG HiChIP

HiChIP (*9*) was optimized and performed in biological replicates for all conditions. Briefly, 5×10^6 Tregs were crosslinked for each FoxP3 HiChIP replicate, 2×10^6 million cells for H3K27Ac and Yy1 HiChIPs respectively. Additionally, 5×10^6 million Tregs and Tconvs were crosslinked to include a negative IgG isotype control HiChIP samples. Low input H3K27Ac HiChIP was performed on 2×10^5 crosslinked WT-GFP and KO-GFP Tregs respectively. DNA sonication of fragments to 200-600 bp following MboI digestion was performed using the Covaris M220. Sonicated DNA was pre-cleared using Protein A magnetic beads followed by overnight chromatin immunoprecipitations with 3 ug H3K27Ac (Abcam ab4729), 5 ug FoxP3 (Abcam ab150743), 3 ug Yy1 (Abcam ab109237) and 5 ug rabbit IgG (Cell Signaling 2729S), in a final concentration 0.07% SDS. Libraries were prepared using Illumina Nextera DNA library preparation kit, size-selected using low-melt gel extraction and paired-end sequenced using NextSeq 500 Illumina platform.

scATAC-seq of Tregs

Single cell ATAC-seq (scATAC-seq) data were generated as part of another study (KC, in preparation). Briefly, splenic regulatory T cells (CD4+, TCRβ+, FoxP3-IRES-GFP+) and T conventional cells (CD4+, TCRβ+, FoxP3-IRES-GFP-) were sorted from 6-8 week old B6 FoxP3-IRES-GFP mice into DMEM + 2% FCS, then resuspended in PBS + 0.04% BSA. Nuclei isolation, transposition, GEM generation, and library construction targeting capture of ~12000 cells (75% Tregs, 25% Tconv) were carried out as detailed in the Chromium Next GEM Single Cell ATAC manual (10x Genomics). Libraries were pooled and sequenced on an Illumina NovaSeq to a final median depth of approximately 30,000 reads per cell (~80% saturation).

RNA-seq expression data

Treg and Tconv RNA-seq biological replicates were obtained from previous work (*33*). RNA-seq counts were used for all gene quantification. Genes were filtered based on replicated and cell-type expression; > 10 counts in both replicates, and in at least 1 condition.

ChIP-seq data

ChIP-seq datasets (*41*); Runx1, FoxP3, Ets1, Smc1a (Cohesin), (*32*); FoxP3, were mapped to mm10 genome using bowtie2 (*68*). BigWig files were generated using deeptools2 (*69*).

ATAC-seq data

Treg and Tconv ATAC-seq datasets were generated in previous work (*33*).

scATAC-seq analysis

Fastq data were aligned to the mm10 reference genome and quantified per cell using Cell Ranger ATAC software (10x Genomics, v1.2). Peak by cell count matrices were generated by mapping reads to open chromatin regions previously defined by the ImmGen consortium (*33*). Data

analysis was performed using Signac v1.1 **(70).** For quality control, only cells with at least 4000

fragments per cell, greater than 55 percent reads in peaks, TSS enrichment score greater than 2,

nucleosome signal less than 10, and ratio of blacklist-region reads less than 0.05 were retained

for further analysis. Putative doublets identified by ArchR v0.9.3 **(71)** and non-Treg, non-Tconv

contaminant cells were also removed. We used the latent semantic indexing approach as

previously described (*59, 72*). Binarized count matrices were normalized using the term

frequency-inverse document frequency (TF-IDF) transformation and reduced in dimensionality by

singular value decomposition (SVD). As the first SVD component was highly correlated with

sequencing depth, components 2-30 were used to create a two-dimensional UMAP visualization.

A gene accessibility score was determined from Treg scATAC-seq data with ArchR v0.9.3, using

an exponentially weighted function that accounts for the activity of distal OCRs in a distance-

dependent manner (*71*) and computes a single measure of a gene's accessibility. A gene-gene

correlation was next determined from the log-normalized gene scores and visualized using a

force-directed network (edge-weighted using correlation coefficients). The network was filtered to

drop weak correlations (-0.2 > and < 0.2). Network clustering was performed using correlation

coefficients (edge attribute) and FoxP3 EPL intensities (node attribute). Network visualization and

clustering were performed using Cytoscape v3.8.0 (*73*).

CAGE data

Treg CAGE data were downloaded from the FANTOM repository (*74*).

H3K27Ac, FoxP3 and Yy1 HiChIP data processing and enhancer-promoter looping

HiChIP paired-end reads were aligned to the mm10 genome using the HiC-Pro pipeline (*75*).

Default settings were used for removal of duplicated reads, assignment to MboI restriction

fragments and filtering for valid interactions. To assess enhancer-promoter loops (EPLs) in

HiChIP data, we computed +/-250 kb contacts from the gene TSS for 9,058 expressed genes in

Tregs and Tconv using the hicPlotViewpoint (*76*) function at a resolution of 5 kb. To assess a background proximity ligation frequency for each HiChIP experiment, we first generated a ligation probability background by sampling 1000 randomly selected intragenic regions more than 10 kb from the promoter and computing the relative counts using a window of +/-250 kb in each HiChIP experiment. For each respective distance position, a Z-statistic was computed using the HiChIP background estimation and adjusted p-values were calculated to assess statistical significance of the contact. This procedure removed from 50-80% of noisy and biased chromatin interactions. Significant EPLs were considered if demonstrated enrichment across replicates, passing an FDR of 5% for H3K27Ac EPLs (n=115,538) and 10% for Yy1 (12,426 EPLs), and enriched with respect to the negative control IgG HiChIP. In a similar manner, FoxP3 EPLs were also further analyzed using using a reproducible set of overlapping 13,982 FoxP3 ChIP-seq peaks (*32, 41*) as anchors. All FoxP3 EPLs (10% FDR) were supported by FoxP3 ChIP-seq binding for at least one end of the FoxP3 loop. 1D H3K27Ac HiChIP analysis was performed using all reads (including re-ligation and dangling-ends, but excluding extrachromosomal contacts) and converted into tabular format (bedpe, using cLoops2 (*77*)) for downstream analysis. For each H3K27Ac EPL, 1D HiChIP signal was established for the both TSS and for distal anchor, and these summed. Per-gene 1D H3K27Ac cumulative signal was calculated by summing 1D signal corresponding to all respective EPLs for each gene. Compartment analysis of WT-GFP and KO-GFP HiChIP was performed at 250 kb resolution using the eigenvector function on KR normalized maps (*78*).

HiChIP experimental metrics

Sample	Total pairs processed	Unique pairs aligned	Cis_intrachromosomal %	Pearson (Inter-replicate)
H3K27Ac Treg R1	172312254	68%	72%	0.85
H3K27Ac Treg R2	140199897	68%	72%	
H3K27Ac Tcon R1	133066435	69%	72%	0.84
H3K27Ac Tcon R2	114716225	70%	72%	
Foxp3 Treg R1	53374194	63%	69%	0.82
Foxp3 Treg R2	51874296	62%	69%	
Yy1 Treg R1	37509324	61%	68%	0.8
Yy1 Treg R2	30881902	62%	68%	
IgG	8328069	59%	69%	
H3K27Ac WT GFP R1	121381491	58%	70%	0.85
H3K27Ac WT GFP R2	139482053	53%	69%	
H3K27Ac KO GFP R1	99594436	63%	72%	0.8
H3K27Ac KO GFP R2	108433482	60%	72%	

Cumulative EPL intensities determined for genes

To measure the overall effect of the enhancer-promoter structure for a given gene, we calculated a single cumulative value by summing the respective HiChIP EPL intensities (H3K27Ac, FoxP3, Yy1). Additionally, the degree in connectivity was calculated by summing the total number of EPLs per gene and compared both with genome-wide expression and binned mRNA quantiles.

Super interactive promoter analysis

To identify Super interactive promoters from Treg and Tconv H3K27Ac or FoxP3 HiChIP, we followed a similar strategy employed by (*48, 79*). Briefly, for each gene we summed the $\log_{10}$ FDR for respective EPLs to derive a single cumulative score per gene. Plots of the ranked cumulative interaction scores in each cell type and condition were plotted. Significant SIPs were determined based on the point of inflection within the curve (slope is equal to 1).

Differential H3K27Ac EPL analysis

Differential H3K27Ac EPLs were identified using biological replicate EPL counts of Treg and Tconv EPLs (P.adj < 0.05, Fold change > 2) determined using diffloop (*80*). Cumulative H3K27Ac EPL differences and comparisons with gene expression between T cells were mapped for all genes and specifically highlighted for the up- and down- regulated Treg signature program (*39*).

Yy1 and FoxP3 comparative analysis

Yy1 EPL and cumulative intensities were compared with FoxP3 respectively to identify TF-specific interactions (2-Fold, P<0.05). Gene ontology analysis was performed for genes with differences between FoxP3 and Yy1 cumulative EPL intensities (2-Fold) using Metascape (*81*).

Deriving a FoxP3-dependent signature from RNA-seq data

Data from GSE154680 (*52*) were downloaded. Raw read counts tables were normalized by median of ratios method with DESeq2 (*82*) and converted to GCT and CLS formats. Genes with a minimum average read count of 35 in at least one group and a coefficient of variation intra-replicate less than 0.30-0.45 were retained. An uncorrected t-test was used to compute differential gene expression between the different groups from the normalized read counts dataset. The FoxP3 dependent gene signature list was computed by merging the differential expressed genes from the aTreg and rTreg datasets comparing Foxp3.GFP.DTR WT versus Foxp3.GFP KO and Fold change >2 or <0.5 and p-value < 0.01.

Promoter, enhancer, and dual modes in FoxP3 looping

FoxP3 ChIP-seq binding data was used to parse FoxP3 EPLs into those with FoxP3 occupancy at promoter, enhancer or both (dual mode) ends of enhancer-promoter loops. Genomic heatmaps (*69*) were generated for each group either centered on FoxP3 bound positions in FoxP3 EPLs or centered on the strongest Treg ATAC-seq OCR for FoxP3-negative sides of the EPL. Gene

ontology analysis of FoxP3 modes were generated using Metascape (*81*). *De novo* motif analysis was performed using HOMER for FoxP3 bound positions of promoter and enhancer EPL modalities. Significant motifs were determined using as background comparisons using equal numbers (10,000) in TSS and non-TSS OCRs. Motif significance was represented for promoter and enhancer FoxP3 modes using Circos plots (*83*).

Il2 and *Il2ra* centric loop analysis

To resolve higher-resolution sets of chromatin loops, we first defined potential anchors from OCRs with accessibility in Tconv or Tregs (297,076 OCRs). HiChIP loops were defined from our OCR anchor set using Hichipper (*84*) (-peak-pad 250 bp), which adopts a restriction fragment bias-aware approach for preprocessing of HiChIP data to call loops. Briefly, anchors were padded 250 base pairs in either direction and were considered if they overlapped MboI motifs. Padding anchors boosts the number of paired-end tags that can be mapped to loops respectively. We removed self-ligation loops and filtered loops that may be biased due to proximity (< 3kb) or low paired-end tag counts using the Mango (*85*) correction step (stage 5). This correction aggregates paired-end tag counts across all samples and retains significant loops (FDR < 0.01). A union set of loops across all data was generated; 1) loops with a minimum of 2 counts in both biological replicates, 2) supported in at least 1 sample across all HiChIP experiments, and 3) greater than 2-Fold compared to IgG HiChIP controls. We mapped the hichipper defined loops based on overlapping HiChIP EPLs for FoxP3, H3K27Ac and Yy1 centering on the *Il2ra* and *Il2* locus.

Data visualization

Analysis was performed using R-3.5.2 with all plots generated using ggplot2 (https://ggplot2.tidyverse.org) (*86*). Statistical tests are described in their respective sections. Browser and genome views of ATAC-seq and ChIP-seq data were generated using UCSC (*87*) and IGV (*88*). EPL arc plots were generated using DNARchitect (*89*). HiChIP heatmaps were

generated using HiC-Pro by merging valid pairs of replicates and processed as .hic files using the hicpro2juicebox function (*75*) and visualized using Juicebox tools (*78*).

ACKNOWLEDGEMENTS

We thank: Drs. Caleb Lareau, Rick Young, and Nicole El-Ali for insightful discussions; K. Hattori, C. Araneo, K. Seddu, and the HMS Biopolymers team, for help with mice, cell sorting and sequencing. This work was funded by grants from the NIH to CB&DM (AI116834, AI150686), JL was supported by an INSERM Poste d'Accueil and an Arthur Sachs scholarship, RR by NIH supplement AI116834-03S1, and KC by NIGMS-T32GM007753 and a Harvard Stem Cell Institute MD/PhD Training Fellowship.

Data and materials availability: All data and materials are available upon request from the corresponding author. All data needed to evaluate the conclusions in the paper are present in the paper.

REFERENCES

1. R. Schleif, DNA looping. *Science.* **240**, 127-128 (1988).

2. M. Ptashne, How eukaryotic transcriptional activators work. *Nature.* **335**, 683-689 (1988).

3. M. Levine, C. Cattoglio, R. Tjian, Looping back to leap forward: transcription enters a new era. *Cell.* **157**, 13-25 (2014).

4. E. E. M. Furlong, M. Levine, Developmental enhancers and chromosome topology. *Science.* **361**, 1341-1345 (2018).

5. S. Schoenfelder, P. Fraser, Long-range enhancer-promoter contacts in gene expression control. *Nat Rev Genet.* (2019).

6. A. Field, K. Adelman, Evaluating enhancer function and transcription. *Annu. Rev. Biochem.* **89**, 213-234 (2020).

7. A. J. Rubin *et al.*, Lineage-specific dynamic and pre-established enhancer-promoter contacts cooperate in terminal differentiation. *Nat Genet.* **49**, 1522-1528 (2017).

8. B. M. Javierre *et al.*, Lineage-specific genome architecture links enhancers and non-coding disease variants to target gene promoters. *Cell.* **167**, 1369-1384 (2016).

9. M. R. Mumbach *et al.*, Enhancer connectome in primary human cells identifies target genes of disease-associated DNA elements. *Nat. Genet.* **49**, 1602-1612 (2017).

10. K. Monahan, A. Horta, S. Lomvardas, LHX2- and LDB1-mediated trans interactions regulate olfactory receptor choice. *Nature.* **565**, 448-453 (2019).

11. F. Jin *et al.*, A high-resolution map of the three-dimensional chromatin interactome in human cells. *Nature.* **503**, 290-294 (2013).

12. S. Cruz-Molina *et al.*, PRC2 facilitates the regulatory topology required for poised enhancer function during pluripotent stem cell differentiation. *Cell Stem Cell.* **20**, 689-705 (2017).

13. F. Comoglio *et al.*, Thrombopoietin signaling to chromatin elicits rapid and pervasive epigenome remodeling within poised chromatin architectures. *Genome Res.* (2018).

14. T. Fukaya, B. Lim, M. Levine, Enhancer control of transcriptional bursting. *Cell.* **166**, 358-368 (2016).

15. W. Su, S. Jackson, R. Tjian, H. Echols, DNA looping between sites for transcriptional activation: self-association of DNA-bound Sp1. *Genes Dev.* **5**, 820-826 (1991).

16. D. Campigli Di Giammartino *et al.*, KLF4 is involved in the organization and regulation of pluripotency-associated three-dimensional enhancer networks. *Nat. Cell Biol.* **21**, 1179-1190 (2019).

17. A. S. Weintraub *et al.*, YY1 is a structural regulator of enhancer-promoter loops. *Cell.* **171**, 1573-1588 (2017).

18. S. Sakaguchi, T. Yamaguchi, T. Nomura, M. Ono, Regulatory T cells and immune tolerance. *Cell.* **133**, 775-787 (2008).

19. S. Z. Josefowicz, L. F. Lu, A. Y. Rudensky, Regulatory T cells: mechanisms of differentiation and function. *Annu. Rev. Immunol.* **30**, 531-564 (2012).

20. M. Panduro, C. Benoist, D. Mathis, Tissue Tregs. *Annu. Rev Immunol.* **34**, 609-633 (2016).

21. A. R. Muñoz-Rojas, D. Mathis, Tissue regulatory T cells: regulatory chameleons. *Nat Rev Immunol.* https://doi.org/10.1038/s41577-021-00519-w (2021).

22. W. Lin *et al.*, Regulatory T cell development in the absence of functional Foxp3. *Nat. Immunol.* **8**, 359-368 (2007).

23. M. A. Gavin *et al.*, Foxp3-dependent programme of regulatory T-cell differentiation. *Nature.* **445**, 771-775 (2007).

24. L. M. Charbonnier *et al.*, Functional reprogramming of regulatory T cells in the absence of Foxp3. *Nat. Immunol.* **20**, 1208-1219 (2019).

25. D. Zemmour *et al.*, Single-cell analysis of FOXP3 deficiencies in humans and mice unmasks intrinsic and extrinsic CD4[+] T cell perturbations. *Nat Immunol.* **22**, 607-619 (2021).

26. Y. Wu *et al.*, FOXP3 controls regulatory T cell function through cooperation with NFAT. *Cell.* **126**, 375-387 (2006).

27. M. Ono *et al.*, Foxp3 controls regulatory T-cell function by interacting with AML1/Runx1. *Nature.* **446**, 685-689 (2007).

28. A. Kitoh *et al.*, Indispensable role of the Runx1-Cbfβ transcription complex for in vivo-suppressive function of FoxP3[+] regulatory T cells. *Immunity.* **31**, 609-620 (2009).

29. W. Fu *et al.*, A multiply redundant genetic switch 'locks in' the transcriptional signature of regulatory T cells. *Nat Immunol.* **13**, 972-980 (2012).

30. D. Rudra *et al.*, Transcription factor Foxp3 and its protein partners form a complex regulatory network. *Nat Immunol.* **13**, 1010-1019 (2012).

31. H. K. Kwon, H. M. Chen, D. Mathis, C. Benoist, Different molecular complexes that mediate transcriptional induction and repression by FoxP3. *Nat Immunol.* **18**, 1238-1248 (2017).

32. R. M. Samstein *et al.*, Foxp3 exploits a pre-existent enhancer landscape for regulatory T cell lineage specification. *Cell.* **151**, 153-166 (2012).

33. H. Yoshida *et al.*, The cis-Regulatory Atlas of the Mouse Immune System. *Cell.* **176**, 897-912 (2019).

34. L. A. Schubert *et al.*, Scurfin (FOXP3) acts as a repressor of transcription and regulates T cell activation. *J. Biol. Chem.* **276**, 37672-37679 (2001).

35. E. Bettelli, M. Dastrange, M. Oukka, Foxp3 interacts with nuclear factor of activated T cells and NF-kappa B to repress cytokine gene expression and effector functions of T helper cells. *Proc Natl Acad Sci U S A.* **102**, 5138-5143 (2005).

36. A. Arvey *et al.*, Inflammation-induced repression of chromatin bound by the transcription factor Foxp3 in regulatory T cells. *Nat Immunol.* **15**, 580-587 (2014).

37. C. Chen *et al.*, Transcriptional regulation by Foxp3 is associated with direct promoter occupancy and modulation of histone acetylation. *J Biol. Chem.* **281**, 36828-36834 (2006).

38. B. Li *et al.*, FOXP3 interactions with histone acetyltransferase and class II histone deacetylases are required for repression. *Proc Natl Acad Sci U S A.* **104**, 4571-4576 (2007).

39. J. A. Hill *et al.*, Foxp3 transcription-factor-dependent and -independent regulation of the regulatory T cell transcriptional signature. *Immunity.* **27**, 786-800 (2007).

40. A. Ferraro *et al.*, Interindividual variation in human T regulatory cells. *Proc Natl Acad Sci U S A.* **111**, E1111-E1120 (2014).

41. Y. Kitagawa *et al.*, Guidance of regulatory T cell development by Satb1-dependent super-enhancer establishment. *Nat Immunol.* **18**, 173-183 (2017).

42. I. Juric *et al.*, MAPS: Model-based analysis of long-range chromatin interactions from PLAC-seq and HiChIP experiments. *PLoS Comput. Biol.* **15**, e1006982 (2019).

43. S. Bhattacharyya, V. Chandra, P. Vijayanand, F. Ay, Identification of significant chromatin contacts from HiChIP data by FitHiChIP. *Nat. Commun.* **10**, 4221 (2019).

44. F. A. Klein *et al.*, FourCSeq: analysis of 4C sequencing data. *Bioinformatics.* **31**, 3085-3091 (2015).

45. F. Grubert *et al.*, Landscape of cohesin-mediated chromatin loops in the human genome. *Nature.* **583**, 737-743 (2020).

46. H. K. Long, S. L. Prescott, J. Wysocka, Ever-changing landscapes: Transcriptional enhancers in development and evolution. *Cell.* **167**, 1170-1187 (2016).

47. R. Siersbaek *et al.*, Dynamic rewiring of promoter-anchored chromatin loops during adipocyte differentiation. *Mol. Cell.* **66**, 420-435 (2017).

48. M. Song *et al.*, Cell-type-specific 3D epigenomes in the developing human cortex. *Nature.* **587**, 644-649 (2020).

49. N. Sugimoto *et al.*, Foxp3-dependent and -independent molecules specific for CD25+CD4+ natural regulatory T cells revealed by DNA microarray analysis. *Int. Immunol.* **18**, 1197-1209 (2006).

50. J. A. Beagan *et al.*, YY1 and CTCF orchestrate a 3D chromatin looping switch during early neural lineage commitment. *Genome Res.* **27**, 1139-1152 (2017).

51. S. S. Hwang *et al.*, YY1 inhibits differentiation and function of regulatory T cells by blocking Foxp3 expression and activity. *Nat Commun.* **7**, 10789 (2016).

52. J. van der Veeken *et al.*, The transcription factor Foxp3 shapes regulatory T cell identity by tuning the activity of trans-acting intermediaries. *Immunity.* **53**, 971-984 (2020).

53. R. N. Ramirez, K. Chowdhary, C. Benoist, Indirect regulation is not the dominant mode of action for the transcription factor FoxP3 in Treg cells. *Submitted.* (2021).

54. E. Bettelli *et al.*, Reciprocal developmental pathways for the generation of pathogenic effector TH17 and regulatory T cells. *Nature.* **441**, 235-238 (2006).

55. E. Lieberman-Aiden *et al.*, Comprehensive mapping of long-range interactions reveals folding principles of the human genome. *Science.* **326**, 289-293 (2009).

56. S. Ma *et al.*, Chromatin potential identified by shared single-cell profiling of RNA and chromatin. *Cell.* **183**, 1103-1116 (2020).

57. A. T. Satpathy *et al.*, Massively parallel single-cell chromatin landscapes of human immune cell development and intratumoral T cell exhaustion. *Nat. Biotechnol.* **37**, 925-936 (2019).

58. J. D. Buenrostro *et al.*, Single-cell chromatin accessibility reveals principles of regulatory variation. *Nature.* **523**, 486-490 (2015).

59. D. A. Cusanovich *et al.*, A Single-Cell Atlas of In Vivo Mammalian Chromatin Accessibility. *Cell.* **174**, 1309-1324 (2018).

60. C. P. Fulco *et al.*, Activity-by-contact model of enhancer-promoter regulation from thousands of CRISPR perturbations. *Nat. Genet.* **51**, 1664-1669 (2019).

61. C. Buecker, J. Wysocka, Enhancers as information integration hubs in development: lessons from genomics. *Trends Genet.* **28**, 276-284 (2012).

62. M. Lagha, J. P. Bothma, M. Levine, Mechanisms of transcriptional precision in animal development. *Trends Genet.* **28**, 409-416 (2012).

63. Y. Zheng *et al.*, Genome-wide analysis of Foxp3 target genes in developing and mature regulatory T cells. *Nature.* **445**, 936-940 (2007).

64. A. Marson *et al.*, Foxp3 occupancy and regulation of key target genes during T-cell stimulation. *Nature.* **445**, 931-935 (2007).

65. J. Zeitlinger, Seven myths of how transcription factors read the cis-regulatory code. *Curr. Opin. Syst. Biol.* **23**, 22-31 (2020).

66. V. K. Chaudhri *et al.*, Charting the cis-regulome of activated B cells by coupling structural and functional genomics. *Nat Immunol.* **21**, 210-220 (2020).

67. A. Liston *et al.*, Differentiation of regulatory Foxp3[+] T cells in the thymic cortex. *Proc Natl. Acad Sci U S. A.* **105**, 11903-11908 (2008).

68. B. Langmead, S. L. Salzberg, Fast gapped-read alignment with Bowtie 2. *Nat Methods.* **9**, 357-359 (2012).

69. F. Ramirez *et al.*, deepTools2: a next generation web server for deep-sequencing data analysis. *Nucleic Acids Res.* **44**, W160-W165 (2016).

70. T. Stuart, A. Srivastava, C. Lareau, R. Satija, Multimodal single-cell chromatin analysis with Signac. *bioRxiv.* https://doi.org/10.1101/2020.11.09.373613 (2020).

71. J. M. Granja *et al.*, ArchR: An integrative and scalable software package for single-cell chromatin accessibility analysis. *bioRxiv.* https://doi.org/10.1101/2020.04.28.066498 (2020).

72. T. Stuart *et al.*, Comprehensive integration of single-cell data. *Cell.* **177**, 1888-1902 (2019).

73. P. Shannon *et al.*, Cytoscape: a software environment for integrated models of biomolecular interaction networks. *Genome Res.* **13**, 2498-2504 (2003).

74. M. Lizio *et al.*, Update of the FANTOM web resource: expansion to provide additional transcriptome atlases. *Nucleic Acids Res.* **47**, D752-D758 (2019).

75. N. Servant *et al.*, HiC-Pro: an optimized and flexible pipeline for Hi-C data processing. *Genome Biol.* **16**, 259 (2015).

76. J. Wolff *et al.*, Galaxy HiCExplorer: a web server for reproducible Hi-C data analysis, quality control and visualization. *Nucleic Acids Res.* **46**, W11-W16 (2018).

77. Y. Cao *et al.*, cLoops2: a full-stack comprehensive analytical tool for chromatin interactions. *bioRxiv.* https://www.biorxiv.org/content/10.1101/2021.07.20.453068v1 (2021).

78. N. C. Durand *et al.*, Juicer provides a one-click system for analyzing loop-resolution Hi-C experiments. *Cell Syst.* **3**, 95-98 (2016).

79. D. Hnisz *et al.*, Super-enhancers in the control of cell identity and disease. *Cell.* **155**, 934-947 (2013).

80. C. A. Lareau, M. J. Aryee, diffloop: a computational framework for identifying and analyzing differential DNA loops from sequencing data. *Bioinformatics.* **34**, 672-674 (2018).

81. Y. Zhou *et al.*, Metascape provides a biologist-oriented resource for the analysis of systems-level datasets. *Nat. Commun.* **10**, 1523 (2019).

82. M. I. Love, W. Huber, S. Anders, Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol.* **15**, 550 (2014).

83. M. Krzywinski *et al.*, Circos: an information aesthetic for comparative genomics. *Genome Res.* **19**, 1639-1645 (2009).

84. C. A. Lareau, M. J. Aryee, hichipper: a preprocessing pipeline for calling DNA loops from HiChIP data. *Nat Methods*. **15**, 155-156 (2018).

85. D. H. Phanstiel, A. P. Boyle, N. Heidari, M. P. Snyder, Mango: a bias-correcting ChIA-PET analysis pipeline. *Bioinformatics.* **31**, 3092-3098 (2015).

86. H. Wickham, *ggplot2: Elegant Graphics for Data Analysis* (Springer-Verlag, New York, 2009).

87. W. J. Kent *et al.*, The human genome browser at UCSC. *Genome Res.* **12**, 996-1006 (2002).

88. J. T. Robinson *et al.*, Integrative genomics viewer. *Nat. Biotechnol.* **29**, 24-26 (2011).

89. R. N. Ramirez, K. Bedirian, S. M. Gray, A. Diallo, DNA Rchitect: an R based visualizer for network analysis of chromatin interaction data. *Bioinformatics.* **36**, 644-646 (2020).

FIGURE LEGENDS

Figure 1. T-cell specific enhancer-promoter loops (EPLs) sync with the transcriptional regulation of the Treg gene program.

A. Profiling the genome-wide enhancer-promoter architectures of mouse regulatory T cells *in vivo* with H3K27Ac HiChIP. A focused view of the *Lrrc32* locus shows a null distribution based on random proximity-biased ligations (95% CI) and those called as significant H3K27Ac HiChIP enhancer-promoter loops (FDR 5%) in Tregs.

B. Scatter density plot of Treg H3K27Ac HiChIP EPLs and the relative cumulative intensities (Pearson r=0.78, $\log_{10}$).

C. Ranking plots of super-interactive promoters determined for H3K27Ac Treg (top, n=523 genes) and Tconv (bottom, n=520 genes) using the summed cumulative H3K27Ac EPL FDRs respectively.

D. T cell-specific enhancer-promoter configurations of the *Ikzf2* locus in Treg (top) and Tconv (bottom) cells. Differential EPLs are considered as >2 fold change and adj.P<0.05. IgG control HiChIP and null distributions (95% CI) are also shown for Tregs and Tconvs. ATAC-seq genomic tracks and RNA expression for *Ikzf2* are shown for Tregs and Tconvs respectively.

E. Fold change vs. mean intensity (MA) plot of differential H3K27Ac HiChIP EPLs between Treg (red) and Tconv (blue) cell-types. EPLs with differential intensity (adj.P<0.05, FC>2) are highlighted with numbers shown.

F. Comparison of 3D H3K27Ac HiChIP EPL intensities (x-axis, fold change in Treg vs. Tconv) and 1D H3K27Ac HiChIP EPL intensities (y-axis, fold change in Treg vs. Tconv).

G. Comparison of RNA expression (x-axis, fold change in Treg vs. Tconv) and cumulative H3K27Ac EPL intensities (y-axis, fold change in Treg vs. Tconv). Genes are colored as Up- (red) or Down- (blue) Treg signature membership.

Figure 2. Genome-wide Treg FoxP3 enhancer-promoter looping generally associates with H3K27Ac-marked architectures.

A. Treg FoxP3 (top) and Treg H3K27Ac (bottom) enhancer-promoter loops shown for the target gene *Entpd1.* Highlighted areas emphasize the agreement between FoxP3 and H3K27Ac EPLs.

B. Rank plot of the FoxP3-specific super-interactive promoters (n=320 genes).

C. FoxP3 (top), Treg H3K27Ac (middle), and Tconv H3K27Ac (bottom) enhancer-promoter loops for the *Lef1* locus.

D-F. Boxplots of FoxP3 cumulative EPL intensities for the Treg (D), FoxP3 transactivated (E) and Core signatures (F). Significance was determined for Treg Up ($p<0.05$), FoxP3 transactivated Up ($p<0.001$), and Core Treg ($p<0.01$) genesets using a t-test.

H. Scatter plot of FoxP3 HiChIP EPL intensities (x-axis, $\log_2$) and Treg H3K27Ac HiChIP EPL intensities (y-axis, $\log_2$). FoxP3-specific EPLs are highlighted.

I. Scatter plot of FoxP3 HiChIP EPL intensities (x-axis, $\log_2$) and Tconv H3K27Ac HiChIP EPL intensities (y-axis, $\log_2$). FoxP3-specific EPLs are highlighted.

J. Fold change distributions comparing H3K27Ac HiChIP EPL intensities (y-axis, Treg vs. Tconv) for each FoxP3 EPL bin (x-axis). Significance was determined using a t-test ($p<0.001$).

Figure 3. FoxP3 enhancer-promoter looping directly modulates FoxP3-dependent transcriptional program.

A. FACS plots for WT-GFP (top) and KO-GFP (bottom) sorted Treg populations (gated on the CD4+TCRB+ population).

B. WT-GFP and KO-GFP intensity and chromatin compartment H3K27Ac HiChIP maps.

C. Genome-wide scatter plot of H3K27Ac HiChIP genome compartment scores (Pearson 0.98).

D. WT-GFP (red) and KO-GFP (purple) H3K27Ac HiChIP intensity maps (left) and EPLs (right) of FoxP3 targets *Ikzf2* and *Lrrc32*. Arrows indicate EPLs FC>1.5 and P<0.05).

E. Comparison of 3D H3K27Ac HiChIP EPL intensities (x-axis, fold change in KO-GFP vs. WT-GFP) and 1D H3K27Ac HiChIP EPL intensities (y-axis, fold change in KO-GFP vs. WT-GFP). *Ikzf2* and *Lrrc32* specific H3K27Ac EPLs are indicated.

F. Scatter plot of WT-GFP and KO-GFP H3K27Ac HiChIP EPL intensities for FoxP3-specific EPLs.

G. Fold change distributions comparing H3K27Ac HiChIP EPL intensities (y-axis, KO-GFP vs. WT-GFP) for each FoxP3 EPL bin (x-axis). Significance was determined using a t-test (p<0.01).

H. Cumulative distribution frequency plot comparing KO-GFP and WT-GFP H3K27Ac EPL intensities for FoxP3 Transactivated (Up, p<0.001) and Core Treg (p<0.01) signatures. Significance was determined using a two-sided KS.test.

Figure 4. Parsing the FoxP3-dependent coregulated network with single-cell chromatin accessibility.

A. Schema of splenic Treg profiling with single-cell ATAC-seq. UMAP of splenic Tregs representing the FoxP3-dependent 'Up' signature gene score and gene-gene correlation heatmap (Pearson).

B. Gene scores were calculated from Treg scATAC-seq for the FoxP3-dependent signature and represented as a force-directed network. Gene-gene correlations (edges) and FoxP3 EPL cumulative intensities for each gene (node) in the network reflected by colored spectra. Indicated clusters were determined using both edge and node attributes.

C. Box plot comparing FoxP3 HiChIP cumulative EPL intensities for Cluster1 (orange) and Cluster2 (green) genes. Significance was determined between groups using a t-test (p<0.01).

D. FoxP3 and H3K27Ac HiChIP EPLs are shown for Cluster1 gene *Nrp1* and Cluster2 gene *Ccr8*.

E. Violin plots of FoxP3 ChIP-seq intensities between Cluster1 and Cluster2 genesets. ChIP-seq signal was measured in a 5kb window around the TSS.

F. Violin plots of Treg H3K27Ac cumulative EPL intensities between Cluster1 and Cluster2 genesets.

G. Volcano plots comparing the transcript expression of resting and activated Tregs (WT-GFP vs KO-GFP). Gene clusters from (B) are highlighted.

H. scATAC gene activity score correlations (Pearson) between FoxP3 and individual genes from Cluster1 and Cluster2 identified in (B).

Figure 5. FoxP3 positioning at enhancer-promoter loops distinguishes gene and TF-motif components

A. FoxP3 HiChIP EPL proportions based on FoxP3 binding modes fixed on either the promoter, enhancer or dual sides of FoxP3 loops.

B. FoxP3 ChIP-seq heatmaps centered on FoxP3 bound position within EPLs for promoter, enhancer and dual modes.

C. Gene ontology enrichments based on FoxP3 HiChIP EPL mode. Size of circle indicates number of genes and color represents significance (p-value).

D. Circos plot of TF motif enrichments for promoter and enhancer FoxP3 HiChIP EPLs. Arc thickness defines TF motif significance (-$\log_{10}$ p-value).

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

SUPPLEMENTAL FIGURE LEGENDS

Figure S1. Mapping the H3K27Ac enhancer-promoter architecture of conventional and regulatory T cells.

A. H3K27Ac HiChIP EPL biological replicate correlations for Treg (top, Pearson=0.85) and Tconv (bottom, Pearson=0.84).

B. Quality control analysis of H3K27Ac Treg (top) and Tconv (bottom) HiChIP using IgG HiChIP control. Density (left) and scatter (right) plots compare H3K27Ac and IgG HiChIP intensities among EPLs for each respective cell-type.

C. Distribution of H3K27Ac HiChIP EPL intensities ($\log_{10}$) and relative loop size (Kbp) in Tregs, with a median H3K27Ac EPL length of 49.5 kb.

D. Genome-wide comparative analysis of cumulative H3K27Ac HiChIP EPL intensities ($\log_{10}$) and mRNA expression ($\log_{10}$) for 9,058 genes (r=0.07, top). Distribution of cumulative H3K27Ac HiChIP EPL intensities ($\log_{10}$) for genes binned as quantiles based on low (10-20), middle (50-60), and high (90-100) expression in Tregs (bottom).

E. Vignettes of the T cell-specific enhancer-promoter structures for Treg signature genes *Il2rb* (Up-regulated) and *Pde3b* (Down-regulated).

F. Fold change vs. p-value (volcano) plot comparing H3K27Ac HiChIP EPLs in conventional and regulatory T cells. EPLs with differential intensity (adj.P<0.05, FC>2) are highlighted with numbers shown.

G. Fold change vs. cumulative H3K27Ac HiChIP EPL intensities, highlighting the Treg Up- and Down- gene signature (red and blue, respectively). Significance in the EPL intensity deviations were determined using a one-way Chi-square test in Up ($P<10^{-5}$) and Down ($P<10^{-5}$) gene subsets respectively.

Figure S2. Determining Treg FoxP3 enhancer-promoter loops using HiChIP.

A. Genome-wide chromosomal view of FoxP3 HiChIP biological replicates with a focused inset on chr1 and the correlations (Pearson) for FoxP3 HiChIP replicates and IgG control HiChIP (right).

B. Density (left) and scatter (right) plot comparisons of FoxP3 HiChIP and IgG intensities based on all unfiltered EPLs.

C. Density (left) and scatter (right) plot comparisons of FoxP3 HiChIP and IgG intensities for high-confidence EPLs following analysis using the proximity-biased null distribution.

D. Comparison of FoxP3 ChIP-seq datasets from (*41*) (x-axis, $\log_{10}$) and (*32*) (y-axis, $\log_{10}$). Lines (>3 ChIP-seq counts) emphasize the FoxP3 ChIP-seq peaks concordant in both datasets.

E. Scatter plot comparison of FoxP3 (x-axis) and IgG (y-axis) HiChIP intensities for high-confidence and FoxP3 ChIP-seq supported FoxP3 HiChIP EPLs.

F. Distribution of FoxP3 HiChIP EPL intensities ($\log_{10}$) and length of FoxP3 EPLs (Kb), with a median size 20 Kb.

G. Genomic view of H3K27Ac (Tconv and Treg), FoxP3 and Yy1 loops for the *Il2ra* locus. Highlighted regions mark positions of FoxP3-specific HiChIP loops. TF ChIP-seq and CAGE data below are profiled in Tregs.

H. Genomic view of H3K27Ac (Tconv and Treg) and Treg FoxP3 loops for the *Tcf7* locus.

I. Genomic view of H3K27Ac (Tconv and Treg) and FoxP3 and Yy1 loops for the *Il2* locus. TF ChIP-seq and CAGE data below are profiled in Tregs.

Figure S3. Generation and validation of FoxP3-deficient mice

A. Position and consequence of the mutation in FoxP3-deficient *Foxp3fs327-gfp/Doi* allele generated by CRISPR mutagenesis of the *Foxp3-gfp/Kuch* reporter allele (*54*). RBR: Runx-binding region; FKHD: Forkhead domain

B. Survival curve of $B6.Foxp3.fs327\text{-}gfp$ mice.

C. Images of WT and $B6.Foxp3.fs327\text{-}gfp$ mice.and their spleens

D. FoxP3 expression detected by intracellular staining and flow cytometry in CD4+ spleen Treg and Treg-like cells. Note the very slight shift in $Foxp3.fs327\text{-}gfp$ relative to unstained control.

E. Breeding strategy to obtain comparable Treg-like cells in which the $Foxp3.fs327$WT or *Foxp3-gfp/Kuch* alleles are active, in heterozygous females protected from inflammation by the normal *Foxp3-Thy1.1* reporter allele.

F. Reproducibility in H3K27Ac HiChIP EPL intensities between biological replicates for WT-GFP (top) and KO-GFP (bottom). The left panels show H3K27Ac EPLs for the *Ikzf2* locus and those for *Lef1* on the right.

G. WT-GFP (red) and KO-GFP (purple) H3K27Ac HiChIP intensity maps (left) and EPLs (right) of FoxP3 targets *Lef1* and *Add3*. Arrows indicate EPLs FC>1.5 and P<0.05).

Figure S4. Transcriptional specificities detailed by enhancer-promoter topologies of FoxP3 and Yy1 in regulatory T cells.

A. Treg Yy1 (top), Treg FoxP3 (middle), and Treg H3K27Ac (bottom) enhancer-promoter loops for the *Pim1* locus. Highlighted areas show differences of Yy1 and FoxP3 EPLs in Tregs.

B. Scatter plot comparing Yy1 (x-axis, $\log_{10}$) and FoxP3 (y-axis, $\log_{10}$) HiChIP EPL intensities (Pearson corr. 0.28).

C. Gene-level cumulative EPL intensities comparing Yy1 (x-axis, $\log_{10}$) and FoxP3 (y-axis, $\log_{10}$). Gene ontology enrichments are shown for genes enriched (>2 Fold) in FoxP3 HiChIP (top, orange) or Yy1 HiChIP (bottom, green) EPL intensities.

D. FoxP3-dependent signature genes (n=576) (*52*) shown for comparisons within resting (x-axis, Treg GFP-DTR-WT vs Foxp3-GFPKO) and activated Tregs (y-axis, Treg GFP-DTR-WT vs Foxp3-GFPKO). Genes colored red are upregulated, those in blue are downregulated FoxP3-dependent genes.

Figure S5. FoxP3 EPL modalities reveal differences in TF motif enrichments.

A. Heatmaps of FoxP3 HiChIP EPLs based on promoter, enhancer and dual modes. Treg ATAC-seq (top) and H3K27Ac ChIP-seq (bottom) signals are shown for each mode. Enhancer sides are positioned to the left and promoter to the right for each group respectively.

B. Ranking of TF motif enrichments (-$\log_{10}$ P-value) of promoter-side (top) and enhancer-side (bottom) FoxP3 EPLs.

Figure S1

Figure S2

Figure S3

A

FoxP3 coding region in *Foxp3^ires-gfpKuch*

337 431

ZF LZ RBR FKHD Ires GFP c

- A G insertion at the position coding for AA 327: **Asn > Lys**
- Frameshift mutation with a predicted Stop stop at AA 361 (exon 12)

Exon 11   Exon 12

WT EFFHNMDYFKYHNMRPPFTYATLIRW   AILEAPERQRTLNEIYHWFTRMFAYFRNHPATWK

327fs-gfp EFFHKHGLLQVPQYATPFHLCHPYPM   GHPGSPGEAEDTQ*

mRNA source: NM_054039.2

*: stop codon

B

WT n=10
327fs-gfp males n=10

Probability of survival
Age (days)

C

WT   *327fs-gfp*

25 day-old males

D

WT littermate (B6 iFOXP3-ires-GFP)
327fs-gfp
Isotype control

FoxP3   *Gated on TCRb^+CD4^+GFP^+*

E

Heterozygous *327fs-gfp* female   X   Male

Chr X [*327fs-gfp* B6]   Chr X: *Foxp3-Thy1.1*

Heterozygous *FoxP3-gfp* female   X   Male

Chr X [*Foxp3-gfp* B6]   Chr X: *Foxp3-Thy1.1*

Female pups   Female pups

KO FOXP3   WT FOXP3

Chr X [*327fs-gfp* / *Foxp3-Thy1.1*]   Chr X [*Foxp3-gfp* / *Foxp3-Thy1.1*]

GFP^+ Tregs (females)

7.0 2.8   2.8 5.2

Thy1.1
GFP

F

H3k27Ac HiChIP WT-GFP   R1   R2   *Ikzf2*   *Lef1*

H3k27Ac HiChIP KO-GFP   R1   R2   *Ikzf2*   *Lef1*

G

WT-GFP   KO-GFP   *Lef1*

WT-GFP   KO-GFP   *Add3*

Figure S4

A

Legend:
- Yy1 HiChIP
- FoxP3 HiChIP
- Treg H3K27Ac HiChIP
- Null distribution (95% CI)

FoxP3 ChIP-seq 0-8

HiChIP intensity

Treg ATAC-seq 0.1-0.6

200 kb   200 kb

Pim1

B

C

D

From van der Veeken *et al* GSE 154680

Figure S5

A

Promoter-side		Enhancer-side		Dual-sided	
FoxP3: −	+	+	−	+	+
Enhancer	Promoter	Enhancer	Promoter	Enhancer	Promoter

ATAC-seq

H3K27Ac ChIP-seq

B

SUPPLEMENTAL TABLE LEGENDS

TableS1 T cell EPLs. Tconv and Treg H3K27Ac EPLs, Treg FoxP3 EPLs and Yy1 EPLs are listed per gene for genomic position and intensity for each respective HiChIP experiment. NA's indicate absence of EPL for the respective experiment.

TableS2 Cumulative EPL intensity and T cell Super Interactive Promoters (SIP). EPL cumulative intensities for H3K27Ac, FoxP3 and Yy1 experiments. Significant H3K27Ac SIPs are listed for Tconvs and Tregs.

TableS3 FoxP3-dependent coregulated network. Gene and respective coordinate information for the genes selected in fig. S3D and displayed in Fig. 4. Cluster designation and FoxP3 EPL intensities are included. RNA expression values, fold change and p-values are included for Treg RNA-seq datasets from GSE154680.

TableS4 Gene ontology enrichments of FoxP3 EPLs. Top gene ontology enrichments and respective genes are listed for each FoxP3 EPL class.