1

Crowd-sourced observations of a polyphagous moth reveal evidence of allochronic

speciation varying along a latitudinal gradient

'

Emily N. Black[a], Jarrett D. Blair[a], Karin R.L. van der Burg[a], Katie E. Marshall[a*]

[a] Department of Zoology, University of British Columbia, Vancouver, British Columbia, Canada

*Corresponding author: Dr. Katie Marshall, kmarshall@zoology.ubc.ca

Keywords: allochronic speciation, citizen science, *Hyphantria cunea*, latitudinal biodiversity gradient, Lepidoptera, macroecology, temporal niche

Abstract

1. Allochronic speciation, where reproductive isolation between populations of a species is facilitated by a difference in reproductive timing, depends on abiotic factors such as seasonality and biotic factors such as diapause intensity. These factors are strongly influenced by latitudinal trends in climate, so we hypothesized that there is a relationship between latitude and divergence among populations separated by life history timing.

2. *Hyphantria cunea* (the fall webworm), a Lepidopteran defoliator with red and black colour morphs, is hypothesized to be undergoing an incipient allochronic speciation. However, given their broad geographic range, the strength of allochronic speciation may vary across latitude. We annotated >11,000 crowd-sourced observations of fall webworm to model geographic distribution, phenology, and differences in colour phenotype between morphs across North America.

3. We found that red and black morph life history timing differs across North America, and the phenology of morphs diverges more in warmer climates at lower latitudes. We also found some evidence that the colour phenotype of morphs also diverges at lower latitudes, suggesting reduced gene flow between colour morphs.

4. Our results demonstrate that seasonality in lower latitudes may increase the strength of allochronic speciation in insects, and that the strength of sympatric speciation can vary along a latitudinal gradient. This has implications for our understanding of broad-scale speciation events and trends in global biodiversity.

Introduction

Speciation occurs when barriers to interbreeding evolve between members of a species. This can occur in two contrasting ways: allopatric speciation, where new species arise due to geographic isolation between populations, or sympatric speciation, where new species arise without geographic isolation (Dieckmann and Doebeli, 1999). Sympatric speciation often has an ecological component leading to reproductive isolation such as differences in abiotic tolerance, host plant use, or other changes to the niche of populations (Sobel et al., 2010). However, the presence of one of these ecological shifts does not necessitate a speciation event, as the magnitude of the ecological differences can determine the extent of divergence (Crespi et al., 2004; Nosil et al., 2009). These ecological shifts are not static and often depend on abiotic and biotic factors that vary in space and time.

Sympatric speciation can also occur due to offsets in life history timing between populations, formally known as allochronic speciation ("allochronic" refers to the ecological separation of populations in time). These offsets may lead to the non-overlap of reproductive forms, or the formation of hybrids between populations that occupy a temporal space resulting in lower fitness (e.g., larvae that do not enter diapause in time for winter), which reinforces divergence (Dambroski and Feder, 2007; Yang et al., 2017). This type of speciation occurs in a variety of taxa, including insects (Dambroski and Feder, 2007; Godefroid et al., 2016; Hall and Willis, 2006). For example, *Rhagoletis pomonella* (the apple-maggot fly) experienced a host-plant shift with the introduction of apple trees to North America, which fruited earlier than the native hawthorn. This non-overlap of host plant phenology imposed a temporal reproductive barrier between *R. pomonella* host plant races, leading to allochronic speciation (Mattsson et al., 2015). Other studies indicate that a change in voltinism (or number of generations per year) in an insect

population can facilitate divergence, such as in the European corn borer (*Ostrinia nubilalis*)

(Coates et al., 2004; Wadsworth et al., 2020, 2013). Phytophagous insects have been suggested

as good candidates to study speciation and the factors contributing to its occurrence as the

exceptional diversity of this group has been attributed to a high incidence of sympatric speciation

(Berlocher and Feder, 2003; Mittelbach et al., 2007).

Though allochronic speciation occurs in temporally separated populations that exist together

geographically, the progress of this speciation can still be impacted by geographic or latitudinal

effects. One of the most general rules in biogeography is that biodiversity is higher in tropical

regions for most taxa, possibly due to greater niche specialization and higher rate of molecular

evolution in warmer climates (Gillooly et al., 2005; Hillebrand, 2004; Mittelbach et al., 2007).

As a result, Mittelbach et al. (2007) postulates that divergence among populations of a species

should also parallel the latitudinal gradient of biodiversity. Allochronic speciation may similarly

be more effective at facilitating population divergence in lower latitudes, as warmer climates

lead to greater temporal niche space. Temporal niche space is defined as the length of time in

which abiotic and biotic conditions are favourable for growth and reproduction, which is closely

tied to seasonality (or cyclical changes in climate within a year; Bradshaw, 1974). Competition

within the temporal niche facilitates diversification of life history patterns; however, the degree

of diversification is limited by the size of the niche and the rate of evolution. When the

phenological niche is larger (such as at lower latitudes), less overlap between the phenology of

competing populations is expected, and vice versa (Post, 2019). Given that seasonality at lower

latitudes facilitates longer growth and reproductive seasons, this suggests there is greater

temporal niche space available for insects in these regions, allowing for less overlap in

phenology between competing populations. This may facilitate greater diversification among

populations undergoing allochronic speciation, and increase the strength of speciation at lower

latitudes.

By contrast, there is evidence that speciation rates may increase at higher latitudes in particular

situations. In a review of 309 bird and mammal sister species, Weir & Schluter (2007) found a

significant increase in speciation and extinction rates towards the poles, and North American

insect species richness may increase at middle latitudes (Weiser et al., 2022). Differences in

diapause initiation and termination are strong drivers of allochronic speciation in insects, and

diapause timing and physiology is expected to be more important in temperate regions

experiencing more extreme winter seasonality (Dambroski and Feder, 2007; Inskeep et al.,

2021). This may increase the strength of allochronic speciation at higher latitudes, especially in

insects who undergo facultative diapause and have greater plasticity in life history timing (Chen

et al., 2014). This leads to two contradictory possibilities: allochronic speciation may be more

effective at lower latitudes because of greater temporal niche availability, or it may be more

effective at higher latitudes due to the greater importance of climate asynchrony and diapause

intensity. Here, we use the fall webworm (*Hyphantria cunea*, Lepidoptera: Erebidae*)* to

investigate these contrasting hypotheses, and we explore how allochronic speciation proceeds

along a latitudinal gradient.

The fall webworm is a widespread generalist moth defoliator native to North America, whose

gregarious larvae live in conspicuous webs on the branches of bushes and trees (Loewy, 2013;

Schowalter and Ring, 2017). The fall webworm appears in discrete generations, ranging from

one generation per year (univoltine) in northern regions to up to four or five generations per year

(multivoltine) in southern regions (Yang et al., 2017). Fall webworm larvae occur in two distinct

colour morphs: a red morph and a black morph which exist sympatrically (Jaenike and Selander,

1980). The red morph has a distinct red head capsule with a brown, tan, or black body, and the

black morph has a black head capsule with a grey or black body. An intermediate maroon morph

exists but is only found in northwestern North America. Little is known about this colour

morph's phylogenetic relatedness to the red and black morphs, but it is likely geographically

separated from the other morphs by the Rocky Mountains (Yang et al., 2017).

The red and black morphs have both physiological and behavioural differences (Oliver, 1964),

and genetic analyses show clear separation of the two groups, suggesting reproductive isolation

with little gene flow between them despite significant range overlap (Jaenike and Selander, 1980;

Loewy, 2013; Vidal et al., 2019; Yang et al., 2017). However, the two morphs are sexually

compatible and produce viable F1 and F2 offspring in the lab (Hattori and Ito, 1973; Morris,

1963). These results suggest that the two fall webworm colour morphs are currently undergoing

a sympatric speciation event. The black and red morph moth flights do not occur together, their

voltinism differs, and the critical photoperiod for pupal diapause is longer in the black morph

form, indicating that allochronic speciation may be responsible for the morph's current

reproductive isolation (Takeda, 2005; Yang et al., 2017). Since the fall webworm is extremely

polyphagous (Loewy, 2013) and the colour morphs share many of the same host plants (Oliver,

1964), changes in host plant preference are an unlikely cause for this speciation event. Yang et

al. (2017) suggest allochronic speciation of the fall webworm is driven by a difference in

diapause intensity, development rate, and voltinism between morphs, which evolved to reduce

intraspecific competition. However, they also find significant variation in voltinism and diapause

timing in both morphs across their latitudinal range (Yang et al., 2017). Given there are limited

genetic analyses investigating the degree of difference between red and black fall webworm

morphs across latitude (Vidal et al., 2019; Yang et al., 2017), differences in phenology between

morphs can serve as an indicator of the strength of allochronic speciation acting on fall

webworm across their range.

Given the fall webworm's broad distribution, crowd-sourced (i.e., citizen) science websites such

as iNaturalist offer a useful alternative to field studies when investigating life history trends

across a large geographic range. These websites (e.g., iNaturalist, eBird, eButterfly) are

applications where users upload records or photographs of organisms they observe, which can

then be annotated, sorted, and downloaded by researchers. Not only do crowd-sourced science

sites reduce the cost and time required for field studies, but they produce large and high-quality

datasets which engage the public in academic research (Jaskuła et al., 2021). iNaturalist has been

used to study differences in phenology (Barve et al., 2020) and variation in colour phenotype on

a large geographic scale (Lehtinen et al., 2020; Tseng et al., 2022), and can be used to investigate

the same phenomena in the fall webworm: a conspicuous caterpillar with thousands of

iNaturalist records across its range. Although crowd-sourced records are not as rigorous as field

samples, their geographic, temporal, and taxonomic scope make them suitable resources for

macroecological studies.

Here we use crowd-sourced iNaturalist records to investigate whether the strength of allochronic

speciation varies across a latitudinal gradient using the fall webworm (*Hyphantria cunea*) as a

model species. The fall webworm is found over a broad latitudinal range, and is conspicuous due

to its large nests, so it is frequently photographed by community scientists. The fall webworm is

also currently undergoing allochronic speciation (Yang et al., 2017) with a limited effect of host

plant specialization as it is extremely polyphagous (Vidal et al., 2019; Warren and Tadic, 1970).

We tested two competing hypotheses: either that allochronic speciation facilitates greater

differences between groups at higher latitudes due to greater influence of diapause intensity and

climate asynchrony, or allochronic speciation facilitates greater differences between groups at

lower latitudes where there is greater temporal niche space. Due to an absence of genetic data,

we used phenological overlap and phenotype measures to evaluate the strength of speciation

across the fall webworm range. We find that there is less phenological overlap between red and

black morph larvae at lower latitudes and in warmer climates. We also find that the colour

phenotype of red and black morph larvae diverges more at lower latitudes, supporting a decrease

in gene flow facilitated by allochronic speciation.

Methods

Data collection and standardization

We decided to focus on the fall webworm larval life stage because colour morph is only reliably

discernable from larval head capsule, not adult colouration (Yang et al., 2017). All data for

geographic, phenology, and phenotype measures was sourced from images uploaded to

iNaturalist (http://inaturalist.org/), a repository of community-collected observations. We found

24,991 observations of *H. cunea* across all years and locations, of which 19,333 were "research

grade" (having two matching independent identifications by users) (Barve et al., 2020). Using

research grade observations helps prevent the inclusion of species commonly misidentified as

fall webworm (e.g., *Malacosoma americanum*, *Estigmene acrea*) (Organic Garden Info, n.d.;

University of New Hampshire, n.d.).

We examined all research grade images of *H. cunea* uploaded from 2018 – 2020 because

previous years had too little data to be suitable for analysis, and 2021 records were not

completed at the time of analysis. We reviewed each photo manually, blinded to observation

location, to identify the larval colour morph and whether the larval head capsule was visible (as

head capsule is the most reliable indicator of colour morph). In total, we reviewed 11,674

iNaturalist observations. We used iNaturalist's built-in feature "Observation Fields" to store

information with the photograph being classified such as whether the head capsule was visible

("Head capsule visible?") and colour morph ("Colour"). The intermediate maroon morph was

classified as red in the "Colour" field (Yang et al., 2017). No colour morph can be reliably

discerned from photos of nests or moths, so we did not include these photographs in

categorization or analysis.

All analyses were performed using R version 4.2.2 (R Core Team, 2021). The dataset, all

associated code, and full-sized figures are published on the Open Science Framework

(https://osf.io/x64gp/). Raw data from iNaturalist includes observations from North America

(which excludes invasive ranges in Asia and Europe; Wu et al., 2019; Yang et al., 2017), and

contains the fields: photo ID, user ID, date and time of observation, observation URL, longitude

and latitude, state, county, country, scientific name, and the custom Observation Fields "Head

Capsule Visible?" and "Colour".

Each fall webworm observation was also assigned to their corresponding plant hardiness zone.

Plant hardiness zones are geographic regions binned by their average annual minimum

temperature and assigned a reference number (1-13, with lower numbers indicating colder

climates) which act as an indicator of overall climatic conditions (Natural Resources Canada,

2012). Because phenology is affected by climate (Hodgson et al., 2011), plant hardiness zones

serve as a method to bin fall webworm into groups that should experience similar seasonality,

and thus, similar life history timing. Plant hardiness zones also scale roughly with latitude, so

they can act as an indicator for trends in seasonality and phenology across latitudes. A plant

hardiness zone map for the United States was obtained from the United States Department of

Agriculture (USDA Agricultural Research Services, n.d.). To generate a map of plant hardiness

zones for Canada using the same categorization as the USA, BioSIM climate generation software

was used to create a map of the average annual minimum temperature across Canada (Régnière

et al., 2014). This map was then binned into plant hardiness zones based on USA categories.

Using the *tidyverse* package (Wickham et al., 2019), fall webworm observations were filtered to

those with an assigned colour morph and visible head capsule for all analyses. Only larval

populations from the eastern United States and Canada (-100 to $0°W$ longitude) were used in

analyses so geographically isolated populations like those in Colorado (Vidal et al., 2019) and

northwestern North America were excluded. After filtering, 5,176 observations remained and

were used for analysis.

Geography

Maps of North American fall webworm distribution from 2018-2020 were generated using the

Tmap package (Figure 1; Tennekes, 2018). To determine if the colour morphs were

geographically separated from one another in their eastern range, we compared the latitudinal,

longitudinal, and elevational (calculated using the *Elevatr* package; Hollister, 2021) distribution

of each colour morph in each year (Figure 2). Differing abiotic tolerance may cause the morphs

to have different geographic distributions; however, we still expect them to be found together

across their range, consistent with a sympatric form of speciation. Two-way ANOVA was used

to determine if colour morph and year recorded had a significant effect on latitude, longitude,

and/or elevation, and Tukey post-hoc tests were used to compare among groups. Not all

observations could be assigned a valid elevation, so sample size was reduced to 4,652

observations when comparing elevation of colour morphs.

Phenology

Because phenology is climate-dependent (Hodgson et al., 2011), we binned fall webworm

observations using plant hardiness zones (Natural Resources Canada, n.d.). We also grouped

observations by year, as phenology may change even in the same plant hardiness zone as weather

varies from year to year (Scranton and Amarasekare, 2017; Tobin et al., 2008). Morph

abundance was calculated by week rather than by day to minimize daily variation in sampling

effort by community scientists (Roy et al., 2016). Given the red morph has a more limited

latitudinal distribution than the black morph (Figure 2), we did not include plant hardiness zones

containing less than 10 observations of the red morph in any given year in the analysis. This

reduced our dataset to 4,866 observations in plant hardiness zones 5-9 for phenology measures.

Although the fall webworm appears in discrete generations each summer (Yang et al., 2017),

these are difficult to parse and compare using crowd-sourced data. So, we investigated

differences in phenology using two separate methods. First, we compared the earliest and latest

10% of observation dates for each colour morph. Using the earliest and latest 10% of observation

dates ensures comparisons between morphs are reflective of overall larval population timing

rather than individual outliers. If divergence between morphs increases with increasing temporal

niche space, we expect differences between the earliest and/or latest observation dates of the

morphs will increase in larger plant hardiness zones. Alternatively, if divergence between the

morphs increases at higher latitudes, we expect differences between observation dates of the

morphs will decrease in larger plant hardiness zones. Therefore, we selected the earliest and

latest 10% of observation dates for the black and red morph in each plant hardiness zone and

year, and calculated the overall mean difference between red and black morph timing and the

standard error of the mean difference. We then used a 3-way ANOVA and general linear model

to test whether the black and red morph showed a difference in timing of their earliest and latest

observances, and whether this was driven by year and plant hardiness zone.

Secondly, we used the "DTS" statistic from the *twosamples* package to quantify the degree of

phenological overlap between colour morphs each year (Dowd, 2022, 2020). The DTS statistic

calculates the difference between the distributions of two groups using the weighted area

between their cumulative distribution functions (aka CDFs; Figure S2). Here we used it to

quantify the degree of phenological difference between red and black morph larvae using the

cumulative distribution function of their observations in each year. For each year and plant

hardiness zone, we calculated the DTS statistic comparing the red and black morphs. If

divergence between morphs increases with increasing temporal niche space, we would expect to

see the DTS statistic increase with plant hardiness zone. Alternatively, if divergence increases at

higher latitudes, we expect to see the DTS statistic decreasing with increasing plant hardiness

zone. Therefore, we used a general linear model to test whether plant hardiness zone, year, or

their interaction significantly predicted DTS statistic. We compared models with both

polynomial and linear predictors as initial plotting indicated the relationship between DTS

statistic and plant hardiness zone may be curvilinear (Fig. 4) and selected best fit models by AIC

score (Mazerolle, 2020). Since the fall webworm reproductive form (moths) does not co-occur

with larvae, differences in phenological overlap of fall webworm larval generations is a predictor

of differences in moth generations (Loewy, 2013).

Colour phenotype

We examined the colour data from iNaturalist images to determine if there is phenotypic

divergence between the black and red colour morphs across plant hardiness zones. From a

standardized dataset containing 4,594 images, those that were low resolution, had poor exposure,

were taken from a far distance, or had an object obstructing the webworm were excluded (Figure

S3D), leaving 3,789 images. The colour data for the entire fall webworm (including head capsule

and body) was extracted using Adobe Photoshop 2021. Individual larvae were selected from

each image using Photoshop's object detection tool (Figure S3A). If Photoshop's object

detection tool was unable to properly select the webworm, the freehand lasso tool was used

instead (Figure S3B, C). These colour records were further filtered to those from 2018-2020, east

of -100°W longitude, and in plant hardiness zones 5-9, leaving 3,475 photos for analysis.

To test whether colour data could predict colour morph, we measured the mean, standard

deviation, minimum value, maximum value, kurtosis, and skewness of each RGB histogram

using image attributes. Colour histograms of images were generated in Photoshop and histogram

data was extracted from webworm selections as a .csv file using Photoshop's "Record

Measurement" feature (Figure S4). Using R, we calculated photograph properties manually from

exported histogram data. These variables were then normalised (i.e., centered and scaled; Kuhn,

2021) in R in preparation for linear discriminant analysis (LDA; Ripley and Venables, 2002).

LDA was performed to find a variable that best separated the red and black colour morphs based

on the RGB histogram data (Table S1). This variable was named "LD1". To determine if the

colour (LD1) of each colour morph depends on plant hardiness zone and year, we performed an

ANCOVA. We also performed two simple linear regressions to determine the effect of plant

hardiness zone and year on the colour of each colour morph individually. If divergence between

morphs increases with increasing temporal niche space, we expect the colour phenology of the

morphs to diverge in larger plant hardiness zones. If divergence increases at higher latitudes, we

expect colour phenology of the morphs to diverge in smaller plant hardiness zones.

Results

Differences between geographic distribution of morphs

We found that red and black morph larvae occurred together across much of their Eastern range

(Figure 1; Figure S1), consistent with a sympatric mode of speciation. Larvae in Colorado and

northwestern North America were completely geographically isolated from Eastern populations

(Figure 1).

While the two morphs are generally found sympatrically, differential abiotic tolerance may

change colour morph distribution within their eastern range (east of -100°W longitude). Black

morph populations were found an average of 2.3 ± 1.4 (SD) degrees further north than the red

morph populations ($F_{1,5170}=93.00$, $P<0.0001$; Figure 2). However, differences in latitudinal range

differed by year ($F_{4,5170}=39.12$, $P<0.0001$), as the average ranges of the morphs were not

significantly different in 2020 (Tukey Post-Hoc, $P=0.06$). We also found that red morph

populations occurred an average of 4.1 ± 1.5 (SD) degrees further west than the black morphs

($F_{1,5170}=104.30$, $P<0.0001$; Figure 2). However, the difference between the longitudinal ranges of

morphs also varied among years ($F_{4,5170}=11.87$, $P<0.0001$), as although the average longitudinal

distributions of morphs in 2020 were significantly different, the magnitude of the difference in

longitude was smaller than in previous years (Tukey HSD, $P<0.0001$). There was no difference

in the elevation of the red and black colour morphs across all years ($F_{1,4644}$=0.06, P=0.8; Figure

2).

Differences between phenology of morphs across plant hardiness zones

To find evidence for a difference in phenology between morphs across latitude, we tested for

differences in each plant hardiness zone and year between the earliest 10% of observation dates

of each morph, and the latest 10% of observation dates of each morph. Previous studies show the

black morph appears earlier than the red and remains later in the fall (Yang et al., 2017). Overall,

we found the earliest 10% of black morph observations occurred an average of 22 ± 3.35

(standard error of mean difference) days earlier than the earliest 10% of red morph observations

($F_{1,521}$=156.7, P<0.0001; Figure 4B). We found differences between the earliest 10% of morph

dates differed by plant hardiness zone. Black morph observations tended to appear earlier across

increasing plant hardiness zone (general linear model, β =-23 days/zone) than the red morph

(general linear model, β =-18 days/zone), resulting in greater differences between morphs in

warmer plant hardiness zones at lower latitudes ($F_{1,521}$=876.8, P<0.0001; Figure 4A).

Differences between earliest observance dates of the morphs across plant hardiness zones were

not driven by year ($F_{2,521}$=1.81, P=0.1; Figure 4A). We found the latest 10% of black morph

larvae occurred an average of 3 ± 2.27 (standard error of the mean difference) days later than the

red morph; however, this difference was not statistically significant ($F_{1,517}$=3.77, P=0.05; Figure

4A). Differences between the 10% of latest dates in the black and red morph were driven by

plant hardiness zone. Black morph observations tended to appear later across increasing plant

hardiness zones (general linear model, β = 17 days/zone) than the red morph (general linear

model, β =10 days/zone; $F_{2,517}$=410.2, P<0.0001; Figure 4B); however, since the latest observed

dates of the morphs converged in plant hardiness zone 7, there is no trend in the difference

between morphs across latitudes. Differences between morph latest observation dates across

plant hardiness zones did not differ by year ($F_{2,517}$=0.74, P=0.6; Figure 4B).

We also tested for differences between phenology of morphs by comparing their temporal

abundance patterns in each year. Using the DTS statistic, we quantified the difference between

the cumulative distribution functions of the morphs across each plant hardiness zone and year

(see Supplemental Figure 2 for cumulative distribution functions of morphs). The relationship

between differences in morph temporal abundance and plant hardiness zone was best fit by a

linear regression according to AIC (Mazerolle, 2020). The regression showed differences in the

temporal abundance of the morphs increased with increasing plant hardiness zones ($F_{1,9}$=25.11,

P<0.001; Figure 5). Trends in temporal overlap across plant hardiness zones did not differ by

year ($F_{2,9}$ =0.15, P=0.9; Figure 5). DTS analysis was also run on all observations with and

without head capsule visible, and no qualitative differences in results were observed.

Differences in colour morphology of morphs across plant hardiness zones

If there are differences in the effectiveness of allochronic speciation across plant hardiness zones,

we would expect to see differences in colour morphology across plant hardiness zones. Of

photos used in colour analysis, 3082 were of black morph webworms while 393 were of red

morph webworms. To test the relationship between colour and latitude, we used an LDA model

to create a single summary variable (LD1) that best separated the red and black morphs. Among

the coefficients of linear discriminants, the mean green value, mean red value, and red standard

deviation had the highest absolute values (-2.60, 1.75, and -0.97 respectively) (Table S1).

We found that the colour morphs were sorted well by linear discriminant analysis of their

colouration ($F_{1,3467}$=925.97, P<0.0001; Figure 6). We also found that trends in colouration across

plant hardiness zones differed between red and black morphs ($F_{1,3467}$=7.37, P<0.01). When

modelling only red morph colouration, the red larvae show a slight increase in LD1 values as

plant hardiness zone increases (linear regression, β=0.1, R^2=0.004; Figure 6); however, this was

not significant ($F_{1,387}$=0.39, P=0.5). Year of observation did not affect trends in red morph

colouration across plant hardiness zones ($F_{2,387}$=1.20, P=0.3). However, LD1 values decrease

slightly with increasing plant hardiness zone when modelling only black morph larvae (linear

regression, β=-0.09, R^2=0.016; $F_{1,3078}$= 42.57, P<0.0001; Figure 6). Trends in black morph

colouration across plant hardiness zones were also driven by year, with 2019 showing greater

decline in LD1 values than other years ($F_{2,3078}$=3.7, P<0.05).

Discussion

Here we show that the black morph appears earlier than the red morph in plant hardiness zones

which occur at lower latitudes (Figure 4A). We also find that differences between the phenology

of the morphs across their entire reproductive season are greater in plant hardiness zones at lower

latitudes where the length of the reproductive season is longer (Figure 5). The colouration of the

black morph also diverges from the red morph in plant hardiness zones at lower latitudes (Figure

6). These results suggest that greater differences in phenology at lower latitudes where temporal

niche space is larger may be facilitating decreased gene flow between morphs, and that the

strength of allochronic speciation is varying along a latitudinal gradient.

While the colour morphs of reproductive adult fall webworms cannot be reliably differentiated

(Yang et al., 2017), differences between the emergence dates of red and black morph fall

webworm larvae suggests there is greater temporal separation of the morphs at lower latitudes.

We found that differences between the earliest observation dates of colour morph larvae were

larger in warmer plant hardiness zones at lower latitudes (Figure 4A). There was no difference in

the latest observation dates of the morphs, nor was there a consistent trend in differences across

plant hardiness zones (Figure 4B). Since the fall webworm overwinters as a pupa and emerges as

a reproductive adult (Li et al., 2001), differences between the emergence timing of colour morph

moths may decrease mating between morphs in the first generation (Yang et al., 2017). As well,

differences in emergence date between morphs were consistent with differences in diapause

intensity found in previous studies of fall webworm (Yang et al., 2017). Differences between red

and black morph diapause intensity may lead to hybrids between morphs being poorly

temporally adapted to their environment, reinforcing genetic separation (Ording et al., 2009;

Yang et al., 2017). These results support our hypothesis that greater temporal niche space at

lower latitudes, not winter seasonality and diapause effects at higher latitudes, drives a latitudinal

gradient in allochronic speciation. Larger temporal niche space at lower latitudes allows for

greater temporal niche partitioning between populations, including changes in seasonal onset and

decline. This phenomenon may be facilitating greater differences in emergence date of the colour

morphs, and changing the strength of allochronic speciation across latitudes (Post, 2019).

The phenology of fall webworm larval colour morphs diverged more in warmer climates,

suggesting allochronic speciation driven by a difference in phenology operates more effectively

at lower latitudes. Evaluating the yearly phenology of the colour morphs was important to

determine whether temporal isolation between morphs is maintained across the entire

reproductive season (Yang et al., 2017). We found that temporal patterns of abundance of the

colour morphs were less similar in climates at lower latitudes (Figure 5). Qualitatively, we also

observed fall webworm voltinism was more similar at higher latitudes (Figure 3); however, we

could not reliably measure the number of fall webworm generations using this data. These

patterns may be driven by trends in temporal niche space, as larger temporal niche space in

warmer climates can facilitate greater niche partitioning (Post, 2019). The importance of

generational overlap on the progress of allochronic speciation across a broad range has been

demonstrated in other insects (Coates et al., 2004; Ording et al., 2009). For example, populations

of the geometrid winter moth (*Inurois sp.*) undergoing allochronic speciation showed increased

genetic differentiation when population phenology overlapped less, which was facilitated by a

difference in seasonality (Yamamoto and Sota, 2009). While our phenological results support a

latitudinal gradient in the strength of allochronic speciation driven by temporal niche space,

genetic studies on the relatedness of the morphs across a latitudinal range are needed to lend

further support to this hypothesis.

Since colour morph is only reliably discernable from larval observations (Yang et al., 2017), we

could not model the phenology of the reproductive form (the moth). Using larval phenology to

predict allochronic speciation assumes that both the red and black morph moths appear in the

same fixed interval after their larval occurrence. Timing of the life cycle stages of the morphs

may have diverged, causing moth phenology to disagree with the larval phenology presented

here. Figure 3 is consistent with phenology modelling of moths presented by Yang et al. (2017),

except at the highest latitude regions where they find univoltine black and red morph moths do

not have overlapping generations, suggesting greater differences in phenology than observed

here (Figure 5). The discrepancy between our results and those of Yang et al. (2017) at the

highest latitudes should be explored, as our interpretation relies on larval occurrence being a

reliable indicator of moth phenology.

Crowd-sourced data from iNaturalist lent itself well to our broad-scale analysis of fall webworm

phenology and phenotype, as it allowed us to investigate macroecological trends across a large

geographic and temporal scale (Barve et al., 2020; Bonney et al., 2009). While the lack of

consistency in sampling methods of iNaturalist images may have caused increased variation in

the data, the volume and spread of data available allowed us to explore questions which cannot

be answered using field studies alone. Crowd-sourced photographs may be biased towards days

and seasons with favourable weather conditions and urban environments with high observer

density, and poor image quality may make colour morph identification more difficult (Roy et al.,

2016). To address this, images were manually reviewed for species and colour morph, only

images in which the head capsule was clearly visible were used, and we calculated larval

abundance and phenology by summarizing observations weekly. As well, biases towards

observations in urban areas should not affect our results since our comparisons between

sympatric morphs do not depend on observer density. However, the accuracy of our phenology

and phenotype measures may still have been affected, and investigations using a single image or

small subset of observations from crowd-sourced datasets such as the one used here may be

unreliable. In future studies using crowd-sourced science, consideration should be given to how

to minimize confounding factors due to uneven sampling, such as using crowd-sourced science

data in tandem with field surveys (Lehtinen et al., 2020). However, the accessibility of

iNaturalist data, ease of collection, volume of data available, and usefulness of crowd-sourced

data for broad-scale research makes it a promising avenue for data collection, and a useful

resource for investigating macroecological trends across latitudes.

Phenotypic results showed colouration of red and black morph caterpillars diverged in warmer

climates. The colour phenotype of the black morph appears to shift away from the colour

phenotype of the red morph at warmer plant hardiness zones. Interestingly, this shift in colour

only appears in the black morph, as red morph webworms do not experience any significant

colour shift across their entire range (Figure 6). If the colour shift were a result of increasing

melanism in higher latitude populations, as observed in other ectotherms (Clusella et al., 2007),

we would expect to observe this shift in both colour morphs. However, changes in colouration

across the fall webworm range were small, and qualitatively we observed variation in colour

phenotype even among individuals of the same region. So, these results lend some support to our

hypothesis that the strength of allochronic speciation is higher at lower latitudes, but further

investigation using field and genetic studies is needed.

While a genetic comparison of colour morphs across latitudes could determine the degree

speciation between morphs and support phenotypic results presented here, a genetic dataset on a

geographic scale large enough to address this question does not exist (Yang et al., 2017). We

attempted to quantify the genetic relatedness of the morphs across their range, but available data

was not sufficient to answer this question (unpublished data). Morphological and phenotypic

comparisons such as the ones performed here can offer valuable information on the relatedness

of populations undergoing speciation prior to investigating population genetic structure. For

example, the bean leaf beetle (*Cerotoma trifurcate*) has four distinct colour phenotypes with a

previously unknown evolutionary relationship. Morphological analyses of the colour phenotypes

were used to generate an initial hypothesis regarding their phylogenetic separation, and genetic

work confirmed weak reproductive separation between phenotypes (Tiroesele et al., 2018).

Trends in the colour morphology and phenology of the fall webworm across their latitudinal

range inform hypotheses on the strength of allochronic speciation on a latitudinal gradient;

however, we encourage further investigation into population genetics of the fall webworm to

confirm the results presented here.

Conclusion

In this study we tested two competing hypotheses: the first that allochronic speciation facilitates

divergence between populations more effectively at higher latitudes, or the second that

allochronic speciation facilitates divergence between populations more effectively at lower

latitudes, using the fall webworm (*Hyphantria cunea*) as a model species. Our analyses of

phenology and phenotypic differences are consistent with the current allochronic speciation in

the fall webworm operating more effectively at lower latitudes where there is greater temporal

niche space, rather than at higher latitudes due to the effects of winter seasonality and differences

in diapause. Further study is needed as some factors such as limited access to genetic data,

variable sampling methods in crowd-sourced science data, and using only larval phenology

measures may have confounded our results. Advances in macroecology and crowd-sourced data

provide an avenue for further study of ecological speciation over large geographic areas.

Furthermore, investigating the interplay of latitudinal effects and ecological speciation can

change our understanding of how speciation operates under different conditions and help to

understand past, present, and future trends in biodiversity. Since the great diversity of

phytophagous insects has been attributed to high rates of sympatric speciation, greater

exploration of allochronic speciation and other forms of ecological speciation in widespread

species could elucidate this evolutionary history (Berlocher and Feder, 2003; Mittelbach et al.,

2007).

Acknowledgements

E.N.B is supported by an NSERC CGS-M fellowship. J.D.B. is supported by an NSRC PGS-D

fellowship. K.v.d.B. is supported by NSF IOS 1930829. K.E.M is supported by an NSERC

Discovery Grant. We humbly thank J. Stireman at the University of Denver for his preliminary

review of our results, and Dr. S. Otto and three anonymous reviewers for helpful comments that

significantly improved the manuscript. We also thank each community member who submitted

their photo to iNaturalist to aid in our data collection. We are appreciative of our friends and

family who supported us throughout this project.

Data Accessibility

All code, raw data, and full-sized figures are available from the OSF Repository:

https://osf.io/x64gp/

Competing Interest

The authors have no competing interests to report.

Author's Contribution (CRediT Author Statement)

Emily Black: Conceptualization, Data Collection, Methodology, Investigation, Formal Analysis,

Visualization, Writing – Original Draft, Writing- Review and Editing, Project Administration.

Jarrett Blair: Methodology, Investigation, Formal Analysis, Visualization, Writing- Original

Draft, Writing – Review and Editing. **Karin van der Burg:** Data Collection (unpublished),

Formal Analysis (unpublished), Writing – Original Draft, Writing – Review and Editing, **Katie**

506   **Marshall:** Conceptualization, Methodology, Investigation, Writing – Original Draft, Writing –

507   Review and Editing, Supervision, Project Administration, Project Funding.

References

Barve V v, Brenskelle L, Li D, Stucky BJ, Barve N v, Hantak MM, et al. Methods for broad-scale plant phenology assessments using citizen scientists' photographs. Appl Plant Sci 2020;8. https://doi.org/10.1002/aps3.11315.

Berlocher SH, Feder JL. Sympatric speciation in phytophagous insects: Moving beyond controversy? Annu Rev Entomol 2003;47:773–815. https://doi.org/10.1146/ANNUREV.ENTO.47.091201.145312.

Bonney R, Cooper CB, Dickinson J, Kelling S, Phillips T, Rosenberg K v., et al. Citizen Science: A Developing Tool for Expanding Science Knowledge and Scientific Literacy. Bioscience 2009;59:977–84. https://doi.org/10.1525/BIO.2009.59.11.9.

Bradshaw WE. Phenology and Seasonal Modeling in Insects. In: Canadell JG, Díaz S, Heldmaier G, Jackson RB, Levia DF, Schulze E-D, et al., editors. Ecological Studies, vol. 8, Berlin: Springer; 1974, p. 127–37. https://doi.org/10.1007/978-3-642-51863-8_11.

Chen C, Wei XT, Xiao HJ, He HM, Xia QW, Xue F sen. Diapause induction and termination in Hyphantria cunea (Drury) (Lepidoptera: Arctiinae). PLoS One 2014;9. https://doi.org/10.1371/JOURNAL.PONE.0098145.

Clusella ST, van Wyk JH, Spotila JR. Thermal melanism in ectotherms. J Therm Biol 2007;32:235–45. https://doi.org/10.1016/J.JTHERBIO.2007.01.013.

Coates BS, Sumerford D v., Hellmich RL. Geographic and voltinism differentiation among North American Ostrinia nubilalis (European corn borer) mitochondrial cytochrome c oxidase haplotypes. Journal of Insect Science 2004;4. https://doi.org/10.1093/JIS/4.1.35.

Crespi BJ, Morris DC, Mound LA. Evolution of ecological and behavioural diversity: Australian Acacia thrips as model organisms. Canberra: Australian Biological Resources Study; 2004.

Dambroski HR, Feder JL. Host plant and latitude-related diapause variation in Rhagoletis pomonella: a test for multifaceted life history adaptation on different stages of diapause development. J Evol Biol 2007;20:2101–12. https://doi.org/10.1111/J.1420-9101.2007.01435.X.

Dieckmann U, Doebeli M. On the origin of species by sympatric speciation. Nature 1999;400:354–7. https://doi.org/10.1038/22521.

Dowd C. Package "twosamples" Type Package Title Fast Permutation Based Two Sample Tests 2022.

Dowd C. A New ECDF Two-Sample Test Statistic. ArXiv 2020. https://doi.org/10.48550/arxiv.2007.01360.

Extreme Minimum Temperature Models. Natural Resources Canada n.d. http://planthardiness.gc.ca/?m=17&lang=en (accessed November 20, 2022).

Fall Webworm. Organic Garden Info n.d. http://www.organicgardeninfo.com/fall-webworm.html (accessed October 30, 2022).

Fall webworm & Eastern Tent Caterpillar [fact sheet]. University of New Hampshire n.d. https://extension.unh.edu/resource/fall-webworm-eastern-tent-caterpillar-fact-sheet (accessed October 30, 2022).

Gillooly JF, Allen AP, West GB, Brown JH. The rate of DNA evolution: Effects of body size and temperature on the molecular clock. Proc Natl Acad Sci U S A 2005;102:140–5. https://doi.org/10.1073/pnas.0407735101.

Godefroid M, Rocha S, Santos H, Paiva MR, Burban C, Kerdelhué C, et al. Climate constrains range expansion of an allochronic population of the pine processionary moth. Divers Distrib 2016;22:1288–300. https://doi.org/10.1111/ddi.12494.

Hall MC, Willis JH. Divergent selection on flowering time contributes to local adaptation in Mimulus guttatus populations. Evolution (N Y) 2006;60:2466–77. https://doi.org/10.1111/j.0014-3820.2006.tb01882.x.

Hattori I, Ito Y. Status black-headed and red-headed types of Hyphantria cunea (Drury) (Lepidoptera : Arctiidae) : II. External characters of the two types and their hybrids. Appl Entomol Zool 1973;8:172–82. https://doi.org/10.1303/AEZ.8.172.

Hillebrand H. On the generality of the latitudinal diversity gradient. American Naturalist 2004;163:192–211. https://doi.org/10.1086/381004.

Hodgson JA, Thomas CD, Oliver TH, Anderson BJ, Brereton TM, Crone EE. Predicting insect phenology across space and time. Glob Chang Biol 2011;17:1289–300. https://doi.org/10.1111/j.1365-2486.2010.02308.x.

Hollister J. Access Elevation Data from Various APIs [R package elevatr version 0.3.4] 2021.

Inskeep KA, Doellman MM, Powell THQ, Berlocher SH, Seifert NR, Hood GR, et al. Divergent diapause life history timing drives both allochronic speciation and reticulate hybridization in an adaptive radiation of Rhagoletis flies. Mol Ecol 2021. https://doi.org/10.1111/MEC.15908.

Jaenike J, Selander RK. On the question of host races in the fall webworm, Hyphantria cunea. Entomol Exp Appl 1980;27:31–7. https://doi.org/10.1111/j.1570-7458.1980.tb02943.x.

Jaskuła R, Kolanowska M, Michalski M, Schwerk A. From phenology and habitat preferences to climate change: Importance of citizen science in studying insect ecology in the continental scale with american red flat bark beetle, cucujus clavipes, as a model species. Insects 2021;12:369. https://doi.org/10.3390/INSECTS12040369/S1.

Kuhn M. Classification and Regression Training [R package caret version 6.0-88] 2021.

Lehtinen RM, Carlson BM, Hamm AR, Riley AG, Mullin MM, Gray WJ. Dispatches from the neighborhood watch: Using citizen science and field survey data to document color morph frequency in space and time. Ecol Evol 2020;10:1526. https://doi.org/10.1002/ECE3.6006.

Li YP, Goto M, Ito S, Sato Y, Sasaki K, Goto N. Physiology of diapause and cold hardiness in
the overwintering pupae of the fall webworm Hyphantria cunea (Lepidoptera: Arctiidae) in
Japan. J Insect Physiol 2001;47:1181–7. https://doi.org/10.1016/S0022-1910(01)00099-3.

Loewy K. Host Use and Geographic Variation in Fall Webworms (Hyphantria cunea). Electronic
Theses and Dissertations 2013.

Map Downloads. USDA Agricultural Research Services n.d.
https://planthardiness.ars.usda.gov/pages/map-downloads (accessed November 20, 2022).

Mattsson M, Hood GR, Feder JL, Ruedas LA. Rapid and repeatable shifts in life-history timing
of Rhagoletis pomonella (Diptera: Tephritidae) following colonization of novel host plants in the
Pacific Northwestern United States. Ecol Evol 2015;5:5823–37.
https://doi.org/10.1002/ECE3.1826.

Mazerolle MJ. AICcmodavg: Model Selection and Multimodel Inference Based on (Q)AIC(c)
2020.

Mittelbach GG, Schemske DW, Cornell H v., Allen AP, Brown JM, Bush MB, et al. Evolution
and the latitudinal diversity gradient: Speciation, extinction and biogeography. Ecol Lett
2007;10:315–31. https://doi.org/10.1111/j.1461-0248.2007.01020.x.

Morris RF. Synonymy and color variation in the fall webworm, Hyphantria cunea Drury
(Lepidoptera: Arctiidae). Can Entomol 1963;95:1217–23. https://doi.org/10.4039/ENT951217-
11.

Nosil P, Harmon LJ, Seehausen O. Ecological explanations for (incomplete) speciation. Trends
Ecol Evol 2009;24:145–56. https://doi.org/10.1016/j.tree.2008.10.011.

Oliver AD. A behavioral study of two races of the fall webworm, Hyphantria cunea,
(Lepidoptera: Arctiidae) in Louisiana. Ann Entomol Soc Am 1964;57:192–4.
https://doi.org/10.1093/aesa/57.2.192.

Ording GJ, Mercader RJ, Aardema ML, Scriber JM. Allochronic isolation and incipient hybrid
speciation in tiger swallowtail butterflies. Global Change Ecology 2009;162:523–32.

Post E. The Phenological Niche. Time in Ecology, Princeton University Press; 2019, p. 67–93.
https://doi.org/10.1515/9780691185491-006/HTML.

R Core Team. R Core Team (2021). R: A language and environment for statistical computing. R
Foundation for Statistical Computing, Vienna, Austria URL Http://WwwR-ProjectOrg/ 2021.

Régnière J, St-Amant R, Béchard A. BioSIM 10 – User's manual. 2014.

Ripley BD, Venables B. MASS [R package MASS version 7.3-54]. Modern Applied Statistics
with S. Fourth Edition, New York: Springer; 2002.

Roy HE, Baxter E, Saunders A, Pocock MJO. Focal Plant Observations as a Standardised
Method for Pollinator Monitoring: Opportunities and Limitations for Mass Participation Citizen
Science. PLoS One 2016;11:e0150794. https://doi.org/10.1371/JOURNAL.PONE.0150794.

Schowalter TD, Ring DR. Biology and management of the fall webworm, Hyphantria cunea (Lepidoptera: Erebidae). J Integr Pest Manag 2017;8:7. https://doi.org/10.1093/jipm/pmw019.

Scranton K, Amarasekare P. Predicting phenological shifts in a changing climate. Proc Natl Acad Sci U S A 2017;114:13212–7. https://doi.org/10.1073/pnas.1711221114.

Sobel JM, Chen GF, Watt LR, Schemske DW. The biology of speciation. Evolution (N Y) 2010;64:295–315. https://doi.org/10.1111/j.1558-5646.2009.00877.x.

Takeda M. Differentiation in life cycle of sympatric populations of two forms of Hyphantria moth in central Missouri. Entomol Sci 2005;8:211–8. https://doi.org/10.1111/j.1479-8298.2005.00117.x.

Tennekes M. Tmap: Thematic maps in R. J Stat Softw 2018;84. https://doi.org/10.18637/JSS.V084.I06.

Tiroesele B, Skoda SR, Hunt TE, Lee DJ, Ullah MI, Molina-Ochoa J, et al. Morphological and Genetic Analysis of Four Color Morphs of Bean Leaf Beetle. Journal of Insect Science 2018;18. https://doi.org/10.1093/JISESA/IEY016.

Tobin PC, Nagarkatti S, Loeb G, Saunders MC. Historical and projected interactions between climate change and insect voltinism in a multivoltine species. Glob Chang Biol 2008;14:951–7. https://doi.org/10.1111/J.1365-2486.2008.01561.X.

Tseng M, Bevanda C, Bhatti SS, Black EN, Chang E, Chiang J, et al. Effects of temperature on monarch caterpillar pigment variation in nature. Insect Conserv Divers 2022. https://doi.org/10.1111/ICAD.12608.

USDA Plant Hardiness Zone Map. USDA Agricultural Research Service 2012. https://planthardiness.ars.usda.gov/ (accessed November 20, 2022).

Vidal MC, Quinn TW, Stireman JO, Tinghitella RM, Murphy SM. Geography is more important than host plant use for the population genetic structure of a generalist insect herbivore. Mol Ecol 2019;28:4317–34. https://doi.org/10.1111/mec.15218.

Wadsworth CB, Okada Y, Dopman EB. Phenology-dependent cold exposure and thermal performance of Ostrinia nubilalis ecotypes. BMC Evol Biol 2020;20:1–14. https://doi.org/10.1186/s12862-020-1598-6.

Wadsworth CB, Woods WA, Hahn DA, Dopman EB. One phase of the dormancy developmental pathway is critical for the evolution of insect seasonality. J Evol Biol 2013;26:2359–68. https://doi.org/10.1111/jeb.12227.

Warren LO, Tadic M. fall webworm, Hyphantria cunea (Durry). Ark Agr Exp Sta Bull 1970;v. Nov 1970, 759.

Weir JT, Schluter D. The latitudinal gradient in recent speciation and extinction rates of birds and mammals. American Association for the Advancement of Science 2007;315:1574–6. https://doi.org/10.1126/science.ll39024.

Weiser MD, Siler CD, Smith SN, Marshall KE, McLaughlin JF, Miller MJ, et al. Robust
metagenomic evidence that local assemblage richness increases with latitude in ground-active
invertebrates of North America. Oikos 2022;2022. https://doi.org/10.1111/OIK.08791.

Wickham H, Averick M, Bryan J, Chang W, McGowan L, François R, et al. Welcome to the
Tidyverse. J Open Source Softw 2019;4:1686. https://doi.org/10.21105/joss.01686.

Wu N, Zhang S, Li X, Cao Y, Liu X, Wang Q, et al. Fall webworm genomes yield insights into
rapid adaptation of invasive species. Nat Ecol Evol 2019;3:105–15.
https://doi.org/10.1038/s41559-018-0746-5.

Yamamoto S, Sota T. Incipient allochronic speciation by climatic disruption of the reproductive
period. Proceedings of the Royal Society B: Biological Sciences 2009;276:2711–9.
https://doi.org/10.1098/RSPB.2009.0349.

Yang F, Kawabata E, Tufail M, Brown JJ, Takeda M. r/K-like trade-off and voltinism
discreteness: The implication to allochronic speciation in the fall webworm, Hyphantria cunea
complex (Arctiidae). Ecol Evol 2017;7:10592–603. https://doi.org/10.1002/ece3.3334.

Figure Captions

Figure 1. Distribution of red and black morph fall webworm observations on iNaturalist from
2018-2020 in Canada and the United States. Each point corresponds to a single iNaturalist
observation. USDA plant hardiness zones indicated on map to show differences in geographic
distribution among morphs. Only observations east of -100°W longitude were used in further
analysis. A smaller value of plant hardiness zone correlates to colder climates existing at higher
latitudes. n=4472 for black morph, n=775 for red morph.

Figure 2. The red and black fall webworm colour morphs have different distributions across
latitude and longitude but not elevation in their eastern North American range (east of -100°W
longitude). Violin plots indicate scaled density of observations at each value. Asterisks indicate
groups are significantly different according to Tukey Post-Hoc test, and "ns" indicates no
significance. The geographical range of the black morph averages 2.3 degrees further north
(P<0.0001) and 4.1 degrees further east (P<0.0001) than the red morph, but there was no
difference in average elevation between the two colour morphs (P=0.8). n=5176 for latitude and
longitude, n=4652 for elevation.

Figure 3. Phenology of red and black morph fall webworm across years and plant hardiness
zones. Points indicate scaled larval abundance in each week of the year, while lines show trends
in abundance across years calculated using the "loess" function.

Figure 4. Difference between mean earliest 10% and latest 10% of observation dates between red
and black morphs across plant hardiness zones. Points show mean of earliest or latest dates, and
error bars show standard error of the mean. **A)** The earliest 10% of black morph observations
occurred an average of 22 ± 3.35 (standard error of the mean difference) days earlier than the red
morph (P<0.0001). Differences between mean dates of the morphs increased in larger plant
hardiness zones (P<0.0001). n=529. **B)** There was no significant difference between the average
latest observance date of the red and black morphs (P=0.05). The black morph tended to occur
later than the red in increasing plant hardiness zone (P<0.0001); however, there were no trends in
the differences between morphs across zones. n=525.

Figure 5. Differences between cumulative distribution functions of red and black colour morphs
increase at warmer plant hardiness zones. Points indicate value of DTS function (overlap
between CDFs) of colour morph populations at each plant hardiness zone and year, lines indicate
linear regression of points in each year. Differences between temporal abundance of the morphs
increased at warmer plant hardiness zones (P<0.001). n=15.

Figure 6. Colouration of black and red morph fall webworm larvae diverges more at warmer
plant hardiness zones. Line type indicates year, while line colour indicates larval colour morph.
Points indicate mean LD1 values of morphs in each plant hardiness zone and year. Grey shaded

region shows 95% confidence interval of each year and morph. Red morph LD1 values do not
change across zones (P=0.5) while black morph LD1 values decrease at warmer plant hardiness
zones (P<0.0001). n=3475.