Non-conventional serine protease activity of the CXC chemokine-cleaving streptococcal

enzyme, SpyCEP

Max Pearson[1,2], Carl Haslam[3], Andrew Fosberry[3], Emma J Jones[3], Mark Reglinski[1], Robert J.
Edwards[4], Richard Ashley Lawrenson[1], Danuta Mossakowska[3,5], James Edward Pease[6], Shiranee
Sriskandan[1, 2]

[1]Department of Infectious Disease, Imperial College London, London W12 0NN, UK

[2]MRC Centre for Molecular Bacteriology & Infection, Imperial College London, London, SW7 2AZ, UK

[3]GlaxoSmithKline R&D, Gunnels Wood Road, Stevenage, Hertfordshire, SG1 2NY, UK

[4]Department of Medicine, Imperial College London, W12 0NN, UK

[5]Malopolska Centre of Biotechnology, Jagiellonian University, 30-387 Kraków, Poland

[6]National Heart and Lung Institute, Imperial College London, London, SW7 2AZ, UK

Character count: 46516

<u>Abstract</u>

The *Streptococcus pyogenes* cell envelope protease (SpyCEP) is a vital virulence factor in

streptococcal pathogenesis. Despite its key role in disease progression and strong association with

invasive disease, little is known about the enzymatic function beyond the ELR[+] CXC chemokine

substrate range. We utilised multiple SpyCEP constructs to interrogate the protein domains and

catalytic residues necessary for enzyme function. We leveraged high-throughput mass spectrometry

to describe the Michaelis-Menton parameters of active SpyCEP, revealing a Michaelis-Menton

constant (K_M) of 53.49 nM and a turnover of 1.34 molecules per second, for the natural chemokine

substrate CXCL8.

Unexpectedly, we found that an N-terminally-truncated SpyCEP C-terminal construct consisting of

only the H279 and S617 catalytic dyad had specific CXCL8 cleaving activity, albeit with a reduced

substrate turnover of 2.45 molecules per hour, representing a ~2000-fold reduction in activity. In

contrast, the K_M of the C-terminal SpyCEP construct and full-length enzyme did not differ. We

conclude that the SpyCEP C-terminus plays a key role in substrate binding and recognition with key

implications for both current and future streptococcal vaccine designs.

Word count: 175

<u>Introduction</u>

Group A Streptococcus (GAS) or *Streptococcus pyogenes* is a leading human pathogen that

manifests clinically as a broad spectrum of diseases, ranging from less severe, usually self-limiting

infections to life-threatening invasive diseases such as necrotizing fasciitis and toxic shock syndrome.

While much of the global health burden can be attributed to streptococcal auto-immune sequelae

such as rheumatic heart disease, invasive diseases contribute greatly to *S. pyogenes* associated

global mortality. Invasive infections account for an estimated 163,000 deaths worldwide per year

and at least 663,000 new cases (Carapetis et al., 2005) with mortality rates remaining high despite

intervention; indeed, 20% of patients with invasive *S. pyogenes* disease die within 7 days of infection

(Ralph and Carapetis, 2013).

Several virulence factors contribute to pathogenesis in invasive *S. pyogenes* disease, chief

among which is <u>*Streptococcus pyogenes* c</u>ell <u>e</u>nvelope <u>p</u>rotease (SpyCEP), an immune-modulatory

cell wall-associated protease. SpyCEP is responsible for the rapid and efficient cleavage of a distinct

group of CXC chemokines comprising CXCL1, CXCL2, CXCL3, CXCL5, CXCL6, CXCL7 and CXCL8 both

locally at the site of an infection, and systemically, (Edwards et al., 2005, Kurupati et al., 2010,

Turner et al., 2009a, Zingaretti et al., 2010). The ELR$^+$ chemokines, named for their conserved N-

terminal glutamate- leucine- arginine motifs, specifically act upon neutrophils eliciting their

recruitment and activation. SpyCEP inactivates these chemokines by cleaving the chemokine C-

terminal α-helix, releasing a 13 amino acid peptide in the case of CXCL8 (Edwards et al., 2005).

Specificity for binding the neutrophil chemokine receptors CXCR1 and CXCR2 is conferred to a large

extent by the chemokine N-terminal ELR motif (Middleton et al., 1997), although the chemokine C-

terminus is necessary for efficient receptor binding and activation (Goldblatt et al., 2019) in addition

to chemokine translocation from tissues to endothelial lumen (Middleton et al., 1997). As such,

SpyCEP cleavage of CXC chemokines results in a reduction of CXCR1 and CXCR2-mediated neutrophil

chemotaxis and subsequent paucity of neutrophils at the site of *S. pyogenes* infection (Edwards et

al., 2005). In addition to CXC chemokines, SpyCEP has recently been shown to cleave the human

anti-microbial peptide, LL-37 (Biswas et al., 2021). Whilst cleavage does not affect the antimicrobial

action of the peptide, it was reported to reduce LL-37 specific neutrophil chemotaxis (Biswas et al.,

2021).

SpyCEP is expressed by *S. pyogenes* as a 1647 amino acid, 180 kDa subtilisin-like serine

protease, the crystal structure of which was recently solved to 2.8 Å resolution (Jobichen et al.,

2018) and further refined to 2.2 Å resolution (McKenna et al., 2020). It is a member of the S8

subtilase family, members of which are characterised by a catalytic triad consisting of an aspartate,

histidine and serine residue each surrounded by a region of highly conserved amino acids (Siezen,

1999). SpyCEP is unique among streptococcal proteases in that, during maturation, it is

autocatalytically cleaved between residues Q244 and S245 into 2 distinct polypeptides, a 30 kDa N-

terminal polypeptide and a 150 kDa C-terminal polypeptide. The two polypeptides harbour the

separate residues required for the formation of the catalytic site (Zingaretti et al., 2010); the N-

terminal fragment contains the catalytic D151 and the C-terminal fragment contains the catalytic

H279 and S617 residues. Upon cleavage, the two polypeptides re-associate non-covalently to

reconstitute the active enzyme (Zingaretti et al., 2010). The recent crystal structures have shed

further light upon the structure of SpyCEP, describing 9 separate domains, the first 5 of which

contain the catalytic triad necessary for enzymatic activity (Jobichen et al., 2018).

SpyCEP has been included as a target antigen in several recent *S. pyogenes* vaccine designs due to its

cell surface expression, highly conserved nature, and central role in *S. pyogenes* pathogenesis.

Immunisation with SpyCEP successfully elicits a SpyCEP-specific neutralising antibody response,

providing protection against systemic bacterial dissemination and reducing disease severity in *S.*

pyogenes intramuscular, skin infection models and non-human primate infection models (Rivera-

Hernandez et al., 2019, Bensi et al., 2012, Pandey et al., 2015, Pandey et al., 2016, Rivera-Hernandez

et al., 2016, Turner et al., 2009b). Data that demonstrate vaccine dependence on enzyme inhibition

highlight the importance of understanding the enzymatic activity of SpyCEP and the potential to

improve upon vaccine or inhibitor design.

Despite this, little is known about the catalytic properties of the enzyme, except preliminary

structure-function studies (Kaur et al., 2010, Jobichen et al., 2018, McKenna et al., 2020, Zingaretti et

al., 2010). Published data largely focuses on the impact of SpyCEP on streptococcal pathogenesis and

disease progression. In this study we generated multiple SpyCEP constructs to confirm domains

necessary for catalytic activity. We then used mass spectrometry to determine the enzyme kinetics

of multiple SpyCEP constructs for the natural substrate CXCL8 .

Methods

Cloning and purification of recombinant SpyCEP constructs

Codon-optimized SpyCEP gene constructs were expressed in *Escherichia coli* using synthetic gene

sequences (GenScript) from Spy_0416 in the SF370 *S. pyogenes* M1 genome (Abate et al., 2013,

Zingaretti et al., 2010) representing full-length enzyme, SpyCEP$^{34-1613}$, the N- terminal polypeptide,

SpyCEP$^{34-244}$, and C-terminal polypeptide SpyCEP$^{245-1613}$. Constructs were also generated with alanine

substitutions to replace catalytic residues in the N-terminal fragment (SpyCEP$^{34-244}$ D151A), and the

C-terminal fragment (SpyCEP$^{245-1613}$ S617A). To enable downstream protein purification, both the

full-length enzyme and C-terminal polypetides were synthesized with a C-terminal 6X-Histidine tag

and the N-terminal polypeptides were synthesized with an N-terminal FLAG tag and TEV linker.

All SpyCEP constructs were cloned into the vector pET-24B and expressed in BL21 (DE3) competent

E. coli, cultured in Terrific Broth (TB) medium supplemented with 50 µg/mL kanamycin at 37°C, 200

rpm, for 3 hours. The cultures were induced with 0.5 mM Isopropyl β-D-1-thiogalactopyranoside

(IPTG), cooled to 15$^{\text{o}}$C and incubated at 200 rpm for 16 hours before lysis by sonication on ice. Full-

length and C-terminal contructs were purified by Ni-IDA affinity chromatography (GenScript) as per

the manufacturer's instructions. N-terminal constructs were purified by anti-flag M2 agarose resin

chromatography (Sigma-Aldrich, A2220) as per manufacturers instruction, and further concentrated

and purified by SP ion-exchange (Sigma-Aldrich, 17-1152-01) and Q FF ion-exchange

chromatography (GE Healthcare, 17-5156-01) as per manufacturer's instruction. All SpyCEP

constructs were subsequently concentrated and purified by size exclusion chromatography on a

HiLoad Superdex 200 or 75 prep grade column (GE Healthcare) dependent on molecular weight.

The resultant panel of recombinantly expressed SpyCEP constructs is shown in Table 1. The re-

association of the SpyCEP N and C-termini to create full-length constructs was carried out by

equimolar co-incubation at 37^{o}C for 30 minutes in 40 mM Tris-HCL pH 7.5, 0.1 mM CHAPS, 1 mM

DTT, 0.1% BSA, 75 mM NaCl.

Table 1. SpyCEP constructs used in this study. Constructs were expressed recombinantly in *E. coli*
except for s.pSpyCEP that was expressed in, and purified from *S. pyogenes* .

Construction of recombinant *S. pyogenes* expressing histidine-tagged SpyCEP
148

To permit release of soluble His-tagged SpyCEP by *S. pyogenes*, the C-terminal anchor domain of

SpyCEP was replaced by a stop codon in strain H292, a strain that makes abundant SpyCEP (Turner et

al., 2009a). A 549 base pair region immediately upstream of the SpyCEP cell wall anchor motif was

amplified from *S. pyogenes* H292 genomic DNA using the primers

:5'GGGAATTCTGTTGTCAGGTAACAGTCTTATCTTGCC -3' and

5'CCGAATTCACAACACTAGGCTTTTGCTGAGGTCGTTG -3'. *EcoRI* restriction sites were incorporated at

the terminal ends of the primer sequences. The amplified DNA was cloned into the homologous

recombination plasmid pUCMUT to produce the vector pUCMUT$_{CEP}$ which was transformed into One

Shot TOP10 Chemically Competent *E. coli* (Thermo Fisher Scientific) according to the manufacturer's

instructions. The nucleotide sequence of the SpyCEP C-terminus was subsequently further modified

to encode a hexa-histidine tag by inverse PCR using pUCMUT$_{CEP}$ as the template and the primers 5'-

TATCCTAGGTAGTGTTGTGAATTCGTAATCATGGTCATAG-3' and 5'-

TATCCTAGGATGATGATGATGATGATGGGCTTTTGCTGAGGTCGTTG-3'. The amplification was

preformed using GoTaq Long PCR Master Mix (Promega) according to the manufacturer's

instructions. *AvrII* restriction sites, incorporated at the terminal ends of the primer sequences,

facilitated re-circulation of the amplified plasmid. The presence of a hexa-his sequence in the

modified pUCMUT$_{CEP}$ construct (denoted pUCMUT$_{CEP-HIS}$) was confirmed by Sanger sequencing using

the pUCMUT sequencing primers 5'- GACAGCAACATCTTTGTGAAAGATGG-3' and 5'-

CATTAATGCAGCTGGCACGAC-3'. The pUCMUT$_{CEP-HIS}$ construct was introduced into H292 by

electroporation and crossed into the chromosome by homologous recombination as previously

described (Lynskey et al., 2013) to generate strain H1317. Secretion of His- tagged SpyCEP into the

culture supernatant of H1317 was confirmed by western blotting (data not shown). To purify

SpyCEP, *S. pyogenes* H1317 was grown in Todd-Hewitt broth (Oxoid) overnight at 37^{o}C with 5% C0$_2$

The culture was pelleted at 2500 xg for 10 minutes and the supernatant sterilised using Amicon 0.22

μM filters, then concentrated using 15 ml Amicon 100 kDa centrifuge filter columns and purified by

nickel column affinity chromatography (Novagen His-Bind Resin) as per the manufacturer's

instructions.

Sodium dodecyl sulphate–polyacrylamide gel electrophoresis (SDS-page)

To visualise chemokine cleavage, 5 μM recombinant human CXCL1 and CXCL8 (R&D Systems) was

incubated with full-length recombinant SpyCEP or C-terminal SpyCEP at a molar ratio of 1:5, 50, 500,

5000 in favour of chemokine for 2 hours at 37^{O}C. SpyCEP[DASA], C-terminal[SA] and N-terminal

constructs[34-244] were included as controls and assayed at the highest 1:5 molar ratio. Reactions were

halted by the addition of Dithiothreitol (DTT) to a final concentration of 100 mM, 4X Bolt LDS sample

buffer and heating to 70°C for 10 minutes. Samples were separated on pre-cast 4-12% MES buffered

Bolt Bis-Tris gradient gels (Invitrogen) by SDS-PAGE gel electrophoresis at 165 V for 35 minutes with

SeeBlue Plus 2 (Invitrogen) used for molecular weight ladder. Gels were stained with PageBlue

protein staining solution (Thermo Fisher Scientific) overnight and de-stained in deionised water.

Multiplex fluorescent western blotting

To visualise generation of both intact CXCL8 and the larger (N-terminal) CXCL8 cleavage product, a

rabbit antiserum was raised against the neo-epitope (anti-ENWVQ) that is exposed following SpyCEP

cleavage of CXCL8 (Edwards et al., 2007). SpyCEP constructs were incubated at 37 ^{o}C with 937.5 nM

of recombinant human CXCL8 (R&D Systems) at a 1:50 molar ratio in favour of CXCL8. Digests were

halted and separated by SDS-PAGE gel electrophoresis as described above. Proteins were

transferred by iBlot2 (Thermo Fisher Scientific) onto 0.22 μm nitrocellulose membranes as per

manufacturer's instructions. Membranes were subsequently blocked for 1 hour at room

temperature in blocking buffer (PBS with 5% skimmed milk powder (Sigma-Aldrich) and 0.1%

Tween), then blotted overnight at 4 ^{o}C with 2 primary antibodies, 1 μg/ml mouse anti-human CXCL8

(R&D Systems) and 1:1000 rabbit anti-ENWVQ. Membranes were washed in wash buffer (PBS with

0.05% Tween) and incubated with 1:7500 goat anti-rabbit IgG A680nm and 1:7500 goat anti-mouse

IgG A790nm for 1 hour at room temperature before being visualised on LiCor Odyssey Fc

(Invitrogen).

SpyCEP activity against LL-37 was assessed by an 16 hour, 37 ^{o}C incubation of 5.56 μM human LL-37

(R&D Systems) with SpyCEP constructs at a 1:10 molar ratio in favour of LL-37. Reactions were

stopped, separated and blotted onto 0.22 μM nitrocellulose as above and incubated overnight at 4 ^{o}C

in blocking buffer supplemented with 2 μg/ml polyclonal sheep IgG anti-LL-37 (R&D Systems).

Membranes were washed in wash buffer (PBS with 0.05% Tween) and incubated in blocking solution

with rabbit anti-sheep IgG (Abcam) at 1:40,000 dilution for 1 hour at room temperature and

visualised on LiCor Odyssey Fc (Invitrogen).

Enzyme linked immunosorbent assay

Catalytic activities of SpyCEP constructs were measured through detection of remaining intact CXCL8

or CXCL1 substrate following incubation with SpyCEP by ELISA (R&D Systems human CXCL8 and

CXCL1 DuoSet ELISA) as per manufacturer's instructions. For cleavage time courses 50 pM or 100 pM

SpyCEP were incubated with an equal volume of 20 nM human CXCL8 and CXCL1 respectively (R&D

Systems) and incubated at room temperature for 30 minutes. Reactions were halted at defined

timepoints with the addition of concentration of Pefabloc (Sigma-Aldrich) to a final concentration of

2 mg/ml (8.34 mM).

Linear regression analyses of the initial five time points of CXCL1 and CXCL8 cleavage (0, 1, 2, 3 and 4

minutes) was utlised to determine the maximal rate of SpyCEP activity.

Mass spectrometry analysis of CXCL8 cleavage

Analysis of CXCL8 cleavage was assayed on a SCIEX API6500 triple quadrupole electrospray mass

spectrometer coupled to a high-throughput robotic sample preparation and injection system,

RapidFire200 (Agilent Technologies). CXCL8 substrate, the 13 amino acid CXCL8 SpyCEP cleavage

product RVVEKFLKRAENS, and a heavy atom substituted internal standard of CXCL8 SpyCEP cleavage

product RV[$U_{13}C_5$ $_{15}$N-VAL]-EKF-[$U_{-13}C_6$ $_{15}$N-Leu]- KRAENS) were monitored by mass spectrometry. The

mass spectrometer was operated in positive electrospray MRM mode, and transitions (Q1/Q3) for

each species were optimised to give m/z as follows: CXCL8, 1048.7/615.3, CXCL8 cleavage product

526.2/211.1, internal standard 530.5/211.1. A dwell time of 50 ms was used for the MRM

transitions. The mass spectrometer was operated with a spray voltage of 5500 V and at a source

temperature of 650 °C.

To assay CXCL8 cleavage dynamically, chemokine and SpyCEP constructs were loaded into a 384 well

plate to the following final concentrations: 6.25-2000nM chemokine, 250 pM SpyCEP or 40 nM C-

terminal SpyCEP. Reactions were stopped at desired time points, 0-240 minutes, by the addition of

stop solution (1% formic acid) supplemented with 1 µM heavy atom substituted CXCL8 internal

standard and centrifuged 2000 xg for 10 minutes. Assay plates were transferred onto the

RapidFire200 integrated to the API6500 mass spectrometer. Samples were aspirated under vacuum

directly from 384-well assay plates for 0.6 s. The samples were then loaded onto a C18 solid-phase

extraction cartridge to remove non-volatile buffer salts, using HPLC -grade water supplemented with

0.1% (v/v) formic acid at a flow rate of 1.5 mL/min for 4 s. The retained analytes were eluted to the

mass spectrometer by washing the cartridge with acetonitrile HPLC-grade water (8:2, v/v) with 0.1%

(v/v) formic acid at 1.25 mL/min for 4 s. The cartridge was re-equilibrated with HPLC-grade water

supplemented with 0.1% (v/v) formic acid for 0.6 s at 1.5 mL/min. Results were normalised to the

fixed internal standard and converted to molarity by interpolation from a standard curve of known

cleaved CXCL8 concentrations.

Kinetic analysis

Linear regression analyses of the initial five time points of full length SpyCEP (0, 1, 2, 3, 4 minutes)

and C-terminal SpyCEP (0, 15, 30, 45, 60 minutes) CXCL8 reactions were plotted against substrate

concentration and kinetics derived from the Michaelis-Menton equation $Y = V_{max}*X/(K_M + X)$ and the

K_{cat} equation $Y = ET*k_{cat}*X/(K_M + X)$ where Et = enzyme concentration as fitted by Prism 8.0.2

(GraphPad).

Results

Screening of human CXCL8 cleavage activity using fluorescent western blotting

To initially assess the activity of our recombinant SpyCEP constructs we screened them for CXCL8

cleavage activity using two-colour multiplex western blotting. Human CXCL8 was incubated for 2

hours at 37 °C with SpyCEP constructs, the reaction products were separated by SDS-PAGE and

immunoblotted using seperate antibodies that detect either intact or cleaved CXCL8. Detection of

full-length CXCL8 (8 kDa, green bands) or the CXCL8 neo epitope (ENWVQ), exposed after SpyCEP

cleavage, (6 kDa, red bands) was evident with this system (Figure 1). Both native *S. pyogenes* SpyCEP

and recombinant full-length SpyCEP successfully cleaved CXCL8 to completion (Figure 1). As

expected, no cleavage was observed when using the catalytically dead mutant, SpyCEP^{DASA}. As has

been previously reported (Zingaretti et al., 2010), the SpyCEP N- and C-termini, when independently

expressed and purified, can be re-associated to form an active enzyme, which successfully cleaved

CXCL8. The N-terminal fragment of SpyCEP alone could not cleave CXCL8. Unexpectedly however,

the C-terminal fragment of SpyCEP was able to cleave CXCL8, albeit not to completion. This

independent catalytic activity was negated by mutation of the catalytic S617 to alanine (C-terminal^{SA}

construct, Figure 1). Cleavage of CXCL8 by the SpyCEP C-terminal fragment was enhanced when re-

associated with the catalytically inert N-terminal mutant (N-terminal^{DA} construct); indeed, activity

was equivalent to that observed with the re-associated enzyme under these conditions, with

cleavage of CXCL8 to near completion. However, the N-terminal^{DA} construct was unable to restore

catalytic activity to the C-terminal^{SA} construct when the two were combined.

The results demonstrated for the first time that the SpyCEP C-terminal fragment alone is sufficient

for catalytic cleavage of CXCL8 in this assay system. The data also established that, although the

presence of the SpyCEP N-terminal fragment enhanced enzymatic activity, the N-terminal residue

D151 was dispensable for enzymatic activity. We further sought to examine whether the C-terminal

displayed activity against the newly described substrate, LL-37 (Biswas et al., 2021). Western blot

analysis confirmed cleavage of LL-37 by full-length recombinant SpyCEP, demonstrated by a

reduction in band size, however the C-terminal fragment was unable to cleave LL-37 despite a high

molar ratio of enzyme to LL-37 (1: 10) and a prolonged 16-hour incubation at 37 °C (Figure EV1).

Figure 1. Cleavage activity of recombinant SpyCEP constructs assayed by immunoblot. Two colour
immunoblot showing cleavage of 150 ng CXCL8 incubated for 2 hours at 37 °C either alone (2[nd] lane)
or with panel of SpyCEP constructs at a 1:50 molar ratio (SpyCEP:CXCL8). Green bands represent
intact CXCL8 (anti-CXCL8 antibody); red bands represent cleaved CXCL8 (anti- ENWVQ). A 17 kDa
molecular weight marker is shown in blue, 8 kDa and 6 kDa molecular weights are highlighted by
arrows. Figure is representative of 2 independent immunoblots.

Differntial cleavage of CXCL8 and CXCL1 by full-length and C-terminal SpyCEP
290

To determine whether the catalytic activity of the C-terminal SpyCEP fragment was reproducible

over a shorter incubation period, we assessed CXCL8 cleavage by ELISA, where CXCL8 cleavage is

detected through a reduction in substrate concentration. SpyCEP C-terminal constructs were

incubated with CXCL8 at molar ratios ranging from 1:5 – 1:250 (enzyme: chemokine) over a 60

minute timecourse at room temperature. Full-length recombinant SpyCEP and the inactive C-

terminal fragment, C-terminal[SA], were included as controls at a molar ratio of 1:1000 and 1:5,

respectively. Under these conditions, near complete CXCL8 cleavage was observed for full-length

SpyCEP and a dose-dependent increase in catalytic activity was observed for the C-terminal fragment

(Figure 2). After 5 minutes incubation full-length SpyCEP cleaved over 50% of the starting CXCL8

input, with only 6.25% of CXCL8 remaining after 1 hour. At the highest concentration of C-terminal

SpyCEP tested, a 1:5 molar ratio, the C-terminal fragment alone cleaved 9% of the starting CXCL8 by

5 minutes, 25% by 30 minutes and 42% by 60 minutes. Indeed, at the lowest concentration tested, a

1:250 molar ratio, the C-terminal of SpyCEP cleaved 9% of CXCL8 input by 1 hour. As demonstrated

by immunofluorescent western blotting, the serine residue at position 617 was vital for SpyCEP

catalytic function as no CXCL8 cleavage was observed using the C-terminal[SA] construct, even when

employed at the highest enzyme: chemokine ratio. After a 1 hour incubation, full length SpyCEP and

C-terminal contructs (assayed using a molar enzyme: chemokine ratio of 1:5, 1:25 and 1:50) cleaved

significantly more CXCL8 compared to the C-terminal[SA] construct.

Figure 2. Cleavage activity of SpyCEP and C-terminal SpyCEP[245-1613] constructs using CXCL8 ELISA.
SpyCEP constructs were co-incubated with CXCL8. Graphs show residual CXCL8 after a 60-minute
room temperature incubation, using full length SpyCEP at a 1:1000 ratio to CXCL8; C-terminal at 1:5
– 1:250 molar ratio to CXCL8; and C-terminal S617A mutant at a 1:5 molar ratio to CXCL8. Reactions
were halted at specified timepoints by the addition of Pefabloc to a final concentration of 2 mg/ml.
N=6 experimental replicates for each construct, data points show means, error bars represent SD. ns
$p > 0.05$, * $p \leq 0.05$, **** $p \leq 0.0001$, at 60 minutes as determined by ordinary one-way ANOVA.

Further SDS-PAGE analysis of the catalytic activity of the C-terminal SpyCEP construct additionally

showed that C-terminal activity was not restricted to the CXCL8 substrate. Over 2 hours at 37 ^{o}C

using a 1:5 molar ratio of enzyme: substrate, the SpyCEP C-terminal construct was capable of

cleaving human CXCL1 to near completion (Figure EV2).

To further assess the activity of SpyCEP and to interrogate reaction rates against separate

chemokines, full-length SpyCEP was incubated with CXCL1 and CXCL8 and the remaining chemokine

levels determined by ELISA. Human CXCL1 or human CXCL8 were incubated with SpyCEP over a 30

minute, room temperature timecourse, at 1:200 or 1:400 molar ratios respectively (enzyme:

chemokine). When incubated in equal volumes, 50 pM SpyCEP rapidly and efficiently cleaved 20 nM

CXCL8, with ~ 15% of the chemokine input remaining after 10 minutes of incubation (Figure 3A). This

contrasted with CXCL1 cleavage, that required 100 pM SpyCEP to cleave just 25% of the chemokine

input over the same 10-minute period. Indeed, by 30 minutes SpyCEP had yet to cleave half of the

starting CXCL1 (Figure 3B). Utilising a linear regression of the initial 5 timepoints, where SpyCEP

activity is maximal, we found that 5 fmol of SpyCEP was able to cleave 284 fmol of CXCL8 per

minute, and 10 fmol of SpyCEP was capable of cleaving 62 fmol of CXCL1 per minute. These data

confirmed the activity of recombinant SpyCEP and highlighted differential cleavage efficiency across

the CXC substrate range – a feature which has been previously recognised but not quantified

(Goldblatt et al., 2019).

Figure 3. Cleavage of CXCL8 and CXCL1 by recombinant full length SpyCEP. 30-minute, room
temperature cleavage time course of: A. 20nM CXCL8 by an equal volume of 50pM SpyCEP and B.
20nM CXCL1 by an equal volume of 100pM SpyCEP. Cleavage reactions were halted at specified
timepoints by the addition of Pefabloc to a final concentration of 2 mg/ml and the remaining
chemokine measured by ELISA. N=6 experimental replicates for each data point, error bars represent
SD of mean values.

Mass spectrometry-derived kinetics of active SpyCEP constructs
344

To directly compare the active full-length and C-terminal SpyCEP constructs and to understand

relative catalytic efficiencies, a mass spectrometry approach to continuously assay the generation of

the 13 amino acid peptide cleaved from the native substrate CXCL8, following incubation with

enzyme, was employed. A range of CXCL8 concentrations (6.25-2000 nM) were incubated with a

fixed concentration of enzyme, either 250 pM for SpyCEP or 40 nM for the C-terminal SpyCEP[245-1613]

and the production of the 13 amino acid peptide monitored over time. 250 pM full length SpyCEP

cleaved CXCL8 to near completion over 30 minutes when substrate concentrations were less than

250 nM; incomplete CXCL8 cleavage was observed when substrate concentration was in excess of

250 nM (Figure EV3A. In contrast, 40 nM of the C-terminal SpyCEP[245-**1613**] cleaved CXCL8 to near

completion over 4 hours when the CXCL8 concentration was 250 nM or less; incomplete CXCL8

cleavage was again observed when substrate concentration were over 250 nM (Figure EV3B). Under

these conditions the C-terminal SpyCEP fragment maintained measurable catalytic activity

thoughout, though with reduced efficacy compared to the full-length construct. A 160-fold increase

in enzyme concentration and additional 3.5 hours reaction time were required to cleave a

comparable amount of CXCL8.

Linear regression analyses of the initial 5 time points, where the rate of cleaved CXCL8 production

was linear, were used to derive Michaelis-Menton plots (Figure 4) and K_M and K_{cat} values for each

construct (Table 2).

Figure 4. Michaelis-Menton analysis of full length and C-terminal SpyCEP construct cleavage of

CXCL8. Data points represent the mean change in cleaved CXCL8 production over time (M s^{-1}) of, **A.**

full-length SpyCEP and **B.** C-terminal SpyCEP). Error bars represent the standard error of the mean of

5 reactions.

Table 2. Kinetic parameters of full length and C-terminal SpyCEP construct activity in cleavage of

CXCL8

K_{cat} and K_M ± SD and K_{cat} / K_M for full length SpyCEP and C-terminal SpyCEP derived from Michaelis-

Menton graphs $Y = V_{max}*X/(K_M + X)$ and K_{cat} equation $Y = ET*k_{cat}*X/(K_M + X)$ where Et = enzyme

concentration, 2.5×10^{-10} for full length SpyCEP and 4×10^{-8} for C-terminal SpyCEP

K_M values for full length SpyCEP and the C-terminal fragment were 53.49 nM and 40.98 nM

respectively, suggesting nanomolar affinity of the SpyCEP constructs for CXCL8. The similarity in the

K_M values between the C- terminal SpyCEP fragment and the full-length enzyme point to a similar

ability to bind substrate. The K_{cat} values (indicative of the number of molecules of cleaved CXCL8

produced per second) however, differed substantially. The full length enzyme cleaved 1.34

molecules of CXCL8 per second compared to the C-terminal construct which cleaved 2.45 molecules

per hour, a 1,970-fold reduction in activity.

Discussion

SpyCEP is a serine protease and a leading virulence factor of *S. pyogenes,* with a narrow range of

substrate specificity, restricted to the family of ELR$^+$ CXC chemokines which modulate neutrophil

mediated immune responses and the newly identified substrate, LL-37 (Biswas et al., 2021).

Autocatalytic processing of SpyCEP results in the generation of two individual fragments that re-

assemble to form an active enzyme (Zingaretti et al., 2010). Here, we describe the enzyme kinetics of

full-length SpyCEP and report the K_M of the enzyme for its natural substrate to be remarkably low,

just 53.49 nM, consistent with high efficiency. Furthermore, we demonstrate that the C-terminal

SpyCEP fragment can catalyse the cleavage of CXCL1 and CXCL8 independent of the N-terminal

fragment. Indeed, when K_M values were compared, they were found to be similar, suggesting that

substrate binding may be confined to the C-terminal domain of SpyCEP. The enzymatic activity of the

C-terminal SpyCEP fragment was, however, markedly reduced compared to full-length SpyCEP. The

N-terminal and N-terminal^{DA} constructs were equally able to restore full catalytic activity of the C-

terminal SpyCEP fragment. Collectively, this suggests that although the aspartate 151 of the N-

terminal fragment may be dispensable for catalysis, the domain itself is important for optimal

enzyme activity.

Serine proteases are ubiquitous and comprise up to one third of all proteolytic enzymes currently

described. They are currently categorised by the MEROPS database (Rawlings et al., 2012) into 13

distinct clans, being differentiated into groups of proteins based on their evolution from the same

common ancestor, with SpyCEP belonging to the S8 family of the SB clan. S8 serine proteases are

typified by a classical catalytic triad composed of serine, histidine and aspartic acid residues that

together contribute to the hydrolysis of a peptide bond within the substrate. It is recognised that a

number of serine protease clans employ a variation on the S8 catalytic triad, utilising instead a triad

of serine, histidine, and glutamic acid, or serine, glutamic acid, and aspartic acid residues. Other

clans utilise catalytic dyads of lysine and histidine or histidine and serine for proteolytic activity. Our

data suggest that SpyCEP activity can reside in a catalytic dyad of histidine and serine, albeit at a

reduced efficacy. It is likely this large gulf in efficiency explains the failure of previous studies to

detect catalytic activity within the isolated C-terminal SpyCEP fragment (Zingaretti et al., 2010).

Within serine proteases there are additional features, beyond the catalytic triad residues, which can

contribute to activity. The oxyanion hole for example, a pocket in the active site composed of

backbone amide NH groups, may provide substrate stabilisation and help drive catalysis (Hedstrom,

2002). Additional residues located in close proximity to the catalytic pocket can also mitigate a loss

of activity resulting from a missing residue, and water also has the potential to moonlight as a

missing functional group (Hedstrom, 2002). These 'stand ins' can provide a possible substitute

machinery to help drive catalytic function. Indeed, some studies have shown that, even with all

three catalytic triad residues removed, serine proteases are still capable of catalysis at rates 1000-

fold greater than the background rate of hydrolysis (Corey and Craik, 1992, Hedstrom, 2002, Paul

and James, 1988).

Mass spectrometry-based kinetics showed that full-length active SpyCEP has a K_M of 53.49 nM and

K_{cat} of 1.34 molecules per second; values which are in agreement with our initial ELISA-based kinetic

assessment (McKenna et al., 2020). In contrast, the C-terminal fragment of SpyCEP has a K_M of 40.98

nM and K_{cat} of 0.00068 molecules per second. These constructs both demonstrated low, nanomolar

K_M values, suggesting a high binding efficiency of SpyCEP for its natural substrate CXCL8, likely

conferred by the C-terminal domain. This is in keeping with the fact that low nanomolar

concentrations of CXCL8 are optimal for neutrophil recruitment (Goldblatt et al., 2019). Although K_M

values are often reported in the μM – mM range, nanomolar K_M values are not without precedent

for other serine proteases; human Kallikrein 6 has a reported K_M of 300 nM and Factor Xa, a

constituent of the prothrombinase complex, has a K_M of 150 nM for prothrombin (Angelo et al.,

2006, Luettgen et al., 2011). Enzyme specificity, a constant which measures the cleavage efficiency

of enzymes, (K_{cat}/K_M), for full length SpyCEP was estimated to be 2.46 x10^7 M^{-1} s^{-1}, a value in the

order of magnitude typical for serine proteases (Hedstrom, 2002). The specificity constant of the C-

terminal fragment was ~1500-fold less, 1.67×10^4 M^{-1} s^{-1}, and K$_{cat}$ ~2000 fold less, a reduction that is

in line with previously reported aspartic acid mutants from a systematic mutational study of the

Bacillus amyloliquefaciens subtilisin catalytic triad (Paul and James, 1988).

The kinetic assays attributed a marked increase in CXCL8 turnover to the additional presence of the

N-terminal fragment. Although we did not specifically evaluate the role of the aspartate residue at

position 151 on the kinetics of SpyCEP activity, this residue did not contribute appreciably to

cleavage of CXCL8 when evaluated using immunoblotting. This raises a question as to whether the N-

terminal fragment confers some additional structural contribution to enzyme activity. Our data

strongly suggest that substrate binding is likely to be attributed to the C-terminal fragment, a finding

consistent with related cell envelope proteinases of *Lactococci* (Siezen, 1999) and the closely related

streptococcal protein, C5a peptidase (Kagawa et al., 2009).

The implications of our findings relating to activity of the C-terminal SpyCEP fragment in *S. pyogenes*

pathogenesis are currently unclear; without the N-terminus, the enzymatic activity detected may be

too low to be of consequence for chemokine cleavage *in vivo,* and in nature both N- and C-terminal

fragments are likely to co-exist. SpyCEP has been the focus of *S. pyogenes* vaccine development

since 2006 (Rodriguez-Ortega et al., 2006), used either in isolation or combination with other

antigenic targets (Bensi et al., 2012, McKenna et al., 2020, Pandey et al., 2016, Reglinski et al., 2016,

Rivera-Hernandez et al., 2016). Vaccine-induced SpyCEP specific antibodies appear not to act

through traditional opsonic means (Rivera-Hernandez et al., 2019) and so likely act through

inhibition of SpyCEP activity. A majority of vaccine preparations evaluated have been based on

'CEP5'; a polypeptide spanning residues 35 – 587 of SpyCEP which contains the N- terminal fragment

and only part of the C terminal fragment (Turner et al., 2009b). Our findings relating to enzyme

function highlight the possibility that antibodies targeting the C-terminal fragment of SpyCEP are

more likely to provide greater neutralizing activity, and potentially improve vaccine efficacy.

Funding

This work was funded by a Biotechnology and Biological Sciences Research Council (BBSRC) NPIF

iCASE studentship (MP) and a Wellcome Trust Collaborative award 215539: 'Understanding and

exploiting Group A streptococcal anti-chemotactic proteases in vaccines for infection' (SS and JP).

Acknowledgements
The authors are grateful for the support of a GSK Golden Triangle Discovery Partnership in Academia

Collaboration (DPAC) that permitted development of the mass spectrometry high throughput

cleavage assay and reagent provision. SS acknowledges the Imperial College London NIHR

Biomedical Research Centre.

Author contributions
MP, CH, AF, DM, JP and SS designed the study. MP, CH, MR, RJE, RAL collected the data. MP and CH

analysed the data. MP, CH, AF, DM, JP and SS interpreted the data. MP collected and prepared the

figures. MP, MR, JP and SS drafted the manuscript. MP, CH, AF, MR, RJE, RAL, DM, JP, SS, revised the

manuscript content.

Conflict of interest

All authors declare there is no conflict of interest.

Expanded View figures

Figure EV1. Western blot of SpyCEP specific LL3-7 cleavage. Immunoblot of 500 ng human LL-37
incubated for 16 hours at 37 °C either alone (lane 7) or with a panel of SpyCEP constructs at a 10:1
molar ratio in favour of LL-37. 4.5 kDa full length LL-37 band and 3.5 kDa cleaved LL-37 band are
indicated by arrows and were detected with 2 µg/ml sheep IgG polyclonal LL-37 antibody and rabbit
anti-sheep IgG antibody (1:40,000). Figure is representative of two experiments.

Figure EV2. Comparative cleavage of CXCL1 and CXCL8 by C-terminal SpyCEP. SDS-page analysis of
A. 5 µM human CXCL1 (all lanes) or B. 5 µM human CXCL8 (all lanes) cleavage by C-terminal SpyCEP
at a 1:5 – 1:5000 molar ratio (SpyCEP: Chemokine). SpyCEP[DASA], C-terminal[SA] and N-terminal controls
were all assayed at the highest 1:5 molar ratio.

Figure EV3. CXCL8 cleavage by active recombinant SpyCEP constructs using mass spectrometry.

A. Production of the 13 amino acid species cleaved from CXCL8 by 250 pM of full length SpyCEP over
30 minutes at room temperature, following incubation with 6.25 nM – 2000 nM CXCL8, 1: 250 –
8000 molar ratio (SpyCEP: CXCL8). **B**. Production of the 13 amino acid species by 40 nM SpyCEP C-
terminal fragment over a 4 hour room temperature incubation, following incubation with 6.25 nM –
2000 nM CXCL8, 1: 0.15 – 50 molar ratio (SpyCEP: CXCL8). N=5 experimental replicates per data
point, error bars represent SD. Figure is representative of two experiments

References

ABATE, F., MALITO, E., FALUGI, F., MARGARIT, Y. R. I. & BOTTOMLEY, M. J. 2013. Cloning, expression, purification, crystallization and preliminary X-ray diffraction analysis of SpyCEP, a candidate antigen for a vaccine against Streptococcus pyogenes. *Acta Crystallogr Sect F Struct Biol Cryst Commun,* 69**,** 1103-6.

ANGELO, P. F., LIMA, A. R., ALVES, F. M., BLABER, S. I., SCARISBRICK, I. A., BLABER, M., JULIANO, L. & JULIANO, M. A. 2006. Substrate specificity of human kallikrein 6: salt and glycosaminoglycan activation effects. *J Biol Chem,* 281**,** 3116-26.

BENSI, G., MORA, M., TUSCANO, G., BIAGINI, M., CHIAROT, E., BOMBACI, M., CAPO, S., FALUGI, F., MANETTI, A. G., DONATO, P., SWENNEN, E., GALLOTTA, M., GARIBALDI, M., PINTO, V., CHIAPPINI, N., MUSSER, J. M., JANULCZYK, R., MARIANI, M., SCARSELLI, M., TELFORD, J. L., GRIFANTINI, R., NORAIS, N., MARGARIT, I. & GRANDI, G. 2012. Multi high-throughput approach for highly selective identification of vaccine candidates: the Group A Streptococcus case. *Mol Cell Proteomics,* 11**,** M111.015693.

BISWAS, D., AMBALAVANAN, P., RAVINS, M., ANAND, A., SHARMA, A., LIM, K. X. Z., TAN, R. Y. M., LIM, H. Y., SOL, A., BACHRACH, G., ANGELI, V. & HANSKI, E. 2021. LL-37-mediated activation of host receptors is critical for defense against group A streptococcal infection. *Cell Rep,* 34**,** 108766.

CARAPETIS, J. R., STEER, A. C., MULHOLLAND, E. K. & WEBER, M. 2005. The global burden of group A streptococcal diseases. *Lancet Infect Dis,* 5**,** 685-94.

COREY, D. R. & CRAIK, C. S. 1992. An investigation into the minimum requirements for peptide hydrolysis by mutation of the catalytic triad of trypsin. *Journal of the American Chemical Society,* 114**,** 1784-1790.

EDWARDS, R. J., TAYLOR, G. W., FERGUSON, M., MURRAY, S., RENDELL, N., WRIGLEY, A., BAI, Z., BOYLE, J., FINNEY, S. J., JONES, A., RUSSELL, H. H., TURNER, C., COHEN, J., FAULKNER, L. & SRISKANDAN, S. 2005. Specific C-terminal cleavage and inactivation of interleukin-8 by invasive disease isolates of Streptococcus pyogenes. *J Infect Dis,* 192**,** 783-90.

EDWARDS, R. J., WRIGLEY, A., BAI, Z., BATEMAN, M., RUSSELL, H., MURRAY, S., LU, H., TAYLOR, G. W., BOOBIS, A. R. & SRISKANDAN, S. 2007. C-terminal antibodies (CTAbs): a simple and broadly applicable approach for the rapid generation of protein-specific antibodies with predefined specificity. *Proteomics,* 7**,** 1364-72.

GOLDBLATT, J., LAWRENSON, R. A., MUIR, L., DATTANI, S., HOFFLAND, A., TSUCHIYA, T., KANEGASAKI, S., SRISKANDAN, S. & PEASE, J. E. 2019. A Requirement for Neutrophil Glycosaminoglycans in Chemokine:Receptor Interactions Is Revealed by the Streptococcal Protease SpyCEP. *J Immunol,* 202**,** 3246-3255.

HEDSTROM, L. 2002. Serine Protease Mechanism and Specificity. *Chemical Reviews,* 102**,** 4501-4524.

JOBICHEN, C., TAN, Y. C., PRABHAKAR, M. T., NAYAK, D., BISWAS, D., PANNU, N. S., HANSKI, E. & SIVARAMAN, J. 2018. Structure of ScpC, a virulence protease from Streptococcus pyogenes, reveals the functional domains and maturation mechanism. *Biochem J,* 475**,** 2847-2860.

KAGAWA, T. F., O'CONNELL, M. R., MOUAT, P., PAOLI, M., O'TOOLE, P. W. & COONEY, J. C. 2009. Model for substrate interactions in C5a peptidase from Streptococcus pyogenes: A 1.9 A crystal structure of the active form of ScpA. *J Mol Biol,* 386**,** 754-72.

KAUR, S. J., NERLICH, A., BERGMANN, S., ROHDE, M., FULDE, M., ZAHNER, D., HANSKI, E., ZINKERNAGEL, A., NIZET, V., CHHATWAL, G. S. & TALAY, S. R. 2010. The CXC chemokine-degrading protease SpyCep of Streptococcus pyogenes promotes its uptake into endothelial cells. *J Biol Chem,* 285**,** 27798-805.

KURUPATI, P., TURNER, C. E., TZIONA, I., LAWRENSON, R. A., ALAM, F. M., NOHADANI, M., STAMP, G. W., ZINKERNAGEL, A. S., NIZET, V., EDWARDS, R. J. & SRISKANDAN, S. 2010. Chemokine-cleaving Streptococcus pyogenes protease SpyCEP is necessary and sufficient for bacterial dissemination within soft tissues and the respiratory tract. *Mol Microbiol,* 76**,** 1387-97.

LUETTGEN, J. M., KNABB, R. M., HE, K., PINTO, D. J. P. & RENDINA, A. R. 2011. Apixaban inhibition of factor Xa: Microscopic rate constants and inhibition mechanism in purified protein systems and in human plasma. *Journal of Enzyme Inhibition and Medicinal Chemistry,* 26**,** 514-526.

LYNSKEY, N. N., GOULDING, D., GIERULA, M., TURNER, C. E., DOUGAN, G., EDWARDS, R. J. & SRISKANDAN, S. 2013. RocA Truncation Underpins Hyper-Encapsulation, Carriage Longevity and Transmissibility of Serotype M18 Group A Streptococci. *PLOS Pathogens,* 9**,** e1003842.

MCKENNA, S., MALITO, E., ROUSE, S. L., ABATE, F., BENSI, G., CHIAROT, E., MICOLI, F., MANCINI, F., GOMES MORIEL, D., GRANDI, G., MOSSAKOWSKA, D., PEARSON, M., XU, Y., PEASE, J., SRISKANDAN, S., MARGARIT, I., BOTTOMLEY, M. J. & MATTHEWS, S. 2020. Structure, dynamics and immunogenicity of a catalytically inactive CXC chemokine-degrading protease SpyCEP from Streptococcus pyogenes. *Computational and Structural Biotechnology Journal,* 18**,** 650-660.

MIDDLETON, J., NEIL, S., WINTLE, J., CLARK-LEWIS, I., MOORE, H., LAM, C., AUER, M., HUB, E. & ROT, A. 1997. Transcytosis and surface presentation of IL-8 by venular endothelial cells. *Cell,* 91**,** 385-95.

PANDEY, M., LANGSHAW, E., HARTAS, J., LAM, A., BATZLOFF, M. R. & GOOD, M. F. 2015. A synthetic M protein peptide synergizes with a CXC chemokine protease to induce vaccine-mediated protection against virulent streptococcal pyoderma and bacteremia. *J Immunol,* 194**,** 5915-25.

PANDEY, M., MORTENSEN, R., CALCUTT, A., POWELL, J., BATZLOFF, M. R., DIETRICH, J. & GOOD, M. F. 2016. Combinatorial Synthetic Peptide Vaccine Strategy Protects against Hypervirulent CovR/S Mutant Streptococci. *J Immunol,* 196**,** 3364-74.

PAUL, C. & JAMES, A. W. 1988. Dissecting the catalytic triad of a serine protease. *Nature,* 332**,** 564.

RALPH, A. P. & CARAPETIS, J. R. 2013. Group a streptococcal diseases and their global burden. *Curr Top Microbiol Immunol,* 368**,** 1-27.

RAWLINGS, N. D., BARRETT, A. J. & BATEMAN, A. 2012. MEROPS: the database of proteolytic enzymes, their substrates and inhibitors. *Nucleic acids research,* 40**,** D343-D350.

REGLINSKI, M., LYNSKEY, N. N., CHOI, Y. J., EDWARDS, R. J. & SRISKANDAN, S. 2016. Development of a multicomponent vaccine for Streptococcus pyogenes based on the antigenic targets of IVIG. *J Infect,* 72**,** 450-9.

RIVERA-HERNANDEZ, T., CARNATHAN, D. G., JONES, S., CORK, A. J., DAVIES, M. R., MOYLE, P. M., TOTH, I., BATZLOFF, M. R., MCCARTHY, J., NIZET, V., GOLDBLATT, D., SILVESTRI, G. & WALKER, M. J. 2019. An Experimental Group A Streptococcus Vaccine That Reduces Pharyngitis and Tonsillitis in a Nonhuman Primate Model. *MBio,* 10.

RIVERA-HERNANDEZ, T., PANDEY, M., HENNINGHAM, A., COLE, J., CHOUDHURY, B., CORK, A.
J., GILLEN, C. M., GHAFFAR, K. A., WEST, N. P., SILVESTRI, G., GOOD, M. F., MOYLE, P.
M., TOTH, I., NIZET, V., BATZLOFF, M. R. & WALKER, M. J. 2016. Differing Efficacies of
Lead Group A Streptococcal Vaccine Candidates and Full-Length M Protein in
Cutaneous and Invasive Disease Models. *MBio,* 7.
RODRIGUEZ-ORTEGA, M. J., NORAIS, N., BENSI, G., LIBERATORI, S., CAPO, S., MORA, M.,
SCARSELLI, M., DORO, F., FERRARI, G., GARAGUSO, I., MAGGI, T., NEUMANN, A.,
COVRE, A., TELFORD, J. L. & GRANDI, G. 2006. Characterization and identification of
vaccine candidate proteins through analysis of the group A Streptococcus surface
proteome. *Nat Biotechnol,* 24**,** 191-7.
SIEZEN, R. J. 1999. Multi-domain, cell-envelope proteinases of lactic acid bacteria. *Antonie*
Van Leeuwenhoek, 76**,** 139-55.
TURNER, C. E., KURUPATI, P., JONES, M. D., EDWARDS, R. J. & SRISKANDAN, S. 2009a.
Emerging role of the interleukin-8 cleaving enzyme SpyCEP in clinical Streptococcus
pyogenes infection. *J Infect Dis,* 200**,** 555-63.
TURNER, C. E., KURUPATI, P., WILES, S., EDWARDS, R. J. & SRISKANDAN, S. 2009b. Impact of
immunization against SpyCEP during invasive disease with two streptococcal species:
Streptococcus pyogenes and Streptococcus equi. *Vaccine,* 27**,** 4923-9.
ZINGARETTI, C., FALUGI, F., NARDI-DEI, V., PIETROCOLA, G., MARIANI, M., LIBERATORI, S.,
GALLOTTA, M., TONTINI, M., TANI, C., SPEZIALE, P., GRANDI, G. & MARGARIT, I.
2010. Streptococcus pyogenes SpyCEP: a chemokine-inactivating protease with
unique structural and biochemical features. *Faseb j,* 24**,** 2839-48.

Figure 1. Cleavage activity of recombinant SpyCEP constructs assayed by immunoblot. Two colour immunoblot showing cleavage of 150 ng CXCL8 incubated for 2 hours at 37 °C either alone (2nd lane) or with panel of SpyCEP constructs at a 1:50 molar ratio (SpyCEP:CXCL8). Green bands represent intact CXCL8 (anti-CXCL8 antibody); red bands represent cleaved CXCL8 (anti-ENWVQ). A 17 kDa molecular weight marker is shown in blue, 8 kDa and 6 kDa molecular weights are highlighted by arrows. Figure is representative of 2 independent immunoblots.

Figure 2. Cleavage activity of SpyCEP and C-terminal SpyCEP[245-1613] constructs using CXCL8 ELISA. SpyCEP constructs were co-incubated with CXCL8. Graphs show residual CXCL8 after a 60-minute room temperature incubation, using full length SpyCEP at a 1:1000 ratio to CXCL8; C-terminal at 1:5 – 1:250 molar ratio to CXCL8; and C-terminal S617A mutant at a 1:5 molar ratio to CXCL8. Reactions were halted at specified timepoints by the addition of Pefabloc to a final concentration of 2 mg/ml. N=6 experimental replicates for each construct, data points show means, error bars represent SD. ns $p > 0.05$, * $p \le 0.05$, **** $p \le 0.0001$, at 60 minutes as determined by ordinary one-way ANOVA.

Figure 3. Cleavage of CXCL8 and CXCL1 by recombinant full length SpyCEP. 30-minute, room temperature cleavage time course of: A. 20nM CXCL8 by an equal volume of 50pM SpyCEP and B. 20nM CXCL1 by an equal volume of 100pM SpyCEP. Cleavage reactions were halted at specified timepoints by the addition of Pefabloc to a final concentration of 2 mg/ml and the remaining chemokine measured by ELISA. N=6 experimental replicates for each data point, error bars represent SD of mean values.

Figure 4. Michaelis-Menton analysis of full length and C-terminal SpyCEP construct cleavage of CXCL8. Data points represent the mean change in cleaved CXCL8 production over time (M s^{-1}) of, **A.** full-length SpyCEP and **B.** C-terminal SpyCEP). Error bars represent the standard error of the mean of 5 reactions.

SpyCEP construct	SpyCEP construct description
SpyCEP	Full-length, SpyCEP$^{34\text{-}1613}$-6His
SpyCEP^{DASA}	Full length double mutant, SpyCEP$^{34\text{-}1613 \ D151A, \ S617A}$-6His
N-terminal (N)	N-terminal fragment, FLAG-TEV-SpyCEP$^{34\text{-}244}$
C-terminal (C)	C-terminal fragment, SpyCEP$^{245\text{-}1613}$-6His
N-terminal^{DA} (N^{DA})	N-terminal mutant fragment, FLAG-TEV-SpyCEP$^{34\text{-}244 \ D151A}$
C-terminal^{SA} (C^{SA})	C-terminal mutant fragment, SpyCEP$^{245\text{-}1613 \ S617A}$-6His
s.pSpyCEP	SpyCEP from *S. pyogenes* supernatant, SpyCEP$^{34\text{-}1613}$-6His

Table 1. SpyCEP constructs used in this study. Constructs were expressed recombinantly in *E. coli* except for s.pSpyCEP that was expressed in, and purified from *S. pyogenes*.

Construct	$K_{cat} \pm$ SD (s^{-1})	$K_M \pm$ SD (nM)	K_{cat}/K_M (M^{-1} s^{-1})
Full-length	1.34 ± 0.61	53.49 ± 10.36	2.46 x10^7
C-terminal	0.00068 ± 0.00003	40.98 ± 7.2	1.67 x10^4

Table 2. Kinetic parameters of full length and C-terminal SpyCEP construct activity in cleavage of CXCL8. K_{cat} and $K_M \pm$ SD and K_{cat} / K_M for full length SpyCEP and C-terminal SpyCEP derived from Michaelis-Menton graphs Y = V_{max}*X/(K_M + X) and K_{cat} equation Y = ET*k_{cat}*X/(K_M + X) where Et = enzyme concentration, 2.5 x10^{-10} for full length SpyCEP and 4 x10^{-8} for C-terminal SpyCEP