Structure-activity studies of *Streptococcus pyogenes* enzyme SpyCEP reveal high affinity

for CXCL8 in the SpyCEP C-terminal

Max Pearson[1,2], Carl Haslam[3], Andrew Fosberry[3], Emma J Jones[3], Mark Reglinski[1], Robert J.

Edwards[4], Richard Ashley Lawrenson[1], Jonathan C Brown[1], Danuta Mossakowska[3,5], James Edward

Pease*[6], Shiranee Sriskandan*[1, 2]

[1]Department of Infectious Disease, Imperial College London, London W12 0NN, UK

[2]Centre for Bacterial Resistance Biology, Imperial College London, London, SW7 2AZ, UK

[3]GlaxoSmithKline R&D, Gunnels Wood Road, Stevenage, Hertfordshire, SG1 2NY, UK

[4]Department of Medicine, Imperial College London, W12 0NN, UK

[5]Malopolska Centre of Biotechnology, Jagiellonian University, 30-387 Kraków, Poland

[6]National Heart and Lung Institute, Imperial College London, London, SW7 2AZ, UK

*Corresponding authors

29 Running title: Enzymatic activity of the streptococcal protease, SpyCEP

30 Character count: 43896

Abstract

The *Streptococcus pyogenes* cell envelope protease (SpyCEP) is vital to streptococcal pathogenesis and disease progression. Despite its strong association with invasive disease, little is known about enzymatic function beyond the ELR[+] CXC chemokine substrate range. As a serine protease, SpyCEP has a catalytic triad consisting of aspartate (D151), histidine (H279), and serine (S617) residues which are all thought to be mandatory for full activity. We utilised a range of SpyCEP constructs to investigate the protein domains and catalytic residues necessary for enzyme function. We designed a high-throughput mass spectrometry assay to measure CXCL8 cleavage and applied this for the first time to study the enzyme kinetics of SpyCEP.

Results revealed a remarkably low Michaelis-Menton constant (K_M) of 82 nM and a turnover of 1.65 molecules per second. We found that an N-terminally-truncated SpyCEP C-terminal construct containing just the catalytic dyad of H279 and S617 was capable of cleaving CXCL8 with a similar K_M of 55 nM, albeit with a reduced substrate turnover of 2.7 molecules per hour, representing a 2,200-fold reduction in activity. We conclude that the SpyCEP C-terminus plays a key role in high affinity substrate recognition and binding, but that the N-terminus is required for full catalytic activity.

Word count: 200

Introduction

Group A Streptococcus (GAS) or *Streptococcus pyogenes* is a leading human pathogen that manifests clinically as a broad spectrum of diseases, ranging from less severe, usually self-limiting infections to life-threatening invasive diseases such as necrotizing fasciitis and toxic shock syndrome. While much of the global health burden of *S. pyogenes* can be attributed to auto-immune sequelae such as rheumatic heart disease, invasive diseases contribute greatly to *S. pyogenes*-associated global mortality. At least 663,000 new cases of invasive infection arise per year, accounting for approximately 163,000 deaths [1] with mortality rates remaining high despite intervention; indeed, 20% of patients with invasive *S. pyogenes* disease die within 7 days of infection onset [2].

Several virulence factors contribute to pathogenesis in invasive *S. pyogenes* disease, chief among which is <u>S</u>treptococcus <u>p</u>yogenes <u>c</u>ell <u>e</u>nvelope <u>p</u>rotease (SpyCEP), an immune-modulatory cell wall-associated protease. SpyCEP is responsible for the rapid and efficient cleavage of a distinct group of CXC chemokines comprising CXCL1, CXCL2, CXCL3, CXCL5, CXCL6, CXCL7, and CXCL8, both locally at the site of an infection, and systemically [3-6]. The ELR$^+$ chemokines, named for their conserved N-terminal glutamate-leucine-arginine motifs, specifically act upon neutrophils eliciting their recruitment and activation. SpyCEP inactivates these chemokines by cleaving the chemokine C-terminal α-helix, releasing a 13 amino acid peptide in the case of CXCL8 [3]. Although chemokine specificity for binding neutrophil chemokine receptors, CXCR1 and CXCR2, is conferred to a large extent by the chemokine N-terminal ELR motif [7], the chemokine C-terminus is necessary for efficient receptor binding and activation [8], in addition to chemokine translocation from tissues to endothelial lumen [7]. As such, SpyCEP cleavage of CXC chemokines results in a reduction of CXCR1 and CXCR2-mediated neutrophil chemotaxis and subsequent paucity of neutrophils at the site of *S. pyogenes* infection [3]. SpyCEP has recently been shown to cleave the human anti-microbial peptide, LL-37 in addition to chemokines [9]. Whilst cleavage does not affect the antimicrobial action of the peptide, it was reported to reduce LL-37 specific neutrophil chemotaxis [9].

SpyCEP is expressed by *S. pyogenes* as a 1647 amino acid, 180 kDa subtilisin-like serine protease, the

crystal structure of which was recently solved to 2.8 Å resolution [10] and further refined to 2.2 Å

resolution [11]. It is a member of the S8 subtilase family, members of which are characterised by a

catalytic triad consisting of an aspartate, histidine, and serine residue, each surrounded by a region

of highly conserved amino acids [12]. SpyCEP is unique among streptococcal proteases in that,

during maturation, it is autocatalytically cleaved between residues Q244 and S245 into 2 distinct

polypeptides, a 30 kDa N-terminal polypeptide and a 150 kDa C-terminal polypeptide. The two

polypeptides harbour the separate residues required for the formation of the catalytic site [6]; the

N-terminal fragment contains the catalytic D151 residue, and the C-terminal fragment contains the

catalytic H279 and S617 residues. Upon cleavage, the two polypeptides re-associate non-covalently

to reconstitute the active enzyme [6]. The recent crystal structures have shed further light upon the

structure of SpyCEP, describing 9 separate domains, the first 5 of which contain the catalytic triad

necessary for enzymatic activity [10].

SpyCEP has been included as a target antigen in several recent *S. pyogenes* vaccine designs due to its

cell surface expression, highly conserved nature, and central role in *S. pyogenes* pathogenesis.

Immunisation with SpyCEP successfully elicits a SpyCEP-specific neutralising antibody response,

providing protection against systemic bacterial dissemination and reducing disease severity in *S.*

pyogenes intramuscular, skin infection models and non-human primate infection models [13-18].

Data that demonstrate vaccine dependence on enzyme inhibition highlight the importance of

understanding the enzymatic activity of SpyCEP and the potential to improve upon vaccine or

inhibitor design [13].

Despite this, little is known about the catalytic properties of the enzyme, except preliminary

structure-function studies [6, 10, 11, 19]. Published data largely focus on the impact of SpyCEP on

streptococcal pathogenesis and disease progression. In this study we generated multiple SpyCEP

constructs to confirm domains necessary for catalytic activity. We then used mass spectrometry to

determine the enzyme kinetics of two SpyCEP constructs for the natural substrate CXCL8.

Methods

Cloning and purification of recombinant SpyCEP constructs in *Escherichia coli*

Codon-optimised SpyCEP gene constructs were expressed in *Escherichia coli* using synthetic gene

sequences (GenScript) from Spy_0416 in the SF370 *S. pyogenes* M1 genome [6, 20] representing full-

length enzyme, SpyCEP[34-1613], the N- terminal polypeptide, SpyCEP[34-244], and C-terminal polypeptide

SpyCEP[245-1613]. Constructs were also generated with alanine substitutions to replace catalytic

residues in the N-terminal fragment (SpyCEP[34-244] D151A), and the C-terminal fragment (SpyCEP[245-

1613] S617A). To enable downstream protein purification, both the full-length enzyme and C-terminal

polypeptides were expressed with a C-terminal 6-histidine tag and the N-terminal polypeptides were

expressed with an N-terminal FLAG tag and TEV linker.

All SpyCEP constructs were cloned into the vector pET-24B and expressed in BL21 (DE3) competent

E. coli, cultured in Terrific Broth medium supplemented with 50 µg/mL kanamycin at 37°C and

shaken at 200 rpm, for 3 hours. The cultures were induced with 0.5 mM Isopropyl β-D-1-

thiogalactopyranoside (IPTG), cooled to 15°C and shaken at 200 rpm for 16 hours before lysis by

sonication on ice. Full-length and C-terminal constructs were purified by Ni-IDA affinity

chromatography (GenScript) as per the manufacturer's instructions. N-terminal constructs were

purified by anti-flag M2 agarose resin chromatography (Sigma-Aldrich) as per manufacturer's

instruction, and further concentrated and purified by SP ion-exchange (Sigma-Aldrich) and Q FF ion-

exchange chromatography (GE Healthcare) as per manufacturer's instruction. All SpyCEP constructs

were subsequently concentrated and purified by size exclusion chromatography on a HiLoad

Superdex 200 or 75 prep grade column (GE Healthcare) dependent on molecular weight.

The resultant panel of recombinantly expressed SpyCEP constructs is shown in Table 1. All of the

resultant protein preparations were found to be >80% pure as determined following analysis by SDS-

PAGE. The re-association of the SpyCEP N and C-termini to create full-length constructs was carried

out by equimolar co-incubation at 37°C for 30 minutes in 40 mM Tris-HCl pH 7.5, 0.1 mM CHAPS, 1

mM DTT, 0.1% BSA, 75 mM NaCl.

SpyCEP construct	SpyCEP construct description
SpyCEP	Full-length, SpyCEP$^{34-1613}$-6His
SpyCEP$^{D151A, S617A}$	Full-length double mutant, SpyCEP$^{34-1613\ D151A,\ S617A}$-6His
N-terminal	N-terminal fragment, FLAG-TEV-SpyCEP$^{34-244}$
C-terminal	C-terminal fragment, SpyCEP$^{245-1613}$-6His
N-terminal^{D151A}	N-terminal mutant fragment, FLAG-TEV-SpyCEP$^{34-244\ D151A}$
C-terminal^{S617A}	C-terminal mutant fragment, SpyCEP$^{245-1613\ S617A}$-6His
s.pSpyCEP	SpyCEP from *S. pyogenes* supernatant, SpyCEP$^{34-1613}$-6His

142 **Table 1. SpyCEP constructs used in this study.** Constructs were expressed recombinantly in *E. coli*
143 except for s.pSpyCEP that was expressed in, and purified from *S. pyogenes*.

145 **Expression of histidine-tagged SpyCEP in *S. pyogenes***
146

To permit release of soluble His-tagged SpyCEP by *S. pyogenes*, the C-terminal anchor domain of

SpyCEP was replaced by a stop codon in *S. pyogenes* strain H292, an *emm*81 strain that makes

abundant SpyCEP [5]. A 549 base pair region immediately upstream of the SpyCEP cell wall anchor

motif was amplified from *S. pyogenes* H292 genomic DNA using the primers

5'GGGAATTCTGTTGTCAGGTAACAGTCTTATCTTGCC -3' and

5'CCGAATTCACAACACTAGGCTTTTGCTGAGGTCGTTG -3'. *EcoRI* restriction sites were incorporated at

the terminal ends of the primer sequences. The amplified DNA was cloned into the homologous

recombination plasmid pUCMUT to produce the vector pUCMUT$_{CEP}$ which was transformed into One

Shot TOP10 Chemically Competent *E. coli* (Thermo Fisher Scientific) according to the manufacturer's

instructions. The nucleotide sequence of the SpyCEP C-terminus was subsequently further modified

to encode a hexa-histidine tag by inverse PCR using pUCMUT$_{CEP}$ as the template and the primers 5'-

TATCCTAGGTAGTGTTGTGAATTCGTAATCATGGTCATAG-3' and 5'-

TATCCTAGGATGATGATGATGATGATGGGCTTTTGCTGAGGTCGTTG-3'. The amplification was

performed using GoTaq Long PCR Master Mix (Promega) according to the manufacturer's

instructions. *AvrII* restriction sites, incorporated at the terminal ends of the primer sequences,

facilitated re-circulation of the amplified plasmid. The presence of a 6-His sequence in the modified

pUCMUT$_{CEP}$ construct (denoted pUCMUT$_{CEP-HIS}$) was confirmed by Sanger sequencing using the

pUCMUT sequencing primers 5'-GACAGCAACATCTTTGTGAAAGATGG-3' and 5'-

CATTAATGCAGCTGGCACGAC-3'. The pUCMUT$_{CEP-HIS}$ construct was introduced into H292 by

electroporation and crossed into the chromosome by homologous recombination as previously

described [21] to generate strain H1317. Secretion of 6-His-tagged SpyCEP into the culture

supernatant of H1317 was confirmed by western blotting (data not shown). To purify SpyCEP, *S.*

pyogenes H1317 was grown in Todd-Hewitt broth (Oxoid) overnight at 37°C with 5% CO_2 The culture

was pelleted at 2500 xg for 10 minutes and the supernatant sterilised using Amicon 0.22 μm filters,

then concentrated using 15 ml Amicon 100 kDa centrifuge filter columns and purified by nickel

column affinity chromatography (Novagen His-Bind Resin) as per the manufacturer's instructions.

Sodium dodecyl sulphate–polyacrylamide gel electrophoresis (SDS-page)

To visualise chemokine cleavage, 50 pmol of recombinant human CXCL1 or CXCL8 (R&D Systems)

was incubated with C-terminal SpyCEP in a final volume of 20 μl, at a molar ratio of 1:5, 50, 500,

5000 in favour of chemokine for 2 hours at 37^{O}C. C-terminal^{S617A} and N-terminal constructs$^{34-244}$ were

included as controls and assayed at the highest 1:5 molar ratio. Reactions were halted by the

addition of Dithiothreitol (DTT) at a final concentration of 100 mM, 4X Bolt LDS sample buffer and

heating to 70°C for 10 minutes. Samples were separated on pre-cast 4-12% MES buffered Bolt Bis-

Tris gradient gels (Invitrogen) by SDS-PAGE gel electrophoresis at 165 V for 35 minutes with SeeBlue

Plus 2 (Invitrogen) used for molecular weight ladder. Gels were stained with PageBlue protein

staining solution (Thermo Fisher Scientific) overnight and de-stained in deionised water.

Multiplex fluorescent western blotting

To visualise generation of both intact CXCL8 and the larger (N-terminal) CXCL8 cleavage product, a

rabbit antiserum was raised against the CXCL8 neo-epitope (anti-ENWVQ) that is exposed following

SpyCEP cleavage of CXCL8 [22]. SpyCEP constructs were incubated at 37 °C with 18.75 pmol of

recombinant human CXCL8 in a final volume of 20 μl, at a 1:50 molar ratio in favour of CXCL8.

Digests were halted and separated by SDS-PAGE gel electrophoresis as described above. Proteins

were transferred by iBlot2 (Thermo Fisher Scientific) onto 0.2 μm nitrocellulose membranes as per

manufacturer's instructions. Membranes were subsequently blocked for 1 hour at room

temperature in blocking buffer (PBS with 5% skimmed milk powder (Sigma-Aldrich) and 0.1%

Tween), then blotted overnight at 4 °C with 2 primary antibodies, 1 μg/ml mouse anti-human CXCL8

(R&D Systems) and 1:1000 rabbit anti-ENWVQ. Membranes were washed in wash buffer (PBS with

0.05% Tween) and incubated with 1:7500 goat anti-rabbit IgG A680nm and 1:7500 goat anti-mouse

IgG A790nm for 1 hour at room temperature before being visualised on LiCor Odyssey Fc

(Invitrogen).

SpyCEP activity against LL-37 was assessed by a 16 hour, 37 °C incubation of 111.1 pmol human LL-37

(R&D Systems) with SpyCEP constructs at a 1:10 molar ratio in favour of LL-37 in a final volume of 20

μl. Reactions were stopped, separated and blotted onto 0.2 μm nitrocellulose as above and

incubated overnight at 4 °C in blocking buffer supplemented with 2 μg/ml polyclonal sheep IgG anti-

LL-37 (R&D Systems). Membranes were washed in wash buffer (PBS with 0.05% Tween) and

incubated in blocking solution with rabbit anti-sheep IgG (Abcam) at 1:40,000 dilution for 1 hour at

room temperature and visualised on LiCor Odyssey Fc (Invitrogen).

Enzyme linked immunosorbent assay (ELISA) analysis of chemokine cleavage

Catalytic activities of SpyCEP constructs were measured through detection of remaining intact CXCL8

or CXCL1 substrate following incubation with SpyCEP by ELISA (R&D Systems human CXCL8 and

CXCL1 DuoSet ELISA) as per manufacturer's instructions. For cleavage time courses of CXCL8 by C-

terminal SpyCEP, 5 pmol CXCL8 were incubated with C-terminal constructs at a range of enzyme:

chemokine molar ratios, 1:5 – 1:250, in a final volume of 100 μl at room temperature for 60 minutes.

Full-length SpyCEP and C-terminal [S617A] were included as controls at a molar ratio of 1:1000 and 1:5,

respectively. To compare SpyCEP cleavage rates of CXCL8 and CXCL1, 5 fmol or 10 fmol SpyCEP were

incubated with 2 pmol human CXCL8 and CXCL1 respectively (R&D Systems), in a final volume of 100

μl and incubated at room temperature for 30 minutes. All reactions were halted at defined

timepoints with the addition of concentration of Pefabloc (Sigma-Aldrich) to a final concentration of

2 mg/ml (8.34 mM).

Linear regression analyses of the initial five time points of CXCL1 and CXCL8 cleavage (0, 1, 2, 3 and 4

minutes) were utilised to determine the maximal rate of SpyCEP activity.

Mass spectrometry analysis of CXCL8 cleavage

Analysis of CXCL8 cleavage was assayed on a SCIEX API6500 triple quadrupole electrospray mass

spectrometer coupled to a high-throughput robotic sample preparation and injection system,

RapidFire200 (Agilent Technologies). CXCL8 substrate, the 13 amino acid CXCL8 SpyCEP cleavage

product RVVEKFLKRAENS, and a heavy atom substituted internal standard of CXCL8 SpyCEP cleavage

product RV [$U_{13}C_5$ $_{15}N$-VAL]-EKF-[$U_{13}C_6$ $_{15}N$-Leu]- KRAENS) were monitored by mass spectrometry.

The mass spectrometer was operated in positive electrospray MRM mode, and transitions (Q1/Q3)

for each species were optimised to give m/z as follows: CXCL8, 1048.7/615.3, CXCL8 cleavage

product 526.2/211.1, internal standard 530.5/211.1. A dwell time of 50 ms was used for the MRM

transitions. The mass spectrometer was operated with a spray voltage of 5500 V and at a source

temperature of 650 °C.

To assay CXCL8 cleavage dynamically, chemokine and SpyCEP constructs were loaded into a 384 well

plate to the following final concentrations: 6.25-2000 nM chemokine, 250 pM SpyCEP or 40 nM C-

terminal SpyCEP in a 20 μl reaction volume. Reactions were carried out at 21°C and stopped at

desired time points, 0-240 minutes, by the addition of stop solution (1% formic acid) supplemented

with 1 µM heavy atom substituted CXCL8 internal standard and centrifuged 2000 xg for 10 minutes.

Assay plates were transferred onto the RapidFire200 integrated to the API6500 mass spectrometer.

Samples were aspirated under vacuum directly from 384-well assay plates for 0.6 s. The samples

were then loaded onto a C18 solid-phase extraction cartridge to remove non-volatile buffer salts,

using HPLC -grade water supplemented with 0.1% (v/v) formic acid at a flow rate of 1.5 mL/min for 4

s. The retained analytes were eluted to the mass spectrometer by washing the cartridge with

acetonitrile HPLC-grade water (8:2, v/v) with 0.1% (v/v) formic acid at 1.25 mL/min for 4 s. The

cartridge was re-equilibrated with HPLC-grade water supplemented with 0.1% (v/v) formic acid for

0.6 s at 1.5 mL/min.

Results were expressed as a ratio of CXCL8 cleavage product (526.2/211.1) intensity areas compared

with the internal standard (530.5/211.1) instensity areas. The results were interpolated from a

standard curve constructed using known amounts of cleaved CXCL8 over the range of 25 – 2000 nM

where there was a linear relationship (R^2=0.99) and with a coefficient of variation < 9% for all

concentrations. Where interpolated values were below 25 nM, the assay LOQ, values were assigned

an arbitrary value of LOQ/2, 12.5 nM.

Kinetic analysis

Linear regression analyses from five time points of full-length SpyCEP (1, 2, 3, 4, 5 minutes) and C-

terminal SpyCEP (15, 30, 45, 60, 75 minutes) CXCL8 reactions were plotted against substrate

concentration and kinetics derived from the Michaelis-Menton equation $Y = V_{max}*X/ (K_M + X)$ and the

K_{cat} equation $Y = ET*k_{cat}*X/(K_M + X)$ where Et = enzyme concentration as fitted by GraphPad Prism v

8.0.2.

Results
259

260 *Screening of CXCL8 cleavage activity using fluorescent western blotting*
261

To initially assess the activity of recombinant SpyCEP constructs we screened constructs for specific

CXCL8 cleavage activity using two-colour multiplex western blotting. Human CXCL8 was incubated

for 2 hours at 37 °C with the different SpyCEP constructs, and the reaction products were separated

by SDS-PAGE, then immunoblotted using separate antibodies that detect either intact or cleaved

CXCL8. Detection of full-length CXCL8 (8 kDa, green bands) or the CXCL8 neo epitope (ENWVQ),

exposed after SpyCEP cleavage, (6 kDa, red bands) was evident with this system (Figure 1).

Recombinant full-length SpyCEP expressed in *E. coli* successfully cleaved CXCL8 to completion, with

no difference observed between CXCL8 cleavage with *S. pyogenes* SpyCEP and *E. coli* SpyCEP (Figure

1). As expected, no CXCL8 cleavage was observed when using the catalytically inactive mutant,

SpyCEP[D151A, S617A]. As has been previously reported [6], the SpyCEP N- and C-termini, when

independently expressed and purified, can be re-associated to form an active enzyme R.E SpyCEP,

which successfully cleaved CXCL8. The N-terminal fragment of SpyCEP alone could not cleave CXCL8.

Unexpectedly, however, the C-terminal fragment of SpyCEP was observed to cleave CXCL8, albeit

not to completion. This independent catalytic activity was negated by mutation of the catalytic S617

to alanine (C-terminal[S617A] construct, Figure 1). Cleavage of CXCL8 by the SpyCEP C-terminal fragment

was enhanced when re-associated with the catalytically inert N-terminal mutant (N-terminal[D151A]

construct); indeed, activity was equivalent to that observed with the re-associated enzyme under

these conditions, with cleavage of CXCL8 to near completion. However, the N-terminal[D151A] construct

was unable to restore catalytic activity to the C-terminal[S617A] construct when the two were

combined.

The results demonstrated for the first time that the SpyCEP C-terminal fragment alone is sufficient

for catalytic cleavage of CXCL8 in this assay system. The data also established that, although the

presence of the SpyCEP N-terminal fragment enhanced enzymatic activity, the N-terminal residue

D151 was dispensable for enzymatic activity. We further sought to examine whether the C-terminal

displayed activity against the newly described substrate, LL-37 [9]. Western blot analysis confirmed

cleavage of LL-37 by full-length recombinant SpyCEP, demonstrated by a reduction in band size,

however the C-terminal fragment was unable to cleave LL-37 despite a high molar ratio of enzyme to

LL-37 (1: 10) and a prolonged 16-hour incubation at 37 °C (Figure S1).

Figure 1. Cleavage activity of recombinant SpyCEP constructs assayed by immunoblot. Two colour
immunoblot showing cleavage of 18.75 pmol CXCL8 when incubated for 2 hours at 37 °C either alone
(2nd lane) or with a panel of SpyCEP constructs at a 1:50 molar ratio (SpyCEP: CXCL8). Green bands
represent intact CXCL8 (anti-CXCL8 antibody); red bands represent cleaved CXCL8 (anti- ENWVQ). A
17 kDa molecular weight marker is shown in blue, 8 kDa and 6 kDa molecular weights are highlighted
by arrows. Figure is representative of 2 independent immunoblots.

Differential cleavage of CXCL8 and CXCL1 by full-length and C-terminal SpyCEP
300

To determine whether the catalytic activity of the C-terminal SpyCEP fragment was reproducible

over a shorter incubation period, we assessed CXCL8 cleavage by ELISA, where CXCL8 cleavage is

detected through a reduction in substrate concentration. SpyCEP C-terminal constructs were

incubated with CXCL8 at molar ratios ranging from 1:5 – 1:250 (enzyme: chemokine) over a 60-

minute time course at room temperature. Full-length recombinant SpyCEP and the inactive C-

terminal fragment, C-terminal[S617A], were included as controls at a molar ratio of 1:1000 and 1:5,

respectively. Under these conditions, near complete CXCL8 cleavage was observed for full-length

SpyCEP and a dose-dependent increase in catalytic activity was observed for the C-terminal fragment

(Figure 2). After 5 minutes incubation full-length SpyCEP cleaved over 50% of the starting CXCL8

input, with only 6.25% of CXCL8 remaining after 1 hour. At the highest concentration of C-terminal

SpyCEP tested, a 1:5 molar ratio, the C-terminal fragment alone cleaved 9% of the starting CXCL8 by

5 minutes, 25% by 30 minutes and 42% by 60 minutes. Indeed, at the lowest concentration tested, a

1:250 molar ratio, the C-terminal of SpyCEP cleaved 9% of CXCL8 input by 1 hour. As demonstrated

by immunofluorescent western blotting, the serine residue at position 617 was vital for SpyCEP

catalytic function, as no CXCL8 cleavage was observed using the C-terminal[S617A] construct, even when

employed at the highest enzyme: chemokine ratio. After a 1-hour incubation, full-length SpyCEP and

C-terminal constructs (assayed using a molar enzyme: chemokine ratio of 1:5, 1:25 and 1:50) cleaved

significantly more CXCL8 compared to the C-terminal[S617A] construct.

Figure 2. Cleavage activity of SpyCEP and C-terminal SpyCEP[245-1613] constructs using CXCL8 ELISA.
SpyCEP constructs were co-incubated with 5 pmol CXCL8. Graphs show residual CXCL8 after a 60-minute room temperature incubation, using full-length SpyCEP at a 1:1000 ratio to CXCL8; the C-terminal SpyCEP construct at a 1:5 – 1:250 molar ratio to CXCL8; and the C-terminal[S617A] mutant at a 1:5 molar ratio to CXCL8. Reactions were halted at specified timepoints by the addition of Pefabloc to a final concentration of 2 mg/ml. N=6 experimental replicates for each construct, data points show means, error bars represent SD. ns $p > 0.05$, * $p \leq 0.05$, **** $p \leq 0.0001$, at 60 minutes as determined by ordinary one-way ANOVA.

SDS-PAGE analysis of the catalytic activity of the C-terminal SpyCEP construct additionally showed

that C-terminal activity was not restricted to the CXCL8 substrate. Over 2 hours at 37 °C using a 1:5

molar ratio of enzyme: substrate, the SpyCEP C-terminal construct was capable of cleaving human

CXCL1 to near completion (Figure S2).

To further assess the activity of SpyCEP and to interrogate reaction rates against separate

chemokines, full-length SpyCEP was incubated with CXCL1 and CXCL8 and the remaining chemokine

levels determined by ELISA. Human CXCL1 or human CXCL8 were incubated with SpyCEP at room

temperature over a 30-minute time course, at 1:200 or 1:400 molar ratios respectively (enzyme:

chemokine). 5 fmol of SpyCEP rapidly and efficiently cleaved 2 pmol CXCL8, with ~ 15% of the

chemokine input remaining after 10 minutes of incubation (Figure 3A). This contrasted with CXCL1

cleavage, that required 10 fmol SpyCEP to cleave just 25% of the chemokine input over the same 10-

minute period. Indeed, by 30 minutes SpyCEP had yet to cleave half of the starting amount of CXCL1

(Figure 3B). Utilising a linear regression of the initial 5 timepoints, where SpyCEP activity is maximal,

we found that 5 fmol of SpyCEP was able to cleave 284 fmol of CXCL8 per minute, and 10 fmol of

SpyCEP was capable of cleaving 62 fmol of CXCL1 per minute. These data confirmed the activity of

recombinant SpyCEP and highlighted differential cleavage efficiency across the CXC substrate range

– a feature which has been previously recognised but not quantified [8].

Figure 3. Cleavage of CXCL8 and CXCL1 by recombinant full-length SpyCEP. 30-minute, room
temperature cleavage time course of: A. 2 pmol CXCL8 with 5 fmol SpyCEP and B. 2 pmol CXCL1 with
10 fmol SpyCEP. Cleavage reactions were halted at specified timepoints by the addition of Pefabloc
to a final concentration of 2 mg/ml and the remaining chemokine measured by ELISA. N=6
experimental replicates for each data point, error bars represent SD of mean values.

Mass spectrometry-derived kinetics of active SpyCEP constructs
365

To directly compare the active full-length and C-terminal SpyCEP constructs and to understand

relative catalytic efficiencies, a mass spectrometry approach to continuously assay the generation of

the 13 amino acid peptide cleaved from the native substrate CXCL8, following incubation with

enzyme, was employed. A range of CXCL8 concentrations (25-2000 nM) were incubated with a fixed

concentration of enzyme, either 250 pM of full length SpyCEP, or 40 nM of the C-terminal SpyCEP[245-

1613] and the production of the 13 amino acid peptide was monitored over time. 250 pM full-length

SpyCEP cleaved CXCL8 to near completion over 30 minutes when substrate concentrations were less

than 250 nM; incomplete CXCL8 cleavage was observed when substrate concentration was in excess

of 250 nM (Figure S3A). In contrast, 40 nM of the C-terminal SpyCEP[245-1613] cleaved CXCL8 to near

completion over 4 hours when the CXCL8 concentration was 250 nM or less; incomplete CXCL8

cleavage was again observed when substrate concentrations were over 250 nM (Figure S3B). Under

these conditions the C-terminal SpyCEP fragment maintained measurable catalytic activity

throughout, though with reduced efficacy compared to the full-length construct. A 160-fold increase

in enzyme concentration and additional 3.5 hours reaction time were required to cleave a

comparable amount of CXCL8.

Linear regression analyses of 5 time points, where the rate of cleaved CXCL8 production was linear,

were used to derive Michaelis-Menton plots (Figure 4) and K_M and K_{cat} values for each construct

(Table 2).

Figure 4. Kinetic activity of full-length and C-terminal SpyCEP construct cleavage of CXCL8. Data
points represent the mean change in cleaved CXCL8 production over time (M s^{-1}) of, **A.** Full-length
SpyCEP and **B.** C-terminal SpyCEP. Error bars represent the standard error of the mean of 5
reactions.

Construct	K_{cat} ± SEM (s^{-1})	K_M ± SEM (nM)	V_{max} ± SEM (M s^{-1})	K_{cat}/K_M (M^{-1} s^{-1})
Full-length	1.65 ± 0.13	82 ± 26	4.14×10^{-10} ± 3.35×10^{-11}	2.02×10^7
C-terminal	0.00075 ± 0.00005	55 ± 17	3.00×10^{-11} ± 2.17×10^{-12}	1.36×10^4

Table 2. Kinetic parameters of full-length and C-terminal SpyCEP construct activity in cleavage of

CXCL8

K_{cat}, K_M and V_{max} ± SEM and K_{cat} / K_M for full-length SpyCEP and C-terminal SpyCEP derived from

Michaelis-Menton graphs Y = V_{max}*X/ (K_M + X) and K_{cat} equation Y = ET*k_{cat}*X/(K_M + X) where Et =

enzyme concentration, 2.5×10^{-10} M for full-length SpyCEP and 4×10^{-8} M for C-terminal SpyCEP.

K_M values for full-length SpyCEP and the C-terminal fragment were 82 nM and 55 nM respectively,

suggesting nanomolar affinity of the SpyCEP constructs for CXCL8. The similarity in the K_M values

between the C- terminal SpyCEP fragment and the full-length enzyme point to a similar ability to

bind substrate. The K_{cat} values (indicative of the number of molecules of cleaved CXCL8 produced per

second) however, differed substantially. The full-length enzyme cleaved 1.65 molecules of CXCL8 per

second compared to the C-terminal construct which cleaved 2.7 molecules per hour, a 2,200-fold

reduction in activity.

Discussion

SpyCEP is a serine protease and a leading virulence factor of *S. pyogenes,* with a narrow range of substrate specificity, restricted to the family of ELR[+] CXC chemokines which modulate neutrophil mediated immune responses [3, 4, 8] and LL-37 [9]. Autocatalytic processing of SpyCEP results in the generation of two individual fragments that re-assemble to form an active enzyme [6]. Here, we describe the enzyme kinetics of full-length SpyCEP and report the K_M of the enzyme for its natural substrate to be remarkably low, just 81.76 nM, consistent with high efficiency. Furthermore, we demonstrate that the C-terminal SpyCEP fragment can catalyse the cleavage of CXCL1 and CXCL8 independent of the N-terminal fragment. Indeed, when K_M values were compared, they were found to be similar, suggesting that substrate binding may be confined to the C-terminal domain of SpyCEP. The enzymatic activity of the C-terminal SpyCEP fragment was, however, markedly reduced compared to full-length SpyCEP. The N-terminal and N-terminal[D151A] constructs were equally able to restore full catalytic activity of the C-terminal SpyCEP fragment. Collectively, this suggests that although the aspartate 151 of the N-terminal fragment may be dispensable for catalysis, the domain itself is important for optimal enzyme activity.

Serine proteases are ubiquitous and comprise up to one third of all proteolytic enzymes currently described. They are currently categorised by the MEROPS database [23] into 13 distinct clans, being differentiated into groups of proteins based on their evolution from the same common ancestor, with SpyCEP belonging to the S8 family of the SB clan. S8 serine proteases are typified by a classical catalytic triad composed of serine, histidine and aspartic acid residues that together contribute to the hydrolysis of a peptide bond within the substrate. It is recognised that a number of serine protease clans employ a variation on the S8 catalytic triad, utilising instead a triad of serine, histidine, and glutamic acid, or serine, glutamic acid, and aspartic acid residues. Other clans utilise catalytic dyads of lysine and histidine or histidine and serine for proteolytic activity. Our data suggest that SpyCEP activity can reside in a catalytic dyad of histidine and serine, albeit at a greatly reduced

efficacy. It is likely this large gulf in efficiency explains the failure of previous studies to detect

catalytic activity within the isolated C-terminal SpyCEP fragment [6].

Within serine proteases there are additional features, beyond the catalytic triad residues, which can

contribute to activity. The oxyanion hole for example, a pocket in the active site composed of

backbone amide NH groups, may provide substrate stabilisation and help drive catalysis [24].

Additional residues located in close proximity to the catalytic pocket can also mitigate a loss of

activity resulting from a missing residue, and water also has the potential to moonlight as a missing

functional group [24]. These 'stand ins' can provide a possible substitute machinery to help drive

catalytic function. Indeed, some studies have shown that, even with all three catalytic triad residues

removed, serine proteases are still capable of catalysis at rates 1000-fold greater than the

background rate of hydrolysis [24-26].

Mass spectrometry-based kinetics showed that full-length active SpyCEP has a K_M of 82 nM and K_{cat}

of 1.65 molecules per second; values which are in agreement with our initial ELISA-based kinetic

assessment [11]. In contrast, the C-terminal fragment of SpyCEP has a K_M of 55 nM and K_{cat} of

0.00075 molecules per second. These constructs both demonstrated low, nanomolar K_M values,

suggesting a high binding efficiency of SpyCEP for its natural substrate CXCL8, likely conferred by the

C-terminal domain. This is in keeping with the fact that low nanomolar concentrations of CXCL8 are

optimal for neutrophil recruitment [8]. Although K_M values are often reported in the $\mu M - mM$

range, nanomolar K_M values are not without precedent for other serine proteases; human kallikrein

6 has a reported K_M of 300 nM and Factor Xa, a constituent of the prothrombinase complex, has a K_M

of 150 nM for prothrombin [27, 28]. Enzyme specificity, a constant which measures the cleavage

efficiency of enzymes, (K_{cat}/K_M), for full-length SpyCEP was estimated to be 2.02 x10^7 M^{-1} s^{-1}, a

value in the order of magnitude typical for serine proteases [24]. The specificity constant of the C-

terminal fragment was ~1500-fold less, 1.36 x10^4 M^{-1} s^{-1}, and K_{cat} 2200 fold less, a reduction that is in

line with previously reported aspartic acid mutants from a systematic mutational study of the

Bacillus amyloliquefaciens subtilisin catalytic triad [26]. The development of an MS-based assay of

SpyCEP activity, that is not dependent on western blotting or ELISA, provides potential for future

high-throughput analysis of SpyCEP activity and detection of SpyCEP inhibitors.

The kinetic assays attributed a marked increase in CXCL8 turnover to the additional presence of the

N-terminal fragment. Although we did not specifically evaluate the role of the aspartate residue at

position 151 on the kinetics of SpyCEP activity, this residue did not contribute appreciably to

cleavage of CXCL8 when evaluated using immunoblotting. This raises a question as to whether the N-

terminal fragment confers some additional structural contribution to enzyme activity. Our data

strongly suggest that substrate binding is likely to be attributed to the C-terminal fragment, a finding

consistent with related cell envelope proteinases of *Lactococci* [12] and the closely related

streptococcal protein, C5a peptidase [29].

The implications of our findings relating to activity of the C-terminal SpyCEP fragment in *S. pyogenes*

pathogenesis are currently unclear; without the N-terminus, the enzymatic activity detected may be

too low to be of consequence for chemokine cleavage *in vivo,* and in nature both N- and C-terminal

fragments are likely to co-exist. SpyCEP has been a focus of *S. pyogenes* vaccine development in

recent years [30], used either in isolation or combination with other antigenic targets [11, 14, 16, 17,

31]. Vaccine-induced SpyCEP specific antibodies appear not to act through traditional opsonic means

[13] and so may act through inhibition of SpyCEP activity. Many vaccine preparations evaluated have

been based on 'CEP5'; a polypeptide spanning residues 35 – 587 of SpyCEP which contains the N-

terminal fragment and only part of the C terminal fragment [18]. Our findings relating to enzyme

function suggest that antibodies targeting the C-terminal fragment of SpyCEP are more likely to

provide greater neutralizing activity, and potentially improve vaccine efficacy.

Funding

This work was funded by a Biotechnology and Biological Sciences Research Council (BBSRC) NPIF

iCASE studentship and a Wellcome Trust Collaborative award 215539: 'Understanding and exploiting

Group A streptococcal anti-chemotactic proteases in vaccines for infection'.

Acknowledgements

The authors are grateful for the support of a GSK Golden Triangle Discovery Partnership in Academia

Collaboration (DPAC) that permitted development of the mass spectrometry high throughput

cleavage assay and reagent provision. SS acknowledges the Imperial College London NIHR

Biomedical Research Centre.

Data availability

No new datasets were generated in this study other than those reported in the manuscript.

Author contributions

MP, CH, AF, DM, JP and SS designed the study. MP, CH, MR, RJE, RAL and JB collected the data. MP

and CH analysed the data. MP, CH, AF, DM, JP and SS interpreted the data. MP collected and

prepared the figures. MP, MR, JP and SS drafted the manuscript. MP, CH, AF, MR, RJE, RAL, DM, JP,

SS, revised the manuscript content.

Conflict of interest

Co-authors Carl Haslam, Andrew Fosberry, Emma J Jones and Danuta Mossakowska were employed

by GlaxoSmithKline R&D at the time this work was conducted. All other authors declare no conflict of

interest.

References

1. Carapetis, J.R., et al., *The global burden of group A streptococcal diseases.* Lancet Infect Dis, 2005. **5**(11): p. 685-94.
2. Ralph, A.P. and J.R. Carapetis, *Group A streptococcal diseases and their global burden.* Curr Top Microbiol Immunol, 2013. **368**: p. 1-27.
3. Edwards, R.J., et al., *Specific C-terminal cleavage and inactivation of interleukin-8 by invasive disease isolates of Streptococcus pyogenes.* J Infect Dis, 2005. **192**(5): p. 783-90.
4. Kurupati, P., et al., *Chemokine-cleaving Streptococcus pyogenes protease SpyCEP is necessary and sufficient for bacterial dissemination within soft tissues and the respiratory tract.* Mol Microbiol, 2010. **76**(6): p. 1387-97.
5. Turner, C.E., et al., *Emerging role of the interleukin-8 cleaving enzyme SpyCEP in clinical Streptococcus pyogenes infection.* J Infect Dis, 2009. **200**(4): p. 555-63.
6. Zingaretti, C., et al., *Streptococcus pyogenes SpyCEP: a chemokine-inactivating protease with unique structural and biochemical features.* FASEB J, 2010. **24**(8): p. 2839-48.
7. Middleton, J., et al., *Transcytosis and surface presentation of IL-8 by venular endothelial cells.* Cell, 1997. **91**(3): p. 385-95.
8. Goldblatt, J., et al., *A Requirement for Neutrophil Glycosaminoglycans in Chemokine:Receptor Interactions Is Revealed by the Streptococcal Protease SpyCEP.* J Immunol, 2019. **202**(11): p. 3246-3255.
9. Biswas, D., et al., *LL-37-mediated activation of host receptors is critical for defense against group A streptococcal infection.* Cell Rep, 2021. **34**(9): p. 108766.
10. Jobichen, C., et al., *Structure of ScpC, a virulence protease from Streptococcus pyogenes, reveals the functional domains and maturation mechanism.* Biochem J, 2018. **475**(17): p. 2847-2860.
11. McKenna, S., et al., *Structure, dynamics and immunogenicity of a catalytically inactive CXC chemokine-degrading protease SpyCEP from Streptococcus pyogenes.* Computational and Structural Biotechnology Journal, 2020. **18**: p. 650-660.
12. Siezen, R.J., *Multi-domain, cell-envelope proteinases of lactic acid bacteria.* Antonie Van Leeuwenhoek, 1999. **76**(1-4): p. 139-55.
13. Rivera-Hernandez, T., et al., *An Experimental Group A Streptococcus Vaccine That Reduces Pharyngitis and Tonsillitis in a Nonhuman Primate Model.* MBio, 2019. **10**(2).
14. Bensi, G., et al., *Multi high-throughput approach for highly selective identification of vaccine candidates: the Group A Streptococcus case.* Mol Cell Proteomics, 2012. **11**(6): p. M111.015693.
15. Pandey, M., et al., *A synthetic M protein peptide synergizes with a CXC chemokine protease to induce vaccine-mediated protection against virulent streptococcal pyoderma and bacteremia.* J Immunol, 2015. **194**(12): p. 5915-25.
16. Pandey, M., et al., *Combinatorial Synthetic Peptide Vaccine Strategy Protects against Hypervirulent CovR/S Mutant Streptococci.* J Immunol, 2016. **196**(8): p. 3364-74.
17. Rivera-Hernandez, T., et al., *Differing Efficacies of Lead Group A Streptococcal Vaccine Candidates and Full-Length M Protein in Cutaneous and Invasive Disease Models.* MBio, 2016. **7**(3).

18. Turner, C.E., et al., *Impact of immunization against SpyCEP during invasive disease*
with two streptococcal species: Streptococcus pyogenes and Streptococcus equi.
Vaccine, 2009. **27**(36): p. 4923-9.

19. Kaur, S.J., et al., *The CXC chemokine-degrading protease SpyCep of Streptococcus*
pyogenes promotes its uptake into endothelial cells. J Biol Chem, 2010. **285**(36): p.
27798-805.

20. Abate, F., et al., *Cloning, expression, purification, crystallization and preliminary X-ray*
diffraction analysis of SpyCEP, a candidate antigen for a vaccine against
Streptococcus pyogenes. Acta Crystallogr Sect F Struct Biol Cryst Commun, 2013.
69(Pt 10): p. 1103-6.

21. Lynskey, N.N., et al., *RocA Truncation Underpins Hyper-Encapsulation, Carriage*
Longevity and Transmissibility of Serotype M18 Group A Streptococci. PLOS
Pathogens, 2013. **9**(12): p. e1003842.

22. Edwards, R.J., et al., *C-terminal antibodies (CTAbs): a simple and broadly applicable*
approach for the rapid generation of protein-specific antibodies with predefined
specificity. Proteomics, 2007. **7**(9): p. 1364-72.

23. Rawlings, N.D., A.J. Barrett, and A. Bateman, *MEROPS: the database of proteolytic*
enzymes, their substrates and inhibitors. Nucleic acids research, 2012. **40**(Database
issue): p. D343-D350.

24. Hedstrom, L., *Serine Protease Mechanism and Specificity.* Chemical Reviews, 2002.
102(12): p. 4501-4524.

25. Corey, D.R. and C.S. Craik, *An investigation into the minimum requirements for*
peptide hydrolysis by mutation of the catalytic triad of trypsin. Journal of the
American Chemical Society, 1992. **114**(5): p. 1784-1790.

26. Paul, C. and A.W. James, *Dissecting the catalytic triad of a serine protease.* Nature,
1988. **332**(6164): p. 564.

27. Angelo, P.F., et al., *Substrate specificity of human kallikrein 6: salt and*
glycosaminoglycan activation effects. J Biol Chem, 2006. **281**(6): p. 3116-26.

28. Luettgen, J.M., et al., *Apixaban inhibition of factor Xa: Microscopic rate constants*
and inhibition mechanism in purified protein systems and in human plasma. Journal
of Enzyme Inhibition and Medicinal Chemistry, 2011. **26**(4): p. 514-526.

29. Kagawa, T.F., et al., *Model for substrate interactions in C5a peptidase from*
Streptococcus pyogenes: A 1.9 A crystal structure of the active form of ScpA. J Mol
Biol, 2009. **386**(3): p. 754-72.

30. Rodriguez-Ortega, M.J., et al., *Characterization and identification of vaccine*
candidate proteins through analysis of the group A Streptococcus surface proteome.
Nat Biotechnol, 2006. **24**(2): p. 191-7.

31. Reglinski, M., et al., *Development of a multicomponent vaccine for Streptococcus*
pyogenes based on the antigenic targets of IVIG. J Infect, 2016. **72**(4): p. 450-9.