Title:

Genome editing of *Capsaspora owczarzaki* suggests an ancestral function of the Hippo

signaling effector YAP/TAZ/Yorkie in cytoskeletal dynamics but not proliferation

Authors:

Jonathan E Phillips[1], Maribel Santos[1], Mohammed Kanchwala[2], Chao Xing[2], Duojia Pan[1*]

Affiliations:

[1]Department of Physiology, Howard Hughes Medical Institute, University of Texas

Southwestern Medical Center, Dallas, Texas 75390, USA

[2]Eugene McDermott Center for Human Growth & Development, Departments of

Bioinformatics and Clinical Sciences, University of Texas Southwestern Medical Center,

Dallas, Texas 75390, USA

* correspondence: duojia.pan@utsouthwestern.edu

Abstract:

Many genes that function in animal development are present in the close unicellular relatives of animals, but little is known regarding the premetazoan function of these genes. Here, we develop techniques for genetic manipulation in the filasterean *Capsaspora owczarzaki* and use these tools to characterize the *Capsaspora* ortholog of the Hippo signaling nuclear effector YAP/TAZ/Yorkie (coYki). In contrast to its potent oncogene activity in metazoans, we show that coYki is dispensable for cell proliferation but regulates cytoskeletal dynamics and the morphology of multicellular aggregates in *Capsaspora*. Our results suggest an ancestral role for the Hippo pathway in cytoskeletal regulation, which was later co-opted to regulate cell proliferation in animals.

One Sentence Summary: Gene disruption in a unicellular holozoan suggests an ancestral role for YAP/TAZ/Yorkie in cytoskeletal regulation and cell aggregation but not cell proliferation.

Main text:

The vast morphological diversity observed in animals is generated by developmental programs mediated by a surprisingly small set of conserved signaling pathways (*1*). An exciting and unexpected finding from comparative genomic analysis indicates that many components of these pathways are present in the closest unicellular relatives of animals (*2*). This finding raises the question of whether these genes show similar functions in animals and their unicellular relatives, and, if not, what the ancestral functions of these

genes were, and how they evolved to function in multicellular processes in animals.

Answering this question could illuminate the roots of animal multicellularity and provide

novel perspective on how to manipulate the activity of these biomedically important

pathways.

One such signaling pathway is the Hippo pathway, which coordinates cell proliferation,

differentiation, and survival in animals. Initially defined as a tumor suppressor pathway

that restricts tissue growth in *Drosophila* development, Hippo signaling plays a conserved

role in organ size control and regeneration in mammals *(3, 4)*. This pathway comprises a

core kinase cascade involving sequential activation of two kinases, Hpo/MST and

Wts/LATS, which culminates in the phosphorylation and inactivation of the potent growth-

stimulatory transcriptional coactivator Yorkie/YAP/TAZ (*5*). This core kinase cascade is

in turn regulated by diverse upstream inputs, most notably mechanical force, the state of

the actin cytoskeleton, and cell-cell or cell-substrate adhesion as part of a

mechanotransduction pathway that relays mechanical and architectural cues to gene

expression (*6*). Echoing its pervasive function in metazoan growth control, the Hippo

pathway is required for contact-inhibition of proliferation in cultured mammalian cells (*7*),

and defective Hippo signaling leading to YAP/TAZ oncogene activation is a major driver

of human cancers (*8*).

The Hippo pathway was at one point thought to be specific to animals. However, recent

genome sequencing efforts have revealed that a core Hippo pathway and many known

upstream regulators are encoded in the genomes of the closest unicellular relatives of

animals such as choanoflagellates and filasterians (*9*). The choanoflagellates are

flagellated filter feeders that can form multicellular rosette structures by incomplete

cytokinesis (*10*), whereas filasterians are amoeboid organisms characterized by actin-

rich filopodial projections (Fig. 1, A and B) and represent the most basal known unicellular

organism encoding all core components of the Hippo pathway (*9*). *Capsaspora*

owczarzaki is a filasterean originally isolated as a putative endosymbiont in the freshwater

snail *B. glabrata* that can attack and kill the parasitic schistosome *S. mansoni* (*11*).

Capsaspora can be cultured easily in growth medium containing protein source and

serum in adherent or shaking culture and can be transiently transfected, allowing for the

examination of protein localization using fluorescent fusion proteins (*12*). Although

Capsaspora tends to grow as a unicellular organism in laboratory culture conditions,

under certain conditions *Capsaspora* cells can adhere to each other to form multicellular

aggregates (Fig. 1, C to E, (*13*)), suggesting that multicellular behaviors may have been

selected for in the lineage leading to *Capsaspora*. Previously, we have shown that

overexpression of *Capsaspora* Hippo pathway components such as CoYki and CoHpo in

Drosophila elicits similar phenotypes as their *Drosophila* counterpart (*9*), suggesting that

a biochemically active Hippo signaling pathway likely evolved well before the emergence

of Metazoa. Besides the Hippo pathway, the *Capsaspora* genome encodes homologues

of other key regulators of animal development such as components of the integrin

adhesome (*14*), cadherins (*15*), tyrosine kinases such as Src and Abl (*16*), NF-kB (*17*)

and p53 (*18*). Thus, *Capsaspora* offers a unique opportunity to elucidate the ancestral

function of these important developmental regulators in a close unicellular relative of

animals.

In an effort to make *Capsaspora* a tractable system for evolutionary cell biology studies, we developed genetic tools for overexpression and loss-of-function analysis. We first established a method for engineering stable *Capsaspora* cell lines by transfecting a vector encoding the fluorescent protein mScarlet and resistance to the antibiotic Geneticin (GenR). After transfection, antibiotic selection, and generation of clonal lines by serial dilution, we were able to generate uniformly mScarlet-positive clonal populations of cells (Fig. 1F). Besides GenR, transfection of *Capsaspora* with genes encoding resistance to nourseothricin (NatR) or hygromycin (HygR) followed by selection with the respective antibiotic could also generate drug-resistant transformants (fig. S1). These results demonstrate that stable *Capsaspora* cell lines expressing multiple transgenes can be generated.

We next sought to develop tools for loss-of-function analysis in *Capsaspora*. After unsuccessful attempts with CRISPR-Cas9-mediated genome editing, possibly due to the lack of nonhomologous end joining machinery components in *Capsaspora*, we used homologous recombination for gene disruption by replacing each allele of a diploid *Capsaspora* gene with a distinct selectable antibiotic marker (Fig. 1G). This strategy allowed us to generate a cell line in which both *coYki* alleles were disrupted (fig. S2), with no detectable expression of the deleted region of *coYki* (Fig. 1, H to J). These results demonstrate the generation of a homozygous *coYki* disruption mutant (subsequently *coYki -/-*) and establish a method for the generation of mutants by gene targeting in *Capsaspora.*

To determine the effect of loss-of-coYki on cell proliferation, we examined the proliferation

of *Capsaspora* cells in adherent or shaking culture. In contrast to what may be expected

based on previous studies in animal cells, we observed no significant difference in cell

proliferation between WT and *coYki -/-* cells in either condition (Fig. S3). We next sought

to examine cell proliferation within *Capsaspora* aggregates. Previously, aggregates have

been generated by gentle shaking of cell cultures (*13*). To make aggregates more

amenable to extended imaging *in situ*, we developed a condition that induced robust

aggregate formation by plating *Capsaspora* cells in low-adherence wells (Fig. 1, C to E).

Under this condition, most cells coalesced into aggregates over 2-3 days, leaving few

isolated cells in the culture. Treatment with the calcium chelator EGTA resulted in rapid

dissociation of the aggregates (fig. S4), suggesting that calcium-dependent cell adhesion

mediates aggregate integrity. We used EdU incorporation to label proliferating cells in

aggregates, and observed no significant difference in the percent of EdU-positive cells

between WT and *coYki -/-* genotypes (Fig. 2, A and B). Unlike clones of cultured

mammalian cells that often display increased cell proliferation at the clonal boundary (*19*),

distribution of EdU+ cells was homogeneous within *Capsaspora* aggregates (Fig. 2A),

suggesting that a mechanism akin to contact-mediated inhibition of proliferation is absent

in *Capsaspora*. Together, these results suggest that the rate of proliferation of

Capsaspora is independent of YAP/TAZ/Yorkie or contact inhibition.

While examining *Capsaspora* aggregates, we observed a profound phenotype in the

shape of the *coYki -/-* aggregates. After aggregate induction by plating cells in low-

adherence conditions, WT cell aggregates showed a round morphology (Fig. 2C). In

contrast, *coYki -/-* aggregates were asymmetric and less circular than WT aggregates

(Fig. 2D). Computational analysis of aggregate morphology and size revealed that WT

and *coYki -/-* aggregate size was not significantly different (Fig. 2E), whereas *coYki -/-*

aggregates were significantly less circular than WT aggregates (Fig. 2F). Optical

sectioning of aggregates expressing mScarlet showed that WT aggregates were thicker

perpendicular to the culture surface and more spherical than *coYki -/-* aggregates (Fig.

2G). Time-lapse microscopy of aggregation over a period of six days showed that,

whereas WT cells accreted into round aggregates that fused with other aggregates when

in close proximity but never underwent fission, *coYki -/-* aggregates were dynamically

asymmetrical, showed a progressive reduction in circularity, and occasionally underwent

fission (movie S1). Taken together, these findings uncover a critical role for coYki in social

interactions underlying aggregate morphology, but not cell proliferation, in *Capsaspora*.

Cell-cell adhesion and cell-substrate adhesion affect the morphology and stability of

tumor cell aggregates (*20, 21*). We therefore tested whether cell-cell adhesion and cell-

substrate adhesion differ in WT *and coYki -/- cells.* A prediction of differential cell-cell

adhesion is that mosaic aggregates of WT and *coYki -/-* cells may show cell sorting within

aggregates (*22*). To test this prediction, we induced aggregate formation after mixing

mCherry-labeled WT or *coYki -/-* cells with unlabeled WT cells at 1:9 ratio. The distribution

of mCherry-labeled WT or *coYki -/-* cells within such mixed aggregates was similar and

no cell sorting was evident (fig. S5)**,** indicating that individual *coYki -/-* cells are competent

to adhere to and integrate within aggregates and suggesting that the aberrant morphology

of *coYki -/-* aggregates is an emergent property of a large number of *coYki -/-* cells. To

further test whether cell-cell adhesion differs in WT and *coYki -/-* cells, we vigorously

vortexed *Capsaspora* culture to separate individual cells, allowed cell clumps to form by

cell-cell adhesion under gentle rotation, and then counted the number of cells in each

clump. Numbers of cells per clump were similar for WT and *coYki -/-* cells (fig. S6A),

suggesting that the occurrence of cell-cell adhesion is similar in these genotypes. We

next tested whether cell-substrate adhesion differed in WT and *coYki -/-* cells by agitating

cultures of adherent cells and counting the number of cells that disassociate from the

culture surface. After agitation, significantly more *coYki -/-* cells remained attached to the

culture surface compared to WT cells (fig. S6B), indicating that coYki negatively regulates

cell-substrate adhesion. Together, our data suggest that coYki affects the morphology of

cell aggregates by affecting cell-substrate adhesion but not cell-cell adhesion.

To characterize properties of *coYki -/-* cells that may contribute to abnormal aggregate

morphology, we used time-lapse microscopy to examine the dynamic morphology of

individual WT and *coYki -/-* cells in adherent culture. Interestingly, the cortices of *coYki -*

/- cells were much more dynamic than those of WT cells. Whereas *coYki -/-* cells display

extensive membrane protrusions and retractions, such dynamic membrane structures

were rarely observed in WT cells (movie S2). To characterize the differences in cell

membrane dynamics, we quantified the occurrence of cell protrusions in WT cells, *coYki*

-/- cells, and *coYki -/-* cells expressing a *coYki* rescue transgene*.* WT cells rarely showed

protrusions. In contrast, *coYki -/-* cells showed an average of 3.1 protrusions per minute

(Fig. 3A), a phenotype that was rescued in *coYki -/-* cells expressing a *coYki* transgene

(Fig. 3A, movie S3). These results suggest that coYki affects the cortical dynamics of

Capsaspora cells.

In principle, the extensive membrane protrusions made by *coYki -/-* cells may represent

F-actin-rich pseudopodia, which drive amoeboid motility, or, alternatively, F-actin-poor

blebs, which are actin-depleted extensions of plasma membrane generated by

disassociation of the membrane from the cell cortex or cortical rupture (*23*). To distinguish

between these possibilities, we stably expressed a fusion protein of the F-actin-binding

Lifeact peptide (*24*) with mScarlet in WT and *coYki -/-* cells and examined F-actin

dynamics by time-lapse microscopy in adherent cells. At the basal side of WT cells,

Lifeact-mScarlet was often enriched at the cell edge corresponding to the direction of cell

movement, suggesting that *Capsaspora* undergoes F-actin-mediated amoeboid

locomotion (movie S4)**.** As cells move, some Lifeact-mScarlet signal appears as

stationary foci on the basal surface, suggesting the existence of actin-rich structures that

mediate cell-substrate interaction in *Capsaspora* (Fig. 3B, movie S4). Optical sections at

mid-height of WT cells showed that F-actin was enriched at a circular cell cortex, where

transient bursts of small F-actin puncta were often observed (Fig. 3B, movie S5). In

contrast, cortical shape was less circular in *coYki -/-* cells, and F-actin concentration along

the cortex was less uniform (Fig. 3B). During the formation of the membrane protrusions

produced by *coYki -/-* cells, F-actin was initially depleted as compared to the rest of the

cell (Fig. 3C, movie S6). After the membrane protrusions had fully extended, F-actin

accumulated first at the distal end and then along the entire cortex of the protrusion,

followed by the regression of the protrusion. A similar sequence of cytoskeletal events

occurs in blebbing mammalian cells (*25*). These results suggest that the protrusions

observed in *coYki -/-* cells are actin-depleted blebs, and that after bleb formation, F-actin

re-accumulates within the bleb leading to cortical reformation and bleb regression.

To determine whether the abnormal cortical cytoskeleton is specific to *coYki -/-* cells in

adherent culture, we examined *coYki -/-* cells within aggregates by time-lapse

microscopy. Lifeact-mScarlet-expressing cells were mixed with cells stably expressing

the green fluorescent protein Venus so that Lifeact-mScarlet-expressing cells were well-

spaced and able to be imaged individually. As in adherent cells, *coYki -/-* cells within

aggregates showed dynamic formation of actin-depleted blebs (movie S7). Thus, *coYki -*

/- cells show aberrant cortical cytoskeleton and excessive membrane blebbing in both

isolated adherent cells and within aggregates.

Loss of coYki causes both bleb-like protrusions and abnormal aggregate morphology,

suggesting that the cellular phenotypes observed in *coYki -/-* cells (membrane blebbing

and related cortical cytoskeletal defects) may underlie the aberrant aggregate

morphology. We tested this hypothesis by treating *coYki -/-* cells with Blebbistatin, a

myosin II inhibitor named after its bleb-inhibiting activity in mammalian cells (*26*).

Strikingly, not only did low concentrations of Blebbistatin suppress the formation of blebs

in *coYki -/-* cells (Fig. 3D, movie S8), Blebbistatin also rescued the abnormal morphology

of *coYki -/-* aggregates (Fig. 3E). Whereas *coYki -/-* cell aggregates were asymmetric and

showed low circularity (Fig. 3E'''), *coYki -/-* aggregates treated with blebbistatin were

circular and resembled WT aggregates (Fig. 3E'''' and 3F). These results suggest a

causal link between the membrane/cytoskeletal defects observed in individual *coYki -/-*

cells and the abnormal morphology of *coYki -/-* aggregates.

To characterize the transcriptional targets of coYki that may underlie the

morphological/cytoskeletal defects of *coYki -/-* cells, we performed RNAseq on WT and

coYki -/- cells in adherent culture. This analysis revealed 1205 differentially expressed

genes, including 397 downregulated and 808 upregulated genes in the *coYki* mutant.

Functional enrichment analysis of these two gene sets revealed distinct enrichment of

functional categories (Fig. 4, A and B). Genes predicted to encode actin-binding proteins

were enriched in the set of genes upregulated in the *coYki -/-* mutant, suggesting that

these genes may play a role in the aberrant cytoskeletal dynamics observed in *coYki -/-*

cells. To further inquire into the potential function of coYki-regulated genes, we searched

the 1205 differentially expressed genes against a previously reported *Capsaspora*

phylome (*27*) to identify 638 *Capsaspora* genes with predicted human or mouse

orthologs. Ingenuity Pathway Analysis of this mammalian ortholog set showed that the

two most significantly enriched functional categories corresponded to cell movement and

cell migration (Fig. 4C). Notably, regulation of cell migration was reported as the most

enriched functional category among YAP target genes in glioblastoma cells (*28*), and cell

motility was an enriched category in an integrative analysis of gene regulatory networks

downstream of YAP/TAZ utilizing transcriptomic and cistromic data from multiple human

tissues (*29*). However, in contrast to functional enrichment studies in metazoan systems

examining genes regulated by Yorkie/YAP/TAZ (*28, 30, 31*), no enrichment was detected

in functional categories of cell proliferation or the cell cycle (Fig. 4C). This finding is

consistent with the lack of a proliferative phenotype in *coYki -/-* cells.

In summary, we show that coYki regulates cytoskeletal dynamics and potentially other

cortical actin-related processes but is dispensable for the proliferation of *Capsaspora*

cells. Such coYki-dependent cortical cytoskeletal function, in turn, regulates the

morphology of *Capsaspora* aggregates in a manner that may involve tuning the strength

of cell-substrate adhesion. Although the natural history of *Capsaspora*, including the

selective advantage of cell aggregation, is poorly understood, our findings suggest that

cytoskeletal regulation represents an ancestral function of the Hippo pathway predating

the origin of Metazoa, and this pathway was later co-opted to control cell proliferation

after the unicellular-to-multicellular transition. Given that the cortical cytoskeleton is both

an important upstream regulator and a downstream effector of Hippo signaling in animal

cells (*32, 33*), the co-option of the Hippo pathway for cell proliferation control in early

metazoans may have provided a convenient mechanism that couples cell proliferation

with cytoskeletal and mechanical properties of the cells. Besides illuminating an ancestral

function of the Hippo pathway, the genetic tools we have developed for modifying the

Capsaspora genome provide an unprecedented opportunity to interrogate the

premetazoan functions of other developmental regulators that are present in the genomes

of close unicellular relatives of animals.

References and Notes:

1.   A. Pires-daSilva, R. J. Sommer, The evolution of signalling pathways in animal
development. *Nat Rev Genet* **4**, 39-49 (2003).

2.   N. Ros-Rocher, A. Perez-Posada, M. M. Leger, I. Ruiz-Trillo, The origin of
animals: an ancestral reconstruction of the unicellular-to-multicellular transition.
Open Biol **11**, 200359 (2021).

3.   S. Wu, J. Huang, J. Dong, D. Pan, hippo encodes a Ste-20 family protein kinase
that restricts cell proliferation and promotes apoptosis in conjunction with
salvador and warts. *Cell* **114**, 445-456 (2003).

4.   K. F. Harvey, C. M. Pfleger, I. K. Hariharan, The Drosophila Mst ortholog, hippo,
restricts growth and cell proliferation and promotes apoptosis. *Cell* **114**, 457-467
(2003).

5.   J. Huang, S. Wu, J. Barrera, K. Matthews, D. Pan, The Hippo signaling pathway
coordinately regulates cell proliferation and apoptosis by inactivating Yorkie, the
Drosophila Homolog of YAP. *Cell* **122**, 421-434 (2005).

6.   Y. Zheng, D. Pan, The Hippo Signaling Pathway in Development and Disease.
Dev Cell **50**, 264-282 (2019).

7.   B. Zhao *et al.*, Inactivation of YAP oncoprotein by the Hippo pathway is involved
in cell contact inhibition and tissue growth control. *Genes Dev* **21**, 2747-2761
(2007).

8.   Y. Wang *et al.*, Comprehensive Molecular Characterization of the Hippo
Signaling Pathway in Cancer. *Cell Rep* **25**, 1304-1317 e1305 (2018).

9.   A. Sebe-Pedros, Y. Zheng, I. Ruiz-Trillo, D. Pan, Premetazoan origin of the hippo
signaling pathway. *Cell Rep* **1**, 13-20 (2012).

10.   S. R. Fairclough, M. J. Dayel, N. King, Multicellular development in a
choanoflagellate. *Curr Biol* **20**, R875-876 (2010).

11.   H. H. Stibbs, A. Owczarzak, C. J. Bayne, P. DeWan, Schistosome sporocyst-
killing Amoebae isolated from Biomphalaria glabrata. *J Invertebr Pathol* **33**, 159-
170 (1979).

12.   H. Parra-Acero *et al.*, Transfection of Capsaspora owczarzaki, a close unicellular
relative of animals. *Development* **145**, (2018).

13.   A. Sebe-Pedros *et al.*, Regulated aggregative multicellularity in a close unicellular
299   relative of metazoa. *Elife* **2**, e01287 (2013).

300   14.   A. Sebe-Pedros, A. J. Roger, F. B. Lang, N. King, I. Ruiz-Trillo, Ancient origin of
301   the integrin-mediated adhesion and signaling machinery. *Proc Natl Acad Sci U S*
302   *A* **107**, 10142-10147 (2010).

303   15.   S. A. Nichols, B. W. Roberts, D. J. Richter, S. R. Fairclough, N. King, Origin of
304   metazoan cadherin diversity and the antiquity of the classical cadherin/beta-
305   catenin complex. *Proc Natl Acad Sci U S A* **109**, 13046-13051 (2012).

306   16.   H. Suga *et al.*, Genomic survey of premetazoans shows deep conservation of
307   cytoplasmic tyrosine kinases and multiple radiations of receptor tyrosine kinases.
308   *Sci Signal* **5**, ra35 (2012).

309   17.   L. M. Williams, T. D. Gilmore, Looking Down on NF-kappaB. *Mol Cell Biol* **40**,
310   (2020).

311   18.   A. Sebe-Pedros, B. M. Degnan, I. Ruiz-Trillo, The origin of Metazoa: a unicellular
312   perspective. *Nat Rev Genet* **18**, 498-512 (2017).

313   19.   H. W. Fisher, J. Yeh, Contact inhibition in colony formation. *Science* **155**, 581-
314   582 (1967).

315   20.   A. F. Blandin *et al.*, Glioma cell dispersion is driven by alpha5 integrin-mediated
316   cell-matrix and cell-cell interactions. *Cancer Lett* **376**, 328-338 (2016).

317   21.   L. Saias, A. Gomes, M. Cazales, B. Ducommun, V. Lobjois, Cell-Cell Adhesion
318   and Cytoskeleton Tension Oppose Each Other in Regulating Tumor Cell
319   Aggregation. *Cancer Res* **75**, 2426-2433 (2015).

320   22.   R. A. Foty, M. S. Steinberg, The differential adhesion hypothesis: a direct
321   evaluation. *Dev Biol* **278**, 255-263 (2005).

322   23.   G. Charras, E. Paluch, Blebs lead the way: how to migrate without lamellipodia.
323   *Nat Rev Mol Cell Biol* **9**, 730-736 (2008).

324   24.   J. Riedl *et al.*, Lifeact: a versatile marker to visualize F-actin. *Nat Methods* **5**, 605-
325   607 (2008).

326   25.   G. T. Charras, C. K. Hu, M. Coughlin, T. J. Mitchison, Reassembly of contractile
327   actin cortex in cell blebs. *J Cell Biol* **175**, 477-490 (2006).

328   26.   A. F. Straight *et al.*, Dissecting temporal and spatial control of cytokinesis with a
329   myosin II Inhibitor. *Science* **299**, 1743-1747 (2003).

330   27.   A. Sebe-Pedros *et al.*, High-Throughput Proteomics Reveals the Unicellular
331   Roots of Animal Phosphosignaling and Cell Differentiation. *Dev Cell* **39**, 186-197
332   (2016).

333   28.   C. Stein *et al.*, YAP1 Exerts Its Transcriptional Control via TEAD-Mediated
334   Activation of Enhancers. *PLoS Genet* **11**, e1005465 (2015).

335   29.   M. Paczkowska *et al.*, Integrative pathway enrichment analysis of multivariate
336   omics data. *Nat Commun* **11**, 735 (2020).

337   30.   A. Ikmi *et al.*, Molecular evolution of the Yap/Yorkie proto-oncogene and
338   elucidation of its core transcriptional program. *Mol Biol Evol* **31**, 1375-1390
339   (2014).

340   31.   F. Zanconato *et al.*, Genome-wide association between YAP/TAZ/TEAD and AP-
341   1 at enhancers drives oncogenic growth. *Nat Cell Biol* **17**, 1218-1227 (2015).

342   32.   H. Deng *et al.*, Spectrin couples cell shape, cortical tension, and Hippo signaling
343   in retinal epithelial morphogenesis. *J Cell Biol* **219**, (2020).

344   33.   B. G. Fernandez *et al.*, Actin-Capping Protein and the Hippo pathway regulate F-
345   actin and tissue growth in Drosophila. *Development* **138**, 2337-2346 (2011).

346   34.   H. Parra-Acero *et al.*, Integrin-Mediated Adhesion in the Unicellular Holozoan
347   Capsaspora owczarzaki. *Curr Biol* **30**, 4270-4275 e4274 (2020).

348   35.   D. S. Bindels *et al.*, mScarlet: a bright monomeric red fluorescent protein for
349   cellular imaging. *Nat Methods* **14**, 53-56 (2017).

36. D. M. Veltman, G. Akar, L. Bosgraaf, P. J. Van Haastert, A new set of small,
extrachromosomal expression vectors for Dictyostelium discoideum. *Plasmid* **61**,
110-118 (2009).

37. S. S. Lehman *et al.*, Versatile nourseothricin and streptomycin/spectinomycin
resistance gene cassettes and their use in chromosome integration vectors. *J
Microbiol Methods* **129**, 8-13 (2016).

38. S. H. Park *et al.*, Generation and application of new rat monoclonal antibodies
against synthetic FLAG and OLLAS tags for improved immunodetection. *J
Immunol Methods* **331**, 27-38 (2008).

39. S. J. Charette, P. Cosson, Preparation of genomic DNA from Dictyostelium
discoideum for PCR analysis. *Biotechniques* **36**, 574-575 (2004).

40. S. Andrews, FastQC: a quality control tool for high throughput sequence data. .
Available online at: http://www.bioinformatics.babraham.ac.uk/projects/fastqc,
(2010).

41. S. Wingett, FastQ Screen: quality control tool to screen a library of sequences in
FastQ format against a set of sequence databases. . *Available online at:
http://www.bioinformatics.babraham.ac.uk/projects/fastq_screen*, (2011).

42. A. Dobin *et al.*, STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15-
21 (2013).

43. Y. Liao, G. K. Smyth, W. Shi, featureCounts: an efficient general purpose
program for assigning sequence reads to genomic features. *Bioinformatics* **30**,
923-930 (2014).

44. M. D. Robinson, D. J. McCarthy, G. K. Smyth, edgeR: a Bioconductor package
for differential expression analysis of digital gene expression data. *Bioinformatics*
26, 139-140 (2010).

45. J. Huerta-Cepas, S. Capella-Gutierrez, L. P. Pryszcz, M. Marcet-Houben, T.
Gabaldon, PhylomeDB v4: zooming into the plurality of evolutionary histories of a
genome. *Nucleic Acids Res* **42**, D897-902 (2014).

46.   E. Paradis, K. Schliep, ape 5.0: an environment for modern phylogenetics and
evolutionary analyses in R. *Bioinformatics* **35**, 526-528 (2019).

47.   A. Kramer, J. Green, J. Pollard, Jr., S. Tugendreich, Causal analysis approaches
in Ingenuity Pathway Analysis. *Bioinformatics* **30**, 523-530 (2014).

48.   W. Huang da, B. T. Sherman, R. A. Lempicki, Bioinformatics enrichment tools:
paths toward the comprehensive functional analysis of large gene lists. *Nucleic*
Acids Res **37**, 1-13 (2009).
385

Acknowledgements:

We thank the McDermott Sequencing Core and the Next-Generation Sequencing Core

for RNA sequencing, and Dr. Iñaki Ruiz-Trillo for sharing information on *Capsaspora*

vectors.

Funding: This work was supported in part by National Institute of Health grant

EY015708 to D.P. During this work J.E.P was a Howard Hughes Medical Institute fellow

of the Life Sciences Research Foundation. D.P. is an investigator of the Howard

Hughes Medical Institute.

Author Contributions: JEP and DP conceived the study and designed experiments.

JEP and MS performed experiments. JEP, MK, and CX analyzed RNAseq data. JEP,

DP, and MK wrote the original manuscript draft. JEP, MS, DP, MK, and CX revised and

edited the manuscript.

Competing interests: Authors declare that they have no competing interests.

Data and materials availability: RNAseq data generated in this study have been

deposited into the NCBI SRA (Bioproject PRJNA759885). The following plasmids have

been deposited at Addgene: pJP72 (#176479), pJP102 (#176480), pJP103 (#176481),

and pJP118 (#176494).

Supplementary Materials

Materials and methods

Figs. S1 to S6

References 34 to 48

Movies S1 to S8

Data S1 to S3

Fig. 1. Stable transgene expression and gene knockout in *Capsaspora*. (A) A solitary

Capaspora cell showing thin F-actin-enriched projections. Phalloidin and DAPI are used

to stain F-actin and DNA, respectively. **(B)** Cells at higher densities can contact other cells

through projections. Scale bar is 5 microns. **(C-D)** *Capsapsora* cells were diluted to 7.5 x

10^5 cells/mL and inoculated into either a standard polystyrene culture plate **(C)** or a low-

adherence plate **(D).** At 3 days, cells in standard plates grow as a monolayer whereas

cells in low-adherence plates form round aggregates. **(E)** An aggregate stained for F-actin

shows actin-rich connections between cells within an aggregate. Scale bar is 10 microns.

(**F)** WT *Capsaspora* cells and a clonal population of cells stably expressing mScarlet were

imaged to show fluorescence. Scale bar is 20 microns. **(G)** Strategy for disrupting the

coYki gene by homologous recombination. GenR, geneticin resistance cassette; TBD,

Tead-binding domain; WW, WW domain. **(H)** Strategy for demonstrating disruption of

coYki by qPCR. Amplicon 1 corresponds to a region of the *coYki* gene that should be

absent in a homozygous mutant, whereas Amplicon 2 corresponds to a segment of the

gene 3' of the region targeted for deletion. **(I-J)** qPCR of Amplicons 1 and 2 in WT and

putative *coYki -/-* cells. The differences between WT and *coYki -/-* for both Amplicons 1

and 2 are significant (t-test).

Fig. 2. Loss of coYki results in no apparent effect on cell proliferation but alters the morphology of multicellular aggregates. (A) To examine proliferation within aggregates, EdU was used to label proliferating cells within aggregates grown in low-adherence plates. **(B)** The percent of cells positive for EdU within aggregates was quantified. Each red circle indicates measurement results for a single aggregate, and mean $\pm$ SEM is shown in black. The difference between WT ant *coYki -/-* cells is not significant (t-test). **(C-D)** WT or *coYki -/-* cells were innoculated into low-adherence plates, and cell aggregates were imaged at 5 days. Scale bar is 75 microns. **(E-F)** Aggregate size and circularity was measured from aggregate images using ImageJ. The difference in aggregate circularity between WT and *coYki -/-* is significant (t-test). Bars indicate the mean $\pm$ SEM (n=3 with 15 aggregates measured for each independent experiment), and dots indicate values for individual aggregates. **(G)** Orthogonal views of WT and *coYki -/-* aggregates are shown. Cell aggregates stably expressing mScarlet were imaged live by confocal microscopy. Red lines show the location of the culture surface. Scale bar is 50 microns.

Fig. 3. Loss of coYki results in bleb-like distortions at the cell cortex. (A) *Capsaspora*

cells on a glass surface were imaged by time-lapse microscopy and individual cells were

scored for number of protrusions per minute. A protrusion is defined as a localized

extension of the cell boundary that disrupts a previously inert region of the cortex and/or

changes the direction of movement of the cell. Differences between WT and *coYki -/-* are

significant (one-way ANOVA, Tukey's test). Values from 3 independent experiments with

20 cells measured for each condition per experiment are shown. Red dots indicate the

measurement from an individual cell, and black bars indicate the mean $\pm$ SEM of the

means from each independent experiment. **(B)** Cells on a glass surface stably expressing

Lifeact-mScarlet were imaged focusing on the base of the cell at the substrate ("basal")

or at the mid-height of the cell ("medial"). Arrowhead shows an example of Lifeact signal

at the base of the cell that remains stationary as the cell moves (see Movie S4). Scale

bar is 5 microns. **(C)** Time series of a *coYki -/-* cell stably expressing Lifeact-mScarlet on

a glass surface. White arrows indicate Lifeact-depleted bleb-like protrusions, and yellow

arrowheads indicate areas of cell cortex constriction, which correlate with increased

Lifeact signal. Scale bar is 5 microns. **(D)** Cells on a glass surface were treated with

DMSO or 1 µM blebbistatin for 1 hour, and then cells were imaged by time lapse

microscopy and the number of protrusions per cell was quantified. **(E)** Cells were

inoculated into low-adherence plates with DMSO **(E',E''')** or 1 µM blebbistatin **(E'',E'''')**,

and aggregates were imaged after 5 days. Scale bar is 75 microns. **(F)** Circularity of

aggregates was measured with ImageJ using images of aggregates from the indicated

conditions. Black bars indicate the mean $\pm$ SEM (n=4 with 15 aggregates measured for

each independent experiment), and red dots indicate the measurements for individual

aggregates. Differences in circularity between WT and *coYki -/-* aggregates treated with

DMSO are significant (one-way ANOVA, Dunnett's test), whereas differences between

WT aggregates treated with DMSO and *coYki -/-* aggregates treated with blebbistatin are

not. "Bleb" indicates blebbistatin.

486
487

A

Enriched categories, genes downregulated in *coYki -/-*

B

Enriched categories, genes upregulated in *coYki -/-*

C

**IPA analysis of mammalian orthologs:
Diseases and Biological Functions enrichment**

Fig. 4. Functional enrichment of genes differentially expressed in *coYki -/-* cells.

Top enriched categories from the sets of genes significantly downregulated **(A)** or upregulated **(B)** in *coYki -/-* cells compared to WT are shown. **(C)** Ingeneuity Pathway Analysis was performed on a set of predicted human/mouse orthologs of *Capsaspora* genes with significant changes in expression in *coYki -/-* cells. Enriched diseases and biological functions with false discovery rate (FDR) $\leq$ 0.05 are shown.

501 # Supplementary Materials for

503 Genome editing of *Capsaspora owczarzaki* suggests an ancestral function

504 of the Hippo signaling effector YAP/TAZ/Yorkie in cytoskeletal dynamics but

505 not proliferation

506
507 Jonathan E Phillips, Mohammed Kanchwala, Chao Xing, and Duojia Pan

508
509 Correspondence to: duojia.pan@utsouthwestern.edu
510
511

512 **This PDF file includes:**

513
514   Materials and Methods
515   Figs. S1 to S6
516   Captions for Movies S1 to S8
517   Captions for Data S1 to S3

518
519 **Other Supplementary Materials for this manuscript include the following:**

520
521   Movies S1 to S8
522   Data S1 to S3:

523
524   Data S1. Sequences of synthesized gene fragments used in this study
525   Data S2. Sequences of oligonucleotides used in this study
526   Data S3. Plasmids constructed for this study

527
528

Materials and Methods

Cell culture

Capsaspora cells were maintained in ATCC Medium 1034 (modified PYNFH medium)

supplemented with 25 μg/mL Ampicillin (referred to below as "growth medium") in a 23°C

incubator. Cells were kept in either 25 cm^2 cell culture flasks in 8 mL medium or in 75 cm^2

cell culture flasks in 15 ml medium. To induce cell aggregation, cells were resuspended

to 7.5 x 10^5 cells/ml in growth medium, and 1 ml of this cell suspension was added per

well to a 24-well ultra-low attachment plate (Sigma CLS3473). To test the effect of

blebbistatin on aggregate morphology, a 20 mM stock solution of blebbistatin (Sigma

B0560) in DMSO was made, and cells were resuspended to 7.5 x 10^5 cells/ml in growth

medium with 1 μm blebbistatin or DMSO only as a vehicle control. Aggregates were then

generated as described above and imaged with a Nikon Eclipse Ti inverted microscope

with NIS-Elements acquisition software.

To characterize cell proliferation during growth on a solid substrate, *Capsaspora* cells

were diluted to 1 x 10^5 cells/ml in growth medium, and 800 μl of this dilution was pipetted

into multiple wells in a 24-well polystyrene cell culture plate. To prevent edge effects,

wells at the edge of the plate were not used and were instead filled with 800 μl of water.

The plates were then kept in in a 23°C incubator. Each day, 1.5 μl of a 500 mM EDTA

solution was added to a well, and cells in the well were resuspended by pipetting up and

down for one minute. Resuspended cells were then transferred to a 1.5 mL tube,

vortexed, and then the cell density was determined by hemocytometer. To characterize

cell proliferation in shaking culture, cells were diluted to 1 x 10^5 cells/ml in growth medium,

and 20 ml of this dilution was added to a 125 ml Erlenmeyer culture flask. Cultures were

then incubated in an orbital shaker at 150 rpm and 23°C. Cell density was determined

daily by transferring 200 µl of culture to a 1.5 ml tube, adding 1.5 µl of 500 mM EDTA

solution to disassociate any aggregated cells, vortexing, and counting by hemocytometer.

<u>Fixation and staining of aggregates</u>

To fix and stain aggregates, aggregates were generated as described above, and then

aggregates were collected from a 24-well plate by gentle pipetting. 500 µl of aggregate

suspension was added dropwise to 9 ml of PM PFA (100 mM PIPES ph 6.9, 0.1 mM

$MgSO_4$, 4% PFA filtered through a 0.45 µm syringe filter) in a 15 mL polystyrene tube,

and tubes were left undisturbed for 1 hour to allow fixation. Aggregates were then

centrifuged at 1000 x g for 3 minutes and resuspended in 500 µl of PEM buffer (100 mM

PIPES pH 6.9, 1 mM EGTA, 0.1 mM $MgSO_4$). After transferring aggregates to a

microcentrifuge tube, aggregates were centrifuged for 2000 x g for 2 minutes, the

supernatant was removed, and aggregates were resuspended in 300 µl of PEM.

As we found that aggregates were fragile during continued centrifugation-resuspension

cycles, we developed a protocol where solution changes could be done without

centrifugation using 24-well plate inserts with an 8 micron pore membrane at the insert

base (Corning 3422). Before use with aggregates, 100 µl PEM buffer was added to a well

in a 24-well plate, 300 µl of PEM was added to a membrane insert, and the insert was

placed in the well and allowed to drain (contact between the membrane and the solution

within the well is critical for draining). Aggregates in 300 µl of PEM were added to the

insert, and the insert was moved to a new well with 100 µl of PEM and allowed to drain.

As a wash step, the membrane insert was then moved to a well containing 1200 µl of

PEM, allowing the membrane insert to fill with PEM buffer, and after a 5 minute incubation,

the membrane was moved to a new well with 100 µl of PEM and allowed to drain. The

membrane insert was then moved to a well containing 1200 µl of PEM

block/permeabilization solution (100 mM PIPES pH 6.9, 1 mM EGTA, 0.1 mM $MgSO_4$,

1% BSA, 0.3% Triton X-100) and incubated for 30 minutes at room temperature. The

membrane insert was moved to a well with 100 µl of PEM and allowed to drain and then

moved into a well containing 1 mL PEM with DAPI (2 µg/mL) and Phalloidin (0.1 µM) and

incubated for 15 minutes at room temperature. Aggregates were then washed once more

in 1200 µl PEM. To mount aggregates for imaging, the membrane insert was placed in a

new well with 100 µl of PEM buffer and was allowed to drain, aggregates were

resuspended in the residual buffer (approximately 50 µl) by gentle pipetting, and then

aggregates were pipetted onto a microscope slide and allowed to settle for 5 minutes.

Kimwipes were then placed in contact with the edge of the liquid on the slide, allowing for

removal of excess buffer by capillary action. 15 µl of Fluoromount G (SouthernBiotech)

was pipetted onto the aggregates, which were then covered with a 22 x 40 mm cover

glass and sealed with nail polish. Aggregates were subsequently imaged with a Zeiss

LSM 880 confocal microscope. Staining of adherent cells with phalloidin was done as

previously described (*34*).

<u>EdU labeling of aggregates</u>

To examine cell proliferation within aggregates by EdU labeling, aggregates were

generated in ultra-low adherence 24-well plates as described above. 3 days after

aggregation was initiated, 100 μl of growth medium containing EdU was added to

aggregates to result in a 150 μM EdU concentration. Aggregates were incubated with

EdU for 4 hours and then fixed and resuspended in 300 μl PEM buffer as described

above. Aggregates were then processed for imaging following the manufacturer's

protocol (ThermoFisher C10339) using the 24-well plate membrane insert method

described above to transfer aggregates between solutions. For all steps, wells with 1 mL

of solution were used for washing or staining cells in membrane inserts, and wells with

100 μl of solution were used to drain wells in preparation for transfer into the next solution.

Molecular Biology

For the sequences of synthesized DNA fragments and oligonucleotides used in this study,

see Data S1 and S2, respectively. For a list of plasmids generated for this study, see Data

S3. To generate a vector for mScarlet expression in *Capsaspora* (pJP71), a DNA

construct containing an open reading frame encoding the mScarlet protein (*35*) codon-

optimized for expression in *Capsaspora* and under control of the previously described

EF1-α promoter and terminator from the pONSY-mCherry vector (*12*) was constructed by

gene synthesis (sJP1) and cloned into the EcoRV site of the pUC57-mini vector

(Genscript). A KpnI site and an AflII site were incorporated at the beginning and end of

the coding sequence, respectively, allowing for the construction of N or C-terminal protein

fusion constructs by Gibson assembly. To generate a plasmid for expression of mScarlet

and a Geneticin resistance gene (pJP72), the NeoR coding sequence from the

Dictyostelium pDM323 vector (*36*) was codon optimized for *Capsaspora*, the optimized

sequence was synthesized with the promoter and terminator from the *Capsaspora* actin

ortholog gene CAOG_06018, and the synthesized fragment (sJP2) was cloned into the

SmaI site of pJP71 by Gibson assembly. To generate plasmids expressing mScarlet and

either nourseothricin resistance (pJP102) or hygromycin resistance (pJP103), an identical

strategy to that used in constructing pJP72 was used, except either a synthesized codon-

optimized nourseothricin acetyltransferase gene (sJP3) based on the sequence of

pUC18T-mini-Tn7T-nat (*37*) or a synthesized codon-optimized hygromycin B

phosphotransferase gene (sJP4) based on the sequence of *Dictyostelium* vector pDM358

(*36*) was used.

To generate a plasmid encoding coYki and Hygromycin resistance for rescue of *coYki -/-*

phenotypes (pJP119), a DNA fragment encoding coYki and a c-terminal OLLAS epitope

tag (*38*) was generated by PCR from a synthesized a gene encoding the coYki protein

(sJP5) using primers oJP101 and oJP102. To limit recombination of the transgene with

the endogenous coYki loci, we recoded the coYki reading frame using synonymous codon

replacement, resulting in a coYki gene with 73 percent nucleotide similarity to the WT

coYki gene but encoding an identical polypeptide. This PCR product was then cloned into

a KpnI and AflII digest of pJP103 using Gibson assembly.

To generate a plasmid encoding a Lifeact-mScarlet fusion protein and hygromycin

resistance (pJP118), a DNA fragment optimized for expression in *Capsaspora* encoding

the Lifeact peptide (*24*) (sJP6) was synthesized and cloned into the KpnI site of pJP103

by Gibson assembly. To generate a plasmid expressing Venus and Geneticin resistance

(pJP114), the Venus reading frame was amplified from $p_{EF1\alpha}$ -CoH2B:Venus (Addgene

#111877, (*12*)) using primers oJP103 and oJP104 and cloned into a KpnI and AflII digest

of pJP72 by Gibson assembly.

Capsaspora Transfection

24-well plates were prepared for transfection by inserting one sterile 12mm circular glass

coverslip in each well to aid in cell adhesion during transfection. *Capsaspora* cells at

exponential growth phase were collected by pipetting medium over attached cells,

resuspended to 7.5 x 10^5 cells/ml in growth medium, and then 800 µl of this cell culture

was added per well to the prepared 24-well plate. Cells were incubated at 23°C overnight.

The following day growth medium was removed from cells and was replaced by 800 µl of

transfection medium (Scheider's Drosophila Medium (ThermoFisher 21720024) with 10%

FBS (Thermofisher 26140079) supplemented with 25 µg/mL Ampicillin). After a 10-minute

incubation at room temperature, transfection medium was removed from the cells and

replaced with 500 of fresh transfection medium. Transfection mixes were then prepared:

100 µl of Opti-MEM I Reduced Serum Medium (ThermoFisher) was added to a 1.5ml

tube, and 1 µg of transfecting DNA was added to this medium (for multiple plasmids, an

equivalent amount of each plasmid by mass was used). 3 µl of *Trans*IT-X2® transfection

reagent (Mirus Bio) was then added to the tube, and the solution was immediately mixed

by pipetting up and down. Transfection mixes were incubated at room temperature for 5

minutes, and then 70 µl of the transfection mix was added dropwise to one well in the 24-

well plate. Cells were incubated at 23°C for 24 hours, and then the medium was removed

and replaced with 800 µl of growth medium. To image fluorescent transient transfectants,

cells were then resuspended by pipetting up and down, and 300 µl of resuspended cells

in growth medium were transferred to a well of an 8-well chambered coverglass slide

(Nunc). Cells were then imaged 48 hours after addition of growth medium using a Zeiss

LSM 880 confocal microscope.

<u>Stable expression of transgenes in clonal *Capsaspora* cell lines</u>

Transforming plasmids were linearized by digestion with either ScaI-HF® or AseI

restriction enzymes (NEB), which cut within the ampicillin resistance gene, and then

purified from solution and resuspended in nuclease-free water. *Capsaspora* cells were

transfected with the linearized plasmids using the *Trans*IT-X2® transfection reagent as

described above. Two days after growth medium was added to cells following

transfection, the growth medium was removed and replaced with 800 µl of growth medium

supplemented with selective drugs. Due to observations of variability in cell viability

between transfections, three drug concentrations were tested in parallel for each

transfection: for Geneticin (ThermoFisher), cells were treated at 40, 60, or 80 µg/ml; for

Nourseothricin (GoldBio), cells were treated at 50, 75, or 100 µg/ml; for Hygromycin B

(Sigma), cells were treated at 150, 200, or 250 µg/ml. Cells were grown in selective

medium for 2 weeks, and medium was changed every 3 days. Following selection, clonal

cell populations were generated by resuspending cells in growth medium by pipetting,

diluting cells to 3 cells/mL in growth medium, and adding 100 µl of this dilution to 200 µl

of growth medium per in a well in a 48-well plate (this procedure generated approximately

10 wells with cell growth for each 48-well plate). To image fluorescence in stably

transfected lines, clonal cell populations were transferred into wells in a 4-well glass

bottom chamber slide (Nunc), incubated at 23°C for 24 hours, and cells were then imaged

using a Nikon Eclipse Ti inverted microscope with NIS-Elements acquisition software.

<u>Disruption of the Capsaspora *coYki* gene by gene targeting</u>

To disrupt *coYki,* we attempted to delete a 228 bp segment of the coYki open reading

frame that encodes the predicted Tead-binding domain (*9*) using a PCR-generated gene

targeting construct. We designed a pair of primers that amplify the drug resistance

markers described above, including the actin (CAOG_06018) promoter and terminator,

with 90 bp of homology adjacent to the sequence targeted for deletion at the 5' end of the

primers (oJP105 and oJP106). These primers were used to amplify the Geneticin

resistance cassette from pJP72 by PCR, the resulting PCR product was gel purified and

resuspended in nuclease-free water, and WT *Capsaspora* cells were transfected with this

DNA as described above. Clonal populations of drug-resistant transformants were

generated as described above, each clone was grown to confluency in one well in a 24-

well plate, cells were collected by pipetting up and down, and genomic DNA was prepared

following the protocol described below. To genotype potential mutants, we performed

diagnostic PCRs on genomic DNA: to test for presence of the WT allele, we used a

forward primer with homology to the genome 5' of the *coYki* sequence targeted for

deletion (oJP107) and a reverse primer with homology within the sequence targeted for

deletion (oJP108). To detect successful deletion, we used oJP107 as a forward primer

and a reverse primer with homology to the geneticin resistance gene (oJP109). 40

percent of analyzed clones (6 of 15 tested clones) showed a PCR product indicative of

disruption of the *coYki* allele (fig. S2B). However, all clones showed a band indicative of

an intact WT *coYki* gene. We therefore reasoned that, at least for the culture conditions

used during transfection, *Capsaspora* cells may be diploid.

After the isolation of clonal lines that were heterozygous for *coYki* disruption with the

Geneticin resistance marker as indicated by diagnostic PCR, we attempted to disrupt the

remaining *coYki* allele using nourseothricin resistance as a marker. pJP102 was used as

a template for PCR using primers oJP105 and oJP106 to generate a nourseothricin

resistance cassette flanked by homology arms targeting *coYki* for disruption. This

construct was used to transfect a Geneticin-resistant heterozygous coYki disruption

mutant, and drug selection of transfectants was done as described above using

simultaneous selection with 60 µg/ml Geneticin and 75 µg/ml nourseothricin. After

generating clonal populations of drug-resistant transformants, diagnostic PCR was done

to detect the intact *coYki* allele as described above or a deletion allele containing the

nourseothricin resistance gene using primers oJP107 and oJP110. Two cell lines that

showed absence of the WT *coYki* allele and presence of both disruption alleles encoding

either Geneticin or nourseothricin resistance were used for subsequent phenotypic

studies.

To quantify *coYki* gene expression in a putative homozygous *coYki* disruption mutant

(*coYki-/-*) by qPCR, 5×10^6 WT or *coYki-/-* cells were collected from culture flasks while

in growth medium by pipetting up and down, and RNA was isolated using a RNeasy Mini

Plus kit (Qiagen). cDNA was made using the iScript cDNA Synthesis Kit (BioRad), qPCR

reactions were made using iQ™ SRBR® Green Supermix (BioRad), and qPCR was

performed using a CFX96 Touch Real-time PCR detection system (BioRad) with the

following primers: oJP111 and oJP112 to detect GAPDH, oJP113 and oJP114 to detect

a region of coYki within the sequence targeted for deletion, and oJP115 and oJP116 to

detect a region coYki 3' of the sequence targeted for deletion.

<u>Preparing *Capsaspora* genomic DNA</u>

Capsaspora genomic DNA was prepared for PCR analysis following a procedure

previously developed for *Dictyostelium discoideum* (*39*). Cells grown in a 24-well or 48-

well plate in growth medium were collected by pipetting up and down, pipetted into a 1.5

mL tube, centrifuged at 4000 x g for 5 minutes, and resuspended in 20 µl of nuclease-

free water. 20 µl of Lysis buffer (50 mM KCl, 10 mM TRIS pH 8.3, 2.5 mM MgCl$_2$, 0.45%

NP40, 0.45% Tween 20, and 800 µg/ml Proteinase K added fresh from a 20 mg/ml stock)

was added to cells, which were then incubated at room temperature for 5 minutes. Tubes

were then placed at 95°C for 5 minutes. After cooling to room temperature, 1 µl of this

sample was then used as a template in a 20-µl diagnostic PCR.

<u>Quantification of aggregate size and circularity</u>

ImageJ was used to quantify aggregate size and circularity using aggregate images.

Images were converted to 8-bit format, processed twice using the "Smooth" function, and

a Threshold was adjusted for each individual image. The Analyze Particles command was

then run with a gate for particle size at 400-infinity micron^2 and the "exclude on edges"

option selected. The returned values for aggregate area and circularity were then used

for further analysis. Values corresponding to more than one adjacent aggregate

interpreted by the algorithm as a single particle were discarded.

<u>Assaying cell-cell adhesion and cell-substrate adhesion</u>

To examine cell-cell adhesion, cells were collected from a culture flask by pipetting growth

media over the culture surface, diluted to 1 x 10^6 cells/mL in growth medium, and vortexed

to disrupt cell clumps. 1 mL volumes of cell culture were then transferred to 1.5 mL tubes

and incubated on a Labquake^{TM} rotator (Thermo) for one hour at room temperature.

Cultures were then examined by hemocytometer, and the number of cells in each clump

of cells was counted for at least 35 cell clumps for each independent experiment.

To examine cell-substrate adhesion, cells were diluted to 5 x 10^5 cells/mL in growth

medium, and 3 mL volumes of culture were transferred per well to 6-well tissue culture-

treated polystyrene plates (Sigma CLS3506) and incubated at 23°C for 48 hours. For

agitation, plates were then placed on an orbital shaker and shaken at 140 RPM for 10

minutes, and then the medium from each well ("disassociated fraction") was transferred

to a separate tube and 3 mL of new growth medium was added to each well. 3 µL of 500

mM EDTA was then added to each well to detach cells from the culture surface, medium

was pipetted up and down over the culture surface 20 times, and this resuspension

("adherent fraction") was transferred to a separate tube. Cell densities of the collected

fractions were determined by hemocytometer counts, total cell amounts were calculated

by adding the cell numbers for disassociated and adherent fractions for each condition,

and the percent of adherent cells for each condition was then calculated. As a control to

examine cell-substrate adhesion in the absence of agitation, the above protocol was

followed, except the orbital shaker agitation step was omitted.

<u>Time-lapse Microscopy</u>

To image cells by time-lapse transmitted light or fluorescence microscopy, cells were

resuspended to 1 x 10^8 cells/ml in growth medium, and 10 µl of this cell suspension was

added as a spot in the center of a well in a 4-well chambered coverglass slide (Nunc).

After a 1 hour incubation at room temperature to allow the cells to settle, the 10 µl volume

of medium was removed, and 1 ml of growth medium was added to the well, resulting in

a spot of cells in the center of the well. Cells were incubated at room temperature for 24

hours, a field of view with well-spaced cells was located, and then cells were imaged by

time-lapse microscopy using a Nikon Eclipse Ti inverted microscope with NIS-Elements

acquisition software. To image the effect of blebbistatin on individual cells, similar imaging

of cells was done, except that cells were treated with either DMSO or 1 µm bebbistatin

(Sigma B0560) for one hour before imaging. To image aggregates by time-lapse

microscopy, aggregates were generated in ultra-low attachment plates as described

above. During aggregate formation, 4-well chambered coverglass slides (Nunc) were

coated with UltraPure agarose (ThermoFisher) by making a 1% agarose in nuclease-free

water mixture, microwaving to dissolve the agarose, adding 800 µl of molten agarose per

well, and removing the molten agarose after 10 seconds by aspiration. Coated wells were

then allowed to dry at room temperature in a cell culture hood, resulting in a thin coating

of agarose. This coating functions to prevent aggregate adhesion to the glass surface.

After aggregate formation, aggregates were resuspended in the well by gentle pipetting,

and then an 800 µl volume of resuspended aggregates was transferred into one agarose-

coated well in a 4-well chambered coverglass slide. Aggregates were then imaged using

a Zeiss LSM 880 confocal microscope.

<u>RNA-seq</u>

To perform RNA-seq on *Capsaspora* cells, WT or *coYki -/-* cells were collected from

growth flasks by removing growth medium, adding fresh growth medium, and

resuspending cells attached to the flask by pipetting the medium over the surface. Cells

were then diluted to 2×10^5 cells/mL, and for each genotype two 75 cm^2 culture flasks

were prepared by adding 16 mL of culture dilution to each flask. Culture flasks were

incubated at 23°C for 2 days, and then cells were collected by pipetting growth medium

over attached cells. Cells were collected by centrifugation at 2100 x g for 5 minutes, and

then all cells for each genotype were combined, resuspended in 500 µl of growth medium,

and transferred to a microcentrifuge tube. RNA was then prepared with the RNEasy Plus

Mini Kit (Qiagen) using the QIAshredder spin column (Qiagen) to homogenize the lysate.

RNA samples were run on an Agilent Tapestation 4200 to determine level of degradation,

thus ensuring only high quality RNA was used (RIN Score 8 or higher). A Qubit® 4.0

Fluorimeter (ThermoFisher) was used to determine the concentration prior to staring

library prep. One microgram of total DNAse-treated RNA was then prepared with the

TruSeq Stranded mRNA Library Prep Kit (Illumina). Poly-A RNA was purified and

fragmented before strand specific cDNA synthesis. cDNA was then A-tailed and indexed

adapters were ligated. After adapter ligation, samples were PCR-amplified and purified

with AmpureXP beads, then validated again on the Agilent Tapestation 4200. Before

being normalized and pooled, samples were quantified by Qubit then run on an Illumina

NextSeq 500 using V2.5 reagents. Three biological replicates were sequenced for each

genotype, with 22-31 million reads generated per sample. The fastq files were checked

for quality using fastqc (v0.11.2)(*40*) and fastq_screen (v0.4.4)(*41*). Fastq files were

mapped to GCF_000151315.2 reference assembly using STAR (*42*). Read counts were

then generated using featureCounts (*43*). Trimmed Mean of M-values normalization and

differential expression analysis were performed using edgeR (*44*) (false discovery rate

(FDR) <=0.05, absolute log2(fold change) >= 0.5, log(counts per million) >=0).

Phylogenetic trees were downloaded from PhyloDB (*45*) corresponding to *Capsaspora*

(PhyID 101) and pairwise distances for each gene were extracted using an R package

ape (*46*). Closest Human and mouse orthologues were then extracted and used to

annotate the *Capsaspora* genes. These annotated gene names were then used with IPA

(*47*) (QIAGEN Inc., https://www.qiagenbioinformatics.com/products/ingenuity-pathway-

analysis) to get significantly enriched pathways in "Diseases and Bio Functions" category.

To identify functional enrichment in the sets of genes upregulated or downregulated in

coYki -/- cells, DAVID Functional Annotation Tool (*48*) was used to identify functional

categories with FDR < 0.05 using all *Capsaspora* genome genes as the gene population

background.

Statistics

All statistics were done using Prism (GraphPad software, San Diego, CA). Student's t-

test was done to compare differences between two groups, and one-way analysis of

variance (ANOVA) with a post-hoc test was used to compare differences among groups

greater than two.

<u>Gene nomenclature</u>

In the manuscript we use italicized text to refer to a gene (e.g., *coYki)*, unitalicized text

to refer to a protein (e.g., coYki), "-/-" to indicate a homozygous deletion mutant (e.g.,

coYki -/-), and ">" to indicate a promoter driving the expression of a gene (e.g., *EF1 >*

coYki).

Fig. S1. Transfection of *Capsaspora* with genes encoding antibiotic resistance markers results in viable cell populations after drug treatment. Cells were transfected with plasmids encoding mScarlet with no antibiotic marker (pJP71), GenR (pJP72), NatR (pJP102), or HygR (pJP103) genes and were treated with the indicated antibiotics 2 days after transfection for the indicated time period. Representative images

863   of the population are shown. Round, phase-bright cells are visible in populations

864   transfected with resistance markers but not cells transfected with mScarlet.

A

B

C

Fig. S2. Diagnostic PCR of putative coYki deletion clonal cell lines. (A) Diagram of

primer pairs used to evaluate the presence of the WT *coYki* allele (*coYki* WT), a deletion

of a region of *coYki* using the GenR marker (Δ*coYki::GenR*), or a deletion of a region of

coYki using the NatR marker (Δ*coYki::NatR*). TBD, Tead-binding domain; WW, WW

domain; GenR, Geneticin resistance; NatR, nourseothricin resistance. **(B)** WT cells were

transfected with a gene-targeting construct encoding GenR designed to delete the TBD

in the *coYki* gene by homologous recombination (see figure 1G in main text). Following

transfection and drug selection, clonal populations of transfectants were generated and

analyzed by diagnostic PCR using the indicated primer pairs. "*" indicates a clone

showing a PCR product indicative of heterozygous coYki disruption. **(C)** A clonal cell line

with a diagnostic PCR result indicating the disruption of a coYki allele with GenR was

transfected with a gene-targeting construct encoding NatR designed to delete the TBD in

the *coYki* gene. After selection with both Geneticin and nourseothricin, clonal populations

of transfectants were generated and analyzed by diagnostic PCR using the indicated

primer pairs. "*" indicates a clone showing a PCR product indicative of homozygous coYki

disruption.

Fig. S3. WT and *coYki -/-* cells proliferate at a similar rate. (A) To examine proliferation

on a solid substrate, cells were grown in wells in a 24-well plate, and each day cells within

one well were resuspended and counted. **(B)** To examine proliferation in shaking culture,

cells were grown in flasks on an orbital shaker, and an aliquot of cells was collected and

counted daily. Values are mean $\pm$ SEM (n $\geq$ 3). Absence of error bars indicates that error

is smaller than the plot symbol.

Fig. S4. Calcium is required for adhesion in *Capsapsora* cell aggregates.

Capsaspora aggregates were treated with water (vehicle control), the calcium chelator

EGTA, or EGTA with an excess of calcium. At 30 minutes aggregates treated with EGTA

had begun to disassociate, with further disassociation evident at 24 hours. Simultaneous

addition of an excess of calcium blocked this disassociation.

Fig. S5. *coYki -/-* cells adhere to and integrate within WT aggregates like WT cells.

(A-D) Cells expressing mScarlet in the WT background **(A-B)** or in the *coYki -/-* mutant

background **(C-D)** were allowed to aggregate as a homogeneous population **(A, C)** or

were mixed with 90% WT cells and allowed to aggregate **(B, D)**. Individual mScarlet-

labeled *coYki -/-* cells associate with WT cells within aggregates **(D')** in an organization

like that of cells from the WT background labeled with mScarlet **(B')**. Scale bar is 75

microns.

A

B

Fig. S6. coYki affects cell-substrate adhesion but shows no effect on cell-cell

adhesion. (A) To examine cell-cell adhesion, cells in suspension were gently rotated for

an hour to stimulate clump formation through cell-cell adhesion. Cultures were then

examined by hemocytometer and the number of cells per clump was counted. Each red

circle indicates the number of cells in a single clump, and error bars indicate the mean ±

SEM of the average number of cells per clump from 3 independent experiments. Absence

of error bars indicates that error is smaller than the plot symbol. The difference in average

number of cells per clump between WT and *coYki -/-* is not significant (t-test). **(B)** To

examine cell-substrate adhesion, adherent cell cultures were either agitated on a rotary

shaker for 10 minutes or left untreated, and then the number of adhered and unadhered

cells in each culture was counted and the percent of cells adhered to the culture substrate

was calculated. The difference between WT and *coYki -/-* is significant for the percent of

cells attached to substrate after agitation ($p < 0.0001$, t-test).

Movie S1. Aggregation of WT or *coYki -/-* cells over 6 days on a low-adherence

surface. Time units are given in days.

Movie S2. WT and *coYki -/-* cells on a glass surface.

Movie S3. *coYki -/-* cells expressing a *coYki* rescue transgene*.*

Movie S4. Basal region of *Capsaspora* cells expressing Lifeact-mScarlet.

Movie S5. Medial region of WT or *coYki -/-* cells expressing Lifeact-mScarlet.

Movie S6. Actin dynamics in a single *coYki -/-* cell expressing Lifeact-mScarlet.

Movie S7. WT or *coYki -/-* cells expressing Lifeact-mScarlet within an aggregate.

Movie S8. Treatment of *coYki -/-* cells expressing Lifeact-mScarlet with blebbistatin

reduces blebbing.

Data S1. Sequences of synthesized gene fragments used in this study

Data S2. Sequences of oligonucleotides used in this study

Data S3. Plasmids constructed for this study