Bacteriophage cocktails displayed protective effects against soft rot-causing

Pectobacterium **sp.**

Aeron Jade S. Parena[a,b,c*], Benji Brayan I. Silva[b,d], Rae Mark L. Mercado[a,b], Adelbert Adrian

A. Sendon[b], Freddiewebb B. Signabone[e], Johnny F. Balidion[b] and Jaymee R. Encabo[a,*]

[a]*Microbiology Division, Institute of Biological Sciences, College of Arts and Sciences,*

University of the Philippines Los Baños, Los Baños, Laguna 4031, Philippines

[b]*Host-Pathogen Interaction Laboratory, Institute of Weed Science, Entomology and Plant*

Pathology, College of Agriculture and Food Science, University of the Philippines Los Baños,

Los Baños, Laguna 4031, Philippines

[c]*Institute of Microbiology and Immunology, National Yang Ming Chiao Tung University,*

Taipei City 112, Taiwan (R. O. C.)

[d]*International Degree Program in Animal Vaccine Technology, International College,*

National Pingtung University of Science and Technology, Pingtung 912, Taiwan

[e]*National Crop Protection Center, College of Agriculture and Food Science, University of the*

Philippines Los Baños, Los Baños, Laguna 4031, Philippines

*Corresponding Authors:

jrencabo@up.edu.ph

asparena@ym.edu.tw

ORCID:

Aeron Jade S. Parena: 0000-0003-4056-4460

Benji Brayan I. Silva: 0000-0002-1376-3053

Jaymee R. Encabo: 0000-0001-8835-6201

Bacteriophage cocktails displayed protective effects against soft rot-causing

***Pectobacterium* sp.**

Abstract:

Soft rot caused by *Pectobacterium* spp. is responsible for significant losses in vegetable

production worldwide. Methods for the effective control of this disease are limited and are

primarily based on good agricultural practices. The use of phages as biocontrol agents appears

to be a promising alternative to combat phytopathogens. In this study, we investigated the

efficacy of lytic phages against soft rot caused by *Pectobacterium carotovorum* subsp.

carotovorum. Designated as PPc_A3, PPc_D1, and PPc_J3, three bacteriophage isolates,

which were recovered from symptomatic tissues and environmental samples, were observed to

effectively lyse *P. carotovorum* subsp. *carotovorum*. PPc_A3 belongs to the *Podoviridae*

family, while phages PPc_D1 and PPc_J3 belong to the *Myoviridae* family based on the

morphological features of the virions as observed using transmission electron microscopy. The

optimal multiplicity of infection (MOI) differed greatly among the three phages. All survived

incubations at 30°C, 40°C and 50°C and at pH ranging from 3.0 to 9.0, but were all

inactivated at 60°C and at pH 12. Both monophage and cocktail preparations were effective in

inhibiting the growth of *P. carotovorum* subsp. *carotovorum* in *in vitro* challenge tests. In the

semi-*in planta* assays, monophage treatments resulted in significant reduction of tissue

maceration in potato slices, while treatment with cocktail preparations completely inhibited the

development of soft rot disease. Overall, these results demonstrate the efficacy of cocktail

formulations of phages PPc_A3, PPc_D1, and PPc_J3 for the biocontrol of soft rot disease

caused by *P. carotovorum* subsp. *carotovorum.*

Keywords: *Pectobacterium* sp., bacterial soft rot, bacteriophages, biocontrol, agricultural

management

1. Introduction

Bacterial soft rot disease causes one of the most damage among vegetable and fruit crops around the world (Bhat et al., 2010). Annual post-harvest losses due to bacterial soft rot vary from 15-30% of the harvested crops and at least 30-50% of the 65 million tons of vegetables in public markets (Agrios, 2005; Bhat et al., 2010; Rahman et al., 2012). Soft rot disease results from the infections by several types of bacteria, but most commonly by members of the soft rot *Pectobacteriaceae* (SRP) (Adeolu et al., 2016; Agyemang et al., 2020; A. O. Charkowski, 2018; Pérombelon, 2002; Wallace, 1948). During infection, SRPs produce enzymes like pectate lyase, cellulase and polygalacturonase, which are capable of breaking down the plant cell wall leading to plant tissue maceration and decay (A. Charkowski et al., 2012; Li et al., 2019; Mattinen et al., 2007; Muturi et al., 2019). *Pectobacterium* species have the broadest host range among soft rot bacteria, and they are considered as the leading cause of soft rot disease in various vegetable crops (Davidsson et al., 2013; J.-A. Lim et al., 2017).

Reported practices to manageo soft rot disease include cutting-off and disposal of infected tissues during or after harvest, wrapping with paper, treatment with hot water; and/or the use of bactericides, such as alum powder (aluminum potassium sulfate), formaldehyde, copper sulfate, and sodium or calcium hypochlorite (Bayogan et al., 2009; Bhat et al., 2010; R. Czajkowski et al., 2011; Liao et al., 2003). However, these methods only slow down the development of soft rot and do not effectively control the disease. Copper- and antibiotic-resistant phytopathogenic bacterial strains were also reported (Behlau et al., 2011; Cooksey, 1990; McManus et al., 2002; Petriccione et al., 2017; Sundin & Bender, 1993). Due to the limited success of the abovementioned control methods, and the currently growing global threat of antimicrobial resistance, attention has been growing about the use of bacteriophages

or phages (viruses of bacteria) as a biocontrol agents against pathogenic bacteria (Buttimer et

al., 2017; Frampton et al., 2012; J.B. Jones et al., 2007; Jeffrey B. Jones et al., 2012).

Bacteriophages (phages) are viruses that can infect, replicate and lyse bacterial cells, When

environmental conditions are favorable for infection, they can effectively reduce bacterial

populations, especially in the case of lytic phages that undergo rapid replication (Hadas et al.,

1997; Sulakvelidze et al., 2001). Phages generally have high host-specificity, and they have no

direct negative effects on non-host cells. This allows the use of single-phage suspensions or

phage cocktail formulations to target only pathogenic bacteria. Such characteristic suggests the

use of lytic phages to be a promising alternative control method against bacterial soft-rot

(Buttimer et al., 2017; J.-A. Lim et al., 2017). The first *Pectobacterium* spp. phage, My1, was

reported in 2012 and several additional novel phage isolates were subsequently reported

(Buttimer et al., 2018, 2020; Kalischuk et al., 2015; Lee et al., 2012; J.-A. Lim et al., 2014,

2015, 2017; J. A. Lim et al., 2013; Pedersen et al., 2020; Voronina et al., 2019).

In this study, virion morphologies, growth characteristics and stabilities of three lytic

bacteriophages infecting *P. carotovorum* subsp. *carotovorum* were investigated. The lytic

activity and efficacy of the phages to prevent tissue maceration caused by *P. carotovorum*

subsp. *carotovorum* in potato slices using monophage and cocktail suspensions were also

determined. Based on *in vitro* and semi *in planta* assays, we found these lytic phages as

promising and effective candidates for the biocontrol of bacterial soft rot caused by *P.*

carotovorum subsp. *carotovorum*.

2. Materials and Methods

2.1. Bacterial Strains and Media

Pectobacterium carotovorum subsp. *carotovorum* BIOTECH 1752, obtained from the Philippine National Collection of Microorganisms (National Institute of Molecular Biology and Biotechnology, UPLB) was used as the primary host for phage isolation. A collection of 48 *Pectobacterium* spp. isolated from vegetable growing areas and market centers in the Philippines (Host-Pathogen Interaction Laboratory, IWEP, UPLB) and eight other representative strains from other genera (Microbiology Division, IBS, UPLB), were used for the host range assay (Supplementary Table 1). For routine tests and maintenance, bacteria were grown in Tryptone Soya Agar (TSA, HiMedia®) for 16-18 hrs at 30°C. For liquid preparations, bacterial cultures were grown in Tryptone Soya Broth (TSB, HiMedia®) for 12-18 hrs at 30°C with 125 rpm agitation.

2.2. Bacteriophage Isolation and Characterization

2.2.1. Isolation and Purification of Bacteriophages

Bacteriophages were isolated from rhizosphere soil, symptomatic tissues, and sewage water as described by Czajkowski et al. (2014). Twenty grams (or ml) of each sample was homogenized in 20 ml saline-magnesium (SM) buffer ($MgSO_4$, NaCl, $CaCl_2$, and Tris-HCl, pH 7.5), and incubated at 30°C for 12-16 hrs. Five hundred μl of log phase *P. carotovorum* subsp. *carotovorum* culture was added to each enrichment set-up and was incubated as above. Enrichment cultures were centrifuged at 3,000 rpm for 15 minutes and the supernatants were chloroform-treated in 1:20 chloroform/supernatant ratio for at least 10 min. The resulting lysates were screened for the presence of bacteriophages by spot test assay (Mazzocco et al., 2009). Overlay mixtures, containing 4.5 ml soft agar (TSA), 0.5 ml 20 mM $CaCl_2$, and 300 μl of log phase *P. carotovorum* subsp. *carotovorum* culture were poured on Petri plates containing pre-solidified TSA. Five μl of the phage lysates were spotted on the overlay and the spots were absorbed into the agar for 10-15 min prior to incubation of plates at 30°C for 12-16 hrs. Zones of lysis on the bacterial lawn indicated the presence of phages.

Phage purification was done using the double layer plaque assay (Kropinski et al., 2009). Ten-
fold dilutions were prepared from each positive lysate. In separate sterile 1.5-ml
microcentrifuge tubes, 300 μl of diluted lysate suspensions were mixed with 300 μl log phase
culture of *P. carotovorum* subsp. *carotovorum* in TSB. These were pre-incubated for 10 min at
4°C to allow adsorption of the phages. Molten overlay (4.5 ml) mixed with 0.5 ml 20 mM $CaCl_2$
was added to the suspensions and the resulting mixtures were poured on to Petri plates
containing pre-solidified TSA. The overlays were allowed to solidify for 15-20 min and the
plates were incubated at 30°C for 12-16 hrs. Well-isolated plaques were picked using sterile
pipette tips and the phages were suspended in 500 μl SM buffer. Bacteriophage isolates were
purified by successive single plaque isolation for at least five times or until uniform plaque
morphology was observed.

2.2.2. Plaque Formation Using Double Layer Plaque Assay
Plaque morphologies were characterized after a series of double layer plaque assays as
described above (Jurczak-Kurek et al., 2016). The transparency and morphology of the plaques
formed by each bacteriophage isolate was described and plaque diameter was measured using
a digital caliper.

2.2.3. Bacteriophage Lysate Preparation and Titration
Double layer plaque assay plates with pure phage plaques were flooded with 3-5 ml of SM
buffer to obtain the bacteriophage lysate. Plates were gently shaken for 15-20 min and the
overlay-SM buffer mixtures were collected and centrifuged at 8,000 rpm for 10 minutes. The
supernatants were chloroform treated in 1:20 chloroform/lysate ratio and were maintained at
4°C until titration. Lysates were subjected to double layer plaque assay (or spot test) for titration
as described above. Phage titer was determined and expressed as plaque forming units (PFU
ml^{-1}).

2.2.4. Determination of Virion Morphology Using Transmission Electron Microscopy (TEM)

High titer lysates (at least 10^{10} PFU ml^{-1}) of the three phage isolates were submitted to the

Electron Microscopy Laboratory of the Research Institute for Tropical Medicine (Muntinlupa

City, Philippines). The phage suspension was placed on carbon-coated copper grids, and 2%

aqueous uranyl acetate was added for three minutes to negatively stain the virions. Phages were

visualized by transmission electron microscopy (TEM; JEOLJEM-1220 Electron Microscope),

and the images were scanned with 800 ms exposure, 1.4 gain, and 1.0 bin with no sharpening,

normal contrast, and 1.00 gamma.

2.2.5. Optimal Multiplicity of Infection (MOI)

The optimal ratio of phage particles to the number of bacterial cells, also called the optimal

multiplicity of infection (MOI), was determined for the three phage isolates. *P. carotovorum*

subsp. *carotovorum* was infected with each phage isolate at four different PFU ml^{-1}/cell ml^{-1}

ratios (MOI): 0.01, 0.1, 1.0, and 10. Following overnight incubation at 30 °C, the suspensions

were chloroform-treated for 10 minutes (1:20) and were centrifuged at 8,000 rpm for 10

minutes. The supernatants were subjected to double layer plaque assay to determine phage titer.

The MOI demonstrating the highest phage titer or PFU ml^{-1} was considered optimal.

2.2.6. Latent Period and Phage Burst Size

One-step growth experiments were conducted to determine phage latent period and burst size

following previously described protocol (R. Czajkowski et al., 2014). Cells of *P. carotovorum*

subsp. *carotovorum* at log phase were harvested by centrifugation at 12,000 rpm for 15 min at

10°C, and were resuspended in TSB to an OD$_{600}$ of 0.5 (c. 5 x 10^8 CFU mL^{-1}). Subsequently,

100 μl of 20mM CaCl$_2$ and 100 μl of the phage isolate were added into the bacterial culture at

a MOI of 0.1 (10^7 PFU mL^{-1}/ 10^8 CFU mL^{-1}). The phages were allowed to adsorb by incubation

at 4°C for 10 min and the suspensions were further incubated at 30°C for 10 min. The

suspensions were centrifuged at 12,000 rpm for 15 min at 10°C to remove unadsorbed phages.

The resulting pellet was resuspended in 10 ml fresh TSB. The mixtures were incubated at 30°C

for 60 min with shaking at 125 rpm. Aliquots were taken from the mixtures every 10 min and

the titer was determined using the Spot Test method. The burst size was calculated using the

formula: $burst\ size = \frac{PFU_{final} - PFU_{initial}}{PFU_{initial}}$ and the experiment was repeated twice with the

same set-up and the results were averaged.

2.2.7. Adsorption Rate

Phage adsorption rate was determined using previously described protocol (R. Czajkowski et

al., 2014). One ml of *P. carotovorum* subsp. *carotovorum* (ca. 10^8 cfu mL^{-1}) was infected with

each phage isolate by adding 100 μl of phage lysate (ca. 10^8 pfu mL^{-1}) to reach MOI 0.1 and

the resulting suspensions were incubated at 30°C for 20 min. After 0 (control), 1, 2, 4, 10 and

20 min, 50 μl aliquots were taken from each suspension and added to 450 μl TSB. Ten-fold

dilutions in SM buffer were prepared and the titer of the unadsorbed phages were determined

using the Spot Test method as described above. The experiment was repeated twice with the

same set-up and the results were averaged. Phage adsorption was calculated as follows:

$\%adsorption = \frac{control\ titer - residual\ titer}{residual\ titer} x100.$

2.2.8. Thermal Resistance and pH Stability

Thermal resistance of each of the three phages was determined at 30, 40, 50 and 60 °C. Phages

(10^8 PFU ml^{-1}) in SM buffer were incubated for 1 hr at each temperature condition and the

surviving PFU was determined using double layer plaque assay (R. Czajkowski et al., 2014).

The effect of pH on phage stability was determined using phages (10^8 PFU ml^{-1}) in SM buffer

within a pH range of 3.0 to 12.0. The experiment was carried out by incubating the phage

suspensions at RT for 16-24 hrs and the surviving PFU was determined using double layer

plaque assay.

2.2.9. Host Range Determination

The host range of the three phage isolates was determined using a collection of 48

Pectobacterium spp. isolates and eight other representative strains from different genera

(Supplementary Table 1). *Pectobacterium* spp. were isolated in the years 2017-2019 from

symptomatic tissues of various vegetable crops.

For the spot test assay, bacterial cultures were grown in TSB at 30°C for 12-16 hrs with agitation

at 125 rpm. From the resulting bacterial cultures, 300 μl of each was inoculated into the overlay

agar and was subsequently poured into a prepared TSA plate and allowed to solidify at room

temperature for 30 min. Five μl of the phage lysate (10^9 PFU ml^{-1}) was spotted onto the top agar

layer, and the plates were allowed to dry at room temperature for 10-15 min. Following

overnight incubation at 30°C, plates were inspected for lysis zones.

2.3. Lytic Activity and Biocontrol Assays

2.3.1. In Vitro Challenge Test

Bacterial inactivation using monophage (PPc_A3, PPc_D1, and PPc_J3) and cocktail

treatments (PPc_A3+D1, PPc_A3+J3, PPc_D1+J3, and PPc_A3+D1+J3) was determined

following the method described by Chen et al. (2018). For monophage treatments, log phase

culture of *P. carotovorum* subsp. *carotovorum* (OD_{600} of 0.5, c. 5 x 10^8 cfu mL^{-1}) was mixed

with the phage lysate at optimal MOI. For cocktail preparations, low (0.001) and high (100)

MOI were used for the experiment. The bacteria-phage mixtures were incubated at 30°C with

shaking at 125 rpm and the OD readings at 600nm were recorded (SPECTROstar^{Nano}) at 15-

minute intervals for eight hrs to monitor the changes in bacterial density.

2.3.2 Semi-In Planta Assays

2.3.2.1 Phage Biocontrol Using Monophage and Cocktail Treatments on Tissue Slices

The efficacy of monophage and cocktail treatments against soft rot caused by *P. carotovorum*

subsp. *carotovorum* was evaluated using a potato slice assay (Robert Czajkowski et al., 2015).

Potato tubers were surface sterilized by soaking in 70% ethanol for 10 minutes, washed with

sterile distilled water, and air dried under the laminar flow hood. The tubers were cut into 7.0

mm-thick transverse disks using sterile knives and three wells (5-mm diameter) were made on

each slice using a sterile cork borer. For monophage treatments, log phase culture of *P.*

carotovorum subsp. carotovorum was mixed with the phage lysates at optimal MOI. For

cocktail preparations, low (0.001) and high (100) MOIs were used for the experiment. Three

potato slices were used for each treatment. Wells were filled with 20 µl of the bacteria-phage

mixtures, 20 µl of sterile SM buffer as negative control, and 20 µl of *P. carotovorum* subsp.

carotovorum (10^8 cells ml^{-1}) as positive control. The diameter of the rotten tissues around the

wells inoculated with bacteria and wells co-inoculated with bacteria and phages were measured

after incubation at 30°C for 72 hrs. Treatment means were compared using one-way ANOVA

with Tukey's post-test ($\alpha = 0.05$).

3. Results

3.1. Isolation and Characterization of Bacteriophages

3.1.1. Isolation of Lytic Bacteriophages

Bacteriophages PPc_A3, PPc_D1 and PPc_J3 were isolated from rhizosphere soil, symptomatic tissues and sewage water, respectively, using *P. carotovorum* subsp. *carotovorum.* All three phages formed clear plaques typical for lytic phages, with varying sizes (Table 1).

3.1.2. Virion Morphology

Examination using TEM revealed that the three phages belong to families *Podoviridae* and *Myoviridae* of Order *Caudovirales*: (Table 1, Figure 1). Phage PPc_A3 belongs to the *Podoviridae* family, with an icosahedral head of 80 nm and short non-contractile tail of 20 nm (Table 1). Phages PPc_D1 and PPc_J3 are members of the *Myoviridae* family, with icosahedral heads (75 nm) and rigid contractile tails (95-135 nm) (Table 1, Figure 1). Based on the proposed unified phage naming, the phages should be formally named vB_PcaP-A3, vB_PcaM-D1 and vB_PcaM-J3 (Kropinski et al., 2009) .

3.1.3. Characterization of Bacteriophages

All three phages exhibited strong adsorption to *P. carotovorum* subsp. *carotovorum,* and approximately 90% of the phage particles adsorbed to the bacteria within 5 minutes (Figure 2). PPc_A3 and PPc_D1 showed similar infection kinetics with a latent period of 10 min and a burst size of 142-151 progeny phages/cell, whereas PPc_J3 showed a longer latent period of 20 min and a smaller burst size of 42 progeny phages/cell (Table 1, Figure 2A & 2B). Among the three phages, PPc_A3 has the lowest optimal MOI (0.01), followed by PPc_D1 (1.0) and PPc_J3 (10.0) (Table 1). All of the three phages retained infectivity at 30°C, 40°C, and 50°C, and in the pH range 3-9 (Figure 2C).

 3.1.4.   Host Range

Spot test assay revealed that the three phages showed antibacterial activity specific for *P.*

carotovorum subsp. *carotovorum.* None of the three phages was able to lyse the 48

Pectobacterium spp. isolates and the other eight representative strains from other genera.

3.2. Lytic Activity and Biocontrol Assays

3.2.1.   Phage Activity In Vitro

Killing curves established for PPc_A3, PPc_D1 and PPc_J3 showed that all three phages were

highly effective against *P. carotovorum* subsp. *carotovorum* at optimal MOI (Figure 3A). The

bacterial cell density remained less than 0.05 (OD_{600}) throughout the 8-hr observation period.

The absence of any increase in OD_{600} reading also suggests that there is no observed phage

resistance development of the host at least in the period of observation (Muturi et al., 2019).

These results indicate that monophage treatments of PPc_A3, PPc_D1 and PPc_J3 based on

their optimal MOIs were equally effective in controlling *P. carotovorum* subsp. *carotovorum*

in vitro.

For the cocktail treatments, *in vitro* challenge tests were performed using low MOI (0.001) and

high MOI (100). Each phage was used at the same concentration and the phage cocktails

consisting of two or three phages were combined and evaluated. At MOI of 0.001, the killing

curves of the phage cocktails composed of two or three phages maintained its bacterial

inactivation with an $OD_{600} < 0.05$ for 8 hrs (Figure 4A). It appears that a phage cocktail

containing any of the three phages had the ability to control bacterial growth at low MOI during

the 8-hr observation period. At MOI 100, killing curves showed that all cocktail treatments

were highly effective against *P. carotovorum* subsp. *carotovorum* (Figure 4A). The bacterial

cell density remained less than 0.05 (OD_{600}) throughout the duration of the experiment.

3.2.2.   Phage Biocontrol Using Monophage and Cocktail Treatments on Tissue Slices

The average diameter of rotting tissue was significantly lower in all monophage treatments than

in the set-up inoculated with *P. carotovorum* subsp. *carotovorum* only (Figure 3B-D). The set-

up co-inoculated with *P. carotovorum* subsp. *carotovorum* and PPc_A3 exhibited the lowest

diameter of rotting tissue among the three treatments (Figure 3B-D). Phage PPc_A3 was able

to reduce potato tuber tissue maceration to at least 60%, while phages PPc_D1 and PPc_J3 were

able to reduce tissue maceration to at least 50% of that observed for the control potato slices

inoculated with *P. carotovorum* subsp. *carotovorum* only (Figure 3B-D). All of the phage

cocktail treatments completely prevented tissue maceration at low and high MOI (Figure 4B-

D).

4. Discussion

In this study, three lytic bacteriophages infecting *P. carotovorum* subsp. *carotovorum* designated as PPc_A3, PPc_D1, and PPc_J3, were isolated from various environmental samples in the Philippines such as rhizosphere soil, symptomatic tissues, and sewage water. We examined their morphological characteristics and found that PPc_A3 belongs to *Family Podoviridae* similar to phages PP1 and PP2 (J.-A. Lim et al., 2017; J. A. Lim et al., 2013). PPc_D1 and PPc_J3 are members of the *Family Myoviridae* like phages PM1 and PM2 (J.-A. Lim et al., 2014, 2015).

The latent period and burst size of each phage were also determined based on their one-step growth curves, as well as their individual adsorption rates, optimal MOI, and stability on varying temperature and pH levels. PPc_A3 has the lowest optimal MOI of 0.01. It has a latent period of 10 min and a calculated burst size value of 142 PFU/cell. PPc_D1 has an optimal MOI of 0.1, latent period of 10 min and a burst size of 151 PFU/cell. PPc_J3 has a high optimal MOI of 10, longest latent period of 30 min, and the lowest calculated burst size of 42 PFU/cell. Since phages displaying shorter latent periods and higher burst sizes are reported to provide better efficiency than phages with longer latent periods in cultures where the density of the host is high, PPc_J3 showed less promising growth characteristics compared to PPc_A3 and PPc_D1 (Abedon et al., 2003). All three, however, showed strong adsorption to when about 90% of the phages were adsorbed within 5 min after incubation. When their stabilities on varying temperature and pH levels were tested, all three survived incubations at 30°C-50°C and at pH 3-9 but were inactivated at 60°C and at pH 12. These are important environmental factors in biological control applications. Since temperatures under field conditions do not exceed 50°C and pH levels usually range from pH 3.3-4.3, results indicate that PPc_A3, PPc_D1 and PPc_J3 can be suitable for future field applications (Navarrete et al., 2017).

All three phage isolates also showed specificity to their isolation host, *P. carotovorum* subsp. *carotovorum.* Phages are generally generally known to have narrow host ranges, typically limited to strains within a certain species of their bacterial host (Ross et al., 2016). This can allow the formulation

of phage cocktails targeting bacterial species of interest only, which may be a certain bacterial plant

pathogen or a specific bacterium in a microbial community that negatively impacts plant growth or

health conditions. Developing a phage-based biocontrol product that can eliminate multiple genera, or

even different related species could be challenging,a but combining phages specific to target hosts may

overcome this problem (Buttimer et al., 2017; Frampton et al., 2012). The specificity of phages also

reduces the risk of harming the natural microbiota, especially those non-target bacteria that are

potentially beneficial, and limits the emergence of phage-resistant bacteria (Loc-Carrillo & Abedon,

2011).

Additionally, the efficacy of the three phage isolates, in both single-phage and cocktail suspensions

(combinations of two and three phages), were assessed through *in vitro* and semi-*in plant*a bacterial

challenge tests, and we found that both single phages and phage cocktails completely inactivate the

growth of *P. carotovorum* subsp. *carotovorum*. However, phages in cocktail suspensions were more

effective over individual application in terms of their protective effect against soft rot by *P. carotovorum*

subsp. *carotovorum* as indicated by the complete maceration reduction in both MOI (0.001 and 100)

tested in the study. Recent reports of phage cocktail formulations against *P. carotovorum* subsp.

carotovorum have also described efficient inhibition of bacterial growth both *in vitro* and *in vivo*

(Carstens et al., 2019; Muturi et al., 2019; Zaczek-Moczydłowska et al., 2020). Having a higher MOI

has also been reported to increase the chance for the bacteriophages to infect all host cells, effectively

killing all host cells after one round of replication (Alves et al., 2016; Thomas, 2001).

However, the use of higher MOI could also possibly increase the emergencebacterial variants that are

resistant to these phages(Chen et al., 2018). This is due to the rapid evolution of the bacterial population

due to the applied selection pressure that favors the propagation of phage-resistant clones over phage-

sensitive clones (Castillo et al., 2015; Chen et al., 2018; Hyman & Abedon, 2010; Middelboe, 2000;

Middelboe et al., 2001). This implies that the use of heavy phage loads is an important selection factor

that should be taken into account in phage therapy and that lower MOIs should also be used when

testing protective effect and bacterial inactivation by phages, particularly in actual plant tissues (Chen

et al., 2018). To this end, further experimental studies to optimize the phage treatment parameters and to identify the minimum effective phage number for potential field and post-harvest application should be conducted.

Data statement

All datasets presented in this paper are included in the article.

CRediT authorship contribution statement

Aeron Jade S. Parena: Methodology, Validation, Formal Analysis, Investigation, Data Curation, Writing-Original Draft, Writing-Review & Editing, Visualization. **Benji Brayan I. Silva:** Conceptualization, Methodology, Validation, Formal Analysis, Writing-Review & Editing. **Rae Mark L. Mercado:** Methodology, Validation, Investigation, Data Curation. **Adrian Adelbert A. Sendon:** Methodology, Validation, Investigation. **Freddiewebb B. Signabon:** Methodology, Validation, Investigation. **Johnny F. Balidion:** Conceptualization, Supervision, Project Administration, Funding Acquisition. **Jaymee R. Encabo:** Conceptualization, Data Curation, Writing-Review & Editing, Supervision, Project administration, Funding acquisition.

Funding

This work was supported by the Department of Science and Technology-Philippine Council for Agriculture, Aquatic and Natural Resources Research and Development (DOST-PCAARRD), Project Code N91312A.

Declaration of competing interest

The authors declare that the research was conducted with no known competing financial interests or personal relationships that could be construed as a potential conflict of interest.

Acknowledgements

We thank Ms. Janren Sarah T. Macaraig and Mr. Benjamin V. Cunanan for the technical assistance with phage lysate preparations, and Mr. Aristeo P. Averion and Envirokonsult Equipment and Services, Inc. for the sewage water samples used for bacteriophage isolation.

References

Abedon, S. T., Hyman, P., & Thomas, C. (2003). Experimental Examination of Bacteriophage Latent-Period Evolution as a Response to Bacterial Availability. *Applied and Environmental Microbiology*, *69*(12). https://doi.org/10.1128/AEM.69.12.7499-7506.2003

Adeolu, M., Alnajar, S., Naushad, S., & Gupta, R. S. (2016). Genome-based phylogeny and taxonomy of the '*Enterobacteriales*': proposal for *Enterobacteriales* ord. nov. divided into the families *Enterobacteriaceae*, *Erwiniaceae* fam. nov., *Pectobacteriaceae* fam. nov., *Yersiniaceae* fam. nov., *Hafniaceae* fam. nov., *Morganellaceae* fam. nov., and *Budviciaceae* fam. nov. *International Journal of Systematic and Evolutionary Microbiology*, *66*(12), 5575–5599. https://doi.org/10.1099/ijsem.0.001485

Agrios, G. N. (2005). Plant Diseases Caused by Prokaryotes: Bacteria and Mollicutes. In G. N. AGRIOS (Ed.), *Plant Pathology (Fifth Edition)* (Fifth Edition, pp. 615–703). Academic Press. https://doi.org/10.1016/B978-0-08-047378-9.50018-X

Agyemang, P. A., Kabir, M. N., Kersey, C. M., & Dumenyo, C. K. (2020). The bacterial soft rot pathogens, *Pectobacterium carotovorum* and *P. atrosepticum*, respond to different classes of virulence-inducing host chemical signals. *Horticulturae*, *6*(1). https://doi.org/10.3390/horticulturae6010013

Alves, D. R., Perez-Esteban, P., Kot, W., Bean, J. E., Arnot, T., Hansen, L. H., Enright, M. C., & Jenkins, A. T. A. (2016). A novel bacteriophage cocktail reduces and disperses *Pseudomonas aeruginosa* biofilms under static and flow conditions. *Microbial Biotechnology*, *9*(1). https://doi.org/10.1111/1751-7915.12316

Bayogan, E. R. V, Jimenez, E. F., Boteng, J. D., Bautista, O. K., & Macario, C. B. (2009). Improving

Postharvest Cabbage (*Brassica oleracea* L. var. *capitata*) Quality Using Alum and Newspaper

Wrap. *BANWA: A Multidisciplinary Journal*, *1774-2016–141617*, 11.

https://doi.org/10.22004/ag.econ.145554

Behlau, F., Canteros, B. I., Minsavage, G. V., Jones, J. B., & Graham, J. H. (2011). Molecular

Characterization of Copper Resistance Genes from *Xanthomonas citri* subsp. *citri* and

Xanthomonas alfalfae subsp. *citrumelonis*. *Applied and Environmental Microbiology*, *77*(12).

https://doi.org/10.1128/AEM.03043-10

Bhat, K. A., Masood, S. D., Bhat, N. A., Bhat, M. A., Razvi, S. M., Mir, M. R., Akhtar, S., Wani, N.,

& Habib, M. (2010). Current Status of Post-Harvest Soft Rot in Vegetables: A Review. *Asian*

Journal of Plant Sciences, *9*(4), 200–208. https://doi.org/10.3923/ajps.2010.200.208

Buttimer, C., Lucid, A., Neve, H., Franz, C., O'Mahony, J., Turner, D., Lavigne, R., & Coffey, A.

(2018). *Pectobacterium atrosepticum* Phage vB_PatP_CB5: A Member of the Proposed Genus

'Phimunavirus.' *Viruses*, *10*(8). https://doi.org/10.3390/v10080394

Buttimer, C., Lynch, C., Hendrix, H., Neve, H., Noben, J.-P., Lavigne, R., & Coffey, A. (2020).

Isolation and Characterization of *Pectobacterium* Phage vB_PatM_CB7: New Insights into the

Genus Certrevirus. *Antibiotics*, *9*(6). https://doi.org/10.3390/antibiotics9060352

Buttimer, C., McAuliffe, O., Ross, R. P., Hill, C., O'Mahony, J., & Coffey, A. (2017). Bacteriophages

and bacterial plant diseases. In *Frontiers in Microbiology* (Vol. 8, Issue JAN). Frontiers Media

S.A. https://doi.org/10.3389/fmicb.2017.00034

Carstens, A. B., Djurhuus, A. M., Kot, W., & Hansen, L. H. (2019). A novel six-phage cocktail reduces

Pectobacterium atrosepticum soft rot infection in potato tubers under simulated storage

conditions. *FEMS Microbiology Letters*, *366*(9). https://doi.org/10.1093/femsle/fnz101

Castillo, D., Christiansen, R. H., Dalsgaard, I., Madsen, L., & Middelboe, M. (2015). Bacteriophage

Resistance Mechanisms in the Fish Pathogen *Flavobacterium psychrophilum*: Linking Genomic

Mutations to Changes in Bacterial Virulence Factors. *Applied and Environmental Microbiology*,

81(3), 1157–1167. https://doi.org/10.1128/AEM.03699-14

Charkowski, A., Blanco, C., Condemine, G., Expert, D., Franza, T., Hayes, C., Hugouvieux-Cotte-

Pattat, N., Solanilla, E. L., Low, D., Moleleki, L., Pirhonen, M., Pitman, A., Perna, N., Reverchon,

S., Palenzuela, P. R., Francisco, M. S., Toth, I., Tsuyumu, S., Van Der Waals, J., … Yedidia, I.

(2012). The role of secretion systems and small molecules in soft-rot *Enterobacteriaceae*

pathogenicity. In *Annual Review of Phytopathology* (Vol. 50, pp. 425–449).

https://doi.org/10.1146/annurev-phyto-081211-173013

Charkowski, A. O. (2018). The Changing Face of Bacterial Soft-Rot Diseases. *Annual Review of*

Phytopathology, *56*(1). https://doi.org/10.1146/annurev-phyto-080417-045906

Chen, L., Yuan, S., Liu, Q., Mai, G., Yang, J., Deng, D., Zhang, B., Liu, C., & Ma, Y. (2018). In Vitro

Design and Evaluation of Phage Cocktails Against *Aeromonas salmonicida*. *Frontiers in*

Microbiology, *9*. https://doi.org/10.3389/fmicb.2018.01476

Cooksey, D. A. (1990). Genetics of Bactericide Resistance in Plant Pathogenic Bacteria. *Annual Review*

of Phytopathology, *28*(1). https://doi.org/10.1146/annurev.py.28.090190.001221

Czajkowski, R., Ozymko, Z., & Lojkowska, E. (2014). Isolation and characterization of novel soilborne

lytic bacteriophages infecting *Dickeya* spp. biovar 3 ('D. solani'). *Plant Pathology*, *63*(4).

https://doi.org/10.1111/ppa.12157

Czajkowski, R., Pérombelon, M. C. M., van Veen, J. A., & van der Wolf, J. M. (2011). Control of

blackleg and tuber soft rot of potato caused by *Pectobacterium* and *Dickeya* species: a review.

Plant Pathology, *60*(6). https://doi.org/10.1111/j.1365-3059.2011.02470.x

Czajkowski, Robert, Ozymko, Z., de Jager, V., Siwinska, J., Smolarska, A., Ossowicki, A., Narajczyk,

M., & Lojkowska, E. (2015). Genomic, Proteomic and Morphological Characterization of Two

Novel Broad Host Lytic Bacteriophages ΦPD10.3 and ΦPD23.1 Infecting Pectinolytic

Pectobacterium spp. and *Dickeya* spp. *PLOS ONE*, *10*(3).

https://doi.org/10.1371/journal.pone.0119812

Davidsson, P. R., Kariola, T., Niemi, O., & Tapio Palva, E. (2013). Pathogenicity of and plant immunity

to soft rot pectobacteria. In *Frontiers in Plant Science* (Vol. 4, Issue JUN). Frontiers Research

Foundation. https://doi.org/10.3389/fpls.2013.00191

Frampton, R. A., Pitman, A. R., & Fineran, P. C. (2012). Advances in bacteriophage-mediated control

of plant pathogens. In *International Journal of Microbiology*.

https://doi.org/10.1155/2012/326452

Hadas, H., Einav, M., Fishov, I., & Zaritsky, A. (1997). Bacteriophage T4 Development Depends on

the Physiology of its Host *Escherichia coli*. *Microbiology*, *143*(1), 179–185.

https://doi.org/10.1099/00221287-143-1-179

Hyman, P., & Abedon, S. T. (2010). *Bacteriophage Host Range and Bacterial Resistance* (pp. 217–

248). https://doi.org/10.1016/S0065-2164(10)70007-1

Jones, J.B., Jackson, L. E., Balogh, B., Obradovic, A., Iriarte, F. B., & Momol, M. T. (2007).

Bacteriophages for Plant Disease Control. *Annual Review of Phytopathology*, *45*(1), 245–262.

https://doi.org/10.1146/annurev.phyto.45.062806.094411

Jones, Jeffrey B., Vallad, G. E., Iriarte, F. B., Obradović, A., Wernsing, M. H., Jackson, L. E., Balogh,

B., Hong, J. C., & Momol, M. T. (2012). Considerations for using bacteriophages for plant disease

control. *Bacteriophage*, *2*(4), e23857. https://doi.org/10.4161/bact.23857

Jurczak-Kurek, A., Gąsior, T., Nejman-Faleńczyk, B., Bloch, S., Dydecka, A., Topka, G., Necel, A.,

Jakubowska-Deredas, M., Narajczyk, M., Richert, M., Mieszkowska, A., Wróbel, B., Węgrzyn,

G., & Węgrzyn, A. (2016). Biodiversity of bacteriophages: morphological and biological

properties of a large group of phages isolated from urban sewage. *Scientific Reports*, *6*(1).

https://doi.org/10.1038/srep34338

Kalischuk, M., Hachey, J., & Kawchuk, L. (2015). Complete Genome Sequence of Phytopathogenic

Pectobacterium atrosepticum Bacteriophage Peat1. *Genome Announcements*, *3*(4).

https://doi.org/10.1128/genomeA.00760-15

Kropinski, A. M., Mazzocco, A., Waddell, T. E., Lingohr, E., & Johnson, R. P. (2009). *Enumeration of*

Bacteriophages by Double Agar Overlay Plaque Assay. https://doi.org/10.1007/978-1-60327-

164-6_7

Lee, J.-H., Shin, H., Ji, S., Malhotra, S., Kumar, M., Ryu, S., & Heu, S. (2012). Complete Genome

Sequence of Phytopathogenic *Pectobacterium carotovorum* subsp. *carotovorum* Bacteriophage

PP1. *Journal of Virology*, *86*(16). https://doi.org/10.1128/JVI.01283-12

Li, X., Fu, L., Chen, C., Sun, W., Tian, Y., & Xie, H. (2019). Characteristics and Rapid Diagnosis of

Pectobacterium carotovorum ssp. Associated With Bacterial Soft Rot of Vegetables in China.

Plant Disease, *104*(4), 1158–1166. https://doi.org/10.1094/PDIS-05-19-1033-RE

Liao, C.-H., McEvoy, J., & Smith, J. (2003). Control of bacterial soft rot and foodborne human

pathogens on fresh fruits and vegetables. *Advances in Plant Disease Management*.

Lim, J.-A., Heu, S., Park, J., & Roh, E. (2017). Genomic characterization of bacteriophage

vB_PcaP_PP2 infecting *Pectobacterium carotovorum* subsp. *carotovorum*, a new member of a

proposed genus in the subfamily *Autographivirinae*. *Archives of Virology*, *162*(8).

https://doi.org/10.1007/s00705-017-3349-6

Lim, J.-A., Lee, D. H., & Heu, S. (2015). Isolation and Genomic Characterization of the T4-Like

Bacteriophage PM2 Infecting *Pectobacterium carotovorum* subsp. *carotovorum*. *The Plant*

Pathology Journal, *31*(1). https://doi.org/10.5423/PPJ.NT.09.2014.0099

Lim, J.-A., Shin, H., Lee, D. H., Han, S.-W., Lee, J.-H., Ryu, S., & Heu, S. (2014). Complete genome

sequence of the *Pectobacterium carotovorum* subsp. *carotovorum* virulent bacteriophage PM1.

Archives of Virology, *159*(8). https://doi.org/10.1007/s00705-014-2005-7

Lim, J. A., Jee, S., Lee, D. H., Roh, E., Jung, K., Oh, C., & Heu, S. (2013). Biocontrol of *Pectobacterium*

carotovorum subsp. *carotovorum* using bacteriophage PP1. *Journal of Microbiology and*

Biotechnology, *23*(8), 1147–1153. https://doi.org/10.4014/jmb.1304.04001

Loc-Carrillo, C., & Abedon, S. T. (2011). Pros and cons of phage therapy. *Bacteriophage*, *1*(2).

https://doi.org/10.4161/bact.1.2.14590

Mattinen, L., Nissinen, R., Riipi, T., Kalkkinen, N., & Pirhonen, M. (2007). Host-extract induced

changes in the secretome of the plant pathogenic bacterium *Pectobacterium atrosepticum*.

Proteomics, *7*(19), 3527–3537. https://doi.org/10.1002/pmic.200600759

Mazzocco, A., Waddell, T. E., Lingohr, E., & Johnson, R. P. (2009). *Enumeration of Bacteriophages*

Using the Small Drop Plaque Assay System. https://doi.org/10.1007/978-1-60327-164-6_9

McManus, P. S., Stockwell, V. O., Sundin, G. W., & Jones, A. L. (2002). Antibiotic use in plant

agriculture. *Annual Review of Phytopathology*, *40*(1).

https://doi.org/10.1146/annurev.phyto.40.120301.093927

Middelboe, M. (2000). Bacterial Growth Rate and Marine Virus–Host Dynamics. *Microbial Ecology*,

40(2), 114–124. https://doi.org/10.1007/s002480000050

Middelboe, M., Hagström, A., Blackburn, N., Sinn, B., Fischer, U., Borch, N. H., Pinhassi, J., Simu,

K., & Lorenz, M. G. (2001). Effects of Bacteriophages on the Population Dynamics of Four

Strains of Pelagic Marine Bacteria. *Microbial Ecology*, *42*(3), 395–406.

https://doi.org/10.1007/s00248-001-0012-1

Muturi, P., Yu, J., Maina, A. N., Kariuki, S., Mwaura, F. B., & Wei, H. (2019). Bacteriophages Isolated

in China for the Control of *Pectobacterium carotovorum* Causing Potato Soft Rot in Kenya.

Virologica Sinica, *34*(3), 287–294. https://doi.org/10.1007/s12250-019-00091-7

Navarrete, I. A., Gabiana, C. C., Dumo, J. R. E., Salmo, S. G., Guzman, M. A. L. G., Valera, N. S., &

Espiritu, E. Q. (2017). Heavy metal concentrations in soils and vegetation in urban areas of

Quezon City, Philippines. *Environmental Monitoring and Assessment*, *189*(4).

https://doi.org/10.1007/s10661-017-5849-y

Pedersen, J. S., Carstens, A. B., Djurhuus, A. M., Kot, W., Neve, H., & Hansen, L. H. (2020).

Pectobacterium Phage Jarilo Displays Broad Host Range and Represents a Novel Genus of

Bacteriophages Within the Family *Autographiviridae*. *PHAGE*, *1*(4).

https://doi.org/10.1089/phage.2020.0037

Pérombelon, M. C. M. (2002). Potato diseases caused by soft rot erwinias: an overview of pathogenesis.

Plant Pathology, *51*(1), 1–12. https://doi.org/10.1046/j.0032-0862.2001.Shorttitle.doc.x

Petriccione, M., Zampella, L., Mastrobuoni, F., & Scortichini, M. (2017). Occurrence of copper-

resistant *Pseudomonas syringae* pv. *syringae* strains isolated from rain and kiwifruit orchards also

infected by P. s. pv. actinidiae. *European Journal of Plant Pathology*, *149*(4).

https://doi.org/10.1007/s10658-017-1246-1

Rahman, M. M., Ali, M. E., Khan, A. A., Akanda, A. M., Uddin, M. K., Hashim, U., & Abd Hamid, S.

B. (2012). Isolation, Characterization, and Identification of Biological Control Agent for Potato

Soft Rot in Bangladesh. *The Scientific World Journal*, *2012*, 1–6.

https://doi.org/10.1100/2012/723293

Ross, A., Ward, S., & Hyman, P. (2016). More Is Better: Selecting for Broad Host Range

Bacteriophages. *Frontiers in Microbiology*, *7*. https://doi.org/10.3389/fmicb.2016.01352

Sulakvelidze, A., Alavidze, Z., & Morris, J. G. (2001). Bacteriophage Therapy. *Antimicrobial Agents*

and Chemotherapy, *45*(3), 649–659. https://doi.org/10.1128/AAC.45.3.649-659.2001

Sundin, G. W., & Bender, C. L. (1993). Ecological and genetic analysis of copper and streptomycin

resistance in *Pseudomonas syringae* pv. *syringae*. *Applied and Environmental Microbiology*,

59(4), 1018–1024. https://doi.org/10.1128/aem.59.4.1018-1024.1993

Thomas, E. (2001). Multiplicity of Infection. In *Encyclopedia of Genetics*. Elsevier.

https://doi.org/10.1006/rwgn.2001.0858

Voronina, M. V., Bugaeva, E. N., Vasiliev, D. M., Kabanova, A. P., Barannik, A. P., Shneider, M. M.,

Kulikov, E. E., Korzhenkov, A. A., Toschakov, S. V., Ignatov, A. N., & Miroshnikov, K. A.

(2019). Characterization of *Pectobacterium carotovorum* subsp. *carotovorum* Bacteriophage

PP16 Prospective for Biocontrol of Potato Soft Rot. *Microbiology*, *88*(4).

https://doi.org/10.1134/S0026261719040118

Wallace, G. B. (1948). Bacterial Soft Rot of Vegetables. *The East African Agricultural Journal*, *14*(1).

https://doi.org/10.1080/03670074.1948.11664644

Zaczek-Moczydłowska, M. A., Young, G. K., Trudgett, J., Plahe, C., Fleming, C. C., Campbell, K., &

O' Hanlon, R. (2020). Phage cocktail containing *Podoviridae* and *Myoviridae* bacteriophages

inhibits the growth of *Pectobacterium* spp. under in vitro and in vivo conditions. *PLOS ONE*,

15(4). https://doi.org/10.1371/journal.pone.0230842

Tables and Figures

Table 1. Characteristics and biological properties of the three lytic bacteriophages isolated using *P. carotovorum* subsp. *carotovorum.*

Phage	Order	Family	Head diameter (nm)	Tail Length (nm)	Plaque Morphology		Optimal MOI	Latent Period (min)	Burst Size (PFU/cell)
					Transparency	Diameter (mm)			
PPc_A3	*Caudovirales*	*Podoviridae*	81.07 ± 7.04	19.58 ± 2.92	clear	2.0 - 3.0	0.01	10	142
PPc_D1	*Caudovirales*	*Myoviridae*	77.50 ± 4.68	98.15 ± 4.09	clear	1.0 - 1.9	0.1	10	151
PPc_J3	*Caudovirales*	*Myoviridae*	76.17 ± 2.81	137.08 ± 12.78	clear	< 1.0	10	30	42

Figure 1. Transmission electron micrographs (TEM) of PPc_A3, PPc_D1, and PPc_J3 virions.

High titer phage lysates were negatively stained with 2% aqueous uranyl acetate and were visualized

by transmission electron microscopy (TEM; JEOL JEM-1220 Electron Microscope). The images were

scanned with 800 ms exposure, 1.4 gain, and 1.0 bin with no sharpening, normal contrast, and 1.00

gamma. Bar represents 50nm.

Figure 2. Growth characteristics of phages PPc_A3, PPc_D1, and PPc_J3 and their stability on

varying temperature and pH levels. (A) One-step growth curve of each phage grown together with

log phase *P. carotovorum* subsp. *carotovorum* in TSB for 60 min at 30 °C. **(B)** Adsorption of each

phage to log phase *P. carotovorum* subsp. *carotovorum* in TSB for 60 min at 30 °C. **(C)** Stability of the

phages at various temperature and pH levels.

Figure 3. Bacterial killing curves of phages PPc_A3, PPc_D1, and PPc_J3 and their protective

effect against *P. carotovorum* subsp. *carotovorum* on potato tuber slices based on optimal MOI.

(A) Killing curves of phages PPc_A3, PPc_D1, PPc_J3 against *P. carotovorum* subsp. *carotovorum*

measured as change in absorbance at OD=600 nm based on optimal MOI **(B)** Protective effect of each

phage on potato tuber slices against *P. carotovorum* subsp. *carotovorum* determined by measuring the

diameter of rotting tissue (in mm) after 72-hr incubation at 30°C. **(C)** Mean diameter of rotting tissue.

(D) Percent tissue maceration reduction in tissue slices co-inoculated with *P. carotovorum* subsp.

carotovorum and PPc_A3, PPc_D1 or PPc_J3. The data in **(A)** is mean ± SEM in duplicates (n=2).

Semi-*in planta* assay **(B-D)** was representative of two independent experiments and was performed in

triplicates (n=3) with three technical replicates each (n=3). Each dot in **(C-D)** is an individual potato

slice (ie. replicate). Statistical significance was determined with one-way ANOVA with Tukey's post-

test, ****p < 0.0001.

Figure 4. Suppression of *P. carotovorum* subsp. *carotovorum* by phage cocktails *in vitro* and semi-in planta. **(A)** Lytic activities of cocktails of two (PPc_A3+PPc_D1, PPc_A3+PPc_J3, and PPc_D1+PPc_J3) and three (PPc_A3+PPc_D1+PPc_J3) phages at low and high MOI against *P. carotovorum* subsp. *carotovorum* measured as change in absorbance at OD=600 nm. **(B)** Protective

effect of the phage cocktails on potato tuber slices against *P. carotovorum* subsp. *carotovorum*

determined by measuring the diameter of rotting tissue (in mm) after 72-hr incubation at 30°C. **(C)**

Mean diameter of rotting tissue. **(D)** Percent tissue maceration reduction in tissue slices co-inoculated

with *P. carotovorum* subsp. *carotovorum* and phage cocktails. The data in **(A)** is mean ± SEM in

triplicates (n=3). Representative images from the semi-*in planta* assay **(B-D)** were performed in two

independent experiments, each with duplicates (n=2) and three technical replicates each (n=3). Each

dot in **(C-D)** is an individual potato slice.