<u>Lateral Line Ablation by Ototoxic Compounds Results in Distinct Rheotaxis Profiles in Larval</u>

<u>Zebrafish</u>

Kyle C Newton[1,5]*, Dovi Kacev[2], Simon R O Nilsson[3], Allison L Saettele[1], Sam A Golden[3], and

Lavinia Sheets[1,4]*

[1] Department of Otolaryngology, Washington University School of Medicine, St. Louis, MO, USA

[2] Scripps Institution of Oceanography, University of California San Diego, La Jolla, CA, USA

[3] Department of Biological Structure, University of Washington, Seattle, WA, USA

[4] Department of Developmental Biology, Washington University School of Medicine, St. Louis,

MO, USA

* Correspondence:

Kyle C Newton (kylecnewton@gmail.com; kyle.newton@oregonstate.edu)

[5] K C Newton's current affiliation: Department of Fisheries, Wildlife and Conservation Sciences,

Coastal Oregon Marine Experiment Station, Oregon State University, Hatfield Marine Science

Center, Newport, OR, USA

Lavinia Sheets (sheetsl@wustl.edu)

Keywords: zebrafish, lateral line, neuromast, sensory hair cell, rheotaxis, pose estimation,

animal tracking, behavior classification, behavioral profile, machine learning, machine vision

ABSTRACT

The zebrafish lateral line is an established model for hair cell organ damage, yet few studies link mechanistic disruptions to changes in biologically relevant behavior. We used larval zebrafish to determine how damage via ototoxic compounds impact rheotaxis. Larvae were treated with $CuSO_4$ or neomycin to disrupt lateral line function then exposed to water flow stimuli. Their swimming behavior was recorded on video then DeepLabCut and SimBA software were used to track movements and classify rheotaxis behavior, respectively. Lateral line-disrupted fish performed rheotaxis, but they swam greater distances, for shorter durations, and with greater angular variance than controls. Furthermore, spectral decomposition analyses confirmed that lesioned fish exhibited ototoxic compound-specific behavioral profiles with distinct changes in the magnitude, frequency, and cross-correlation between fluctuations in linear and angular movements. Our observations demonstrate that lateral line input is needed for fish to hold their station in flow efficiently and reveals that commonly used lesion methods have unique effects on rheotaxis behavior.

INTRODUCTION

The lateral line is a sensory system used by fishes and amphibians to detect water flow. The functional units of the lateral line are neuromasts; bundles of sensory hair cells located externally along the head and body that mechanotransduce low frequency ($\leq$ 200 Hz) water flow stimuli into electrochemical signals for interpretation by the central nervous system (reviewed in [1]). The lateral line is known to partially mediate rheotaxis, a multimodal behavior ([2]) that integrates input from visual ([3], [4], [5]), vestibular ([6], [7]), tactile ([3], [8], [9], [10]), and lateral line systems ([9], [10], [8]) to facilitate fish orientation and movement of fish with respect to water flow ([9], [10]).

Although the contribution from the lateral line is well established, there is conflicting evidence on whether it is essential for rheotaxis in fish ([2], [5], [9], [10], [11], [12]). Inconsistent methodologies used on distinct fish species that differentially rely on the lateral line to mediate

swimming behaviors obfuscate the relationship between the lateral line and rheotaxis. In

addition, a recent review hypothesized that differences in the spatial characteristics and velocity

of the flow stimuli used to assay rheotaxis likely resulted in the disparate results reported in

these studies (see Table 1 in [13]).

Zebrafish hair cells demonstrate unique regenerative features that make them an

important model for hearing loss research ([14], [15], [16], [17], [18], [19]). Many studies have focused on

mechanistic disruption of hair cell activity, but few have explored the association between

disruption and hair cell mediated behavior. Previous researchers have developed several

different assays to study rheotaxis in larval fishes (e.g., [5], [20], [21], [22]). However, we sought to

develop a behavioral assay and analytical methodology that is sensitive, spatiotemporally

scalable, and robust enough to be used on a variety of species, ontogenetic stages, sensory

modalities, and behavioral responses that are pertinent to biomedical and ecological research.

We therefore investigated how the lateral line contributes to rheotaxis in larval zebrafish

and developed a standardized assay that could identify subtle differences in rheotaxis behavior

([12]). We compared the rheotaxis response of fish with an intact lateral line to those with lateral

line hair cells ablated by two commonly used compounds: copper sulfate ($CuSO_4$; [17], [23], [24], [25])

and neomycin ([5], [15], [26]). We hypothesized that different ototoxic compounds might produce

distinct changes in rheotaxis behavior because they injure lateral lines via different cellular

mechanisms ([27]), and that these behavioral changes could be empirically quantified using

machine vision and learning technology.

RESULTS & DISCUSSION

To determine the contribution of the lateral line to rheotaxis in fishes, we used $CuSO_4$

and neomycin to ablate the lateral line neuromasts of larval zebrafish (6-7 days post-fertilization

(dpf)), then video recorded the swimming behavior of individual larvae in a microflume under no-

flow and flow stimulus conditions (Fig. 1). Our procedures eliminated visual and linear

acceleration cues (see methods) and young larval zebrafish cannot detect horizontal angular

Figure 1. Experimental microflume used to conduct rheotaxis assays under IR illumination. a) The microflume (220 x 100 x

40 mm) with a removeable working section (30 x 30 x 10 mm) was 3D printed from translucent resin and placed on top of an

infrared (850 nm) LED array. b) Schematic of experimental set-up. The IR light passed through the flume and the overhead camera

recorded rheotaxis trials at either 200 or 60 fps onto SD cards. The timing and duration for the camera and flume pump onset and

offset of was controlled by an Arduino and pump voltage (i.e., water flow velocity = 9.74 mm s^{-1}) was controlled by a rheostat. Each

trial was monitored via the live camera feed displayed on the PC and all videos were copied in duplicate onto a 12TB RAID array.

Dots on fish larva indicate seven body positions tracked by DeepLabCut.

velocity cues (i.e., yaw; [31], [32]); however, it was not possible to selectively block tactile cues in a

non-invasive manner. To confirm ablation of lateral line neuromasts, a subset of all fish that had

undergone behavioral testing were fixed, then proceed for immunolabeling of hair cells and

innervating afferent neurons (Fig. 2). We observed near total hair cell loss in both anterior and

posterior lateral line neuromasts with CuSO$_4$ treatment (Fig. 2 a, b, d, e, g) and a significant

impact on the morphology and reduction in hair cell number per neuromast following neomycin

treatment (Fig. 2 a, c, d, f, g). These results support that the ototoxic compound-treated fish

used in our behavior assay had a total absence or severe impairment of lateral line function.

Figure 2. Confirmation of neuromast hair cell loss following CuSO$_4$ or neomycin treatment. a-f) Representative confocal max

intensity projection images of the: a-c) mid posterior lateral line (MidLL) fourth neuromast (L4); and d-f) second anterior supraorbital

(SO2) neuromast from the fish cohorts used for behavior experiments. Hair cells were labeled with an antibody against Otoferlin

(HCS1; green a-f, gray a'-f'). Afferent neurons were labeled with an antibody against Calbindin (magenta), and cell nuclei were

labeled with DAPI (blue). g) Quantification of the grand mean (± SEM) number of hair cells per neuromast in intact (CTL), CuSO4-

and neomycin-treated fish. Each dot represents the mean number of hair cells from the MidLL (L3, L4, and L5) or SO (left and right)

neuromasts from an individual fish. Data were collected from fish used in three experimental behavior trials; 4-6 fish per condition

per trial. Significance values: ** < 0.01, ***< 0.001, **** < 0.0001

We used machine vision and learning software for 3D pose estimation, movement

tracking, and annotation of videos for positive rheotaxis events (as defined in methods) when

fish oriented toward (0° ± 45°) and actively swam into the oncoming flow (e.g., Supplementary

Fig. 1). We standardized our analyses by comparing rheotaxis data acquired during flow to the

swimming behavior of fish when they were randomly oriented at 0° ± 45° under no flow (see

data analysis). Under no flow conditions, we determined that each treatment group of fish was

randomly distributed (Fig. 3; Supplementary Table 1) and had no natural proclivity to orient their

bodies to 0° ± 45° (Supplementary Table S2).

Lateral Line Ablation Alters but Does Not Eliminate Rheotaxis Behavior

114   We predicted that fish treated with the minimum dosage of $CuSO_4$ or neomycin

115   necessary to ablate lateral line hair cells would not perform rheotaxis as well as control fish with

116   an intact lateral line. Surprisingly, fish with lesioned lateral line organs could still orient into

117   oncoming flow like control fish (Fig. 3; Supplementary Tables 1-2), indicating that the effects of

118   ablation were subtle, and the lateral line was not essential for rheotaxis behavior.

119   By analyzing the kinematic components of rheotaxis movement, stark behavioral

changes were observed among lesioned fish. Comparisons between lateral line-ablated and

control groups showed significant differences in the mean body angle or angular variance (Fig.

3, Supplementary Table 3), mean duration of rheotaxis events, mean number of rheotaxis

events, total distance traveled, and latency to the onset of rheotaxis (Fig. 4, Supplementary

Tables 4-7). These observations demonstrate that rheotaxis occurs but is altered in lateral line-

ablated groups, which contrasts with previous reports on larval zebrafish ([5], [22], [28]) and supports

the idea that the lateral line is not required for rheotaxis in fishes ([2], [11], [12], [29], [30]).

127   We posit that our focus on acute rheotaxis behavior (~20 s) and the fine scale

spatiotemporal sampling of machine vision technology allowed us to detect subtle changes in

behavior that were overlooked in previous rheotaxis studies ([2], [5], [12], [29], [30]). Our methods

eliminated turbulent water flow, optic flow ([3]), and certain vestibular cues (yaw: [31], [32]; linear

acceleration). However, we did not eliminate tactile cues because we could not prevent fish

from contacting the substrate and our flow rate was sufficient to displace substrate coupled fish

along the bottom and against the rear mesh of arena. Therefore, we propose that lateral line

ablated fish might have used tactile cues to gain an external frame of reference and perform

rheotaxis ([8], [10]), but explicitly testing this idea would require designing an experimental

apparatus that enables the video capture of fish movements along the vertical plane (Z-axis),

which is not possible in our translucent micro flume.

Ototoxic Compounds Differentially Influence the Distribution of Mean Body Angles During Flow

To identify differences in body orientation, the mean body angle for fish in the presence

or absence of flow stimuli were compared among treatment groups (Fig. 3). Since flow

originated at the top of the flume (located at 0°), mean body angle was determined relative to

that of the oncoming water stimulus. The angular variance of each group is represented by the

inverse of the mean length of the resultant vector (i.e., short vectors = high variance, and vice

versa; Fig. 3). Under no flow, fish from each treatment group swam with a random orientation,

indicated by a grand mean body angle that was statistically different from 0° and length of the

resultant vector close to zero (Fig. 3a, Supplementary Table 2). When flow was applied, all

groups exhibited significant alignment into the oncoming flow stimulus because the grand mean

body angles were clustered at 0° and length of the resultant vectors were close to one (Fig. 3b-

c).

During rheotaxis, post-hoc comparisons within treatment groups showed a significant

difference in the homogeneity of distributions between initial and final portions of the flow

stimulus, indicating that overall orientation behavior within groups was not consistent for the

duration of flow presentation (Fig. 3b-c, Supplementary Table 3). Comparisons among groups

within the initial 10s portion of flow showed a significant difference in the distribution of mean

body angles between control and $CuSO_4$ fish, and between control and neomycin fish (Fig. 3b,

Supplementary Table 3). There was an interaction between treatment and stimulus where the

distribution of mean body angle in $CuSO_4$-ablated fish during the initial stimulus bin was different

than that of neomycin-ablated fish during the final stimulus bin (Supplementary Table 3),

suggesting that these lesion methods differentially impacted the rheotaxis behavior of fish.

Figure 3. The mean resultant vectors of fish treatment groups before and during water flow stimulus indicates fish with

lateral-line organs ablated by CuSO₄ or neomycin can still perform rheotaxis. Each dot outside the circles represents the mean

body angle of an individual fish for the 10 s duration of the no flow and flow stimulus conditions. The grand mean vector for each

group is represented by a summary vector with an angle, *theta*, and a mean resultant length, *rho*, where the length of the vector

represents the distribution of individual angles around the mean angle of the group. The length of the vector ranges from zero for

uniform distributions, to one for distributions perfectly aligned with the mean angle. Consequently, the angular variance (1- *rho*) is

inversely related to vector length. Under no flow (t = 0-10 s), groups of lateral line intact (control, n = 248) and lesioned (CuSO4, n =

204; neomycin, n = 222; 18 experimental sessions) fish have a random distribution of individual mean body angles. Under flow, all

groups show a statistically significant orientation to 0° ± 45°, but the distributions of the individual mean angles within the groups

differ between the initial (t = 10-20 s) and final (t = 20-30 s) stimulus bins. Distributions with the same lowercase letter indicate

groups that do not differ statistically.

Lateral Line-Ablated Fish Performed Rheotaxis for Shorter Durations but Travelled Longer

Distances

Although an intact lateral line was not required to perform rheotaxis, lesioned fish

behaved differently than non-lesioned fish in flow stimulus, indicating that lack of input from

neuromast hair cells affected rheotaxis behavior. To quantify differences in the gross metrics of

rheotaxis behavior among groups, we used generalized linear mixed models (GLMMs) that

accounted for the random effects of individual variation to compare the mean duration and

mean number of rheotaxis events, total distance traveled, and latency between flow

presentation and behavior onset. We standardized the data by comparing events of rheotaxis to

events when fish were randomly oriented at 0° ± 45° under no flow. Hereafter both conditions

are referred to as "events".

Without flow, the mean duration of random orientation events did not differ among

treatment groups (Fig. 4a, Supplementary Table 4). With flow, the mean duration of rheotaxis

events increased for each group, and was greatest in the controls, less in neomycin-, and least

in $CuSO_4$-treated fish (Fig. 4a, Supplementary Table 4). An interaction between stimulus and

treatment was seen during the initial 10s of flow in the $CuSO_4$ treatment cohort, but during the

final 10s the differences among all treatments became significant (Supplementary Table 4).

Additionally, there was a significant effect of stimulus and an effect of treatment on the mean

number of events where the mean was greatest in neomycin-treated, less in controls, and least

in CuSO$_4$-treated fish under no flow and flow conditions (Fig. 4b, Supplementary Table 5).

Figure 4. Lateral line ablated fish performed rheotaxis for shorter mean durations yet travelled greater total distances

compared to controls. Red diamonds in each plot indicate the mean ± SE values. a) Lateral line intact (gray = control, n=248) fish

have a longer mean duration of rheotaxis events during flow stimulus than lesioned fish (blue = CuSO$_4$, n = 204; green = neomycin,

n = 222; 18 experimental sessions). b) The mean number of 0° orientation and rheotaxis events was greatest for neomycin fish and

the least for CuSO$_4$ fish under no flow and flow conditions, respectively. c) Under no flow, neomycin fish traveled a greater total

distance than control and CuSO$_4$ fish; but under flow, neomycin and CuSO$_4$ fish traveled a greater total distance than control fish. d)

Compared to control and neomycin fish, CuSO$_4$ fish had the longest mean latency to the onset of the first rheotaxis event after flow

stimulus presentation. Lines indicate statistical comparisons between control and treatment groups (see Supplementary Tables 4-7).

The effects of treatment are indicated by long color-coded bars with branches, whereas interactions are indicated with short bars.

There was a significant effect of stimulus in a-c (not shown for clarity). Significance values: * = 0.05, ** = 0.01

The total distance travelled during events was influenced by the flow stimulus and type of lesion,

where neomycin-treated fish travelled further than control and $CuSO_4$-treated fish in the

presence and absence of flow (Fig. 4c, Supplementary Table 6). Compared to control and

neomycin fish, $CuSO_4$-lesioned fish had a longer latency between flow onset and rheotaxis

initiation (Fig. 4d, Supplementary Table S7). Altogether, our data show that an intact lateral line

allowed fish to sustain rheotaxis into oncoming flow for longer durations yet travel shorter

distances, suggesting that ototoxic compounds reduced the economy of movement of larval

zebrafish in response to flow. These results support the idea that the lateral line allows

epibenthic fish under flow conditions to hold their station with respect to the substrate ([2], [33])

215   Interestingly, $CuSO_4$ and neomycin ablation affected rheotaxis behavior in different

ways. $CuSO_4$ exposure decreased activity, as evidenced by fewer rheotaxis events with greater

latency between flow delivery and behavior initiation. This contrasts with the neomycin-exposed

fish that exhibited frequent bursts of rheotaxis and travelled longer distances. While zebrafish

larvae have been shown to be relatively resistant to the concentration and exposure time of

$CuSO_4$ used ([34]), subtle differences in behavior could have been a consequence of nonspecific

neural toxicity. Alternatively, the distinct effects of $CuSO_4$ and neomycin treatments on the

spatiotemporal nature of rheotaxis might have been due, in part, to their different mechanisms

of neuromast ablation. In $CuSO_4$-treated fish, the hair cells were completely ablated, and the

supporting cells and afferent neurons were severely damaged ([17]). However, on occasion a few

hair cells would remain in neomycin-treated fish (Fig. 2c, f). If residual hair cells in the neomycin

group retained some functionality despite severe morphological damage, then it is possible that

their sensitivity might have been amplified through efferent modulation ([35]), or influenced by

intact supporting cells (Fig. 2c, f) and recruited from a "silent" state (e.g.,[36]), to compensate for

reduced sensory input thus resulting in the observed difference in behavior between lesioned

groups.

Overall Spatial Use of the Arena During Rheotaxis Differs Among Treatments

During flow, we observed that lateral line-lesioned fish were often pushed against the

back of the testing arena, whereas intact fish were often swimming at the front of the arena near

the flow source. To quantify this observation, density plots of the location of fish in the arena

throughout the duration of the experiment were generated (Fig. 5 (2D); Supplementary Fig. 2

(1D)) then compared among treatments. The total density of fish in 2D space was parsed out

over five regions of interest (ROI, Fig. 5a) that were determined by the size of fish, their

orientation during rheotaxis, and the dimensions of the flow field.

Under no flow, fish that were placed into the flume generally avoided the middle as they

explored the boundaries of the arena; however, $CuSO_4$-treated fish preferred the middle more

and explored the boundaries less than the control or neomycin-treated fish (Fig. 5a,

Supplementary Fig. 2). Under flow, generalized linear models (Supplementary Table 8) indicate

that the spatial use of fish during rheotaxis differed among all three treatment groups. (Fig. 5d).

Fish from all treatment groups performed rheotaxis with a pronounced preference for the sides

(Fig. 5d) that mimicked the small lateral gradient in the laminar flow field (Supplementary Fig.

3a). The reduced velocity of the laminar flow gradient along the sides was created by a minute

boundary layer of null flow adjacent to the wall that gradually increased (over ~2-5 mm, front to

rear) until it became part of the freestream flow field (i.e., blue arrows in Supplementary Fig. 3).

The gradient provided a refuge (e.g., [37]) where intact and lesioned fish could swim into the flow

with reduced energetic cost ([38]). As our definition of rheotaxis (see methods; Supplementary

Fig. 1d, e) specifies that only fish that moved their tail and had forward body translation into the

flow, fish that were in null water flow (i.e. too close to the wall) were excluded from the rheotaxis

dataset. Furthermore, control fish frequently performed rheotaxis while maintaining position in

Figure 5. Under flow, lateral line intact (control) fish used the front part of the arena near the source of the flow, whereas lesioned (CuSO4, neomycin) fish use the back portion of the arena. a-b) For all possible mean body angles (0°- 360°): a) all treatment groups of fish show similar density, or total spatial use, of the arena under no flow conditions; b) however, the controls (control, n = 248) used the front of the arena and the lesioned (CuSO4, n = 204; neomycin, n = 222; 18 experimental sessions) fish used the back of the arena under flow. c-d) Filtering the data for mean body angles required for rheotaxis (0° ± 45°): c) all groups clustered along the left and right sides of the arena under no flow; d) but under flow, the controls used the front-left and the lesioned use the back-right portions of the arena. Dotted lines and labels in capital letters indicate spatial regions of interest (ROI). For clarity, statistical comparisons among treatment and ROIs were not included in the figure; however, there were significant fixed effects for CuSO4 and neomycin treatments, the back ROI, and interactions between treatment and ROI (Supplementary Table 8).

the flow near the upper-left corner of the arena whereas lesioned fish primarily occupied the

back-right corner (Fig. 5d, Supplementary Fig. 3b). These observations indicate that an intact

lateral line enabled larval zebrafish to better station hold while occupying the areas of strongest

flow (Supplementary Fig. 3b) and avoid being swept backwards against the rear mesh. The

propensity of lesioned fish to use the rear of the arena while performing rheotaxis in the

absence of visual ([3]) and horizontal vestibular ([31], [32]) cues suggests that these larvae might

have used tactile cues to provide the external frame of reference necessary to orient and swim

against flow ([2], [8], [10]). Unfortunately, our flume design and camera setup prevented us from

exploring this possibility.

Lateral Line Ablation Reduces the Proportion of Individual Fish that Perform Rheotaxis

During flow presentation, the control group had the greatest proportion of individual fish

that performed rheotaxis during the experiment (as defined in Supplementary Fig. 1) followed by

neomycin-treated then CuSO$_4$-treated groups (Fig. 6). Intact fish plateaued relatively quickly

compared to lesioned fish, but the values for all three groups converged during the final few

seconds of flow presentation.

Figure 6. Intact lateral line enabled a greater proportion of fish to perform rheotaxis within the first 15 seconds of stimulus

onset. Time series of the proportion of individual fish within each group that performed rheotaxis during flow presentation. A greater

Spectral Decomposition Shows that Lateral Line Ablation Impacts the Overall Trend and

Periodic Fluctuation in Linear and Angular Movement

To uncover the impact that lateral line ablation had on the swimming kinematics of fish,

we analyzed how the magnitude of the linear (relative distanced moved, relative velocity,

relative acceleration) and angular (mean body angle, mean length of the resultant vector)

components of movement changed over time. We observed that all treatment groups swam in

the burst-and-glide style that is characteristic of larval zebrafish with intact lateral lines, but with

noticeable differences in the quality of their movement. Because the lateral line mediates station

holding behavior ([2], [33], [38]), we postulated that, under flow, the oscillations in relative linear and

angular movements for lateral line-intact fish would be smaller in magnitude than those of

lesioned fish. The observed time series data (Fig. 7a, d, g; Supplementary Fig. 4a, d) had a

"seismic" appearance where noise masked the underlying signal. Therefore, we removed the

random noise (Supplementary Fig. 4g-k) and decomposed the observed datasets into their

fundamental large- and small-scale components: the overall trend in movement magnitude

during the entire experiment (Fig. 7b, e, h; Supplementary Fig. 4b, e) and the periodicity, or

recurring fluctuations in movement magnitude, that occurred during any given second of the

experiment (Fig. 7c, f, i). Because the relative movement (Fig. 7), velocity and acceleration

(Supplementary Fig. 4c, f) data showed similar periodic fluctuations, only relative movement is

shown in Fig. 7 for visual clarity.

During rheotaxis, the trend among groups was that CuSO$_4$ ablation reduced the

magnitude of relative movement (Fig. 7b) but not the relative velocity or acceleration of fish

(Supplementary Fig. 4c, f) compared to control and neomycin groups. CuSO$_4$- and neomycin-

treated fish had a trend of reduced ability to achieve and maintain orientation within flow relative

to controls (Fig. 7e). Lesioned fish had mean resultant vectors (*rho*) of longer length meaning

that they had less variance (1- *rho*) in their mean body angle compared to controls (Fig. 7h).

These data suggest that an intact lateral line allowed control fish to better detect changes in

water flow, regularly make small magnitude course corrections to their body angle (Fig. 7h),

rapidly orient into flow with greater accuracy (Fig. 7e), which resulted in a greater proportion of

non-lesioned fish performing rheotaxis compared to lesioned fish (Fig. 6).

Figure 7. The overall trends and periodic fluctuations in the linear (relative distance moved) and angular (mean body angle,

mean length of the resultant vector) motion parameters of rheotaxis behavior differ among treatment groups. (Note: the

relative velocity and acceleration periodicity data mimicked the patterns observed in relative movement; see Supplementary Fig. 3).

Gray = control (n = 248), blue = CuSO₄ (n = 204), green = neomycin (n = 222). Spectral decomposition of the observed data (a, d, g)

removed the noise (Supplementary Fig. 3g, h, i) to reveal the overall underlying trends (b, e, h) and the periodicity, or recurring

fluctuations (c, f, i) that occurred during any given 1 s of the experiment. The periodicity waveform peaks (c, f, i) indicate the average

amount (amplitude), number, direction (positive = increasing; negative = decreasing), and order of occurrence for these cyclic

fluctuations as a function of unit time (1 s). The overall trends were that $CuSO_4$ treated fish had the least relative movement (b),

while the control fish more rapidly oriented to 0° (e) and swam with more angular variance (h; 1 – mean length of the resultant

vector) compared to lesioned fish. The periodic fluctuation in relative distance moved (c) was greatest in $CuSO_4$ treated fish

compared to control or neomycin treated fish. However, the fluctuation in mean body angle (f) was greatest in neomycin treated fish

compared to control and $CuSO_4$ fish, while the fluctuation in mean length of the resultant vector (i.e the angular variance; i) was

greatest in control fish compared to lesioned fish.

Among treatment groups, there were clear differences in the magnitude of changes in

the periodic cycles of linear (Fig. 7c; Supplementary Fig. 4c, f) and angular (Fig. 7f, i)

movements during rheotaxis. The linear data show that $CuSO_4$-treated fish had fluctuations of

the greatest magnitude in relative distance moved, velocity, and acceleration compared to those

of control and neomycin-treated fish (Fig. 7c; Supplementary Fig. 4c, f; Supplementary Table 9).

In $CuSO_4$-treated fish, the amplitude of the waveform peaks during the sampling period was

large and decreased rapidly, but in control and neomycin-treated fish the peaks were small and

increased gradually (Fig. 7c; Supplementary Fig. 4c, f). The large fluctuations of $CuSO_4$-treated

fish could serve to compensate for their delayed response to flow (Fig. 4d), but when their trend

of lesser relative movement (Fig. 7b) is considered, it results in the greatest relative increase in

distance traveled among groups (Fig. 4c). These empirical data support the qualitative

observations that $CuSO_4$-treated fish performed rheotaxis with delayed responses and erratic

linear movements, whereas control and neomycin-ablated fish performed rheotaxis with low

latency responses and graded linear movements.

The periodic fluctuation in mean body angle among treatment groups showed an initial

small turn to the left (negative values) followed by a series of right (positive values) and left

turns that were of relatively small magnitude in controls and $CuSO_4$-treated fish, and relatively

large magnitude in neomycin-treated fish (Fig. 7f; Supplementary Table 9). The amplitude of the

waveform peaks in control and $CuSO_4$-treated fish were relatively small and gradually increased

during the sampling period, but those of neomycin-treated fish were relatively large throughout.

Therefore, control and CuSO$_4$-treated fish performed rheotaxis with a graded response in mean

body angle, but neomycin-treated fish performed rheotaxis with large erratic changes in mean

body angle. We also examined the periodicity in mean resultant length, which reflects the

fluctuation in angular variance as fish performed rheotaxis. In control fish, the waveform had

peaks of small amplitude that rapidly increased during the sampling period, but in lesioned fish

the waveform was comprised of small peaks of consistent amplitude (Fig. 7i). Intact fish showed

a graded response where small initial changes in angular variance preceded much larger

subsequent changes that coincided with the flat portion of the peaks in mean body angle (Fig.

7f), which indicates that control fish regularly made small changes in their body angle and

maintained this new heading before making another angular adjustment. Conversely, lesioned

fish had very little fluctuation in angular variance (Fig. 7i) once a mean body angle was chosen,

regardless of whether the change in mean body angle was relatively small (e.g., CuSO$_4$) or

large (e.g., neomycin; Fig. 7f).

These data support that, during rheotaxis, fish with an intact lateral line organ made

small adjustments to their linear movement, velocity, acceleration, and mean body angle with a

high degree of variability. This graded, flexible, and finely tuned behavioral response to flow

resulted in a greater ability of control fish to rapidly attune and maintain fidelity to the oncoming

flow vector (Fig. 7e). Conversely, CuSO$_4$- and neomycin-ablation caused fish to swim into flow

with erratic linear and angular movements, respectively, which ultimately undermined their

ability to maintain a 0° heading into the flow (Fig. 7e).

Spectral Analysis Reveals that Lateral Line Ablation Impacts the Temporal Fluctuation of Linear

and Angular Movement

To determine the impact that lateral line ablation had on the frequency of the fluctuations

in linear and angular movement, we decomposed each periodicity dataset (Fig. 7c, f, i;

Supplementary Fig. 4c, f) into a power spectrum that depicted the spectral density (i.e., number

of changes in movement per frequency) over a continuous range of frequencies (Fig. 8a-e). For

each parameter, the frequency and amplitude of three most dominant peaks were summed to

calculate the net shifts in frequency and spectral density among treatment groups

(Supplementary Table 10). Relative to controls, the power spectra of $CuSO_4$- and neomycin-

treated fish showed a net shift to lower dominant frequencies (downshift) and a net increase in

the density of the dominant peaks for relative movement, velocity, and acceleration (Fig. 8a-c,

Supplementary Table 10), which indicates that lateral line-ablated fish performed rheotaxis with

greater numbers of linear movements to compensate for their overall reduction in fluctuation

frequency. The rheotaxis behavior of $CuSO_4$-treated fish showed less frequent fluctuations in

relative movement compared to controls and neomycin-treated fish (Fig 8a), but the fluctuations

in relative velocity and acceleration for $CuSO_4$-treated fish and controls occurred over a wider

range of frequencies than those of neomycin-treated fish (Fig 8b-c). Specifically, the relative

movement spectra of the control and $CuSO_4$-treated fish had broad clusters of dominant peaks

that gradually shifted to higher frequencies (upshifted) for relative velocity and acceleration.

However, the spectra of neomycin-treated fish were dominated by a single high-density peak

that consistently occurred at 0.21 Hz for relative movement, velocity, and acceleration (Fig. 8a-

c). These data reveal that lateral line ablation results in rheotaxis behavior characterized by

fluctuations in relative movement, velocity and acceleration that occur less frequently, but that

neomycin ablation has the additional effect of "temporally restricting" the frequency of

fluctuations in linear movement to occur around a consistent dominant frequency.

Power spectra density curves indicated that intact fish performed rheotaxis with fewer

numbers of fluctuations in mean body angle that occurred less frequently, whereas lesioned fish

performed rheotaxis with a greater numbers of high frequency fluctuations in mean body angle.

The spectrum of controls shows five broadly spaced peaks, but those of the lesioned fish have

fewer peaks that are clustered around the dominant frequencies (Fig. 8e) indicating that intact

fish performed rheotaxis with greater variety in the frequency of fluctuations in their angular

variance compared to lesioned fish (Fig. 8e). We observed the mean body angle of $CuSO_4$- and

neomycin-treated fish showed a net upshift in the dominant frequencies, a net increase in

dominant peak density, and temporal restriction, or greater clustering of power spectra into

distinct peaks, relative to that of controls (Fig. 8d, Supplementary Table 10). By contrast, the

spectrum of control fish was relatively flat with low density peaks that lacked distinct clustering

into dominant frequencies (Fig. 8d) Additionally, there were two divergent patterns seen in the

power spectra for mean resultant vector length in lesioned fish: $CuSO_4$-treated fish had a net

downshift in dominant frequencies and a net increase in peak density, whereas neomycin-

treated fish had a net upshift in frequency and a net decrease in peak density relative to those

of controls (Fig. 8e, Supplementary Table 10). Thus, $CuSO_4$-lesioned fish performed rheotaxis

with greater number of fluctuations in angular variance that occurred less frequently, but

neomycin-lesioned fish performed rheotaxis with fewer numbers of fluctuations in angular

variance that occurred more frequently compared to intact fish. These results show that lateral

line ablation in larval zebrafish resulted in rheotaxis behavior with greater numbers of

fluctuations in mean body angle, fewer numbers of fluctuations in angular variance, and a

greater clustering of angular movements into fewer peaks.

Figure 8. Power spectra density curves show that an intact lateral line allowed fish to make fewer yet more temporally variable changes in relative linear and angular movement. Because frequency and period are inversely related, the low frequency peaks to left of the periodograms indicate cycles with longer periods, and vice versa The amplitude of the peaks indicates the spectral density, or the number of movement events at a given frequency that occurred during the experiment. The peaks with the greatest amplitude indicate the fundamental or dominant frequencies of fluctuation in the periodicity data. The frequency and amplitude of three most dominant peaks were summed to calculate the net shifts in frequency and power. Relative to controls (gray, n =248), lesioned (blue = $CuSO_4$, n=204; green = neomycin, n = 222) fish had a net downshift in the three dominant frequencies of (a) relative movement, (b) velocity, and (c) acceleration and a net upshift in (d) mean body angle of larval zebrafish during rheotaxis.

For the dominant frequencies of (e) mean resultant length, there was a net downshift and upshift for $CuSO_4$- and neomycin-treated

fish, respectively. Furthermore, relative to lateral line intact fish, the peaks of lesioned fish are clustered into fewer peaks of greater

amplitude over a relatively narrow range, which indicates that lateral line ablation increased the number yet reduced the temporal

variation of changes in movement.

We interpret the relatively flat and low-density power spectra observed in control fish

(Fig. 8a-d) to indicate that an intact lateral line allowed fish to respond to flow with fewer overall

fluctuations in linear and angular movement over a broader range of fluctuation frequencies

compared to lesioned fish. It is sensible that the mechanosensory input from lateral line hair

cells gave intact fish a greater ability to respond to flow with greater efficiency, efficacy, and

economy of movement. Conversely, fewer peaks of greater density observed in lateral line

lesioned fish indicates that losing the ability to detect water flow led to more changes in linear or

angular movement that occurred over a restricted range of frequencies thus reducing the

effectiveness of their movements in response to flow.

Cross Correlation Between Linear and Angular Movements Shows Distinct Rheotaxis Profiles

Among Groups

In the absence of visual cues, intact larval zebrafish detected the presence of flow

through a change in linear or angular displacement ([22]) and responded by either turning right,

left, or moving forward into the flow. Because these movements relied on input from the lateral

line, we investigated the effect of ototoxic compounds on the cross correlation between linear

and angular movements. The correlograms depict the degree, direction, and relative timing

between an above average increase in relative movement (Fig. 9a-b), velocity (Fig. 9c-d), or

acceleration (Fig. 9e-f) and an above average change in mean body angle (Fig. 9a, c, e) or

mean resultant length (Fig. 9b, d, f). By focusing on the strongest cross-correlations, treatment-

specific patterns emerged in the direction and relative timing between large changes in linear

and angular movements during rheotaxis.

The rheotaxis movements of control fish in response to flow were smoother, less erratic,

and more effective at station holding within the flow field compared to those of lesioned fish. The

rheotaxis profile of control fish was such that the relative timing and direction of changes in

mean body angle depended upon the time derivative of the linear movement, but changes

angular variance were consistent across all linear movements. For example, intact fish tended

to change their mean body angle to the *left* of flow *before* a large change in relative movement

(Fig. 9a), to the *right* of flow *simultaneous* with changes in velocity (Fig. 9c), and to the *left* of

flow *after* changes in acceleration (Fig. 9e). However, the strongest correlation between a

reduction in angular variance always occurred *prior* to large changes in relative movement (Fig.

9a), velocity (Fig. 9c), and acceleration (Fig. 9e). In lesioned fish, the timing and direction of the

strongest cross-correlations shifted the rheotaxis profile of these fish away from that of controls

in a predictable, consistent, and treatment-specific manner. $CuSO_4$-treated fish tended to

change their mean body angle to the *right* of the oncoming flow vector *after* large changes in

movement (Fig. 9a), velocity (Fig. 9c), and acceleration (Fig. 9e). However, neomycin-treated

fish tended to change their body angle to the *left* of the flow *after* changes in movement (Fig.

9a), velocity (Fig. 9c), and acceleration (Fig. 9e). As in controls, lesioned fish always had a

reduced angular variance correlated with changes in their linear movements, but the relative

timing of the cross correlation shifted according to the ototoxic compound used and the time

derivative of the linear movement. In $CuSO_4$-treated fish, the strongest correlation between a

reduction in angular variance tended to occur *prior* to changes in movement (Fig. 9b), and *after*

changes in velocity (Fig. 9d) or acceleration (Fig. 9f). Conversely, in neomycin-treated fish,

reductions in angular variance occurred *after* changes in movement (Fig. 9b), and *before*

changes in velocity (Fig. 9d) or acceleration (Fig. 9). These data revealed that control fish with

an intact lateral line detected oncoming flow and adjusted their mean body angle *prior* to

Figure 9. Cross correlations between linear and angular movement data indicate ototoxic compound-specific changes to

the rheotaxis behavioral profile of fish. The correlograms depict how an above average increase in relative movement (a, b),

relative velocity (c, d), or relative acceleration (e, f) were significantly cross correlated with above average increases or decreases in

the mean body angle (a, c, e) or mean resultant length (b, d, f). In the figures for mean body angle (a,c,e), the positive and negative

peaks indicate that fish were oriented to the right or left of the oncoming flow vector, respectively. In the figures for mean resultant

length (b, d, f), the positive and negative peaks indicate fish that had a lesser or greater variance of the mean body angle,

respectively. The X-axis indicates the relative timing, or lag, of the cross correlation between the angular parameter with respect to

an above average increase in the linear parameter (zero = simultaneous occurrence; negative = angular change occurs *before*

linear change; and positive = angular change occurs *after* linear change). *For example, interpret panel 10a as:* In control (gray, n =

248) fish, above average increases in relative movement are significantly correlated with changes in mean body angle to the left of

the flow vector that previously occurred. In CuSO₄-treated (blue, n =204) fish, above average increases in relative movement are

significantly correlated with above average changes in mean body angle that subsequently occurred. In neomycin-treated (green, n

= 222) fish, above average increases in relative movement are significantly correlated with below average changes in mean body

angle that subsequently occurred.

making a change in their relative movement, while lateral line lesioned fish changed their mean

body angle *after* a change in their relative linear movement. Our results support the reported

mechanism of water flow detection in larval zebrafish where they use their lateral line hair cells

to sense the vorticity, or curl, created by gradients in the flow field the determine flow direction

by measuring the temporal change in vorticity as they experience flow ([22]).

It appears that an intact lateral line allowed fish to rapidly detect flow and adjust their

heading prior to swimming into the flow. However, in the absence of lateral line and visual cues,

and the inability of 6-7dpf zebrafish larvae to detect horizontal angular cues (yaw) by the

vestibular system ([31], [32]), it is possible that lesioned fish might have relied on tactile cues, such

as physical displacement along the substrate, to gain an external frame of reference necessary

to orient and swim into flow. Intact fish also tended to change their body angle to the left of the

flow vector, which may reflect lateral line handedness where larval zebrafish prefer to use the

left side of their lateral line to detect flow stimuli in a manner like that of blind cavefish ([39]).

Lateral line ablation shifted the relative timing of the cross correlations between angular and

linear movement and might have impaired any potential lateral line handedness, which may

have further contributed to the erratic rheotaxis behavior observed in lesioned fish.

In summary, lateral line ablation of larval zebrafish resulted in distinct, treatment-specific

rheotaxis profiles that differed from that of intact fish in the following ways: 1) delayed relative

timing between changes in mean body angle and all linear movements, 2) changed mean body

angle in response to flow, and 3) shifted timing between reductions in angular variance and

changes in linear movement.

CONCLUSIONS

In this study, ablating the lateral line of larval zebrafish with two commonly used ototoxic

compounds impacted their ability to produce the fine adjustments required to station hold in

response to water flow. Lateral line ablation inhibited the ability of fish to discriminate subtle

distinctions in flow, resulting in more intense overcorrections and decreased economy of motion.

Our data support the hypotheses that the lateral line mediates station holding behavior ([2], [33], [38]),

but is not required for rheotaxis behavior ([2], [11], [12], [29], [30]) in larval zebrafish in non-uniform laminar

flow. We posit that the physical displacement of lesioned fish along the substrate may have

provided sufficient tactile cues ([8], [10]) necessary for lateral line ablated fish to perform rheotaxis.

We propose that the greater angular variance observed in intact fish might indicate that

these fish were regularly sampling the velocity gradients of the flow stimuli ([22]) from a variety of

body angles so that they could reduce their response latency, quickly orient with respect to

fluctuating flow stimuli, and maintain their overall mean body angle with greater fidelity to the

flow vector than lesioned fish. During flow, intact fish had recurring fluctuations in relative

movement, velocity, and mean body of lower magnitude, and fluctuations in relative movement,

velocity, acceleration, and mean body angle of controls that were fewer in number yet occurred

over a wider range of temporal frequencies compared to lesioned fish. Thus, the sensory cues

detected by the lateral line allowed control fish to respond to water flow with less intensity and

greater temporal variation in their movements, resulting in greater economy of movement.

This is the first study to demonstrate that two ototoxic compounds commonly used to

ablate the lateral line impacts the behavioral profiles and mechanism of rheotaxis in zebrafish.

We propose a novel functional assay for understanding the behavioral impacts of sensory hair

cell ototoxicity, which may be used to supplement future studies exploring lateral line injury,

protection, and recovery. Furthermore, the simplicity of the equipment and precision of the

machine learning analyses used in this assay make it amenable to adaptation for detecting

subtle behavioral changes in a wide variety of animal models.

METHODS

Ethics Statement

This study was performed with the approval of the Institutional Animal Care and Use

Committee of Washington University School of Medicine in St. Louis and in accordance with

NIH guidelines for use of zebrafish.

Zebrafish

Adult zebrafish were raised under standard conditions at 27-29°C in the Washington

University Zebrafish Facility. The wild type line AB* was used for all experiments unless

otherwise stated. Embryos were raised in incubators at 28°C in E3 media (5 mM NaCl, 0.17 mM

KCl, 0.33 mM $CaCl_2$, 0.33 mM $MgCl_2$; ([40])) with a 14:10 hr light:dark cycle. After 4 days post

fertilization (dpf), larvae were raised in 100-200 mL E3 media in 250-mL plastic beakers and fed

rotifers daily. Sex was not considered because it cannot be determined in zebrafish larvae at the

developmental stage used in this study.

Lateral Line Ablation

At 6 or 7dpf, ~15 larval zebrafish were placed into each well of a flat bottom 6-well

polystyrene plate (#351146, Falcon) in 8 mL of E3 media. Treatment animals were placed into

50 μM neomycin trisulfate salt hydrate (N1876-25G, Sigma Aldrich) or 10 μM $CuSO_4$ (451657-

10G, Sigma Aldrich) solutions made in 8 mL of E3 media. The concentration of neomycin used

(50 μM) was chosen to introduce the maximum amount of lateral line hair cell loss with minimal

damage to muscle fibers ([41]). The plate was placed into an incubator at 29°C and exposed to

the treatment for 30 min (neomycin) or 60 min ($CuSO_4$). After exposure, the fish were removed

from treatment, rinsed 3X in media, placed into 8 mL of clean media and allowed to recover in

the incubator for 120 min ($CuSO_4$) or 150 min (neomycin) to standardize the total experimental

time to 180 min. Control fish received no chemical treatments yet underwent the same

procedures as the $CuSO_4$ and neomycin fish. At the end of treatment, the larvae were removed

from the incubator and immediately began rheotaxis behavior trials.

Immunohistochemistry

Lateral line ablation was confirmed via immunohistochemistry on a subset of control and

lesioned fish used in the behavior experiments. Larvae were sedated on ice (5 min, 0°C) then

fixed overnight at 4°C in PO_4 buffer with 4% paraformaldehyde, 4% sucrose, and 0.2 mM $CaCl_2$.

Larvae were rinsed 3X with PBS and blocked for 2 hr at room temperature in PBS buffer with

5% horse serum, 1% Triton-X, and 1% DMSO. Primary antibodies for Otoferlin (HCS-1, mouse

IgG2a, 1:500, Developmental Studies Hybridoma Bank,) and Calbindin (mouse IgG1, 1:1000,

Cat#:214011, Synaptic Systems) were diluted in 1x PBS buffer with 2% horse serum and 0.1%

Triton-X, then incubated with the larvae overnight at 4°C with rotation. Larvae were rinsed 5X

with PBS solution, then placed in secondary antibody (goat anti-mouse IgG2a, Alexa 488,

1:1000, ThermoFisher; goat anti-mouse IgG1, Alexa 647, 1:1000, ThermoFisher) diluted in PBS

with 2% horse serum and incubated for 2 hr at 22°C with rotation. Fish were rinsed 3X with PBS

then incubated with DAPI (1:2000, Invitrogen) in PBS for 20 min at 22°C to label cell nuclei.

Larvae were rinsed 2X with PBS then mounted onto glass slides with elvanol (13% w/v polyvinyl

alcohol, 33% w/v glycerol, 1% w/v DABCO (1,4 diazobicylo[2,2,2] octane) in 0.2 M Tris, pH 8.5)

and #1.5 cover slips.

Confocal Imaging and Hair Cell Quantification

Immunolabeled z-stack images were acquired via an ORCA-Flash 4.0 V3 camera

(Hamamatsu) using a Leica DM6 Fixed Stage microscope with an X-Light V2TP spinning disc

confocal (60 µm pinholes) controlled by Metamorph software. The region of interest tool in

Metamorph was used to select specific neuromasts (~700 x 700 px) from the surrounding area.

Z-stack images of 100 ms exposure were acquired through a 63X/1.4 N.A. oil immersion lens,

0.5 µm z-step size. Excitation for DAPI (405 nm) Alexa 488, and Alexa 647 was provided by

89 North LDI-7 Laser Diode Illuminator on the lowest power setting (20%) that could acquire

images and minimize photobleaching. Confocal images were processed in FIJI (ImageJ, NIH)

software to create maximal intensity z-stack projections with minor exposure and contrast

adjustments. Hair cells were quantified by scrolling through z-stacks and counting only cells that

had both DAPI and HCS-1 labels present. Any hair cells that had pyknotic nuclei (indicated by

condensed DAPI labeling) were not included in the counts. For each fish, the mean number of

hair cells per neuromast for the midline was calculated among multiple locations (L3-5), and for

the supra orbital (SO1, SO2) neuromasts the mean was calculated between the left and right

locations to account for interindividual variation. Hair cell count distributions were tested for

normality; statistical significance was determined using the Kruskal-Wallis test with Dunn's

multiple comparisons post hoc tests using GraphPad Prism 9.2.0.

Experimental Apparatus for Rheotaxis Behavior

A microflume (220 x 100 x 40 mm; Fig. 1a) was constructed in two pieces of translucent

clear resin (RS-F2-GPCL-04, Formlabs) using a high-resolution 3D printer (Form 2, Formlabs)

and joined with two-stage epoxy. Flow collimators (~40mm long) made of ~3mm diameter

plastic straws were placed immediately upstream of the working section in a portion of the flume

bounded on either side by plastic mesh (25 µm) that was cemented to the wall with epoxy (Fig.

1a). Silicone sealant was used to cement a 1 mm-thick layer of plastic on top of flume to prevent

water spillage from the high-pressure side of the flume by covering the pump outflow, first bend,

and flow collimators up to the working section. The low-pressure portion of the flume

downstream from the working section was open at the top to facilitate the addition and removal

of larval zebrafish and water when necessary. Larvae were isolated within the upper 10 mm of

the flume working section to enable reliable video recording by a square (30 x 30 x 10 mm)

arena that was 3D printed from clear resin. Plastic mesh (25 µm) was cemented with epoxy to

the upstream and downstream sides of the removable arena which allowed water to flow

through the working section after the arena was friction seated into the channel at the top of the

flume and immediately adjacent to the flow collimators. An Arduino (UNO R3, Osepp) with a

digital display used custom scripts to control the onset and offset of the pump and camera to

ensure the consistent duration of water flow and video recordings. Flow of constant velocity was

provided by a 6V bow thruster motor (#108-01, Raboesch) inserted into the flume (Fig. 1b) and

modulated by an inline rheostat between the Aurdino and pump. Methylene blue tests indicated

that the flow field was laminar, non-uniform, and stable after < 250 ms of pump onset (data not

shown; Supplementary Fig. 3). The flow field had a lateral gradient along the Y-axis that was

created by a minute boundary layer of null flow immediately adjacent to the walls and gradually

increased (over ~2-5 mm, front to rear) until it became part of the freestream flow field (i.e., blue

arrows in Supplementary Fig. 3).

The flume was placed onto a diffused array of 196 LEDs that emitted infrared (IR, 850

nm) light up through a layer of diffusion material (several Kimwipes© sealed in plastic) and the

translucent flume (Fig. 1b). A monochromatic high-speed camera (SC1 without IR filter,

Edgertronic.com) with a 60 mm manual focus macro lens (Nikon) was placed on a tripod directly

over the flume to record behavioral trials at f16, 1/1000 s, ISO 20000, and 60 or 200 frames s^{-1}

(fps). Videos were saved onto 64 GB SD cards and archived on a 12 TB RAID array (see

below) All source files, scripts, details of apparatus construction, and SOP provided online in the

Open Source Framework repository, DOI:10.17605/OSF.IO/RVYFZ.).

Lateral Line Isolation and Rheotaxis Trials

To isolate the contribution of the lateral line to rheotaxis, we conducted the behavioral

trials under infrared light to eliminate visual cues (Fig. 1a) and reduced linear acceleration cues

to the vestibular system by using a flow stimulus that rapidly accelerated (< 250 ms) to a

constant maximum velocity (9.74 mm s^{-1}). Angular acceleration was not a factor because larval

zebrafish at 6-7dpf cannot detect the angular motion of yaw in the horizontal plane ([31], [32]).

Tactile cues were not possible to selectively block in a non-invasive manner.

The flume was filled with E3 media (28°C) and the arena placed within the flume. A

thermometer was placed into the open portion of the flume to monitor heat generated by the IR

lights and miniature ice packs (2 x 2 cm; -20°C) were used to maintain a consistent temperature

range of 27-29°C. Under IR illumination, a single larval zebrafish was transferred by pipette from

the 6-well plate to the arena within the flume. The swimming activity of the fish was monitored

for ~10 s to ensure that it exhibited the burst-and-glide behavior indicative of normal larval

zebrafish swimming ([42]). The Arduino was used to begin the trial by using a custom script that

triggered the camera to record 10 s of baseline swimming behavior without flow then activate

the pump and record 20 s of swimming behavior under flow. After 30 s had elapsed (no flow =

0-10 s; initial flow = 10-20 s; and final flow = 20-30 s), the pump and camera turned off and the

larvae was removed from the arena. Cohorts of five individual fish from each group (control,

neomycin, or $CuSO_4$) were tested before switching to a new group of fish. This process was

repeated for up to four iterations during each experimental session (n = 15-20 fish per treatment

per session) for a total of 18 sessions (control, n = 248; $CuSO_4$, n = 204; neomycin, n = 222).

3D Markerless Pose Estimation (DeepLabCut)

Equipment: Our behavioral data acquisition and analysis computer ran on the Windows

10 operating system and was based on a Dell Precision 3630 workstation with an Intel Xeon E-

2246G processor, 64 GB RAM, multiple 2TB SSD hard drives, EVGA GFORCE RTX 2080Ti

video card (GPU), dual 24" 4K monitors, Dell Thunderbolt 3 PCIe Card, OWC Mercury Elite Pro

Dock (TB3RSDK24T) - 24TB Thunderbolt 3 Dock and Dual-Drive RAID configured as 12TB

RAID 1.

Installation: Our GPU required Tensorflow 1.12 with the NVIDIA CUDA package to be

installed prior to multi-animal DeepLabCut2.2b8 (maDLC; [43], [44]), Python 3.6, and necessary

dependencies (https://github.com/DeepLabCut/DeepLabCut/blob/master/docs/installation.md).

Detailed tutorials for using maDLC with a single animal are available online

(https://github.com/DeepLabCut/DeepLabCut/blob/master/docs/maDLC_AdvUserGuide.md);

however, the pertinent details of our procedure are as follows.

Dataset Curation: To reduce computational load, all video files were downsampled to

1000 x 1000 px, dead pixels (i.e., black spots) and extraneous portions of the video were

cropped out using the video editor function.

Project Creation: A single animal maDLC project was created and the project

config.yaml file was modified to create and draw a skeleton interconnecting seven unique body

parts (left and right eyes, swim bladder, four points along the tail; Fig. 1a) on each larva. A

curated set of ten videos provided representative examples of target positive rheotaxis

behaviors across experimental treatments, 20 frames were extracted from each video, and the

seven body parts labeled on each frame. The annotated frames were checked for accuracy and

multiple additional skeletal connections between the seven body parts were added to increase

maDLC learning speed and model accuracy.

Pose Estimation: The training dataset used cropped images (400 x 400 px) to reduce

computational loading, Resnet-50 pre-trained network weights, and imgaug data augmentation.

We trained the network until all the loss parameters plateaued at 100,000 iterations then it was

evaluated (PCK values close to 1, RMSE values low) and cross-validated using the default

parameters.

Identity Tracking: The curated videos were analyzed, and the detections assembled into

tracklets using the box method because it provided the best results for our test subjects. The

original videos were overlaid with the newly labeled body parts to correct outliers and

misidentification of body parts in the tracklet data files.

Post Processing: The results were plotted for each video and a new labeled video was

created to double check for labeling accuracy and ensure there was no need to augment the

data set with additional labeled frames. Novel videos were batch processed and analyzed up

through the plot trajectories and label videos step.

Supervised Behavioral Annotation, Classification, & Analysis (SimBA)

Definition of Rheotaxis Behavior: We created a custom Python feature extraction script

(file online) that defined positive rheotaxis as when the larvae swam into the oncoming water

flow at an angle of 0° ± 45° for at least 100 ms (Supplemental Fig. 1). The tail movement and

forward body translation components were used to distinguish active swimming behavior into

the flow from passive backward drift with a body angle of 0° ± 45°.

Installation: SimBAxTF-development version 68 ([45]), Python 3.6, Git, FFmpeg, and all

dependencies. (https://github.com/sgoldenlab/simba/blob/master/docs/installation.md).

Dataset Curation: The pre-processed video files used for maDLC analysis were

converted to AVI format using the SimBA video editor function.

Project Creation: A new SimBA project was created according to the Scenario 1 tutorial:

(https://github.com/sgoldenlab/simba/blob/master/docs/Scenario1.md). The user defined pose

configuration and DLC-multi animal options were selected prior to generating the project config

file. The user-defined pose configuration nomenclature was modified to match the body part and

individual animal labels used in the maDLC config.yaml file. A curated set of ten representative

rheotaxis behavior videos from each treatment group, along with their associated final tracklet

data files created by maDLC, were imported and their frames extracted.

Load Project: The project was loaded and the video parameters (frame rate (fps),

resolution, and pixel measurements (px/mm)) set for each curated video. Outlier correction was

achieved using the swim bladder and tail-3 body parts, movement criterion set to 0.7, and

location criterion set to 1.5 ([45]). Features were extracted using the custom Python feature

extraction script file. Rheotaxis behavior was labeled (i.e., annotations were created for

predictive classifiers) for each curated video, save one that was set aside for validation. The

default training, hyperparameters, and evaluation settings were used to create the model ([45]).

Throughout the earmarked video, the interactive plot function was used to validate the model by

determining the probability threshold for the accurate prediction and minimum duration of a

rheotaxis event. These values became the discrimination threshold (0.5) and minimum behavior

bout length (100 ms) settings used to run the machine model. Naïve videos were processed,

behaviors annotated, and machine results data files archived for analysis.

Data Analysis

Rheotaxis only occurred when fish swam at angles 0° ± 45° under flow conditions;

therefore, the rheotaxis data were compared to data when fish randomly swam at 0° ± 45°

under no flow. However, the entire 0-360° body angle datasets were used to determine if the

fish from each treatment group were randomly distributed under no flow (Fig. 3), and to

determine the X-Y (2D) spatial use under no flow (Fig. 6a) and flow (Fig. 6b) conditions. Time

series of rheotaxis data sampled at 200 and 60 Hz (fps) were quantized into their lowest

common bin size of 100 ms.

Data wrangling and cleaning was performed in R ([46]) with the packages *tidyverse* ([47]),

dplyr ([48]), *plyr* ([49]), and *readbulk* ([50]). Figures and graphs were created with packages *circular*

([51]), *ggplot2* ([52]), and *viridis* ([53]). The Rayleigh statistical tests (V-test) of uniformity for circular

data in a specified mean direction (mu = 0°) and the Watson-Wheeler tests for differences in the

grand mean body angle or angular variance (the test does not specify which parameter differs)

were performed with the package *CircMLE* ([54]). The mean duration, number, total distance

travelled and mean latency to the onset of rheotaxis events were calculated for 10 s bins (i.e.,

none, initial, and final flow) using SimBA. These data were analyzed in R using the generalized

linear mixed models (GLMM) and post hoc t-tests (Satterthwaite method) in the packages

lmerTest ([55]) and *lme4* ([56]). The significance values for fixed effects of these GLMM tests are not

reported by those packages and were run separately using the package *stats* ([46]) for type III

ANOVAs. The package *zoo* ([57]) was used to convert rheotaxis data into times series and the

package *spectral* ([58]) was used to perform the spectral decomposition analysis.

Declaration of interests:

The authors declare no competing financial or non-financial interests.

Acknowledgments: We want to thank Valentin Militchin for the design and construction of the

electronic components of the experimental apparatus, David Lee for editorial support, and Mark

Warchol for feedback on the manuscript. This work was supported by the National Institute on

Deafness and Other Communication Disorders R01DC016066 (L.S.), NIDA R00DA045662

(S.A.G.), NIDA P30 DA048736 (S.R.O.N. and S.A.G.), NARSAD Young Investigator Award

27082 (S.A.G.).

Data Availability

The datasets generated and analyzed during the current study are available in the Open

Science Framework repository, DOI 10.17605/OSF.IO/RVYFZ.

Code Availability

The R code generated for the analyses during the current study are available in the Open

Science Framework repository, DOI 10.17605/OSF.IO/RVYFZ.

REFERENCES

Chagnaud, B. P. & Coombs, S. in *The Lateral Line System* (eds Sheryl Coombs, Horst Bleckmann, Richard R. Fay, & Arthur N. Popper) 151-194 (Springer New York, 2014).

Van Trump, W. J. & McHenry, M. J. The lateral line system is not necessary for rheotaxis in the Mexican blind cavefish (Astyanax fasciatus). *Integr Comp Biol* **53**, 799-809, doi:10.1093/icb/ict064 (2013).

Lyon, E. P. ON RHEOTROPISM. I. — RHEOTROPISM IN FISHES. *American Journal of Physiology-Legacy Content* **12**, 149-161, doi:10.1152/ajplegacy.1904.12.2.149 (1904).

ARNOLD, G. P. RHEOTROPISM IN FISHES. *Biological Reviews* **49**, 515-576, doi:https://doi.org/10.1111/j.1469-185X.1974.tb01173.x (1974).

Suli, A., Watson, G. M., Rubel, E. W. & Raible, D. W. Rheotaxis in larval zebrafish is mediated by lateral line mechanosensory hair cells. *PLoS One* **7**, e29727, doi:10.1371/journal.pone.0029727 (2012).

Pavlov, D. & Tyuryukov, S. The role of lateral-line organs and equilibrium in the behavior and orientation of the dace, Leuciscus leuciscus, in a turbulent flow. *JOURNAL OF ICHTHYOLOGY C/C OF VOPROSY IKHTIOLOGII* **33**, 45-45 (1993).

Pavlov, D. S. & Tjurjukov, S. N. Reactions of dace to linear accelerations. *Journal of Fish Biology* **46**, 768-774, doi:https://doi.org/10.1111/j.1095-8649.1995.tb01600.x (1995).

John C. Montgomery, C. F. B. A. G. C. The lateral line can mediate rheotaxis in fish. *Nature*, 960-963 (1997).

Baker, C. F. & Montgomery, J. C. Lateral line mediated rheotaxis in the Antarctic fish Pagothenia borchgrevinki. *Polar Biology* **21**, 305-309, doi:10.1007/s003000050366 (1999).

Baker, C. F. & Montgomery, J. C. The sensory basis of rheotaxis in the blind Mexican cave fish, Astyanax fasciatus. *Journal of Comparative Physiology A* **184**, 519-527, doi:10.1007/s003590050351 (1999).

Dykgraaf, S. Untersuchungen über die Funktion der Seitenorgane an Fischen. *Zeitschrift für vergleichende Physiologie* **20**, 162-214, doi:10.1007/BF00340757 (1933).

Bak-Coleman, J., Court, A., Paley, D. A. & Coombs, S. The spatiotemporal dynamics of rheotactic behavior depends on flow speed and available sensory information. *J Exp Biol* **216**, 4011-4024, doi:10.1242/jeb.090480 (2013).

Coombs, S., Bak-Coleman, J. & Montgomery, J. Rheotaxis revisited: a multi-behavioral and multisensory perspective on how fish orient to flow. *J Exp Biol* **223**, doi:10.1242/jeb.223008 (2020).

Coffin, A. B., Brignull, H., Raible, D. W. & Rubel, E. W. in *The Lateral Line System* (eds Sheryl Coombs, Horst Bleckmann, Richard R. Fay, & Arthur N. Popper) 313-347 (Springer New York, 2014).

Harris, J. A. *et al.* Neomycin-Induced Hair Cell Death and Rapid Regeneration in the Lateral Line of Zebrafish (Danio rerio). *Journal of the Association for Research in Otolaryngology* **4**, 219-234, doi:10.1007/s10162-002-3022-x (2003).

Ma, E. Y., Rubel, E. W. & Raible, D. W. Notch Signaling Regulates the Extent of Hair Cell Regeneration in the Zebrafish Lateral Line. *The Journal of Neuroscience* **28**, 2261-2273, doi:10.1523/jneurosci.4372-07.2008 (2008).

Hardy, K. *et al.* Functional development and regeneration of hair cells in the zebrafish lateral line. *J Physiol* **599**, 3913-3936, doi:10.1113/JP281522 (2021).

Kniss, J. S., Jiang, L. & Piotrowski, T. Insights into sensory hair cell regeneration from the zebrafish lateral line. *Current Opinion in Genetics & Development* **40**, 32-40, doi:https://doi.org/10.1016/j.gde.2016.05.012 (2016).

19   Pickett, S. B. & Raible, D. W. Water Waves to Sound Waves: Using Zebrafish to Explore
Hair Cell Biology. *J Assoc Res Otolaryngol* **20**, 1-19, doi:10.1007/s10162-018-00711-1
(2019).
20   Olive, R. *et al.* Rheotaxis of Larval Zebrafish: Behavioral Study of a Multi-Sensory
Process. *Front Syst Neurosci* **10**, 14, doi:10.3389/fnsys.2016.00014 (2016).
21   Olszewski, J., Haehnel, M., Taguchi, M. & Liao, J. C. Zebrafish larvae exhibit rheotaxis
and can escape a continuous suction source using their lateral line. *PLoS One* **7**,
e36661, doi:10.1371/journal.pone.0036661 (2012).
22   Oteiza, P., Odstrcil, I., Lauder, G., Portugues, R. & Engert, F. A novel mechanism for
mechanosensory-based rheotaxis in larval zebrafish. *Nature* **547**, 445-448,
doi:10.1038/nature23014 (2017).
23   Hernández, P. P., Moreno, V., Olivari, F. A. & Allende, M. L. Sub-lethal concentrations of
waterborne copper are toxic to lateral line neuromasts in zebrafish (Danio rerio). *Hearing*
Research **213**, 1-10, doi:https://doi.org/10.1016/j.heares.2005.10.015 (2006).
24   Linbo, T. L., Stehr, C. M., Incardona, J. P. & Scholz, N. L. Dissolved copper triggers cell
death in the peripheral mechanosensory system of larval fish. *Environmental Toxicology*
and Chemistry **25**, 597-603, doi:https://doi.org/10.1897/05-241R.1 (2006).
25   Olivari, F. A., Hernández, P. P. & Allende, M. L. Acute copper exposure induces
oxidative stress and cell death in lateral line hair cells of zebrafish larvae. *Brain*
Research **1244**, 1-12, doi:https://doi.org/10.1016/j.brainres.2008.09.050 (2008).
26   Williams, J. A. & Holder, N. Cell turnover in neuromasts of zebrafish larvae. *Hearing*
Research **143**, 171-181, doi:https://doi.org/10.1016/S0378-5955(00)00039-3 (2000).
27   Coffin, A. B., Rubel, E. W. & Raible, D. W. Bax, Bcl2, and p53 differentially regulate
neomycin- and gentamicin-induced hair cell death in the zebrafish lateral line. *J Assoc*
Res Otolaryngol **14**, 645-659, doi:10.1007/s10162-013-0404-1 (2013).
28   Buck, L. M., Winter, M. J., Redfern, W. S. & Whitfield, T. T. Ototoxin-induced cellular
damage in neuromasts disrupts lateral line function in larval zebrafish. *Hear Res* **284**,
67-81, doi:10.1016/j.heares.2011.12.001 (2012).
29   Bak-Coleman, J. & Coombs, S. Sedentary behavior as a factor in determining lateral line
contributions to rheotaxis. *J Exp Biol* **217**, 2338-2347, doi:10.1242/jeb.102574 (2014).
30   Elder, J. & Coombs, S. The influence of turbulence on the sensory basis of rheotaxis. *J*
Comp Physiol A Neuroethol Sens Neural Behav Physiol **201**, 667-680,
doi:10.1007/s00359-015-1014-7 (2015).
31   Beck, J. C., Gilland, E., Tank, D. W. & Baker, R. Quantifying the Ontogeny of Optokinetic
and Vestibuloocular Behaviors in Zebrafish, Medaka, and Goldfish. *Journal of*
Neurophysiology **92**, 3546-3561, doi:10.1152/jn.00311.2004 (2004).
32   Mo, W., Chen, F., Nechiporuk, A. & Nicolson, T. Quantification of vestibular-induced eye
movements in zebrafish larvae. *BMC Neurosci* **11**, 110, doi:10.1186/1471-2202-11-110
(2010).
33   Liao, J. C. The role of the lateral line and vision on body kinematics and hydrodynamic
preference of rainbow trout in turbulent flow. *J Exp Biol* **209**, 4077-4090,
doi:10.1242/jeb.02487 (2006).
34   Hernandez, P. P. *et al.* Sublethal concentrations of waterborne copper induce cellular
stress and cell death in zebrafish embryos and larvae. *Biological Research* **44**, 7-15
(2011).
35   Toro, C. *et al.* Dopamine Modulates the Activity of Sensory Hair Cells. *J Neurosci* **35**,
16494-16503, doi:10.1523/JNEUROSCI.1691-15.2015 (2015).
36   Zhang, Q. *et al.* Synaptically silent sensory hair cells in zebrafish are recruited after
damage. *Nat Commun* **9**, 1388, doi:10.1038/s41467-018-03806-8 (2018).

37 Johansen, J. L., Fulton, C. J. & Bellwood, D. R. Avoiding the flow: refuges expand the
swimming potential of coral reef fishes. *Coral Reefs* **26**, 577-583, doi:10.1007/s00338-
007-0217-y (2007).
38 Webb, P. W. Entrainment by river chub nocomis micropogon and smallmouth bass
micropterus dolomieu on cylinders. *J Exp Biol* **201 (Pt 16)**, 2403-2412 (1998).
39 Burt de Perera, T. B. & Braithwaite, V. A. Laterality in a non-visual sensory modality--the
lateral line of fish. *Curr Biol* **15**, R241-242, doi:10.1016/j.cub.2005.03.035 (2005).
40 Nusslein-Volhard, C. & Dahm, R. *Zebrafish*. (Oxford University Press, 2002).
41 Han, E. et al (2020 Analysis of behavioral changes in zebrafish (Danio rerio) larvae
caused by aminoglycoside-induced damage to the lateral line and muscles.
Neurotoxicology 78, 134-142, doi: 10.1016/j.neuro.2020.03.005
42 Budick, S. A. & O'Malley, D. M. Locomotor repertoire of the larval zebrafish: swimming,
turning and prey capture. *Journal of Experimental Biology* **203**, 2565-2579,
doi:10.1242/jeb.203.17.2565 (2000).
43 Mathis, A. *et al.* DeepLabCut: markerless pose estimation of user-defined body parts
with deep learning. *Nature Neuroscience* **21**, 1281-1289, doi:10.1038/s41593-018-0209-
y (2018).
44 Nath, T. *et al.* Using DeepLabCut for 3D markerless pose estimation across species and
behaviors. *Nature Protocols* **14**, 2152-2176, doi:10.1038/s41596-019-0176-0 (2019).
45 Nilsson, S. R. O. *et al.*, doi:10.1101/2020.04.19.049452 (2020).
894
46 R Core Team (2020). R: A language and environment for statistical computing. R
Foundation for Statistical Computing, Vienna, Austria. URL https://www.R-project.org/.
47 Wickham et al., (2019). Welcome to the tidyverse. Journal of Open Source Software,
4(43), 1686, https://doi.org/10.21105/joss.01686
48 Wickham, H., François, R., Henry, L, and Müller, K. (2021). dplyr: A Grammar of Data
Manipulation. R package version 1.0.3. https://CRAN.R-project.org/package=dplyr
49 Wickham, H. (2011). The Split-Apply-Combine Strategy for Data Analysis. Journal of
Statistical Software, 40(1), 1-29. URL http://www.jstatsoft.org/v40/i01/.
50 Kieslich, P. J., and Henninger, F. (2016). Readbulk: An R package for reading and
combining multiple data files. https://doi.org/10.5281/zenodo.596649
51 Agostinelli C. and Lund, U. (2017). R package 'circular': Circular Statistics (version 0.4-
93). URL https://r-forge.r-project.org/projects/circular/
52 Wickham, H. ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag New York,
2016.
53 Garnier, S. (2018). viridis: Default Color Maps from 'matplotlib'. R package version
0.5.1.https://CRAN.R-project.org/package=viridis
54 Fitak, R. R. and Johnsen, S. (2017). Bringing the analysis of animal orientation data full
circle: model-based approaches with maximum likelihood. Journal of Experimental
Biology, 220, 3878-3882. doi:10.1242/jeb.167056
55 Kuznetsova A, Brockhoff PB, Christensen RHB (2017). "lmerTest Package: Tests in
Linear Mixed Effects Models." Journal of Statistical Software_, *82*(13), 1-26. doi:
10.18637/jss.v082.i13 (URL:https://doi.org/10.18637/jss.v082.i13).
56 Bates, D., Maechler, M., Bolker, B., Walker, S. (2015). Fitting Linear Mixed-Effects
Models Using lme4. Journal of Statistical Software, 67(1), 1-48.
doi:10.18637/jss.v067.i01.
57 Zeileis, A. and Grothendieck, G. (2005). zoo: S3 Infrastructure for Regular and Irregular
Time Series. Journal of Statistical Software, 14(6), 1-27. doi:10.18637/jss.v014.i06
58 Seilmayer, M. (2021). spectral: Common Methods of Spectral Data Analysis. R package
version 2.0. https://CRAN.R-project.org/package=spectral
924

SUPPLEMENTARY INFORMATION

Supplementary Figure 1. Definition of positive rheotaxis behavior. a) Larval zebrafish in the microflume arena performing
rheotaxis under flow as defined by multiple conditions, including b) the water flow stimulus was on; c) the fish body angle was
oriented to 0° ± 45° (green shaded wedge); d) the tail moved laterally every 100 ms; and e) the body of the fish had forward
translation every 100 ms. Note that conditions (d) and (e) were used to discriminate between fish displaying positive rheotaxis and
those passively drifting backward with body angles of 0° ± 45°.

Supplemental Figure 2. Intact lateral line allowed fish to hold their position near the source of the flow. a) The total spatial

use of the arena in the X- dimension (left to right) does not differ among treatments (gray = control, blue = CuSO$_4$, green =

neomycin) or flow conditions (none = dotted lines, flow = solid lines). All fish preferred to occupy the right versus the left side of the

arena. b) in the Y- dimension (front to back) under no flow conditions, the CuSO$_4$ treated fish occupied the center of the arena more

than the control or neomycin treated fish. However, under flow conditions, the lateral line intact (control; gray solid line) fish occupied

the front of the arena, whereas lesioned fish (blue and green solids lines) predominantly occupied the back of the arena.

flow rates (mm s⁻¹)

943 **Supplementary Figure 3. Visualization of methylene blue dye tests within the experimental arena shows a laminar yet non-**
944 **uniform flow field**. a) The vectors are color coded according to the mean of the b) cross-sectional flow values (mm s^{-1}) from top to
945 bottom. Each cross-sectional flow value (b) is the mean of five trials. Fish typically occupied the dark shaded area in the bottom two-
946 thirds of the water column and very rarely swam near the surface.

Supplementary Figure 4. The overall trends and periodic fluctuations in the relative velocity and acceleration in rheotaxis behavior differed among treatment groups. Gray = control (n = 248), blue = CuSO$_4$ (n = 204), green = neomycin (n =222). Spectral decomposition of the observed data (a, d) removed the noise (h, i) to reveal the overall underlying trends (b, e) and the periodicity, or recurring fluctuations (c, f) that occurred during any given 1 s of the experiment. The periodicity waveform peaks indicate the average amount (amplitude), number, direction (positive = increasing; negative = decreasing), and order of occurrence for these cyclic fluctuations as a function of unit time (1 s). The overall trends were that there were no differences in relative (b) velocity or (e) acceleration among groups. The periodic fluctuation in relative velocity (c) and acceleration (f) was greatest in CuSO$_4$ treated fish compared to control or neomycin treated fish. The noise for relative movement (g), mean body angle (j), and mean resultant length (k) are shown.

	No Flow Stimulus 1-10s *theta, rho,* ang var	Flow Stimulus 11-20s *theta, rho,* ang var	Flow Stimulus 21-30s *theta, rho,* ang var
Control	135.5° 0.084 0.916	357.9° 0.687 0.313	1.7° 0.688 0.312
CuSO4	84.8 ° 0.099 0.901	10.8° 0.530 0.470	3.1° 0.611 0.389
Neomycin	86.7° 0.090 0.900	6.1° 0.656 0.344	6.7° 0.655 0.345

960 **Supplementary Table 1.** Grand mean body angle vector parameters. *Theta* = group mean body angle; *rho* = mean length of
961 resultant vector, where 0 = uniform distribution of individual mean body angles, 1 = perfect alignment individual mean body angles;
962 angular variance = 1-*rho*.

963
964
965

	No Flow Stimulus 1-10s test stat, p-value	Flow Stimulus 11-20s test stat, p-value	Flow Stimulus 21-30s test stat, p-value
Control	-0.0595 0.9079	0.686 **< 0.001**	0.6678 **< 0.001**
CuSO4	0.009 0.4276	0.5204 **< 0.001**	0.6105 **< 0.001**
Neomycin	0.0052 0.4566	0.6522 **< 0.001**	0.6507 **< 0.001**

967 **Supplementary Table 2.** Rayleigh test of uniformity (V-test) for an expected grand mean body angle, *mu* = 0°. Among all
968 treatments, groups under no flow were not significantly aligned to 0°, whereas groups under flow conditions were significantly
969 aligned to 0°.

970
971

	CTL-10s p-value	CTL-20s p-value	Cu-10s p-value	Cu-20s p-value	Neo-10s p-value	Neo-20s p-value
CTL-10s test stat	-	0.4747	**2.58e-05**	0.119	**0.0357**	>0.10
CTL-20s test stat	1.49	-	**7.041e-05**	**0.0508**	0.07419	0.4502
Cu-10s test stat	21.128	19.122	-	0.04975	0.05861	**0.0090**
Cu-20s test stat	4.2573	5.9597	6.0014	-	0.311	0.1921
Neo-10s test stat	6.6655	5.2022	5.6737	2.3357	-	0.7364
Neo-20s test stat	0.1251	1.5961	9.4268	3.3000	0.61196	-

973
974 **Supplementary Table 3**. Watson Wheeler test for differences in either grand mean body angle or the distribution of the individual
975 mean angles (*the test does not specify*) among treatment groups under flow conditions.
976

GLMM	Estimate	Std. Error	df	t value	Pr(>|t|)
Stimulus (No flow v Flow 10s)	1.14950	0.10746	1390	10.697	< 2e-16 ***
Stimulus (No flow v Flow 20s)	1.62296	0.10746	1390	15.102	< 2e-16 ***
Treatment (Control v $CuSO_4$)	0.05485	0.11893	2068	0.461	0.64473
Treatment (Control v Neomycin)	0.04574	0.11507	2068	0.398	0.69103
Stim*Treat (CTL-10s v $CuSO_4$-10s)	-0.34246	0.16277	1390	-2.104	0.03556 *
Stim*Treat (CTL-20s v $CuSO_4$-20s)	-0.45260	0.16277	1390	-2.781	0.00550 **
Stim*Treat (CTL-10s v Neo-10s)	-0.29562	0.15749	1390	-1.877	0.06072 .
Stim*Treat (CTL-20s v Neo-20s)	-0.36519	0.15749	1390	-2.319	0.02055 *

ANOVA (III)	Sum Sq	Mean Sq	NumDF	DenDF	F value	Pr(>F)
Stimulus	660.88	330.44	2	1390	216.7689	< 2.2e-16 ***
Treatment	29.83	14.92	2	695	9.7842	6.449e-05 ***
Stim*Treat	15.40	3.85	4	1390	2.5249	0.03926 *

979
980 **Supplementary Table 4**. Generalized Linear Mixed Model with Satterthwaite's method of testing for differences among treatments
981 in the mean duration of rheotaxis events. Type III ANOVA yielded significance values for fixed effects and interactions because the
982 LME4 package in R does not identify them in its output. Significance codes: '***' 0.001, '**' 0.01, '*' 0.05, '.' 0.1
983

GLMM	Estimate	Std. Error	df	t value	Pr(>|t|)
Stimulus (No flow v Flow 10s)	2.65530	0.15538	1390	17.089	< 2e-16 ***
Stimulus (No flow v Flow 20s)	2.91667	0.15538	1390	18.771	< 2e-16 ***
Treatment (Control v $CuSO_4$)	-0.52696	0.21170	1614	-2.489	0.0129 *
Treatment (Control v Neomycin)	0.46123	0.20484	1614	2.252	0.0245 *
Stim*Treat (CTL-10s v $CuSO_4$-10s)	-0.57687	0.23535	1390	-2.451	0.0144 *
Stim*Treat (CTL-20s v $CuSO_4$-20s)	0.05882	0.23535	1390	0.250	0.8027
Stim*Treat (CTL-10s v Neo-10s)	0.09687	0.22772	1390	0.425	0.6706
Stim*Treat (CTL-20s v Neo-20s)	0.05290	0.22772	1390	0.232	0.8163

ANOVA (III)	Sum Sq	Mean Sq	NumDF	DenDF	F value	Pr(>F)
Stimulus	3488.9	1744.46	2	1390	547.3568	< 2.2e-16 ***
Treatment	167.1	83.57	2	695	26.2224	1.049e-11 ***
Stim*Treat	40.0	9.99	4	1390	3.1356	0.01403 *

986
987 **Supplementary Table 5**. Generalized Linear Mixed Model with Satterthwaite's method of testing for differences among treatments
988 in the mean number of rheotaxis events. Type III ANOVA yielded significance values for fixed effects and interactions because the
989 LME4 package in R does not identify them in its output. Significance codes: '***' 0.001, '**' 0.01, '*' 0.05, '.' 0.1
990

GLMM	Estimate	Std. Error	df	t value	Pr(>|t|)
Stimulus (No flow v Flow 10s)	0.70177	0.16338	1721	4.295	**1.84e-05 *****
Stimulus (No flow v Flow 20s)	0.65449	0.16338	1721	4.006	**6.44e-05 *****
Treatment (Control v CuSO$_4$)	0.18141	0.11436	695	1.586	0.11313
Treatment (Control v Neomycin)	0.30691	0.11102	706	2.764	**0.00585 ****

ANOVA (III)	Sum Sq	Mean Sq	NumDF	DenDF	F value	Pr(>F)
Stimulus	243.652	81.037	2	1565	38.8494	**< 2.2e-16 *****
Treatment	8.117	3.796	2	699	3.8913	**0.02086 ***

994 **Supplementary Table 6**. Generalized Linear Mixed Model with Satterthwaite's method of testing for differences among treatments

995 in the total distance travelled during rheotaxis events. GLMM model had no interaction between stimulus and treatment. Type III

996 ANOVA yielded significance values for fixed effects because the LME4 package in R does not identify them in its output.

997 Significance codes: '***' 0.001, '**' 0.01, '*' 0.05, '.' 0.1

GLMM	Estimate	Std. Error	df	t value	Pr(>|t|)
Stimulus (No flow v Flow 10s)	8.460e+00	3.333e-01	2.085e+03	25.386	**< 2e-16 *****
Stimulus (No flow v Flow 20s)	1.762e+01	3.333e-01	2.085e+03	52.882	**< 2e-16 *****
Treatment (Control v $CuSO_4$)	-1.384e+00	3.569e-01	2.085e+03	-3.877	**0.000109 *****
Treatment (Control v Neomycin)	8.802e-03	2.085e+03	2.085e+03	0.025	0.979669
Stim*Treat (CTL-10s v $CuSO_4$-10s)	.904e-01	5.048e-01	2.085e+03	0.972	0.331412
Stim*Treat (CTL-20s v $CuSO_4$-20s)	1.015e+00	5.048e-01	2.085e+03	2.011	**0.044503 ***
Stim*Treat (CTL-10s v Neo-10s)	-4.100e-02	4.884e-01	2.085e+03	-0.084	0.933106
Stim*Treat (CTL-20s v Neo-20s)	8.078e-02	4.884e-01	2.085e+03	0.165	0.868645

1000

ANOVA (III)	Sum Sq	Mean Sq	NumDF	DenDF	F value	Pr(>F)
Stimulus	112476	56238	2	2089	3834.664	**< 2.2e-16 *****
Treatment	345	173	2	2089	11.764	**8.306e-06 *****

1001
1002 **Supplementary Table 7**. Generalized Linear Mixed Model with Satterthwaite's method of testing for differences among treatments
1003 in the mean latency to the onset of first rheotaxis event. GLMM model had no interaction between stimulus and treatment. Type III
1004 ANOVA yielded significance values for fixed effects and interactions because the LME4 package in R does not identify them in its
1005 output. Significance codes: '***' 0.001, '**' 0.01, '*' 0.05, '.' 0.1
1006

GLM	Estimate	Std. Error	t value	Pr(>|t|)
Treatment (Control v CuSO$_4$)	31.9668	4.4820	7.132	**1.21e-12 *****
Treatment (Control v Neomycin)	14.6849	4.0870	3.593	**0.000332 *****
ROI (Back v Center)	4.6943	4.7014	0.998	0.318122
ROI (Back v Front)	-6.3503	4.7014	-1.351	0.176878
ROI (Back v Left)	-13.8344	4.7014	-2.943	**0.003278 ****
ROI (Back v Right)	7.7325	4.7014	-1.645	0.100126
ROI*Treat (CTL-Back v CuSO$_4$-Center)	10.2432	6.3386	1.616	0.106188
ROI*Treat (CTL-Back v Neo-Center)	733.35	5.7799	3.287	**0.001023 ****
ROI*Treat (CTL-Back v CuSO$_4$-Front)	18.9995	6.3386	-4.265	**2.05e-05 *****
ROI*Treat (CTL-Back v Neo-Front)	-27.0351	5.7799	0.146	0.884161
ROI*Treat (CTL-Back v CuSO$_4$-Left)	-23.8271	6.3386	-3.759	**0.000174 *****
ROI*Treat (CTL-Back v Neo-Left)	-13.3969	5.7799	-2.318	**0.020519 ***
ROI*Treat (CTL-Back v CuSO$_4$-Right)	-21.5175	6.3386	-3.395	**0.000695 *****
ROI*Treat (CTL-Back v Neo-Right)	-10.7854	5.7799	-1.866	**0.062127 .**

1007

ANOVA (III)	Sum Sq	Mean Sq	Df	F value	Pr(>F)
Treatment	172565	86283	2	49.727	**< 2.2e-16 *****
ROI	778750	194687	4	112.204	**< 2.2e-16 *****
Treatment*ROI	135133	16892	8	9.735	**1.93e-13 *****

1008
1009 **Supplementary Table 8**. Generalized Linear Model tests for differences in the two-dimensional X-Y spatial use among treatments.
1010 GLM model included fixed effects of stimulus and treatment, and the effect of the interaction between stimulus and treatment.
1011 ANOVA yielded significance values for fixed effects and interactions because the LME4 package in R does not identify them in its
1012 output. Significance codes: '***' 0.001, '**' 0.01, '*' 0.05, '.' 0.1

Variable	Control Min	Control Max	Control Range	CuSO₄ Min	CuSO₄ Max	CuSO₄ Range	Cu-Ctl ΔRange	Cu-Ctl ΔRange (factor)	Neomycin Min	Neomycin Max	Neomycin Range	Neo-Ctl ΔRange	Neo-Ctl ΔRange (factor)
mov	-0.0431	0.0583	0.1014	-0.0724	0.1547	0.2270	0.1257	**2.2400**	-0.0414	0.0441	0.0855	-0.0158	*0.8437*
vel	-0.7762	0.8416	1.6178	-0.7991	1.7910	2.5900	0.9723	**1.6010**	-0.7129	0.7997	1.5126	-0.1051	*0.9350*
acc	-11.8950	12.3300	24.2250	-15.8962	12.1951	28.0914	3.8664	**1.1596**	-8.1738	9.7733	17.9471	-6.2779	*0.7409*
angle	-0.0158	0.0107	0.0265	-0.0087	0.0145	0.0231	-0.0034	*0.8729*	-0.0162	0.0256	0.0418	0.0153	**1.5766**
res L	-0.0052	0.0075	0.0127	-0.0038	0.0023	0.0061	-0.0066	*0.4832*	-0.0026	0.0028	0.0054	-0.0074	*0.4219*

Supplementary Table 9 (Based on Fig. 8c, g, k, o, s). Lateral line ablation by drug treatment (Cu, Neo) shifts the range in overall amplitude of linear (*mov, vel, acc*) and angular (*angle, res L*) seasonality data relative to those of the control (Ctl) group. Movement parameters in which the amplitude increased in treatment fish relative to controls are indicated in **bold**, whereas those that decreased are in *italics*.

		Control	CuSO₄	Cu-Ctl	Cu-Ctl	Neomycin	Neo-Ctl	Neo-Ctl	Control	CuSO₄	Cu-Ctl	Cu-Ctl	Neomycin	Neo-Ctl	Neo-Ctl
Variable	Dominant Peak	Freq (1/s)	Freq	ΔFreq (1/s)	Net Shift Freq 1-3	Freq	ΔFreq (1/s)	Net Shift Freq 1-3	Power	Power	ΔPower (factor)	Net Shift Pwr 1-3	Power	ΔPower (factor)	Net Shift Pwr 1-3
mov	1st	0.1567	0.1528	*-0.0039*	*down*	0.2118	*0.0551*	*down*	0.0169	0.0448	**2.6475**	**up**	0.0337	**1.9964**	**up**
mov	2nd	0.0267	0.1042	*0.0775*	-	0.0139	*-0.0128*	-	0.0101	0.0326	**3.2379**	-	0.0139	**1.3814**	-
mov	3rd	0.2533	0.0208	*-0.2325*	-	0.1632	*-0.0901*	-	0.0082	0.0287	**3.5016**	-	0.0129	**1.5777**	-
vel	1st	0.2533	0.1563	*-0.0971*	*down*	0.2118	*-0.0415*	*down*	1.8444	3.1181	**1.6906**	**up**	6.9068	**3.7447**	**up**
vel	2nd	1.8444	0.2604	*-1.5840*	-	0.4792	*-1.3652*	-	1.5272	2.5076	**1.6419**	-	1.4930	**0.9776**	-
vel	3rd	0.3100	0.2118	*-0.0982*	-	0.1632	*-0.1468*	-	0.7717	1.9820	**2.5683**	-	1.3884	**1.7991**	-
acc	1st	0.2500	0.2604	*0.0104*	*down*	0.2118	*-0.0382*	*down*	363.7884	502.2213	**1.3805**	**up**	938.4990	**2.5798**	**up**
acc	2nd	0.4800	0.3056	*-0.1744*	-	0.4410	*-0.0390*	-	244.0334	402.0855	**1.6477**	-	393.4430	**1.6123**	-
acc	3rd	0.3100	0.1458	*-0.1642*	-	0.2847	*-0.0253*	-	204.4871	304.1066	**1.4872**	-	346.3958	**1.6940**	-
angle	1st	0.0400	0.0382	**-0.0018**	**up**	0.0174	**-0.0226**	**up**	0.0029	0.0042	**1.4571**	**up**	0.0040	**1.3874**	**up**
angle	2nd	0.0700	0.1528	**0.0828**	-	0.1250	**0.0550**	-	0.0026	0.0014	**0.5371**	-	0.0028	**1.0712**	-
angle	3rd	0.1167	0.2083	**0.0917**	-	0.1944	**0.0778**	-	0.0009	0.0009	**1.0510**	-	0.0014	**1.6840**	-
res L	1st	0.0600	0.0313	*-0.0288*	*down*	0.1319	**0.0719**	**up**	0.0002	0.0002	**1.3167**	**up**	0.0001	*0.4215*	*down*
res L	2nd	0.3433	0.0625	*-0.2808*	-	0.2396	**-0.1038**	-	0.0001	0.0001	**1.7516**	-	0.0000	*0.7071*	-
res L	3rd	0.1667	0.1319	*-0.0347*	-	0.4097	**0.2431**	-	0.0001	0.0001	**1.3909**	-	0.0000	*0.6849*	-

Supplementary Table 10 (based on Fig. 9). The primary, secondary, and tertiary dominant frequencies (*Freq*) of the linear (*mov, vel, acc*) and angular (*angle, res L*) movement power spectra shift up or down (*±ΔFreq*) in each of the drug treatments (Cu, Neo) compared to the control (Ctl) group. The peak power (*Pwr*) for all dominant frequencies increases (*+ΔPwr*) in nearly all cases. Values for the net shift in frequency and power were determined by adding the values of the +/- relative shift (ΔFrequency, ΔPower) of all three dominant frequencies and generalizing the overall shift in frequency and power as "up" or "down". The trend of CuSO4 and neomycin ablation results in net downshifts in the frequency and net upshift in power of relative movement, velocity and acceleration and a net upshift in the frequency and power of the mean body angle. However, CuSO4 ablation results in a net downshift in the frequency and net upshift in power, whereas neomycin results in a net upshift in the frequency and net downshift in power. Movement parameters in which the amplitude increased in treatment fish relative to controls are indicated in **bold**, whereas those that decreased are in *italics*.

Linear parameter reference	Angular parameter CCF	Control Lag: (+/-) before, after	Control Heading: right, left Variance: more, less	CuSO$_4$ Lag: (+/-) before, after	CuSO$_4$ Heading: right, left Variance: more, less	Neomycin Lag: (+/-) before, after	Neomycin Heading: right, left Variance: more, less
movement	body angle	before	left	**after**	**left**	*after*	*right*
movement	resultant length	before	less	**before**	**less**	*after*	*less*
velocity	body angle	simultaneous	right	*after*	*right*	**after**	**left**
velocity	resultant length	before	less	*after*	*less*	**before**	**less**
acceleration	body angle	**after**	**left**	*after*	*right*	**after**	**left**
acceleration	resultant length	**before**	**less**	*after*	*less*	**before**	**less**

Supplementary Table 11 (based on Fig. 10). The method, and perhaps mechanism, of lateral line ablation by. CuSO$_4$ and neomycin results in two distinct type of rheotaxis kinematics that are different from control fish. Each method has the opposite effect on the linear and angular movement parameters during rheotaxis and neither matches the phenotype of fish with an intact lateral line. Behavioral phenotypes in which an above average increase in the indicated linear variable was strongly correlated with a leftward heading that will occurr later and a decrease in angular variance that occurred previously are indicated in **bold**. Phenotypes in which an above average increase in the indicated linear variable was strongly correlated with a rightward heading and a decrease in angular variance that will occur later are indicated in *italics*.