SARS-CoV-2 RdRp uses NDPs as a substrate and is able to incorporate NHC

into RNA from diphosphate form molnupiravir

Maofeng Wang, Cancan Wu, Nan Liu, Fengyu Zhang, Hongjie Dong, Shuai Wang,

Min Chen, Xiaoqiong Jiang & Lichuan Gu*

State Key Laboratory of Microbial Technology, Shandong University, 72 Binhai Road,

Qingdao 266237, P.R. China

*To whom correspondence should be addressed: L.G. (e-mail: lcgu@sdu.edu.cn)

Abstract

The coronavirus disease 2019 (COVID-19) has been ravaging throughout the world for more than two years and has severely impaired both human health and the economy. The causative agent, severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) employs the viral RNA-dependent RNA polymerase (RdRp) complex for genome replication and transcription, making RdRp an appealing target for antiviral drug development. Here, we reveal that RdRp can recognize and utilize nucleoside diphosphates (NDPs) as a substrate to synthesize RNAs with an efficiency of about two thirds of using nucleoside triphosphates (NTPs) as a substrate. NDPs incorporation is also template-specific and has high fidelity. Moreover, RdRp can incorporate β-d-N4-hydroxycytidine (NHC) into RNA while using diphosphate form molnupiravir (MDP) as a substrate. We also observed that MDP is a better substrate for RdRp than the triphosphate form molnupiravir (MTP).

Introduction

The pandemic of coronavirus disease 2019 (COVID-19) has severely impacted human health and the global economy since the outbreak in late 2019[1,2]. The causative agent, severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) is homologous to 2002 SARS-CoV with a genome sequence similarity of 79%[3,4], forming a sister clade to SARS-CoV and considered a newly β-coronavirus[5]. As of the date of writing, more than 491.75 million infections and more than 6.17 million deaths have been confirmed (https://www.who.int/emergencies/diseases/novel-coronavirus-2019/situation-reports). Although a variety of vaccines have been developed to contain the pandemic, the rapidly spreading mutant strains presses need to develop effective medication or treatment to counter the virus[6-9].

SARS-CoV-2 is a positive-sense single-stranded RNA virus[5] with a large genome of approximately 30 kb organized in 14 open reading frames[4,10-12]. Replication of the genome and transcription of genes are all dependent on a protein complex known as RNA-dependent RNA polymerase (RdRp). The RdRp of SARS-CoV-2 contains a catalytic subunit (non-structural protein 12, nsp12) and two accessory subunits (nsp7 and nsp8)[13,14]. Processivity is enhanced by the presence of nsp 7/8 and bound to nsp 12 in a 1:2:1 stoichiometry[15,16]. Due to the significant role of RdRp and the lack of homologues in humans, RdRp has become the most appealing target for anti-coronavirus drug development[17,18].

To date, the most promising RdRp inhibitors are nucleotide/nucleoside analogs (NA). NA prodrugs are designed to be metabolized to the active 5'-triphosphate form (5'-TP)

once inside cells[19], thus they can compete with endogenous nucleotides for

incorporation into newly synthesized viral RNAs, resulting in viral genomes lethal

mutation or inactivation[20]. Remdesivir is a broad antiviral drug approved for the

treatment of SARS-CoV-2, which can be incorporated into RNA to sterically hinder

RdRp, thereby blocking viral RNA synthesis[21]. Other NAs (favipiravir, ribavirin) are

integrated into nascent viral RNA at high rates but are not recognized as endogenous

nucleotides in subsequent rounds of replication, increasing the mutation rate and

resulting in an inviable genome, a process known as "lethal mutagenesis"[22-24]. The

recently approved NA antiviral drug molnupiravir significantly increases the

frequency of viral RNA mutations and impairs SARS-CoV-2 in both animal models

and humans[25,26].

By far none of these RdRp targeting drugs is effective enough to fulfill the need for

treatment of COVID-19. To develop a better RdRp targeting drug we need to better

understanding of both the properties of RdRp and how these inhibitors function.

Previous studies showed that some DNA polymerases[27,28] and HIV reverse

transcriptase[29] can use deoxynucleoside diphosphates (dNDPs) as a substrate to

synthesize DNA, and *Escherichia coli* RNA polymerase[30] use nucleoside

diphosphates (NDPs) to synthesize RNA. Although both dNDPs and NDPs showed

very weak activity in these experiments, the data prompted us to test if NDPs are an

active substrate for RdRp. In this study, we found that SARS-CoV-2 RdRp can

recognize and utilize NDPs as a substrate to synthesize RNAs with an efficiency of

about two thirds of using nucleoside triphosphates (NTPs) as a substrate. In addition,

we proved that molnupiravir is not only fully functionable in its diphosphate form but

also more active than in triphosphate form. This could imply a new strategy for

designing COVID-19 treatment nucleoside analogue drugs with more efficacy and

fewer side effects.

Results

RdRp synthesizes RNA by using both NTPs and NDPs as a substrate

Since RdRp is by far one of the most attractive targets for anti-coronavirus drug

development, a thorough characterization of RdRp polymerase activity will, without

doubt, greatly contribute to discovering the effective strategy for drug screening[17,18].

The RdRp of SARS-CoV-2 was purified by assembling nsp12, nsp7 and nsp8 (Fig.1a).

Based on previous research, SARS-CoV-2 RdRp polymerase activity was measured

by conducting a conventional RNA elongation assay[31]. When NTPs was added,

elongation catalyzed by RdRp complex occurred on the RNA duplex, resulting in an

intact double-stranded RNA product (Fig.1c).

To test if RdRp uses NDPs or NMPs as a substrate, we performed an RNA extension

assay by using NTPs, NDPs or NMPs as substrate, respectively. The result revealed

that RdRp synthesized RNA products of the same length regardless of NTPs or NDPs

as substrate (Fig.1d). To examine the template specificity of this RNA synthesis, we

challenged 13+23 nt RNA with various NDPs (Fig.1b). Only the cognate ADP

promoted RNA extension, indicating that NDPs incorporation is template-specific

(Fig.1e).

This result also raised the question of whether the RNA made from NDPs has the

right sequence. Hence, two experiments were performed: single-clonal sequencing of

ligated plasmids, and high-throughput sequencing of PCR products. First, the cDNA

was synthesized through reverse transcription of the RNA products, and then

amplified by 10 cycles of PCR with primers containing the restriction sites of XhoI

and BamHI. PCR products were then ligated into the vector pET-15b and used to

transform *E. coli* DH5α strain. Plasmids extracted from the positive clones were

sequenced by Sanger sequencing. The sequencing results indicated that the RNA

synthesized from NDPs has the right sequence (Fig.1f). Second, the cDNA amplified

by 10 cycles of PCR was sequenced by high-throughput sequencing (Fig.1g). At least

50,000 sequences were obtained for each product (SI). The ADP, GDP, CDP and UDP

(purchased from Sigma) used in these experiments were characterized by mass

spectrometry respectively to make sure these NDPs were not contaminated by NTPs

(S1).

NDPs are nearly two thirds as efficient as NTPs as a substrate of RdRp

Previous studies indicated that although some DNA polymerases and RNA

polymerases are able to use diphosphate form substrate. The activity of the

diphosphate form substrate, however, is much lower than that of the triphosphate form

substrate[28-30]. To determine if this is also the case for RdRp we performed a kinetics

study by using NTPs and NDPs as a substrate respectively. The 13+23 nt RNA duplex

was used as a template and the elongation of product over time was observed (Fig.2a).

Kinetics curves indicated that the relative incorporation efficiency of NDPs is about

two thirds (67.94% ± 2.66%) of that of NTPs (Fig.2b). These data suggested that the

incorporation rate for NDPs is slower than that of NTPs. Subsequently, we decided to

test if all four NDPs equally contribute to the slower incorporation rate. A 13+26 nt

RNA duplex was synthesized for the assay testing the incorporation efficiency for

each NDP (Fig.2c). Each reaction system contains three NTPs and the NDP of interest

(Fig.2d). Unexpectedly, our data showed that substitution of any single NTP by NDP

has no observable effect on the efficiency of RNA synthesis (Fig.2e). This may imply

that the speed of RNA synthesis is only affected by continuous NDP incorporation but

not the intermittent insertion.

SARS-CoV-2 RdRp incorporates NHC into RNA by using diphosphate form

molnupiravir as a substrate

Since RdRp has become one of the most important targets for the development of

antiviral drugs, many nucleoside analogue drugs have been constructed to target

RdRp for COVID-19 treatment[24,32-34]. Among all these chemicals reported by far,

molnupiravir has been regarded as a promising drug candidate. The molecular

mechanism of molnupiravir was determined recently. Once ingested by patients,

molnupiravir undergoes stepwise phosphorylation to yield the active nucleoside

triphosphate analogue (β-d-N4-hydroxycytidine (NHC) triphosphate, or MTP). The

RdRp then uses MTP as substrate and incorporates NHC monophosphate into RNA at

the position where the nucleotide in the template is G or A, resulting in "error

catastrophe" and virus death[26,35,36] (Fig.3a).

Although the mechanism of molnupiravir has been well established, our finding of-

RdRp's ability to utilize NDPs as a substrate still raises a question: is NHC

diphosphate (MDP) also an active form and recognized by RdRp as a substrate? To

address this question, the 13+26 nt RNA duplex was used to perform the RNA

extension assay with NDPs and MDP as the substrates in comparison with the

extension using NTPs and MTP as the substrates. The 13+26 nt RNA duplex allows

ten nucleotides (nt) of extension. The 13+26 nt RNA duplex contanis two adjacent G

and two adjacent A, allowing NHC incorporation into the RNA (Fig.3b). In this case,

when cytidine triphosphate (CTP) or uridine triphosphate (UTP) is replaced by MDP,

the incorporation of M does not hinder the incorporation of the next subsequent

nucleotide, and RdRp still completes the extension reaction. Similar results were

obtained using MTP as the substrate (Fig.3c). Therefore, it is reasonable to speculate

that both MTP and MDP are active forms of monulpiravir in humans.

MDP is a better substrate than MTP

Since both MTP and MDP are active substrates of RdRp we performed an assay to

figure out which one is the better substrate. By observing their efficiency in reaction,

we found that when the template is G, the relative incorporation efficiency of MTP is

0.78, while MDP is 0.87 in comparison to CTP. When the template is A, the relative

incorporation efficiency of MTP is 0.58, while MDP is 0.77 in comparison to UTP

(Fig.3d). In order to obtain a reliable result, the experiment was repeated six times

with two batches of RdRp. The results indicated MDP is the better substrate for RdRp

in comparison to MTP. MTP and MDP were characterized by HPLC and mass

spectrometry to make sure they are the right pure compounds (S2).

Next, we tried to determine if MDP is also more efficient in introducing mutations

than MTP. The products of RdRp-catalyzed extension reactions containing MTP or

MDP using 13+29 nt RNA as a template (Fig. 3e) were reverse-transcribed and

PCR-amplified, and the number of mutations was counted by high-throughput

sequencing.　The statistical results were shown in Table 1. Similar mutation ratios

were found when the same concentrations of MTP and MDP were added. Higher ratio

of MTP or MDP to CTP associates with higher mutation rate, with MDP resulting in

slightly more mutations.

Table.1

substrate	NTP	50 μM CTP 50 μM MTP	50 μM CTP 50 μM MDP	5 μM CTP 50 μM MTP	5 μM CTP 50 μM MDP	0.8 μM CTP 50 μM MTP	0.8 μM CTP 50 μM MDP
all sequences	67883	64316	61673	59588	70906	59129	56214
no mutation （TCCCTTT）	67105	63626	61054	58386	69443	57831	54945
% of no mutation	98.85%	98.93%	99.00%	97.98%	97.94%	97.80%	97.74%
mutations*	778	690	619	1202	1463	1298	1269
% of mutation	1.15%	1.07%	1.00%	2.02%	2.06%	2.20%	2.26%

*The mutation resulting from the incorporation of M is shown as: TCCCTTC,

TTCCTTT, TACCTTT, TCCCCTT, TCACTTT, TCCCTCT, TCCCTAT etc.

NDPs are a good substrate for SARS-CoV-2 RdRp but not other virus RNA

polymerases

To test if other virus RNA polymerases also use NDPs as a substrate, we purified

SARS-CoV RdRp, NS5 protein of ZIKA virus and T7 RNA polymerase (RNAP) and

measured their activities when NDPs were used as s substrate. Results show that

amount these RNA polymerases tested SARS-CoV-2 RdRp had the highest activity

when using NDPs as a substrate. SARS-CoV RdRp could also use NDPs as a

substrate, but the product band seam a double RNA strand shorter than expectation

implying stalled extension. NS5 of ZIKA virus did not produce a well-defined product

band when using NDPs as a substrate (Fig.4a). The transcription of T7 RNAP was

measured based on fluorescence technology. We found that except for the case in

which ATP was replaced by ADP and weak transcription occurred, any other NTP

replaced by NDP would inhibit the transcription completely. (Fig.4b).

Discussion

Since its outbreak in late 2019, the COVID-19 pandemic has been plaguing people for

two years with no sign of abating[3,11]. Although many kinds of vaccines have been

developed to suppress the pandemic, they are not sufficient to withhold the spread of

SARS-CoV-2[37]. The development of potent antivirals against SARS-CoV-2 is still an

urgent need[6,38].

Since RNA-dependent RNA polymerase (RdRp) is critical for viral genome

replication and transcription and conserved in RNA virus species, RdRp has become

one of the most appealing targets for antivirals development[17,31,38,39]. Thoroughly

characterization of RdRp both structurally and biochemically would no doubt be

beneficial to drug development.

The substrate usage of RdRp is astonishing. To the best of our knowledge, most RNA

polymerases, including DNA-dependent and RNA-dependent RNA polymerases, use

nucleoside triphosphates (NTPs) as a substrate to synthesize RNA under the guidance

of a template strand. We found that SARS-CoV-2 RdRp synthesizes RNA with

comparable efficiency using NTPs and NDPs as a substrate, and other RNA viruses

that can utilize NDPs as a substrate have also been found. We speculate that the

ability to use NDPs as a substrate may be an advantage of RNA virus during infection.

During later stages of infection, host cells may not be able to produce enough ATP

and other NTPs. By using NDPs as a substrate, SARS-CoV-2 can continue the process

of genome replication and assembly of progeny viruses, while most metabolic

activities stop.

It has long been a general knowledge that to become active against virus nucleoside

analogue drugs must undergo stepwise addition of phosphate groups to become the

triphosphate form. In order to produce this outcome, this type of antivirals must be

recognized by three kinds of kinases[40]. This raises a concern that the triphosphate

form of antivirals may be incorporated into host mRNA[36]. Even worse, mutagenic

ribonucleoside analogs could be reduced into the 2'-deoxyribonucleotide form by host

ribonucleotide reductase and then incorporated into DNA[35,36]. Our finding that RdRp

also uses MDP as a substrate could largely resolve these concerns. Since the

diphosphate form is also active, there would be no issue regarding whether the

prodrugs can become triphosphate form inside host cells. Furthermore, nucleoside

diphosphate analogues, which can evade nucleoside diphosphate kinase, would cease

to have the possibility of being incorporated into mRNA or DNA. The best strategy is

to use the membrane-permeable nucleotide diphosphate analog for COVID-19

treatment[40,41]. The advantage of this strategy is that the drug is not only able to bypass

all the phosphorylation steps and has no risk of incorporation into host mRNA or

DNA (Fig.5).

Materials and Methods

Protein expression and purification

Preparation of SARS-CoV-2 RdRp, composed of nsp12, nsp7 and two copied of the

nsp8 subunits, was carried out as described in ref.[31]. The SARS-CoV-2 *nsp12* and

nsp7 gene were cloned into a modified pET-21b vector with the C-terminus

possessing a 6×His-tag. The *nsp8* gene was cloned into the modified pET-32a vector

with the N-terminus possessing a trx-His$_6$-tag and PreScission Protease site. The

nsp12-pET-21b and *nsp8*-pET-32a plasmids were transformed into *E. coli* BL21

(DE3) and the transformed cells were cultured at 37 °C in LB with a final

concentration of 100 μg/ml ampicillin. The *nsp7*-pET-21b plasmid was transformed

into *E. coli* Rosetta-gami2 (DE3) and the transformed cells were cultured at 37 °C in

LB contanning a final concentration of 100 μg/ml ampicillin and 25 μg/ml

chloramphenicol.

Bacterial cultures were incubated by shaking at 200 rpm, 37 °C to an OD600 of 0.8,

then the temperature was lowered to 16 °C and a final concentration of 0.2 mM of

isopropyl β-D-1-thiogalactopyranoside (IPTG) was added to induce protein

expression for 20 h. Subsequently, the cells were harvested by centrifugation at 5000

×*g* for 15 min at 4 °C. The pellet was resuspended in lysis buffer containing 25 mM

TrisHCl pH 8.0, 250 mM NaCl, 4 mM MgCl$_2$, 10% Glycerol and homogenized with a

high-pressure cell disrupter at 4 °C. The lysate was centrifuged at 25,200 ×*g* for 50

min at 4 °C, and the supernatant was then loaded onto Ni-NTA affinity

chromatography column for purification. Nsp12, nsp7 were eluted with elution buffer

(25 mM TrisHCl pH 8.0, 200 mM NaCl, 4 mM $MgCl_2$, 250 mM Imidazole). Nsp8

was subject to on-column tag cleavage by PPase (PreScission Protease) and then

eluted by the lysis buffer. Next, all proteins were purified by ion exchange

chromatography (Source 15Q HR 16/10, GE Healthcare) and size exclusion

chromatography (Superdex 200 10/300GL, GE Healthcare) with 25 mM TrisHCl pH

8.0, 200 mM NaCl, 4 mM $MgCl_2$.

For nsp12-nsp7-nsp8 complex assembly, protein samples were mixed with the molar

ratio of nsp7:nsp8:nsp12=2:2:1 at 4 °C overnight. The incubated RdRp complex was

then concentrated with a 100 kDa molecular weight cut-off centrifugal filter unit

(Millipore Corporation) and then further purified by size exclusion chromatography

using a Superdex 200 10/300 GL column in 25 mM TrisHCl pH 8.0, 200 mM NaCl, 4

mM $MgCl_2$. SARS-CoV RdRp was purified by the same procedure. The NS5 RdRp

domain of ZIKV was purified as previously described[42].

Preparation of fluorescently labeled RNA for polymerase activity

RNA template-product duplex was designed according to the published SARS-CoV-2

RNA extension assays[31]. FAM labeled 13 nt oligonucleotide with the sequence of

FAM - GCUAUGUGAGAUU and the 23 nt complementary RNA strand with the

sequence of AGUUAACUUUAAUCUCACAUAGC were synthesized for

polymerase activity assay. A 26 nt RNA strand with the sequence

UAGCUGGCCUUAAAAUCUCACAUAGC was synthesized for substrate efficiency

assay. And a 29 nt RNA strand with the sequence AGUUAACUUUAAAGGGAAUCUCACAUAGC were used for the detection of mutation rates due to the incorporation of MTP and MDP into RNA. All unmodified and 5' FAM-labeled RNA oligonucleotides were purchased from Tsingke Biotechnology Co.,Ltd. The RNA strands were mixed in equal molar ratio in DEPC water, annealed by heating it to 95 °C for 10 min and gradually cooling to room temperature to make the RNA duplexes.

RdRp polymerase activity assays

DEPC-treated water was used in the preparation of all solutions. RdRp at final concentration of 2 μM was incubated with 200 nM 13+23 nt RNA duplex and 50 μM NTPs (or NDPs / NMPs) in a 20 μl reaction buffer containing 20 mM TrisHCl pH8.0, 10 mM KCl, 10 mM MgCl$_2$, 0.01% Triton-X100, 1 mM DTT for 1 min at 37 °C[31], the reactions were stopped with 2 × stop buffer (10 M urea, 50 mM EDTA). Eventually, glycerol was added to a concentration of 6.5%. 10 μl RNA product for each reaction was resolved on 20% denaturing polyacrylamide-urea gels and imaged with a Tanon-5200 Multi Fluorescence Imager. The experiments described were performed in triplet, unless specified otherwise.

RNA extension assays with NDPs as substrate

RdRp of different viruses at final concentration of 2 μM was incubated with 200 nM 13+23 nt RNA duplex and 50 μM NTPs (or NDPs/NMPs/ADP/GDP/CDP/UDP) in a 20 μl reaction buffer containing 20 mM TrisHCl pH8.0, 10 mM KCl, 10 mM MgCl$_2$, 0.01% Triton-X100, 1 mM DTT for 1 min at 37 °C.

Sanger sequencing

The RNA product, primer1 (CCGCTCGAGCGGAGTTAACTTT) and dNTPs were gently mixed at 65 °C for 10 min. The mixture was cooled down to 24 °C then 10× MMulV buffer and MMulV reverse transcriptase were added and incubated for 2 h to obtain cDNA. Thereafter, the reverse transcriptase was inactivated at 70 °C for 15 min. Primer2 (CGCGGATCCGCGGCTATGTGAGAT) was then added, and the product was amplified by PCR catalyzed by the high-fidelity DNA polymerase pfu-Phusion. The PCR product was digested with XhoI and BamHI and ligated into pET-15b vector, and transformed into *E-coli* DH5α to obtain clones. Plasmids from positive clones were then sequenced with T7 universal primers.

High-throughput sequencing

The dsRNA products were mixed with primer1 (acgatgcaaagtctcgacaaatggtcgataccaattcaCCGCTCGAGCGGAGTTAACTTT) and dNTPs at 65 °C for 10 min and then cooled down to 24 °C for primer annealing. 10 × MMulV buffer, RNase inhibitor, and MMulV reverse transcriptase were added to the above system at 24 °C for 2 h to obtain cDNA, and then the enzymes were inactivated at 70 °C for 15 min. Subsequently, primer2 (cagataaactataattcctaatcgcgaggtggcactgcaaCGCGGATCCGCGGCTATGTGAGA) and high-fidelity DNA polymerase pfu-Phusion were added to perform PCR amplification. PCR products were then purified by agarose gel electrophoresis for high-throughput sequencing (Sangon Biotech).

Substrate efficiency assay

RNA extension was performed by incubating 2 μM RdRp, 250 nM 13+23 nt RNA

duplex, 50 μM substrate (which are NTPs, NDPs) in a 20 μl reaction system at 37 °C

for 0, 10, 20, 30, 40, 50, 60 s. RdRp catalyzed reactions each containing three NTPs

and one NDP (ADP GTP CTP UTP, ATP GDP CTP UTP, ATP GTP CDP UTP and

ATP GTP CTP UDP) and 13+26 nt RNA duplex was performed for 1 min. The

reactions were stopped with 2 × stop buffer. Grayscale analysis was then carried out

using ImageJ and ilustrated by using OriginPro 8 (OriginLab, USA).

RNA extension assays with MDP as substrate

A 13+26 nt RNA duplex was used in this assay (Fig.5e). The 5' end of the RNA

product strand was labeled with a 6-carboxyfluorescein (FAM) group, which allows

us to monitor RNA elongation. RdRp can incorporate NHC monophosphate into RNA

at the position where the nucleotide in the template is G or A. MDP and MTP were

purchased from MedChemExpress.

For RNA extension RdRp at final concentration of 2 μM was incubated with 200 nM

13+26 nt RNA duplex and 50 μM ATP GTP UTP or CTP GTP UTP with or without

50 μM MTP/MDP in 20 μl reaction buffer for 1min at 37 °C. The reactions were

stopped with 2 × stop buffer. 10 μl RNA product for each reaction was resolved on

20% denaturing polyacrylamide-urea gels and imaged with a Tanon-5200 Multi

Fluorescence Imager to confirm the extension. Grayscale analysis was then carried

out using ImageJ and ilustrated by using OriginPro 8 (OriginLab, USA).

Incorporation mutation rate detection

RdRp at final concentration of 2 μM was incubated with 200 nM 13+29 nt RNA

duplex and 50 μM ATP GTP UTP, 800 nM or 5 μM or 50 μM CTP and 50 μM

MTP/MDP in 20 μl reaction buffer for 1min at 37 ℃. Mutation rates were determined

by High-throughput sequencing as mentioned above.

Fluorescence assay for T7 RNA polymerase transcription

Transcription of T7 RNA polymerase was measured by a fluorescence assay which

employs a pair of nucleic acid probes with fluorescent and quenching groups

respectively to detect mRNA production. When the probes pair with the mRNA FAM

and quenching group gets closer and the fluorescence produced by FAM will be

quenched. The change of the fluorescent signal is proportional to the mRNA

production.

Data availability

All data generated or analyzed during this study are included in this published article

(and its Supplementary Information files).

Acknowledgements The authors thank Yan Zhang at Zhejiang University for offering

SARS-CoV-2 RdRp gene, Peihui Wang at Cheeloo College of Medicine, Shandong

University for offering SARS-CoV RdRp gene, Haitao Yang at ShanghaiTech

University for offering ZIKV NS5 gene, Wuxi Biortus Biosciences Co. Ltd for

providing SARS-CoV-2 RdRp expressed in insect cells, Carina Muyao Gu at

CBe-Learn School for helping with the English expression of the manuscript. This

work was supported by Shandong Provincial Key Research and Development

Program (2020CXGC011305).

Author contributions M.W. and L.G. conceived, designed the experiments and analyzed the data. M.W. performed functional assays of RdRp. M.W., C.W., and X.J. participated in expression, purification of the SARS-CoV-2 RdRp complex and other proteins. N.L., F.Z., H.D., S.W., M.C. helped with the experiments and participated in figure and manuscript preparation. M.W. and L.G. wrote the manuscript.

Competing interests The authors declare no competing interests.

References

1   Chen, Y., Liu, Q. & Guo, D. Emerging coronaviruses: Genome structure, replication, and pathogenesis. *J Med Virol* **92**, 418-423, doi:10.1002/jmv.25681 (2020).

2   Zhu, N. *et al.* A Novel Coronavirus from Patients with Pneumonia in China, 2019. *N Engl J Med* **382**, 727-733, doi:10.1056/NEJMoa2001017 (2020).

3   Lu, R. *et al.* Genomic characterisation and epidemiology of 2019 novel coronavirus: implications for virus origins and receptor binding. *The Lancet* **395**, 565-574, doi:10.1016/s0140-6736(20)30251-8 (2020).

4   Chan, J. F. *et al.* Genomic characterization of the 2019 novel human-pathogenic coronavirus isolated from a patient with atypical pneumonia after visiting Wuhan. *Emerg Microbes Infect* **9**, 221-236, doi:10.1080/22221751.2020.1719902 (2020).

5   Coronaviridae Study Group of the International Committee on Taxonomy of, V. The species Severe acute respiratory syndrome-related coronavirus: classifying 2019-nCoV and naming it SARS-CoV-2. *Nat Microbiol* **5**, 536-544,

389   doi:10.1038/s41564-020-0695-z (2020).

390 6 Liu, C. *et al.* Research and Development on Therapeutic Agents and Vaccines

391   for COVID-19 and Related Human Coronavirus Diseases. *ACS Cent Sci* **6**,

392   315-331, doi:10.1021/acscentsci.0c00272 (2020).

393 7 Harvey, W. T. *et al.* SARS-CoV-2 variants, spike mutations and immune

394   escape. *Nat Rev Microbiol* **19**, 409-424, doi:10.1038/s41579-021-00573-0

395   (2021).

396 8 Chen, J., Gao, K., Wang, R. & Wei, G. W. Prediction and mitigation of

397   mutation threats to COVID-19 vaccines and antibody therapies. *Chem Sci* **12**,

398   6929-6948, doi:10.1039/d1sc01203g (2021).

399 9 Chen, J., Gao, K., Wang, R. & Wei, G. W. Revealing the Threat of Emerging

400   SARS-CoV-2 Mutations to Antibody Therapies. *J Mol Biol* **433**, 167155,

401   doi:10.1016/j.jmb.2021.167155 (2021).

402 10 Rudd, C. E. GSK-3 Inhibition as a Therapeutic Approach Against SARs CoV2:

403   Dual Benefit of Inhibiting Viral Replication While Potentiating the Immune

404   Response. *Front Immunol* **11**, 1638, doi:10.3389/fimmu.2020.01638 (2020).

405 11 Wu, F. *et al.* A new coronavirus associated with human respiratory disease in

406   China. *Nature* **579**, 265-269, doi:10.1038/s41586-020-2008-3 (2020).

407 12 Zhang, Y. Z. & Holmes, E. C. A Genomic Perspective on the Origin and

408   Emergence of SARS-CoV-2. *Cell* **181**, 223-227,

409   doi:10.1016/j.cell.2020.03.035 (2020).

410 13 Wu, C. *et al.* Analysis of therapeutic targets for SARS-CoV-2 and discovery of

potential drugs by computational methods. *Acta Pharm Sin B* **10**, 766-788,

doi:10.1016/j.apsb.2020.02.008 (2020).

14 Subissi, L. *et al.* One severe acute respiratory syndrome coronavirus protein

complex integrates processive RNA polymerase and exonuclease activities.

Proc Natl Acad Sci U S A **111**, E3900-3909, doi:10.1073/pnas.1323705111

(2014).

15 Kirchdoerfer, R. N. & Ward, A. B. Structure of the SARS-CoV nsp12

polymerase bound to nsp7 and nsp8 co-factors. *Nat Commun* **10**, 2342,

doi:10.1038/s41467-019-10280-3 (2019).

16 Hillen, H. S. *et al.* Structure of replicating SARS-CoV-2 polymerase. *Nature*

584, 154-156, doi:10.1038/s41586-020-2368-8 (2020).

17 McKee, D. L., Sternberg, A., Stange, U., Laufer, S. & Naujokat, C. Candidate

drugs against SARS-CoV-2 and COVID-19. *Pharmacol Res* **157**, 104859,

doi:10.1016/j.phrs.2020.104859 (2020).

18 Chu, C. K. *et al.* Antiviral Activity of Nucleoside Analogues against

SARS-coronavirus (SARS-CoV). **17**, 285-289,

doi:10.1177/095632020601700506 (2006).

19 Stein, D. S. & Moore, K. H. P. Phosphorylation of Nucleoside Analog

Antiretrovirals: A Review for Clinicians. *Pharmacotherapy: The Journal of*

Human Pharmacology and Drug Therapy **21**, 11-34,

doi:https://doi.org/10.1592/phco.21.1.11.34439 (2001).

20 Agostini, M. L. *et al.* Small-Molecule Antiviral β-d-N (4)-Hydroxycytidine

Inhibits a Proofreading-Intact Coronavirus with a High Genetic Barrier to

Resistance. *Journal of virology* **93**, e01348-01319, doi:10.1128/JVI.01348-19

(2019).

21   Picarazzi, F., Vicenti, I., Saladini, F., Zazzi, M. & Mori, M. Targeting the

RdRp of Emerging RNA Viruses: The Structure-Based Drug Design Challenge.

Molecules **25**, doi:10.3390/molecules25235695 (2020).

22   Crotty, S. *et al.* The broad-spectrum antiviral ribonucleoside ribavirin is an

RNA virus mutagen. *Nature Medicine* **6**, 1375-1379, doi:10.1038/82191

(2000).

23   Shannon, A. *et al.* Rapid incorporation of Favipiravir by the fast and

permissive viral RNA polymerase complex results in SARS-CoV-2 lethal

mutagenesis. *Nat Commun* **11**, 4682, doi:10.1038/s41467-020-18463-z (2020).

24   Wang, M. *et al.* Remdesivir and chloroquine effectively inhibit the recently

emerged novel coronavirus (2019-nCoV) in vitro. *Cell Res* **30**, 269-271,

doi:10.1038/s41422-020-0282-0 (2020).

25   Sheahan, T. P. *et al.* An orally bioavailable broad-spectrum antiviral inhibits

SARS-CoV-2 in human airway epithelial cell cultures and multiple

coronaviruses in mice. *Sci Transl Med* **12**, eabb5883,

doi:10.1126/scitranslmed.abb5883 (2020).

26   Kabinger, F. *et al.* Mechanism of molnupiravir-induced SARS-CoV-2

mutagenesis. *Nat Struct Mol Biol* **28**, 740-746,

doi:10.1038/s41594-021-00651-0 (2021).

27 Yang, G., Franklin, M., Li, J., Lin, T. C. & Konigsberg, W. Correlation of the

Kinetics of Finger Domain Mutants in RB69 DNA Polymerase with Its

Structure. *Biochemistry* **41**, 2526-2534, doi:10.1021/bi0119924 (2002).

28 Burke, C. R. & Luptak, A. DNA synthesis from diphosphate substrates by

DNA polymerases. *Proc Natl Acad Sci U S A* **115**, 980-985,

doi:10.1073/pnas.1712193115 (2018).

29 Garforth, S. J., Parniak, M. A. & Prasad, V. R. Utilization of a

deoxynucleoside diphosphate substrate by HIV reverse transcriptase. *PLoS*

One **3**, e2074, doi:10.1371/journal.pone.0002074 (2008).

30 Gottesman, M. E. & Mustaev, A. Ribonucleoside-5'-diphosphates (NDPs)

support RNA polymerase transcription, suggesting NDPs may have been

substrates for primordial nucleic acid biosynthesis. *J Biol Chem* **294**,

11785-11792, doi:10.1074/jbc.RA119.009074 (2019).

31 Yin, W. *et al.* Structural basis for inhibition of the RNA-dependent RNA

polymerase from SARS-CoV-2 by remdesivir. *Science* **368**, 1499-1504,

doi:10.1126/science.abc1560 (2020).

32 Choy, K. T. *et al.* Remdesivir, lopinavir, emetine, and homoharringtonine

inhibit SARS-CoV-2 replication in vitro. *Antiviral Res* **178**, 104786,

doi:10.1016/j.antiviral.2020.104786 (2020).

33 Williamson, B. N. *et al.* Clinical benefit of remdesivir in rhesus macaques

infected with SARS-CoV-2. *Nature* **585**, 273-276,

doi:10.1038/s41586-020-2423-5 (2020).

34   Xie, X. *et al.* A nanoluciferase SARS-CoV-2 for rapid neutralization testing

and screening of anti-infective drugs for COVID-19. *Nat Commun* **11**, 5214,

doi:10.1038/s41467-020-19055-7 (2020).

35   Zhou, S. *et al.* beta-d-N4-hydroxycytidine Inhibits SARS-CoV-2 Through

Lethal Mutagenesis But Is Also Mutagenic To Mammalian Cells. *J Infect Dis*

224, 415-419, doi:10.1093/infdis/jiab247 (2021).

36   Malone, B. & Campbell, E. A. Molnupiravir: coding for catastrophe. *Nat*

Struct Mol Biol **28**, 706-708, doi:10.1038/s41594-021-00657-8 (2021).

37   Noreen, S., Maqbool, I. & Madni, A. Dexamethasone: Therapeutic potential,

risks, and future projection during COVID-19 pandemic. *Eur J Pharmacol*

894, 173854, doi:10.1016/j.ejphar.2021.173854 (2021).

38   Quimque, M. T. J. *et al.* Virtual screening-driven drug discovery of

SARS-CoV2 enzyme inhibitors targeting viral attachment, replication,

post-translational modification and host immunity evasion infection

mechanisms. *J Biomol Struct Dyn* **39**, 4316-4333,

doi:10.1080/07391102.2020.1776639 (2021).

39   Jiang, Y., Yin, W. & Xu, H. E. RNA-dependent RNA polymerase: Structure,

mechanism, and drug discovery for COVID-19. *Biochem Biophys Res*

Commun **538**, 47-53, doi:10.1016/j.bbrc.2020.08.116 (2021).

40   Meier, C. Nucleoside diphosphate and triphosphate prodrugs - An unsolvable

task? *Antivir Chem Chemother* **25**, 69-82, doi:10.1177/2040206617738656

(2017).

41   Gollnest, T. *et al.* Membrane-permeable Triphosphate Prodrugs of Nucleoside

Analogues. *Angew   Chem   Int   Ed   Engl*   **55**,   5255-5258,

doi:10.1002/anie.201511808 (2016).

42   Ferrero, D. S. *et al.* Supramolecular arrangement of the full-length Zika virus

NS5. *PLoS Pathog* **15**, e1007656, doi:10.1371/journal.ppat.1007656 (2019).

 Figure legends

Fig.1 RdRp uses both NTPs and NDPs as a substrate. **a** Size-exclusion chromatogram

of the SARS-CoV-2 nsp12-nsp7-nsp8(RdRp) complex. RdRp was also characterized

by SDS-PAGE. **b** The 13+23 nt RNA template-product duplex. The direction of RNA

extension is shown. The color of the depicted circles indicates the experimental

design: blue, RNA template strand; red, RNA product strand. The 5'end of the RNA

product contains a FAM fluorescent label. **c** Incubation of the RdRp with RNA duplex

and NTPs leads to RNA extension. Nsp12 alone has almost no polymerase activity,

and requires nsp7 and nsp8 to form an RdRp complex to have polymerase activity. **d**

RdRp was incubated with the 13+23 nt duplex in the presence of NMPs, NDPs or

NTPs respectively. These are showing duplicate experiments performed with different

batches RdRp. The positions of the template RNA and the full-length extension

product are indicated on denaturing gel. **e** Electrophoretic separation of reaction

products of single nucleoside diphosphates as substrates. The positions of the original

RNA and extension products are indicated. **f** Part of the Sanger sequencing

chromatogram of the RT-PCR products corresponding to the extended 10 nt RNA on

the 13+23 nt RNA duplex with NDPs as substrate. The dash line circle represents the

10 nt extension. **g** Schematic of high-throughput sequencing of RdRp reaction

products.

Fig.2 The activities of NDPs and NTPs are comparable. **a** Primer-extension reactions

on 13+23 nt RNA templates sampled at indicated time points. **b** Quantification of

elongation products for (**a**). The experiment was performed in triplet. **c** The 13+26 nt

RNA template-product duplex. The direction of RNA extension is shown. The 5'end

of the RNA product was labeled by FAM. **d** RdRp catalyzed reactions containing

different substrates. Lane 1: no substrate; lane 2: NTPs; lane 3: ATPGTPCTP; lane4:

ATPGTPCTP UDP; lane5: GTPCTPUTP; lane6: GTPCTPUTP ADP; lane7:

ATPCTPUTP; lane8: ATPCTPUTP GDP; lane9: ATPGTPUTP; lane10: ATPGTPUTP

CDP. The positions of the 13+26 nt RNA and extension products are indicated. **e**

Quantification of elongation products for (**d**).

Fig.3 RdRp incorporates NHC into RNA with MDP as a substrate. **a** Metabolism of

nucleoside analogues molnupiravir in host cell. A two-step model of

molnupiravir-induced RNA mutagenesis. **b** The 13+26 nt RNA allows for RNA

extension by 13 nucleotides. NHC monophosphate can be incorporated into growing

RNA instead of C or U. **c** RdRp catalyzed reactions containing different substrates.

Lane 1: no substrate; lane 2: NTPs; lane 3: ATPGTPUTP; lane4: ATPGTPUTP MTP;

lane5: ATPGTPUTP MDP; lane6: ATPGTPCTP; lane7: ATPGTPCTP MTP; lane8:

ATPGTPCTP MDP. The positions of the 13+26 nt RNA and extension products are

indicated. RNA elongation stalls at the expected positions when the cognate NTP is

absent from the reaction. **d** Quantification of elongation products for (**c**). **e** The 13+29

nt RNA allows for RNA extension by 16 nucleotides. NHC monophosphate can pair

with G or A for incorporation into growing RNA.

Fig.4 Substrate properties of other RNA viruses **a** RdRp of SARS-CoV-2 and

SARS-CoV and NS5 of ZIKV were incubated with the 13+23 nt RNA duplex in the

presence of NTPs or NDPs respectively. The positions of the template RNA and the

full-length extension product are indicated. **b** The transcription of T7 RNA

polymerase was measured by monitoring the signal of a fluorescent probe. Fluorescent signal continuously changed when transcription occurred, indicating the presence of RNA product. However, when NDPs were used as a substrate, the fluorescent signal remained unchanged, indicating the lack of RNA product. Substitution of ATP by ADP led to weak transcription, any other NTP replaced by NDP would inhibit the transcription completely.

Fig.5 Nucleotide diphosphate analog prodrugs designed by our strategy would have less potential to cause side effects. The terminal phosphate group (β-phosphate) is modified by lipophilic masks to give the prodrug more membrane permeability. Inside cell the masks are removed by enzymatic or chemical reactions. The ribose moiety and the base group should be designed in such a way that the diphosphate form drug can evade NDPK thus eliminating the risk of being incorporated into host mRNA or DNA (Note: NDPs and dNDPs are phosphorylated by the same NDPK).

a

NTPs   NDPs

0   10 20 30 40 50 60   0   10 20 30 40 50 60   s

← 23nt
← 13nt

b

—•— NTP as substrate
—■— NDP as substrate

Elongation %

Time/s

c

FAM
5' GCUAUGUGAGAUU   polymerization
3' CGAUACACUCUAAAAUUCCGGUCGAU 5'

d

1   2   3   4   5   6   7   8   9   10

← 26nt
← 19nt
← 17nt
← 15nt
← 13nt

e

Relative incorporation efficiency

2-NTP 4-ATPGTPCTPUDP 6-GTPCTPUTPADP 8-ATPCTPUTPGDP 10-ATPGTPUTPCDP

a

b

a

SARS-CoV-2   SARS-CoV   ZIKV NS5
no   NTP   NDP   no   NTP   NDP   no   NTP   NDP

← 23nt
] Stalled product
← 13nt

b